



Vol.7 No.2 | Juli 2024

JURNAL BIOMEDIKA DAN KESEHATAN

Publikasi dari Fakultas Kedokteran Universitas Trisakti

Editorial

Eyelid Infection (Blepharitis) Problem In The Elderly
Husnun Amalia, Megawati Yulia Wina Pratiwi, Ita Tazkiatul Izzati Mustopa

Original Article

Factors Affecting Stunted Status in Children Under Two Years at Karya Mulia Public Health Center 2023

Dery Wahyudi, Namira Alifah Fahiratunnisa, Muhammad Wildan Refaldi et al

Massive Transfusion and Intensive Management after Hysterectomy in Placenta Accreta
Alfi Marita Trsitiarti, Eric Edwin Yuliantara

Enhancing Salivary DNA Preservation via DNase I Inactivation: Role of Temperature and EDTA

Zulham Yamamoto, Nurul Sulviani, Sry Suryani Widjaja

The Relationship between Spiritual Quotient with Stress Levels in Medical Students in Semarang

Nurahmat Yanisa Irfandi, Ratih Widayati, Wijayanti Fuad

The Effect of Mung Bean Sprout Extract (*Phaseolus radiatus* L.) on Catalase Levels in Male Wistar Rats Induced Paraquat Herbicide

Angga Pria Sundawa, Reza Adityas Trisnasdi, Annisa Nurul Hikmah

The Analysis of Work Fatigue and Proposed Improvement Using the Bourdon Wiersma Method and New Seven Tools

Vera Devani, Abim Wahyu Maypando

Impact of Cotrimoxazole on the Development of Chicken Embryo Neural Tube

Alifia Shafanaura Pamuji, Cut Fauziah, Yanto Sandy Tjang et al

The Effect of Preoperative Oral Glucose Administration on Blood Glucose Levels in Diabetes Mellitus Patients in General Anesthesia Surgery

Dhiny Yolanda Harahap, Achsanuddin Hanafie, Andriamuri Primaputra Lubis et al

Matos-Carvalho Index as a Comparison to other Discriminant Indexes in Initial Beta Thalassemia Screening

Mulyadi Mulyadi, Mulyati Mulyati, Tri Ratnaningsih et al

Dietary Arrangements and Exercise Activities were Associated with Glycemic Control in Diabetes Patients at Grogol Petamburan Subdistrict Public Health Center

Sirly Hidayanti, Kartini Kartini

The Relationship Between Self-Esteem and Emotional Disorders in Adolescents at Senior High School

Rifat Adi Hendrianto, Erita Istriana

Case Report

Cholestatic Jaundice Due To Biliary Atresia With Cytomegalovirus And Malaria Infection: Blood Transfusion-Transmitted Infection?

Mario Mario, Yasmine Mashabi, Nany Hairunisa

Eyelid Dermoid Cyst: A Case Report

Riani Witjaksana, Husnun Amalia, Nany Hairunisa et al

Review Article

Microbiology Examination for Diagnosis of Mycobacterium other than Tuberculosis (MOTT) Infection

Arleen Devita, Ade Dharmawan

Emerging Threats in the Age of Pandemics: A Focus on COVID-19 and the Novel Sub-Variant EG 5 ("Eris"): Review Article

Raghda Alsayed, Hamsa Thamer, Seenar Hameed et al

Dewan Redaksi



Ketua Penyunting (Editor-in-Chief)

Dr. dr. Husnun Amalia, Sp.M
Departemen Ilmu Penyakit Mata, Fakultas Kedokteran
Universitas Trisakti, Indonesia

Wakil Ketua Penyunting (Deputy Editor-in-Chief)

Dr. Drs. ML. Edy Parwanto, M.Biomed
Departemen Biologi Kedokteran, Fakultas Kedokteran
Universitas Trisakti, Indonesia

Penyunting Ahli (Associate Editor)

dr. Nany Hairunisa, MCHSc
Departemen Ilmu Kedokteran Kerja, Fakultas Kedokteran
Universitas Trisakti, Indonesia

Dewan Penyunting (Editorial Boards)

Prof. Dr. dr. Adi Hidayat, MS (Indonesia)
Dr. dr. Yenny, Sp.FK (Indonesia)
dr. Laksmi Maharani, Sp. OG (Indonesia)
dr. Monica Dwi Hartanti, M.Biomed, PhD (Indonesia)
Dr. dr. Raditya Wratsangka, Sp.O.G, Subsp. Obginsos (Indonesia)
Dr. Siti Sugih Hartiningsih, S.Si, M.Kes (Indonesia)
dr. Dito Anurogo, M.Sc (Indonesia)
Prof. Dr. Emad Yousif (Irak)

Editor Produksi

Afton Muhandis, S.I.Kom

Alamat Korespondensi

Fakultas Kedokteran Universitas Trisakti
Jalan Kyai Tapa Np. 260 (Kampus B) Grogol, Jakarta 11440
Telp. 021-5672731 ext. 2502 | Fax. 021-5660706
www.jbiomedkes.org | E-mail: jbiomedkes@trisakti.ac.id

Penerbit

Fakultas Kedokteran Universitas Trisakti

Petunjuk Penulisan

Format penyusunan manuskrip

Manuskrip diketik pada kertas berukuran A4 (210 x 297 mm) dengan batas tepi 254 mm (*margin Normal*), huruf diketik dengan tipe huruf (*font*) *Times New Roman*, besar huruf (*font size*) 12 point dengan menggunakan spasi rangkap 2 (*double space*). Setiap bagian dari manuskrip dimulai pada halaman baru dengan urutan sebagai berikut: halaman judul, abstrak dan kata kunci (*keywords*), teks keseluruhan, ucapan terima kasih, daftar pustaka, tabel dan gambar (setiap tabel dan gambar pada halaman terpisah). Nomor halaman dicantumkan secara berurutan dimulai dari halaman judul pada sudut sebelah kanan bawah. Manuskrip sebaiknya ditulis maksimal 16 halaman.

Halaman judul

Halaman judul mencakup: a) judul manuskrip yang dibuat sesingkat mungkin, spesifik informatif dan ringkasan judul tidak lebih dari 40 karakter (hitung huruf dan spasi) yang dicantumkan dibawah judul, b) nama penulis disusun berurutan dengan nama mahasiswa sebagai pengarang pertama, diikuti oleh Pembimbing sebagai pengarang kedua. Nama penulis ditulis lengkap tanpa gelar dan dicantumkan seperti aslinya, tidak dibalik seperti pada daftar pustaka dan sitasi, c) alamat setiap penulis, nama departemen dan lembaga afiliasi penulis, d) nama dan alamat penulis untuk korespondensi serta nomor telepon, nomor faksimili, alamat email. Judul penelitian dibuat jelas, singkat, spesifik, informatif, dan sesuai dengan topik manuskrip. Jumlah kata tidak lebih dari 12 kata agar mudah dan cepat dipahami pembaca.

Abstrak dan kata kunci

Abstrak berjumlah 200-250 kata ditulis dalam bahasa Indonesia dan Inggris. Abstrak berisikan latar belakang termasuk tujuan penelitian, metode, hasil, dan kesimpulan. Kata kunci dicantumkan di bawah abstrak pada halaman yang sama sebanyak 4-6 kata. Bagian abstrak merupakan ringkasan dari isi makalah yang dibuat secara singkat, informatif, dengan menekankan pada aspek baru dan penting dari penelitian.

Teks

Teks makalah manuskrip dibagi dalam beberapa bagian dengan judul sebagai berikut: ***Pendahuluan, Metode, Hasil, Pembahasan, Kesimpulan dan saran.***

Pendahuluan

a. Latar belakang merupakan bagian yang menjelaskan alasan mengapa masalah ini penting untuk diteliti. Bagian ini memuat penjelasan mengapa masalah itu dipandang menarik, penting, dan perlu diteliti untuk mencari pemecahannya. Penjelasan dapat diperoleh dari penelusuran pustaka yang berkaitan erat dengan

masalah yang diteliti.

b. Keaslian penelitian dikemukakan dengan menunjukkan bahwa masalah yang dihadapi belum pernah dipecahkan oleh peneliti terdahulu atau dinyatakan dengan tegas perbedaan penelitian ini dengan penelitian terdahulu.

c. Tujuan penelitian yang menjelaskan hasil yang akan dicapai.

Metode

Metode penelitian berisi uraian terpadu dan sistematis mengenai bagaimana penelitian akan dilaksanakan. Metode terdiri dari :

a. Desain

b. Populasi / sampel (subjek) penelitian

Diuraikan kriteria inklusi dan eksklusi subjek penelitian, cara pemilihan sampel (subjek penelitian) secara random atau non-random, serta besar sampel yang akan di pilih. Teknik pemilihan sampel harus dijelaskan secara rinci. Bila perlu dibuat alur pemilihan sampel.

c. Bahan dan alat serta pengukuran

Bahan dan alat yang harus disajikan pada laporan terbatas pada bahan (materi) dan alat utama yang diperlukan untuk penelitian dan harus disebutkan spesifikasinya. Prosedur pengukuran perlu dijelaskan sesuai dengan tahapan yang dilakukan.

d. Alur kerja penelitian

Jalannya penelitian perlu dijelaskan mengenai jenis pendekatan yang dipakai untuk mendapatkan data, melalui pendekatan laboratorium, klinik, komunitas, observasi, dll.

e. Analisis data

Perlu dijelaskan jenis teknik statistik yang digunakan untuk menjawab masalah dan mencapai tujuan penelitian. Data yang diperoleh dapat dianalisis menggunakan teknik statistik secara parametrik dan non-parametrik.

Hasil

Suatu hasil penelitian hendaknya disajikan dengan jelas, logis, runut, sehingga mudah untuk dimengerti. Hasil penelitian sebaiknya ditampilkan selain dalam bentuk narasi dapat pula berupa gambar, tabel, foto, dan grafik sehingga memudahkan untuk dipahami. Hasil dan interpretasi analisis statistik dituliskan secara jelas dalam uraian hasil penelitian.

Pada tahap awal disajikan distribusi karakteristik subjek penelitian, yang biasanya dibuat pada sebuah tabel. Kemudian disajikan temuan penting yang diperoleh, kalau cukup banyak sebaiknya pada sebuah tabel. Bila terbatas misalkan hanya satu atau dua temuan cukup dalam bentuk narasi/teks.

Tabel, bagan/gambar, grafik dibuat dengan jelas, diberi nomor urut serta keterangan yang jelas. Keterangan

tabel diletakkan di atas tabel dan keterangan gambar diletakkan di bawah gambar. Maksimal tabel dan gambar 5. Semua tabel, grafik dan gambar diberi nomor dan keterangan yang jelas. Setiap tabel dianalisis dan diinterpretasi secara sistematis, dan hasilnya ditulis di bawah tabel tersebut. Perhitungan statistik detail tidak perlu ditulis dalam bagian hasil ini. Bila perhitungan statistik dianggap perlu ditulis, maka sebaiknya diletakkan dalam lampiran saja.

Pembahasan

Langkah awal harus diuraikan temuan penting yang diperoleh dari penelitian sesuai dengan tujuan penelitian. Kemudian bandingkan hasil penelitian yang diperoleh dengan hasil-hasil penelitian sebelumnya. Perlu dijelaskan kesesuaian dan ketidaksesuaian hasil penelitian yang didapat terhadap kerangka teori atau hasil penelitian lain yang telah dilakukan sebelumnya. Selanjutnya menggunakan teori-teori yang ada uraikan mekanisme terjadinya hasil penelitian tersebut. Bagian pembahasan juga menjelaskan mengenai kelemahan dan kelebihan penelitian yang telah dilakukan. Uraikan implikasi dari hasil penelitian yang diperoleh.

Kesimpulan

Kesimpulan hendaknya dibuat dalam bentuk narasi dan menguraikan secara singkat, jelas, padat menurut urutan yang sistematis. Bagian ini memuat tentang hasil penelitian yang telah diperoleh untuk menjawab tujuan penelitian. Saran menguraikan perlunya dilakukan penelitian lebih lanjut untuk memperbaiki kelemahan/keterbatasan dari penelitian yang telah dilakukan.

Ucapan terima kasih

Ditujukan kepada pihak-pihak yang memberikan bantuan dana dan dukungan antara lain dukungan dari bagian dan lembaga, para professional yang memberikan kontribusi dalam penyusunan makalah, dan untuk penguji I maupun penguji II. Pembimbing tidak perlu dicantumkan pada Ucapan Terima Kasih karena sudah dicantumkan sebagai penulis.

Daftar Referensi

Daftar referensi/rujukan hanya mencatumkan rujukan yang telah digunakan dan ditulis menurut sistem Vancouver.

Online Submission

Penulis dapat mengirim naskah manuskrip melalui *online submission* di *website* Jurnal Biomedika dan Kesehatan.

Langkah online submission:

1. Pastikan naskah yang akan diunggah sudah mengikuti semua petunjuk penulisan
2. Lakukan pendaftaran author di : <https://jbiomedkes.org/index.php/jbk/user/register>
3. Setelah terdaftar silakan unggah naskah manuskrip dan isi form yang terdapat di dalam website, dan ikuti langkah selanjutnya.

Daftar Cek Pengiriman Naskah Manuskrip

- ☐ Naskah manuskrip belum pernah dipublikasikan sebelumnya, juga tidak dalam pengajuan ke jurnal lain.
- ☐ File manuskrip harus berformat OpenOffice, Ms. Word atau RTF dokumen, *font* 12, *Times New Roman*, *double spacing*.
- ☐ Halaman judul harus memuat jelas judul, nama lengkap penulis tanpa gelar, departemen penulis, universitas, alamat lengkap, nomor telepon dan email.
- ☐ Pelaporan data manuskrip dari penelitian yang melibatkan manusia dan hewan memerlukan persetujuan formal (kaji etik) oleh dewan peninjau atau komisi etik institusi yang bersangkutan.
- ☐ Daftar rujukan memuat semua rujukan yang terdapat di dalam manuskrip dan ditulis sesuai urutan pengutipannya menggunakan sistem Vancouver.

Daftar Isi



Jurnal Biomedika dan Kesehatan - Vol. 7 No. 2 Juli 2024

Editorial

- Eyelid Infection (Blepharitis) Problem In The Elderly 133-137
Husnun Amalia, Megawati Yulia Wina Pratiwi, Ita Tazkiatul Izzati Mustopa

Original Article

- Factors Affecting Stunted Status in Children Under Two Years at Karya 138-144
Mulia Public Health Center 2023
Dery Wahyudi, Namira Alifah Fahiratunnisa, Muhammad Wildan Refaldi, Syarifah Nurul Yanti Rizki Syahab Asseggaf

- Massive Transfusion and Intensive Management after Hysterectomy in 145-151
Placenta Accreta
Alfi Marita Trsitiarti, Eric Edwin Yuliantara

Zxz

- Enhancing Salivary DNA Preservation via DNase I Inactivation: Role of 152-162
Temperature and EDTA
Zulham Yamamoto, Nurul Sulviani, Sry Suryani Widjaja

- The Relationship between Spiritual Quotient with Stress Levels in Medical 163-170
Students in Semarang
Nurahmat Yanisa Irfandi, Ratih Widayati, Wijayanti Fuad

- The Effect of Mung Bean Sprout Extract (*Phaseolus radiatus* L.) on 171-178
Catalase Levels in Male Wistar Rats Induced Paraquat Herbicide
Angga Pria Sundawa, Reza Adityas Trisnasdi, Annisa Nurul Hikmah

- The Analysis of Work Fatigue and Proposed Improvement Using the 179-189
Bourdon Wiersma Method and New Seven Tools
Vera Devani, Abim Wahyu Maypando

- Impact of Cotrimoxazole on the Development of Chicken Embryo Neural 190-198
Tube
Alifia Shafanaura Pamuji, Cut Fauziah, Yanto Sandy Tjang, Maria Selvester Thadeus

- The Effect of Preoperative Oral Glucose Administration on Blood Glucose 199-207
Levels in Diabetes Mellitus Patients in General Anesthesia Surgery
Dhiny Yolanda Harahap, Achsanuddin Hanafie, Andriamuri Primaputra Lubis, Rina Amelia

- Matos-Carvalho Index as a Comparison to other Discriminant Indexes in 208-217
Initial Beta Thalassemia Screening
Mulyadi Mulyadi, Mulyati Mulyati, Tri Ratnaningsih, Nur Imma Harahap, Indra Lesmana, Niken Satuti Nur Handayani

Dietary Arrangements and Exercise Activities were Associated with Glycemic Control in Diabetes Patients at Grogol Petamburan Subdistrict Public Health Center <i>Sirly Hidhayanti, Kartini Kartini</i>	218-227
The Relationship Between Self-Esteem and Emotional Disorders in Adolescents at Senior High School <i>Rifat Adi Hendrianto, Erita Istriana</i>	228-236
Case Report Cholestatic Jaundice Due To Biliary Atresia With Cytomegalovirus And Malaria Infection: Blood Transfusion-Transmitted Infection? <i>Mario Mario, Yasmine Mashabi, Nany Hairunisa</i>	237-244
Eyelid Dermoid Cyst: A Case Report <i>Riani Witjaksana, Husnun Amalia, Nany Hairunisa, Erlani Kartadinata, Anggraeni Adhiwardan, Noviani Prasetyaningsih</i>	245-250
Review Article Microbiology Examination for Diagnosis of Mycobacterium other than Tuberculosis (MOTT) Infection <i>Arleen Devita, Ade Dharmawan</i>	251-261
Emerging Threats in the Age of Pandemics: A Focus on COVID-19 and the Novel Sub-Variant EG 5 ("Eris"): Review Article <i>Raghda Alsayed, Hamsa Thamer, Seenar Hameed, Mohammed Kadhom, Nany Hairunisa, Husnun Amalia, Yasmine Mashabi, Dina Ahmed, Sarah Mahdi, Amani Husain, Israa Salman, Emad Yousif</i>	262-267

CASE REPORT

Eyelid Dermoid Cyst: A Case Report

Palpebral Kista Dermoid

Riani Witjaksana¹, Husnun Amalia¹, Nany Hairunisa², Erlani Kartadinata¹, Anggraeni Adhiwardani¹, Noviani Prasetyaningsih¹

¹Ophthalmology Department, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

²Occupational Medicine Department, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

 riani.witjaksana@trisakti.ac.id

 <https://doi.org/10.18051/JBiomedKes.2024.v7.245-250>

ABSTRACT

Dermoid and epidermoid cysts are prevalent benign orbital tumors in childhood, representing 3-9% of pediatric ocular tumors. Occasionally, dermoid cysts can attain significant size, leading to cosmetic deformities. A precise surgical approach to dermoid cysts can lead to considerable cosmetic improvement for the patient.

A 19-year-old female presented with swelling in the outer corner of her left eye, accompanied by minor discomfort in the affected area. Upon ophthalmologic examination 1.5x1 cm masses were detected in the upper left eyelid near the lateral canthus Clinical diagnosis of Dermoid cyst left eye upper eyelid made. Subsequently, a surgical excision under general anesthesia was scheduled. During the procedure, the cyst was successfully excised while preserving the cyst wall integrity.

The conclusion is imperative for ophthalmologists to recognize the necessity of precisely performing surgical excision of these cysts to achieve favorable functional and aesthetic outcomes.

Keywords: Dermoid Cyst; Epidermoid Cyst; Surgical Management; Pediatric Ocular Tumors; Eyelid.

ABSTRAK

Epidermoid dan dermoid merupakan tumor jinak pada orbita yang paling sering ditemukan pada anak dengan prevalensi 3-9%. Pada keadaan tumor yang besar dapat menyebabkan gangguan kosmetik pada penderitanya. Tindakan operatif sebaiknya dilakukan secara terencana agar didapatkan hasil kosmetik yang baik untuk Pasien

Perempuan 19 tahun datang dengan keluhan pembengkakan pada bagian atas kelopak atas mata kiri. Pada pemeriksaan didapatkan benjolan dengan ukuran 1.5x1 cm pada lateral kelopak mata dekat kantung lateral. Pasien didiagnosis dengan kista dermoid palpebra superior mata kiri. Pasien dilakukan Tindakan operatif dalam anestesi umum. Kista demoid dieksisi secara utuh.

Kesimpulannya adalah sangat penting bagi dokter mata untuk melakukan tindakan eksisi terencana untuk mencapai hasil estetik dan fungsi yang baik.

Kata Kunci: Kista Dermoid; Kista Epidermoid; Penatalaksanaan Operatif; Tumor Pada Anak; Kelopak Mata.

INTRODUCTION

Dermoid and epidermoid cysts are relatively common benign orbital tumors of childhood, constituting 5-9% of pediatric ocular tumors and accounting for 3-9% of all orbital masses.¹⁻⁴ These cysts typically arise from the sequestration of surface ectodermal elements during the closure of fetal suture lines in embryogenesis, often co-occurring with other congenital abnormalities. They can manifest in various locations within the orbit, such as the supratemporal and lacrimal sac areas.²⁻⁴ Superficial cysts tend to become symptomatic during childhood, whereas deeper orbital dermoids may not become clinically apparent until adulthood.^{1-4,6-7}

Orbital dermoid cysts most frequently appear in the lateral brow near the Fronto-Zygomatic Suture (FTZ), presenting as smooth, painless, oval masses that grow slowly. These masses may exhibit mobility or may be fixed to the periosteum at the underlying suture.^{1,5,8-9}

Surgical excision stands as the primary treatment for dermoid cysts, supported by numerous case reports confirming this approach through histopathological analysis. Overall, complete surgical excision remains the cornerstone for managing eyelid dermoid cysts, emphasizing meticulous removal to achieve favorable cosmetic results and minimize the likelihood of recurrence.⁵

CASE REPORT

A 19-year-old female presented to the outpatient department with complaints of swelling in the upper eyelid of her left eye, accompanied by mild intermittent pain in the affected area. The parents reported noticing this change when she was a year old, with intermittent growth observed over the past few years. There were no reports of diplopia on vertical gaze.

The ophthalmic examination revealed normal visual acuity in both eyes (1.0), normal intraocular pressure (13,4 mmHg), and unremarkable findings on both biomicroscopic and ophthalmologic assessments. Eye movements were within normal limits. Examination of the left eye identified a 1.5 cm x 1cm protrusion located 1/3 laterally from the upper eyelid margin, 0.5 cm above it. A tender, cystic structure measuring 0.5cm was palpable and fixed to the subcutaneous tissue. A clinical diagnosis of a dermoid cyst in the left upper lid was established, with X-ray imaging performed to assess orbital involvement and bone erosion. A surgical excision under general anesthesia was scheduled.

On 15 July 2023, an excision surgery was conducted wherein a horizontal incision was performed along the eyelid crease, leading to the dissection of the skin. The cyst wall was then located, and through careful blunt dissection, it was separated from its surrounding attachment. Subsequently, precise sharp dissection was carried out until the surface of the cyst was reached, allowing for the complete removal of the cyst without causing any rupture to its wall. The wound was then closed using a Nylon 6-0 suture. The excised specimen was sent for histopathological examination which revealed the presence of squamous epithelium lining with dermal components such as hair follicles, sebaceous, and sweat glands, thereby confirming the diagnosis of a dermoid cyst. Postoperatively, the patients exhibited a favorable cosmetic outcome, with no signs of inflammatory response during the subsequent follow-up examination. Wound closure was performed with Nylon 6-0 suture.




Figure 1. Pre-operative




Figure 2. Post-operative: A. day-1, B and C. 1 month.

DISCUSSION

Dermoid cyst represents the frequently used clinical term for benign cystic teratoma, choriostoma resulting from the entrapment of surface ectoderm into underlying mesenchyme along the embryonic line of fusion.¹ These cysts emerge as a consequence of the proliferation of epidermal cells within a restricted region of the dermis.¹⁰ The majority of dermoid cysts identified in ophthalmic clinical scenarios are superficial, typically presenting in early childhood as noticeable swellings in the eyebrow or eyelid.^{1,3,4} Orbital dermoid cysts can be categorized according to several criteria including juxta-sutural, sutural, or soft tissue location; superficial or deep positioning; intraosseous or extraosseous presence; and intraorbital or extraorbital localization. Periocular and orbital dermoid cysts are commonly divided into superficial and deep lesion categories.^{1,4,10} Certain inherited syndromes have been linked to epidermoid cysts such as Gardner syndrome, basal cell nevus syndrome, and pachyonychia congenital.² These cysts, which are smooth, painless, and either mobile or partially mobile, typically appear in the FTZ region. Symptoms such as proptosis, displacement, ptosis, or diplopia, which vary based on the depth and size of the cysts. Dermoid cysts in the orbit are commonly linked with the Fronto-Zygomatic and Fronto-Ethmoidal sutures, which play a significant role in their developmental process.^{5,6,8,9}

Deep orbital dermoid cysts are seldom observed, challenging to diagnose, and necessitate more elaborate surgical procedures such as orbitotomy. Typically, the identification of an orbital

dermoid is established before surgery through the assessment of the cyst's location, texture, and characteristic features on imaging modalities like X-ray or CT-scan.^{6-8,10}

Orbital dermoids require surgical excision due to their tendency to increase in size and release their contents into surrounding tissues. The cyst's contents are highly irritating and can lead to a severe inflammatory response, which results in fibrosis. It is advised to completely remove the dermoid together with its cyst wall through a surgical method that allows for ideal access to all parts of the lesion.^{1,5,11} While the primary aim of surgery is the total removal of epidermal and dermal component, some patients may manifest with the recurrent intraorbital lesion, introducing the challenging aspect of a delicate, scarred cyst all adhering closely to normal orbital structures. Aggressive removal of the recurrent lining may result in functional deficits postoperatively. In certain instances, although not ideal, surgical fistulization or marsupialization may be considered as a feasible alternative.⁵

In 1988, Kronish and Dorzbach provided a summary of previously proposed methods for removing dermoid cysts, which include: (1) performing a direct incision directly over the mass, (2) making incisions above, below, or through the eyebrow, (3) utilizing a superomedial approach through a Lynch incision, (4) performing medial lid splitting, (5) conducting a lateral canthotomy, and (6) making an incision in the upper lid crease.¹

An incision in the eyelid crease is recommended for the management of superficial supertemporal dermoid cysts. This particular incision provides favorable exposure, helps avoid vital structures, and results in an improved cosmetic outcome. When dealing with cysts located anterior to the FTZ, a superior eyelid crease incision offers sufficient exposure and favorable cosmetic results compared to an incision directly above the lesion.^{4,5} Cysts located well above the FTZ necessitate an infra-brow or low intra-brow incision to reduce the risk of frontal branch of the facial nerve damage associated with an eyelid crease approach, and to prevent inadvertent transection of the frontalis muscle that may occur with an incision directly over the cysts.^{5,13} Intraorbital cysts that expand from the lateral orbital wall sutures may present in early childhood but are more frequently diagnosed in adolescence and adulthood, as proptosis resulting from cyst expansion becomes more apparent.^{1,4,5} The majority of the lateral intraorbital dermoid cysts demonstrate osseous involvement, potentially leading to pressure erosion or an atypical morphology, emphasizing the significance of understanding the interaction between the cyst and bone in surgical planning. Complete excision of all epidermal elements of the cyst is imperative, and in instances where there exists a close association between the cyst wall and the bone, bone burring might be deemed necessary.^{4,8,9}

Aspiration of cyst walls was contraindicated owing to the potential recurrence of the cyst and the provocation of a severe inflammatory response. Poorly planned, delayed, partial removal or inadvertent rupture of the cyst could trigger an inflammatory reaction and lead to lasting consequences such as the development of an orbiculate fistula.^{1,8,9}

Complications associated with epidermoid cysts include infection, malignant transformation, and rupture. The frequency of malignant transformation from an epidermoid cyst to cutaneous squamous cell carcinoma is typically low, varying from 0.011% to 0.045%. In our case, there were no visible clinical signs of infection.^{14,5}

CONCLUSION

Depression and obesity are two prevalent disorders with a substantial influence on worldwide health and are closely linked and important health conditions. Various factors, including biological, psychological, and behavioral aspects, can impact the intricate relationship between depression and obesity, and play a role in maintaining the two-way relationship.

Interactions between modern antidepressants and newer antipsychotic medications are often significant in clinical practice. Weight gain is a common side effect of antidepressants and antipsychotics. Therefore, it is crucial to educate patients about drug side effects, hunger regulation, the importance of family support, and regular exercise from the start of treatment for depression and obesity. It is also crucial for psychiatrists and dietitians to collaborate to get the best therapeutic outcomes for patients with obesity and depression.

ACKNOWLEDGEMENT

All authors would like to Cilincing General Hospital for their support throughout this case report.

AUTHORS CONTRIBUTION

All authors contributed to this article.

FUNDING

This case report is not funded by any institution.

CONFLICT OF INTEREST

We declare that we have no conflicts of interest to disclose.

REFERENCES

1. Chopra V, Damodaran K. A case of dermoid cyst. *IP International Journal of Ocular Oncology and Oculoplasty*. 2022; 8(4): 270-272. Available from: <http://doi.org/10.18231/j.ijoo.2022.2022.059>
2. Korn BS, Burkat CN, Carter KD, et al. Congenital Orbital Anomalies. 2019-2020 Basic and Clinical Science Course, Section 07: Oculofacial Plastic and Orbital Surgery. San Francisco: American Academy of Ophthalmology; 2020.p.61-3.
3. Herd RW, Archer SM, Braverman RS, et al. Orbital Disorder. 2019-2020 Basic and Clinical Science Course, section 6: Pediatric Ophthalmology and Strabismus.. San Francisco: American Academy of Ophthalmology; 2020. p.251-4
4. Can yRajesh E, Charumathi R. Eyelid dermoid cyst: Case report a rare manifestation. *Pan Am J Ophthalmol* [serial online]. 2022 Jan [Cited 2023 Sept 15];4:3. Available from: <https://www.thepajo.org/text.asp?2022/4/1/3/335853>
5. Bajric J, Harris G. Surgical Technique: The spectrum of orbital dermoid cysts and their surgical management. *The International Journal on Orbital Disorders, Oculoplastic and Lacrimal Surgery*. 2019;39(4):266-75. doi: 10.1080/01676830.2019.1631357

6. Sezenoz AS, Arat YO, Tepeoglu M. An unusual eyelid mass: Tarsal dermoid cyst. *Saudi J Ophthalmol.* 2015;29:312-4
7. Bansal R, Honavar SG, Talloju SS, et.al. Orbital dermoid cyst. *Indian J Ophthalmol.* 2022;70(2):709. doi: 10.4103/ijo.IJO_145_22.
8. Leatherbarrow B. *Orbital Surgery: Orbital Disorders. Oculoplastic Surgery.* New York: Thieme Medical Publishers; 2020.p.505-07.
9. Howard BE, Masood MM, Clark JM, et.al. Intraorbital Dermoid Cyst With t: clinicalZygomaticofrontal Suture Erosion in an Adult. *The Journal of Craniofacial Surgery.* 2019; 30:514-5. doi: 10.1097/scs.0000000000005005
10. Balasudaram P, Garg A, Prabhakar A, et al. Evolution of epidermoid cyst into dermoid cyst: embryological explanation and radiological-pathological correlation. *Neuroradiol J.* 2019;32(2):92-7. Available from: <http://doi.org/10.1177/1971400918821086>
11. Pushker N, Meel R, Kumar A, et al. Orbital and periorbital dermoid/epidermoid cyst: a series of 280 cases and a brief review. *Can J Ophthalmol.* 2020;55(2):176-71. Available from: <http://doi.org/10.1016/j.cjo.2019.08005>
12. Mathur R, Saxena S. Epidermoid Cyst of The Eyelid-A Case Report And Review of Literature. *IOSR Journal of Dental and Medical Sciences.* Vol 17:1,2018. p.01-03.
13. Wilde C, Vahdani K, Thaung C, et.al. Persistent Orbital Inflammation following Complete excision of deep dermoid cysts. *Orbit* 2023; 42(4):434-36. Available from: <http://doi.org.10.1080/1676830.2022.2025855>
14. Shah K, Thacker M, Mehta K, et.al. A Case of Epidermoid Cyst of Eyelid. *GAIMS J Med Sci* 2023;3(2):84-86. Available from: <https://doi.org/10.5281/zenodo.8194726>
15. AlRubaian A, Alkatan HM, Al-Faky YH, et.al. Tarsal-related cysts: Different diagnoses with similar presentations. *Saudi J Ophthalmol.* 2019 Jul-Sep;33(3):209-213. Doi: 10.1016/j.sjopt.2019.01.004



This work is licensed under a Creative Commons Attribution Non-Commercial 4.0 International License

Eyelid Dermoid Cyst: A Case Report

by Anggraeni Adiwardhani FK

Submission date: 15-Jun-2026 09:08AM (UTC+0700)

Submission ID: 2980825967

File name: 14._572_CR_245-250.pdf (307.25K)

Word count: 2401

Character count: 13647

CASE REPORT

Eyelid Dermoid Cyst: A Case Report

Palpebral Kista Dermoid

Riani Witjaksana^{1✉}, Husnun Amalia¹, Nany Hairunisa², Erlani Kartadinata¹, Anggraeni Adhiwardani¹, Noviani Prasetyaningsih¹

¹Ophthalmology Department, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

²Occupational Medicine Department, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

✉ riani.witjaksana@trisakti.ac.id

<https://doi.org/10.18051/JBiomedKes.2024.v7.245-250>

ABSTRACT

Dermoid and epidermoid cysts are prevalent benign orbital tumors in childhood, representing 3-9% of pediatric ocular tumors. Occasionally, dermoid cysts can attain significant size, leading to cosmetic deformities. A precise surgical approach to dermoid cysts can lead to considerable cosmetic improvement for the patient.

A 19-year-old female presented with swelling in the outer corner of her left eye, accompanied by minor discomfort in the affected area. Upon ophthalmologic examination 1.5x1 cm masses were detected in the upper left eyelid near the lateral canthus Clinical diagnosis of Dermoid cyst left eye upper eyelid made. Subsequently, a surgical excision under general anesthesia was scheduled. During the procedure, the cyst was successfully excised while preserving the cyst wall integrity. The conclusion is imperative for ophthalmologists to recognize the necessity of precisely performing surgical excision of these cysts to achieve favorable functional and aesthetic outcomes.

Keywords: Dermoid Cyst; Epidermoid Cyst; Surgical Management; Pediatric Ocular Tumors; Eyelid.

ABSTRAK

Epidermoid dan dermoid merupakan tumor jinak pada orbita yang paling sering ditemukan pada anak dengan prevalensi 3-9%. Pada keadaan tumor yang besar dapat menyebabkan gangguan kosmetik pada penderitanya. Tindakan operatif sebaiknya dilakukan secara terencana agar didapatkan hasil kosmetik yang baik untuk Pasien

Perempuan 19 tahun datang dengan keluhan pembengkakan pada bagian atas kelopak atas mata kiri. Pada pemeriksaan didapatkan benjolan dengan ukuran 1,5x1 cm pada lateral kelopak mata dekat kantung lateral. Pasien didiagnosis dengan kista dermoid palpebra superior mata kiri. Pasien dilakukan Tindakan operatif dalam anestesi umum. Kista dermoid dieksisi secara utuh.

Kesimpulannya adalah sangat penting bagi dokter mata untuk melakukan tindakan eksisi terencana untuk mencapai hasil estetik dan fungsi yang baik.

Kata Kunci: Kista Dermoid; Kista Epidermoid; Penatalaksanaan Operatif; Tumor Pada Anak; Kelopak Mata.

INTRODUCTION

Dermoid and epidermoid cysts are relatively common benign orbital tumors of childhood, constituting 5-9% of pediatric ocular tumors and accounting for 3-9% of all orbital masses.^{1,4} These cysts typically arise from the sequestration of surface ectodermal elements during the closure of fetal suture lines in embryogenesis, often co-occurring with other congenital abnormalities. They can manifest in various locations within the orbit, such as the supratemporal and lacrimal sac areas.^{2,4} Superficial cysts tend to become symptomatic during childhood, whereas deeper orbital dermoids may not become clinically apparent until adulthood.^{1,4,6-7}

Orbital dermoid cysts most frequently appear in the lateral brow near the Fronto-Zygomatic Suture (FTZ), presenting as smooth, painless, oval masses that grow slowly. These masses may exhibit mobility or may be fixed to the periosteum at the underlying suture.^{1,5,8-9}

Surgical excision stands as the primary treatment for dermoid cysts, supported by numerous case reports confirming this approach through histopathological analysis. Overall, complete surgical excision remains the cornerstone for managing eyelid dermoid cysts, emphasizing meticulous removal to achieve favorable cosmetic results and minimize the likelihood of recurrence.⁵

CASE REPORT

A 19-year-old female presented to the outpatient department with complaints of swelling in the upper eyelid of her left eye, accompanied by mild intermittent pain in the affected area. The parents reported noticing this change when she was a year old, with intermittent growth observed over the past few years. There were no reports of diplopia on vertical gaze.

The ophthalmic examination revealed normal visual acuity in both eyes (1.0), normal intraocular pressure (13.4 mmHg), and unremarkable findings on both biomicroscopic and ophthalmologic assessments. Eye movements were within normal limits. Examination of the left eye identified a 1.5 cm x 1 cm protrusion located 1/3 laterally from the upper eyelid margin, 0.5 cm above it. A tender, cystic structure measuring 0.5 cm was palpable and fixed to the subcutaneous tissue. A clinical diagnosis of a dermoid cyst in the left upper lid was established, with X-ray imaging performed to assess orbital involvement and bone erosion. A surgical excision under general anesthesia was scheduled.

On 15 July 2023, an excision surgery was conducted wherein a horizontal incision was performed along the eyelid crease, leading to the dissection of the skin. The cyst wall was then located, and through careful blunt dissection, it was separated from its surrounding attachment. Subsequently, precise sharp dissection was carried out until the surface of the cyst was reached, allowing for the complete removal of the cyst without causing any rupture to its wall. The wound was then closed using a Nylon 6-0 suture. The excised specimen was sent for histopathological examination which revealed the presence of squamous epithelium lining with dermal components such as hair follicles, sebaceous, and sweat glands, thereby confirming the diagnosis of a dermoid cyst. Postoperatively, the patients exhibited a favorable cosmetic outcome, with no signs of inflammatory response during the subsequent follow-up examination. Wound closure was performed with Nylon 6-0 suture.




Figure 1. Pre-operative




Figure 2. Post-operative: A. day-1, B and C. 1 month.

DISCUSSION

Dermoid cyst represents the frequently used clinical term for benign cystic teratoma, choriostoma resulting from the entrapment of surface ectoderm into underlying mesenchyme along the embryonic line of fusion.¹ These cysts emerge as a consequence of the proliferation of epidermal cells within a restricted region of the dermis.¹⁰ The majority of dermoid cysts identified in ophthalmic clinical scenarios are superficial, typically presenting in early childhood as noticeable swellings in the eyebrow or eyelid.^{1,3,4} Orbital dermoid cysts can be categorized according to several criteria including juxta-sutural, sutural, or soft tissue location; superficial or deep positioning; intraosseous or extraosseous presence; and intraorbital or extraorbital localization. Periocular and orbital dermoid cysts are commonly divided into superficial and deep lesion categories.^{1,4,10} Certain inherited syndromes have been linked to epidermoid cysts such as Gardner syndrome, basal cell nevus syndrome, and pachyonychia congenita.² These cysts, which are smooth, painless, and either mobile or partially mobile, typically appear in the FTZ region. Symptoms such as proptosis, displacement, ptosis, or diplopia, which vary based on the depth and size of the cysts. Dermoid cysts in the orbit are commonly linked with the Fronto-Zygomatic and Fronto-Ethmoidal sutures, which play a significant role in their developmental process.^{5,6,8,9}

Deep orbital dermoid cysts are seldom observed, challenging to diagnose, and necessitate more elaborate surgical procedures such as orbitotomy. Typically, the identification of an orbital

dermoid is established before surgery through the assessment of the cyst's location, texture, and characteristic features on imaging modalities like X-ray or CT-scan.^{6-8,10}

Orbital dermoids require surgical excision due to their tendency to increase in size and release their contents into surrounding tissues. The cyst's contents are highly irritating and can lead to a severe inflammatory response, which results in fibrosis. It is advised to completely remove the dermoid together with its cyst wall through a surgical method that allows for ideal access to all parts of the lesion.^{15,11} While the primary aim of surgery is the total removal of epidermal and dermal component, some patients may manifest with the recurrent intraorbital lesion, introducing the challenging aspect of a delicate, scarred cyst all adhering closely to normal orbital structures. Aggressive removal of the recurrent lining may result in functional deficits postoperatively. In certain instances, although not ideal, surgical fistulization or marsupialization may be considered as a feasible alternative.⁵

In 1988, Kronish and Dorzbach provided a summary of previously proposed methods for removing dermoid cysts, which include: (1) performing a direct incision directly over the mass, (2) making incisions above, below, or through the eyebrow, (3) utilizing a superomedial approach through a Lynch incision, (4) performing medial lid splitting, (5) conducting a lateral canthotomy, and (6) making an incision in the upper lid crease.¹

An incision in the eyelid crease is recommended for the management of superficial supertemporal dermoid cysts. This particular incision provides favorable exposure, helps avoid vital structures, and results in an improved cosmetic outcome. When dealing with cysts located anterior to the FTZ, a superior eyelid crease incision offers sufficient exposure and favorable cosmetic results compared to an incision directly above the lesion.^{4,5} Cysts located well above the FTZ necessitate an infra-brow or low intra-brow incision to reduce the risk of frontal branch of the facial nerve damage associated with an eyelid crease approach, and to prevent inadvertent transection of the frontalis muscle that may occur with an incision directly over the cysts.^{5,13} Intraorbital cysts that expand from the lateral orbital wall sutures may present in early childhood but are more frequently diagnosed in adolescence and adulthood, as proptosis resulting from cyst expansion becomes more apparent.^{14,5} The majority of the lateral intraorbital dermoid cysts demonstrate osseous involvement, potentially leading to pressure erosion or an atypical morphology, emphasizing the significance of understanding the interaction between the cyst and bone in surgical planning. Complete excision of all epidermal elements of the cyst is imperative, and in instances where there exists a close association between the cyst wall and the bone, bone burring might be deemed necessary.^{4,8,9}

Aspiration of cyst walls was contraindicated owing to the potential recurrence of the cyst and the provocation of a severe inflammatory response. Poorly planned, delayed, partial removal or inadvertent rupture of the cyst could trigger an inflammatory reaction and lead to lasting consequences such as the development of an orbiculate fistula.^{1,8,9}

Complications associated with epidermoid cysts include infection, malignant transformation, and rupture. The frequency of malignant transformation from an epidermoid cyst to cutaneous squamous cell carcinoma is typically low, varying from 0.01% to 0.045%. In our case, there were no visible clinical signs of infection.^{14,5}

CONCLUSION

Depression and obesity are two prevalent disorders with a substantial influence on worldwide health and are closely linked and important health conditions. Various factors, including biological, psychological, and behavioral aspects, can impact the intricate relationship between depression and obesity, and play a role in maintaining the two-way relationship.

Interactions between modern antidepressants and newer antipsychotic medications are often significant in clinical practice. Weight gain is a common side effect of antidepressants and antipsychotics. Therefore, it is crucial to educate patients about drug side effects, hunger regulation, the importance of family support, and regular exercise from the start of treatment for depression and obesity. It is also crucial for psychiatrists and dietitians to collaborate to get the best therapeutic outcomes for patients with obesity and depression.

ACKNOWLEDGEMENT

All authors would like to Cilincing General Hospital for their support throughout this case report.

5

AUTHORS CONTRIBUTION

All authors contributed to this article.

FUNDING

This case report is not funded by any institution.

CONFLICT OF INTEREST

12

We declare that we have no conflicts of interest to disclose.

REFERENCES

1. ChopraV, Damodaran K. A case of dermoid cyst. IP International Journal of Ocular Oncology and Oculoplasty.2022; 8(4): 270-272. Available from: <http://doi.org/10.18231/ijoo.2022.2022.059>
2. Korn BS, Burkat CN, Carter KD, et.al. Congenital Orbital Anomalies. 2019-2020 Basic and Clinical Science Course, Section 07: Oculofacial Plastic and Orbital Surgery. San Francisco: American Academy of Ophthalmology; 2020.p.61-3.
3. Hered RW, Archer SM, Braverman RS, et al. Orbital Disorder.2019-2020 Basic and Clinical Science Course, section 6: Pediatric Ophthalmology and Strabismus.. San Francisco: American Academy of Ophthalmology;2020. p.251-4
4. Can yRajesh E, Charumathi R. Eyelid dermoid cyst: Case report a rare manifestation. Pan Am J Ophthalmol [serial online].2022 Jan [Cited 2023 Sept 15];4:3. Available from: <https://www.thepajo.org/text.asp?2022/4/1/3/335853>
5. Bajric J, Harris G. Surgical Technique: The spectrum of orbital dermoid cysts and their surgical management. The International Journal on Orbital Disorders, Oculoplastic and Lacrimal Surgery.2019;39(4):266-75. doi:10.1080/01676830.2019.1631357

6. Sezenoz AS, Arat YO, Tepeoglu M. An unusual eyelid mass: Tarsal dermoid cyst. *Saudi J Ophthalmol.* 2015;29:312-4
7. Bansal R, Honavar SG, Talloju SS, et.al. Orbital dermoid cyst. *Indian J Ophthalmol.* 2022;70(2):709. doi: 10.4103/ijoo.IJO_145_22.
8. Leatherbarrow B. *Orbital Surgery: Orbital Disorders. Oculoplastic Surgery.* New York: Thieme Medical Publishers; 2020.p.505-07.
9. Howard BE, Masood MM, Clark JM, et.al. Intraorbital Dermoid Cyst With t: clinicalZygomaticofrontal Suture Erosion in an Adult. *The Journal of Craniofacial Surgery.* 2019; 30:514-5. doi: 10.1097/scs.0000000000005005
10. Balasudaram P, Garg A, Prabhakar A, et al. Evolution of epidermoid cyst into dermoid cyst: embryological explanation and radiological-pathological correlation. *Neuroradiol J.* 2019;32(2):92-7. Available from: <http://doi.org/10.1177/1971400918821086>
11. Pushker N, Meel R, Kumar A, et al. Orbital and periorbital dermoid/epidermoid cyst: a series of 280 cases and a brief review. *Can J Ophthalmol.* 2020;55(2):176-71. Available from: <http://doi.org/10.1016/j.cjo.2019.08005>
12. Mathur R, Saxena S. Epidermoid Cyst of The Eyelid-A Case Report And Review of Literature. *IOSR Journal of Dental and Medical Sciences.* Vol 17:1,2018. p.01-03.
13. Wilde C, Vahdani K, Thaug C, et.al. Persistent Orbital Inflammation following Complete excision of deep dermoid cysts. *Orbit* 2023; 42(4):434-36. Available from: <http://doi.org/10.1080/1676830.2022.2025855>
14. Shah K, Thacker M, Mehta K, et.al. A Case of Epidermoid Cyst of Eyelid. *GAIMS J Med Sci* 2023;3(2):84-86. Available from: <https://doi.org/10.5281/zenodo.8194726>
15. AlRubaian A, Alkatan HM, Al-Faky YH, et.al. Tarsal-related cysts: Different diagnoses with similar presentations. *Saudi J Ophthalmol.* 2019 Jul-Sep;33(3):209-213. Doi: 10.1016/j.sjopt.2019.01.004



This work is licensed under a Creative Commons Attribution Non-Commercial 4.0 International License

Eyelid Dermoid Cyst: A Case Report

ORIGINALITY REPORT

13%	9%	7%	3%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	"Ocular Adnexal Lesions", Springer Science and Business Media LLC, 2019 Publication	2%
2	www.healio.com Internet Source	1%
3	www.unboundmedicine.com Internet Source	1%
4	www.iosrjournals.org Internet Source	1%
5	Atut Cicih Mayasari, Nugroho Abikusno, Laksmi Maharani, Raditya Wratsangka. "Painless Placental Abruptio with 80% Retroplacental Bleeding: Case Report", Jurnal Biomedika dan Kesehatan, 2024 Publication	1%
6	Submitted to Universitas Trisakti Student Paper	1%
7	doi.org Internet Source	1%
8	Marrakchi Salma, El Aouadi Salma, El Harras Yahya, Sassi Samia et al. "Orbital dermoid cyst in children: A case report", Radiology Case Reports, 2025 Publication	1%
9	jbiomedkes.org Internet Source	1%

10	"Ocular Pathology", Springer Science and Business Media LLC, 2025	1 %
Publication		

11	Submitted to Universitas Pasundan	1 %
Student Paper		

12	nagasaki-u.repo.nii.ac.jp	1 %
Internet Source		

13	jsmc.univsul.edu.iq	1 %
Internet Source		

Exclude quotes	Off	Exclude matches	< 10 words
Exclude bibliography	On		