tankings Journal Value Country Rankings Viz Tools

Custom email Custom email made easy made easy

i

Tatiana Boyko Ta

Acta Oc

₃ Scandinavica ∂




SCOPE


Acta Odontologica Scandinavica publishes papers conveying new knowledge within all areas of oral health and disease sciences, except dental biomaterials, which will be referred to Acta Biomaterialia Odontologica Scandinavica. Original research papers as well as critical state-of-the-art reviews relevant to the diagnosis, epidemiology, health service, prevention, etiology, pathogenesis and treatment of diseases affecting tissues of the oral cavity and associated structures will be considered for publication. All manuscripts including scientific progress are reviewed by members of a panel of international referees, with preference given to analytical studies based on hypotheses. Papers on cause and effect or explanatory/associative relationships for experimental or observational studies are also favored and simple descriptive studies have a low priority for publication. Acta Odontologica Scandinavica has been published since 1939 and is sponsored by the Dental Associations and Dental Schools in Denmark, Finland, Iceland, Norway and Sweden with editorial responsibility alternating between these countries.

Buy vitamins and supplements











← Show this widget in your own website

Just copy the code below and paste within your html code:

<a href="https://www.scimaç



Metrics based on Scopus® data as of March 2025

G gabriela león 3 years ago

Good evening, greetings from Gabriela and Magaly from the University of Cuenca, Ecuador. We would like to know if you accept to publish this topic :

Stability of anterior open bite treatment: a literature review

We look forward to your prompt response.

Thank you in advance for your help

reply



Melanie Ortiz 3 years ago

Dear Gabriela,

Thank you for contacting us.

We are sorry to tell you that SCImago Journal & Country Rank is not a journal. SJR is a portal with scientometric indicators of journals indexed in Elsevier/Scopus.

SCImago Team

We suggest you visit the journal's homepage (See submission/author guidelines) or contact the journal's editorial staff , so they could inform you more deeply.

Best Regards, SCImago Team

S sadi shirshab 3 years ago

Hi

dear sir

I am a specialist dentist, Dr. Sadi Shirshab Thiab.. I have research that I completed with my own efforts. I want to publish it in your magazine if it is included in the Scopus. I would like the approval of the publication to be quick. Thank you.

Sadi

reply



Melanie Ortiz 3 years ago

Dear Sadi,

Thank you for contacting us.

We are sorry to tell you that SCImago Journal & Country Rank is not a journal. SJR is a portal with scientometric indicators of journals indexed in Elsevier/Scopus. We suggest you visit the journal's homepage (See submission/author guidelines) or contact the journal's editorial staff , so they could inform you more deeply. Best Regards, SCImago Team

Email (will not be published)	
I'm not a robot reCAPTCHA Privacy - Terms Submit	

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.


Developed by:

Powered by:





Follow us on @ScimagoJR



Legal Notice

Privacy Policy

^

^

ACTA ODONTOLOGICA SCANDINAVICA

ABOUT - CONTENT - INSTRUCTIONS FOR AUTHORS - POLICIES AND GUIDELINES - CONTACT

Q SEARCH

HOME / Editorial Team

Editorial Team

Editor-in-Chief

Palle Holmstrup

Professor em, DrOdont, OdontDr (hc), PhD, DDS,

Section for Oral Biology and Immunopathology, University of Copenhagen, Faculty of Health and Medical Sciences, Department of Odontology, University of Copenhagen, Denmark

Research Area: Periodontology

Associate Editors

Lina Stangvaltaite-Mouhat

Seniorforsker/Senior Researcher, DDS, MPH, PhD Tannhelsetjenestens kompetansesenter Øst, i Norge Research area: Cariology and Endodontics

Professor

Department of Clinical Dentistry. University of Bergen, Norway

Research area: Community Dentistry

Mats Jontell

Professor em. OdontDr. FDSRCSEd

Department of Oral Medicine and Pathology, Institute of Odontology, Sahlgrenska Academy, University of Gothenburg, Sweden Research area: Oral Medicine and Pathology

Editorial Board

Iva Z. Alajbeg - School of Dental Medicine, University of Zagreb, Croatia

Gregory S. Antonarakis - Dental School, University of Geneva, Switzerland

Lene Baad-Hansen - Department of Dentistry and Oral Health, University of Aarhus, Denmark

Daniel Belstrøm - Department of Odontology, University of Copenhagen, Denmark,

Jon Dahl - Department of Dentistry, University of Oslo, Norway

Gunnar Dahlén - Department of Odontology, University of Gothenburg, Sweden

Göran Dahllöf - Department of Dental Medicine, University of Stockholm, Sweden

Christian Damgaard - Department of Odontology, University of Copenhagen, Denmark

Morten Enersen - Department of Dentistry, University of Oslo, Norway

Anders Gustafsson - Department of Dental Medicine, University of Stockholm, Sweden

Elin Hadler-Olsen - Department of Medical Biology, University of Tromsø, Norway

Bengt Hasseus - Department of Odontology, University of Gothenburg, Sweden

Keith Horner - School of Dentistry, University of Manchester, UK

Lise-Lotte Kirkevang - Department of Dentistry and Oral Health, University of Aarhus, Denmark

Björn Klinge - Department of Dental Medicine, University of Stockholm, Sweden

Thomas Kvist - Department of Odontology, University of Gothenburg, Sweden Satu Lahti - Department of Community Dentistry, University of Turku, Finland

Louise Hauge Matzen - Department of Dentistry and Oral Health, University of Aarhus, Denmark

Anne Marie Lynge Pedersen - Department of Odontology, University of Copenhagen, Denmark

Fernanda Cristina Petersen - Department of Dentistry, University of Oslo, Norway

Bjarni Elvar Pétursson - Department of Odontology, University of Reykjavík, Iceland

Kaisu Pienihäkkinen - Department of Community Dentistry, University of Turku, Finland

Pertti Pirttiniemi - Department of Medicine, University of Oulu, Finland

Kasper Rosing - Department of Odontology, University of Copenhagen Amer Sehic - Department of Dentistry, University of Oslo, Norway

Rasa Skudutyte-Rysstad - Faculty of Dentistry, University of Oslo, Norway

Indre Stankeviciene - Vilnius University Hospital, Lithuania

Marte-Mari Uhlen-Strand - Tannhelsetienestens Kompetansesenter Øst. Norway

Vaska Vandevska-Radunovich - Department of Dentistry, University of Oslo, Norway

Mieszko Wieckiewicz - Department of Experimental Dentistry, Wrocław Medical University, Poland



Acta Odontologica Scandinavica publishes original research papers as well as critical reviews relevant to the diagnosis, epidemiology, health service, prevention, aetiology, pathogenesis, pathology, physiology, microbiology, development and treatment of diseases affecting tissues of the oral cavity and associated structures including papers on cause and effect or explanatory/associative relationships for experimental or observational studies.



Acta Odontologica Scandinavica, Online ISSN: 1502-3850
Published by Medical Journals Sweden AB (559296-5023) under the terms of the Creative Commons CC-BY 4.0 license. Kungsängsvägen 27, 753 23 Uppsala, Sweden



HOME / ARCHIVES / Vol. 84: (2025) Acta Odontologica Scandinavica

Vol. 84: (2025) Acta Odontologica Scandinavica

RESEARCH ARTICLE

Expectations of orthodontic treatment among 7–12 year-old children – a cross sectional study

Anne-Maria Aulu, Tuija Pikkusaari, David Rice, Heidi Arponen

Published: 2025-06-18

☐ PDF ☐ HTML ☐ XML ☐ APPENDIX

Effect of occlusal reduction on post-operative pain of symptomatic and asymptomatic molar teeth

Aysenur Kamacı Esen, Fatma Furuncuoğlu, Fatima Betul Basturk, Muhammet Nuri Taşcıoğlu, Masoud Parirokh

Published: 2025-06-11

Impact of molar incisor hypomineralization on oral health-related quality of life, dental fear and anxiety in Swedish

Adnan Hajdarević, Birgitta Jälevik, Emina Čirgić, Agneta Robertson, Nina Sabel

Published: 2025-06-11

Danish endodontic practice-based research network: follow-up data

Lise-Lotte Kirkevang, Lotte Hein Sørensen

343-348

Published: 2025-06-10

⚠ PDF 🖹 HTML 🖹 XML

 ${\bf Clinical\ efficacy\ of\ diode\ laser\ for\ pulpotomy\ in\ primary\ teeth:\ a\ meta-analysis\ of\ randomised\ controlled\ trials}$

Yue Gao, Mina Hu, Iian Xu

Published: 2025-06-10

A PDF
B HTML
B XML

Associations between odontogenic and sinus pathologies – a low-dose CBCT study

Veli-Pekka Riekki, Mika T. Nevalainen, Marianne Haapea, Annina Sipola, Soili Kallio-Pulkkinen, Michaela K. Bode

Published: 2025-06-03

⚠ PDF 🖹 HTML 🖹 XML

 $\ \, \text{Evidence of mutations in tumour suppressor genes among oral cancer in Naswar, smokeless to bacco users } \,$ Fatima Iqbal, Sajjad Ahmad, Hoor Maryam , Humaira Amin

Published: 2025-06-03



publishes original research papers as well as critical reviews relevant to the diagnosis, epidemiology, health service, prevention, aetiology, pathogenesis, pathology, physiology, microbiology, development and treatment of diseases affecting tissues of the oral cavity and associated structures including papers on cause and effect or explanatory/associative relationships for experimental or observational





A PDF B HTML B XML Self-reported oral health outcomes after switching to a novel nicotine pouch technology: a pilot study Giusy Rita Maria La Rosa, Karl Fagerström, Sebastiano Antonio Pacino, Jan Kowalski, Renata Górska, Stefan Gospodaru, 292-298 Gheorghe Bordeniuc, Valeriu Fala, Amaliya Amaliya, Iain Chapple, Riccardo Polosa Published: 2025-05-27 PDF
B HTML
B XML A retrospective study analysing outcomes of the coronectomy procedure at a university dental teaching clinic in Umar Sharif, Stein Atle Lie, Cecilie Gudveig Gjerde Published: 2025-05-26 ∄ PDF 🖹 HTML 🖹 XML Impact of fixed appliance treatment on root resorption in root canal-treated teeth: a systematic review and meta-Yongfang Feng, Rong Wang, Yumin Zhou, Shengnan Zhan Published: 2025-05-20 Comparison of the effectiveness of different irrigation activation methods on biofilm removal in lateral canals – an

in vitro study Uğur Aydın, Bilal Turan, Emre Çulha, Muazzez Naz Baştürk Özer, Melih Özdemir

Published: 2025-05-20

∄ PDF 🖹 HTML 🖹 XML

Quality of life among adults with less severe malocolusions seeking orthodontic treatm

Quality of life among adults with less severe malocclusions seeking orthodontic treatment Narmina Sandlund, Idil Burt, Robert Schibbye 251-257

Published: 2025-05-13

Is the presence of accessory mandibular canals associated with the dimensions of the mandibular canal?

Emre Sözen, Hasan Akpınar

Published: 2025-05-13

 $Association\ between\ subjective\ orthodontic\ treatment\ need,\ oral\ health-related\ quality\ of\ life,\ and\ occlusal\ deviations\ in\ a\ cohort\ of\ 17-year-old\ adolescents\ in\ Finland$

Heidi Arponen, Eija Latvakoski, Linnea Närhi, Anna-Sofia Silvola

Published: 2025-05-13

E-cadherin expression in the tongue epithelium of streptozotocin-induced diabetic rats: an exploratory study

Hassan Hamed Kaabi, Abdullah Mohamed Alsoghier, Islam Abdulrahim Alredah, Rayan Mohammed Alqahtani, Ibrahim Suliman Alsanie, Hanan Abdulgafour Balto

Published: 2025-05-13


A PDF B HTML B XML

Plasma levels of immune system activation markers Neopterin and Kynurenine-to-Tryptophan Ratio, and oral


health among community-dwelling adults in Norway: a population-based, cohort study

Sarala Banjara, Ellen Berggreen, Jannicke Igland, Anne-Kristine Astrøm, Øivind Midttun, Dagmar Bunæs, Gerhard Sulo

Published: 2025-05-13



Oral health-related quality of life in children attending university special needs and paediatric dental clinics in Trinidad and Tobago: A parental perspective Ramaa Balkaran , Satu Lahti, Visha Ramroop, Jorma I. Virtanen ⚠ PDF 📑 HTML 🗎 XML The effect of different variables on push-out tests in 3D-printed oval and round-shaped root canals: a methodological study Published: 2025-02-13 Who requires dental treatment under general anesthesia due to pain and severe dental anxiety? Findings from panoramic X-ray images and anamnesis Vilde Aardal , Caroline Hol , Anne Rønneberg , Sudan Prasad Neupane, Tiril Willumsen Published: 2025-02-04 PDF ☐ HTML ☐ XML ☐ APPENDIX Parental knowledge, awareness, and attitudes towards children's oral habits: a descriptive cross-sectional study Sarah Almugairin, Alanoud Alwably, Nada Alayed, Alanoud Algazlan, Hadeel Alrowaily, Elzahraa Eldwakhly, Abdullazez Published: 2025-01-30 $Translation, cross-cultural \ adaptation, and \ psychometric \ properties \ of the \ Finnish \ version \ of the \ Malocclusion \ Impact \ Questionnaire \ (MIQ)$ Lucas Arrais Campos, Terhi Kaikkonen, Kaisa Ylitervo, Leena Ylikontiola, Anna-Sofia Silvola Published: 2025-01-30 ☐ PDF ☐ APPENDIX 1 ☐ APPENDIX 2 ☐ HTML ☐ XML Patient-reported outcomes of esthetics, function and oral hygiene with single dental implants 10–15 years after placement: a cross-sectional study Viveca Wallin Bengtsson, Christel Lindahl, Sven Scholander Published: 2025-01-21 Oral lesions and disorders and their prevalence arising from the use of illicit drugs in a prison population Marta Relvas, Luciana Rocha, Paulo Rompante, Filomena Salazar, Luís Monteiro, Ana Sofia Vinhas, Maria Gonçalves Published: 2025-01-21 ☐ PDF ☐ HTML ☐ XML Oral health-related quality of life of community-dwellers in the canton of Bern, Switzerland Roberta Borg-Bartolo, Andrea Roccuzzo, Christian Tennert, Maria Prasinou, Maurus Jäggi, Pedro Molinero-Mourelle, Michael M. Bornstein, Guglielmo Campus Published: 2025-01-16 A PDF B HTML B XML Diagnosis of approximal caries in children with convolutional neural networks based detection algorithms on radiographs: A pilot study Zeynep Seyda Yavsan, Hediye Orhan, Enes Efe, Emrehan Yavsan Bublished: 2025 01 05





☐ PDF ☐ APPENDICES

 $Effects \ of \ different \ or al \ barrier \ membranes \ on \ the \ efficacy \ and \ safety \ of \ guided \ bone \ regeneration \ in \ patients \ with$ dental implants: a systematic review and meta-analysis Haozhou Tang, Yonghui Zhang 318-331 Published: 2025-06-10 ∄ PDF ${\bf Manifestations\ of\ the\ oral\ mucosa\ and\ salivary\ glands\ in\ irritable\ bowel\ syndrome\ and\ microscopic\ colitis\ -\ A}$ Hanna Göthlin, Bengt Hasséus, Klas Sjöberg, Maria Bankvall Published: 2025-06-11 ∄ PDF 🖹 HTML 🖹 XML $Effects \ of \ total \ temporoman dibular \ joint \ replacement \ with \ all oplastic \ prosthesis - a \ systematic \ review$ Pär Morin, Jani Talvilahti, Mattias Ulmner, Per Alstergren, Eva Nordendahl, Aron Naimi-Akbar 258-265 ∄ PDF 🖺 HTML 🖹 XML Efficacy of nonsurgical periodontal treatment on patients with periodontitis and type 2 diabetes mellitus: a systematic review and Bayesian network meta-analysis Xuejie Xie, Jun Xu, Yiming Li, Li Tang, Gulinuer Awuti Published: 2025-05-13 A PDF A HTML A XML Prevalence of bruxism in patients affected by epilepsy: a systematic review and meta-analysis ${\sf Giuseppe\ Minervini,\ Rocco\ Franco,\ Marco\ Di\ Blasio,\ Mirko\ Martelli,\ Marco\ Gargari,\ Patrizio\ Bollero,\ Marco\ Cicciù}$ 155-164 PDF ☐ HTML ☐ XML Application of machine learning in dentistry: insights, prospects and challenges Lin Wang, Yanyan Xu, Weiqian Wang, Yuanyuan Lu Published: 2025-03-27 Ď PDF 🖹 HTML 🖺 XML Is root canal treatment and an indirect coronal restoration of a mandibular first molar cost-effective compared to extraction and an implant-supported crown? A decision analytic approach Nikki Savolainen, Fredrik Frisk, Thomas Kvist Published: 2025-02-27

Ď PDF 🗎 HTML 🗎 XML

Acta Odontologica Scandinavica. Online ISSN: 1502-3850
Published by Medical Journals Sweden AB (559296-5023) under the terms of the <u>Creative Commons CC-BY 4.0</u> license. Kungsängsvägen 27, 753 23 Uppsala, Sweden
Contact us.

Platform & workflow by OJS / PKP

MJS PUBLISHING

ORIGINAL ARTICLE

Dimensionality of the Fonseca Anamnestic Index and validation of its short-form derivative

Adrian Ujin Yap^{a,b,c,d,e,f}, Indrayadi Gunardi⁹, Darren Zong Ru Lee^h and Carolina Marpaung⁶

^aDivision of Dentistry, Ng Teng Fong General Hospital, Singapore, Singapore; ^bFaculty of Dentistry, National University Health System, Singapore, Singapore; ^cNational Dental Research Institute, Singapore, Singapore; ^dNational Dental Center, Singapore, Singapore; ^eDuke-NUS Medical School, Singapore Health Services, Singapore, Singapore; ^fDepartment of Prosthodontics, Faculty of Dentistry, Universitas Trisakti, Indonesia; ^gDepartment of Oral Medicine, Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia; ^hSchool of Health and Social Sciences, Nanyang Polytechnic, Singapore, Singapore

ARSTRACT

Objectives: Recently, the Short-Form Fonseca Anamnestic Index (SFAI) was shown to have high diagnostic accuracy when compared to the Diagnostic Criteria for Temporomandibular Disorders (DC/TMD) in patient samples. This study investigated the dimensionality of the parent instrument (Fonseca Anamnestic Index [FAI]) and validated its main component using Rasch analysis in non-patient populations.

Methods: FAI data from a total of 901 participants from Singapore and Indonesia with a mean age 19.30 ± 1.48 years (65.0% women) were examined. Of these, 53.8% were FAI positive and 46.2% were FAI negative. Principal Component Analysis (PCA) was performed to extract the main component of the FAI using an eigenvalue > 1.0 and direct oblimin rotation with item loading of > 0.40. Rasch analysis was subsequently carried out on the items of the main component.

Results: The FAI was found to be multidimensional with the main component involving items F1, F2, F3, F6, and F7 which were the items of the SFAI. The SFAI had moderate internal consistency (Cronbach alpha = 0.63) and fitted the Rasch model with person and item infit/outfit mean square (MnSq) values of 0.98/0.96 and 1.00/0.96 logits respectively. The infit/outfit MnSq of the SFAI items ranged from 0.82 to 1.06 logits with Item F2 (side-movement difficulty) being the most difficult and item F3 (muscle pain) the easiest.

Conclusions: The FAI is multidimensional with the main component comprising the five items of the SFAI that fitted the Rasch model. With its good Rasch validity, separation, and reliability, the SFAI is a promising tool for TMD screening.

ARTICLE HISTORY

Received 3 September 2024 Accepted 10 January 2025

KEYWORDS

Temporomandibular disorders; general population; questionnaires; validity; methods

Introduction

Temporomandibular disorders (TMDs) are a diverse group of neuromuscular and musculoskeletal conditions affecting the temporomandibular joints (TMJs), muscles of masticatory, and adjoining structures. In addition to TMJ and masticatory muscle pain, the other signs/symptoms of TMDs include headaches, neck pain, TMJ noises, jaw functional limitations, as well as abnormal jaw movements [1, 2]. The complex etiology of TMDs involves a myriad of biopsychosocial risk factors including sex hormones, trauma, parafunctions, stress and emotions [1, 2]. Women, particularly those aged between 20 and 40 years, appear to be at greater risk of TMDs [1–3]. While the prevalence of TMDs established by formalized diagnostic criteria ranged from 6.0 to 15.8%, the occurrence of TMD signs/symptoms in the general population was substantially greater (up to 75.0%) based on clinical evaluations/self-reported questionnaires [4, 5]. The high variability in TMD prevalence observed may be explained by the different instruments/protocols employed for identifying and diagnosing TMDs.

Formalized diagnostic criteria, like the Research Diagnostic and Diagnostic Criteria for TMDs (RDC/TMD and DC/TMD), generally involve comprehensive symptom history taking, protocolized physical examination, and defined algorithms for rendering specific TMD diagnosis [6, 7]. Though reliable and valid, the use of RDC/TMD and DC/TMD is not pragmatic for clinical triage and epidemiological studies involving large samples due to their complicated and time-consuming procedures. Instruments for screening the presence/absence of TMDs or 'TMD screeners' need to be cheap, easy, fast to dispense (ideally self-administered), reliable, and valid. Gonzalez et al. had reviewed the shortcomings of past TMD screeners and current ones include the TMD Pain Screener (TPS), Three Questions for TMDs (3Q/TMD), and Fonseca Anamnestic Index (FAI) [8-10]. The 6-item TPS is a component of the DC/TMD, but is specifically designed to identify painful TMDs. In contrast, the 3-item 3Q/ TMD and the 10-item FAI are capable of detecting both painrelated and intra-articular conditions. However, the diagnostic accuracy of the 3Q/TMD might be constrained due to its limited

CONTACT Dr. Carolina Marpaung a carolina@trisakti.ac.id Department of Prosthodontics, Faculty of Dentistry, Universitas Trisakti, JI Kyai Tapa no 260, Jakarta 1140, Indonesia

number of questions [8, 9]. Among the three screening instruments, the FAI is the most widely used and has been translated into numerous languages, including Chinese, Malay, Turkish, Arabic, Spanish, and Polish [11–16].

The psychometric properties of the FAI are well-reported and it was shown to yield consistent outcomes with other TMD screeners such as the American Association of Orofacial Pain Questionnaire as well as formalized TMD diagnostic criteria [11–20]. More recently, the Short-Form Fonseca Anamnestic Index (SFAI), comprising of five TMD-specific items, demonstrated high diagnostic accuracy when related to both the RDC/TMD and DC/TMD [20, 21], raising concerns over the possible over-estimation of TMD prevalence by the FAI due to its multidimensionality (measurement of more than one dimension of a construct) and inclusion of non-TMD-specific symptoms and risk factors, specifically headache, neck pain, parafunctions, malocclusion, and emotional tension [22].

The objectives of this study were to explore the dimensionality of the FAI and to validate its main component using Rasch analysis. The latter is a modern statistical technique founded on the item-response theory that is commonly employed in medicine for instrument development and refinement [23]. The Rasch model stipulates that item responses are the result of linear probabilistic interactions between the person's 'ability' and the item's 'difficulty' [24]. It transforms ordinal data into interval-scaled ones that are conveyed in 'log odds/logits' and hierarchically arranged. In addition, the Rasch model indicates fit statistics specifying how well discrete items and individual respondents describe/match the group [25]. The null hypotheses were: (1) The FAI is not multidimensional, and (2) items of its main component do not fit the Rasch model and are not valid in terms of person ability and item difficulty.

Materials and methods

Data for this study were gathered from health surveys approved by the relevant institutional review boards (reference numbers: SHS2018005 and 377-S1/KEPK/FKG/8-2020). The participants were recruited from young adults attending a polytechnic in Singapore and University in Indonesia. A minimum sample of 371 participants was established using a sample size calculator (https://www.calculator.net) founded on a 95% confidence level, 5% confidence interval, a combined student population of 34,700, and 42% mild-to-severe TMD based on the FAI [26]. Individuals must be aged 18-44 years and proficient in English or Bahasa Indonesia (BI) to qualify for the study. Those with a history of recent (prior 2 weeks) orofacial trauma/surgical procedures, debilitating psychological, and systemic conditions were excluded. Participation in the study was voluntary with no incentives offered. Informed consent was obtained before completing questionnaires that included demographic information and the English or Indonesian language versions of the FAI. The Indonesian FAI (FAI-I) was developed following the International Network for Orofacial Pain and Related Disorders Methodology (INfORM) protocol and has good reliability and validity [27].

The FAI comprised of four pain-related questions (TMJ, masticatory muscle, neck pain, and headaches), three function-related questions (TMJ sounds, opening, and side-movement difficulty), and three questions on TMD risk factors (parafunctional habits, malocclusion, and emotional tension). Participants rated the FAI items on a 3-point response scale with 'no', 'sometimes', and 'yes' being awarded 0, 5, and 10 points, respectively. Total sum scores are calculated and TMD severity is graded as follows: 'no' (0-15 points), 'mild' (20-40 points), 'moderate' (45-65 points), or 'severe' (70-100 points) [10, 17]. For the SFAI, sum scores are only totaled for items F1, F2, F3, F6, and F7. The absence and presence of TMDs are indicated by scores of $\leq 15 \text{ points}$ and $\geq 20 \text{ points}$, respectively [20, 21].

Statistical and Rasch analyses were conducted using the SPSS statistics software version 24.0 (IBM Corporation, Armonk, New York, USA) and Winsteps Version 4.3.4 (Linacre, Beaverton, Oregon, USA) [28]. Principal Component Analysis (PCA) was utilized to establish the dimensionality of the FAI. To confirm the applicability of PCA, sampling adequacy was assessed through the Kaiser-Meyer-Olkin (KMO) test, and the significance of item correlations was examined using Bartlett's test of sphericity (BTS). While KMO values ≥ 0.60 are considered satisfactory, BTS p-values of < 0.05indicate adequate item correlations and suitability for factor extraction. An eigenvalue > 1.0 and direct oblimin rotation with a threshold item loading of > 0.40 were applied for extracting the components of the FAI [29]. The main component that explained most of variance was isolated and its internal consistency (Cronbach alpha [α]) was calculated and categorized as follows: very low ($\alpha \le 0.30$); low (0.30 < $\alpha \le 0.60$); moderate (0.60 < $\alpha \le 0.75$); high (0.75 < $\alpha \le 0.90$); and very high ($\alpha > 0.90$) [30].

To validate the main component of the FAI, Rasch analysis was performed, focusing on person/item fit (the alignment of individual responses and test items with the expected measurement model), separation (the ability to distinguish between different levels of the trait being measured), reliability statistics (the consistency of the measurement across respondents and items), and point-measure correlations (the relationship between individual item scores and the overall trait being measured). Fit statistics, including infit (sensitivity to unexpected responses to items close to a respondent's ability level) and outfit (sensitivity to unexpected responses to items far from a respondent's ability level) values, were reported as mean-square (MnSq) statistics. Acceptable MnSq ranges for polytomous models (response options ≥3) were 0.6–1.4 logits, with standardized fit statistics (Zstd) within ± 1.9 [29]. Items with MnSq values outside this range (< 0.6 or > 1.4 logits) were excluded. Item difficulty was assessed using θ values, and participants' ability was analyzed using the Wright person-item map [24, 25]. Probability curves for the three response categories were plotted, and differential item functioning (DIF) was checked by country (Singapore/Indonesia) and gender (female/male) [31, 32].

Results

Out of 1219 eligible young adults screened, 318 declined study involvement ensuing in a response rate of 73.9%. The mean

Table 2. Summary of factor loadings of the items in the main component of the FAI (SFAI).

Items	Component 1
F1. Do you have difficulty opening your mouth wide?	0.72
F2. Do you have difficulty moving your jaw to the sides?	0.67
F7. Have you ever noticed any noise in your TMJ while chewing or opening your mouth?	0.66
F6. Do you have ear aches or pain in that area (temporomandibular joint)?	0.60
F3. Do you feel fatigue or muscle pain when you chew?	0.59
Eigenvalue	2.10
% of variance explained	41.94
Cronbach alpha (KR-20)	0.63

Values ≥ 0.40 were considered for relevant factor loadings after oblique rotation. FAI: Fonseca Anamnestic Index; SFAI: Short-Form Fonseca Anamnestic Index; TMJ: Temporomandibular joints.

age of the 901 participants, comprising 400 Singaporeans and 501 Indonesians, was 19.30 ± 1.48 years and 65.0% were women. Of these, 46.2% (n = 416) were FAI-negative and 53.8% (n = 485) were FAI-positive. Mild, moderate, and severe TMD was present in 43.2%, 9.9%, and 0.8% of the participants accordingly (Supplementary Table 1). Based on the SFAI, TMD was present in 10.8% of the cohort and absent in 89.2%.

Items of the FAI demonstrated adequate sampling and appropriate correlations for PCA with a KMO score 0.77 and a significant BTS (p < 0.001). Table 1 displays the outcome of PCA. The FAI was found to be multidimensional with three components explaining 50.9% of the scale variance. Items F1, F2, F3, F6, and F7, which were the items of the SFAI, formed the main component and accounted for 27.1% of the difference. While the second component consisted of items F4, F5, and F10, the third component involved items F8 and F9. The internal consistency of the main component of SFAI was moderate with a Cronbach alpha (α) of 0.63 (Table 2).

Table 3 shows the outcomes of Rasch analysis including fit index for persons/items and the degree of difficulty for the

SFAI items. Mean person (participant) ability was -1.48 ± 0.94 logits with infit and outfit MnSq values of 0.98 and 0.96 logits. Mean item difficulty was 0.00 ± 0.75 logits with infit and outfit MnSq values of 1.00 and 0.96 logits. While Rasch person separation was low (0.00), Rasch item separation was high (8.05). Similarly, Rasch person reliability was also low (0.00), but Rasch item reliability was excellent (0.98). All five items of the SFAI fitted the Rasch model. Infit MnSq values ranged from 0.87 to 1.06 logits, whereas outfit values varied from 0.82 to 1.03 logits. Infit and outfit Zstd values were within the span of ± 1.9. Point-measure correlations were all positive and fluctuated between 0.50 and 0.71. Item F3 (Do you feel fatigue or muscle pain when you chew?) was observed to be the easiest ($\theta = -0.80$) and item F2 (Do you have difficulty moving your jaw to the sides?) was deemed the hardest ($\theta = 1.25$). Table 4 indicates the differential item functioning for country and gender. Country bias was observed for item F3 and gender bias was noted for items F3 (muscle pain) and F6 (TMJ pain) (p < 0.05).

The Wright person-item map is presented in Figure 1 with person and item measures indicated on the left and right sides, respectively. Person measure is ordered based on ability (greater TMD severity on top) whilst item measure is ordered according to difficulty (more challenging on top). Person measure extended from -3.75 to 3.75 logits whereas item measure went from -0.80 to 1.25 logits. The spread of person ability surpassed that of item difficulty and was partial toward the lower end of TMD severity. The mean (M) item difficulty was higher than the mean (M) person ability suggesting that the items generally have a lower prospect of being endorsed aptly by the study sample. Figure 2 displays the probability curves for the three response categories. All response categories had distinct peaks with Rasch-Andrich thresholds ranging from -0.69 to 0.69 specifying that they were the most probable response category and applicable for defining the measured variables.

Table 1. Dimensionality of the FAI (n = 901).

ltems ^a			Component	
	_	1	2	3
F1	Do you have difficulty opening your mouth wide?	0.83		
F2	Do you have difficulty moving your jaw to the sides?	0.75		
F7	Have you ever noticed any noise in your TMJ while chewing or opening your mouth?	0.60		
F6	Do you have ear aches or pain in that area?	0.41		
F3	Do you feel fatigue or muscle pain when you chew?	0.41		
F4	Do you have frequent headaches?		0.83	
F5	Do you have neck pain or a stiff neck?		0.77	
F10	Do you consider yourself a tense (nervous) person?		0.50	
F8	Do you have any habits such as clenching or grinding your teeth?			0.76
F9	Do you feel that your teeth do not come together well?			0.73
Eigenvalues		2.71	1.37	1.01
Scale variance (%)		27.14	13.74	10.05
Total scale variance (%)			50.93	

Values in bold highlight factor loadings that correspond to the specific component most. Component loadings < 0.40 were omitted. FAI: Fonseca Anamnestic Index; TMJ: Temporomandibular joints.

Table 3. Fit index for persons/items and degree of item difficulty (logits).

Item	Measure			Misfit	order	
		Infit MnSq	Infit Zstd	Outfit MnSq	Outfit Zstd	Point-measure correlation
Summary of persons	-1.48 ± 0.94	0.98	0.09	0.96	0.11	-
Mean ± SD						
Summary of items	0.00 ± 0.75	1.00	0.12	0.96	-0.20	-
Mean ± SD						
Item statistics	(θ)					
F1	0.13	0.87	-1.92	0.93	-0.84	0.62
F2	1.25	1.00	0.00	0.82	-1.20	0.50
F3	-0.80	1.04	0.64	1.01	0.21	0.68
F6	0.18	1.06	0.87	1.03	0.35	0.58
F7	-0.76	1.06	1.02	1.03	0.49	0.71

MnSq: mean-square; Zstd: Z-standardized; θ : theta (item's difficulty index).

Discussion

The dimensionality of the FAI was ascertained and its main component was validated using Rasch analysis. As the FAI was multidimensional and the items of its main component (SFAI) conformed to the Rasch model in terms of fit, separation, and reliability statistics, both null hypotheses were rejected. This study was the first to jointly appraise two language versions of the FAI allowing for more precise estimates and generalization of results for Southeast Asian non-clinical community samples. The Indonesian FAI had suitable test-retest reliability (intraclass correlation coefficient = 0.72) and validity when related to the Oral Health Impact Profile-14 (correlation coefficient = 0.47) [33]. Young adults were selected for the study as they represent the peak incidence age for TMDs and constitute the majority of TMD patients [1, 2, 34]. Among the young adults examined, 53.8% were FAI-positive with the majority (46.2%) having mild TMD. The finding corroborated those of similar Asian studies reporting FAI-based TMD prevalence ranging from 46.8 to 53.3% [35, 36]. These prevalence rates were considerably greater than those established using formalized diagnostic criteria. Based on the SFAI, TMD was present in 10.8% of the participants which was consistent with the prevalence range (6.0-15.8%) determined with the RDC/TMD [2]. The aforementioned, reinforced concerns over the over-estimation of TMD prevalence with the FAI.

Multidimensionality of the FAI was confirmed with PCA, though a bifactor structure had also been reported [15]. The main component of the FAI consisted of the five items of the

Table 4. Differential item functioning for country and gender (n = 901).

Item	Cou	Country		nder
	X ²	Prob	X ²	Prob
F1	0.880	0.348	2.082	0.149
F2	0.289	0.590	1.842	0.174
F3	4.104	0.042*	20.101	0.000*
F6	1.395	0.237	3.893	0.048*
F7	3.627	0.056	0.768	0.380

^{*}Results of Chi-square test (p < 0.05).




Figure 1. Wright person-item map for the items of the main component of the FAI (SFAI).

Right side indicates item difficulty while left side indicates person ability. M: mean; S: 1 SD; T: 2 SD; FAI: Fonseca Anamnestic Index; SFAI: Short-Form Fonseca Anamnestic Index.

SFAI and verified the work of Rodrigues-Bigaton et al. who concluded that the main dimension of the FAI involved items F1, F2, F3, F6, and F7 [22]. These SFAI items had moderate internal consistency ($\alpha = 0.63$) and were subsequently subjected to Rasch measurements. The second component encompassed items F4 (headache), F5 (neck pain), and F10 (emotional tension). Head and neck pain often co-exist arising from trigeminocervical relationships and are associated with psychosocial factors including emotional distress [37, 38]. The last component entailed items F8 (parafunctional habits) and F9 (malocclusion) that are TMD risk factors. Sánchez-Torrelo et al. also identified three components in the FAI, though the distribution of items differed slightly [12]. While systematic reviews indicated that parafunctions such as awake/sleep bruxism could be related to TMDs, no 'clinically relevant' associations were established between dental malocclusion and TMDs in population-based and clinical studies [39-41].

Classical test methods have been applied to practically all studies appraising the psychometric properties of the FAI [11–17]. However, the classical test theory has some conceptual restrictions. These include the supposition that variance in responses is due to person ability, reliability remains constant for different person ability, and all items contribute equally to the total score [42, 43]. In contrast to the classical test approach, Rasch analysis supports the examination of fit, separation, and

reliability statistics, and hierarchical structures, and provides distribution as well as probability graphics [44]. MnSq fit statistics specify the magnitude of randomness with values < 0.6 and > 1.4 logits indicating unfavorable data predictability and unpredictability, respectively. Both infit (inlier-sensitive) and outfit (outlier-sensitive) MnSq for person ability, item difficulty, and all five items of the SFAI were within the ideal range of 0.6-1.4 logits and were productive for measurement. Furthermore, infit/outfit Zstd values were within the range of ±1.9 indicating that the data had reasonable predictability [31]. The low person separation (< 2) implies that the SFAI is not sufficiently sensitive to distinguish individuals with mild and severe TMD. This was consistent with the purpose of the SFAI which was to detect the presence or absence of TMDs. The high item separation (>3) suggests that the person sample was large enough to confirm the construct validity (item difficulty hierarchy) of the SFAI. The low person reliability observed can be attributed to the very low occurrence of moderate-to-severe TMD (7.7%) among the young adults examined. The excellent reliability of SFAI indicates that it had a large item difficulty range and a reproducible item difficulty hierarchy. Point-measure correlations were all positive denoting that all SFAI item-level scoring concurred with the latent variables.

Item F3 (muscle pain) and item F2 (side-movement difficulty) were considered the least and most difficult questions by the




Figure 2. Probability curve to show the operation of the three response categories (category 1 = no, category 2 = sometimes, and category 3 = yes).

participants. The findings are plausible considering that the mean glide distance during mastication is under 1.5 mm and participants may have problems recognizing limited lateral jaw movements [45]. As regards differential item functioning, country bias was present for item F3 and gender bias was present for items F3 and F6 (TMJ pain). The country bias for item F3 can be ascribed to the higher frequency of selfreported muscle fatigue/pain in Indonesian young adults (39.1%) compared to their Singaporean (19.6%) counterparts. Similarly, the gender bias could be attributed to the higher risk of TMDs amongst women and their susceptibility to chronic pain conditions [1, 46]. While muscle pain was reported by 35.0% of women and 22.2% of men, TMJ pain was reported by 19.1% of women and 12.7% of men. The greater pain sensitivity in women had been attributed to sociocultural, psychological, and experiential gender differences among other factors [46].

As shown in the Wright person-item map, the spread of person ability was greater than that of item difficulty and was partial to the lower end of TMD severity. This can again be ascribed to very low prevalence of moderate-to-severe TMD and the absence of TMD in the vast majority of subjects (89.2%) based on the SFAI. As the mean item difficulty was higher than mean person ability, items of the SFAI had a lower prospect of being endorsed appropriately by community samples without TMDs. This was in agreement with the greater positive predictive values (99.4–99.5%) than negative predictive values (41.7–83.3%) when the SFAI was referenced to the DC/TMD. The SFAI is thus more proficient at identifying the presence than the absence of TMDs [21].

This study, like all others, has its limitations. First, as a selfadministered instrument, the FAI is susceptible to various biases, including those related to sampling techniques, social desirability, recall periods, participants' understanding, and selective memory [47]. However, Rasch analysis helps mitigate some of these concerns by rigorously evaluating the psychometric properties of questionnaires, reducing measurement error, and ensuring that the data align with the underlying model. Second, only the English and Indonesian versions of the FAI were investigated. As dimensionality of the FAI may be dependent on instrument translation and the study population, future work should incorporate other language versions of the FAI and racial groups. Third, the study sample involved only young adults in the community. The study should be extended to older adults as well as the elderly to determine the reliability and validity of the SFAI in the general population. Lastly, it is important to note that the SFAI serves only as a TMD screener. The use of formalized diagnostic protocols and criteria, as well as adjunctive diagnostic imaging, are necessary for rendering definitive TMD diagnoses. Notwithstanding, the SFAI, with its simplicity, low cost, good validity, and reliability, is useful for identifying the presence of TMDs in clinical practice. Given the high prevalence of TMDs in the general population, clinicians should screen all patients for TMDs and document any pre-existing signs or symptoms before initiating treatment, particularly in light of the litigious nature of modern society.

Conclusion

Within the limitation of this study, the following conclusions were drawn:

- 1. The FAI was multidimensional with the main component consisting of items F1, F2, F3, F6, and F7 which were the items of the SFAI.
- The SFAI fitted the Rasch model and all its items were productive for measurement and had reasonable predictability.
- The SFAI had high item separation confirming its construct validity and excellent item reliability with a large item difficulty hierarchy.
- Item F3 (muscle pain) and item F2 (side-movement difficulty) were deemed the least and most difficult.
- 5. The SFAI can be used to screen for TMDs in clinical practice.

Acknowledgement

The authors would like to thank the School of Health and Social Sciences, Nanyang Polytechnic and the Faculty of Dentistry, Universitas Trisakti for supporting this research.

Author contribution

Yap AU contributed to conceptualization, data curation, formal analysis, investigation, methodology, project administration, resources, supervision, validation, visualization, and writing the original manuscript. Gunardi I contributed to data curation, formal analysis, methodology, resources, software, validation. and review/editing of the manuscript. Lee DZR contributed to data curation, formal analysis, methodology, resources, software, validation, and review/editing of the manuscript.

Marpaung C contributed to conceptualization, data curation, investigation, methodology, project administration, resources, supervision, and review/editing of the manuscript.

Conflict of interest

The authors have no conflict of interest to declare relating to this work.

Funding source

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

- [1] List T, Jensen RH. Temporomandibular disorders: old ideas and new concepts. Cephalalgia. 2017;37:692–704. https://doi. org/10.1177/0333102416686302
- [2] National Academies of Sciences, Engineering, and Medicine, Health and Medicine Division, Board on Health Care Services, Board on Health Sciences Policy, Committee on Temporomandibular

- Disorders (TMDs). From research discoveries to clinical treatment. In: Yost O, Liverman CT, English R, Mackey S, Bond EC, editors. Temporomandibular disorders: priorities for research and care. Washington, DC: National Academies Press (US); 2020: 24–29.
- [3] Bueno CH, Pereira DD, Pattussi MP, Grossi PK, Grossi ML. Gender differences in temporomandibular disorders in adult populational studies: a systematic review and meta-analysis. J Oral Rehabil. 2018;45:720–9. https://doi.org/10.1111/joor.12661
- [4] Manfredini D, Guarda-Nardini L, Winocur E, Piccotti F, Ahlberg J, Lobbezoo F. Research diagnostic criteria for temporomandibular disorders: a systematic review of axis I epidemiologic findings. Oral Surg Oral Med Oral Pathol Oral Radiol Endod. 2011;112:453–62. https:// doi.org/10.1016/j.tripleo.2011.04.021
- [5] Ryan J, Akhter R, Hassan N, Hilton G, Wickham J, Ibaragi S. Epidemiology of temporomandibular disorder in the general population: a systematic review. Adv Dent Oral Health. 2019;10:555787. https://doi.org/10.19080/ADOH.2019.10.555787
- [6] Dworkin SF, LeResche L. Research diagnostic criteria for temporomandibular disorders: review, criteria, examinations and specifications, critique. J Craniomandib Disord. 1992;6:301–55.
- [7] Schiffman E, Ohrbach R, Truelove E, et al. Diagnostic criteria for temporomandibular disorders (DC/TMD) for clinical and research applications: recommendations of the International RDC/TMD Consortium Network and Orofacial Pain Special Interest Group. J Oral Facial Pain Headache. 2014;28:6–27. https://doi.org/10.11607/ jop.1151
- [8] Gonzalez YM, Schiffman E, Gordon SM, et al. Development of a brief and effective temporomandibular disorder pain screening questionnaire: reliability and validity. J Am Dent Assoc. 2011;142:1183–91. https://doi.org/10.14219/jada.archive.2011.0088
- [9] Lövgren A, Visscher CM, Häggman-Henrikson B, Lobbezoo F, Marklund S, Wänman A. Validity of three screening questions (3Q/ TMD) in relation to the DC/TMD. J Oral Rehabil. 2016;43:729–36. https://doi.org/10.1111/joor.12428
- [10] Fonseca DM, Bonfante G, Valle AL, Freitas SFT. Diagnosis by anamnesis of craniomandibular dysfunction [Diagnóstico pela anamneses da disfunção craniomandibular]. Rev Gaúcha Odontol. 1994;42:23–32.
- [11] Zhang MJ, Yap AU, Lei J, Fu KY. Psychometric evaluation of the Chinese version of the Fonseca anamnestic index for temporomandibular disorders. J Oral Rehabil. 2020;47:313–18. https://doi. org/10.1111/joor.12893
- [12] Sánchez-Torrelo CM, Zagalaz-Anula N, Alonso-Royo R, et al. Transcultural adaptation and validation of the Fonseca Anamnestic Index in a Spanish population with temporomandibular disorders. J Clin Med. 2020;9:3230. https://doi.org/10.3390/jcm9103230
- [13] Alyessary AS, Yap AU, Almousawi A. The Arabic Fonseca Anamnestic Index: psychometric properties and use for screening temporomandibular disorders in prospective orthodontic patients. Cranio. 2023;41:131–8. https://doi.org/10.1080/08869634.2020.1827627
- [14] Yap AU, Apipi M, Ismail SM, Yusof ZYM, Kadir K. Psychometric properties of the Malay Fonseca Anamnestic Index for temporomandibular disorders. Cranio. 2024;42:801–8. https://doi.org/10.1080/08869634. 2022.2059130.
- [15] Arikan H, Citaker S, Ucok C. Psychometric properties of the Fonseca Anamnestic Index (FAI) for temporomandibular disorders: Turkish version, responsiveness, reliability, and validity study. Disabil Rehabil. 2024;46:1408–15. https://doi.org/10.1080/09638288.2023.2199221
- [16] Gałczyńska-Rusin M, Pobudek-Radzikowska M, Czajka-Jakubowska A. Polish language adaptation and validation of the Fonseca Anamnestic Index for individuals with temporomandibular disorders. Dent Med Probl. 2024;61(5):705–11. https://doi.org/10.17219/ dmp/177287
- [17] Topuz MF, Oghan F, Ceyhan A, Ozkan Y, Erdogan O, Musmul A, et al. Assessment of the severity of temporomandibular disorders in females: validity and reliability of the Fonseca anamnestic index. Cranio. 2023;41:84–7. https://doi.org/10.1080/08869634.20 20.1814652

- [18] Berni KC, Dibai-Filho AV, Rodrigues-Bigaton D. Accuracy of the Fonseca anamnestic index in the identification of myogenous temporomandibular disorder in female community cases. J Body Mov Ther. 2015;19:404–9, https://doi.org/10.1016/i.ibmt.2014.08.001
- [19] Pastore GP, Goulart DR, Pastore PR, Prati AJ, de Moraes M. Comparison of instruments used to select and classify patients with temporomandibular disorder. Acta Odontol Latinoam. 2018;31:16–22.
- [20] Pires PF, de Castro EM, Pelai EB, de Arruda ABC, Rodrigues-Bigaton D. Analysis of the accuracy and reliability of the Short-Form Fonseca Anamnestic Index in the diagnosis of myogenous temporomandibular disorder in women. Braz J Phys Ther. 2018;22:276–82. https://doi. org/10.1016/i.bipt.2018.02.003
- [21] Yap AU, Zhang MJ, Lei J, Fu KY. Diagnostic accuracy of the short-form Fonseca Anamnestic Index in relation to the diagnostic criteria for temporomandibular disorders. J Prosthet Dent. 2022;128:977–83. https://doi.org/10.1016/j.prosdent.2021.02.016
- [22] Rodrigues-Bigaton D, de Castro EM, Pires PF. Factor and Rasch analysis of the Fonseca anamnestic index for the diagnosis of myogenous temporomandibular disorder. Braz J Phys Ther. 2017;21:120–6. https://doi.org/10.1016/j.bjpt.2017.03.007
- [23] Belvedere SL, de Morton NA. Application of Rasch analysis in health care is increasing and is applied for variable reasons in mobility instruments. J Clin Epidemiol. 2010;63:1287–97. https://doi. org/10.1016/j.jclinepi.2010.02.012
- [24] Rasch G. Probabilistic models for some intelligence and attainment tests. Chicago, IL: Mesa Press; 1993.
- [25] Wright BD, Stone MH. Best test design: Rasch measurement. Chicago, IL: MESA Press: 1993.
- [26] Natu VP, Yap AU, Su MH, Irfan Ali NM, Ansari A. Temporomandibular disorder symptoms and their association with quality of life, emotional states and sleep quality in South-East Asian youths. J Oral Rehabil. 2018;45:756–63. https://doi.org/10.1111/joor.12692
- [27] Yap AU, Marpaung C, Gunardi I. Psychometric properties of the Indonesian Fonseca anamnestic index and the presence/severity of temporomandibular disorders among Indonesian young adults. Cranio. 2024;42:379–86. https://doi.org/10.1080/08869634.20
- [28] Linacre JM. Winsteps Rasch measurement computer program user's quide. Beaverton, OR: Winsteps.com; 2021.
- [29] Field A. Discovering statistics using SPSS. 3rd ed. London: Sage Publications: 2009.
- [30] Cunha CM, Almeida Neto OP, Stackfleth R. Main psychometric evaluation methods of measuring instrument's reliability. Rev Atenção Saúde. 2016;14:98–103. https://doi.org/10.13037/ras. vol14n47 3391
- [31] Linacre MJ. What do infit and outfit, mean-square and standardized mean? [Internet]. [cited 2024 Sept 3]. Available from: https://rasch. org/rmt/rmt162f.htm
- [32] Dabaghi S, Esmaielzadeh F, Rohani C. Application of Rasch analysis for development and psychometric properties of adolescents' quality of life instruments: a systematic review. Adolesc Health Med Ther. 2020;11:173–97. https://doi.org/10.2147/AHMT.S265413
- [33] Slade GD. Derivation and validation of a short-form oral health impact profile. Community Dent Oral Epidemiol. 1997;25:284–90. https://doi.org/10.1111/j.1600-0528.1997.tb00941.x
- [34] Yap AU, Cao Y, Zhang MJ, Lei J, Fu KY. Age-related differences in diagnostic categories, psychological states and oral health-related quality of life of adult temporomandibular disorder patients. J Oral Rehabil. 2021;48:361–8. https://doi.org/10.1111/joor.13121
- [35] Zafar MS, Fareed WM, Taymour N, Khurshid Z, Khan AH. Self-reported frequency of temporomandibular disorders among undergraduate students at Taibah University. J Taibah Univ Med Sci. 2017;12:517–22. https://doi.org/10.1016/j.jtumed.2016.12.004
- [36] Habib SR, Al Rifaiy MQ, Awan KH, Alsaif A, Alshalan A, Altokais Y. Prevalence and severity of temporomandibular disorders among university students in Riyadh. Saudi Dent J. 2015;27:125–30. https://doi.org/10.1016/j.sdentj.2014.11.009

- [37] Friedman MH, Nelson AJ. Head and neck pain review: traditional and new perspectives. J Orthop Sports Phys Ther. 1996;24:268–78. https://doi.org/10.2519/jospt.1996.24.4.268
- [38] Hasvold T, Johnsen R, Førde OH. Non-migrainous headache, neck or shoulder pain, and migraine differences in association with background factors in a city population. Scand J Prim Health Care. 1996;14:92–9. https://doi.org/10.3109/02813439608997077
- [39] Jiménez-Silva A, Peña-Durán C, Tobar-Reyes J, Frugone-Zambra R. Sleep and awake bruxism in adults and its relationship with temporomandibular disorders: a systematic review from 2003 to 2014. Acta Odontol Scand. 2017;75:36–58. https://doi.org/10.1080/00016 357.2016.1247465
- [40] de Oliveira Reis L, Ribeiro RA, Martins CC, Devito KL. Association between bruxism and temporomandibular disorders in children: a systematic review and meta-analysis. Int J Paediatr Dent. 2019;29:585–59. https://doi.org/10.1111/jpd.12496
- [41] Manfredini D, Lombardo L, Siciliani G. Temporomandibular disorders and dental occlusion. A systematic review of association studies: end of an era? J Oral Rehabil. 2017;44:908–23. https://doi.org/10.1111/ joor.12531

- [42] Lozano Rojas OM, Rojas Tejada AJ, Pérez Meléndez C. Development of a specific health-related quality of life test in drug abusers using the Rasch rating scale model. Eur Addict Res. 2009;15:63–70. https:// doi.org/10.1159/000189784
- [43] Wong HM, McGrath CP, King NM. Rasch validation of the early child-hood oral health impact scale. Community Dent Oral Epidemiol. 2011;39:449–57. https://doi.org/10.1111/j.1600-0528.2011.00614.x
- [44] Prieto L, Alonso J, Lamarca R. Classical test theory versus Rasch analysis for quality of life questionnaire reduction. Health Qual Life Outcomes. 2003;1:27. https://doi.org/10.1186/1477-7525-1-27
- [45] Rilo B, Fernández-Formoso N, Mora MJ, Cadarso-Suárez C, Santana U. Distance of the contact glide in the closing masticatory stroke during mastication of three types of food. J Oral Rehabil. 2009;36:571–6. https://doi.org/10.1111/j.1365-2842.2009.01956.x
- [46] Mogil JS, Bailey AL. Sex and gender differences in pain and analgesia. Prog Brain Res. 2010;186:141–57. https://doi.org/10.1016/ B978-0-444-53630-3.00009-9
- [47] Althubaiti A. Information bias in health research: definition, pitfalls, and adjustment methods. J Multidiscip Healthc. 2016;9:211–17. https://doi.org/10.2147/JMDH.S104807

Dimensionality of the Fonseca Anamnestic Index and validation of its short-form derivative

by Carolina Damayanti Marpaung

Submission date: 01-Aug-2025 04:34PM (UTC+0700)

Submission ID: 2579109261

File name: ACTA_ODONT_Dimensionality_of_the_Fonseca_drg_Maya.pdf (275.57K)

Word count: 6412 Character count: 33479





ORIGINAL ARTICLE

Dimensionality of the Fonseca Anamnestic Index and validation of its short-form derivative

Adrian Ujin Yapab,cde. 60, Indrayadi Gunardi 60, Darren Zong Ru Lee 60 and Carolina Marpaung 60

"Division of Dentistry, Ng Teng Fong General Hospital, Singapore, Singapore; "Faculty of Dentistry, National University Health System, Singapore, Singapore; "National Dental Research Institute, Singapore, Singapore; "National Dental Center, Singapore, Singapore; "Duke-NUS Medical School, Singapore Health Services, Singapore; Singapore; 'Department of Prosthodontics, Faculty of Dentistry, Universitas Trisakti, Indonesia; "Department of Oral Medicine, Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia; "School of Health and Social Sciences, Nanyang Polytechnic, Singapore, Singapore

ABSTRACT

Objectives: Recently, the Short-Form Fonseca Anamnestic Index (SFAI) was shown to have high diagnostic accuracy when compared to the Diagnostic Criteria for Temporomandibular Disorders (DC/TMD) in patient samples. This study investigated the dimensionality of the parent instrument (Fonammestic Index [FAII) and validated its main component using Rasch analysis in non-patient populations.

Methods: FAI data from a total of 901 participants from Singapore and Indonesia with a mean age 19.30 ± 1.48 years (65.0% women) were examined. Of these, 53.8% were FAI positive and 46.2% were FAI negative.

Methods: FAI data from a total of 901 participants from Singapore and Indonesia with a mean age 19.30 ± 1.48 years (65.0% women) were examined. Of these, 53.8% were FAI positive and 46.2% were FAI negative. Principal Component Analysis (PCA) was performed to extract the main component of the FAI using an eigenvalue > 1.0 and direct oblimin rotation with item loading of > 0.40. Rasch analysis was subsequently carried out on the items of the main component.

Results: The FAI was found to be multidimensional with the main component involving items F1, F2, F3, F6, and F7 which were the items of the SFAI. The SFAI had moderate internal consistency (Cronbach alpha = 0.63) and fitted the Rasch model with person and item infit/outfit mean square (MnSq) values of 0.98/0.96 and 1.00/0.96 logits respectively. The infit/outfit MnSq of the SFAI items ranged from 0.82 to 1.06 logits with Item F2 (side-movement difficulty) being the most difficult and item F3 (muscle pain) the easiest.

Conclusions: The FAI is multidimensional with the main component comprising the five items of the SFAI that fitted the Rasch model. With its good Rasch validity, separation, and reliability, the SFAI is a promising tool for TMD screening.

ARTICI F HISTORY

Received 3 September 2024 Accepted 10 January 2025

KEYWORDS Temporomandibular disorders; general population; questionnaires; validity; methods

Introduction

Temporomandibular disorders (TMDs) are a diverse group of neuromuscular and musculoskeletal conditions affecting the temporomandibular joints (TMJs), muscles of masticatory, and adjoining structures. In addition to TMJ and masticatory muscle pain, the other signs/symptoms of TMDs include headaches, neck pain, TMJ noises, jaw functional limitations, as well as abnormal jaw movements [1, 2]. The complex etiology of TMDs involves a myriad of biopsychosocial risk factors including sex hormones, trauma, parafunctions, stress and emotions [1, 2]. Women, particularly those aged between 20 and 40 years, appear to be at greater risk of TMDs [1-3]. While the prevalence of TMDs established by formalized diagnostic criteria ranged from 6.0 to 15.8%, the occurrence of TMD signs/symptoms in the general population was substantially greater (up to 75.0%) based on clinical evaluations/self-reported questionnaires [4, 5]. The high variability in TMD prevalence observed may be explained by the different instruments/protocols employed for identifying and diagnosing TMDs.

Formalized diagnostic criteria, like the Research Diagnostic and Diagnostic Criteria for TMDs (RDC/TMD and DC/TMD), generally involve comprehensive symptom history taking, protocolized physical examination, and defined algorithms for rendering specific TMD diagnosis [6, 7]. Though reliable and valid, the use of RDC/TMD and DC/TMD is not pragmatic for clinical triage and epidemiological studies involving large samples due to their complicated and time-consuming procedures. Instruments for screening the presence/absence of TMDs or 'TMD screeners' need to be cheap, easy, fast to dispense (ideally self-administered), reliable, and valid. Gonzalez et al. had reviewed the shortcomings of past TMD screeners and current ones include the TMD Pain Screener (TPS), Three Questions for TMDs (3Q/TMD), and Fonseca Anamnestic Index (FAI) [8-10]. The 6-item TPS is a component of the DC/TMD, but is specifically designed to identify painful TMDs. In contrast, the 3-item 3Q/ TMD and the 10-item FAI are capable of detecting both painrelated and intra-articular conditions. However, the diagnostic accuracy of the 3Q/TMD might be constrained due to its limited

CONTACT Dr. Carolina Marpaung a carolina @trisakti.acid Department of Prosthodontics, Faculty of Dentistry, Universitas Trisakti, JI Kyai Tapa no 260, Jakarta 1140, Indonesia

Indonesia

Supplemental data for this article can be accessed online at https://doi.org/10.2340/aos.v84.42960

2005 The Author's) Published by MIS Dublished by Behalf of Acto Odostelopics Specification Specific Title is

2 0225 The Author(s). Published by MJS Publishing on behalf of Acta Odontological Scandinavica Society. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/40/), allowing third parties to copy and redistribute the material in any medium or format and to remix, transform, and build upon the material, with the condition of proper attribution to the original work.

number of questions [8, 9]. Among the three screening instruments, the FAI is the most widely used and has been translated into numerous languages, including Chinese, Malay, Turkish, Arabic, Spanish, and Polish [11–16].

The psychometric properties of the FAI are well-reported and it was shown to yield consistent outcomes with other TMD screeners such as the American Association of Orofacial Pain Questionnaire as well as formalized TMD diagnostic criteria [11–20]. More recently, the Short-Form Fonseca Anamnestic Index (SFAI), comprising of five TMD-specific items, demonstrated high diagnostic accuracy when related to both the RDC/TMD and DC/TMD [20, 21], raising concerns over the possible over-estimation of TMD prevalence by the FAI due to its multidimensionality (measurement of more than one dimension of a construct) and inclusion of non-TMD-specific symptoms and risk factors, specifically headache, neck pain, parafunctions, malocclusion, and emotional tension [22].

The objectives of this study were to explore the dimensionality of the FAI and to validate its main component using Rasch analysis. The latter is a modern statistical technique founded on the item-response theory that is commonly employed in medicine for instrument development and refinement [23]. The Rasch model stipulates that item responses are the result of linear probabilistic interactions between the persons' ability' and the item's 'difficulty' [24]. It transforms ordinal data into interval-scaled ones that are conveyed in 'log odds'/logits' and hierarchically arranged. In addition, the Rasch model indicates fit statistics specifying how well discrete items and individual respondents describe/match the group [25]. The null hypotheses were: (1) The FAI is not multidimensional, and (2) items of its main component do not fit the Rasch model and are not valid in terms of person ability and item difficulty.

Materials and methods

Data for this study were gathered from health surveys approved by the relevant institutional review boards (reference numbers: SHS2018005 and 377-S1/KEPK/FKG/8-2020). The participants were recruited from young adults attending a polytechnic in Singapore and University in Indonesia. A minimum sample of 371 participants was established using a sample size calculator (https://www.calculator.net) founded on a 95% confidence level, 5% confidence interval, a combined student population of 34,700, and 42% mild-to-severe TMD based on the FAI [26]. Individuals must be aged 18-44 years and proficient in English or Bahasa Indonesia (BI) to qualify for the study. Those with a history of recent (prior 2 weeks) orofacial trauma/surgical procedures, debilitating psychological, and systemic conditions were excluded. Participation in the study was voluntary with no incentives offered. Informed consent was obtained before completing questionnaires that included demographic information and the English or Indonesian language versions of the FAI. The Indonesian FAI (FAI-I) was developed following the International Network for Orofacial Pain and Related Disorders Methodology (INfORM) protocol and has good reliability and validity [27].

The FAI comprised of four pain-related questions (TMJ, masticatory muscle, neck pain, and headaches), three function-related questions (TMJ sounds, opening, and side-movement difficulty), and three questions on TMD risk factors (parafunctional habits, malocclusion, and emotional tension). Participants rated the FAI items on a 3-point response scale with 'no', 'sometimes', and 'yes' being awarded 0, 5, and 10 points, respectively. Total sum scores are calculated and TMD severity is graded as follows: 'no' (0–15 points), 'mild' (20–40 points), 'moderate' (45–65 points), or 'severe' (70–100 points) [10, 17]. For the SFAI, sum scores are only totaled for items F1, F2, F3, F6, and F7. The absence and presence of TMDs are indicated by scores of ≤15 points and ≥ 20 points, respectively [20, 21].

Statistical and Rasch analyses were conducted using the SPSS statistics software version 24.0 (IBM Corporation, Armonk, New York, USA) and Winsteps Version 4.3.4 (Linacre, Beaverton, Oregon, USA) [28]. Principal Component Analysis (PCA) was utilized to establish the dimensionality of the FAI. To confirm the applicability of PCA, sampling adequacy was assessed through the Kaiser-Meyer-Olkin (KMO) test, and the significance of item correlations was examined using Bartlett's test of sphericity (BTS). While KMO values \geq 0.60 are considered satisfactory, BTS p-values of < 0.05 indicate adequate item correlations and suitability for factor extraction. An eigenvalue > 1.0 and direct oblimin rotation with a threshold item loading of > 0.40 were applied for extracting the components of the FAI [29]. The main component that explained most of variance was isolated and its internal consistency (Cronbach alpha $[\alpha]$) was calculated and categorized as follows: very low ($\alpha \le 0.30$); low (0.30 < $\alpha \le 0.60$); moderate (0.60 < $\alpha \le 0.75$); high (0.75 < $\alpha \le 0.90$); and very high ($\alpha > 0.90$) [30].

To validate the main component of the FAI, Rasch analysis was performed, focusing on person/item fit (the alignment of individual responses and test items with the expected measurement model), separation (the ability to distinguish between different levels of the trait being measured), reliability statistics (the consistency of the measurement across respondents and items), and point-measure correlations (the relationship between individual item scores and the overall trait being measured). Fit statistics, including infit (sensitivity to unexpected responses to items close to a respondent's ability level) and outfit (sensitivity to unexpected responses to items far from a respondent's ability level) values, were reported as mean-square (MnSq) statistics. Acceptable MnSq ranges for polytomous models (response options \geq 3) were 0.6-1.4 logits, with standardized fit statistics (Zstd) within ± 1.9 [29]. Items with MnSq values outside this range (<0.6 or > 1.4 logits) were excluded. Item difficulty was assessed using θ values, and participants' ability was analyzed using the Wright person-item map [24, 25]. Probability curves for the three response categories were plotted, and differential item functioning (DIF) was checked by country (Singapore/Indonesia) and gender (female/male) [31, 32].

Results

Out of 1219 eligible young adults screened, 318 declined study involvement ensuing in a response rate of 73.9%. The mean

Table 2. Summary of factor loadings of the items in the main component of

Items	Component
F1. Do you have difficulty opening your mouth wide?	0.72
F2. Do you have difficulty moving your jaw to the sides?	0.67
F7. Have you ever noticed any noise in your TMJ while	0.66
chewing or opening your mouth?	
F6. Do you have ear aches or pain in that area (temporomandibular joint)?	0.60
F3. Do you feel fatigue or muscle pain when you chew?	0.59
Eigenvalue	2.10
% of variance explained	41.94
Cronbach alpha (KR-20)	0.63

Values ≥ 0.40 were considered for relevant factor loadings after oblique rotation. FAI: Fonseca Anamnestic Index; SFAI: Short-Form Fonseca Anamnestic Index; TMJ: Temporomandibular joints.

age of the 901 participants, comprising 400 Singaporeans and 501 Indonesians, was 19.30 \pm 1.48 years and 65.0% were women. Of these, 46.2% (n = 416) were FAI-negative and 53.8% (n = 485) were FAI-positive. Mild, moderate, and severe TMD was present in 43.2%, 9.9%, and 0.8% of the participants accordingly (Supplementary Table 1). Based on the SFAI, TMD was present in 10.8% of the cohort and absent in 89.2%.

Items of the FAI demonstrated adequate sampling and appropriate correlations for PCA with a KMO score 0.77 and a significant BTS (p < 0.001). Table 1 displays the outcome of PCA. The FAI was found to be multidimensional with three components explaining 50.9% of the scale variance. Items F1, F2, F3, F6, and F7, which were the items of the SFAI, formed the main component and accounted for 27.1% of the difference. While the second component consisted of items F4, F5, and F10, the third component involved items F8 and F9. The internal consistency of the main component of SFAI was moderate with a Cronbach alpha (a) of 0.63 (Table 2).

Table 3 shows the outcomes of Rasch analysis including fit index for persons/items and the degree of difficulty for the

SFAI items. Mean person (participant) ability was -1.48 ± 0.94 logits with infit and outfit MnSq values of 0.98 and 0.96 logits. Mean item difficulty was 0.00 ± 0.75 logits with infit and outfit MnSq values of 1.00 and 0.96 logits. While Rasch person separation was low (0.00), Rasch item separation was high (8.05). Similarly, Rasch person reliability was also low (0.00), but Rasch item reliability was excellent (0.98). All five items of the SFAI fitted the Rasch model. Infit MnSq values ranged from 0.87 to 1.06 logits, whereas outfit values varied from 0.82 to 1.03 logits. Infit and outfit Zstd values were within the span of \pm 1.9. Point-measure correlations were all positive and fluctuated between 0.50 and 0.71. Item F3 (Do you feel fatigue or muscle pain when you chew?) was observed to be the easiest ($\theta = -0.80$) and item F2 (Do you have difficulty moving your jaw to the sides?) was deemed the hardest ($\theta = 1.25$). Table 4 indicates the differential item functioning for country and gender. Country bias was observed for item F3 and gender bias was noted for items F3 (muscle pain) and F6 (TMJ pain) (p < 0.05).

The Wright person-item map is presented in Figure 1 with person and item measures indicated on the left and right sides, respectively. Person measure is ordered based on ability (greater TMD severity on top) whilst item measure is ordered according to difficulty (more challenging on top). Person measure extended from -3.75 to 3.75 logits whereas item measure went from -0.80 to 1.25 logits. The spread of person ability surpassed that of item difficulty and was partial toward the lower end of TMD severity. The mean (M) item difficulty was higher than the mean (M) person ability suggesting that the items generally have a lower prospect of being endorsed aptly by the study sample. Figure 2 displays the probability curves for the three response categories. All response categories had distinct peaks with Rasch-Andrich thresholds ranging from -0.69 to 0.69 specifying that they were the most probable response category and applicable for defining the measured variables.

Table 1. Dimensionality of the FAI (n = 901).

Items ^a			Component	
		1	2	3
F1	Do you have difficulty opening your mouth wide?	0.83		
F2	Do you have difficulty moving your jaw to the sides?	0.75		
F7	Have you ever noticed any noise in your TMJ while chewing or opening your mouth?	0.60		
F6	Do you have ear aches or pain in that area?	0.41		
F3	Do you feel fatigue or muscle pain when you chew?	0.41		
F4	Do you have frequent headaches?		0.83	
F5	Do you have neck pain or a stiff neck?		0.77	
F10	Do you consider yourself a tense (nervous) person?		0.50	
F8	Do you have any habits such as clenching or grinding your teeth?			0.76
F9	Do you feel that your teeth do not come together well?			0.73
Eigenvalues		2.71	1.37	1.01
Scale variance (%)		27.14	13.74	10.05
Total scale variance (%)			50.93	

Values in bold highlight factor loadings that correspond to the specific component most. Component loadings < 0.40 were omitted. FAI: Fonseca Anamnestic Index; TMJ: Temporomandibular joints.

Table 3. Fit index for persons/items and degree of item difficulty (logits).

Item	measure	Measure Misfit order				
		Infit MnSa	Infit Zstd	Outfit MnSa	Outfit Zstd	Point-measure correlation
Summary of persons	-1.48 ± 0.94	0.98	0.09	0.96	0.11	-
Mean ± SD						
Summary of items	0.00 ± 0.75	1.00	0.12	0.96	-0.20	-
Mean ± SD						
Item statistics	(θ)					
F1	0.13	0.87	-1.92	0.93	-0.84	0.62
F2	1.25	1.00	0.00	0.82	-1.20	0.50
F3	-0.80	1.04	0.64	1.01	0.21	0.68
F6	0.18	1.06	0.87	1.03	0.35	0.58
F7	-0.76	1.06	1.02	1.03	0.49	0.71

MnSq: mean-square; Zstd: Z-standardized; θ: theta (item's difficulty index).

Discussion

The dimensionality of the FAI was ascertained and its main component was validated using Rasch analysis. As the FAI was multidimensional and the items of its main component (SFAI) conformed to the Rasch model in terms of fit, separation, and reliability statistics, both null hypotheses were rejected. This study was the first to jointly appraise two language versions of the FAI allowing for more precise estimates and generalization of results for Southeast Asian non-clinical community samples. The Indonesian FAI had suitable test-retest reliability (intraclass correlation coefficient = 0.72) and validity when related to the Oral Health Impact Profile-14 (correlation coefficient = 0.47) [33]. Young adults were selected for the study as they represent the peak incidence age for TMDs and constitute the majority of TMD patients [1, 2, 34]. Among the young adults examined, 53.8% were FAI-positive with the majority (46.2%) having mild TMD. The finding corroborated those of similar Asian studies reporting FAI-based TMD prevalence ranging from 46.8 to 53.3% [35, 36]. These prevalence rates were considerably greater than those established using formalized diagnostic criteria. Based on the SFAI, TMD was present in 10.8% of the participants which was consistent with the prevalence range (6.0-15.8%) determined with the RDC/TMD [2]. The aforementioned, reinforced concerns over the over-esti $mation\ of\ TMD\ prevalence\ with\ the\ FAI.$

Multidimensionality of the FAI was confirmed with PCA, though a bifactor structure had also been reported [15]. The main component of the FAI consisted of the five items of the

Table 4. Differential item functioning for country and gender (n = 901).

Cor	intry	Ger	nder
X ²	Prob	X ²	Prob
0.880	0.348	2.082	0.149
0.289	0.590	1.842	0.174
4.104	0.042*	20.101	0.000*
1.395	0.237	3.893	0.048*
3.627	0.056	0.768	0.380
	X ² 0.880 0.289 4.104 1.395	0.880 0.348 0.289 0.590 4.104 0.042* 1.395 0.237	X² Prob X² 0.880 0.348 2.082 0.289 0.590 1.842 4.104 0.042* 20.101 1.395 0.237 3.893

^{*}Results of Chi-square test (p < 0.05).




Figure 1. Wright person-item map for the items of the main component of the FAI (SFAI).

Right side indicates item difficulty while left side indicates person ability. M: mean; S: 1 SD; T: 2 SD; FAI: Fonseca Anamnestic Index; SFAI: Short-Form Fonseca Anamnestic Index.

concluded that the main dimension of the FAI involved items F1, F2, F3, F6, and F7 [22]. These SFAI items had moderate internal consistency ($\alpha = 0.63$) and were subsequently subjected to Rasch measurements. The second component encompassed $items\,F4\,(headache), F5\,(neck\,pain), and\,F10\,(emotional\,tension).$ Head and neck pain often co-exist arising from trigeminocervical relationships and are associated with psychosocial factors including emotional distress [37, 38]. The last component entailed items F8 (parafunctional habits) and F9 (malocclusion) that are TMD risk factors. Sánchez-Torrelo et al. also identified three components in the FAI, though the distribution of items differed slightly [12]. While systematic reviews indicated that parafunctions such as awake/sleep bruxism could be related to TMDs, no 'clinically relevant' associations were established between dental malocclusion and TMDs in population-based and clinical studies [39-41].

Classical test methods have been applied to practically all studies appraising the psychometric properties of the FAI [11–17]. However, the classical test theory has some conceptual restrictions. These include the supposition that variance in responses is due to person ability, reliability remains constant for different person ability, and all items contribute equally to the total score [42, 43]. In contrast to the classical test approach, Rasch analysis supports the examination of fit, separation, and

SFAI and verified the work of Rodrigues-Bigaton et al. who reliability statistics, and hierarchical structures, and provides distribution as well as probability graphics [44]. MnSq fit statistics specify the magnitude of randomness with values $< 0.6\,$ and > 1.4 logits indicating unfavorable data predictability and unpredictability, respectively. Both infit (inlier-sensitive) and outfit (outlier-sensitive) MnSq for person ability, item difficulty, and all five items of the SFAI were within the ideal range of 0.6-1.4 logits and were productive for measurement. Furthermore, infit/outfit Zstd values were within the range of ±1.9 indicating that the data had reasonable predictability [31]. The low person separation (<2) implies that the SFAI is not sufficiently sensitive to distinguish individuals with mild and severe TMD. This was consistent with the purpose of the SFAI which was to detect the presence or absence of TMDs. The high item separation (>3) suggests that the person sample was large enough to confirm the construct validity (item difficulty hierarchy) of the SFAI. The low person reliability observed can be attributed to the very low occurrence of moderate-to-severe TMD (7.7%) among the young adults examined. The excellent reliability of SFAI indicates that it had a large item difficulty range and a reproducible item difficulty hierarchy. Point-measure correlations were all positive denoting that all SFAI item-level scoring concurred with the latent variables.

> Item F3 (muscle pain) and item F2 (side-movement difficulty) were considered the least and most difficult questions by the




Figure 2. Probability curve to show the operation of the three response categories (category 1 = no, category 2 = sometimes, and category 3 = yes).

participants. The findings are plausible considering that the mean glide distance during mastication is under 1.5 mm and participants may have problems recognizing limited lateral jaw movements [45]. As regards differential item functioning, country bias was present for item F3 and gender bias was present for items F3 and F6 (TMJ pain). The country bias for item F3 can be ascribed to the higher frequency of selfreported muscle fatigue/pain in Indonesian young adults (39.1%) compared to their Singaporean (19.6%) counterparts. Similarly, the gender bias could be attributed to the higher risk of TMDs amongst women and their susceptibility to chronic pain conditions [1, 46]. While muscle pain was reported by 35.0% of women and 22.2% of men, TMJ pain was reported by 19.1% of women and 12.7% of men. The greater $pain sensitivity in women \, had \, been \, attributed \, to \, sociocultural, \,$ psychological, and experiential gender differences among other factors [46].

As shown in the Wright person-item map, the spread of person ability was greater than that of item difficulty and was partial to the lower end of TMD severity. This can again be ascribed to very low prevalence of moderate-to-severe TMD and the absence of TMD in the vast majority of subjects (89.2%) based on the SFAL As the mean item difficulty was higher than mean person ability, items of the SFAl had a lower prospect of being endorsed appropriately by community samples without TMDs. This was in agreement with the greater positive predictive values (99.4–99.5%) than negative predictive values (41.7–83.3%) when the SFAl was referenced to the DC/TMD. The SFAl is thus more proficient at identifying the presence than the absence of TMDs [21].

This study, like all others, has its limitations. First, as a selfadministered instrument, the FAI is susceptible to various biases, including those related to sampling techniques, social desirability, recall periods, participants' understanding, and selective memory [47]. However, Rasch analysis helps mitigate some of these concerns by rigorously evaluating the psychometric properties of questionnaires, reducing measurement error, and ensuring that the data align with the underlying model. Second, only the English and Indonesian versions of the FAI were investigated. As dimensionality of the FAI may be dependent on instrument translation and the study population, future work should incorporate other language versions of the FAI and racial groups. Third, the study sample involved only young adults in the community. The study should be extended to older adults as well as the elderly to determine the reliability and validity of the SFAI in the general population. Lastly, it is important to note that the SFAI serves only as a TMD screener. The use of formalized diagnostic protocols and criteria, as well as adjunctive diagnostic imaging, are necessary for rendering definitive TMD diagnoses. Notwithstanding, the SFAI, with its simplicity, low cost, good validity, and reliability, is useful for identifying the presence of TMDs in clinical practice. Given the high prevalence of TMDs in the general population, clinicians should screen all patients for TMDs and document any pre-existing signs or symptoms before initiating treatment, particularly in light of the litigious nature of

Conclusion

Within the limitation of this study, the following conclusions were drawn:

- The FAI was multidimensional with the main component consisting of items F1, F2, F3, F6, and F7 which were the items of the SFAI.
- The SFAI fitted the Rasch model and all its items were productive for measurement and had reasonable predictability.
- The SFAI had high item separation confirming its construct validity and excellent item reliability with a large item difficulty hierarchy.
- Item F3 (muscle pain) and item F2 (side-movement difficulty) were deemed the least and most difficult.
- 5. The SFAI can be used to screen for TMDs in clinical practice.

Acknowledgement

The authors would like to thank the School of Health and Social Sciences, Nanyang Polytechnic and the Faculty of Dentistry, Universitas Trisakti for supporting this research.

Author contribution

Yap AU contributed to conceptualization, data curation, formal analysis, investigation, methodology, project administration, resources, supervision, validation, visualization, and writing the original manuscript. Gunardi I contributed to data curation, formal analysis, methodology, resources, software, validation. and review/editing of the manuscript. Lee DZR contributed to data curation, formal analysis, methodology, resources, software, validation, and review/editing of the manuscript.

Marpaung C contributed to conceptualization, data curation, investigation, methodology, project administration, resources, supervision, and review/editing of the manuscript.

Conflict of interest

The authors have no conflict of interest to declare relating to this work

Funding source

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

- List T, Jensen RH. Temporomandibular disorders: old ideas and new concepts. Cephalalgia. 2017;37:692–704. https://doi. org/10.1177/0333102416686302
- 2] National Academies of Sciences, Engineering, and Medicine, Health and Medicine Division, Board on Health Care Services, Board on Health Sciences Policy, Committee on Temporomandibular

- Disorders (TMDs). From research discoveries to clinical treatment. In: Yost O, Liverman CT, English R, Mackey S, Bond EC, editors. Temporomandibular disorders: priorities for research and care. Washington, D.C. National Academies Press (US); 2020: 24-29.
- [3] Bueno CH, Pereira DD, Pattussi MP, Grossi PK, Grossi ML. Gender differences in temporomandibular disorders in adult populational studies: a systematic review and meta-analysis. J Oral Rehabil. 2018;45:720–9. https://doi.org/10.1111/joor.12661
- [4] Manfredini D, Guarda-Nardini L, Winocur E, Piccotti F, Ahlberg J, Lobbezoo F. Research diagnostic criteria for temporomandibular disorders: a systematic review of axis I epidemiologic findings. Oral Surg Oral Med Oral Pathol Oral Radiol Endod. 2011;112:453–62. https:// doi.org/10.1016/i.triplep.2011.04.021
- [5] Ryan J, Akhter R, Hassan N, Hilton G, Wickham J, Ibaragi S. Epidemiology of temporomandibular disorder in the general population: a systematic review. Adv Dent Oral Health. 2019;10:555787. https://doi.org/10.19080/ADOH.2019.10.555787
- [6] Dworkin SF, LeResche L. Research diagnostic criteria for temporomandibular disorders: review, criteria, examinations and specifications, critique. J Craniomandib Disord. 1992;6:301–55.
- [7] Schiffman E, Ohrbach R, Truelove E, et al. Diagnostic criteria for temporomandibular disorders (DC/TMD) for clinical and research applications: recommendations of the International RDC/TMD Consortium Network and Orofacial Pain Special Interest Group, J Oral Facial Pain Headache. 2014;28:6–27. https://doi.org/10.11607/ jop.1151
- [8] Gonzalez YM, Schiffman E, Gordon SM, et al. Development of a brief and effective temporomandibular disorder pain screening questionnaire: reliability and validity. J Am Dent Assoc. 2011;142:1183–91. https://doi.org/10.14219/jada.archive.2011.0088
- [9] Lövgren A, Visscher CM, Häggman-Henrikson B, Lobbezoo F, Marklund S, Wänman A. Validity of three screening questions (3Q/ TMD) in relation to the DC/TMD. J Oral Rehabil. 2016;43:729–36. https://doi.org/10.1111/joor.12428
- [10] Fonseca DM, Bonfante G, Valle AL, Freitas SFT. Diagnosis by anamnesis of craniomandibular dysfunction [Diagnóstico pela anamneses da disfunção craniomandibular]. Rev Gaúcha Odontol. 1994;42:23–32.
- [11] Zhang MJ, Yap AU, Lei J, Fu KY. Psychometric evaluation of the Chinese version of the Fonseca anamnestic index for temporomandibular disorders. J Oral Rehabil. 2020;47:313–18. https://doi. org/10.1111/joor.12893
- Sánchez-Torrelo CM, Zagalaz-Anula N, Alonso-Royo R, et al. Transcultural adaptation and validation of the Fonseca Anamnestic Index in a Spanish population with temporomandibular disorders. J Clin Med. 2020;9:3230. https://doi.org/10.3390/jcmp103230
 Alyessary AS, Yap AU, Almousawi A. The Arabic Fonseca Anamnestic
- [13] Alyessary AS, Yap AU, Almousawi A. The Arabic Fonseca Anamnestic Index: psychometric properties and use for screening temporomandibular disorders in prospective orthodontic patients. Cranio. 2023;41:121.8 https://doi.org/10.1090/0956624.2020.19275677
- 2023;41:131-8. https://doi.org/10.1080/08869634.2020.1827627
 [14] Yap AU, Apipi M, Ismail SM, Yusof ZYM, Kadir K. Psychometric properties of the Malay Fonseca Anamnestic Index for temporomandibular disorders. Cranio. 2024;42:801-8. https://doi.org/10.1080/08869634. 2022.2059130.
- [15] Arikan H, Citaker S, Ucok C, Psychometric properties of the Fonseca Anamestic Index (FAI) for temporomandibular disorders: Turkish version, responsiveness, reliability, and validity study. Disabil Rehabil. 2024;46:1408–15. https://doi.org/10.1080/09638288.2023.2199221
- [16] Galczyńska-Rusin M, Pobudek-Radzikowska M, Czajka-Jakubowska A. Polish language adaptation and validation of the Fonseca Anamnestic Index for individuals with temporomandibular disorders. Dent Med Probl. 2024;61(5):705–11. https://doi.org/10.17219/ dmp/177287
- [17] Topuz MF, Oghan F, Ceyhan A, Ozkan Y, Erdogan O, Musmul A, et al. Assessment of the severity of temporomandibular disorders in females: validity and reliability of the Fonseca anamnestic index. Cranio. 2023;41:84–7. https://doi.org/10.1080/08869634.20 20.1814652


- [18] Berni KC, Dibai-Filho AV, Rodrigues-Bigaton D. Accuracy of the Fonseca anamnestic index in the identification of myogenous temporomandibular disorder in female community cases. J Body Mov Ther. 2015;19:404–9. https://doi.org/10.1016/j.jbmt.2014.08.001
- [19] Pastore GP, Goulart DR, Pastore PR, Prati AJ, de Moraes M. Comparison of instruments used to select and classify patients with temporomandibular disorder. Acta Odontol Latingam, 2018:31:16–22.
- 20] Pires PF, de Castro EM, Pelai EB, de Arruda ABC, Rodrigues-Bigaton D. Analysis of the accuracy and reliability of the Short-Form Fonseca Anamnestic Index in the diagnosis of myogenous temporomandibular disorder in women. Braz J Phys Ther. 2018;22:276–82. https://doi. org/10.1016/j.bipt.2018.02.003
- Yap AU, Zhang MJ, Lei J, Fu KY. Diagnostic accuracy of the short-form Fonseca Anamnestic Index in relation to the diagnostic criteria for temporomandibular disorders. J Prosthet Dent. 2022;128:977–83. https://doi.org/10.1016/j.prosdept.2021.02.016
- [22] Rodrigues-Bigaton D, de Castro EM, Pires PF. Factor and Rasch analysis of the Fonseca anamnestic index for the diagnosis of myogenous temporomandibular disorder. Braz J Phys Ther. 2017;21:120–6. https://doi.org/10.1016/j.bjpt.2017.03.007
- 23] Belvedere SL, de Morton NA. Application of Rasch analysis in health care is increasing and is applied for variable reasons in mobility instruments. J Clin Epidemiol. 2010;63:1287–97. https://doi. org/10.1016/j.jclinepi.2010.02.012
- [24] Rasch G. Probabilistic models for some intelligence and attainment tests. Chicago, IL: Mesa Press; 1993.
- [25] Wright BD, Stone MH. Best test design: Rasch measurement. Chicago, IL: MESA Press; 1993.
- [26] Natu VP, Yap AU, Su MH, Irfan Ali NM, Ansari A. Temporomandibular disorder symptoms and their association with quality of life, emotional states and sleep quality in South-East Asian youths. J Oral Rehabil. 2018;45:756–63. https://doi.org/10.1111/joor.12692
- [27] Yap AU, Marpaung C, Gunardi I. Psychometric properties of the Indonesian Fonseca anamnestic index and the presence/severity of temporomandibular disorders among Indonesian young adults. Cranio. 2024;42:379–86. https://doi.org/10.1080/08869634.20
- [28] Linacre JM. Winsteps Rasch measurement computer program user's guide. Beaverton, OR; Winsteps.com; 2021.
- [29] Field A. Discovering statistics using SPSS. 3rd ed. London: Sage Publications: 2009.
- [30] Cunha CM, Almeida Neto OP, Stackfleth R. Main psychometric evaluation methods of measuring instrument's reliability. Rev Atenção Saúde. 2016;14:98–103. https://doi.org/10.13037/ras. vol14n47.3391
- [31] Linacre MJ. What do infit and outfit, mean-square and standardized mean? [Internet]. [cited 2024 Sept 3]. Available from: https://rasch. org/rmt/rmt162f.htm
- [32] Dabaghi S, Esmaielzadeh F, Rohani C. Application of Rasch analysis for development and psychometric properties of adolescents' quality of life instruments: a systematic review. Adolesc Health Med Ther. 2020;11:173–97. https://doi.org/10.2147/AHMT.5265413
- 33] Slade GD. Derivation and validation of a short-form oral health impact profile. Community Dent Oral Epidemiol. 1997;25:284–90. https://doi.org/10.1111/j.1600-0528.1997.tb00941.x
- [34] Yap AU, Cao Y, Zhang MJ, Lei J, Fu KY. Age-related differences in diagnostic categories, psychological states and oral health-related quality of life of adult temporomandibular disorder patients. J Oral Rehabil. 2021;48:361–8. https://doi.org/10.1111/joor.13121
- [35] Zafar MS, Fareed WM, Taymour N, Khurshid Z, Khan AH. Self-reported frequency of temporomandibular disorders among undergraduate students at Taibah University. J Taibah Univ Med Sci. 2017;12:517–22. https://doi.org/10.1016/j.jtumed.2016.12.004
- [36] Habib SR, Al Rifaiy MQ, Awan KH, Alsaif A, Alshalan A, Altokais Y. Prevalence and severity of temporomandibular disorders among university students in Riyadh. Saudi Dent J. 2015;27:125–30. https:// doi.org/10.1016/j.sdentj.2014.11.009

144 A. U. YAP ET AL.

- and new perspectives. J Orthop Sports Phys Ther. 1996;24:268–78. https://doi.org/10.2519/jospt.1996.24.4.268
- Hasvold T, Johnsen R, Førde OH. Non-migrainous headache, neck or shoulder pain, and migraine - differences in association with [43] background factors in a city population. Scand J Prim Health Care. 1996;14:92–9. https://doi.org/10.3109/02813439608997077 Jiménez-Silva A, Peña-Durán C, Tobar-Reyes J, Frugone-Zambra R. [44]
- Sleep and awake bruxism in adults and its relationship with temporomandibular disorders: a systematic review from 2003 to 2014. Acta Odontol Scand. 2017;75:36–58. https://doi.org/10.1080/00016 [45]
- de Oliveira Reis L, Ribeiro RA, Martins CC, Devito KL. Association between bruxism and temporomandibular disorders in children: a systematic review and meta-analysis. Int J Paediatr Dent. [46] 2019;29:585–59. https://doi.org/10.1111/ipd.12496
- $Man fredini\,D, Lombardo\,L, Siciliani\,G.\, Temporoman dibular\,disorders$ and dental occlusion. A systematic review of association studies: end of an era? J Oral Rehabil. 2017;44:908–23. https://doi.org/10.1111/ ioor.12531
- [37] Friedman MH, Nelson AJ. Head and neck pain review: traditional [42] Lozano Rojas OM, Rojas Tejada AJ, Pérez Meléndez C. Development of a specific health-related quality of life test in drug abusers using the Rasch rating scale model. Eur Addict Res. 2009;15:63-70. https:// doi.org/10.1159/000189784
 - Wong HM, McGrath CP, King NM. Rasch validation of the early childhood oral health impact scale. Community Dent Oral Epidemiol. 2011:39:449-57. https://doi.org/10.1111/i.1600-0528.2011.00614.x
 - Prieto L, Alonso J, Lamarca R. Classical test theory versus Rasch analysis for quality of life questionnaire reduction. Health Qual Life Outcomes. 2003;1:27. https://doi.org/10.1186/1477-7525-1-27
 - Rilo B, Femández-Formoso N, Mora MJ, Cadarso-Suárez C, Santana U. Distance of the contact glide in the closing masticatory stroke during mastication of three types of food. J Oral Rehabil. 2009;36:571–6. https://doi.org/10.1111/j.1365-2842.2009.01956.x
 - Mogil JS, Bailey AL. Sex and gender differences in pain and anal-gesia. Prog Brain Res. 2010;186:141–57. https://doi.org/10.1016/
 - B978-0-444-53630-3.00009-9 Althubaiti A. Information bias in health research: definition, pitfalls, and adjustment methods. J Multidiscip Healthc. 2016;9:211–17. https://doi.org/10.2147/JMDH.S104807


Dimensionality of the Fonseca Anamnestic Index and validation of its short-form derivative

	ALITY REPORT	ts snort-form de	rivative		
1 SIMILA	8% ARITY INDEX	14% INTERNET SOURCES	19% PUBLICATIONS	8% STUDENT PAPER	RS
PRIMAR	RY SOURCES				
1	Ylitervo, "Transla psychor version Questio	rrais Campos, To Leena Ylikontic Ition, cross-cultu netric propertie of the Malocclus nnaire (MIQ)", A avica, 2025	ola, Anna-Sofia ural adaptatio s of the Finnis sion Impact	a Silvola. n, and sh	2%
2	opende Internet Sour	ntistryjournal.co	om		2%
3	Castro, lanalysis the diag	Rodrigues-Biga Paulo F. Pires. "I of the Fonseca nosis of myoge omandibular dis of Physical Ther	Factor and Ras anamnestic ir nous sorder", Brazil	sch ndex for	2%
4	pmc.nck	oi.nlm.nih.gov			1%
5	WWW.res	searchsquare.co	om		1%
6	pubmed Internet Sour	d.ncbi.nlm.nih.g	OV		1%
7	and forr	Jjin Yap, Carolin m of temporom ms: Functional, social impacts",	andibular discophysical, and	order	1%



Publication

14	Adrian Ujin Yap, Ni Luh Dewi, Carolina Marpaung. "Psychological characteristics of young adults with temporomandibular disorders, somatization and combined conditions: A multidimensional evaluation", Journal of Oral Rehabilitation, 2023	<1%
15	scholarshare.temple.edu Internet Source	<1%
16	drum.lib.umd.edu Internet Source	<1%
17	Submitted to University of Sydney Student Paper	<1%
18	Armelia Sari Widyarman, Muhammad Ihsan Rizal, Moehammad Orliando Roeslan, Carolina Damayanti Marpaung. "Quality Improvement in Dental and Medical Knowledge, Research, Skills and Ethics Facing Global Challenges", CRC Press, 2024 Publication	<1%
19	Hanna Göthlin, Bengt Hasséus, Klas Sjöberg, Maria Bankvall. "Manifestations of the oral mucosa and salivary glands in irritable bowel syndrome and microscopic colitis – A systematic review", Acta Odontologica Scandinavica, 2025 Publication	<1%
20	"Higher Education: Handbook of Theory and Research", Springer Nature, 1999	<1%
21	Adrian Ujin Yap, Ye Cao, Minjuan Zhang, Jie Lei, Kai-Yuan Fu. "Temporomandibular Disorder severity and diagnostic groups: Their associations with sleep quality and impairments", Sleep Medicine, 2021	<1%



Exclude quotes On
Exclude bibliography On

Exclude matches

< 15 words