

[Home](#) / [User](#) / [Author](#) / [Submissions](#) / [#39339](#) / [Summary](#)

#39339 Summary

[Summary](#) | [Review](#) | [Editing](#)

Submission

Authors	Elline Elline, Tien Suwartini, Komariah Komariah, Reynaldo Lambertus
Title	The Characterization Of Hydroxyapatite, Eppigalocatechine-3-Gallate, Hydroxypropyl Methylcellulose As Bioactive Material For Pulp Capping
Original file	39339-88932-2-SM.docx 2024-08-07
Supp. files	None
Submitter	Elline Elline
Date submitted	August 7, 2024 - 08:46 PM
Section	Original Article
Editor	Shaista Afroz Muhammad Firdausy
Abstract Views	20

Status

Status	Published Vol 11, No 2 (2024): December 2024
Initiated	2025-03-05
Last modified	2025-03-05

Submission Metadata

Authors

Name	Elline Elline
Affiliation	Universitas Trisakti
Country	Indonesia
Bio Statement	Konservasi Gigi
Principal contact for editorial correspondence.	
Name	Tien Suwartini
Affiliation	Universitas Trisakti
Country	Indonesia
Bio Statement	Konservasi Gigi
Name	Komariah Komariah
Affiliation	Universitas Trisakti
Country	Indonesia
Bio Statement	–
Name	Reynaldo Lambertus
Affiliation	Universitas Trisakti
Country	Indonesia
Bio Statement	Konservasi Gigi

Title and Abstract

Title	The Characterization Of Hydroxyapatite, Eppigallocatechine-3-Gallate, Hydroxypropyl Methylcellulose As Bioactive Material For Pulp Capping
Abstract	<p>Background: According to a previous study, hydroxyapatite (HA) has the potential to be used in dentistry, but it still needs further research, especially as a biomaterial for pulp capping. The pulp capping treatment can maintain pulp vitality and induce dentin reparative formation. Calcium hydroxide is the gold standard of pulp capping material, but it can cause tunnel defects. Several studies made a crosslink with epigallocatechin-3-gallate (EGCG) and hydroxypropyl methylcellulose (HPMC) to improve the effect of hydroxyapatite as a biomaterial. The HA-EGCG-HPMC combination is expected to have the potential as a bioactive material. In this study, HA-EGCG-HPMC was characterized physicochemically by several criteria, such as the gelation time, pH, and antibacterial effect against <i>Aggregatibacter actinomycetemcomitans</i>. This study aims to determine the physicochemical characteristics of the HA-EGCG-HPMC material as a pulp capping material</p> <p>Method: This research is an in vitro laboratory experimental study design. Hydroxyapatite powder was dissolved with distilled water at concentrations of 4%, 2%, and 1%, with 10µmol/mL EGCG and hydroxypropyl methylcellulose as the carrier agent. Characterization of the samples measured are gelation time, pH, and antibacterial effect by microdilution method.</p> <p>Result: The HA-EGCG-HPMC combination has an ideal gelation time of about 23-31 minutes, and its pH is within the range of 7.28 - 7.33. However, the HA-EGCG-HPMC does not yet have an antibacterial effect against <i>Aggregatibacter actinomycetemcomitans</i>.</p> <p>Conclusion: The hydroxyapatite, EGCG, and hydroxypropyl methylcellulose can be used as a bioactive material, such as pulp capping material seen from physicochemical characterization that can be developed further.</p>

Indexing

Keywords	Characterization, Eppigallocatechine-3-gallate, Hydroxyapatite, Hydroxypropyl methylcellulose, Pulp capping
Language	en

Supporting Agencies

Agencies	Universitas Trisakti
-----------------	----------------------

Indexing

Keywords	Characterization, Eppigallocatechine-3-gallate, Hydroxyapatite, Hydroxypropyl methylcellulose, Pulp capping
Language	en

Supporting Agencies

Agencies	Universitas Trisakti
-----------------	----------------------

References

References	—
-------------------	---

#39339 Review

[Summary](#) | [Review](#) | [Editing](#)

Submission

Authors	Elline Elline, Tien Suwartini, Komariah Komariah, Reynaldo Lambertus
Title	The Characterization Of Hydroxyapatite, Eppigalocatechine-3-Gallate, Hydroxypropyl Methylcellulose As Bioactive Material For Pulp Capping
Section	Original Article
Editor	Shaista Afroz Muhammad Firdausy

Peer Review

Round 1

Review Version	39339-88933-2-RV.docx	2024-11-25
Initiated		2024-11-25
Last modified		2025-03-05
Uploaded file	Reviewer B 39339-95360-1-RV.docx	2024-12-20

Editor Decision

Decision	Accept Submission	2024-12-31
Notify Editor	Editor/Author Email Record	2024-12-31
Editor Version	39339-93069-1-ED.docx	2024-11-25
Author Version	39339-90883-1-ED.docx	2024-11-12
	39339-90883-2-ED.pdf	2024-12-27
	39339-90883-3-ED.pdf	2024-12-27

The Characterization Of Hydroxyapatite, Eppigalocatechine-3-Gallate, Hydroxypropyl Methylcellulose As Bioactive Material For Pulp Capping

Elline Elline*, Tien Suwartini*, Komariah**, Reynaldo Lambertus***

*Faculty of dental medicine, Department of Conservative Dentistry, Universitas Trisakti, Kyai Tapa Grogol No 26, Jakarta Barat, Indonesia

**Faculty of dental medicine, Department of Oral Biology. Sub division Histology, Universitas Trisakti, Kyai Tapa Grogol No 26, Jakarta Barat, Indonesia

***Post Graduate Student of Conservative Dentistry Universitas Trisakti, Kyai Tapa Grogol No 26, Jakarta Barat, Indonesia

Correspondence: elline@trisakti.ac.id

Received 07 August 2024; 1st revision 02 October 2024; 2st revision 27 December 2024; Accepted 31 December 2024; Published online 31 December 2024

Keywords:

Characterization,
Eppigalocatechine-3-
gallate, Hydroxyapatite,
Hydroxypropyl
methylcellulose, Pulp
capping

ABSTRACT

Background: According to a previous study, hydroxyapatite (HA) has the potential to be used in dentistry, but it still needs further research, especially as a biomaterial for pulp capping. The pulp capping treatment can maintain pulp vitality and induce dentin reparative formation. Calcium hydroxide is the gold standard of pulp capping material, but it can cause tunnel defects. Several studies made a crosslink with epigallocatechin-3-gallate (EGCG) and hydroxypropyl methylcellulose (HPMC) to improve the effect of hydroxyapatite as a biomaterial. The HA-EGCG-HPMC combination is expected to have the potential as a bioactive material. In this study, HA-EGCG-HPMC was characterized physicochemically by several criteria, such as the gelation time, pH, and antibacterial effect against *Aggregatibacter actinomycetemcomitans*. This study aims to determine the physicochemical characteristics of the HA-EGCG-HPMC material as a pulp capping material

Method: This research is an *in vitro* laboratory experimental study design. Hydroxyapatite powder was dissolved with distilled water at concentrations of 4%, 2%, and 1%, with 10 $\mu\text{mol/mL}$ EGCG and hydroxypropyl methylcellulose as the carrier agent. Characterization of the samples measured are gelation time, pH, and antibacterial effect by microdilution method.

Result: The HA-EGCG-HPMC combination has an ideal gelation time of about 23-31 minutes, and its pH is within the range of 7.28 - 7.33. However, the HA-EGCG-HPMC does not yet have an antibacterial effect against *Aggregatibacter actinomycetemcomitans*.

Conclusion: The hydroxyapatite, EGCG, and hydroxypropyl methylcellulose can be used as a bioactive material, such as pulp capping material seen from physicochemical characterization that can be developed further

Copyright ©2024 National Research and Innovation Agency. This is an open access article under the CC BY-SA license (<https://creativecommons.org/licenses/by-sa/4.0/>).

doi: <http://dx.doi.org/10.30659/odj.11.2.366-374>

2460-4119 / 2354-5992 ©2024 National Research and Innovation Agency

This is an open access article under the CC BY-SA license (<https://creativecommons.org/licenses/by-sa/4.0/>)

Odonto : Dental Journal accredited as Sinta 2 Journal (<https://sinta.kemdikbud.go.id/journals/profile/3200>)

How to Cite: Elline et al. The Characterization Of Hydroxyapatite, Eppigalocatechine-3-Gallate, Hydroxypropyl Methylcellulose As Bioactive Material For Pulp Capping. Odonto: Dental Journal, v.11, n.2, p. 366-374, December 2024.

INTRODUCTION

Dental caries is the most prevalent disease worldwide, with 2.3 billion people suffering from caries in permanent teeth.¹ The results of Hasil Riset Kesehatan Dasar (RISKESDAS) or Basic Health Research in 2018, the most significant dental problems in Indonesia are cavities and toothache at 57.6%, and 76.2% of children aged 12 years have caries so needs to be a concern. Untreated caries can cause dental pulp necrosis, which causes pulp vitality to be lost and teeth to become brittle.²

The pulp capping treatment is used when the pulp is exposed due to caries, trauma, or iatrogenic causes, such as during caries cleaning or tooth preparation. Direct or indirect pulp capping is usually used in deep cavities with unexposed pulp.³ The commonly used reconstructed pulp cap therapy materials are calcium hydroxide Ca(OH)₂ or MTA. Ca(OH)₂ has a high pH of up to 12.5, which is an antibacterial material that provides hemostasis and stimulates mineralized tissue and the formation of dentin bridges.^{3,4} According to William, under The International Standard (ISO) 9917-1 guidance, all water-based dental cement takes 1 hour to become rigid when stirred, so it can be used for applications.⁵

The limitations of Ca(OH)₂ are that the high pH can irritate the dental pulp and can cause inflammation or necrosis of the exposed pulp surface, poor seal strength and no adhesion quality to the tissue, poor physical properties, and dissolves over time, causing tunnel defects in the reparative dentin to form under the calcium hydroxide, facilitating pulp recontamination.⁶

Mineral trioxide aggregate (MTA) is often used as a recycled pulp cap treatment because it has the advantages of good mechanical properties, good seal formation, antibacterial, biocompatible, and high pH.⁷ However, MTA limitation leads to pulp obliteration, which makes it difficult to treat a root canal.⁸ There are limitations to the materials commonly used in pulp cap treatment, so other materials that function optimally but can reduce the shortcomings of the previous materials are needed.

Hydroxyapatite (HA) has been used extensively in dentistry, including in the field of vital pulp therapy.⁹ HA may form dentinal bridges in pulp capping treatments.^{10,11} The advantage of HA from chicken eggshells is the level of hydroxyapatite (0.0950g/g).¹² *Epigallocatechin-3-gallate (EGCG)* is 50-60% of the content of catechins, which are polyphenolic chemical compounds from tea leaves.¹³ EGCG has antioxidant, antitumor, antibacterial, and anti-inflammatory effects and is good at maintaining cell viability.¹⁴ EGCG is proven to increase the mechanical properties of materials.¹⁵ 10µg/mL EGCG has anti-inflammatory effects without affecting the proliferation and differentiation of pulp cells so that it can be used for pulp capping.¹³

HA and EGCG are to be applied as pulp capping materials, and a suspending agent using a 2% hydroxypropyl methylcellulose (HPMC) polymer is required to form a phase from liquid to gel.¹⁶ The mixture of HA, EGCG, and HPMC carrier agents is expected to support and accelerate the formation of reparative dentin without the occurrence of severe inflammation.

Lactobacillus, Actinomyces, and Olsenella dominate bacteria in deep carious lesions. Actinomyces is the second most common bacterium in deep carious lesions, and it can induce leukocytes to apoptosis and lysis.¹⁷ So, it is expected that the mixture of HA and EGCG can be used as a therapeutic material for recycled pulp caps and can kill actinomyces bacteria.

MATERIALS AND METHODS

1. Formulation of HA-EGCG-HPMC hydrogel

Hydroxyapatite powder (ProDB , PT.Aleesha Berkah Utama, Bekasi, Jawa Barat, Indonesia) is dissolved in deionized water and stirred with a magnetic stirrer at a speed of 350 rpm with a concentration of 1%, 2%, and 4%.¹ 10µg/mL EGCG (Sigma Aldrich, E4268, 80 %, USA).was added to the hydroxyapatite solution. Each sample was stirred until homogeneous at a temperature of 40°C for 30 minutes, and the carrier material 2% HPMC (Benecel, K100M, Ashland, Wilmington,USA) was added.¹⁸

2. Gelation time

Gelation time was performed with an inverted tube method test. Gelation time was determined by changes in sHAe from liquid to gel (sol-gel phase). Three replications were done to measure the average value.

3. pH

The pH was measured using a pHmeter, and it was calibrated first with an acid solution with 4.0 pH and 10,0 base pH. The pH value of HA-EGCG-HPMC was determined after the formulation was mixed at 1,20,40, and 60 minutes with four repetitions each.⁴

4. Antibacterial Test

a. The Microdilution Test Preparation

0,1 ml samples of 4% HA-EGCG-HPMC, 2%HA-EGCG-HPMC, and 1%HA-EGCG-HPMC, positive control (BioMTA+, PPH Cerkamed, Stawola Wola) were prepared into a 96-well plate with four repetitions. Each well was inoculated with 20 µL of the *Aggregatibacter actinomycetemcomitans* microbial suspension with the turbidity of the test colony suspension standardized to 0.5 McFarland (approximately 1.5x10⁸ CFU/mL). Samples were incubated for 24 hours at a temperature of 37°C.⁵

b. The Bacterial Suspension Test Preparation

The *Aggregatibacter actinomycetemcomitans* test colony suspension was made by taking one colony phase from the solid agar culture medium (24 hours of bacteria cultured on the agar medium) into a reaction tube containing 5 mL of physiological NaCl. The tube was then vibrated, and the turbidity of the test colony suspension was standardized to 0.5 McFarland (approximately 1.5x10⁸ CFU/mL). The suspension must be used as an inoculum within 15 minutes.

5. Statistical Analysis

All data obtained were subjected to a Shapiro–Wilk normality test. If the data were normal, a one-way analysis of variance (ANOVA) was used. Otherwise, the data were analyzed using Kruskal–Wallis. The Wilcoxon statistical test measured the pH because the data was not normally distributed.

RESULTS

The gelation time test results obtained with 4% HA+EGCG+HPMC, 2% HA+EGCG+HPMC, and 1% HA+EGCG+HPMC are presented with the mean and standard deviation (Sd) in Table 1.

Table 1. Descriptive gelation time data

Group name	N	Mean ± Sd (minutes)
HA 4%	9	23,78 ± 0,88
HA 2%	9	27,67 ± 0,50
HA 1%	9	30,00 ± 0,71

Based on Table 1., the average gelation time of the HA-EGCG-HPMC mixture at a concentration of 4% obtained a gelation time of 23.78 minutes, 2% HA-EGCG-HPMC obtained a gelation time of 27.67 minutes, and 1% HA-EGCG-HPMC obtained a gelation time of 30 minutes, then continued with a non-parametric statistical test, namely the Kruskal Wallis test ($p = 0.000$). The results of the Kruskal Wallis test followed by the Mann Whitney test showed that the 4% HA-EGCG-HPMC group had a significant difference with 2% HA-EGCG-HPMC ($p=0.045$) and 1% ($p=0.000$), for 1% HA-EGCG-HPMC showed a significant difference with the 2% HA-EGCG-HPMC group with a value of $p=0.045$. Concentrations of 4%, 2%, and 1% showed gelation times below 60 minutes to qualify as pulp cap treatment materials (table 2.).

Table 2. Post-hoc Mann Whitney test of gelation time

Sample 1(HA-EGCG-HPMC)	Sample 2 (HA-EGCG-HPMC)	p
HAP 4%	HAP 2%	*0,045
HAP 4%	HAP 1%	*0,000
HAP 2%	HAP 1%	*0,045
*P<0,05		

The pH test was conducted to determine the basicity contained in the sample. The pH test was carried out using a sample tested using a pH-meter calculated at the time after the material was stirred until homogeneous and measured periodically at 1 minute, 20 minutes, 40 minutes, and 60 minutes, presented with the mean and standard deviation in Table 3.

Table 3. Mean and standard deviation of pH test

Time	Group Name	N	Mean ± Sd
1 minute	HA 4%	9	7,32 ± 0,01
	HA 2%	9	7,30 ± 0,02
	HA 1%	9	7,39 ± 0,01
20 minutes	HA 4%	9	7,32 ± 0,09
	HA 2%	9	7,29 ± 0,01
	HA 1%	9	7,31 ± 0,01
40 minutes	HA 4%	9	7,31 ± 0,01
	HA 2%	9	7,30 ± 0,01
	HA 1%	9	7,29 ± 0,01
60 minutes	HA 4%	9	7,31 ± 0,01
	HA 2%	9	7,30 ± 0,01
	HA 1%	9	7,29 ± 0,01

The test results were continued with Wilcoxon non-parametric ($p < 0.05$). The results of further tests using Post-hoc Mann Whitney obtained 1 minute group (HA 2%) and 1 minute (HA 1%), 20 minutes group (HA 2%) and 20 minutes (HA 4%) ($p=0.04$), 40 minutes group (HA 2%) and 1 minute (HA 1%), 20 minutes group (HA 2%) and 1 minute (HA 1%), 60 minutes group (HA 1%) and 1 minute (HA 1%), 40 minutes group (HA 1%) and 1 minute (HA 1%) ($p=0.00$), group 40 min (HA 1%) and 20 min (HA 4%), group 40 min (HA 1%) and 1 min (HA 4%) ($p=0.005$), group 60 min (HA 2%) and 1 min (HA 1%) ($p=0.02$), group 60 minutes (HA 1%) and 60 minutes (HA 4%) ($p=0.03$), group 60 minutes (HA 1%) and 1 minute (HA 4%) ($p=0.01$), and group 60 minutes (HA 1%) and 20 minutes (HA 4%) ($p=0.01$) significantly different. Based on the results of the material test, it was found that the mixture at concentrations of 4%, 2%, and 1% showed an average value of neutral pH with a range of 7.29-7.31. These results show that the tested HA-EGCG-HPMC material is as expected, which is close to the normal pulp pH of 7.3.

The antibacterial test was conducted in 5 test groups, namely: negative control group (distilled water), 4%HA-EGCG-HPMC, 2%HA 2%-EGCG-HPMC, 1%HA- EGCG-HPMC, and positive control group (4%MTA) samples were mixed with a solution that already contained *Aggregatibacter actinomycetemcomitans* bacteria followed by incubation of samples for 24 hours at 37 °C, then the results were observed by turbidimetric method, and all samples were turbid in color indicating that all samples had bacterial growth, then the samples were applied to MHA agar media to confirm the samples had bacterial growth. The calculation of the number of colonies formed on agar media is presented in Figure 1.




Figure 1. Picture of the number of colonies formed in each sample treatment

The ANOVA test results showed significant differences between the test groups ($p < 0.01$). Further tests were conducted to see the differences between groups. The results of the post-hoc test showed that the number of colonies in the negative group was significantly different from 4%HA-EGCG-HPMC, 2%HA -EGCG-HPMC, 1%HA-EGCG-HPMC, and 4%MTA ($p < 0.01$) with mean colony counts of $1 \times 10^8 \pm 55 \times 10^4$, $2.5 \times 10^7 \pm 28 \times 10^4$, $3.7 \times 10^7 \pm 14 \times 10^4$, $6.4 \times 10^7 \pm 87 \times 10^3$, and $1.3 \times 10^7 \pm 16 \times 10^4$ CFU/mL, respectively. The number of colonies of 4% HA-EGCG-HPMC significantly differed from 1% HA -EGCG-HPMC ($p < 0.001$), which showed that 4% HA-EGCG-HPMC was lower than 1% HA.

number of colonies of 2%HA-EGCG-HPMC was slightly lower than that of 1%HA -EGCG-HPMC ($p<0.001$) and 4%MTA ($p<0.001$), 1%HA-EGCG-HPMC group was higher than 4% MTA group ($p<0.001$) (Figure 2).




Figure 2A-E. Sample test results against the bacteria *A. actinomycetemcomitans* (A. MTA, B. Negative control, C. 1%HA-EGCG-HPMC, D. 2%HA-EGCG-HPMC, E. 4%HA-EGCG-HPMC)

In Figure 2, the density of *A.actinomycetemcomitans* in 4% MTA is the least formed when compared to the control (-), but there is no significant difference in the 4% MTA group with 4% HA-EGCG-HPMC. There was a significant difference in the number of colonies in the 4% HA-EGCG-HPMC group with 1% HA-EGCG-HPMC. Based on the post-hoc test, there is a density of *A. actinomycetemcomitans* in each test sample of the HA-EGCG-HPMC mixture, so it can be concluded that the HA-EGCG-HPMC mixture does not have an antibacterial effect on *A. actinomycetemcomitans* bacteria.

DISCUSSION

The purpose of performing a reconstructed pulp capping is to increase the healing ability of the pulp and promote the formation of a mineralized tissue layer by placing the biomaterial directly on the exposed pulp roof.¹⁹ In this study, a mixture of HA-EGCG-HPMC materials was tested to find alternative materials for pulp capping treatment. First, a characterization test of the hydrogel suspension material with hydroxyapatite, EGCG, and HPMC was carried out, and it was seen that the gelation time was sufficient so that it could be used for clinical applications, whereas slow gelation time made it difficult to apply in a certain area.²⁰ The hydrogel suspension mixture with a concentration of 4% HA-EGCG-HPMC had a gelation time an average of 24-26 minutes, at a concentration of 2% the gelation material at an average time of 27-30 minutes, and at a concentration of 1% the gelation material averaged 29-31 minutes so that the hypothesis was accepted because it had sufficient gelation time for clinical applications, which was under 60 minutes.²¹

In this study, the level of basicity of the HA-EGCG-HPMC material mixture was tested using a pH meter, and samples with three concentration mixtures were shown in Table 5.2. The results show that the material after adding the gelation agent and after the gelation material shows a neutral pH (Table 2), so it is to the pH conditions in the pulp chamber, namely 7.3-7.8.²² According to Elline et al,

pH is essential in determining the morphology and structure of a formulation. The pH can influence the ion balance. The increase or decrease of OH⁻ ion, will affect the Ca²⁺, PO₄³⁻, and (HPO₄)²⁻ ions¹⁸

The results of the antibacterial effect test of the HA-EGCG-HPMC mixture using the dilution method of 4%, 2%, and 1% concentrations of mixed materials mixed with *Aggregatibacter actinomycetemcomitans* bacterial suspensions in 96-well plates and incubated at 37°C for 24 hours, the results showed that all samples contained bacterial growth. Confirmation of bacterial growth and development is achieved by scratching on agar discs to see the growth of the number of bacterial colonies present (Figure 2). Thus, the HA-EGCG-HPMC mixture has no antibacterial effect on *Aggregatibacter actinomycetemcomitans* bacteria. According to a previous study, EGCG would break the bacterial plasma membrane in a certain concentration. For example, the *Streptococcus mutans* plasma membrane will be destroyed with 0,2 mg/ml after EGCG treatment.²³ So, in this study may be the concentration of EGCG has not fulfilled the minimum criteria as a treatment

CONCLUSIONS

Based on the test results, the HA-EGCG-HPMC mixture at a concentration of 4% has a gelation time of 23.78 minutes. It is concluded that this material has a qualification to be used in clinical conditions. It also has a degree of acidity that resembles the vital pulp at 7.30 after gelation. In the antibacterial test results, the 4% HA-EGCG-HPMC mixture had a lower antibacterial effect when compared to 4% MTA. Further research is needed to develop a mixture of HA-EGCG-HPMC materials to increase the ability of antibacterial effects that can support the potential of the material to become a pulp capping material

ACKNOWLEDGMENT

There is financial assistance from FKG Universitas, and the researchers would to acknowledge

REFERENCES

1. Kazemina M, Abdi A, Shohaimi S, Jalali R, Vaisi-Raygani A, Salari N, et al. Dental caries in primary and permanent teeth in children's worldwide, 1995 to 2019: A systematic review and meta-analysis. , Head and Face Medicine. 2020. 16(1);22. Available from : <https://pubmed.ncbi.nlm.nih.gov/33023617/>
2. Karies Gigi Anak di Jakarta Timur P, Putri N, Lusida N, Rosyada D, Ala Al-Maududi A, Studi Kesehatan Masyarakat P, et al. Faktor yang Berhubungan dengan Perilaku Orangtua. 2023. JKKI. 19(1): 11-17. Available from: <https://jurnal.umj.ac.id/index.php/JKK>
3. Alp Ş, Ulusoy N. Current Approaches in Pulp Capping: A Review. Cyprus J Med Sci. 2024. 28;154–60. Available from : <https://cyprusjmedsci.com/articles/current-approaches-in-pulp-capping-a-review/doi/cjms.2023.2022-37>
4. Darmadi EY, Soesilo D. Antimicrobial activity of calcium hydroxide, calcium oxide, and mineral trioxide aggregate paste against α-Hemolytic Streptococcus. CDJ. 2024 Jul 24;14(1):24–8. Available from : <https://e-journal.unair.ac.id/CDJ/article/view/60848>
5. Ha W, Nicholson T, Kahler B, Walsh L. Rheological Characterization as an Alternative Method to Indentation for Determining the Setting Time of Restorative and Endodontic Cements. Materials. 2017.20;10(12):1451. Available from : https://www.researchgate.net/publication/321959668_Rheological_Characterization_as_an_Alternative_Method_to_Indentation_for_Determining_the_Setting_Time_of_Restorative_and_Endodontic_Cements
6. Covaci A, Ciocan LT, Gălbinașu B, Bucur MV, Matei M, Didilescu AC. Dental Pulp Response to Different Types of Calcium-Based Materials Applied in Deep Carious Lesion

- Treatment—A Clinical Study. *J Funct Biomater*. 2022. 2;13(2):51. Available from : <https://www.mdpi.com/2079-4983/13/2/51>
7. Kunert M, Lukomska-Szymanska M. Bio-Inductive Materials in Direct and Indirect Pulp Capping - A Review Article. *Materials*.2020 (13). Available from : <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7085085/>
 8. Eid A, Mancino D, Rekab MS, Haikel Y, Kharouf N. Effectiveness of Three Agents in Pulpotomy Treatment of Permanent Molars with Incomplete Root Development: A Randomized Controlled Trial. *Healthcare*. 2022. 25;10(3):431. Available from : <https://pubmed.ncbi.nlm.nih.gov/35326909/>
 9. James Irawan I, Adi Wicaksono I, Raya Bandung Sumedang Km J. Review Artikel: Penggunaan Material Bioaktif Dalam Terapi Sel Punca Untuk Karies Gigi. *Open Journal Systems STF Muhammadiyah Cirebon* : ojs.stfmuhammadiyahcirebon.ac.id. 2022.7 Available from : <https://www.semanticscholar.org/reader/e6321d801888edcc04f6b93e87249b548a6fb680>
 10. Niazi FH, Luddin N, Harun MH, Hasan A, Kannan TP, Mohamad S, et al. Dentin–Pulp Complex Response in Molars of Rats after Occlusal and Cervical Restorations with Conventional Glass Ionomer Cement and Nano-Hydroxyapatite Silica Glass Ionomer Cement. *Appl. Sci*. 2023. 1;13(5). Available from : <https://www.mdpi.com/2076-3417/13/5/3156>
 11. Rahmitasari F, Cevanti TA, Prabowo PB, Sularsih S, Prananingrum W. The effect of incorporating hydroxyapatite into Type II glass ionomer cement on flexural strength and the examination of fractured surfaces using scanning electron microscopy. *ODJ*. 2023. 29;10(2):152. Available from : <http://jurnal.unissula.ac.id/index.php/odj/article/viewFile/33491/9378>
 12. Elline E, Ismiyatin K. Nanohydroxiapatite Using Chicken Eggshell Waste and Its Characterization. *MJMHS*. 2021.17. Available from : https://medic.upm.edu.my/upload/dokumen/2021092200460413_2021_0338.pdf
 13. Li Y, Zhao Y, Han J, Wang Y, Lei S. Effects of epigallocatechin gallate (EGCG) on the biological properties of human dental pulp stem cells and inflammatory pulp tissue. *Arch Oral Biol*. 2021. 123:105034. Available from : <https://www.sciencedirect.com/science/article/abs/pii/S000399692030412X>
 14. Mokra D, Joskova M, Mokry J. Therapeutic Effects of Green Tea Polyphenol (–)-Epigallocatechin-3-Gallate (EGCG) in Relation to Molecular Pathways Controlling Inflammation, Oxidative Stress, and Apoptosis. *Int. J. Mol. Sci*. 2023.24. Available from : <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9820274/>
 15. Amanda HG, Elline E, Fibryanto E. Synthesis and Physical Characterization of Nano-Hydroxyapatite-Collagen-Epigallocatechin-3-Gallate Hydrogel Composite. *JIDA* 2022.17;5(1):7. Available from : https://www.researchgate.net/publication/361867826_Synthesis_and_Physical_Characterization_of_Nano-Hydroxyapatite-Collagen-Epigallocatechin-3-Gallate_Hydrogel_Composite#fullTextFileContent
 16. Santoso JH, Prahasti AE, Elline. Effect Of 4 Wt% Hydroxyapatite Pasta On Dental Email Hardness. *IJKG*. 2024. 21;20(1):46–52. Available from : <https://e-journal.unmas.ac.id/index.php/interdental/article/view/8580>
 17. Paul O, Arora P, Mayer M, Chatterjee S. Inflammation in Periodontal Disease: Possible Link to Vascular Disease. *Front.Physiol*. 2021. 11; 1-11.Available from : <https://pubmed.ncbi.nlm.nih.gov/33519515/>
 18. Elline E, Ismiyatin K, Indah Budhy T, Bhardwaj A. The potential of eggshell hydroxyapatite, collagen, and EGCG (HAp-Col-EGCG) scaffold as a pulp regeneration material. *Saudi Dent. J*. 2022. 1;34(8):715–22. Available from : <https://pubmed.ncbi.nlm.nih.gov/36570587/>
 19. Islam R, Islam MRR, Tanaka T, Alam MK, Ahmed HMA, Sano H. Direct pulp capping procedures – Evidence and practice. *Jpn. Dent. Sci. Rev*. 2023.59:48–61. Available from : <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9985044/>
 20. El-Sayed NS, Kamel S. Polysaccharides-Based Injectable Hydrogels: Preparation, Characteristics, and Biomedical Applications . *J.Colloids Interfaces*. 2022. 6(78). Available from : <https://www.mdpi.com/2504-5377/6/4/78>
 21. Zhang Y, Huang Y. Rational Design of Smart Hydrogels for Biomedical Applications. Vol. 8, *Front. in Chem*. 2021.8;1-13. Available from : <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7889811/>

22. Singh PH. Theory of Optimum pH for Pulp Vitality: A Clinical and Experimental Theory of Endodontics. 2020. 3;1-3. Available from : https://www.researchgate.net/publication/338402785_Theory_of_Optimum_pH_for_Pulp_Vitality_A_Clinical_and_Experimental_Theory_of_Endodontics
23. Kong C, Zhang H, Li L, Liu Z. Effects of green tea extract epigallocatechin-3-gallate (EGCG) on oral disease-associated microbes: a review. Vol. 14, Journal of Oral Microbiology. Taylor and Francis Ltd.; 2022. 14;1-14. Available from : <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9542882/>

odonto

by Eline Istanto

Submission date: 10-Mar-2025 06:51PM (UTC+0700)

Submission ID: 2610601724

File name: REV1_Odonto_Jurnal_ID39339.pdf (1.12M)

Word count: 3657

Character count: 19703

The Characterization Of Hydroxyapatite, Eppigalocatechine-3-Gallate, Hydroxypropyl Methylcellulose As Bioactive Material For Pulp Capping

Elline Elline^{*}, Tien Suwartini^{*}, Komariah^{**}, Reynaldo Lambertus^{***}

^{*}Faculty of dental medicine, Department of Conservative Dentistry, Universitas Trisakti, Kyai Tapa Grogol No 26, Jakarta Barat, Indonesia

^{**}Faculty of dental medicine, Department of Oral Biology. Sub division Histology, Universitas Trisakti, Kyai Tapa Grogol No 26, Jakarta Barat, Indonesia

^{***}Post Graduate Student of Conservative Dentistry Universitas Trisakti, Kyai Tapa Grogol No 26, Jakarta Barat, Indonesia

Correspondence: elline@trisakti.ac.id

ABSTRACT

Background: According to a previous study, hydroxyapatite (HA) has the potential to be used in dentistry, but it still needs further research, especially as a biomaterial for pulp capping. The pulp capping treatment can maintain pulp vitality and induce dentin reparative formation. Calcium hydroxide is the gold standard of pulp capping material, but it can cause tunnel defects. Several studies made a crosslink with epigallocatechin-3-gallate (EGCG) and hydroxypropyl methylcellulose (HPMC) to improve the effect of hydroxyapatite as a biomaterial. The HA-EGCG-HPMC combination is expected to have the potential as a bioactive material. In this study, HA-EGCG-HPMC was characterized physicochemically by several criteria, such as the gelation time, pH, and antibacterial effect against *Aggregatibacter actinomycetemcomitans*. This study aims to determine the physicochemical characteristics of the HA-EGCG-HPMC material as a pulp capping material.

Method: This research is an in vitro laboratory experimental study design. Hydroxyapatite powder was dissolved with distilled water at concentrations of 4%, 2%, and 1%, with 10µmol/mL EGCG and hydroxypropyl methylcellulose as the carrier agent. Characterization of the samples measured are gelation time, pH, and antibacterial effect by microdilution method.

Result: The HA-EGCG-HPMC combination has an ideal gelation time of about 23-31 minutes, and its pH is within the range of 7.28 - 7.33. However, the HA-EGCG-HPMC does not yet have an antibacterial effect against *Aggregatibacter actinomycetemcomitans*.

Conclusion: The hydroxyapatite, EGCG, and hydroxypropyl methylcellulose can be used as a bioactive material, such as pulp capping material seen from physicochemical characterization that can be developed further.

Keywords: Characterization, Eppigalocatechine-3-gallate, Hydroxyapatite, Hydroxypropyl methylcellulose, Pulp capping

INTRODUCTION

Dental caries is the most prevalent disease worldwide, with 2.3 billion people suffering from caries in permanent teeth.¹ The results of Hasil Riset Kesehatan Dasar (RISKESDAS) or Basic Health Research in 2018, the most significant dental problems in Indonesia are cavities and toothache at 57.6%, and 76.2% of children aged 12 years have caries so needs to be a concern. Untreated caries can cause dental pulp necrosis, which causes pulp vitality to be lost and teeth to become brittle.²

The pulp capping treatment is used when the pulp is exposed due to caries, trauma, or iatrogenic causes, such as during caries cleaning or tooth preparation. Direct or indirect pulp capping is usually used in deep cavities with unexposed pulp.³ The commonly used reconstructed pulp cap therapy materials are calcium hydroxide Ca(OH)₂ or MTA. Ca(OH)₂ has a high pH of up to 12.5, which is an antibacterial material that provides hemostasis and stimulates mineralized tissue and the formation of dentin bridges.^{3,4} According to William, under The International Standard (ISO) 9917-1 guidance, all water-based dental cement takes 1 hour to become rigid when stirred, so it can be used for applications.⁵

The limitations of Ca(OH)₂ are that the high pH can irritate the dental pulp and can cause inflammation or necrosis of the exposed pulp surface, poor seal strength and no adhesion quality to the tissue, poor physical properties, and dissolves over time, causing tunnel defects in the reparative dentin to form under the calcium hydroxide, facilitating pulp recontamination.⁶

Mineral trioxide aggregate (MTA) is often used as a recycled pulp cap treatment because it has the advantages of good mechanical properties, good seal formation, antibacterial, biocompatible, and high pH.⁷ However, MTA limitation leads to pulp obliteration, which makes it difficult to treat a root canal.⁸ There are limitations to the materials commonly used in pulp cap treatment, so other materials that function optimally but can reduce the shortcomings of the previous materials are needed.

Hydroxyapatite (HA) has been used extensively in dentistry, including in the field of vital pulp therapy.⁹ HA may form dentinal bridges in pulp capping treatments.^{10,11} The advantage of HA from chicken eggshells is the level of hydroxyapatite (0.0950g/g).¹² *Epigallocatechin-3-gallate (EGCG)* is 50-60% of the content of catechins, which are polyphenolic chemical compounds from tea leaves.¹³ EGCG has antioxidant, antitumor, antibacterial, and anti-inflammatory effects and is good at maintaining cell viability.¹⁴ EGCG is proven to increase the mechanical properties of materials.¹⁵ 10µg/mL EGCG has anti-inflammatory effects without affecting the proliferation and differentiation of pulp cells so that it can be used for pulp capping.¹³

HA and EGCG are to be applied as pulp capping materials, and a suspending agent using a 2% hydroxypropyl methylcellulose (HPMC) polymer is required to form a phase from liquid to gel.¹⁶ The mixture of HA, EGCG, and HPMC carrier agents is expected to support and accelerate the formation of reparative dentin without the occurrence of severe inflammation.

Lactobacillus, Actinomyces, and Olsenella dominate bacteria in deep carious lesions. Actinomyces is the second most common bacterium in deep carious lesions, and it can induce leukocytes to apoptosis and lysis.¹⁷ So, it is expected that the mixture of HA and EGCG can be used as a therapeutic material for recycled pulp caps and can kill actinomyces bacteria.

MATERIALS AND METHODS

1. Formulation of HA-EGCG-HPMC hydrogel

Hydroxyapatite powder (ProDB, PT. Aleesha Berkah Utama, Bekasi, Jawa Barat, Indonesia) is dissolved in deionized water and stirred with a magnetic stirrer at a speed of 350 rpm with a concentration of 1%, 2%, and 4%.¹ 10µg/mL EGCG (Sigma Aldrich, E4268, 80%, USA) was added to the hydroxyapatite solution. Each sample was stirred until homogeneous at a temperature

of 40°C for 30 minutes, and the carrier material 2% HPMC (Benecel, K100M, Ashland, Wilmington, USA) was added.¹⁸

2. Gelation time

Gelation time was performed with an inverted tube method test. Gelation time was determined by changes in sHAe from liquid to gel (sol-gel phase). Three replications were done to measure the average value.

3. pH

The pH was measured using a pHmeter, and it was calibrated first with an acid solution with 4.0 pH and 10,0 base pH. The pH value of HA-EGCG-HPMC was determined after the formulation was mixed at 1,20,40, and 60 minutes with four repetitions each.⁴

4. Antibacterial Test

a. The Microdilution Test Preparation

0,1 ml samples of 4% HA-EGCG-HPMC, 2%HA-EGCG-HPMC, and 1%HA-EGCG-HPMC, positive control (BioMTA+, PPH Cerkamed, Stawola Wola) were prepared into a 96-well plate with four repetitions. Each well was inoculated with 20 µL of the *Aggregatibacter actinomycetemcomitans* microbial suspension with the turbidity of the test colony suspension standardized to 0.5 McFarland (approximately 1.5x10⁸ CFU/mL). Samples were incubated for 24 hours at a temperature of 37°C.⁵

b. The Bacterial Suspension Test Preparation

The *Aggregatibacter actinomycetemcomitans* test colony suspension was made by taking one colony phase from the solid agar culture medium (24 hours of bacteria cultured on the agar medium) into a reaction tube containing 5 mL of physiological NaCl. The tube was then vibrated, and the turbidity of the test colony suspension was standardized to 0.5 McFarland (approximately 1.5x10⁸ CFU/mL). The suspension must be used as an inoculum within 15 minutes.

5. Statistical Analysis

All data obtained were subjected to a Shapiro–Wilk normality test. If the data were normal, a one-way analysis of variance (ANOVA) was used. Otherwise, the data were analyzed using Kruskal–Wallis. The Wilcoxon statistical test measured the pH because the data was not normally distributed.

RESULTS

The gelation time test results obtained with 4% HA+EGCG+HPMC, 2% HA+EGCG+HPMC, and 1% HA+EGCG+HPMC are presented with the mean and standard deviation (Sd) in Table 1.

Table 1. Descriptive gelation time data

Group name	N	Mean ± Sd (minutes)
HA 4%	9	23,78 ± 0,88
HA 2%	9	27,67 ± 0,50
HA 1%	9	30,00 ± 0,71

Based on Table 1., the average gelation time of the HA-EGCG-HPMC mixture at a concentration of 4% obtained a gelation time of 23.78 minutes, 2% HA-EGCG-HPMC obtained a gelation time of 27.67 minutes, and 1% HA-EGCG-HPMC obtained a gelation time of 30 minutes, then continued with a non-parametric statistical test, namely the Kruskal Wallis test ($p = 0.000$). The results of the Kruskal Wallis test followed by the Mann Whitney test showed that the 4% HA-EGCG-HPMC group had a significant difference with 2% HA-EGCG-HPMC ($p=0.045$) and 1% ($p=0.000$), for 1% HA-EGCG-HPMC showed a significant difference with the 2% HA-EGCG-HPMC group with a value of $p=0.045$. Concentrations of 4%, 2%, and 1% showed gelation times below 60 minutes to qualify as pulp cap treatment materials (table 2.).

Table 2. Post-hoc Mann Whitney test of gelation time

Sample 1(HA-EGCG-HPMC)	Sample 2 (HA-EGCG-HPMC)	p
HAP 4%	HAP 2%	*0,045
HAP 4%	HAP 1%	*0,000
HAP 2%	HAP 1%	*0,045
*P<0,05		

The pH test was conducted to determine the basicity contained in the sample. The pH test was carried out using a sample tested using a pH-meter calculated at the time after the material was stirred until homogeneous and measured periodically at 1 minute, 20 minutes, 40 minutes, and 60 minutes, presented with the mean and standard deviation in Table 3.

Table 3. Mean and standard deviation of pH test

Time	Group Name	N	Mean \pm Sd
1 minute	HA 4%	9	7,32 \pm 0,01
	HA 2%	9	7,30 \pm 0,02
	HA 1%	9	7,39 \pm 0,01
20 minutes	HA 4%	9	7,32 \pm 0,09
	HA 2%	9	7,29 \pm 0,01
	HA 1%	9	7,31 \pm 0,01
40 minutes	HA 4%	9	7,31 \pm 0,01
	HA 2%	9	7,30 \pm 0,01
	HA 1%	9	7,29 \pm 0,01
60 minutes	HA 4%	9	7,31 \pm 0,01
	HA 2%	9	7,30 \pm 0,01
	HA 1%	9	7,29 \pm 0,01

The test results were continued with Wilcoxon non-parametric ($p < 0.05$). The results of further tests using Post-hoc Mann Whitney obtained 1minute group (HA 2%) and 1 minute (HA 1%), 20 minutes group (HA 2%) and 20 minutes (HA 4%) ($p=0.04$), 40 minutes group (HA 2%) and 1 minute (HA 1%), 20 minutes group (HA 2%) and 1 minute (HA 1%), 60 minutes group (HA 1%) and 1 minute (HA 1%), 40 minutes group (HA 1%) and 1 minute (HA 1%) ($p=0, 00$), group 40 min (HA 1%) and 20 min (HA 4%), group 40 min (HA 1%) and 1 min (HA 4%) ($p=0.005$), group 60 min (HA 2%) and 1 min (HA 1%) ($p=0.02$), group 60 minutes (HA 1%) and 60 minutes (HA 4%) ($p=0.03$), group 60 minutes (HA 1%) and 1 minute (HA 4%) ($p=0.01$), and group 60 minutes (HA 1%) and 20 minutes (HA 4%) ($p=0.01$)

significantly different. Based on the results of the material test, it was found that the mixture at concentrations of 4%, 2%, and 1% showed an average value of neutral pH with a range of 7.29-7.31. These results show that the tested HA-EGCG-HPMC material is as expected, which is close to the normal pulp pH of 7.3.

The antibacterial test was conducted in 5 test groups, namely: negative control group (distilled water), 4%HA-EGCG-HPMC, 2%HA 2%-EGCG-HPMC, 1%HA- EGCG-HPMC, and positive control group (4%MTA) samples were mixed with a solution that already contained *Aggregatibacter actinomycetemcomitans* bacteria followed by incubation of samples for 24 hours at 37 °C, then the results were observed by turbidimetric method, and all samples were turbid in color indicating that all samples had bacterial growth, then the samples were applied to MHA agar media to confirm the samples had bacterial growth. The calculation of the number of colonies formed on agar media is presented in Figure 1.




Figure 1. Picture of the number of colonies formed in each sample treatment

The ANOVA test results showed significant differences between the test groups ($p < 0.01$). Further tests were conducted to see the differences between groups. The results of the post-hoc test showed that the number of colonies in the negative group was significantly different from 4%HA-EGCG-HPMC, 2%HA -EGCG-HPMC, 1%HA-EGCG-HPMC, and 4%MTA ($p < 0.01$) with mean colony counts of $1 \times 10^8 \pm 55 \times 10^4$, $2.5 \times 10^7 \pm 28 \times 10^4$, $3.7 \times 10^7 \pm 14 \times 10^4$, $6.4 \times 10^7 \pm 87 \times 10^3$, and $1.3 \times 10^7 \pm 16 \times 10^4$ CFU/mL, respectively. The number of colonies of 4% HA-EGCG-HPMC significantly differed from 1% HA -EGCG-HPMC ($p < 0.001$), which showed that 4% HA-EGCG-HPMC was lower than 1% HA. The number of colonies of 2%HA-EGCG-HPMC was slightly lower than that of 1%HA -EGCG-HPMC ($p < 0.001$) and 4%MTA ($p < 0.001$), 1%HA-EGCG-HPMC group was higher than 4% MTA group ($p < 0.001$) (Figure 2).




Figure 2A-E. Sample test results against the bacteria *A. actinomycetemcomitans* (A. MTA, B. Negative control, C. 1%HA-EGCG-HPMC, D. 2%HA-EGCG-HPMC, E. 4%HA-EGCG-HPMC)

In Figure 2, the density of *A.actinomycetemcomitans* in 4% MTA is the least formed when compared to the control (-), but there is no significant difference in the 4% MTA group with 4% HA-EGCG-HPMC. There was a significant difference in the number of colonies in the 4% HA-EGCG-HPMC group with 1% HA-EGCG-HPMC. Based on the post-hoc test, there is a density of *A. actinomycetemcomitans* in each test sample of the HA-EGCG-HPMC mixture, so it can be concluded that the HA-EGCG-HPMC mixture does not have an antibacterial effect on *A. actinomycetemcomitans* bacteria.

DISCUSSION

The purpose of performing a reconstructed pulp capping is to increase the healing ability of the pulp and promote the formation of a mineralized tissue layer by placing the biomaterial directly on the exposed pulp roof.¹⁹ In this study, a mixture of HA-EGCG-HPMC materials was tested to find alternative materials for pulp capping treatment. First, a characterization test of the hydrogel suspension material with hydroxyapatite, EGCG, and HPMC was carried out, and it was seen that the gelation time was sufficient so that it could be used for clinical applications, whereas slow gelation time made it difficult to apply in a certain area.²⁰ The hydrogel suspension mixture with a concentration of 4% HA-EGCG-HPMC had a gelation time an average of 24-26 minutes, at a concentration of 2% the gelation material at an average time of 27-30 minutes, and at a concentration of 1% the gelation material averaged 29-31 minutes so that the hypothesis was accepted because it had sufficient gelation time for clinical applications, which was under 60 minutes.²¹

In this study, the level of basicity of the HA-EGCG-HPMC material mixture was tested using a pH meter, and samples with three concentration mixtures were shown in Table 5.2. The results show that the material after adding the gelation agent and after the gelation material shows a neutral pH (Table 2), so it is to the pH conditions in the pulp chamber, namely 7.3-7.8.²² According to Elline et al, pH is essential in determining the morphology and structure of a formulation. The pH can influence the ion balance. The increase or decrease of OH⁻ ion, will affect the Ca²⁺, PO₄³⁻, and (HPO₄)²⁻ ions¹⁸

The results of the antibacterial effect test of the HA-EGCG-HPMC mixture using the dilution method of 4%, 2%, and 1% concentrations of mixed materials mixed with *Aggregatibacter actinomycetemcomitans* bacterial suspensions in 96-well plates and incubated at 37°C for 24 hours,

the results showed that all samples contained bacterial growth. Confirmation of bacterial growth and development is achieved by scratching on agar discs to see the growth of the number of bacterial colonies present (Figure 2). Thus, the HA-EGCG-HPMC mixture has no antibacterial effect on *Aggregatibacter actinomycetemcomitans* bacteria. According to a previous study, EGCG would break the bacterial plasma membrane in a certain concentration. For example, the *Streptococcus mutans* plasma membrane will be destroyed with 0,2 mg/ml after EGCG treatment.²³ So, in this study may be the concentration of EGCG has not fulfilled the minimum criteria as a treatment

CONCLUSIONS

Based on the test results, the HA-EGCG-HPMC mixture at a concentration of 4% has a gelation time of 23.78 minutes. It is concluded that this material has a qualification to be used in clinical conditions. It also has a degree of acidity that resembles the vital pulp at 7.30 after gelation. In the antibacterial test results, the 4% HA-EGCG-HPMC mixture had a lower antibacterial effect when compared to 4% MTA. Further research is needed to develop a mixture of HA-EGCG-HPMC materials to increase the ability of antibacterial effects that can support the potential of the material to become a pulp capping material

ACKNOWLEDGMENT

There is financial assistance from FKG Universitas, and the researchers would to acknowledge

REFERENCES

1. Kazemina M, Abdi A, Shohaimi S, Jalali R, Vaisi-Raygani A, Salari N, et al. Dental caries in primary and permanent teeth in children's worldwide, 1995 to 2019: A systematic review and meta-analysis. , Head and Face Medicine. 2020. 16(1);22. Available from : <https://pubmed.ncbi.nlm.nih.gov/33023617/>
2. Karies Gigi Anak di Jakarta Timur P, Putri N, Lusida N, Rosyada D, Ala Al-Maududi A, Studi Kesehatan Masyarakat P, et al. Faktor yang Berhubungan dengan Perilaku Orangtua. 2023. JKKI. 19(1): 11-17. Available from: <https://jurnal.umj.ac.id/index.php/JKKI>
3. Alp Ş, Ulusoy N. Current Approaches in Pulp Capping: A Review. Cyprus J Med Sci. 2024. 28;154–60. Available from : <https://cyprusjmedsci.com/articles/current-approaches-in-pulp-capping-a-review/doi/cjms.2023.2022-37>
4. Darmadi EY, Soesilo D. Antimicrobial activity of calcium hydroxide, calcium oxide, and mineral trioxide aggregate paste against α -Hemolytic Streptococcus. CDJ. 2024 Jul 24;14(1):24–8. Available from : <https://e-journal.unair.ac.id/CDJ/article/view/60848>
5. Ha W, Nicholson T, Kahler B, Walsh L. Rheological Characterization as an Alternative Method to Indentation for Determining the Setting Time of Restorative and Endodontic Cements. Materials. 2017.20;10(12):1451. Available from : https://www.researchgate.net/publication/321959668_Rheological_Characterization_as_an_Alternative_Method_to_Indentation_for_Determining_the_Setting_Time_of_Restorative_and_Endodontic_Cements
6. Covaci A, Ciocan LT, Gălbinașu B, Bucur MV, Matei M, Didilescu AC. Dental Pulp Response to Different Types of Calcium-Based Materials Applied in Deep Carious Lesion Treatment—A Clinical Study. J Funct Biomater. 2022. 2;13(2):51. Available from : <https://www.mdpi.com/2079-4983/13/2/51>
7. Kunert M, Lukomska-Szymanska M. Bio-Inductive Materials in Direct and Indirect Pulp Capping - A Review Article. Materials.2020 (13). Available from : <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7085085/>
8. Eid A, Mancino D, Rekab MS, Haikel Y, Kharouf N. Effectiveness of Three Agents in Pulpotomy Treatment of Permanent Molars with Incomplete Root Development: A

- Randomized Controlled Trial. *Healthcare*. 2022. 25;10(3):431. Available from : <https://pubmed.ncbi.nlm.nih.gov/35326909/>
9. James Irawan I, Adi Wicaksono I, Raya Bandung Sumedang Km J. Review Artikel: Penggunaan Material Bioaktif Dalam Terapi Sel Punca Untuk Karies Gigi. *Open Journal Systems STF Muhammadiyah Cirebon* : ojs.stfmuhammadiyahcirebon.ac.id. 2022.7 Available from : <https://www.semanticscholar.org/reader/e6321d801888edcc04f6b93e87249b548a6fb680>
 10. Niazi FH, Luddin N, Harun MH, Hasan A, Kannan TP, Mohamad S, et al. Dentin–Pulp Complex Response in Molars of Rats after Occlusal and Cervical Restorations with Conventional Glass Ionomer Cement and Nano-Hydroxyapatite Silica Glass Ionomer Cement. *Appl. Sci*. 2023. 1;13(5). Available from : <https://www.mdpi.com/2076-3417/13/5/3156>
 11. Rahmitasari F, Cevanti TA, Prabowo PB, Sularsih S, Prananingrum W. The effect of incorporating hydroxyapatite into Type II glass ionomer cement on flexural strength and the examination of fractured surfaces using scanning electron microscopy. *ODJ*. 2023. 29;10(2):152. Available from : <http://jurnal.unissula.ac.id/index.php/odj/article/viewFile/33491/9378>
 12. Elline E, Ismiyatin K. Nanohydroxiapatite Using Chicken Eggshell Waste and Its Characterization. *MJMHS*. 2021.17. Available from : https://medic.upm.edu.my/upload/dokumen/2021092200460413_2021_0338.pdf
 13. Li Y, Zhao Y, Han J, Wang Y, Lei S. Effects of epigallocatechin gallate (EGCG) on the biological properties of human dental pulp stem cells and inflammatory pulp tissue. *Arch Oral Biol*. 2021. 123:105034. Available from : <https://www.sciencedirect.com/science/article/abs/pii/S000399692030412X>
 14. Mokra D, Joskova M, Mokry J. Therapeutic Effects of Green Tea Polyphenol (–)-Epigallocatechin-3-Gallate (EGCG) in Relation to Molecular Pathways Controlling Inflammation, Oxidative Stress, and Apoptosis. *Int. J. Mol. Sci*. 2023.24. Available from : <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9820274/>
 15. Amanda HG, Elline E, Fibryanto E. Synthesis and Physical Characterization of Nano-Hydroxyapatite-Collagen-Epigallocatechin-3-Gallate Hydrogel Composite. *JIDA* 2022.17;5(1):7. Available from : https://www.researchgate.net/publication/361867826_Synthesis_and_Physical_Characterization_of_Nano-Hydroxyapatite-Collagen-Epigallocatechin-3-Gallate_Hydrogel_Composite#fullTextFileContent
 16. Santoso JH, Prahasti AE, Elline. Effect Of 4 Wt% Hydroxyapatite Pasta On Dental Email Hardness. *IJKG*. 2024. 21;20(1):46–52. Available from : <https://e-journal.unmas.ac.id/index.php/interdental/article/view/8580>
 17. Paul O, Arora P, Mayer M, Chatterjee S. Inflammation in Periodontal Disease: Possible Link to Vascular Disease. *Front.Physiol*. 2021. 11; 1-11. Available from : <https://pubmed.ncbi.nlm.nih.gov/33519515/>
 18. Elline E, Ismiyatin K, Indah Budhy T, Bhardwaj A. The potential of eggshell hydroxyapatite, collagen, and EGCG (HAP-Col-EGCG) scaffold as a pulp regeneration material. *Saudi Dent. J*. 2022. 1;34(8):715–22. Available from : <https://pubmed.ncbi.nlm.nih.gov/36570587/>
 19. Islam R, Islam MRR, Tanaka T, Alam MK, Ahmed HMA, Sano H. Direct pulp capping procedures – Evidence and practice. *Jpn. Dent. Sci. Rev*. 2023.59:48–61. Available from : <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9985044/>
 20. El-Sayed NS, Kamel S. Polysaccharides-Based Injectable Hydrogels: Preparation, Characteristics, and Biomedical Applications . *J.Colloids Interfaces*. 2022. 6(78). Available from : <https://www.mdpi.com/2504-5377/6/4/78>
 21. Zhang Y, Huang Y. Rational Design of Smart Hydrogels for Biomedical Applications. *Vol. 8, Front. in Chem*. 2021.8;1-13. Available from : <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7889811/>
 22. Singh PH. Theory of Optimum pH for Pulp Vitality: A Clinical and Experimental Theory of Endodontics. 2020. 3;1-3. Available from : https://www.researchgate.net/publication/338402785_Theory_of_Optimum_pH_for_Pulp_Vitality_A_Clinical_and_Experimental_Theory_of_Endodontics
 23. Kong C, Zhang H, Li L, Liu Z. Effects of green tea extract epigallocatechin-3-gallate (EGCG) on oral disease-associated microbes: a review. *Vol. 14, Journal of Oral Microbiology*. Taylor and Francis Ltd.; 2022. 14;1-14. Available from : <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9542882/>

ORIGINALITY REPORT

17%	12%	12%	3%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	www.ncbi.nlm.nih.gov Internet Source	3%
2	M. Andi Chandra, Ilham Kunchahyo, Ana Indrayati. "Optimization of Quercetin Gel Formulation using Factorial Design Method and Antibacterial Test against Propionibacterium acnes", Borneo Journal of Pharmacy, 2022 Publication	2%
3	Submitted to University of Wollongong Student Paper	2%
4	www.mdpi.com Internet Source	1%
5	Elline Elline, Kun Ismiyatin, Theresia Indah Budhy, Anuj Bhardwaj. "The Potential of Eggshell Hydroxyapatite, Collagen, and EGCG (HAp-Col-EGCG) Scaffold as A Pulp Regeneration Material", The Saudi Dental Journal, 2022 Publication	1%
6	Juha Lilja, Teri Ridley, George M.W. Cronkite, Hermann J. Enzenhofer, John A. Holmes. "Optimizing sampling effort within a systematic design for estimating abundant escapement of sockeye salmon (Oncorhynchus nerka) in their natal river", Fisheries Research, 2008 Publication	1%

7	doaj.org Internet Source	1 %
8	repository.unair.ac.id Internet Source	1 %
9	e-journal.unair.ac.id Internet Source	1 %
10	He Liu, Jing Lu, Qianzhou Jiang, Markus Haapasalo, Junrong Qian, Franklin R. Tay, Ya Shen. "Biomaterial scaffolds for clinical procedures in endodontic regeneration", <i>Bioactive Materials</i> , 2022 Publication	1 %
11	jcdronline.org Internet Source	1 %
12	Gönül Yenilmez Çiftçi, Gizem Demir Demirel, Gulseren Turhal, Büşra Demirkan, Asuman Demiroglu-Zergeroglu. "2-Aminoanthraquinone substituted cyclotriphosphazene derivatives and their effect on non-small cell lung cancer cell lines", <i>Inorganica Chimica Acta</i> , 2023 Publication	<1 %
13	Issei Noda, Rio Goto, Shunpei Yamashita, Kengo Kawanishi, Masahiro Tsutsumi, Shintarou Kudo. "Assessment of the ulnar nerve stiffness during valgus stress of the elbow using shear wave elastography: an observational cross-sectional study", <i>Journal of Ultrasound</i> , 2024 Publication	<1 %
14	ejournal.upi.edu Internet Source	<1 %
15	www.ektodermaldisplazi.com Internet Source	

<1 %

16

etd.repository.ugm.ac.id

Internet Source

<1 %

17

irep.iium.edu.my

Internet Source

<1 %

18

Gamal Abdel Nasser Atia, Sara Z. Mohamed, Hamdy Abdel Halim, Mohamed M. Ghobashy et al. "Advances in Bioceramic silicates for therapeutic, and regenerative Dentofacial reconstruction", *Ceramics International*, 2024

Publication

<1 %

19

Yingying Bai, Lishan Wang, Rong Xu, Ying Cui. "Mesenchymal stem cells with p38 mitogen-activated protein kinase interference ameliorate mouse ischemic stroke", *Experimental Biology and Medicine*, 2023

Publication

<1 %

20

archive.org

Internet Source

<1 %

21

www.ijmm.org

Internet Source

<1 %

22

d2v96fxpocvxx.cloudfront.net

Internet Source

<1 %

23

eprints.covenantuniversity.edu.ng

Internet Source

<1 %

24

potravinarstvo.com

Internet Source

<1 %

25

repository.unsoed.ac.id

Internet Source

<1 %

26

sciendo.com

Internet Source

<1 %

27

Thomas Reilly, Jan Cabri, Duarte Araújo.
"Science and Football V - The Proceedings of
the Fifth World Congress on Sports Science
and Football", Routledge, 2019

Publication

<1%

28

www.frontiersin.org

Internet Source

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

FINAL GRADE

GENERAL COMMENTS

/100

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8
