



A CSIR Publication

Indian Journal of Biochemistry and Biophysics

www.niscpr.res.in



CSIR-National Institute of Science
Communication And Policy Research
New Delhi, INDIA

Editorial Team

Editor in Chief

Mr. Stefan Dimitrov, Chromatin Biology and Epigenetics Team Head, Izmir Biomedicine and Genome Center, Dokuz Eylul University Health Campus, Izmir 35330, Turkey

Executive Editor

Prof. D Narasimha Rao, Honorary Professor, Department of Biochemistry, Biological Sciences Building, Indian Institute of Science (IISc), Bengaluru-560 012, Karnataka, India

Subject Editors

Mr. Sarfraz Ahmad, Director of Clinical Research, Gynecologic Oncology, AdventHealth Cancer Institute, Florida State University, University of Central Florida, Colleges of Medicine, Orlando, FL-32804, USA, India

Mr. TP Singh, SERB Distinguished Fellow, All India Institute of Medical Sciences (AIIMS), New Delhi-110 029, Delhi, India

Prof. Ryoya Takahashi, Department of Biochemistry, Faculty of Pharmaceutical Sciences, Toho University, Chiba 234-8510, Japan

Mr. Tapas K. Kundu, Director, CSIR-Central Drug Research Institute, (CDRI), Lucknow-226 031, Uttar Pradesh, India

Prof. Sathees C Raghavan, Department of Biochemistry, Biological Sciences Building, Indian Institute of Science (IISc), Bengaluru-560 012, Karnataka, India

Prof. DN Rao, Former Prof. & Head, Department of Biochemistry, All India Institute of Medical Sciences (AIIMS), New Delhi-110 029, Delhi, India

Dr. Vinay K Nandicoori, Director, CSIR-Centre for Cellular & Molecular Biology (CCMB), Hyderabad-500 007, Telangana, India

Mr. KRS Sambhasivarao, Prof. Vice-Chancellor, Mizoram University, Aizawl-796 004, Mizoram, India

Prof. PP Mathur, Vice-Chancellor, Birla Global University, Bhubaneswar-751 029, Odisha, India

Editorial Board

Dr. Bushra Ateeq, Department of Biological Sciences, and Bioengineering, Indian Institute of Technology (IITK), Kanpur-208 016, Uttar Pradesh, India

Prof. Swagata Dasgupta, Department of Chemistry, Associate Dean, Indian Institute of Technology-Kharagpur, Kharagpur-721 302, West Bengal, India

Prof. Rajiv Bhat, School of Biotechnology, Jawaharlal Nehru University (JNU), New Delhi-110 067, Delhi, India

Dr. M Balasubramanyam, Dean of Research Studies Madras, Diabetes Research Foundation (MDRF), Chennai-600 086, Tamil Nadu, India

Prof. Ramagopal Uppaluri, Department of Chemical Engineering, Indian Institute of Technology (IITG), Guwahati-781 039, Assam, India

Prof. M J Swamy, School of Chemistry, Hyderabad Central University, Hyderabad-500 046, Telangana, India

Dr. Ravishankar R, Molecular and Structural Biology, CSIR-Central Drug Research Institute (CDRI), Lucknow-226 031, Uttar Pradesh, India

Prof. Udaykumar Ranga, Molecular Biology and Genetics Unit (MBGU), HIV-AIDS Laboratory, Molecular Biology & Genetics Unit, Jawaharlal Nehru Center for Advanced Scientific Research (JNCASR), Jakkur, Bengaluru-560 064, Karnataka, India

Prof. Roop Mallik, Department of Chemical Sciences, Tata Institute of Fundamental Research (TIFR), Mumbai- 400 005, Maharashtra, India

Prof. Kaiser Jamil, Dean and Director, School of Life Sciences, Centre for Biotechnology and Bioinformatics, Jawaharlal Nehru Institute of Advanced Studies (JNIAS), Hyderabad-500 003, Telangana, India

Prof. Debi Sarkar, Department of Biochemistry, University of Delhi, South Campus, New Delhi-110 021, Delhi, India

Mr. Punyasloke Bhadury, Department of Biological Sciences, Indian Institute of Science Education and Research (IISER), Kolkata-741 246, West Bengal, India

Dr. Mithilesh Mishra, Department of Biological Sciences, Tata Institute of Fundamental Research (TIFR), Mumbai- 400 005, Maharashtra, India

Prof. Manju Bansal, Molecular Biophysics Unit, Indian Institute of Science, Bangalore-560 012, India

Prof. Chandrima Das, Saha Institute of Nuclear Physics (SINP), Kolkata-700 064, West Bengal, India

Mr. Punit Kaur, Head, Department of Biophysics All India Institute of Medical Sciences, (AIIMS) New Delhi-110 029, Delhi, India

Dr. Venkata Dasu Veeranki, Indian Institute of Technology(IIT) Guwahati Guwahati 781039 Assam, India, India

Prof. Subir Kumar Das, College of Medicine & JNM Hospital WBUHS, Kalyani, Nadia 741235, West Bengal.

Prof. N S Puneekar, Department of Biosciences and Bioengineering, Indian Institute of Technology Bombay, Mumbai-400 076, India

Scientific Editor

Dr. N K Prasanna, CSIR-NIScPR

Vol. 60 No. 10: October 2023

Published: 2023-09-27

Papers

Eleutherine bulbosa bulb extract induces apoptosis and inhibits cell migration by downregulating Sonic hedgehog in human tongue cancer cells: An in vitro study

Ferry Sandra, Kelvin Suryajaya, Angliana Chouw, Maria Celinna, Dewi Ranggaini, Kyung Hoon Lee, Cynthia Retna Sartika

763-769

Fabrication of phyco-functionalized zinc oxide nanoparticles and their in vitro evaluation against bacteria and cancer cell line

Geethalakshmi Ramakrishnan, Sirajunnisa Abdul Razack, Logeswari Ravi, Renganathan Sahadevan

770-778

Proteomic analysis of Substantia nigra proteins in STZ-induced Type-II diabetic rats: A possible link with Parkinson's disease

Sanaullah Khan, Habibur Rehman, Kaleemullah Mandokhail, Niamatullah Kakar, Asmatullah Khan, Jameelur Rehman, Fazalur Rehman, Qudratullah, Syed Zahid Ali Shah, Mujeebur Rehman

779-789

Administration of Curcumin, Betanin, and CoQ10 combined with nickel oxide, iron superoxide nanoparticles show preventive effects in breast cancer: Effect on apoptosis pathway and MiR-455 expression

Ghazaleh Ganjipour, Masomeh Heshmati, Mehrdad Hashemi, Maliheh Entezari

790-802

Comparative study of anti-diabetic effects of insulin, epigenin and Salvia mirzayanii extract in streptozotocin-induced diabetic male rats

Rahman Mahdizadehdehosta, Hamid Shahbazmohammadi, Soheila Moein, Neptun Soltani, Mahmoodreza Moein

803-809

Eleutherine bulbosa bulb extract induces apoptosis and inhibits cell migration by downregulating Sonic hedgehog in human tongue cancer cells: An *in vitro* study

Ferry Sandra^{1*}, Kelvin Suryajaya², Angliana Chow³, Maria Celinna⁴, Dewi Ranggaini⁵, Kyung Hoon Lee⁶ & Cynthia Retna Sartika³

¹Department of Biochemistry and Molecular Biology, Division of Oral Biology, Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia

²Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia

³PT Prodia StemCell Indonesia, Jakarta Pusat-10430, Jakarta, Indonesia

⁴The Prodia Education and Research Institute, Jakarta Pusat-10430, Jakarta, Indonesia

⁵Department of Physiology, Division of Oral Biology, Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia

⁶Research Institute, Ballys Co. Ltd, Incheon-22219, Republic of Korea

Received 09 May 2023; revised 18 September 2023

Eleutherine bulbosa (Mill.) Urb. is a medicinal plant which has long been used to treat cancer. *E. bulbosa* bulb extract (EBBE) has been reported to show cytotoxicity towards several types of human cancer. However, the cytotoxic effect of EBBE towards tongue cancer cells has not been investigated. The present study aimed to evaluate the effects of EBBE towards the viability, apoptosis, and migratory activities of tongue cancer cells. Human oral squamous cell carcinoma (HSC)-3 cells were treated with various concentrations of EBBE for 24 h. The number of viable and apoptotic HSC-3 cells were measured using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) and sub-G1 assay, respectively. The migratory activities of HSC-3 cells were assessed using scratch and transwell assay. Sonic hedgehog (SHH) expression was measured using immunoblotting. Upon EBBE treatment, apoptotic HSC-3 cells were significantly higher in a concentration-dependent manner ($P < 0.05$). Meanwhile, viability, migratory activities, and migrated HSC-3 cell number were significantly lower in a concentration-dependent manner ($P < 0.05$). The SHH expression levels in EBBE-treated HSC-3 cells were also lower in a concentration-dependent manner. EBBE reduces HSC-3 cell viability through apoptosis and inhibits its migratory activities by downregulating SHH expression.

Keywords: Apoptosis, Cell migration, *Eleutherine bulbosa*, Sonic hedgehog, Tongue cancer

Oral squamous cell carcinoma (OSCC) is the most common histological type of oral cancer¹ which arises from squamous epithelial cells in the oral cavity. This cancer generally affects the floor of the mouth and the tongue². Various modes of treatment are developed to manage OSCC, including chemotherapy, radiotherapy, surgery, or its combination³. However, these treatments cause numerous side effects.

Discovery and development of new therapeutic agents derived from natural sources, particularly plants with less side effects are recently one of the main focuses on oncology research. Several plants have been reported to have anticancer effects, including

*Brucea javanica*⁴, *Cinnamomum cassia*⁵, *Myristica fragrans*⁶, and *Oroxylum indicum*⁷. Plants have been known as rich sources of various active compounds, including flavonoids, which was reported to inhibit cancer proliferation *via* cell cycle arrest, antiproliferative and antioxidative activity, apoptosis induction, promotion of differentiation, inhibition of angiogenesis and carcinogen metabolic activation, and multidrug resistance modulation⁸. In addition, several non-flavonoid phenolic compounds have also been reported to show anticancer properties, such as caffeic acid⁹.

Eleutherine bulbosa (Mill.) Urb. is a traditional medicinal plant belonging to the Iridaceae family. Its bulbs are commonly used for the treatment of breast cancer, hypertension, stroke, and diabetes mellitus. It has also been used to increase the production of breast milk and treat sexual disorder¹⁰. *E. bulbosa* bulbs have

*Correspondence:
Phone: +628128077780
E-mail: ferry@trisakti.ac.id

also been reported to have antimicrobial¹¹ and antioxidant¹² activity.

Eleutherine bulbosa contains a wide variety of compounds, primarily naphthoquinones, anthraquinones, and naphthalene¹³. The main compounds found in this plant are eleutherin, isoeleutherin, and eleutherol¹⁴. Several compounds obtained from *E. bulbosa* bulbs have been reported to show anticancer activities, such as 6,8-hydroxy-3,4-di-methoxy-1-methyl-anthraquinone-2-carboxylic acid methyl ester¹⁵, eleutherinoside C, and isoeleutherin¹⁶.

Eleutherine bulbosa bulb extract (EBBE) has been reported to show cytotoxicity towards several types of human cancer, such as cervical cancer (line HeLa)¹⁷, colon cancer (line SW480, HCT116, DLD1, WiDr)^{16,18}, breast cancer (line T47D)¹⁹, leukemia (line K562)¹⁵, and retinoblastoma (WERI-Rb-1)²⁰. However, the cytotoxic effect of EBBE towards human tongue cancer cells has not been investigated. Therefore, the present study was conducted to evaluate the effects of EBBE towards the viability, apoptosis, and migratory activities of tongue cancer cells.

Materials and Methods

E. bulbosa bulb extract (EBBE) preparation

E. bulbosa bulbs were obtained from Indonesian Medicinal and Aromatic Crops Research Institute (IMACRI), Indonesia. Plant identification was carried out by expert botanist at Research Center for Plant Conservation and Botanic Gardens, Indonesian Institute of Sciences, Indonesia (No. B-1269/IPH.3/KS/X/2020). EBBE was extracted by maceration method⁴. Briefly, bulbs of *E. bulbosa* were minced and dried. The dried material was extracted with ethanol 70%, filtered and evaporated using a rotary evaporator. The resulting crude EBBE was then stored at 4°C.

HSC-3 cell culture

HSC-3 cell culture was performed as previously described⁴ with modification. The HSC-3 cell line was purchased from Sigma-Aldrich Pte. Ltd. (St. Louis, MO, USA). HSC-3 cells were cultured in a complete medium containing Dulbecco's modified Eagle's medium (DMEM) (Sigma-Aldrich), 10% fetal bovine serum (FBS) (PAN-Biotech, Aidenbach, Germany), 50 U/mL penicillin and 50 µg/mL streptomycin (Sigma-Aldrich) in a humidified 5% CO₂, 37°C incubator. HSC-3 cells were then dissociated with trypsin-ethylenediamine tetraacetic acid (EDTA) solution (Sigma-Aldrich) after reaching

80% confluency. Upon reaching the desired cell number, HSC-3 cells were seeded into 24-well plate and 96-well plate for further assays.

Cell viability assay

The number of viable cells were measured quantitatively using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay as previously described²¹ with modification. Briefly, 5×10^3 HSC-3 cells were seeded into 96-well plates and then treated with EBBE (1, 10, or 100 µg/mL), 1 µM Doxorubicin (Dankos Farma, Jakarta, Indonesia) or medium merely for 24 h. After that, 100 µL of MTT (Sigma-Aldrich) in culture medium was loaded into each well. After incubating the plates for 4 h, culture medium was discarded, and the formed formazan crystals were dissolved in 100 µL dimethylsulfoxide (DMSO). Results were measured at OD₅₇₀ using a microplate reader (Bio-Rad, Hercules, CA, USA). Each experimental group was measured in six replicates. In addition, the number of untreated cells were counted with hemacytometer and used for interpolating OD₅₇₀ values of HSC-3 cells.

Sub-G1 assay

To investigate the cytotoxic effect of EBBE on HSC-3 cells, the number of apoptotic cells were determined using sub-G1 assay as previously described⁹ with modification. Treated and untreated HSC-3 cells were harvested and suspended in 450 µL hypotonic fluorochrome solution containing 50 µg/mL propidium iodide (Sigma-Aldrich), 0.1% Triton X-100 (Sigma-Aldrich), and 0.1% sodium citrate (Wako, Osaka, Japan). After that, cell suspensions were incubated in the dark for 2 h at room temperature. Fluorescence of individual nuclei was measured using FACSCanto II flow cytometer (Becton Dickinson, Franklin Lakes, NJ, USA) at 100,000 events.

Scratch assay

Scratch assay was performed as previously described²². HSC-3 cells (5×10^4) were seeded into 24-well plates and incubated overnight. Scratch was made on each well with micropipette yellow tips. Then, the wells were washed with phosphate-buffered saline (PBS) (Wako) and treated with/without EBBE or 20 µM cyclopamine tartrate (BioVision, Milpitas, CA, USA) (as positive control) for 24 h. The gap closures were observed and documented under an inverted light microscope (Carl Zeiss, Jena, Germany).

Transwell assay

Transwell assay was performed as previously described²³ with modification. HSC-3 cells (5×10^4) were seeded in the 8 μm pore size chamber insert (Merck, Darmstadt, Germany). After that, 800 μL culture medium with/without EBBE or 20 μM cyclopamine tartrate was loaded into each well of the 24-well plate. HSC-3-containing chamber inserts were then placed on the 24-well plate. After incubating the chamber for 24 h at 37°C, migrated cells in the lower surface of the chamber were fixed. The cells were counted and documented under an inverted light microscope.

Immunoblotting

Expression levels of Sonic hedgehog (SHH) protein were measured using immunoblotting method as previously described²⁴ with modification. HSC-3 cells were lysed and homogenized with Laemmli sample buffer (Bio-Rad). The proteins from each experimental group were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and transferred to a polyvinylidene difluoride (PVDF) sheet. After blocking with 5% skim milk solution, the sheet was incubated with rabbit monoclonal anti-SHH antibody as a primary antibody and horseradish peroxidase (HRP)-conjugated goat anti-rabbit antibody as a secondary antibody (Cell Signaling Technology, Danvers, MA, USA). Immunoblot results were analyzed using ImageJ software (National Institute of Health, Bethesda, MD, USA) to measure the density of each SHH band.

Statistical analysis

Statistical analyses were performed using IBM SPSS Statistics version 26.0 (SPSS IBM, Armonk, NY, USA). Shapiro-Wilk test was used as a normality test. Differences of MTT assay, sub-G1 assay, and scratch assay results between experimental groups were analyzed using Kruskal-Wallis H with Mann-Whitney U *post hoc* analysis. Meanwhile, differences of transwell assay results and SHH band density between experimental groups were analyzed using one-way analysis of variances (ANOVA) with Tukey's honestly significant difference (HSD) *post hoc* analysis. All results were expressed as mean \pm standard deviation (SD). $P < 0.05$ were considered as statistically significant.

Results

EBBE reduced the number of viable HSC-3 cells

The number of viable HSC-3 cells cultured in medium containing extract solvent (sham group) for

24 h was $9,607 \pm 13.81$. Treatment with EBBE showed that the viability of HSC-3 cells was significantly lower compared compared with that of the sham group ($P < 0.05$) in a concentration-dependent manner. The number of viable HSC-3 cells treated with 1, 10, and 100 $\mu\text{g/mL}$ EBBE were $9,055 \pm 268.69$, $6,777 \pm 278.04$, and $3,659 \pm 32.86$, respectively. The viability of EBBE-treated HSC-3 cells was significantly higher ($P < 0.05$) than that of Doxorubicin-treated HSC-3 cells (226 ± 32.34) (Fig. 1). These findings suggested that EBBE could reduce the viability of HSC-3 cells.

EBBE induced apoptosis of HSC-3 cells

Sub-G1 assay results showed that the apoptosis percentage of the sham group was $8.27 \pm 0.14\%$. Apoptotic HSC-3 cells were significantly higher compared with the sham group ($P < 0.05$) in concentration-dependent manner after EBBE treatment. Apoptosis percentages of HSC-3 cells treated with 1, 10, and 100 $\mu\text{g/mL}$ EBBE were $9.60 \pm 0.35\%$, $17.78 \pm 0.32\%$, and $38.64 \pm 0.70\%$, respectively. These percentages were significantly lower ($P < 0.05$) than that of Doxorubicin-treated HSC-3 cells ($95.95 \pm 2.60\%$) (Fig. 2). Therefore, these results suggested that EBBE treatment could promote apoptosis of HSC-3 cells.

EBBE inhibited HSC-3 cell migration

The gap closure percentage of HSC-3 cells cultured in medium containing extract solvent after 24 h was




Fig. 1 — EBBE decreased the number of viable HSC-3 cells in a concentration-dependent manner. HSC-3 cells were cultured and treated with/without 1 μM Doxorubicin or various concentrations of EBBE for 24 h as indicated in the panel. The number of viable cells were measured with MTT assay as described in Materials and Methods. The data are expressed as mean \pm SD ($n=6$). * $P < 0.05$ compared to the sham group




Fig. 2 — EBBE increased the percentage of apoptotic HSC-3 cells in a concentration-dependent manner. HSC-3 cells were cultured and treated with/without 1 μM Doxorubicin or various concentrations of EBBE for 24 h as indicated. Treated cells were then subjected to sub-G1 assay as described in Materials and Methods. (A) Histogram of flow cytometric results. (B) Percentage of apoptotic HSC-3 cells. The data are expressed as mean \pm SD (n=18). * $P < 0.05$ compared to the sham group

96.11 \pm 4.17%. After EBBE treatment, the migratory activities of HSC-3 cells were significantly lower compared with that of the sham group ($P < 0.05$) in a concentration-dependent manner. Gap closure percentages of HSC-3 cells treated with 1, 10, and 100 $\mu\text{g/mL}$ EBBE were 89.56 \pm 5.77%, 76.78 \pm 4.87%, and 60.67 \pm 7.62%, respectively. These percentages were significantly higher ($P < 0.05$) than that of cyclopamine tartrate-treated HSC-3 cells (43.00 \pm 4.82%) (Fig. 3).

Transwell assay also showed similar results. The number of migrated HSC-3 cells in the sham group was 134.44 \pm 11.60. The number of migrated HSC-3 cells was significantly lower than that of the sham group ($P < 0.05$) in a concentration-dependent manner after EBBE treatment. The number of migrated HSC-3 cells treated with 1, 10, and 100 $\mu\text{g/mL}$ EBBE were 107.44 \pm 15.21, 95.11 \pm 6.01, and 39.22 \pm 3.56, respectively. These numbers were significantly higher ($P < 0.05$) than that of cyclopamine tartrate-treated HSC-3 cells (17.89 \pm 3.22) (Fig. 4). Overall, these

results indicated that EBBE exhibited an anti-migratory effect towards HSC-3 cells.

EBBE downregulated the expression of SHH

Upon the addition of EBBE, the SHH expression levels in HSC-3 cells were lower in a concentration-dependent manner. Expression level of SHH in HSC-3 cells treated with 1 $\mu\text{g/mL}$ EBBE was significantly higher ($P < 0.05$), while SHH expression levels in HSC-3 cells treated with 10 and 100 $\mu\text{g/mL}$ EBBE were significantly lower ($P < 0.05$) compared with that in the sham group. SHH expression levels in HSC-3 cells treated with 1 and 10 $\mu\text{g/mL}$ EBBE were significantly higher compared with that in cyclopamine tartrate-treated HSC-3 cells ($P < 0.05$). However, no significant difference was observed in the SHH expression levels between cyclopamine tartrate and 100 $\mu\text{g/mL}$ EBBE treatment ($P = 0.142$) (Fig. 5). Thus, these findings suggested that EBBE could decrease the expression of SHH.




Fig. 3 — EBBE inhibited migratory activities of HSC-3 cells. HSC-3 cell culture was scratched and treated with/without EBBE or 20 µM cyclopamine tartrate for 24 h as indicated. Gap closure was measured as described in Materials and Methods. (A) Scratched areas of HSC-3 cell culture. White bar: 100 µM. (B) Gap closure percentage of EBBE-treated HSC-3 cell. The data are expressed as mean ± SD (n=9). **P* <0.05 compared to the sham group




Fig. 4 — EBBE reduced the number of migrated HSC-3 cells. HSC-3 cell culture was treated with/without EBBE or 20 µM cyclopamine tartrate for 24 h as indicated. Migrated cells were measured as described in Materials and Methods. The data are expressed as mean ± SD (n=9). **P* <0.05 compared to the sham group




Fig. 5 — EBBE reduced the expression of SHH. HSC-3 cell culture was treated with/without EBBE or 20 µM cyclopamine tartrate for 24 h as indicated. HSC-3 cells were lysed, homogenized, and immunoblotted using anti-SHH antibody as described in Materials and Methods. The data are expressed as mean ± SD (n=3). **P* <0.05 compared to the sham group

Discussion

In the present study, we investigated the potency of EBBE as anticancer agent for OSCC treatment. EBBE showed a cytotoxic effect towards HSC-3 cells by reducing the viability of HSC-3 cells in a concentration-dependent manner. According to the sub-G1 assay results, the decrease in HSC-3 cell viability was caused by apoptosis. These results were in line with previous studies that EBBE was able to induce apoptosis of cancer cells^{18-20,25}. EBBE was also reported to inhibit cell cycle in several cell lines^{19,25}, indicating that EBBE may induce apoptosis of HSC-3 cells through cell cycle arrest since both processes are interconnected²⁶. Based on a molecular docking study, eleutherol, eleutherin, and isoeleutherin could induce apoptosis by activating caspase-8²⁷.

Cell migration is one of the key processes involved in cancer metastasis²⁸. Scratch and transwell assays demonstrated that EBBE restrained the migration of HSC-3 cells in a concentration-dependent manner. This showed the novel activity of EBBE in impairing HSC-3 cell migration. Therefore, EBBE may have anti-metastasis activity towards tongue cancer cells.

Metastasis of tongue cancer is regulated by several proteins, including SHH. One of the SHH signaling pathways that has been reported to regulate cell migration in tongue cancer is SHH/glioma-associated oncogene homologue (SHH/GLI)²⁹. SHH protein binds and inactivates its receptor, Patched 1 (PTCH1). This interaction causes the activation of transcription factors from the GLI family, which play a major role in regulating cancer migration, invasion, and proliferation³⁰.

The present study revealed that EBBE decreased SHH expression levels along with the increase in EBBE concentrations, suggesting that the inhibitory effect of EBBE towards the expression of SHH may be responsible for its anti-migratory property. To the best of our knowledge, this is the first study demonstrating SHH downregulation in EBBE-treated HSC-3 cells. EBBE may inhibit HSC-3 cell migration via inhibition of SHH/GLI signaling pathway. Possibly, all signaling pathways that are regulated by SHH can be inhibited by EBBE. EBBE may also restrain HSC-3 cell migration through another pathway. In SW480 colon cancer cell line, eleutherinoside C and isoeleutherin obtained from *E. bulbosa* bulbs was reported to inhibit Wnt/ β -catenin pathway¹⁶, which has been reported to play an important role in regulating cancer migration³¹.

Therefore, signaling pathways involved in the inhibition of HSC-3 cell migration by EBBE still need to be explored. In addition, signaling components downstream to SHH and target molecules that are affected by EBBE should be identified. Since EBBE also contains various bioactive compounds, further research is needed to identify which compounds are responsible for apoptosis and anti-migratory activity. Thus, more investigations are required to understand the molecular mechanisms underlying EBBE-induced HSC-3 cell apoptosis and confirm which SHH signaling pathway contributes to the inhibition of cell migration by EBBE.

Conclusion

EBBE reduces HSC-3 cell viability through apoptosis and inhibits its migratory activities by downregulating SHH expression. Taken together, EBBE could be a potential anti-tongue cancer agent, particularly as an antimetastatic agent.

Conflict of interest

All authors declare no conflicts of interest.

References

- 1 Chen YK, Huang HC, Lin LM & Lin CC, Primary oral squamous cell carcinoma: An analysis of 703 cases in southern Taiwan. *Oral Oncol*, 35 (1999) 173.
- 2 Neville BW & Day TA, Oral cancer and precancerous lesions. *CA Cancer J Clin*, 52 (2002) 195.
- 3 Minhas S, Kashif M, Altaf W, Afzal N & Nagi AH, Concomitant-chemoradiotherapy-associated oral lesions in patients with oral squamous-cell carcinoma. *Cancer Biol Med*, 14 (2017) 176.
- 4 Wicaksono BD, Tangkearung E & Sandra F, *Brucea javanica* leaf extract induced apoptosis in human oral squamous cell carcinoma (HSC2) cells by attenuation of mitochondrial membrane permeability. *Indones Biomed J*, 7 (2015) 107.
- 5 Chang WL, Cheng FC, Wang SP, Chou ST & Shih Y, *Cinnamomum cassia* essential oil and its major constituent cinnamaldehyde induced cell cycle arrest and apoptosis in human oral squamous cell carcinoma HSC-3 cells. *Environ Toxicol*, 32 (2017) 456.
- 6 Susianti S, Lesmana R, Salam S, Julaha E, Pratiwi YS, Sylviana N, Goenawan H, Kurniawan A & Supratman U, The effect of nutmeg seed (*M. fragrans*) extracts induces apoptosis in melanoma maligna cell's (B16-F10). *Indones Biomed J*, 13 (2021) 68.
- 7 Poonacha SK, Harishkumar M, Radha M, Varadarajan R, Nalilu SK, Shetty SS, Shetty PK, Chandrashekarappa RB, Sreenivas MG & Bavabeedu SKB, Insight into oroxylinA-7-O- β -d-glucuronide-enriched *Oroxylum indicum* bark extract in oral cancer HSC-3 cell apoptotic mechanism: Role of mitochondrial microenvironment. *Molecules*, 26 (2021) 7430.
- 8 Ren W, Qiao Z, Wang H, Zhu L & Zhang L, Flavonoids: Promising anticancer agents. *Med Res Rev*, 23 (2003) 519.

- 9 Sandra F & Sidharta MA, Caffeic acid induced apoptosis in MG63 osteosarcoma cells through activation of caspases. *Mol Cell Biomed Sci*, 1 (2017) 28.
- 10 Ieyama T, Gunawan-Puteri MDPT & Kawabata J, α -Glucosidase inhibitors from the bulb of *Eleutherine americana*. *Food Chem*, 128 (2011) 308.
- 11 Subramaniam K, Suriyamoorthy S, Wahab F, Sharon FB & Rex GR, Antagonistic activity of *Eleutherine palmifolia* Linn. *Asian Pac J Trop Dis*, 2 (2012) S491.
- 12 Kuntorini EM, Dewi M & Misrina M, Anatomical structure and antioxidant activity of red bulb plant (*Eleutherine americana*) on different plant age. *Biodiversitas*, 17 (2016) 229.
- 13 Insanu M, Kusmardiyani S & Hartati R, Recent studies on phytochemicals and pharmacological effects of *Eleutherine americana* Merr. *Procedia Chem*, 13 (2014) 221.
- 14 Chen Z, Huang H, Wang C, Li Y, Ding J, Sankawa U, Noguchi H & Itaka Y, Hongonin, a new naphthalene derivative from hong-cong, the rhizome of *Eleutherine Americana* Merr. et Heyne (Iridaceae). *Chem Pharm Bull*, 34 (1986) 2743.
- 15 Xu J, Qiu F, Duan W, Qu G, Wang N & Yao X, New bioactive constituents from *Eleutherine americana*. *Front Chem China*, 1 (2006) 320.
- 16 Li X, Ohtsuki T, Koyano T, Kowithayakorn T & Ishibashi M, New Wnt/ β -catenin signaling inhibitors isolated from *Eleutherine palmifolia*. *Chem Asian J*, 4 (2009) 540.
- 17 Mutiah R, Minggarwati TS, Kristanti RA & Susanti E, Compound identification and anticancer activity of ethyl acetate fraction from bawang sabrang (*Eleutherine palmifolia* (L.) Merr.) on HeLa cervical cancer cell line. *Indones J Cancer Chemoprevent*, 10 (2019) 131.
- 18 Lubis IA, Ichwan M, Mustofa M & Satria D, Anticancer activity of *Eleutherine bulbosa* (Mill.) Urb. extract on WiDr cell line *in vitro*, paper presented to 2nd Public Health International Conference (PHICo 2017), Faculty of Public Health – Universitas Sumatera Utara, Medan, Indonesia, 18-19 December 2017.
- 19 Yuniarti A, Sundhani E & Nurulita NA, The potentiation effect of bawangdayak (*Sisyrinchium palmifolium* L.) extract on T47D cell growth inhibition after 5-fluorouracil treatment. *Pharmaciana*, 8 (2018) 195.
- 20 Kamarudin AA, Sayuti NH, Saad N, Razak NAA & Esa NM, Induction of apoptosis by *Eleutherine bulbosa* (Mill.) Urb. bulb extracted under optimised extraction condition on human retinoblastoma cancer cells (WERI-Rb-1). *J Ethnopharmacol*, 284 (2022) 114770.
- 21 Sandra F, Sudiono J, Trisfilha P & Pratiwi D, Cytotoxicity of *Alpinia galanga* rhizome crude extract on NIH-3T3 cells. *Indones Biomed J*, 9 (2017) 23.
- 22 Ariesanti Y, Sandra F, Claresta B & Alvita L, *Coffea canephora* bean extract induces NIH3T3 cell migration. *Indones Biomed J*, 13 (2021) 216.
- 23 Oktaviono YH, Al-Farabi MJ, Suastika LOS, Hartono F, Dirgantara Y & Sandra F, Preliminary study: Purple sweet potato extract seems to be superior to increase the migration of impaired endothelial progenitor cells compared to L-Ascorbic acid. *Sci Pharm*, 87 (2019) 16.
- 24 Rizal MI & Sandra F, *Brucea javanica* leaf extract activates caspase-9 and caspase-3 of mitochondrial apoptotic pathway in human oral squamous cell carcinoma. *Indones Biomed J*, 8 (2016) 43.
- 25 Mutiah R, Listiyana A, Suryadinata A, Annisa R, Hakim A, Anggraini W & Susilowati R, Activity of inhibit the cell cycle and induct apoptosis in HeLa cancer cell with combination of sabrang onion (*Eleutherine palmifolia* (L.) Merr) and starfruit mistletoe (*Macrosolen cochinchinensis* (Lour.) Tiegh). *J App Pharm Sci*, 8 (2018) 122.
- 26 Pucci B, Kasten M & Giordano A, Cell cycle and apoptosis. *Neoplasia*, 2 (2000) 291.
- 27 Quadros Gomes AR, da Rocha Galucio NC, de Albuquerque KCO, Brígido HPC, Varela ELP, Castro ALG, Vale VV, Bahia MO, Rodriguez Burbano RM, de Molfeta FA, Carneiro LA, Percario S & Dolabela MF, Toxicity evaluation of *Eleutherine plicata* Herb. extracts and possible cell death mechanism. *Toxicol Rep*, 8 (2021) 1480.
- 28 Ellis IR, The migration and invasion of oral squamous carcinoma cells: Matrix, growth factor and signalling involvement. *Cancers*, 13 (2021) 2633.
- 29 Fan HX, Wang S, Zhao H, Liu N, Chen D, Sun M & Zheng JH, Sonic hedgehog signaling may promote invasion and metastasis of oral squamous cell carcinoma by activating MMP-9 and E-cadherin expression. *Med Oncol*, 31 (2014) 41.
- 30 Carballo GB, Honorato JR, de Lopes GPF & Spohr TCLSE, A highlight on Sonic hedgehog pathway. *Cell Commun Signal*, 16 (2018) 11.
- 31 Liang S, Zhang S, Wang P, Yang C, Shang C, Yang J & Wang J, LncRNA, TUG1 regulates the oral squamous cell carcinoma progression possibly *via* interacting with Wnt/ β -catenin signaling. *Gene*, 608 (2017) 49.



Digital Receipt

This receipt acknowledges that Turnitin received your paper. Below you will find the receipt information regarding your submission.

The first page of your submissions is displayed below.

Submission author: Ferry Sandra
Assignment title: SIJALI 11
Submission title: Eleutherine bulbosa bulb extract induces apoptosis and inhi...
File name: IJBB_83_auth_corr_1.pdf
File size: 9.06M
Page count: 7
Word count: 3,960
Character count: 22,117
Submission date: 17-Jan-2024 11:17AM (UTC+0700)
Submission ID: 2272330930

Indian Journal of Biochemistry & Biophysics
Vol. 60, October 2023, pp. 703-709
DOI: 10.56042/ijbb.v60i10.1293

NIS&PR
Digital Receipt

Eleutherine bulbosa bulb extract induces apoptosis and inhibits cell migration by downregulating Sonic hedgehog in human tongue cancer cells: An *in vitro* study

Ferry Sandra^{1*}, Kelvin Suryajaya², Angliana Chouw³, Maria Celina⁴, Dewi Ranggani⁵, Kyung Hoon Lee⁶ & Cynthia Retna Sartika⁷

¹Department of Biochemistry and Molecular Biology, Division of Oral Biology, Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia
²Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia
³PT Prodia StemCell Indonesia, Jakarta Pusat-10430, Jakarta, Indonesia
⁴The Prodia Education and Research Institute, Jakarta Pusat-10430, Jakarta, Indonesia
⁵Department of Physiology, Division of Oral Biology, Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia
⁶Research Institute, Ballys Co. Ltd, Incheon-22219, Republic of Korea

Received 09 May 2023; revised 18 September 2023

Eleutherine bulbosa (Mill.) Urb. is a medicinal plant which has long been used to treat cancer. *E. bulbosa* bulb extract (EBBE) has been reported to show cytotoxicity towards several types of human cancer. However, the cytotoxic effect of EBBE towards tongue cancer cells has not been investigated. The present study aimed to evaluate the effects of EBBE towards the viability, apoptosis, and migratory activities of tongue cancer cells. Human oral squamous cell carcinoma (HSC-3) cells were treated with various concentrations of EBBE for 24 h. The number of viable and apoptotic HSC-3 cells were measured using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) and sub-G1 assay, respectively. The migratory activities of HSC-3 cells were assessed using scratch and transwell assay. Sonic hedgehog (SHH) expression was measured using immunoblotting. Upon EBBE treatment, apoptotic HSC-3 cells were significantly higher in a concentration-dependent manner ($P < 0.05$). Meanwhile, viability, migratory activities, and migrated HSC-3 cell number were significantly lower in a concentration-dependent manner ($P < 0.05$). The SHH expression levels in EBBE-treated HSC-3 cells were also lower in a concentration-dependent manner. EBBE reduces HSC-3 cell viability through apoptosis and inhibits its migratory activities by downregulating SHH expression.

Keywords: Apoptosis, Cell migration, *Eleutherine bulbosa*, Sonic hedgehog, Tongue cancer

Oral squamous cell carcinoma (OSCC) is the most common histological type of oral cancer¹ which arises from squamous epithelial cells in the oral cavity. This cancer generally affects the floor of the mouth and the tongue². Various modes of treatment are developed to manage OSCC, including chemotherapy, radiotherapy, surgery, or its combination³. However, these treatments cause numerous side effects.

Discovery and development of new therapeutic agents derived from natural sources, particularly plants with less side effects are recently one of the main focuses on oncology research. Several plants have been reported to have anticancer effects, including *Brucea javanica*⁴, *Cinnamomum cassia*⁵, *Myrsine fragrans*⁶, and *Oroxylum indicum*⁷. Plants have been known as rich sources of various active compounds, including flavonoids, which was reported to inhibit cancer proliferation via cell cycle arrest, antiproliferative and antioxidative activity, apoptosis induction, promotion of differentiation, inhibition of angiogenesis and carcinogen metabolic activation, and multidrug resistance modulation⁸. In addition, several non-flavonoid phenolic compounds have also been reported to show anticancer properties, such as caffeic acid⁹.

Eleutherine bulbosa (Mill.) Urb. is a traditional medicinal plant belonging to the Iridaceae family. Its bulbs are commonly used for the treatment of breast cancer, hypertension, stroke, and diabetes mellitus. It has also been used to increase the production of breast milk and treat sexual disorder¹⁰. *E. bulbosa* bulbs have

*Correspondence:
Phone: +629128077780
E-mail: ferry@trisakti.ac.id

Eleutherine bulbosa bulb extract induces apoptosis and inhibits cell migration by downregulating Sonic hedgehog in human tongue cancer cells: An in vitro study

by Ferry Sandra

Submission date: 17-Jan-2024 11:17AM (UTC+0700)

Submission ID: 2272330930

File name: IJBB_83_auth_corr_1.pdf (9.06M)

Word count: 3960

Character count: 22117



Eleutherine bulbosa bulb extract induces apoptosis and inhibits cell migration by downregulating Sonic hedgehog in human tongue cancer cells: An *in vitro* study

Ferry Sandra^{1*}, Kelvin Suryajaya², Angliana Chouw³, Maria Celinna⁴, Dewi Ranggaim⁵, Kyung Hoon Lee⁶ & Cynthia Retna Sartika³

¹Department of Biochemistry and Molecular Biology, Division of Oral Biology, Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia

²Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia

³PT Prodia StemCell Indonesia, Jakarta Pusat-10430, Jakarta, Indonesia

⁴The Prodia Education and Research Institute, Jakarta Pusat-10430, Jakarta, Indonesia

⁵Department of Physiology, Division of Oral Biology, Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia

⁶Research Institute, Ballys Co. Ltd, Incheon-22219, Republic of Korea

Received 09 May 2023; revised 18 September 2023

Eleutherine bulbosa (Mill.) Urb. is a medicinal plant which has long been used to treat cancer. *E. bulbosa* bulb extract (EBBE) has been reported to show cytotoxicity towards several types of human cancer. However, the cytotoxic effect of EBBE towards tongue cancer cells has not been investigated. The present study aimed to evaluate the effects of EBBE towards the viability, apoptosis, and migratory activities of tongue cancer cells. Human oral squamous cell carcinoma (HSC)-3 cells were treated with various concentrations of EBBE for 24 h. The number of viable and apoptotic HSC-3 cells were measured using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) and sub-G1 assay, respectively. The migratory activities of HSC-3 cells were assessed using scratch and transwell assay. Sonic hedgehog (SHH) expression was measured using immunoblotting. Upon EBBE treatment, apoptotic HSC-3 cells were significantly higher in a concentration-dependent manner ($P < 0.05$). Meanwhile, viability, migratory activities, and migrated HSC-3 cell number were significantly lower in a concentration-dependent manner ($P < 0.05$). The SHH expression levels in EBBE-treated HSC-3 cells were also lower in a concentration-dependent manner. EBBE reduces HSC-3 cell viability through apoptosis and inhibits its migratory activities by downregulating SHH expression.

Keywords: Apoptosis, Cell migration, *Eleutherine bulbosa*, Sonic hedgehog, Tongue cancer

Oral squamous cell carcinoma (OSCC) is the most common histological type of oral cancer¹ which arises from squamous epithelial cells in the oral cavity. This cancer generally affects the floor of the mouth and the tongue². Various modes of treatment are developed to manage OSCC, including chemotherapy, radiotherapy, surgery, or its combination³. However, these treatments cause numerous side effects.

Discovery and development of new therapeutic agents derived from natural sources, particularly plants with less side effects are recently one of the main focuses on oncology research. Several plants have been reported to have anticancer effects, including

*Brucea javanica*⁴, *Cinnamomum cassia*⁵, *Myristica fragrans*⁶, and *Oroxylum indicum*⁷. Plants have been known as rich sources of various active compounds, including flavonoids, which was reported to inhibit cancer proliferation *via* cell cycle arrest, antiproliferative and antioxidative activity, apoptosis induction, promotion of differentiation, inhibition of angiogenesis and carcinogen metabolic activation, and multidrug resistance modulation⁸. In addition, several non-flavonoid phenolic compounds have also been reported to show anticancer properties, such as caffeic acid⁹.

Eleutherine bulbosa (Mill.) Urb. is a traditional medicinal plant belonging to the Iridaceae family. Its bulbs are commonly used for the treatment of breast cancer, hypertension, stroke, and diabetes mellitus. It has also been used to increase the production of breast milk and treat sexual disorder¹⁰. *E. bulbosa* bulbs have

*Correspondence:
Phone: +628128077780
E-mail: ferry@trisakti.ac.id

also been reported to have antimicrobial¹¹ and antioxidant¹² activity.

Eleutherine bulbosa contains a wide variety of compounds, primarily naphthoquinones, anthraquinones, and naphthalene¹³. The main compounds found in this plant are eleutherin, isoeleutherin, and eleutherol¹⁴. Several compounds obtained from *E. bulbosa* bulbs have been reported to show anticancer activities, such as 6,8-hydroxy-3,4-di-methoxy-1-methyl-anthraquinone-2-carboxylic acid methyl ester¹⁵, eleutherinoside C, and isoeleutherin¹⁶.

Eleutherine bulbosa bulb extract (EBBE) has been reported to show cytotoxicity towards several types of human cancer, such as cervical cancer (line HeLa)¹⁷, colon cancer (line SW480, HCT116, DLD1, WiDr)^{16,18}, breast cancer (line T47D)¹⁹, leukemia (line K562)¹⁵, and retinoblastoma (WERI-Rb-1)²⁰. However, the cytotoxic effect of EBBE towards human tongue cancer cells has not been investigated. Therefore, the present study was conducted to evaluate the effects of EBBE towards the viability, apoptosis, and migratory activities of tongue cancer cells.

Materials and Methods

E. bulbosa bulb extract (EBBE) preparation

E. bulbosa bulbs were obtained from Indonesian Medicinal and Aromatic Crops Research Institute (IMACRI), Indonesia. Plant identification was carried out by expert botanist at Research Center for Plant Conservation and Botanic Gardens, Indonesian Institute of Sciences, Indonesia (No. B-1269/IPH.3/KS/X/2020). EBBE was extracted by maceration method⁴. Briefly, bulbs of *E. bulbosa* were minced and dried. The dried material was extracted with ethanol 70%, filtered and evaporated using a rotary evaporator. The resulting crude EBBE was then stored at 4°C.

HSC-3 cell culture

HSC-3 cell culture was performed as previously described⁴ with modification. The HSC-3 cell line was purchased from Sigma-Aldrich Pte. Ltd. (St. Louis, MO, USA). HSC-3 cells were cultured in a complete medium containing Dulbecco's modified Eagle's medium (DMEM) (Sigma-Aldrich), 10% fetal bovine serum (FBS) (PAN-Biotech, Aidenbach, Germany), 50 U/mL penicillin and 50 µg/mL streptomycin (Sigma-Aldrich) in a humidified 5% CO₂, 37°C incubator. HSC-3 cells were then dissociated with trypsin-ethylenediamine tetraacetic acid (EDTA) solution (Sigma-Aldrich) after reaching

80% confluency. Upon reaching the desired cell number, HSC-3 cells were seeded into 24-well plate and 96-well plate for further assays.

Cell viability assay

The number of viable cells were measured quantitatively using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay as previously described²¹ with modification. Briefly, 5×10^3 HSC-3 cells were seeded into 96-well plates and then treated with EBBE (1, 10, or 100 µg/mL), 1 µM Doxorubicin (Dankos Farma, Jakarta, Indonesia) or medium merely for 24 h. After that, 100 µL of MTT (Sigma-Aldrich) in culture medium was loaded into each well. After incubating the plates for 4 h, culture medium was discarded, and the formed formazan crystals were dissolved in 100 µL dimethylsulfoxide (DMSO). Results were measured at OD₅₇₀ using a microplate reader (Bio-Rad, Hercules, CA, USA). Each experimental group was measured in six replicates. In addition, the number of untreated cells were counted with hemacytometer and used for interpolating OD₅₇₀ values of HSC-3 cells.

Sub-G1 assay

To investigate the cytotoxic effect of EBBE on HSC-3 cells, the number of apoptotic cells were determined using sub-G1 assay as previously described⁹ with modification. Treated and untreated HSC-3 cells were harvested and suspended in 450 µL hypotonic fluorochrome solution containing 50 µg/mL propidium iodide (Sigma-Aldrich), 0.1% Triton X-100 (Sigma-Aldrich), and 0.1% sodium citrate (Wako, Osaka, Japan). After that, cell suspensions were incubated in the dark for 2 h at room temperature. Fluorescence of individual nuclei was measured using FACSCanto II flow cytometer (Becton Dickinson, Franklin Lakes, NJ, USA) at 100,000 events.

Scratch assay

Scratch assay was performed as previously described²². HSC-3 cells (5×10^4) were seeded into 24-well plates and incubated overnight. Scratch was made on each well with micropipette yellow tips. Then, the wells were washed with phosphate-buffered saline (PBS) (Wako) and treated with/without EBBE or 20 µM cyclopamine tartrate (BioVision, Milpitas, CA, USA) (as positive control) for 24 h. The gap closures were observed and documented under an inverted light microscope (Carl Zeiss, Jena, Germany).

Transwell assay

Transwell assay was performed as previously described²³ with modification. HSC-3 cells (5×10^4) were seeded in the 8 μm pore size chamber insert (Merck, Darmstadt, Germany). After that, 800 μL culture medium with/without EBBE or 20 μM cyclopamine tartrate was loaded into each well of the 24-well plate. HSC-3-containing chamber inserts were then placed on the 24-well plate. After incubating the chamber for 24 h at 37°C, migrated cells in the lower surface of the chamber were fixed. The cells were counted and documented under an inverted light microscope.

Immunoblotting

Expression levels of Sonic hedgehog (SHH) protein were measured using immunoblotting method as previously described²⁴ with modification. HSC-3 cells were lysed and homogenized with Laemmli sample buffer (Bio-Rad). The proteins from each experimental group were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and transferred to a polyvinylidene difluoride (PVDF) sheet. After blocking with 5% skim milk solution, the sheet was incubated with rabbit monoclonal anti-SHH antibody as a primary antibody and horseradish peroxidase (HRP)-conjugated goat anti-rabbit antibody as a secondary antibody (Cell Signaling Technology, Danvers, MA, USA). Immunoblot results were analyzed using ImageJ software (National Institute of Health, Bethesda, MD, USA) to measure the density of each SHH band.

Statistical analysis

Statistical analyses were performed using IBM SPSS Statistics version 26.0 (SPSS IBM, Armonk, NY, USA). Shapiro-Wilk test was used as a normality test. Differences of MTT assay, sub-G1 assay, and scratch assay results between experimental groups were analyzed using Kruskal-Wallis H with Mann-Whitney U *post hoc* analysis. Meanwhile, differences of transwell assay results and SHH band density between experimental groups were analyzed using one-way analysis of variances (ANOVA) with Tukey's honestly significant difference (HSD) *post hoc* analysis. All results were expressed as mean \pm standard deviation (SD). $P < 0.05$ were considered as statistically significant.

Results

EBBE reduced the number of viable HSC-3 cells

The number of viable HSC-3 cells cultured in medium containing extract solvent (sham group) for

24 h was $9,607 \pm 13.81$. Treatment with EBBE showed that the viability of HSC-3 cells was significantly lower compared with that of the sham group ($P < 0.05$) in a concentration-dependent manner. The number of viable HSC-3 cells treated with 1, 10, and 100 $\mu\text{g/mL}$ EBBE were $9,055 \pm 268.69$, $6,777 \pm 278.04$, and $3,659 \pm 32.86$, respectively. The viability of EBBE-treated HSC-3 cells was significantly higher ($P < 0.05$) than that of Doxorubicin-treated HSC-3 cells (226 ± 32.34) (Fig. 1). These findings suggested that EBBE could reduce the viability of HSC-3 cells.

EBBE induced apoptosis of HSC-3 cells

Sub-G1 assay results showed that the apoptosis percentage of the sham group was $8.27 \pm 0.14\%$. Apoptotic HSC-3 cells were significantly higher compared with the sham group ($P < 0.05$) in concentration-dependent manner after EBBE treatment. Apoptosis percentages of HSC-3 cells treated with 1, 10, and 100 $\mu\text{g/mL}$ EBBE were $9.60 \pm 0.35\%$, $17.78 \pm 0.32\%$, and $38.64 \pm 0.70\%$, respectively. These percentages were significantly lower ($P < 0.05$) than that of Doxorubicin-treated HSC-3 cells ($95.95 \pm 2.60\%$) (Fig. 2). Therefore, these results suggested that EBBE treatment could promote apoptosis of HSC-3 cells.

EBBE inhibited HSC-3 cell migration

The gap closure percentage of HSC-3 cells cultured in medium containing extract solvent after 24 h was




Fig. 1 — EBBE decreased the number of viable HSC-3 cells in a concentration-dependent manner. HSC-3 cells were cultured and treated with/without 1 μM Doxorubicin or various concentrations of EBBE for 24 h as indicated in the panel. The number of viable cells were measured with MTT assay as described in Materials and Methods. The data are expressed as mean \pm SD ($n=6$). $*P < 0.05$ compared to the sham group




Fig. 2 — EBBE increased the percentage of apoptotic HSC-3 cells in a concentration-dependent manner. HSC-3 cells were cultured and treated with/without 1 μ M Doxorubicin or various concentrations of EBBE for 24 h as indicated. Treated cells were then subjected to sub-G1 assay as described in Materials and Methods. (A) Histogram of flow cytometric results. (B) Percentage of apoptotic HSC-3 cells. The data are expressed as mean \pm SD ($n=18$). * $P < 0.05$ compared to the sham group

96.11 \pm 4.17%. After EBBE treatment, the migratory activities of HSC-3 cells were significantly lower compared with that of the sham group ($P < 0.05$) in a concentration-dependent manner. Gap closure percentages of HSC-3 cells treated with 1, 10, and 100 μ g/mL EBBE were 89.56 \pm 5.77%, 76.78 \pm 4.87%, and 60.67 \pm 7.62%, respectively. These percentages were significantly higher ($P < 0.05$) than that of cyclopamine tartrate-treated HSC-3 cells (43.00 \pm 4.82%) (Fig. 3).

Transwell assay also showed similar results. The number of migrated HSC-3 cells in the sham group was 134.44 \pm 11.60. The number of migrated HSC-3 cells was significantly lower than that of the sham group ($P < 0.05$) in a concentration-dependent manner after EBBE treatment. The number of migrated HSC-3 cells treated with 1, 10, and 100 μ g/mL EBBE were 107.44 \pm 15.21, 95.11 \pm 6.01, and 39.22 \pm 3.56, respectively. These numbers were significantly higher ($P < 0.05$) than that of cyclopamine tartrate-treated HSC-3 cells (17.89 \pm 3.22) (Fig. 4). Overall, these

results indicated that EBBE exhibited an anti-migratory effect towards HSC-3 cells.

EBBE downregulated the expression of SHH

Upon the addition of EBBE, the SHH expression levels in HSC-3 cells were lower in a concentration-dependent manner. Expression level of SHH in HSC-3 cells treated with 1 μ g/mL EBBE was significantly higher ($P < 0.05$), while SHH expression levels in HSC-3 cells treated with 10 and 100 μ g/mL EBBE were significantly lower ($P < 0.05$) compared with that in the sham group. SHH expression levels in HSC-3 cells treated with 1 and 10 μ g/mL EBBE were significantly higher compared with that in cyclopamine tartrate-treated HSC-3 cells ($P < 0.05$). However, no significant difference was observed in the SHH expression levels between cyclopamine tartrate and 100 μ g/mL EBBE treatment ($P = 0.142$) (Fig. 5). Thus, these findings suggested that EBBE could decrease the expression of SHH.




Fig. 3 — EBBE inhibited migratory activities of HSC-3 cells. HSC-3 cell culture was scratched and treated with/without EBBE or 20 µM cyclopamine tartrate for 24 h as indicated. Gap closure was measured as described in Materials and Methods. (A) Scratched areas of HSC-3 cell culture. White bar: 100 µm. (B) Gap closure percentage of EBBE-treated HSC-3 cell. The data are expressed as mean ± SD (n=9). * $P < 0.05$ compared to the sham group




Fig. 4 — EBBE reduced the number of migrated HSC-3 cells. HSC-3 cell culture was treated with/without EBBE or 20 µM cyclopamine tartrate for 24 h as indicated. Migrated cells were measured as described in Materials and Methods. The data are expressed as mean ± SD (n=9). * $P < 0.05$ compared to the sham group




Fig. 5 — EBBE reduced the expression of SHH. HSC-3 cell culture was treated with/without EBBE or 20 µM cyclopamine tartrate for 24 h as indicated. HSC-3 cells were lysed, homogenized, and immunoblotted using anti-SHH antibody as described in Materials and Methods. The data are expressed as mean ± SD (n=3). * $P < 0.05$ compared to the sham group

Discussion

In the present study, we investigated the potency of EBBE as anticancer agent for OSCC treatment. EBBE showed a cytotoxic effect towards HSC-3 cells by reducing the viability of HSC-3 cells in a concentration-dependent manner. According to the sub-G1 assay results, the decrease in HSC-3 cell viability was caused by apoptosis. These results were in line with previous studies that EBBE was able to induce apoptosis of cancer cells^{18-20,25}. EBBE was also reported to inhibit cell cycle in several cell lines^{19,25}, indicating that EBBE may induce apoptosis of HSC-3 cells through cell cycle arrest since both processes are interconnected²⁶. Based on a molecular docking study, eleutherol, eleutherin, and isoeleutherin could induce apoptosis by activating caspase-8²⁷.

Cell migration is one of the key processes involved in cancer metastasis²⁸. Scratch and transwell assays demonstrated that EBBE restrained the migration of HSC-3 cells in a concentration-dependent manner. This showed the novel activity of EBBE in impairing HSC-3 cell migration. Therefore, EBBE may have anti-metastasis activity towards tongue cancer cells.

Metastasis of tongue cancer is regulated by several proteins, including SHH. One of the SHH signaling pathways that has been reported to regulate cell migration in tongue cancer is SHH/glioma-associated oncogene homologue (SHH/GLI)²⁹. SHH protein binds and inactivates its receptor, Patched 1 (PTCH1). This interaction causes the activation of transcription factors from the GLI family, which play a major role in regulating cancer migration, invasion, and proliferation³⁰.

The present study revealed that EBBE decreased SHH expression levels along with the increase in EBBE concentrations, suggesting that the inhibitory effect of EBBE towards the expression of SHH may be responsible for its anti-migratory property. To the best of our knowledge, this is the first study demonstrating SHH downregulation in EBBE-treated HSC-3 cells. EBBE may inhibit HSC-3 cell migration via inhibition of SHH/GLI signaling pathway. Possibly, all signaling pathways that are regulated by SHH can be inhibited by EBBE. EBBE may also restrain HSC-3 cell migration through another pathway. In SW480 colon cancer cell line, eleutherinoside C and isoeleutherin obtained from *E. bulbosa* bulbs was reported to inhibit Wnt/ β -catenin pathway¹⁶, which has been reported to play an important role in regulating cancer migration³¹.

Therefore, signaling pathways involved in the inhibition of HSC-3 cell migration by EBBE still need to be explored. In addition, signaling components downstream to SHH and target molecules that are affected by EBBE should be identified. Since EBBE also contains various bioactive compounds, further research is needed to identify which compounds are responsible for apoptosis and anti-migratory activity. Thus, more investigations are required to understand the molecular mechanisms underlying EBBE-induced HSC-3 cell apoptosis and confirm which SHH signaling pathway contributes to the inhibition of cell migration by EBBE.

Conclusion

EBBE reduces HSC-3 cell viability through apoptosis and inhibits its migratory activities by downregulating SHH expression. Taken together, EBBE could be a potential anti-tongue cancer agent, particularly as an antimetastatic agent.

Conflict of interest

All authors declare no conflicts of interest.

References

- Chen YK, Huang HC, Lin LM & Lin CC. Primary oral squamous cell carcinoma: An analysis of 703 cases in southern Taiwan. *Oral Oncol*, 35 (1999) 173.
- Neville BW & Day TA. Oral cancer and precancerous lesions. *CA Cancer J Clin*, 52 (2002) 195.
- Minhas S, Kashif M, Altaf W, Afezal N & Nagi AH. Concomitant-chemoradiotherapy-associated oral lesions in patients with oral squamous-cell carcinoma. *Cancer Biol Med*, 14 (2017) 176.
- Wicaksono BD, Tangkearung E & Sandra F. *Brucea javanica* leaf extract induced apoptosis in human oral squamous cell carcinoma (HSC2) cells by attenuation of mitochondrial membrane permeability. *Indones Biomed J*, 7 (2015) 107.
- Chang WL, Cheng FC, Wang SP, Chou ST & Shih Y. *Cinnamomum cassia* essential oil and its major constituent cinnamaldehyde induced cell cycle arrest and apoptosis in human oral squamous cell carcinoma HSC-3 cells. *Environ Toxicol*, 32 (2017) 456.
- Susianti S, Lesmana R, Salam S, Julaha E, Pratiwi YS, Sylviana N, Goenawan H, Kuniawan A & Supratman U. The effect of nutmeg seed (*M. fragrans*) extracts induces apoptosis in melanoma malignant cell's (B16-F10). *Indones Biomed J*, 13 (2021) 68.
- Poonacha SK, Harishkumar M, Radha M, Varadarajan R, Nalilu SK, Shetty SS, Shetty PK, Chandrashekhara RB, Sreenivas MG & Bavabeedu SKB. Insight into oroxylin A-7-O- β -d-glucuronide-enriched *Oroxylum indicum* bark extract in oral cancer HSC-3 cell apoptotic mechanism: Role of mitochondrial microenvironment. *Molecules*, 26 (2021) 7430.
- Ren W, Qiao Z, Wang H, Zhu L & Zhang L. Flavonoids: Promising anticancer agents. *Med Res Rev*, 23 (2003) 519.

- 9 Sandra F & Sidharta MA, Caffeic acid induced apoptosis in MG63 osteosarcoma cells through activation of caspases. *Mol Cell Biomed Sci*, 1 (2017) 28.
- 10 Ieyama T, Gunawan-Puteri MDPT & Kawabata J, α -Glucosidase inhibitors from the bulb of *Eleutherine americana*. *Food Chem*, 128 (2011) 308.
- 11 Subramaniam K, Suriyamoorthy S, Wahab F, Sharon FB & Rex GR, Antagonistic activity of *Eleutherine palmifolia* Linn. *Asian Pac J Trop Dis*, 2 (2012) S491.
- 12 Kuntorini EM, Dewi M & Misrina M, Anatomical structure and antioxidant activity of red bulb plant (*Eleutherine americana*) on different plant age. *Biodiversitas*, 17 (2016) 229.
- 13 Insanu M, Kusmardiyani S & Hartati R, Recent studies on phytochemicals and pharmacological effects of *Eleutherine americana* Merr. *Procedia Chem*, 13 (2014) 221.
- 14 Chen Z, Huang H, Wang C, Li Y, Ding J, Sankawa U, Noguchi H & Iitaka Y, Hongconin, a new naphthalene derivative from hong-cong, the rhizome of *Eleutherine Americana* Merr. et Heyne (Iridaceae). *Chem Pharm Bull*, 34 (1986) 2743.
- 15 Xu J, Qiu F, Duan W, Qu G, Wang N & Yao X, New bioactive constituents from *Eleutherine americana*. *Front Chem China*, 1 (2006) 320.
- 16 Li X, Ohtsuki T, Koyano T, Kowithayakorn T & Ishibashi M, New Wnt/ β -catenin signaling inhibitors isolated from *Eleutherine palmifolia*. *Chem Asian J*, 4 (2009) 540.
- 17 Mutiah R, Minggarwati TS, Kristanti RA & Susanti E, Compound identification and anticancer activity of ethyl acetate fraction from bawang sabrang (*Eleutherine palmifolia* (L.) Merr.) on HeLa cervical cancer cell line. *Indones J Cancer Chemoprevent*, 10 (2019) 131.
- 18 Lubis IA, Ichwan M, Mustofa M & Satria D, Anticancer activity of *Eleutherine bulbosa* (Mill.) Urb. extract on WiDr cell line *in vitro*, paper presented to 2nd Public Health International Conference (PHICO 2017), Faculty of Public Health - Universitas Sumatera Utara, Medan, Indonesia, 18-19 December 2017.
- 19 Yuniarti A, Sundhani E & Nurulita NA, The potentiation effect of bawangdayak (*Sisyrinchium palmifolium* L.) extract on T47D cell growth inhibition after 5-fluorouracil treatment. *Pharmaciana*, 8 (2018) 195.
- 20 Kamarudin AA, Sayuti NH, Saad N, Razak NAA & Esa NM, Induction of apoptosis by *Eleutherine bulbosa* (Mill.) Urb. bulb extracted under optimised extraction condition on human retinoblastoma cancer cells (WERI-Rb-1). *J Ethnopharmacol*, 284 (2022) 114770.
- 21 Sandra F, Sudiono J, Trisfilha P & Pratiwi D, Cytotoxicity of *Alpinia galanga* rhizome crude extract on NIH-3T3 cells. *Indones Biomed J*, 9 (2017) 23.
- 22 Ariesanti Y, Sandra F, Claresta B & Alvita L, *Coffea canephora* bean extract induces NIH3T3 cell migration. *Indones Biomed J*, 13 (2021) 216.
- 23 Oktaviono YH, Al-Farabi MJ, Suastika LOS, Hartono F, Dirgantara Y & Sandra F, Preliminary study: Purple sweet potato extract seems to be superior to increase the migration of impaired endothelial progenitor cells compared to L-Ascorbic acid. *Sci Pharm*, 87 (2019) 16.
- 24 Rizal MI & Sandra F, *Brucea javanica* leaf extract activates caspase-9 and caspase-3 of mitochondrial apoptotic pathway in human oral squamous cell carcinoma. *Indones Biomed J*, 8 (2016) 43.
- 25 Mutiah R, Listiyana A, Suryadinata A, Annisa R, Hakim A, Anggraini W & Susilowati R, Activity of inhibit the cell cycle and induct apoptosis in HeLa cancer cell with combination of sabrang onion (*Eleutherine palmifolia* (L.) Merr) and starfruit mistletoe (*Macrosolen cochinchinensis* (Lour.) Tiegh). *J App Pharm Sci*, 8 (2018) 122.
- 26 Pucci B, Kasten M & Giordano A, Cell cycle and apoptosis. *Neoplasia*, 2 (2000) 291.
- 27 Quadros Gomes AR, da Rocha Galucio NC, de Albuquerque KCO, Brigido HPC, Varela ELP, Castro ALG, Vale VV, Bahia MO, Rodriguez Burbano RM, de Molfeta FA, Carneiro LA, Percario S & Dolabela MF, Toxicity evaluation of *Eleutherine plicata* Herb. extracts and possible cell death mechanism. *Toxicol Rep*, 8 (2021) 1480.
- 28 Ellis IR, The migration and invasion of oral squamous carcinoma cells: Matrix, growth factor and signalling involvement. *Cancers*, 13 (2021) 2633.
- 29 Fan HX, Wang S, Zhao H, Liu N, Chen D, Sun M & Zheng JH, Sonic hedgehog signaling may promote invasion and metastasis of oral squamous cell carcinoma by activating MMP-9 and E-cadherin expression. *Med Oncol*, 31 (2014) 41.
- 30 Carballo GB, Honorato JR, de Lopes GPF & Spohr TCLSE, A highlight on Sonic hedgehog pathway. *Cell Commun Signal*, 16 (2018) 11.
- 31 Liang S, Zhang S, Wang P, Yang C, Shang C, Yang J & Wang J, LncRNA, TUG1 regulates the oral squamous cell carcinoma progression possibly *via* interacting with Wnt/ β -catenin signaling. *Gene*, 608 (2017) 49.

Eleutherine bulbosa bulb extract induces apoptosis and inhibits cell migration by downregulating Sonic hedgehog in human tongue cancer cells: An in vitro study

ORIGINALITY REPORT

12%

SIMILARITY INDEX

10%

INTERNET SOURCES

15%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

- 1** www.scielo.br Internet Source 2%
- 2** Kon, Takao(Department of Dentistry and Oral Surgery, Hirosaki University Graduate School of Medicine, Hirosaki, Japan, 今, 敬生 and 弘前大学大学院医学研究科). "Role of type I- and type II-interferon in expression of melanoma differentiation-associated gene-5 in HSC-3 oral squamous carcinoma cells", Biomedical Research Foundation, 2014. Publication 1%
- 3** molecular-cancer.biomedcentral.com Internet Source 1%
- 4** Po-Yuan Chen, Kai-Chun Lin, Jing-Pin Lin, Nou-Ying Tang, Jai-Sing Yang, Kung-Wen Lu, Jing-Gung Chung. " Phenethyl Isothiocyanate (PEITC) Inhibits the Growth of Human Oral Squamous Carcinoma HSC-3 Cells through Phase Arrest and Mitochondria-Mediated 1%

Apoptotic Cell Death ", Evidence-Based Complementary and Alternative Medicine, 2012

Publication

5	integrativeonc.org Internet Source	1 %
6	Fukuda, . "IL-23 promotes growth and proliferation in human squamous cell carcinoma of the oral cavity", International Journal of Oncology, 2010. Publication	1 %
7	Afifi, Marwa M., Lauren A. Austin, Megan A. Mackey, and Mostafa A. El-Sayed. "XAV939: From a Small Inhibitor to a Potent Drug Bioconjugate When Delivered by Gold Nanoparticles", Bioconjugate Chemistry, 2014. Publication	1 %
8	advances.umw.edu.pl Internet Source	1 %
9	aacr.silverchair-cdn.com Internet Source	1 %
10	repository.uhamka.ac.id Internet Source	1 %
11	www.koreascience.or.kr Internet Source	1 %

12 Xia Xiao, Huaqin Liu, Xiaodong Li. "Orlistat treatment induces apoptosis and arrests cell cycle in HSC-3 oral cancer cells", Microbial Pathogenesis, 2017
Publication 1 %

13 perpustakaan.poltekkes-malang.ac.id
Internet Source 1 %

14 Zhigang Zeng, Kaiyan Xiao. " Polysaccharide (GLP) Inhibited the Progression of Oral Squamous Cell Carcinoma via the miR-188/BCL9/ -Catenin Pathway ", Advances in Polymer Technology, 2020
Publication 1 %

15 rupress.org
Internet Source 1 %

16 www.pubmedcentral.nih.gov
Internet Source <1 %

Exclude quotes On

Exclude matches < 15 words

Exclude bibliography On

Eleutherine bulbosa bulb extract induces apoptosis and inhibits cell migration by downregulating Sonic hedgehog in human tongue cancer cells: An in vitro study

GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/15

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7



Ferry Sandra <ferry@trisakti.ac.id>

[IJBB] Editor Decision (IJBB#83)

Team IJBB <ijbb@niscpr.res.in>

Sat, Sep 16, 2023 at 3:36 PM

To: Ferry Sandra <ferry@trisakti.ac.id>, Kelvin Suryajaya <kelvin.thong222@gmail.com>, Angliana Chow <angliana@prostem.co.id>, Maria Celinna <maria.celinna@prodia.institute>, Dewi Ranggaini <monica.dewi.r@trisakti.ac.id>, Cynthia Retna Sartika <cynthia@prostem.co.id>

Ferry Sandra, Kelvin Suryajaya, Angliana Chow, Maria Celinna, Dewi Ranggaini, Cynthia Retna Sartika:

We have reached a decision regarding your submission to Indian Journal of Biochemistry and Biophysics (IJBB), "Eleutherine bulbosa Bulb Extract Induces Apoptosis and Inhibits Cell Migration by Downregulating Sonic Hedgehog Gene in Human Tongue Cancer Cells: An in vitro Study(Article Id: IJBB#83)".

Our decision is: Revisions Required

[Indian Journal of Biochemistry and Biophysics \(IJBB\)](#)

**B-IJBB#83 report.docx**

18K



Ferry Sandra <ferry@trisakti.ac.id>

Editor Decision: Revision Required (IJBB#83)

IJBB <ijbb@niscpr.res.in>
To: ferry@trisakti.ac.id

Sat, Sep 16, 2023 at 3:38 PM

Dear Author,

We have reached a decision regarding your submission to Indian Journal of Biochemistry and Biophysics (IJBB), "Eleutherine bulbosa Bulb Extract Induces Apoptosis and Inhibits Cell Migration by Downregulating Sonic Hedgehog Gene in Human Tongue Cancer Cells: An in vitro Study(Article Id: IJBB#83)".

Kindly submit your revised by [tomorrow](#) evening. We have commitment for other papers as well hence delay from your side will lead us to shift your MS to later issues according to the date of revision.

Our decision is: Revisions Required

Thanks & Regards
Ream IJBB**IJBB#83 report.docx**

18K

IJBB#83

Eleutherine bulbosa Bulb Extract Induces Apoptosis and Inhibits Cell Migration by Downregulating Sonic Hedgehog Gene in Human Tongue Cancer Cells

Reviewer Comments

After careful review of the manuscript, it has been found that the article is suitable for publication in IJBB after minor revision. The following comments should be included in the revised manuscript

Comments:

1. Lots of words (especially in References) are merged - may be computer-to-computer use glitch - please make sure the words are appropriately spaced.
2. Throughout the manuscript (including in Figure Legends), please use "h" instead of "hours".
3. Each Figure should list the Abbreviations used in a particular figure (it should stand alone) despite already abbreviated in the text elsewhere.
4. Figures 2 and 3 have two panels - the legend says "A" and "B" whereas the figures denote as "a" and "b" - please be consistent, eitherway is fine.
5. **PI highlights the corrections and resubmit the MS ASAP.**

Editor's note:

- Ensure that it has not been published elsewhere meanwhile, either in part or in full, including the illustrations.
- Read our "Instructions to authors" before preparing your Manuscript and adhere to those instructions strictly. Read some recent papers of IJBB in our repository NOPR (<http://nopr.niscpr.res.in>) which has free access.
- Keywords need not repeat terms that already appear in the title. Avoid generic terms as well.
- Provide a short/running title.
- The abstract should begin with the justification of the present study and its requirement. Why at all this study? What is the demand/necessity?
- Introduction and Methodology sections need to have adequate references.
- Whole text should be in TNR, font 11 size; and Subheadings, font 9. Text should follow in the next line. Refer any latest IJBB paper for reference.
- Ethical statements like Conflict-of-interest and Informed consent (if involved) need to be specified as per Committee on Publication Ethics guidelines, separately prior to acknowledgement.

- **You may drop the surplus ones. Limit the total references to <45 after updation.**
- Avoid citing references from fake journals (Please check if your references are from journals indexed by standard agencies, such as SCI indexed (including SCI-expanded list), Scopus, PubMed, NAAS, etc. You may also visit University Grants Commission (UGC), Govt. of India web page and use the list of approved journals to verify. The following link <https://predatoryjournals.com/publishers/> gives you list of predatory publishers, Journals, etc.
- Format and style of references in the text as well as in the respective reference section is not as per IJBB pattern. While citing refs in the text as superscript there should not be any space between (neither between the sentence and reference number in superscript nor between the ref nos.). For e.g. it should not be like “same species [14, 15].” but like this “same species^{14,15}.” If it is more than two refs, then it needs to be hyphenated. For e.g. “... of all drugs and foreign substances^{1,2,3}.” “... of all drugs and foreign substances¹⁻³.”
- For ‘references’ section, strictly follow the pattern of IJBB [author names (initials without space), ‘&’ before the last author and comma after the end of the author names; title of the paper followed by full stop, source (**abbreviated form of the journal following international standards and in italics without any full stop**) followed by non-italic comma, vol no., year within brackets followed by initial page no. followed by a full stop].
- In references, we give all author names. Et al. will not work in IJBB.
- Also, follow proper style for units: “ml” should be “mL”. Similarly, mL also. Moreover, use proper symbols for temperature degree, minus sign, etc. You may refer any latest paper from IJBB through our open repository NOPR at <http://nopr.niscpr.res.in>.
- When you mention room temperature in your study, you need to be specific at its first occurrence in the text.
- All tables should be in TNR font 9, single line space, no bold letters with proper rows and columns.

You may address all these quickly and send us an error-free final version quickly so that it can be taken up for one of the forthcoming issue.

Sincerely yours,

Team IJBB



Ferry Sandra <ferry@trisakti.ac.id>

[IJBB] Editor Decision (IJBB#83)

Ferry Sandra <ferry@trisakti.ac.id>
To: ijbb@niscpr.res.in

Mon, Sep 18, 2023 at 1:40 PM

Dear Team IJBB,

Thank you for your email. We have uploaded a revised version of our manuscript in your submission system. Hopefully, our manuscript is acceptable for publication in your reputable journal.

Regards,
Ferry Sandra
[Quoted text hidden]

—
Ferry Sandra, D.D.S., Ph.D.
Head of Medical Research Center
Universitas Trisakti

***Eleutherine bulbosa* Bulb Extract Induces Apoptosis and Inhibits Cell Migration by
Downregulating Sonic Hedgehog Gene in Human Tongue Cancer Cells: An *in vitro* Study**

Running title: *Eleutherine bulbosa* Effect on Tongue Cancer Cells

Ferry Sandra^{1*}, Kelvin Suryajaya², Angliana Chow³, Maria Celinna⁴, Dewi Ranggaini⁵, Cynthia
Retna Sartika³

¹Department of Biochemistry and Molecular Biology, Division of Oral Biology, Faculty of Dentistry,
Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia

²Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia

³PT Prodia StemCell Indonesia, Jakarta Pusat-10430, Jakarta, Indonesia

⁴The Prodia Education and Research Institute, Jakarta Pusat-10430, Jakarta, Indonesia

⁵Department of Physiology, Division of Oral Biology, Faculty of Dentistry, Universitas Trisakti,
Jakarta Barat-11440, Jakarta, Indonesia

***Corresponding Author:**

Ferry Sandra

Department of Biochemistry and Molecular Biology, Division of Oral Biology, Faculty of Dentistry,

Universitas Trisakti, Jl. Kyai Tapa No. 260, Jakarta Barat-11440, Jakarta, Indonesia

E-mail address: ferry@trisakti.ac.id

Phone: +628128077780

Abstract

Eleutherine bulbosa (Mill.) Urb. is a medicinal plant which has long been used to treat cancer. *E. bulbosa* bulb extract (EBBE) has been reported to show cytotoxicity towards several types of human cancer. However, the cytotoxic effect of EBBE towards oral cancer cells has not been investigated. The present study aimed to evaluate the effects of EBBE towards the viability, apoptosis, and migratory activities of oral cancer cells. Human oral squamous cell carcinoma (HSC)-3 cells were treated with various concentrations of EBBE for 24 h. The number of viable and apoptotic HSC-3 cells were measured using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) and sub-G1 assay, respectively. The migratory activities of HSC-3 cells were assessed using scratch and transwell assay. Sonic hedgehog (SHH) expression was measured using immunoblotting. Upon EBBE treatment, apoptotic HSC-3 cells were significantly higher in a concentration-dependent manner ($P < 0.05$). Meanwhile, viability, migratory activities, and migrated HSC-3 cell number were significantly lower in a concentration-dependent manner ($P < 0.05$). The SHH expression levels in EBBE-treated HSC-3 cells were also lower in a concentration-dependent manner. EBBE reduces HSC-3 cell viability through apoptosis and inhibits its migratory activities by downregulating SHH expression.

Keywords: HSC-3, MTT, sub-G1, transwell, immunoblotting

Oral squamous cell carcinoma (OSCC) is the most common histological type of oral cancer¹ which arises from squamous epithelial cells in the oral cavity. This cancer generally affects the floor of the mouth and the tongue². Various modes of treatment are developed to manage OSCC, including chemotherapy, radiotherapy, surgery, or its combination³. However, these treatments cause numerous side effects.

Discovery and development of new therapeutic agents derived from natural sources, particularly plants with less side effects are recently one of the main focuses on oncology research. Several plants have been reported to have anticancer effects, including *Brucea javanica*⁴, *Cinnamomum cassia*⁵, *Myristica fragrans*⁶, and *Oroxylum indicum*⁷. Plants have been known as rich sources of various active compounds, including flavonoids, which was reported to inhibit cancer proliferation via cell cycle arrest, antiproliferative and antioxidative activity, apoptosis induction, promotion of differentiation, inhibition of angiogenesis and carcinogen metabolic activation, and multidrug resistance modulation⁸. In addition, several non-flavonoid phenolic compounds have also been reported to show anticancer properties, such as caffeic acid⁹.

Eleutherine bulbosa (Mill.) Urb. is a traditional medicinal plant belonging to the Iridaceae family. Its bulbs are commonly used for the treatment of breast cancer, hypertension, stroke, and diabetes mellitus. It has also been used to increase the production of breast milk and treat sexual disorder¹⁰. *E. bulbosa* bulbs have also been reported to have antimicrobial¹¹ and antioxidant¹² activity.

E. bulbosa contains a wide variety of compounds, primarily naphthoquinones, anthraquinones, and naphthalene¹³. The main compounds found in this plant are eleutherin, isoeleutherin, and eleutherol¹⁴. Several compounds obtained from *E. bulbosa* bulbs have been reported to show anticancer activities, such as 6,8-hydroxy-3,4- di-methoxy-1-methyl-anthraquinone-2-carboxylic acid methyl ester¹⁵, eleutherinoside C, and isoeleutherin¹⁶.

E. bulbosa bulb extract (EBBE) has been reported to show cytotoxicity towards several types of human cancer, such as cervical cancer (line HeLa)¹⁷, colon cancer (line SW480, HCT116, DLD1, WiDr)^{16,18}, breast cancer (line T47D)¹⁹, leukemia (line K562)¹⁵, and retinoblastoma (WERI-Rb-1)²⁰. However, the cytotoxic effect of EBBE towards human oral cancer cells has not been investigated.

Therefore, the present study was conducted to evaluate the effects of EBBE towards the viability, apoptosis, and migratory activities of oral cancer cells.

Materials and Methods

E. bulbosa bulb extract (EBBE) preparation

E. bulbosa bulbs were obtained from Indonesian Medicinal and Aromatic Crops Research Institute (IMACRI), Indonesia. Plant identification was carried out by expert botanist at Research Center for Plant Conservation and Botanic Gardens, Indonesian Institute of Sciences, Indonesia (No. B-1269/IPH.3/KS/X/2020). EBBE was extracted by maceration method⁴. Briefly, bulbs of *E. bulbosa* were minced and dried. The dried material was extracted with ethanol 70%, filtered and evaporated using a rotary evaporator. The resulting crude EBBE was then stored at 4°C.

HSC-3 cell culture

HSC-3 cell culture was performed as previously described⁴ with modification. The HSC-3 cell line was purchased from Sigma-Aldrich Pte. Ltd. (St. Louis, MO, USA). HSC-3 cells were cultured in a complete medium containing Dulbecco's modified Eagle's medium (DMEM) (Sigma-Aldrich), 10% fetal bovine serum (FBS) (PAN-Biotech, Aidenbach, Germany), 50 U/mL penicillin and 50 µg/mL streptomycin (Sigma-Aldrich) in a humidified 5% CO₂, 37°C incubator. HSC-3 cells were then dissociated with trypsin-ethylenediamine tetraacetic acid (EDTA) solution (Sigma-Aldrich) after reaching 80% confluency. Upon reaching the desired cell number, HSC-3 cells were seeded into 24-well plate and 96-well plate for further assays.

Cell viability assay

The number of viable cells were measured quantitatively using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay as previously described²¹ with modification. Briefly, 5×10^3 HSC-3 cells were seeded into 96-well plates and then treated with EBBE (1, 10, or 100 µg/mL), 1 µM Doxorubicin (Dankos Farma, Jakarta, Indonesia) or medium merely for 24 h. After that 100 µL of MTT (Sigma-Aldrich) in culture medium was loaded into each well. After incubating the plates for 4 h, culture medium was discarded, and the formed formazan crystals were dissolved in 100 µL dimethylsulfoxide (DMSO). Results were measured at OD₅₇₀ using a microplate reader (Bio-Rad,

Hercules, CA, USA). Each experimental group was measured in six replicates. In addition, the number of untreated cells were counted with hemacytometer and used for interpolating OD₅₇₀ values of HSC-3 cells.

Sub-G1 assay

To investigate the cytotoxic effect of EBBE on HSC-3 cells, the number of apoptotic cells were determined using sub-G1 assay as previously described⁹ with modification. Treated and untreated HSC-3 cells were harvested and suspended in 450 µL hypotonic fluorochrome solution containing 50 µg/mL propidium iodide (Sigma-Aldrich), 0.1% Triton X-100 (Sigma-Aldrich), and 0.1% sodium citrate (Wako, Osaka, Japan). After that, cell suspensions were incubated in the dark for 2 h at room temperature. Fluorescence of individual nuclei was measured using FACSCanto II flow cytometer (Becton Dickinson, Franklin Lakes, NJ, USA) at 100,000 events.

Scratch assay

Scratch assay was performed as previously described²². HSC-3 cells (5×10^4) were seeded into 24-well plates and incubated overnight. Scratch was made on each well with micropipette yellow tips. Then, the wells were washed with phosphate-buffered saline (PBS) (Wako) and treated with/without EBBE or 20 μ M cyclopamine tartrate (BioVision, Milpitas, CA, USA) (as positive control) for 24 h. The gap closures were observed and documented under an inverted light microscope (Carl Zeiss, Jena, Germany).

Transwell assay

Transwell assay was performed as previously described²³ with modification. HSC-3 cells (5×10^4) were seeded in the 8- μ M pore size chamber insert (Merck, Darmstadt, Germany). After that, 800 μ L culture medium with/without EBBE or 20 μ M cyclopamine tartrate was loaded into each well of the 24-well plate. HSC-3-containing chamber inserts were then placed on the 24-well plate. After incubating the chamber for 24 h at 37°C, migrated cells in the lower surface of the chamber were fixed. The cells were counted and documented under an inverted light microscope.

Immunoblotting

Expression levels of Sonic hedgehog (SHH) protein were measured using immunoblotting method as previously described²⁴ with modification. HSC-3 cells were lysed and homogenized with Laemmli sample buffer (Bio-Rad). The proteins from each experimental group were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and transferred to a polyvinylidene difluoride (PVDF) sheet. After blocking with 5% skim milk solution, the sheet was incubated with rabbit monoclonal anti-SHH antibody as a primary antibody and horseradish peroxidase (HRP)-conjugated goat anti-rabbit antibody as a secondary antibody (Cell Signaling Technology, Danvers, MA, USA). Immunoblot results were analyzed using ImageJ software (National Institute of Health, Bethesda, MD, USA) to measure the density of each SHH band.

Statistical analysis

Statistical analyses were performed using IBM SPSS Statistics version 26.0 (SPSS IBM, Armonk, NY, USA). Shapiro-Wilk test was used as a normality test. Differences of MTT assay, sub-

G1 assay, and scratch assay results between experimental groups were analyzed using Kruskal-Wallis H with Mann-Whitney U *post hoc* analysis. Meanwhile, differences of transwell assay results and SHH band density between experimental groups were analyzed using one-way analysis of variances (ANOVA) with Tukey's honestly significant difference (HSD) *post hoc* analysis. All results were expressed as mean \pm standard deviation (SD). $P < 0.05$ were considered as statistically significant.

Results

EBBE reduced the number of viable HSC-3 cells

The number of viable HSC-3 cells cultured in medium containing extract solvent (sham group) for 24 h was $9,607 \pm 13.81$. Treatment with EBBE showed that the viability of HSC-3 cells was significantly lower compared to the sham group ($P < 0.05$) in a concentration-dependent manner. The number of viable HSC-3 cells treated with 1, 10, and 100 $\mu\text{g/mL}$ EBBE were $9,055 \pm 268.69$, $6,777 \pm 278.04$, and $3,659 \pm 32.86$, respectively. The viability of EBBE-treated HSC-3 cells was significantly higher ($P < 0.05$) than Doxorubicin-treated HSC-3 cells (226 ± 32.34) (Fig. 1). These findings suggested that EBBE could reduce the viability of HSC-3 cells.

EBBE induced apoptosis of HSC-3 cells

Sub-G1 assay results showed that the apoptosis percentage of the sham group was $8.27 \pm 0.14\%$. Apoptotic HSC-3 cells were significantly higher compared to the sham group ($P < 0.05$) in concentration-dependent manner after EBBE treatment. Apoptosis percentages of HSC-3 cells treated with 1, 10, and 100 $\mu\text{g/mL}$ EBBE were $9.60 \pm 0.35\%$, $17.78 \pm 0.32\%$, and $38.64 \pm 0.70\%$, respectively. These percentages were significantly lower ($P < 0.05$) than Doxorubicin-treated HSC-3 cells ($95.95 \pm 2.60\%$) (Fig. 2). Therefore, these results suggested that EBBE treatment could promote apoptosis of HSC-3 cells.

EBBE inhibited HSC-3 cell migration

The gap closure percentage of HSC-3 cells cultured in medium containing extract solvent after 24 h was $96.11 \pm 4.17\%$. After EBBE treatment, the migratory activities of HSC-3 cells were significantly lower compared to the sham group ($P < 0.05$) in a concentration-dependent manner. Gap closure percentage of HSC-3 cells treated with 1, 10, and 100 $\mu\text{g/mL}$ EBBE were $89.56 \pm 5.77\%$,

76.78±4.87%, and 60.67±7.62%, respectively. These percentages were significantly higher ($P<0.05$) than cyclopamine tartrate-treated HSC-3 cells (43.00±4.82%) (Fig. 3).

Transwell assay also showed similar results. The number of migrated HSC-3 cells in the sham group was 134.44±11.60. The number of migrated HSC-3 cells was significantly lower than the sham group ($P<0.05$) in a concentration-dependent manner after EBBE treatment. The number of migrated HSC-3 cells treated with 1, 10, and 100 µg/mL EBBE were 107.44±15.21, 95.11±6.01, and 39.22±3.56, respectively. These numbers were significantly higher ($P<0.05$) than cyclopamine tartrate-treated HSC-3 cells (17.89±3.22) (Fig. 4). Overall, these results indicated that EBBE exhibited an anti-migratory effect towards HSC-3 cells.

EBBE downregulated the expression of SHH

Upon the addition of EBBE, the SHH expression levels in HSC-3 cells were lower in a concentration-dependent manner. Expression level of SHH in HSC-3 cells treated with 1 µg/mL EBBE was significantly higher ($P<0.05$), while SHH expression levels in HSC-3 cells treated with 10 and 100 µg/mL EBBE were significantly lower ($P<0.05$) compared to the sham group. SHH expression levels in HSC-3 cells treated with 1 and 10 µg/mL EBBE were significantly higher compared to cyclopamine tartrate-treated HSC-3 cells ($P<0.05$). However, no significant difference was observed in the SHH expression levels between cyclopamine tartrate and 100 µg/mL EBBE treatment ($P=0.142$) (Fig. 5). Thus, these findings suggested that EBBE could decrease the expression of SHH.

Discussion

In the present study, we investigated the potency of EBBE as anticancer agent for OSCC treatment. EBBE showed a cytotoxic effect towards HSC-3 cells by reducing the viability of HSC-3 cells in a concentration-dependent manner. According to the sub-G1 assay results, the decrease in HSC-3 cell viability was caused by apoptosis. These results were in line with previous studies that EBBE was able to induce apoptosis of cancer cells^{18–20,25}. EBBE was also reported to inhibit cell cycle in several cell lines^{19,25}, indicating that EBBE may induce apoptosis of HSC-3 cells through cell cycle

arrest since both processes are interconnected²⁶. Based on a molecular docking study, eleutherol, eleutherin, and isoeleutherin could induce apoptosis by activating caspase-8²⁷.

Cell migration is one of the key processes involved in cancer metastasis²⁸. Scratch and transwell assays demonstrated that EBBE restrained the migration of HSC-3 cells in a concentration-dependent manner. This showed the novel activity of EBBE in impairing HSC-3 cell migration. Therefore, EBBE may have anti-metastasis activity towards tongue cancer cells.

Metastasis of tongue cancer is regulated by several proteins, including SHH. One of the SHH signaling pathways that has been reported to regulate cell migration in tongue cancer is SHH/glioma-associated oncogene homologue (SHH/GLI)²⁹. SHH protein binds and inactivates its receptor, Patched 1 (PTCH1). This interaction causes the activation of transcription factors from the GLI family, which play a major role in regulating cancer migration, invasion, and proliferation³⁰.

The present study revealed that EBBE decreased SHH expression levels along with the increase in EBBE concentrations, suggesting that the inhibitory effect of EBBE towards the expression of SHH may be responsible for its anti-migratory property. To the best of our knowledge, this is the first study demonstrating SHH downregulation in EBBE-treated HSC-3 cells. EBBE may inhibit HSC-3 cell migration via inhibition of SHH/GLI signaling pathway. Possibly, all signaling pathways that are regulated by SHH can be inhibited by EBBE. EBBE may also restrain HSC-3 cell migration through another pathway. In SW480 colon cancer cell line, eleutherinoside C and isoeleutherin obtained from *E. bulbosa* bulbs was reported to inhibit Wnt/ β -catenin pathway¹⁶, which has been reported to play an important role in regulating cancer migration³¹. Therefore, signaling pathways involved in the inhibition of HSC-3 cell migration by EBBE still need to be explored. In addition, signaling components downstream to SHH and target genes that are affected by EBBE should be identified. Since EBBE also contains various bioactive compounds, further research is needed to identify which compounds are responsible for apoptosis and anti-migratory activity. Thus, more investigations are required to understand the molecular mechanisms underlying EBBE-induced HSC-3 cell apoptosis and confirm which SHH signaling pathway contributes to the inhibition of cell migration by EBBE.

In conclusion, EBBE reduces HSC-3 cell viability through apoptosis and inhibits its migratory activities by downregulating SHH expression. Taken together, EBBE could be a potential anti-tongue cancer agent, particularly as an antimetastatic agent.

Conflict of interest

All authors declare no conflicts of interest.

References

1. Chen YK, Huang HC, Lin LM & Lin CC, Primary oral squamous cell carcinoma: An analysis of 703 cases in southern Taiwan. *Oral Oncol*, 35 (1999) 173.
2. Neville BW & Day TA, Oral cancer and precancerous lesions. *CA Cancer J Clin*, 52 (2002) 195.
3. Minhas S, Kashif M, Altaf W, Afzal N & Nagi AH, Concomitant-chemoradiotherapy-associated oral lesions in patients with oral squamous-cell carcinoma. *Cancer Biol Med*, 14 (2017) 176.
4. Wicaksono BD, Tangkearung E & Sandra F, Brucea javanica leaf extract induced apoptosis in human oral squamous cell carcinoma (HSC2) cells by attenuation of mitochondrial membrane permeability. *Indones Biomed J*, 7 (2015) 107.
5. Chang WL, Cheng FC, Wang SP, Chou ST & Shih Y, Cinnamomum cassia essential oil and its major constituent cinnamaldehyde induced cell cycle arrest and apoptosis in human oral squamous cell carcinoma HSC-3 cells. *Environ Toxicol*, 32 (2017) 456.
6. Susianti S, Lesmana R, Salam S, Julaeha E, Pratiwi YS, Sylviana N, Goenawan H, Kurniawan A & Supratman U, The effect of nutmeg seed (*M. fragrans*) extracts induces apoptosis in melanoma malignant cell's (B16-F10). *Indones Biomed J*, 13 (2021) 68.
7. Kameyanda Poonacha S, Harishkumar M, Radha M, Varadarajan R, Nalilu SK, Shetty SS, Shetty PK, Chandrashekhara RB, Sreenivas MG & Bhandary Bavabeedu SK, Insight into oroxylinA-7-O- β -d-glucuronide-enriched Oroxyllum indicum bark extract in oral cancer HSC-3 cell apoptotic mechanism: Role of mitochondrial microenvironment. *Molecules*, 26 (2021) 7430.
8. Ren W, Qiao Z, Wang H, Zhu L & Zhang L, Flavonoids: Promising anticancer agents. *Med Res Rev*, 23 (2003) 519.

9. Sandra F & Sidharta MA, Caffeic acid induced apoptosis in MG63 osteosarcoma cells through activation of caspases. *Mol Cell Biomed Sci*, 1 (2017) 28.
10. Ieyama T, Gunawan-Puteri MDPT & Kawabata J, α -Glucosidase inhibitors from the bulb of *Eleutherine americana*. *Food Chem*, 128 (2011) 308.
11. Subramaniam K, Suriyamoorthy S, Wahab F, Sharon FB & Rex GR, Antagonistic activity of *Eleutherine palmifolia* Linn. *Asian Pac J Trop Dis*, 2 (2012) S491.
12. Kuntorini EM, Dewi M & Misrina M, Anatomical structure and antioxidant activity of red bulb plant (*Eleutherine americana*) on different plant age. *Biodiversitas*, 17 (2016) 229.
13. Insanu M, Kusmardiyani S & Hartati R, Recent studies on phytochemicals and pharmacological effects of *Eleutherine americana* Merr. *Procedia Chem*, 13 (2014) 221.
14. Chen Z, Huang H, Wang C, Li Y, Ding J, Sankawa U, Noguchi H & Iitaka Y, Hongconin, a new naphthalene derivative from hong-cong, the rhizome of *Eleutherine americana* Merr. et Heyne (Iridaceae). *Chem Pharm Bull*, 34 (1986) 2743.
15. Xu J, Qiu F, Duan W, Qu G, Wang N & Yao X, New bioactive constituents from *Eleutherine americana*. *Front Chem China*, 1 (2006) 320.
16. Li X, Ohtsuki T, Koyano T, Kowithayakorn T & Ishibashi M, New Wnt/ β -catenin signaling inhibitors isolated from *Eleutherine palmifolia*. *Chem Asian J*, 4 (2009) 540.
17. Mutiah R, Minggarwati TS, Kristanti RA & Susanti E, Compound identification and anticancer activity of ethyl acetate fraction from bawang sabrang (*Eleutherine palmifolia* (L.) Merr.) on HeLa cervical cancer cell line. *Indones J Cancer Chemoprevent*, 10 (2019) 131.
18. Lubis IA, Ichwan M, Mustofa M & Satria D, Anticancer activity of *Eleutherine bulbosa* (Mill.) Urb. extract on WiDr cell line in vitro, paper presented to 2nd Public Health International Conference (PHICo 2017), Faculty of Public Health – Universitas Sumatera Utara, Medan, Indonesia, 18-19 December 2017.
19. Yuniarti A, Sundhani E & Nurulita NA, The potentiation effect of bawang dayak (*Sisyrinchium palmifolium* L.) extract on T47D cell growth inhibition after 5-fluorouracil treatment. *Pharmaciana*, 8 (2018) 195.

20. Kamarudin AA, Sayuti NH, Saad N, Razak NAA & Esa NM, Induction of apoptosis by *Eleutherine bulbosa* (Mill.) Urb. bulb extracted under optimised extraction condition on human retinoblastoma cancer cells (WERI-Rb-1). *J Ethnopharmacol*, 284 (2022) 114770.
21. Sandra F, Sudiono J, Trisfilha P & Pratiwi D, Cytotoxicity of *Alpinia galanga* rhizome crude extract on NIH-3T3 cells. *Indones Biomed J*, 9 (2017) 23.
22. Ariesanti Y, Sandra F, Claresta B & Alvita L, *Coffea canephora* bean extract induces NIH3T3 cell migration. *Indones Biomed J*, 13 (2021) 216.
23. Oktaviono YH, Al-Farabi MJ, Suastika LOS, Hartono F, Dirgantara Y & Sandra F, Preliminary study: Purple sweet potato extract seems to be superior to increase the migration of impaired endothelial progenitor cells compared to L-Ascorbic acid. *Sci Pharm*, 87 (2019) 16.
24. Rizal MI & Sandra F, *Brucea javanica* leaf extract activates caspase-9 and caspase-3 of mitochondrial apoptotic pathway in human oral squamous cell carcinoma. *Indones Biomed J*, 8 (2016) 43.
25. Mutiah R, Listiyana A, Suryadinata A, Annisa R, Hakim A, Anggraini W & Susilowati R, Activity of inhibit the cell cycle and induct apoptosis in HeLa cancer cell with combination of sabrang onion (*Eleutherine palmifolia* (L.) Merr) and starfruit mistletoe (*Macrosolen cochinchinensis* (Lour.) Tiegh). *J App Pharm Sci*, 8 (2018) 122.
26. Pucci B, Kasten M & Giordano A, Cell cycle and apoptosis. *Neoplasia*, 2 (2000) 291.
27. Quadros Gomes AR, da Rocha Galucio NC, de Albuquerque KCO, Brígido HPC, Varela ELP, Castro ALG, Vale VV, Bahia MO, Rodriguez Burbano RM, de Molfeta FA, Carneiro LA, Percario S & Dolabela MF, Toxicity evaluation of *Eleutherine plicata* Herb. extracts and possible cell death mechanism. *Toxicol Rep*, 8 (2021) 1480.
28. Ellis IR, The migration and invasion of oral squamous carcinoma cells: Matrix, growth factor and signalling involvement. *Cancers*, 13 (2021) 2633.
29. Fan HX, Wang S, Zhao H, Liu N, Chen D, Sun M & Zheng JH, Sonic hedgehog signaling may promote invasion and metastasis of oral squamous cell carcinoma by activating MMP-9 and E-cadherin expression. *Med Oncol*, 31 (2014) 41.

30. Carballo GB, Honorato JR, de Lopes GPF & Spohr TCLSE, A highlight on Sonic hedgehog pathway. *Cell Commun Signal*, 16 (2018) 11.
31. Liang S, Zhang S, Wang P, Yang C, Shang C, Yang J & Wang J, LncRNA, TUG1 regulates the oral squamous cell carcinoma progression possibly via interacting with Wnt/ β -catenin signaling. *Gene*, 608 (2017) 49.

Figure and Figure Legends




Fig. 1. EBBE decreased the number of viable HSC-3 cells in a concentration-dependent manner. HSC-3 cells were cultured and treated with/without 1 µM Doxorubicin and various concentrations of EBBE for 24 h as indicated in the panel. The number of viable cells were measured with MTT assay as described in Materials and Methods. The data are expressed as mean ± SD (n=6). * $P < 0.05$ compared to the sham group. EBBE: *E. bulbosa* bulb extract.




Fig. 2. EBBE increased the percentage of apoptotic HSC-3 cells in a concentration-dependent manner. HSC-3 cells were cultured and treated with/without 1 µM Doxorubicin and various concentrations of EBBE for 24 h as indicated. Treated cells were then subjected to sub-G1 assay as described in Materials and Methods. (A) Histogram of flow cytometric results. (B) Percentage of apoptotic HSC-3 cells. The data are expressed as mean ± SD (n=18). * $P < 0.05$ compared to the sham group. EBBE: *E. bulbosa* bulb extract.




Fig. 3. EBBE inhibited migratory activities of HSC-3 cells. HSC-3 cell culture was scratched and treated with/without EBBE or 20 µM cyclopamine tartrate for 24 h as indicated. Gap closure was measured as described in Materials and Methods. (A) Scratched areas of HSC-3 cell culture. White bar: 100 µM. (B) Gap closure percentage of EBBE-treated HSC-3 cell. The data are expressed as mean ± SD (n=9). * $P < 0.05$ compared to the sham group. EBBE: *E. bulbosa* bulb extract.




Fig. 4. EBBE reduced the number of migrated HSC-3 cells. HSC-3 cell culture was treated with/without EBBE or 20 μ M cyclopamine tartrate for 24 h as indicated. Migrated cells were measured as described in Materials and Methods. The data are expressed as mean \pm SD (n=9). * P <0.05 compared to the sham group. EBBE: *E. bulbosa* bulb extract.




Fig. 5. EBBE reduced the expression of SHH. HSC-3 cell culture was treated with/without EBBE or 20 μ M cyclopamine tartrate for 24 h as indicated. HSC-3 cells were lysed, homogenized, and immunoblotted using anti-SHH antibody as described in Materials and Methods. The data are expressed as mean \pm SD (n=3). * P <0.05 compared to the sham group. EBBE: *E. bulbosa* bulb extract; SHH: Sonic hedgehog.



Ferry Sandra <ferry@trisakti.ac.id>

Ultra Urgent PRC IJBB#83

IJBB NIScPR <ijbbniscpr@gmail.com>
To: ferry@trisakti.ac.id

Mon, Sep 18, 2023 at 5:05 PM

 **IJBB#83.docx**

Dear Author,

Enclosed please find the edited Proof ready copy of the above MS. A suitably revised manuscript (in this case final version) in accordance with the editorial corrections/suggestions may please be submitted positively by **today** itself for finalization. Kindly use the enclosed Word file only for marking corrections (highlighted with red color) and submission.

- For 'references' section, strictly follow the pattern of IJBB [author names (initials without space), '&' before the last author and comma after the end of the author names; title of the paper followed by full stop, source (**abbreviated form of the journal following international standards and in italics without any full stop**) followed by non-italic comma, vol no., year within brackets followed by initial page no. followed by a full stop].
- In references, we give all author names. Et al. will not work in IJBB.
- Pl check the running title also.

Kindly send the corrected PRC by tomorrow.

Prof DN Rao
Executive editor
dnrao@iisc.ac.in
ijbb@niscpr.res.in

Eleutherine bulbosa Bulb Extract Induces Apoptosis and Inhibits Cell Migration by Down regulating Sonic Hedgehog Gene in Human Tongue Cancer Cells: An *in vitro* Study

Ferry Sandra^{1*}, Kelvin Suryajaya², Angliana Chouw³, Maria Celinna⁴, Dewi Ranggaini⁵ & Cynthia Retna Sartika³

¹Department of Biochemistry and Molecular Biology, Division of Oral Biology, Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia

²Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia

³PT Prodia Stem Cell Indonesia, Jakarta Pusat-10430, Jakarta, Indonesia

⁴The Prodia Education and Research Institute, Jakarta Pusat-10430, Jakarta, Indonesia

⁵Department of Physiology, Division of Oral Biology, Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia

Received 09 May 2023; revised 18 September 2023

Eleutherine bulbosa (Mill.) Urb. is a medicinal plant which has long been used to treat cancer. *E. bulbosa* bulb extract (EBBE) has been reported to show cytotoxicity towards several types of human cancer. However, the cytotoxic effect of EBBE towards oral cancer cells has not been investigated. The present study aimed to evaluate the effects of EBBE towards the viability, apoptosis, and migratory activities of oral cancer cells. Human oral squamous cell carcinoma (HSC)-3 cells were treated with various concentrations of EBBE for 24 h. The number of viable and apoptotic HSC-3 cells were measured using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) and sub-G1 assay, respectively. The migratory activities of HSC-3 cells were assessed using scratch and transwell assay. Sonic hedgehog (SHH) expression was measured using immunoblotting. Upon EBBE treatment, apoptotic HSC-3 cells were significantly higher in a concentration-dependent manner ($P < 0.05$). Meanwhile, viability, migratory activities, and migrated HSC-3 cell number were significantly lower in a concentration-dependent manner ($P < 0.05$). The SHH expression levels in EBBE-treated HSC-3 cells were also lower in a concentration-dependent manner. EBBE reduces HSC-3 cell viability through apoptosis and inhibits its migratory activities by downregulating SHH expression.

Keywords: Apoptosis, Cell migration, *Eleutherine bulbosa*, Sonic hedgehog, Tongue cancer

Oral squamous cell carcinoma (OSCC) is the most common histological type of oral cancer¹ which arises from squamous epithelial cells in the oral cavity. This cancer generally affects the floor of the mouth and the tongue². Various modes of treatment are developed to manage OSCC, including chemotherapy, radiotherapy, surgery, or its combination³. However, these treatments cause numerous side effects.

Discovery and development of new therapeutic agents derived from natural sources, particularly plants with less side effects are recently one of the main focuses on oncology research. Several plants have been reported to have anticancer effects, including *Brucea javanica*⁴, *Cinnamomum cassia*⁵, *Myristica fragrans*⁶, and *Oroxylum indicum*⁷. Plants have been known as rich sources of various active compounds, including

flavonoids, which was reported to inhibit cancer proliferation *via* cell cycle arrest, antiproliferative and antioxidative activity, apoptosis induction, promotion of differentiation, inhibition of angiogenesis and carcinogen metabolic activation, and multidrug resistance modulation⁸. In addition, several non-flavonoid phenolic compounds have also been reported to show anticancer properties, such as caffeic acid⁹.

Eleutherine bulbosa (Mill.) Urb. is a traditional medicinal plant belonging to the Iridaceae family. Its bulbs are commonly used for the treatment of breast cancer, hypertension, stroke, and diabetes mellitus. It has also been used to increase the production of breast milk and treat sexual disorder¹⁰. *E. bulbosa* bulbs have also been reported to have antimicrobial¹¹ and antioxidant¹² activity.

E. bulbosa contains a wide variety of compounds, primarily naphthoquinones, anthraquinones, and naphthalene¹³. The main compounds found in this plant are eleutherin, isoeleutherin, and eleutherol¹⁴. Several

*Correspondence:

Phone: +628128077780

E-mail: ferry@trisakti.ac.id

compounds obtained from *E. bulbosa* bulbs have been reported to show anticancer activities, such as 6,8-hydroxy-3,4-di-methoxy-1-methyl-anthraquinone-2-carboxylic acid methyl ester¹⁵, eleutherinoside C, and isoeleutherin¹⁶.

E. bulbosa bulb extract (EBBE) has been reported to show cytotoxicity towards several types of human cancer, such as cervical cancer (line HeLa)¹⁷, colon cancer (line SW480, HCT116, DLD1, WiDr)^{16,18}, breast cancer (line T47D)¹⁹, leukemia (line K562)¹⁵, and retinoblastoma (WERI-Rb-1)²⁰. However, the cytotoxic effect of EBBE towards human oral cancer cells has not been investigated. Therefore, the present study was conducted to evaluate the effects of EBBE towards the viability, apoptosis, and migratory activities of oral cancer cells.

Materials and Methods

E. bulbosa bulb extract (EBBE) preparation

E. bulbosa bulbs were obtained from Indonesian Medicinal and Aromatic Crops Research Institute (IMACRI), Indonesia. Plant identification was carried out by expert botanist at Research Center for Plant Conservation and Botanic Gardens, Indonesian Institute of Sciences, Indonesia (No. B-1269/IPH.3/KS/X/2020). EBBE was extracted by maceration method⁴. Briefly, bulbs of *E. bulbosa* were minced and dried. The dried material was extracted with ethanol 70%, filtered and evaporated using a rotary evaporator. The resulting crude EBBE was then stored at 4°C.

HSC-3 cell culture

HSC-3 cell culture was performed as previously described⁴ with modification. The HSC-3 cell line was purchased from Sigma-Aldrich Pte. Ltd. (St. Louis, MO, USA). HSC-3 cells were cultured in a complete medium containing Dulbecco's modified Eagle's medium (DMEM) (Sigma-Aldrich), 10% fetal bovine serum (FBS) (PAN-Biotech, Aidenbach, Germany), 50 U/mL penicillin and 50 µg/mL streptomycin (Sigma-Aldrich) in a humidified 5% CO₂, 37°C incubator. HSC-3 cells were then dissociated with trypsin-ethylenediamine tetraacetic acid (EDTA) solution (Sigma-Aldrich) after reaching 80% confluency. Upon reaching the desired cell number, HSC-3 cells were seeded into 24-well plate and 96-well plate for further assays.

Cell viability assay

The number of viable cells were measured quantitatively using 3-(4,5-dimethylthiazol-2-yl)-2,5-

diphenyltetrazolium bromide (MTT) assay as previously described²¹ with modification. Briefly, 5×10^3 HSC-3 cells were seeded into 96-well plates and then treated with EBBE (1, 10, or 100 µg/mL), 1 µM Doxorubicin (Dankos Farma, Jakarta, Indonesia) or medium merely for 24 h. After that 100 µL of MTT (Sigma-Aldrich) in culture medium was loaded into each well. After incubating the plates for 4 h, culture medium was discarded, and the formed formazan crystals were dissolved in 100 µL dimethylsulfoxide (DMSO). Results were measured at OD₅₇₀ using a microplate reader (Bio-Rad, Hercules, CA, USA). Each experimental group was measured in six replicates. In addition, the number of untreated cells were counted with hemacytometer and used for interpolating OD₅₇₀ values of HSC-3 cells.

Sub-G1 assay

To investigate the cytotoxic effect of EBBE on HSC-3 cells, the number of apoptotic cells were determined using sub-G1 assay as previously described⁹ with modification. Treated and untreated HSC-3 cells were harvested and suspended in 450 µL hypotonic fluorochrome solution containing 50 µg/mL propidium iodide (Sigma-Aldrich), 0.1% Triton X-100 (Sigma-Aldrich), and 0.1% sodium citrate (Wako, Osaka, Japan). After that, cell suspensions were incubated in the dark for 2 h at room temperature. Fluorescence of individual nuclei was measured using FACSCanto II flow cytometer (Becton Dickinson, Franklin Lakes, NJ, USA) at 100,000 events.

Scratch assay

Scratch assay was performed as previously described²². HSC-3 cells (5×10^4) were seeded into 24-well plates and incubated overnight. Scratch was made on each well with micropipette yellow tips. Then, the wells were washed with phosphate-buffered saline (PBS) (Wako) and treated with/without EBBE or 20 µM cyclopamine tartrate (BioVision, Milpitas, CA, USA) (as positive control) for 24 h. The gap closures were observed and documented under an inverted light microscope (Carl Zeiss, Jena, Germany).

Transwell assay

Transwell assay was performed as previously described²³ with modification. HSC-3 cells (5×10^4) were seeded in the 8-µm pore size chamber insert (Merck, Darmstadt, Germany). After that, 800 µL culture medium with/without EBBE or 20 µM cyclopamine tartrate was loaded into each well of the

24-well plate. HSC-3-containing chamber inserts were then placed on the 24-well plate. After incubating the chamber for 24 h at 37°C, migrated cells in the lower surface of the chamber were fixed. The cells were counted and documented under an inverted light microscope.

Immunoblotting

Expression levels of Sonic hedgehog (SHH) protein were measured using immunoblotting method as previously described²⁴ with modification. HSC-3 cells were lysed and homogenized with Laemmli sample buffer (Bio-Rad). The proteins from each experimental group were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and transferred to a polyvinylidene difluoride (PVDF) sheet. After blocking with 5% skim milk solution, the sheet was incubated with rabbit monoclonal anti-SHH antibody as a primary antibody and horseradish peroxidase (HRP)-conjugated goat anti-rabbit antibody as a secondary antibody (Cell Signaling Technology, Danvers, MA, USA). Immunoblot results were analyzed using ImageJ software (National Institute of Health, Bethesda, MD, USA) to measure the density of each SHH band.

Statistical analysis

Statistical analyses were performed using IBM SPSS Statistics version 26.0 (SPSS IBM, Armonk, NY, USA). Shapiro-Wilk test was used as a normality test. Differences of MTT assay, sub-G1 assay, and scratch assay results between experimental groups were analyzed using Kruskal-Wallis H with Mann-Whitney U *post hoc* analysis. Meanwhile, differences of transwell assay results and SHH band density between experimental groups were analyzed using one-way analysis of variances (ANOVA) with Tukey's honestly significant difference (HSD) *post hoc* analysis. All results were expressed as mean \pm standard deviation (SD). $P < 0.05$ were considered as statistically significant.

Results

EBBE reduced the number of viable HSC-3 cells

The number of viable HSC-3 cells cultured in medium containing extract solvent (sham group) for 24 h was $9,607 \pm 13.81$. Treatment with EBBE showed that the viability of HSC-3 cells was significantly lower compared to the sham group ($P < 0.05$) in a concentration-dependent manner. The number of viable HSC-3 cells treated with 1, 10, and 100 $\mu\text{g/mL}$

EBBE were $9,055 \pm 268.69$, $6,777 \pm 278.04$, and $3,659 \pm 32.86$, respectively. The viability of EBBE-treated HSC-3 cells was significantly higher ($P < 0.05$) than Doxorubicin-treated HSC-3 cells (226 ± 32.34) (Fig. 1). These findings suggested that EBBE could reduce the viability of HSC-3 cells.

EBBE induced apoptosis of HSC-3 cells

Sub-G1 assay results showed that the apoptosis percentage of the sham group was $8.27 \pm 0.14\%$. Apoptotic HSC-3 cells were significantly higher compared to the sham group ($P < 0.05$) in concentration-dependent manner after EBBE treatment. Apoptosis percentages of HSC-3 cells treated with 1, 10, and 100 $\mu\text{g/mL}$ EBBE were $9.60 \pm 0.35\%$, $17.78 \pm 0.32\%$, and $38.64 \pm 0.70\%$, respectively. These percentages were significantly lower ($P < 0.05$) than Doxorubicin-treated HSC-3 cells ($95.95 \pm 2.60\%$) (Fig. 2). Therefore, these results suggested that EBBE treatment could promote apoptosis of HSC-3 cells.

EBBE inhibited HSC-3 cell migration

The gap closure percentage of HSC-3 cells cultured in medium containing extract solvent after 24 h was $96.11 \pm 4.17\%$. After EBBE treatment, the migratory activities of HSC-3 cells were significantly lower compared to the sham group ($P < 0.05$) in a concentration-dependent manner. Gap closure percentage of HSC-3 cells treated with 1, 10, and 100 $\mu\text{g/mL}$ EBBE were $89.56 \pm 5.77\%$, $76.78 \pm 4.87\%$, and




Fig. 1 — EBBE decreased the number of viable HSC-3 cells in a concentration-dependent manner. HSC-3 cells were cultured and treated with/without 1 μM Doxorubicin and various concentrations of EBBE for 24 h as indicated in the panel. The number of viable cells were measured with MTT assay as described in Materials and Methods. The data are expressed as mean \pm SD ($n=6$). $*P < 0.05$ compared to the sham group




Fig. 2 — EBBE increased the percentage of apoptotic HSC-3 cells in a concentration-dependent manner. HSC-3 cells were cultured and treated with/without 1 μM Doxorubicin and various concentrations of EBBE for 24 h as indicated. Treated cells were then subjected to sub-G1 assay as described in Materials and Methods. (A) Histogram of flow cytometric results. (B) Percentage of apoptotic HSC-3 cells. The data are expressed as mean \pm SD (n=18). * $P < 0.05$ compared to the sham group

60.67 \pm 7.62%, respectively. These percentages were significantly higher ($P < 0.05$) than cyclopamine tartrate-treated HSC-3 cells (43.00 \pm 4.82%) (Fig. 3).

Transwell assay also showed similar results. The number of migrated HSC-3 cells in the sham group was 134.44 \pm 11.60. The number of migrated HSC-3 cells was significantly lower than the sham group ($P < 0.05$) in a concentration-dependent manner after EBBE treatment. The number of migrated HSC-3 cells treated with 1, 10, and 100 $\mu\text{g/mL}$ EBBE were 107.44 \pm 15.21, 95.11 \pm 6.01, and 39.22 \pm 3.56, respectively. These numbers were significantly higher ($P < 0.05$) than cyclopamine tartrate-treated HSC-3 cells (17.89 \pm 3.22) (Fig. 4). Overall, these results indicated that EBBE exhibited an anti-migratory effect towards HSC-3 cells.

EBBE down regulated the expression of SHH

Upon the addition of EBBE, the SHH expression levels in HSC-3 cells were lower in a concentration-dependent manner. Expression level of SHH in HSC-3 cells treated with 1 $\mu\text{g/mL}$ EBBE was significantly higher ($P < 0.05$), while SHH expression levels in HSC-3 cells treated with 10 and 100 $\mu\text{g/mL}$ EBBE were significantly lower ($P < 0.05$) compared to the sham group. SHH expression levels in HSC-3 cells treated with 1 and 10 $\mu\text{g/mL}$ EBBE were significantly higher compared to cyclopamine tartrate-treated HSC-3 cells ($P < 0.05$). However, no significant difference was observed in the SHH expression levels between cyclopamine tartrate and 100 $\mu\text{g/mL}$ EBBE treatment ($P = 0.142$) (Fig. 5). Thus, these findings suggested that EBBE could decrease the expression of SHH.




Fig. 3 — EBBE inhibited migratory activities of HSC-3 cells. HSC-3 cell culture was scratched and treated with/without EBBE or 20 µM cyclopamine tartrate for 24 h as indicated. Gap closure was measured as described in Materials and Methods. (A) Scratched areas of HSC-3 cell culture. White bar: 100 µM. (B) Gap closure percentage of EBBE-treated HSC-3 cell. The data are expressed as mean ± SD (n=9). * $P < 0.05$ compared to the sham group




Fig. 4 — EBBE reduced the number of migrated HSC-3 cells. HSC-3 cell culture was treated with/without EBBE or 20 µM cyclopamine tartrate for 24 h as indicated. Migrated cells were measured as described in Materials and Methods. The data are expressed as mean ± SD (n=9). * $P < 0.05$ compared to the sham group




Fig. 5 — EBBE reduced the expression of SHH. HSC-3 cell culture was treated with/without EBBE or 20 µM cyclopamine tartrate for 24 h as indicated. HSC-3 cells were lysed, homogenized, and immunoblotted using anti-SHH antibody as described in Materials and Methods. The data are expressed as mean ± SD (n=3). * $P < 0.05$ compared to the sham group

Discussion

In the present study, we investigated the potency of EBBE as anticancer agent for OSCC treatment. EBBE showed a cytotoxic effect towards HSC-3 cells by reducing the viability of HSC-3 cells in a concentration-dependent manner. According to the sub-G1 assay results, the decrease in HSC-3 cell viability was caused by apoptosis. These results were in line with previous studies that EBBE was able to induce apoptosis of cancer cells^{18-20,25}. EBBE was also reported to inhibit cell cycle in several cell lines^{19,25}, indicating that EBBE may induce apoptosis of HSC-3 cells through cell cycle arrest since both processes are interconnected²⁶. Based on a molecular docking study, eleutherol, eleutherin, and isoeleutherin could induce apoptosis by activating caspase-8²⁷.

Cell migration is one of the key processes involved in cancer metastasis²⁸. Scratch and transwell assays demonstrated that EBBE restrained the migration of HSC-3 cells in a concentration-dependent manner. This showed the novel activity of EBBE in impairing HSC-3 cell migration. Therefore, EBBE may have anti-metastasis activity towards tongue cancer cells.

Metastasis of tongue cancer is regulated by several proteins, including SHH. One of the SHH signaling pathways that has been reported to regulate cell migration in tongue cancer is SHH/glioma-associated oncogene homologue (SHH/GLI)²⁹. SHH protein binds and inactivates its receptor, Patched 1 (PTCH1). This interaction causes the activation of transcription factors from the GLI family, which play a major role in regulating cancer migration, invasion, and proliferation³⁰.

The present study revealed that EBBE decreased SHH expression levels along with the increase in EBBE concentrations, suggesting that the inhibitory effect of EBBE towards the expression of SHH may be responsible for its anti-migratory property. To the best of our knowledge, this is the first study demonstrating SHH downregulation in EBBE-treated HSC-3 cells. EBBE may inhibit HSC-3 cell migration via inhibition of SHH/GLI signaling pathway. Possibly, all signaling pathways that are regulated by SHH can be inhibited by EBBE. EBBE may also restrain HSC-3 cell migration through another pathway. In SW480 colon cancer cell line, eleutherinoside C and isoeleutherin obtained from *E. bulbosa* bulbs was reported to inhibit Wnt/ β -catenin pathway¹⁶, which has been reported to play an important role in regulating cancer migration³¹.

Therefore, signaling pathways involved in the inhibition of HSC-3 cell migration by EBBE still need to be explored. In addition, signaling components downstream to SHH and target genes that are affected by EBBE should be identified. Since EBBE also contains various bioactive compounds, further research is needed to identify which compounds are responsible for apoptosis and anti-migratory activity. Thus, more investigations are required to understand the molecular mechanisms underlying EBBE-induced HSC-3 cell apoptosis and confirm which SHH signaling pathway contributes to the inhibition of cell migration by EBBE.

Conclusion

EBBE reduces HSC-3 cell viability through apoptosis and inhibits its migratory activities by down regulating SHH expression. Taken together, EBBE could be a potential anti-tongue cancer agent, particularly as an ant metastatic agent.

Conflict of interest

All authors declare no conflicts of interest.

References

- 1 Chen YK, Huang HC, Lin LM & Lin CC, Primary oral squamous cell carcinoma: An analysis of 703 cases in southern Taiwan. *Oral Oncol*,35 (1999) 173.
- 2 Neville BW & Day TA, Oral cancer and precancerous lesions. *CA Cancer J Clin*,52 (2002) 195.
- 3 Minhas S, Kashif M, Altaf W, Afzal N & Nagi AH, Concomitant-chemoradiotherapy-associated oral lesions in patients with oral squamous-cell carcinoma. *Cancer Biol Med*, 14(2017) 176.
- 4 Wicaksono BD, Tangkearung E & Sandra F, Brucea javanica leaf extract induced apoptosis in human oral squamous cell carcinoma (HSC2) cells by attenuation of mitochondrial membrane permeability. *Indones Biomed J*, 7(2015) 107.
- 5 5Chang WL, Cheng FC, Wang SP, Chou ST & Shih Y, Cinnamomum cassia essential oil and its major constituent cinnamaldehyde induced cell cycle arrest and apoptosis in human oral squamous cell carcinoma HSC-3 cells. *Environ Toxicol*, 32 (2017) 456.
- 6 Susianti S, Lesmana R, Salam S, Julaeha E, Pratiwi YS, Sylviana N, Goenawan H, Kurniawan A & Supratman U, The effect of nutmeg seed (*M. fragrans*) extracts induces apoptosis in melanoma maligna cell's (B16-F10). *Indones Biomed J*, 13 (2021) 68.
- 7 Kameyanda Poonacha S, Harishkumar M, Radha M, Varadarajan R, Nalilu SK, Shetty SS, Shetty PK, Chandrashekarappa RB, Sreenivas MG & Bhandary Bavabeedu SK, Insight into oroxylinA-7-O- β -d-glucuronide-enriched *Oroxylum indicum* bark extract in oral cancer HSC-3 cell apoptotic mechanism: Role of mitochondrial microenvironment. *Molecules*, 26 (2021) 7430.
- 8 Ren W, Qiao Z, Wang H, Zhu L & Zhang L, Flavonoids: Promising anticancer agents. *Med Res Rev*, 23 (2003) 519.

- 9 Sandra F&Sidharta MA, Caffeic acid induced apoptosis in MG63 osteosarcoma cells through activation of caspases. *Mol Cell Biomed Sci*, 1(2017) 28.
- 10 Ieyama T, Gunawan-Puteri MDPT& Kawabata J, α -Glucosidase inhibitors from the bulb of *Eleutherine americana*. *Food Chem*, 128 (2011) 308.
- 11 Subramaniam K, Suriyamoorthy S, Wahab F, Sharon FB&Rex GR, Antagonistic activity of *Eleutherine palmifolia* Linn. *Asian Pac J Trop Dis*, 2 (2012) S491.
- 12 Kuntorini EM, Dewi M& Misrina M, Anatomical structure and antioxidant activity of red bulb plant (*Eleutherine americana*) on different plant age. *Biodiversitas*, 17 (2016) 229.
- 13 Insanu M, Kusumardiyani S& Hartati R, Recent studies on phytochemicals and pharmacological effects of *Eleutherine americana* Merr. *Procedia Chem*, 13 (2014) 221.
- 14 Chen Z, Huang H, Wang C, Li Y, Ding J, Sankawa U, Noguchi H & Iitaka Y, Hongconin, a new naphthalene derivative from hong-cong, the rhizome of *Eleutherine americana* Merr. et Heyne (Iridaceae). *Chem Pharm Bull*, 34 (1986) 2743.
- 15 Xu J, Qiu F, Duan W, Qu G, Wang N& Yao X, New bioactive constituents from *Eleutherine americana*. *Front Chem China*, 1 (2006) 320.
- 16 Li X, Ohtsuki T, Koyano T, Kowithayakorn T& Ishibashi M, New Wnt/ β -catenin signaling inhibitors isolated from *Eleutherine palmifolia*. *Chem Asian J*, 4(2009) 540.
- 17 Mutiah R, Minggarwati TS, Kristanti RA& Susanti E, Compound identification and anticancer activity of ethyl acetate fraction from bawang sabrang (*Eleutherine palmifolia* (L.) Merr.) on HeLa cervical cancer cell line. *Indones J Cancer Chemoprevent*, 10 (2019) 131.
- 18 Lubis IA, Ichwan M, Mustofa M & Satria D, Anticancer activity of *Eleutherine bulbosa* (Mill.) Urb. extract on WiDr cell line in vitro, paper presented to 2nd Public Health International Conference (PHICo 2017), Faculty of Public Health – Universitas Sumatera Utara, Medan, Indonesia, 18-19 December 2017.
- 19 Yuniarti A, Sundhani E& Nurulita NA, The potentiation effect of bawang dayak (*Sisyrinchium palmifolium* L.) extract on T47D cell growth inhibition after 5-fluorouracil treatment. *Pharmaciana*, 8(2018) 195.
- 20 Kamarudin AA, Sayuti NH, Saad N, Razak NAA& Esa NM, Induction of apoptosis by *Eleutherine bulbosa* (Mill.) Urb. bulb extracted under optimised extraction condition on human retinoblastoma cancer cells (WERI-Rb-1). *J Ethnopharmacol*, 284 (2022) 114770.
- 21 Sandra F, Sudiono J, Trisfilha P& Pratiwi D, Cytotoxicity of *Alpinia galanga* rhizome crude extract on NIH-3T3 cells. *Indones Biomed J*, 9(2017) 23. <https://doi.org/10.18585/inabj.v9i1.212>.
- 22 Ariesanti Y, Sandra F, Claresta B& Alvita L, Coffea canephora bean extract induces NIH3T3 cell migration. *Indones Biomed J*, 13(2021) 216. <https://doi.org/10.18585/inabj.v13i2.1522>
- 23 Oktaviono YH, Al-Farabi MJ, Suastika LOS, Hartono F, Dirgantara Y& Sandra F, Preliminary study: Purple sweet potato extract seems to be superior to increase the migration of impaired endothelial progenitor cells compared to L-Ascorbic acid. *Sci Pharm*, 87(2019) 16.
- 24 Rizal MI& Sandra F, Brucea javanica leaf extract activates caspase-9 and caspase-3 of mitochondrial apoptotic pathway in human oral squamous cell carcinoma. *Indones Biomed J*, 8 (2016) 43.
- 25 Mutiah R, Listiyana A, Suryadinata A, Annisa R, Hakim A, Angraini W & Susilowati R, Activity of inhibit the cell cycle and induct apoptosis in HeLa cancer cell with combination of sabrang onion (*Eleutherine palmifolia* (L.) Merr) and starfruit mistletoe (*Macrosolen cochinchinensis* (Lour.) Tiegh). *J App Pharm Sci*, 8 (2018) 122.
- 26 Pucci B, Kasten M& Giordano A, Cell cycle and apoptosis. *Neoplasia*, 2(2000) 291.
- 27 Quadros Gomes AR, da Rocha Galucio NC, de Albuquerque KCO, Brígido HPC, Varela ELP, Castro ALG, Vale VV, Bahia MO, Rodriguez Burbano RM, de Molfeta FA, Carneiro LA, Percario S& Dolabela MF, Toxicity evaluation of *Eleutherine plicata* Herb. extracts and possible cell death mechanism. *Toxicol Rep*, 8 (2021) 1480.
- 28 Ellis IR, The migration and invasion of oral squamous carcinoma cells: Matrix, growth factor and signalling involvement. *Cancers*, 13 (2021) 2633.
- 29 Fan HX, Wang S, Zhao H, Liu N, Chen D, Sun M& Zheng JH, Sonic hedgehog signaling may promote invasion and metastasis of oral squamous cell carcinoma by activating MMP-9 and E-cadherin expression. *Med Oncol*, 31 (2014) 41.
- 30 Carballo GB, Honorato JR, de Lopes GPF& Spohr TCLSE, A highlight on Sonic hedgehog pathway. *Cell Commun Signal*, 16 (2018) 11.
- 31 Liang S, Zhang S, Wang P, Yang C, Shang C, Yang J & Wang J, LncRNA, TUG1 regulates the oral squamous cell carcinoma progression possibly via interacting with Wnt/ β -catenin signaling. *Gene*, 608 (2017) 49.



Ferry Sandra <ferry@trisakti.ac.id>

Ultra Urgent PRC IJBB#83

Ferry Sandra <ferry@trisakti.ac.id>
To: ijbbniscpr@gmail.com

Thu, Sep 21, 2023 at 4:45 PM

Dear Team IJBB,

Good day. Herewith I put the Google Drive link for the corrected PRC of manuscript IJBB#83.
<https://docs.google.com/document/d/18S0pXM8C9IgeQ0lj1pa6i0HwnGjC0ILL/edit?usp=sharing&oid=110414335342504619051&rtpof=true&sd=true>

There are some corrections in this PRC:

1. In the title of the manuscript, please delete the word "Gene" (highlighted in red, double strikethrough).
2. In the affiliation list, we added one more author (Kyung Hoon Lee) and his affiliation (highlighted in red).
3. In the 4th paragraph of the Discussion section, we changed the word "genes" to "molecules" (highlighted in red).
4. In Reference no. 18, please delete the 'h' in the word "Publich" (highlighted in red, strikethrough).

Hopefully you can find the PRC well. Thank you. I wish you a nice day.

Regards,
Ferry Sandra
[Quoted text hidden]

--

Ferry Sandra, D.D.S., Ph.D.
Head of Medical Research Center
Universitas Trisakti

Eleutherine bulbosa Bulb Extract Induces Apoptosis and Inhibits Cell Migration by Downregulating Sonic Hedgehog Gene in Human Tongue Cancer Cells: An *in vitro* Study

Ferry Sandra^{1*}, Kelvin Suryajaya², Angliana Chouw³, Maria Celinna⁴, Dewi Ranggaini⁵, **Kyung Hoon Lee⁶** & Cynthia Retna Sartika³

¹Department of Biochemistry and Molecular Biology, Division of Oral Biology, Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia

²Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia

³PT Prodia Stem Cell Indonesia, Jakarta Pusat-10430, Jakarta, Indonesia

⁴The Prodia Education and Research Institute, Jakarta Pusat-10430, Jakarta, Indonesia

⁵Department of Physiology, Division of Oral Biology, Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia

⁶Research Institute, Ballys Co. Ltd, Incheon-22219, Republic of Korea

Received 09 May 2023; revised 18 September 2023

Eleutherine bulbosa (Mill.) Urb. is a medicinal plant which has long been used to treat cancer. *E. bulbosa* bulb extract (EBBE) has been reported to show cytotoxicity towards several types of human cancer. However, the cytotoxic effect of EBBE towards oral cancer cells has not been investigated. The present study aimed to evaluate the effects of EBBE towards the viability, apoptosis, and migratory activities of oral cancer cells. Human oral squamous cell carcinoma (HSC)-3 cells were treated with various concentrations of EBBE for 24 h. The number of viable and apoptotic HSC-3 cells were measured using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) and sub-G1 assay, respectively. The migratory activities of HSC-3 cells were assessed using scratch and transwell assay. Sonic hedgehog (SHH) expression was measured using immunoblotting. Upon EBBE treatment, apoptotic HSC-3 cells were significantly higher in a concentration-dependent manner ($P < 0.05$). Meanwhile, viability, migratory activities, and migrated HSC-3 cell number were significantly lower in a concentration-dependent manner ($P < 0.05$). The SHH expression levels in EBBE-treated HSC-3 cells were also lower in a concentration-dependent manner. EBBE reduces HSC-3 cell viability through apoptosis and inhibits its migratory activities by downregulating SHH expression.

Keywords: Apoptosis, Cell migration, *Eleutherine bulbosa*, Sonic hedgehog, Tongue cancer

Oral squamous cell carcinoma (OSCC) is the most common histological type of oral cancer¹ which arises from squamous epithelial cells in the oral cavity. This cancer generally affects the floor of the mouth and the tongue². Various modes of treatment are developed to manage OSCC, including chemotherapy, radiotherapy, surgery, or its combination³. However, these treatments cause numerous side effects.

Discovery and development of new therapeutic agents derived from natural sources, particularly plants with less side effects are recently one of the main focuses on oncology research. Several plants have been reported to have anticancer effects, including *Brucea javanica*⁴, *Cinnamomum cassia*⁵, *Myristica fragrans*⁶,

and *Oroxylum indicum*⁷. Plants have been known as rich sources of various active compounds, including flavonoids, which was reported to inhibit cancer proliferation *via* cell cycle arrest, antiproliferative and antioxidative activity, apoptosis induction, promotion of differentiation, inhibition of angiogenesis and carcinogen metabolic activation, and multidrug resistance modulation⁸. In addition, several non-flavonoid phenolic compounds have also been reported to show anticancer properties, such as caffeic acid⁹.

Eleutherine bulbosa (Mill.) Urb. is a traditional medicinal plant belonging to the Iridaceae family. Its bulbs are commonly used for the treatment of breast cancer, hypertension, stroke, and diabetes mellitus. It has also been used to increase the production of breast milk and treat sexual disorder¹⁰. *E. bulbosa* bulbs have also been reported to have antimicrobial¹¹ and antioxidant¹² activity.

*Correspondence:
Phone: +628128077780
E-mail: ferry@trisakti.ac.id

E. bulbosa contains a wide variety of compounds, primarily naphthoquinones, anthraquinones, and naphthalene¹³. The main compounds found in this plant are eleutherin, isoeleutherin, and eleutherol¹⁴. Several compounds obtained from *E. bulbosa* bulbs have been reported to show anticancer activities, such as 6,8-hydroxy-3,4-di-methoxy-1-methyl-anthraquinone-2-carboxylic acid methyl ester¹⁵, eleutherinoside C, and isoeleutherin¹⁶.

E. bulbosa bulb extract (EBBE) has been reported to show cytotoxicity towards several types of human cancer, such as cervical cancer (line HeLa)¹⁷, colon cancer (line SW480, HCT116, DLD1, WiDr)^{16,18}, breast cancer (line T47D)¹⁹, leukemia (line K562)¹⁵, and retinoblastoma (WERI-Rb-1)²⁰. However, the cytotoxic effect of EBBE towards human oral cancer cells has not been investigated. Therefore, the present study was conducted to evaluate the effects of EBBE towards the viability, apoptosis, and migratory activities of oral cancer cells.

Materials and Methods

E. bulbosa bulb extract (EBBE) preparation

E. bulbosa bulbs were obtained from Indonesian Medicinal and Aromatic Crops Research Institute (IMACRI), Indonesia. Plant identification was carried out by expert botanist at Research Center for Plant Conservation and Botanic Gardens, Indonesian Institute of Sciences, Indonesia (No. B-1269/IPH.3/KS/X/2020). EBBE was extracted by maceration method⁴. Briefly, bulbs of *E. bulbosa* were minced and dried. The dried material was extracted with ethanol 70%, filtered and evaporated using a rotary evaporator. The resulting crude EBBE was then stored at 4°C.

HSC-3 cell culture

HSC-3 cell culture was performed as previously described⁴ with modification. The HSC-3 cell line was purchased from Sigma-Aldrich Pte. Ltd. (St. Louis, MO, USA). HSC-3 cells were cultured in a complete medium containing Dulbecco's modified Eagle's medium (DMEM) (Sigma-Aldrich), 10% fetal bovine serum (FBS) (PAN-Biotech, Aidenbach, Germany), 50 U/mL penicillin and 50 µg/mL streptomycin (Sigma-Aldrich) in a humidified 5% CO₂, 37°C incubator. HSC-3 cells were then dissociated with trypsin-ethylenediamine tetraacetic acid (EDTA) solution (Sigma-Aldrich) after reaching 80% confluency. Upon reaching the desired cell number, HSC-3 cells were seeded into 24-well plate

and 96-well plate for further assays.

Cell viability assay

The number of viable cells were measured quantitatively using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay as previously described²¹ with modification. Briefly, 5×10^3 HSC-3 cells were seeded into 96-well plates and then treated with EBBE (1, 10, or 100 µg/mL), 1 µM Doxorubicin (Dankos Farma, Jakarta, Indonesia) or medium merely for 24 h. After that 100 µL of MTT (Sigma-Aldrich) in culture medium was loaded into each well. After incubating the plates for 4 h, culture medium was discarded, and the formed formazan crystals were dissolved in 100 µL dimethylsulfoxide (DMSO). Results were measured at OD₅₇₀ using a microplate reader (Bio-Rad, Hercules, CA, USA). Each experimental group was measured in six replicates. In addition, the number of untreated cells were counted with hemacytometer and used for interpolating OD₅₇₀ values of HSC-3 cells.

Sub-G1 assay

To investigate the cytotoxic effect of EBBE on HSC-3 cells, the number of apoptotic cells were determined using sub-G1 assay as previously described⁹ with modification. Treated and untreated HSC-3 cells were harvested and suspended in 450 µL hypotonic fluorochrome solution containing 50 µg/mL propidium iodide (Sigma-Aldrich), 0.1% Triton X-100 (Sigma-Aldrich), and 0.1% sodium citrate (Wako, Osaka, Japan). After that, cell suspensions were incubated in the dark for 2 h at room temperature. Fluorescence of individual nuclei was measured using FACSCanto II flow cytometer (Becton Dickinson, Franklin Lakes, NJ, USA) at 100,000 events.

Scratch assay

Scratch assay was performed as previously described²². HSC-3 cells (5×10^4) were seeded into 24-well plates and incubated overnight. Scratch was made on each well with micropipette yellow tips. Then, the wells were washed with phosphate-buffered saline (PBS) (Wako) and treated with/without EBBE or 20 µM cyclopamine tartrate (BioVision, Milpitas, CA, USA) (as positive control) for 24 h. The gap closures were observed and documented under an inverted light microscope (Carl Zeiss, Jena, Germany).

Transwell assay

Transwell assay was performed as previously

described²³ with modification. HSC-3 cells (5×10^4) were seeded in the 8- μ m pore size chamber insert (Merck, Darmstadt, Germany). After that, 800 μ L culture medium with/without EBBE or 20 μ M cyclopamine tartrate was loaded into each well of the 24-well plate. HSC-3-containing chamber inserts were then placed on the 24-well plate. After incubating the chamber for 24 h at 37°C, migrated cells in the lower surface of the chamber were fixed. The cells were counted and documented under an inverted light microscope.

Immunoblotting

Expression levels of Sonic hedgehog (SHH) protein were measured using immunoblotting method as previously described²⁴ with modification. HSC-3 cells were lysed and homogenized with Laemmli sample buffer (Bio-Rad). The proteins from each experimental group were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and transferred to a polyvinylidene difluoride (PVDF) sheet. After blocking with 5% skim milk solution, the sheet was incubated with rabbit monoclonal anti-SHH antibody as a primary antibody and horseradish peroxidase (HRP)-conjugated goat anti-rabbit antibody as a secondary antibody (Cell Signaling Technology, Danvers, MA, USA). Immunoblot results were analyzed using ImageJ software (National Institute of Health, Bethesda, MD, USA) to measure the density of each SHH band.

Statistical analysis

Statistical analyses were performed using IBM SPSS Statistics version 26.0 (SPSS IBM, Armonk, NY, USA). Shapiro-Wilk test was used as a normality test. Differences of MTT assay, sub-G1 assay, and scratch assay results between experimental groups were analyzed using Kruskal-Wallis H with Mann-Whitney U *post hoc* analysis. Meanwhile, differences of transwell assay results and SHH band density between experimental groups were analyzed using one-way analysis of variances (ANOVA) with Tukey's honestly significant difference (HSD) *post hoc* analysis. All results were expressed as mean \pm standard deviation (SD). $P < 0.05$ were considered as statistically significant.

Results

EBBE reduced the number of viable HSC-3 cells

The number of viable HSC-3 cells cultured in medium containing extract solvent (sham group) for

24 h was $9,607 \pm 13.81$. Treatment with EBBE showed that the viability of HSC-3 cells was significantly lower compared to the sham group ($P < 0.05$) in a concentration-dependent manner. The number of viable HSC-3 cells treated with 1, 10, and 100 μ g/mL EBBE were $9,055 \pm 268.69$, $6,777 \pm 278.04$, and $3,659 \pm 32.86$, respectively. The viability of EBBE-treated HSC-3 cells was significantly higher ($P < 0.05$) than Doxorubicin-treated HSC-3 cells (226 ± 32.34) (Fig. 1). These findings suggested that EBBE could reduce the viability of HSC-3 cells.

EBBE induced apoptosis of HSC-3 cells

Sub-G1 assay results showed that the apoptosis percentage of the sham group was $8.27 \pm 0.14\%$. Apoptotic HSC-3 cells were significantly higher compared to the sham group ($P < 0.05$) in concentration-dependent manner after EBBE treatment. Apoptosis percentages of HSC-3 cells treated with 1, 10, and 100 μ g/mL EBBE were $9.60 \pm 0.35\%$, $17.78 \pm 0.32\%$, and $38.64 \pm 0.70\%$, respectively. These percentages were significantly lower ($P < 0.05$) than Doxorubicin-treated HSC-3 cells ($95.95 \pm 2.60\%$) (Fig. 2). Therefore, these results suggested that EBBE treatment could promote apoptosis of HSC-3 cells.

EBBE inhibited HSC-3 cell migration

The gap closure percentage of HSC-3 cells cultured in medium containing extract solvent after 24 h was $96.11 \pm 4.17\%$. After EBBE treatment, the migratory




Fig. 1 — EBBE decreased the number of viable HSC-3 cells in a concentration-dependent manner. HSC-3 cells were cultured and treated with/without 1 μ M Doxorubicin and various concentrations of EBBE for 24 h as indicated in the panel. The number of viable cells were measured with MTT assay as described in Materials and Methods. The data are expressed as mean \pm SD (n=6). * $P < 0.05$ compared to the sham group




Fig. 2 — EBBE increased the percentage of apoptotic HSC-3 cells in a concentration-dependent manner. HSC-3 cells were cultured and treated with/without 1 μ M Doxorubicin and various concentrations of EBBE for 24 h as indicated. Treated cells were then subjected to sub-G1 assay as described in Materials and Methods. (A) Histogram of flow cytometric results. (B) Percentage of apoptotic HSC-3 cells. The data are expressed as mean \pm SD (n=18). * P < 0.05 compared to the sham group

activities of HSC-3 cells were significantly lower compared to the sham group (P < 0.05) in a concentration-dependent manner. Gap closure percentage of HSC-3 cells treated with 1, 10, and 100 μ g/mL EBBE were $89.56 \pm 5.77\%$, $76.78 \pm 4.87\%$, and $60.67 \pm 7.62\%$, respectively. These percentages were significantly higher (P < 0.05) than cyclopamine tartrate-treated HSC-3 cells ($43.00 \pm 4.82\%$) (Fig. 3).

Transwell assay also showed similar results. The number of migrated HSC-3 cells in the sham group was 134.44 ± 11.60 . The number of migrated HSC-3 cells was significantly lower than the sham group (P < 0.05) in a concentration-dependent manner after EBBE treatment. The number of migrated HSC-3 cells treated with 1, 10, and 100 μ g/mL EBBE were 107.44 ± 15.21 , 95.11 ± 6.01 , and 39.22 ± 3.56 , respectively. These numbers were significantly higher (P < 0.05) than cyclopamine tartrate-treated HSC-3 cells (17.89 ± 3.22) (Fig. 4). Overall, these results

indicated that EBBE exhibited an anti-migratory effect towards HSC-3 cells.

EBBE downregulated the expression of SHH

Upon the addition of EBBE, the SHH expression levels in HSC-3 cells were lower in a concentration-dependent manner. Expression level of SHH in HSC-3 cells treated with 1 μ g/mL EBBE was significantly higher (P < 0.05), while SHH expression levels in HSC-3 cells treated with 10 and 100 μ g/mL EBBE were significantly lower (P < 0.05) compared to the sham group. SHH expression levels in HSC-3 cells treated with 1 and 10 μ g/mL EBBE were significantly higher compared to cyclopamine tartrate-treated HSC-3 cells (P < 0.05). However, no significant difference was observed in the SHH expression levels between cyclopamine tartrate and 100 μ g/mL EBBE treatment ($P = 0.142$) (Fig. 5). Thus, these findings suggested that EBBE could decrease the expression of SHH.




Fig. 3 — EBBE inhibited migratory activities of HSC-3 cells. HSC-3 cell culture was scratched and treated with/without EBBE or 20 µM cyclopamine tartrate for 24 h as indicated. Gap closure was measured as described in Materials and Methods. (A) Scratched areas of HSC-3 cell culture. White bar: 100 µM. (B) Gap closure percentage of EBBE-treated HSC-3 cell. The data are expressed as mean ± SD (n=9). **P* < 0.05 compared to the sham group




Fig. 4 — EBBE reduced the number of migrated HSC-3 cells. HSC-3 cell culture was treated with/without EBBE or 20 µM cyclopamine tartrate for 24 h as indicated. Migrated cells were measured as described in Materials and Methods. The data are expressed as mean ± SD (n=9). **P* < 0.05 compared to the sham group




Fig. 5 — EBBE reduced the expression of SHH. HSC-3 cell culture was treated with/without EBBE or 20 µM cyclopamine tartrate for 24 h as indicated. HSC-3 cells were lysed, homogenized, and immunoblotted using anti-SHH antibody as described in Materials and Methods. The data are expressed as mean ± SD (n=3). **P* < 0.05 compared to the sham group

Discussion

In the present study, we investigated the potency of EBBE as anticancer agent for OSCC treatment. EBBE showed a cytotoxic effect towards HSC-3 cells by reducing the viability of HSC-3 cells in a concentration-dependent manner. According to the sub-G1 assay results, the decrease in HSC-3 cell viability was caused by apoptosis. These results were in line with previous studies that EBBE was able to induce apoptosis of cancer cells^{18–20,25}. EBBE was also reported to inhibit cell cycle in several cell lines^{19,25}, indicating that EBBE may induce apoptosis of HSC-3 cells through cell cycle arrest since both processes are interconnected²⁶. Based on a molecular docking study, eleutherol, eleutherin, and isoeleutherin could induce apoptosis by activating caspase-8²⁷.

Cell migration is one of the key processes involved in cancer metastasis²⁸. Scratch and transwell assays demonstrated that EBBE restrained the migration of HSC-3 cells in a concentration-dependent manner. This showed the novel activity of EBBE in impairing HSC-3 cell migration. Therefore, EBBE may have anti-metastasis activity towards tongue cancer cells.

Metastasis of tongue cancer is regulated by several proteins, including SHH. One of the SHH signaling pathways that has been reported to regulate cell migration in tongue cancer is SHH/glioma-associated oncogene homologue (SHH/GLI)²⁹. SHH protein binds and inactivates its receptor, Patched 1 (PTCH1). This interaction causes the activation of transcription factors from the GLI family, which play a major role in regulating cancer migration, invasion, and proliferation³⁰.

The present study revealed that EBBE decreased SHH expression levels along with the increase in EBBE concentrations, suggesting that the inhibitory effect of EBBE towards the expression of SHH may be responsible for its anti-migratory property. To the best of our knowledge, this is the first study demonstrating SHH downregulation in EBBE-treated HSC-3 cells. EBBE may inhibit HSC-3 cell migration via inhibition of SHH/GLI signaling pathway. Possibly, all signaling pathways that are regulated by SHH can be inhibited by EBBE. EBBE may also restrain HSC-3 cell migration through another pathway. In SW480 colon cancer cell line, eleutherinoside C and isoeleutherin obtained from *E. bulbosa* bulbs was reported to inhibit Wnt/ β -catenin pathway¹⁶, which has been reported to play an important role in regulating cancer migration³¹.

Therefore, signaling pathways involved in the inhibition of HSC-3 cell migration by EBBE still need to be explored. In addition, signaling components downstream to SHH and target molecules that are affected by EBBE should be identified. Since EBBE also contains various bioactive compounds, further research is needed to identify which compounds are responsible for apoptosis and anti-migratory activity. Thus, more investigations are required to understand the molecular mechanisms underlying EBBE-induced HSC-3 cell apoptosis and confirm which SHH signaling pathway contributes to the inhibition of cell migration by EBBE.

Conclusion

EBBE reduces HSC-3 cell viability through apoptosis and inhibits its migratory activities by down regulating SHH expression. Taken together, EBBE could be a potential anti-tongue cancer agent, particularly as an ant metastatic agent.

Conflict of interest

All authors declare no conflicts of interest.

References

- Chen YK, Huang HC, Lin LM & Lin CC, Primary oral squamous cell carcinoma: An analysis of 703 cases in southern Taiwan. *Oral Oncol*, 35 (1999) 173.
- Neville BW & Day TA, Oral cancer and precancerous lesions. *CA Cancer J Clin*, 52 (2002) 195.
- Minhas S, Kashif M, Altaf W, Afzal N & Nagi AH, Concomitant-chemoradiotherapy-associated oral lesions in patients with oral squamous-cell carcinoma. *Cancer Biol Med*, 14 (2017) 176.
- Wicaksono BD, Tangkearung E & Sandra F, *Brucea javanica* leaf extract induced apoptosis in human oral squamous cell carcinoma (HSC2) cells by attenuation of mitochondrial membrane permeability. *Indones Biomed J*, 7 (2015) 107.
- Chang WL, Cheng FC, Wang SP, Chou ST & Shih Y, *Cinnamomum cassia* essential oil and its major constituent cinnamaldehyde induced cell cycle arrest and apoptosis in human oral squamous cell carcinoma HSC-3 cells. *Environ Toxicol*, 32 (2017) 456.
- Susianti S, Lesmana R, Salam S, Julaeha E, Pratiwi YS, Sylviana N, Goenawan H, Kurniawan A & Supratman U, The effect of nutmeg seed (*M. fragrans*) extracts induces apoptosis in melanoma maligna cell's (B16-F10). *Indones Biomed J*, 13 (2021) 68.
- Kameyanda Poonacha S, Harishkumar M, Radha M, Varadarajan R, Nalilu SK, Shetty SS, Shetty PK, Chandrashekarappa RB, Sreenivas MG & Bhandary Bavabeedu SK, Insight into oroxylinA-7-O- β -d-glucuronide-enriched *Oroxylum indicum* bark extract in oral cancer HSC-3 cell apoptotic mechanism: Role of mitochondrial microenvironment. *Molecules*, 26 (2021) 7430.
- Ren W, Qiao Z, Wang H, Zhu L & Zhang L, Flavonoids:

- Promising anticancer agents. *Med Res Rev*, 23 (2003) 519.
- 9 Sandra F & Sidharta MA, Caffeic acid induced apoptosis in MG63 osteosarcoma cells through activation of caspases. *Mol Cell Biomed Sci*, 1 (2017) 28.
 - 10 Ieyama T, Gunawan-Puteri MDPT & Kawabata J, α -Glucosidase inhibitors from the bulb of *Eleutherine americana*. *Food Chem*, 128 (2011) 308.
 - 11 Subramaniam K, Suriyamoorthy S, Wahab F, Sharon FB & Rex GR, Antagonistic activity of *Eleutherine palmifolia* Linn. *Asian Pac J Trop Dis*, 2 (2012) S491.
 - 12 Kuntorini EM, Dewi M & Misrina M, Anatomical structure and antioxidant activity of red bulb plant (*Eleutherine americana*) on different plant age. *Biodiversitas*, 17 (2016) 229.
 - 13 Insanu M, Kusumardiyani S & Hartati R, Recent studies on phytochemicals and pharmacological effects of *Eleutherine americana* Merr. *Procedia Chem*, 13 (2014) 221.
 - 14 Chen Z, Huang H, Wang C, Li Y, Ding J, Sankawa U, Noguchi H & Iitaka Y, Hongconin, a new naphthalene derivative from hong-cong, the rhizome of *Eleutherine americana* Merr. et Heyne (Iridaceae). *Chem Pharm Bull*, 34 (1986) 2743.
 - 15 Xu J, Qiu F, Duan W, Qu G, Wang N & Yao X, New bioactive constituents from *Eleutherine americana*. *Front Chem China*, 1 (2006) 320.
 - 16 Li X, Ohtsuki T, Koyano T, Kowithayakorn T & Ishibashi M, New Wnt/ β -catenin signaling inhibitors isolated from *Eleutherine palmifolia*. *Chem Asian J*, 4 (2009) 540.
 - 17 Mutiah R, Minggarwati TS, Kristanti RA & Susanti E, Compound identification and anticancer activity of ethyl acetate fraction from bawang sabrang (*Eleutherine palmifolia* (L.) Merr.) on HeLa cervical cancer cell line. *Indones J Cancer Chemoprevent*, 10 (2019) 131.
 - 18 Lubis IA, Ichwan M, Mustofa M & Satria D, Anticancer activity of *Eleutherine bulbosa* (Mill.) Urb. extract on WiDr cell line in vitro, paper presented to 2nd Public Health International Conference (PHICo 2017), Faculty of Public Health – Universitas Sumatera Utara, Medan, Indonesia, 18-19 December 2017.
 - 19 Yuniarti A, Sundhani E & Nurulita NA, The potentiation effect of bawang dayak (*Sisyrinchium palmifolium* L.) extract on T47D cell growth inhibition after 5-fluorouracil treatment. *Pharmaciana*, 8 (2018) 195.
 - 20 Kamarudin AA, Sayuti NH, Saad N, Razak NAA & Esa NM, Induction of apoptosis by *Eleutherine bulbosa* (Mill.) Urb. bulb extracted under optimised extraction condition on human retinoblastoma cancer cells (WERI-Rb-1). *J Ethnopharmacol*, 284 (2022) 114770.
 - 21 Sandra F, Sudiono J, Trisfilha P & Pratiwi D, Cytotoxicity of *Alpinia galanga* rhizome crude extract on NIH-3T3 cells. *Indones Biomed J*, 9 (2017) 23.
 - 22 Ariesanti Y, Sandra F, Claresta B & Alvita L, *Coffea canephora* bean extract induces NIH3T3 cell migration. *Indones Biomed J*, 13 (2021) 216.
 - 23 Oktaviono YH, Al-Farabi MJ, Suastika LOS, Hartono F, Dirgantara Y & Sandra F, Preliminary study: Purple sweet potato extract seems to be superior to increase the migration of impaired endothelial progenitor cells compared to L-Ascorbic acid. *Sci Pharm*, 87(2019) 16.
 - 24 Rizal MI & Sandra F, *Brucea javanica* leaf extract activates caspase-9 and caspase-3 of mitochondrial apoptotic pathway in human oral squamous cell carcinoma. *Indones Biomed J*, 8 (2016) 43.
 - 25 Mutiah R, Listiyana A, Suryadinata A, Annisa R, Hakim A, Anggraini W & Susilowati R, Activity of inhibit the cell cycle and induct apoptosis in HeLa cancer cell with combination of sabrang onion (*Eleutherine palmifolia* (L.) Merr) and starfruit mistletoe (*Macrosolen cochinchinensis* (Lour.) Tiegh). *J App Pharm Sci*, 8 (2018) 122.
 - 26 Pucci B, Kasten M & Giordano A, Cell cycle and apoptosis. *Neoplasia*, 2 (2000) 291.
 - 27 Quadros Gomes AR, da Rocha Galucio NC, de Albuquerque KCO, Brígido HPC, Varela ELP, Castro ALG, Vale VV, Bahia MO, Rodriguez Burbano RM, de Molfeta FA, Carneiro LA, Percario S & Dolabela MF, Toxicity evaluation of *Eleutherine plicata* Herb. extracts and possible cell death mechanism. *Toxicol Rep*, 8 (2021) 1480.
 - 28 Ellis IR, The migration and invasion of oral squamous carcinoma cells: Matrix, growth factor and signalling involvement. *Cancers*, 13 (2021) 2633.
 - 29 Fan HX, Wang S, Zhao H, Liu N, Chen D, Sun M & Zheng JH, Sonic hedgehog signaling may promote invasion and metastasis of oral squamous cell carcinoma by activating MMP-9 and E-cadherin expression. *Med Oncol*, 31 (2014) 41.
 - 30 Carballo GB, Honorato JR, de Lopes GPF & Spohr TCLSE, A highlight on Sonic hedgehog pathway. *Cell Commun Signal*, 16 (2018) 11.
 - 31 Liang S, Zhang S, Wang P, Yang C, Shang C, Yang J & Wang J, LncRNA, TUG1 regulates the oral squamous cell carcinoma progression possibly via interacting with Wnt/ β -catenin signaling. *Gene*, 608 (2017) 49.



Ferry Sandra <ferry@trisakti.ac.id>

[IJBB] Editor Decision

Team IJBB <ijbb@niscpr.res.in>

Mon, Sep 25, 2023 at 12:36 PM

To: Ferry Sandra <ferry@trisakti.ac.id>, Kelvin Suryajaya <kelvin.thong222@gmail.com>, Angliana Chow <angliana@prostem.co.id>, Maria Celinna <maria.celinna@prodia.institute>, Dewi Ranggaini <monica.dewi.r@trisakti.ac.id>, Cynthia Retna Sartika <cynthia@prostem.co.id>

Ferry Sandra, Kelvin Suryajaya, Angliana Chow, Maria Celinna, Dewi Ranggaini, Cynthia Retna Sartika:

The editing of your submission, "Eleutherine bulbosa Bulb Extract Induces Apoptosis and Inhibits Cell Migration by Downregulating Sonic Hedgehog Gene in Human Tongue Cancer Cells: An in vitro Study(Article Id: IJBB#83)," is complete. We are now sending it to production.

Submission URL: <https://or.niscpr.res.in/index.php/IJBB/authorDashboard/submission/1293>

[Indian Journal of Biochemistry and Biophysics \(IJBB\)](#)



Ferry Sandra <ferry@trisakti.ac.id>

Ultra Urgent PRC IJBB#83

IJBB NIScPR <ijbbniscpr@gmail.com>
To: ferry@trisakti.ac.id

Mon, Sep 25, 2023 at 12:48 PM


Dear Author,

Enclosed please find the final proof-ready copy of the above MS. In accordance with the editorial corrections/suggestions may please be submitted positively by **today** itself for finalization. Kindly use the enclosed PDF file only for marking corrections and submission. Highlight the corrections made in the MS with yellow color. After publishing the article we do not entertain any kind of corrections.

Thanks & Regards

Team IJBB

[Quoted text hidden]

 **IJBB#83 auth corr.pdf**
1319K

Eleutherine bulbosa bulb extract induces apoptosis and inhibits cell migration by down-regulating sonic hedgehog in human tongue cancer cells: An *in vitro* study

Ferry Sandra^{1*}, Kelvin Suryajaya², Angliana Chouw³, Maria Celinna⁴, Dewi Ranggaini⁵, Kyung Hoon Lee⁶ & Cynthia Retna Sartika³

¹Department of Biochemistry and Molecular Biology, Division of Oral Biology, Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia

²Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia

³PT Prodia StemCell Indonesia, Jakarta Pusat-10430, Jakarta, Indonesia

⁴The Prodia Education and Research Institute, Jakarta Pusat-10430, Jakarta, Indonesia

⁵Department of Physiology, Division of Oral Biology, Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia

⁶Research Institute, Ballys Co. Ltd, Incheon-22219, Republic of Korea

Received 09 May 2023; revised 18 September 2023

Eleutherine bulbosa (Mill.) Urb. is a medicinal plant which has long been used to treat cancer. *E. bulbosa* bulb extract (EBBE) has been reported to show cytotoxicity towards several types of human cancer. However, the cytotoxic effect of EBBE towards oral cancer cells has not been investigated. The present study aimed to evaluate the effects of EBBE towards the viability, apoptosis, and migratory activities of oral cancer cells. Human oral squamous cell carcinoma (HSC)-3 cells were treated with various concentrations of EBBE for 24 h. The number of viable and apoptotic HSC-3 cells were measured using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) and sub-G1 assay, respectively. The migratory activities of HSC-3 cells were assessed using scratch and transwell assay. Sonic hedgehog (SHH) expression was measured using immunoblotting. Upon EBBE treatment, apoptotic HSC-3 cells were significantly higher in a concentration-dependent manner ($P < 0.05$). Meanwhile, viability, migratory activities, and migrated HSC-3 cell number were significantly lower in a concentration-dependent manner ($P < 0.05$). The SHH expression levels in EBBE-treated HSC-3 cells were also lower in a concentration-dependent manner. EBBE reduces HSC-3 cell viability through apoptosis and inhibits its migratory activities by downregulating SHH expression.

Keywords: Apoptosis, Cell migration, *Eleutherine bulbosa*, Sonic hedgehog, Tongue cancer

Oral squamous cell carcinoma (OSCC) is the most common histological type of oral cancer¹ which arises from squamous epithelial cells in the oral cavity. This cancer generally affects the floor of the mouth and the tongue². Various modes of treatment are developed to manage OSCC, including chemotherapy, radiotherapy, surgery, or its combination³. However, these treatments cause numerous side effects.

Discovery and development of new therapeutic agents derived from natural sources, particularly plants with less side effects are recently one of the main focuses on oncology research. Several plants have been reported to have anticancer effects, including

*Brucea javanica*⁴, *Cinnamomum cassia*⁵, *Myristica fragrans*⁶, and *Oroxylum indicum*⁷. Plants have been known as rich sources of various active compounds, including flavonoids, which was reported to inhibit cancer proliferation *via* cell cycle arrest, antiproliferative and antioxidative activity, apoptosis induction, promotion of differentiation, inhibition of angiogenesis and carcinogen metabolic activation, and multidrug resistance modulation⁸. In addition, several non-flavonoid phenolic compounds have also been reported to show anticancer properties, such as caffeic acid⁹.

Eleutherine bulbosa (Mill.) Urb. is a traditional medicinal plant belonging to the Iridaceae family. Its bulbs are commonly used for the treatment of breast cancer, hypertension, stroke, and diabetes mellitus. It has also been used to increase the production of breast milk and treat sexual disorder¹⁰. *E. bulbosa* bulbs have

*Correspondence:
Phone: +628128077780
E-mail: ferry@trisakti.ac.id

also been reported to have antimicrobial¹¹ and antioxidant¹² activity.

Eleutherine bulbosa contains a wide variety of compounds, primarily naphthoquinones, anthraquinones, and naphthalene¹³. The main compounds found in this plant are eleutherin, isoeleutherin, and eleutherol¹⁴. Several compounds obtained from *E. bulbosa* bulbs have been reported to show anticancer activities, such as 6,8-hydroxy-3,4-di-methoxy-1-methyl-anthraquinone-2-carboxylic acid methyl ester¹⁵, eleutherinoside C, and isoeleutherin¹⁶.

Eleutherine bulbosa bulb extract (EBBE) has been reported to show cytotoxicity towards several types of human cancer, such as cervical cancer (line HeLa)¹⁷, colon cancer (line SW480, HCT116, DLD1, WiDr)^{16,18}, breast cancer (line T47D)¹⁹, leukemia (line K562)¹⁵, and retinoblastoma (WERI-Rb-1)²⁰. However, the cytotoxic effect of EBBE towards human oral cancer cells has not been investigated. Therefore, the present study was conducted to evaluate the effects of EBBE towards the viability, apoptosis, and migratory activities of oral cancer cells.

Materials and Methods

E. bulbosa bulb extract (EBBE) preparation

E. bulbosa bulbs were obtained from Indonesian Medicinal and Aromatic Crops Research Institute (IMACRI), Indonesia. Plant identification was carried out by expert botanist at Research Center for Plant Conservation and Botanic Gardens, Indonesian Institute of Sciences, Indonesia (No. B-1269/IPH.3/KS/X/2020). EBBE was extracted by maceration method⁴. Briefly, bulbs of *E. bulbosa* were minced and dried. The dried material was extracted with ethanol 70%, filtered and evaporated using a rotary evaporator. The resulting crude EBBE was then stored at 4°C.

HSC-3 cell culture

HSC-3 cell culture was performed as previously described⁴ with modification. The HSC-3 cell line was purchased from Sigma-Aldrich Pte. Ltd. (St. Louis, MO, USA). HSC-3 cells were cultured in a complete medium containing Dulbecco's modified Eagle's medium (DMEM) (Sigma-Aldrich), 10% fetal bovine serum (FBS) (PAN-Biotech, Aidenbach, Germany), 50 U/mL penicillin and 50 µg/mL streptomycin (Sigma-Aldrich) in a humidified 5% CO₂, 37°C incubator. HSC-3 cells were then dissociated with trypsin-ethylenediamine tetraacetic acid (EDTA) solution (Sigma-Aldrich) after reaching

80% confluency. Upon reaching the desired cell number, HSC-3 cells were seeded into 24-well plate and 96-well plate for further assays.

Cell viability assay

The number of viable cells were measured quantitatively using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay as previously described²¹ with modification. Briefly, 5×10^3 HSC-3 cells were seeded into 96-well plates and then treated with EBBE (1, 10, or 100 µg/mL), 1 µM Doxorubicin (Dankos Farma, Jakarta, Indonesia) or medium merely for 24 h. After that 100 µL of MTT (Sigma-Aldrich) in culture medium was loaded into each well. After incubating the plates for 4 h, culture medium was discarded, and the formed formazan crystals were dissolved in 100 µL dimethylsulfoxide (DMSO). Results were measured at OD₅₇₀ using a microplate reader (Bio-Rad, Hercules, CA, USA). Each experimental group was measured in six replicates. In addition, the number of untreated cells were counted with hemacytometer and used for interpolating OD₅₇₀ values of HSC-3 cells.

Sub-G1 assay

To investigate the cytotoxic effect of EBBE on HSC-3 cells, the number of apoptotic cells were determined using sub-G1 assay as previously described⁹ with modification. Treated and untreated HSC-3 cells were harvested and suspended in 450 µL hypotonic fluorochrome solution containing 50 µg/mL propidium iodide (Sigma-Aldrich), 0.1% Triton X-100 (Sigma-Aldrich), and 0.1% sodium citrate (Wako, Osaka, Japan). After that, cell suspensions were incubated in the dark for 2 h at room temperature. Fluorescence of individual nuclei was measured using FACSCanto II flow cytometer (Becton Dickinson, Franklin Lakes, NJ, USA) at 100,000 events.

Scratch assay

Scratch assay was performed as previously described²². HSC-3 cells (5×10^4) were seeded into 24-well plates and incubated overnight. Scratch was made on each well with micropipette yellow tips. Then, the wells were washed with phosphate-buffered saline (PBS) (Wako) and treated with/without EBBE or 20 µM cyclopamine tartrate (BioVision, Milpitas, CA, USA) (as positive control) for 24 h. The gap closures were observed and documented under an inverted light microscope (Carl Zeiss, Jena, Germany).

Transwell assay

Transwell assay was performed as previously described²³ with modification. HSC-3 cells (5×10^4) were seeded in the 8- μ m pore size chamber insert (Merck, Darmstadt, Germany). After that, 800 μ L culture medium with/without EBBE or 20 μ M cyclopamine tartrate was loaded into each well of the 24-well plate. HSC-3-containing chamber inserts were then placed on the 24-well plate. After incubating the chamber for 24 h at 37°C, migrated cells in the lower surface of the chamber were fixed. The cells were counted and documented under an inverted light microscope.

Immunoblotting

Expression levels of Sonic hedgehog (SHH) protein were measured using immunoblotting method as previously described²⁴ with modification. HSC-3 cells were lysed and homogenized with Laemmli sample buffer (Bio-Rad). The proteins from each experimental group were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and transferred to a polyvinylidene difluoride (PVDF) sheet. After blocking with 5% skim milk solution, the sheet was incubated with rabbit monoclonal anti-SHH antibody as a primary antibody and horseradish peroxidase (HRP)-conjugated goat anti-rabbit antibody as a secondary antibody (Cell Signaling Technology, Danvers, MA, USA). Immunoblot results were analyzed using ImageJ software (National Institute of Health, Bethesda, MD, USA) to measure the density of each SHH band.

Statistical analysis

Statistical analyses were performed using IBM SPSS Statistics version 26.0 (SPSS IBM, Armonk, NY, USA). Shapiro-Wilk test was used as a normality test. Differences of MTT assay, sub-G1 assay, and scratch assay results between experimental groups were analyzed using Kruskal-Wallis H with Mann-Whitney U *post hoc* analysis. Meanwhile, differences of transwell assay results and SHH band density between experimental groups were analyzed using one-way analysis of variances (ANOVA) with Tukey's honestly significant difference (HSD) *post hoc* analysis. All results were expressed as mean \pm standard deviation (SD). $P < 0.05$ were considered as statistically significant.

Results

EBBE reduced the number of viable HSC-3 cells

The number of viable HSC-3 cells cultured in medium containing extract solvent (sham group) for

24 h was $9,607 \pm 13.81$. Treatment with EBBE showed that the viability of HSC-3 cells was significantly lower compared to the sham group ($P < 0.05$) in a concentration-dependent manner. The number of viable HSC-3 cells treated with 1, 10, and 100 μ g/mL EBBE were $9,055 \pm 268.69$, $6,777 \pm 278.04$, and $3,659 \pm 32.86$, respectively. The viability of EBBE-treated HSC-3 cells was significantly higher ($P < 0.05$) than Doxorubicin-treated HSC-3 cells (226 ± 32.34) (Fig. 1). These findings suggested that EBBE could reduce the viability of HSC-3 cells.

EBBE induced apoptosis of HSC-3 cells

Sub-G1 assay results showed that the apoptosis percentage of the sham group was $8.27 \pm 0.14\%$. Apoptotic HSC-3 cells were significantly higher compared to the sham group ($P < 0.05$) in concentration-dependent manner after EBBE treatment. Apoptosis percentages of HSC-3 cells treated with 1, 10, and 100 μ g/mL EBBE were $9.60 \pm 0.35\%$, $17.78 \pm 0.32\%$, and $38.64 \pm 0.70\%$, respectively. These percentages were significantly lower ($P < 0.05$) than Doxorubicin-treated HSC-3 cells ($95.95 \pm 2.60\%$) (Fig. 2). Therefore, these results suggested that EBBE treatment could promote apoptosis of HSC-3 cells.

EBBE inhibited HSC-3 cell migration

The gap closure percentage of HSC-3 cells cultured in medium containing extract solvent after 24 h was




Fig. 1 — EBBE decreased the number of viable HSC-3 cells in a concentration-dependent manner. HSC-3 cells were cultured and treated with/without 1 μ M Doxorubicin and various concentrations of EBBE for 24 h as indicated in the panel. The number of viable cells were measured with MTT assay as described in Materials and Methods. The data are expressed as mean \pm SD ($n=6$). * $P < 0.05$ compared to the sham group




Fig. 2 — EBBE increased the percentage of apoptotic HSC-3 cells in a concentration-dependent manner. HSC-3 cells were cultured and treated with/without 1 μM Doxorubicin and various concentrations of EBBE for 24 h as indicated. Treated cells were then subjected to sub-G1 assay as described in Materials and Methods. (A) Histogram of flow cytometric results. (B) Percentage of apoptotic HSC-3 cells. The data are expressed as mean \pm SD (n=18). * $P < 0.05$ compared to the sham group

96.11 \pm 4.17%. After EBBE treatment, the migratory activities of HSC-3 cells were significantly lower compared to the sham group ($P < 0.05$) in a concentration-dependent manner. Gap closure percentage of HSC-3 cells treated with 1, 10, and 100 $\mu\text{g/mL}$ EBBE were 89.56 \pm 5.77%, 76.78 \pm 4.87%, and 60.67 \pm 7.62%, respectively. These percentages were significantly higher ($P < 0.05$) than cyclopamine tartrate-treated HSC-3 cells (43.00 \pm 4.82%) (Fig. 3).

Transwell assay also showed similar results. The number of migrated HSC-3 cells in the sham group was 134.44 \pm 11.60. The number of migrated HSC-3 cells was significantly lower than the sham group ($P < 0.05$) in a concentration-dependent manner after EBBE treatment. The number of migrated HSC-3 cells treated with 1, 10, and 100 $\mu\text{g/mL}$ EBBE were 107.44 \pm 15.21, 95.11 \pm 6.01, and 39.22 \pm 3.56, respectively. These numbers were significantly higher ($P < 0.05$) than cyclopamine tartrate-treated HSC-3 cells (17.89 \pm 3.22) (Fig. 4). Overall, these results

indicated that EBBE exhibited an anti-migratory effect towards HSC-3 cells.

EBBE downregulated the expression of SHH

Upon the addition of EBBE, the SHH expression levels in HSC-3 cells were lower in a concentration-dependent manner. Expression level of SHH in HSC-3 cells treated with 1 $\mu\text{g/mL}$ EBBE was significantly higher ($P < 0.05$), while SHH expression levels in HSC-3 cells treated with 10 and 100 $\mu\text{g/mL}$ EBBE were significantly lower ($P < 0.05$) compared to the sham group. SHH expression levels in HSC-3 cells treated with 1 and 10 $\mu\text{g/mL}$ EBBE were significantly higher compared to cyclopamine tartrate-treated HSC-3 cells ($P < 0.05$). However, no significant difference was observed in the SHH expression levels between cyclopamine tartrate and 100 $\mu\text{g/mL}$ EBBE treatment ($P = 0.142$) (Fig. 5). Thus, these findings suggested that EBBE could decrease the expression of SHH.




Fig. 3 — EBBE inhibited migratory activities of HSC-3 cells. HSC-3 cell culture was scratched and treated with/without EBBE or 20 µM cyclopamine tartrate for 24 h as indicated. Gap closure was measured as described in Materials and Methods. (A) Scratched areas of HSC-3 cell culture. White bar: 100 µM. (B) Gap closure percentage of EBBE-treated HSC-3 cell. The data are expressed as mean ± SD (n=9). **P* < 0.05 compared to the sham group




Fig. 4 — EBBE reduced the number of migrated HSC-3 cells. HSC-3 cell culture was treated with/without EBBE or 20 µM cyclopamine tartrate for 24 h as indicated. Migrated cells were measured as described in Materials and Methods. The data are expressed as mean ± SD (n=9). **P* < 0.05 compared to the sham group




Fig. 5 — EBBE reduced the expression of SHH. HSC-3 cell culture was treated with/without EBBE or 20 µM cyclopamine tartrate for 24 h as indicated. HSC-3 cells were lysed, homogenized, and immunoblotted using anti-SHH antibody as described in Materials and Methods. The data are expressed as mean ± SD (n=3). **P* < 0.05 compared to the sham group

Discussion

In the present study, we investigated the potency of EBBE as anticancer agent for OSCC treatment. EBBE showed a cytotoxic effect towards HSC-3 cells by reducing the viability of HSC-3 cells in a concentration-dependent manner. According to the sub-G1 assay results, the decrease in HSC-3 cell viability was caused by apoptosis. These results were in line with previous studies that EBBE was able to induce apoptosis of cancer cells^{18-20,25}. EBBE was also reported to inhibit cell cycle in several cell lines^{19,25}, indicating that EBBE may induce apoptosis of HSC-3 cells through cell cycle arrest since both processes are interconnected²⁶. Based on a molecular docking study, eleutherol, eleutherin, and isoeleutherin could induce apoptosis by activating caspase-8²⁷.

Cell migration is one of the key processes involved in cancer metastasis²⁸. Scratch and transwell assays demonstrated that EBBE restrained the migration of HSC-3 cells in a concentration-dependent manner. This showed the novel activity of EBBE in impairing HSC-3 cell migration. Therefore, EBBE may have anti-metastasis activity towards tongue cancer cells.

Metastasis of tongue cancer is regulated by several proteins, including SHH. One of the SHH signaling pathways that has been reported to regulate cell migration in tongue cancer is SHH/glioma-associated oncogene homologue (SHH/GLI)²⁹. SHH protein binds and inactivates its receptor, Patched 1 (PTCH1). This interaction causes the activation of transcription factors from the GLI family, which play a major role in regulating cancer migration, invasion, and proliferation³⁰.

The present study revealed that EBBE decreased SHH expression levels along with the increase in EBBE concentrations, suggesting that the inhibitory effect of EBBE towards the expression of SHH may be responsible for its anti-migratory property. To the best of our knowledge, this is the first study demonstrating SHH downregulation in EBBE-treated HSC-3 cells. EBBE may inhibit HSC-3 cell migration via inhibition of SHH/GLI signaling pathway. Possibly, all signaling pathways that are regulated by SHH can be inhibited by EBBE. EBBE may also restrain HSC-3 cell migration through another pathway. In SW480 colon cancer cell line, eleutherinoside C and isoeleutherin obtained from *E. bulbosa* bulbs was reported to inhibit Wnt/ β -catenin pathway¹⁶, which has been reported to play an important role in regulating cancer migration³¹.

Therefore, signaling pathways involved in the inhibition of HSC-3 cell migration by EBBE still need to be explored. In addition, signaling components downstream to SHH and target molecules that are affected by EBBE should be identified. Since EBBE also contains various bioactive compounds, further research is needed to identify which compounds are responsible for apoptosis and anti-migratory activity. Thus, more investigations are required to understand the molecular mechanisms underlying EBBE-induced HSC-3 cell apoptosis and confirm which SHH signaling pathway contributes to the inhibition of cell migration by EBBE.

Conclusion

EBBE reduces HSC-3 cell viability through apoptosis and inhibits its migratory activities by down regulating SHH expression. Taken together, EBBE could be a potential anti-tongue cancer agent, particularly as an ant metastatic agent.

Conflict of interest

All authors declare no conflicts of interest.

References

- 1 Chen YK, Huang HC, Lin LM & Lin CC, Primary oral squamous cell carcinoma: An analysis of 703 cases in southern Taiwan. *Oral Oncol*, 35 (1999) 173.
- 2 Neville BW & Day TA, Oral cancer and precancerous lesions. *CA Cancer J Clin*, 52 (2002) 195.
- 3 Minhas S, Kashif M, Altaf W, Afzal N & Nagi AH, Concomitant-chemoradiotherapy-associated oral lesions in patients with oral squamous-cell carcinoma. *Cancer Biol Med*, 14 (2017) 176.
- 4 Wicaksono BD, Tangkearung E & Sandra F, *Brucea javanica* leaf extract induced apoptosis in human oral squamous cell carcinoma (HSC2) cells by attenuation of mitochondrial membrane permeability. *Indones Biomed J*, 7 (2015) 107.
- 5 Chang WL, Cheng FC, Wang SP, Chou ST & Shih Y, *Cinnamomum cassia* essential oil and its major constituent cinnamaldehyde induced cell cycle arrest and apoptosis in human oral squamous cell carcinoma HSC-3 cells. *Environ Toxicol*, 32 (2017) 456.
- 6 Susianti S, Lesmana R, Salam S, Julaha E, Pratiwi YS, Sylviana N, Goenawan H, Kumiawan A & Supratman U, The effect of nutmeg seed (*M. fragrans*) extracts induces apoptosis in melanoma maligna cell's (B16-F10). *Indones Biomed J*, 13 (2021) 68.
- 7 KameyandaPoonacha S, Harishkumar M, Radha M, Varadarajan R, Nalilu SK, Shetty SS, Shetty PK, Chandrashekharappa RB, Sreenivas MG & Bavabeedu SKB, Insight into oroxylinA-7-O- β -d-glucuronide-enriched *Oroxylum indicum* bark extract in oral cancer HSC-3 cell apoptotic mechanism: Role of mitochondrial microenvironment. *Molecules*, 26 (2021) 7430.
- 8 Ren W, Qiao Z, Wang H, Zhu L & Zhang L, Flavonoids: Promising anticancer agents. *Med Res Rev*, 23 (2003) 519.

- 9 Sandra F & Sidharta MA, Caffeic acid induced apoptosis in MG63 osteosarcoma cells through activation of caspases. *Mol Cell Biomed Sci*, 1(2017)28.
- 10 Ieyama T, Gunawan-Puteri MDPT & Kawabata J, α -Glucosidase inhibitors from the bulb of *Eleutherine americana*. *Food Chem*, 128 (2011) 308.
- 11 Subramaniam K, Suriyamoorthy S, Wahab F, Sharon FB & Rex GR, Antagonistic activity of *Eleutherine palmifolia* Linn. *Asian Pac J Trop Dis*, 2 (2012) S491.
- 12 Kuntorini EM, Dewi M & Misrina M, Anatomical structure and antioxidant activity of red bulb plant (*Eleutherine americana*) on different plant age. *Biodiversitas*, 17 (2016) 229.
- 13 Insanu M, Kusumardiyani S & Hartati R, Recent studies on phytochemicals and pharmacological effects of *Eleutherine americana* Merr. *Procedia Chem*, 13 (2014) 221.
- 14 Chen Z, Huang H, Wang C, Li Y, Ding J, Sankawa U, Noguchi H & Itaka Y, Hongconin, a new naphthalene derivative from hong-cong, the rhizome of *Eleutherine Americana* Merr. et Heyne (Iridaceae). *Chem Pharm Bull*, 34 (1986) 2743.
- 15 Xu J, Qiu F, Duan W, Qu G, Wang N & Yao X, New bioactive constituents from *Eleutherine americana*. *Front Chem China*, 1 (2006) 320.
- 16 Li X, Ohtsuki T, Koyano T, Kowithayakorn T & Ishibashi M, New Wnt/ β -catenin signaling inhibitors isolated from *Eleutherine palmifolia*. *Chem Asian J*, 4(2009) 540.
- 17 Mutiah R, Minggarwati TS, Kristanti RA & Susanti E, Compound identification and anticancer activity of ethyl acetate fraction from bawang sabrang (*Eleutherine palmifolia* (L.) Merr.) on HeLa cervical cancer cell line. *Indones J Cancer Chemoprevent*, 10 (2019) 131.
- 18 Lubis IA, Ichwan M, Mustofa M & Satria D, Anticancer activity of *Eleutherine bulbosa* (Mill.) Urb. extract on WiDr cell line *in vitro*, paper presented to 2nd Public Health International Conference (PHICo 2017), Faculty of Public Health – Universitas Sumatera Utara, Medan, Indonesia, 18-19 December 2017.
- 19 Yuniarti A, Sundhani E & Nurulita NA, The potentiation effect of bawangdayak (*Sisyrinchium palmifolium* L.) extract on T47D cell growth inhibition after 5-fluorouracil treatment. *Pharmaciana*, 8 (2018) 195.
- 20 Kamarudin AA, Sayuti NH, Saad N, Razak NAA & Esa NM, Induction of apoptosis by *Eleutherine bulbosa* (Mill.) Urb. bulb extracted under optimised extraction condition on human retinoblastoma cancer cells (WERI-Rb-1). *J Ethnopharmacol*, 284 (2022) 114770.
- 21 Sandra F, Sudiono J, Trisfilha P & Pratiwi D, Cytotoxicity of *Alpinia galanga* rhizome crude extract on NIH-3T3 cells. *Indones Biomed J*, 9(2017) 23.
- 22 Ariesanti Y, Sandra F, Claresta B & Alvita L, *Coffea canephora* bean extract induces NIH3T3 cell migration. *Indones Biomed J*, 13(2021) 216.
- 23 Oktaviono YH, Al-Farabi MJ, Suastika LOS, Hartono F, Dirgantara Y & Sandra F, Preliminary study: Purple sweet potato extract seems to be superior to increase the migration of impaired endothelial progenitor cells compared to L-Ascorbic acid. *Sci Pharm*, 87 (2019) 16.
- 24 Rizal MI & Sandra F, *Brucea javanica* leaf extract activates caspase-9 and caspase-3 of mitochondrial apoptotic pathway in human oral squamous cell carcinoma. *Indones Biomed J*, 8 (2016) 43.
- 25 Mutiah R, Listiyana A, Suryadinata A, Annisa R, Hakim A, Anggraini W & Susilowati R, Activity of inhibit the cell cycle and induct apoptosis in HeLa cancer cell with combination of sabrang onion (*Eleutherine palmifolia* (L.) Merr) and starfruit mistletoe (*Macrosolen cochinchinensis* (Lour.) Tiegh). *J App Pharm Sci*, 8 (2018) 122.
- 26 Pucci B, Kasten M & Giordano A, Cell cycle and apoptosis. *Neoplasia*, 2(2000) 291.
- 27 Quadros Gomes AR, da Rocha Galucio NC, de Albuquerque KCO, Brigido HPC, Varela ELP, Castro ALG, Vale VV, Bahia MO, Rodriguez Burbano RM, de Molfeta FA, Carneiro LA, Percario S & Dolabela MF, Toxicity evaluation of *Eleutherine plicata* Herb. extracts and possible cell death mechanism. *Toxicol Rep*, 8 (2021) 1480.
- 28 Ellis IR, The migration and invasion of oral squamous carcinoma cells: Matrix, growth factor and signalling involvement. *Cancers*, 13 (2021) 2633.
- 29 Fan HX, Wang S, Zhao H, Liu N, Chen D, Sun M & Zheng JH, Sonic hedgehog signaling may promote invasion and metastasis of oral squamous cell carcinoma by activating MMP-9 and E-cadherin expression. *Med Oncol*, 31 (2014) 41.
- 30 Carballo GB, Honorato JR, de Lopes GPF & Spohr TCLSE, A highlight on Sonic hedgehog pathway. *Cell Commun Signal*, 16 (2018) 11.
- 31 Liang S, Zhang S, Wang P, Yang C, Shang C, Yang J & Wang J, LncRNA, TUG1 regulates the oral squamous cell carcinoma progression possibly *via* interacting with Wnt/ β -catenin signaling. *Gene*, 608 (2017) 49.



Ferry Sandra <ferry@trisakti.ac.id>

Ultra Urgent PRC IJBB#83

Ferry Sandra <ferry@trisakti.ac.id>
To: ijbbniscpr@gmail.com

Wed, Sep 27, 2023 at 9:44 AM

Dear Team IJBB,

Herewith I send the corrected PRC PDF and declaration letter with signature of all authors. The revised parts are highlighted in yellow. However, the corrections are written in the comments. Hopefully, you can find the PRC and declaration letter well.

Thank you.

Regards,
Ferry Sandra
[Quoted text hidden]
[Quoted text hidden]

2 attachments**IJBB#83 auth corr - commented.pdf**

1373K

**Certificate of Originality.pdf**

244K

Eleutherine bulbosa bulb extract induces apoptosis and inhibits cell migration by down-regulating sonic hedgehog in human tongue cancer cells: An *in vitro* study

Ferry Sandra^{1*}, Kelvin Suryajaya², Angliana Chow³, Maria Celinna⁴, Dewi Ranggaini⁵, Kyung Hoon Lee⁶ & Cynthia Retna Sartika³

¹Department of Biochemistry and Molecular Biology, Division of Oral Biology, Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia

²Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia

³PT Prodia StemCell Indonesia, Jakarta Pusat-10430, Jakarta, Indonesia

⁴The Prodia Education and Research Institute, Jakarta Pusat-10430, Jakarta, Indonesia

⁵Department of Physiology, Division of Oral Biology, Faculty of Dentistry, Universitas Trisakti, Jakarta Barat-11440, Jakarta, Indonesia

⁶Research Institute, Ballys Co. Ltd, Incheon-22219, Republic of Korea

Received 09 May 2023; revised 18 September 2023

Eleutherine bulbosa (Mill.) Urb. is a medicinal plant which has long been used to treat cancer. *E. bulbosa* bulb extract (EBBE) has been reported to show cytotoxicity towards several types of human cancer. However, the cytotoxic effect of EBBE towards oral cancer cells has not been investigated. The present study aimed to evaluate the effects of EBBE towards the viability, apoptosis, and migratory activities of oral cancer cells. Human oral squamous cell carcinoma (HSC)-3 cells were treated with various concentrations of EBBE for 24 h. The number of viable and apoptotic HSC-3 cells were measured using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) and sub-G1 assay, respectively. The migratory activities of HSC-3 cells were assessed using scratch and transwell assay. Sonic hedgehog (SHH) expression was measured using immunoblotting. Upon EBBE treatment, apoptotic HSC-3 cells were significantly higher in a concentration-dependent manner ($P < 0.05$). Meanwhile, viability, migratory activities, and migrated HSC-3 cell number were significantly lower in a concentration-dependent manner ($P < 0.05$). The SHH expression levels in EBBE-treated HSC-3 cells were also lower in a concentration-dependent manner. EBBE reduces HSC-3 cell viability through apoptosis and inhibits its migratory activities by downregulating SHH expression.

Keywords: Apoptosis, Cell migration, *Eleutherine bulbosa*, Sonic hedgehog, Tongue cancer

Oral squamous cell carcinoma (OSCC) is the most common histological type of oral cancer¹ which arises from squamous epithelial cells in the oral cavity. This cancer generally affects the floor of the mouth and the tongue². Various modes of treatment are developed to manage OSCC, including chemotherapy, radiotherapy, surgery, or its combination³. However, these treatments cause numerous side effects.

Discovery and development of new therapeutic agents derived from natural sources, particularly plants with less side effects are recently one of the main focuses on oncology research. Several plants have been reported to have anticancer effects, including

*Brucea javanica*⁴, *Cinnamomum cassia*⁵, *Myristica fragrans*⁶, and *Oroxylum indicum*⁷. Plants have been known as rich sources of various active compounds, including flavonoids, which was reported to inhibit cancer proliferation *via* cell cycle arrest, antiproliferative and antioxidative activity, apoptosis induction, promotion of differentiation, inhibition of angiogenesis and carcinogen metabolic activation, and multidrug resistance modulation⁸. In addition, several non-flavonoid phenolic compounds have also been reported to show anticancer properties, such as caffeic acid⁹.

Eleutherine bulbosa (Mill.) Urb. is a traditional medicinal plant belonging to the Iridaceae family. Its bulbs are commonly used for the treatment of breast cancer, hypertension, stroke, and diabetes mellitus. It has also been used to increase the production of breast milk and treat sexual disorder¹⁰. *E. bulbosa* bulbs have

*Correspondence:
Phone: +628128077780
E-mail: ferry@trisakti.ac.id

also been reported to have antimicrobial¹¹ and antioxidant¹² activity.

Eleutherine bulbosa contains a wide variety of compounds, primarily naphthoquinones, anthraquinones, and naphthalene¹³. The main compounds found in this plant are eleutherin, isoeleutherin, and eleutherol¹⁴. Several compounds obtained from *E. bulbosa* bulbs have been reported to show anticancer activities, such as 6,8-hydroxy-3,4-di-methoxy-1-methyl-anthraquinone-2-carboxylic acid methyl ester¹⁵, eleutherinoside C, and isoeleutherin¹⁶.

Eleutherine bulbosa bulb extract (EBBE) has been reported to show cytotoxicity towards several types of human cancer, such as cervical cancer (line HeLa)¹⁷, colon cancer (line SW480, HCT116, DLD1, WiDr)^{16,18}, breast cancer (line T47D)¹⁹, leukemia (line K562)¹⁵, and retinoblastoma (WERI-Rb-1)²⁰. However, the cytotoxic effect of EBBE towards human oral cancer cells has not been investigated. Therefore, the present study was conducted to evaluate the effects of EBBE towards the viability, apoptosis, and migratory activities of oral cancer cells.

Materials and Methods

E. bulbosa bulb extract (EBBE) preparation

E. bulbosa bulbs were obtained from Indonesian Medicinal and Aromatic Crops Research Institute (IMACRI), Indonesia. Plant identification was carried out by expert botanist at Research Center for Plant Conservation and Botanic Gardens, Indonesian Institute of Sciences, Indonesia (No. B-1269/IPH.3/KS/X/2020). EBBE was extracted by maceration method⁴. Briefly, bulbs of *E. bulbosa* were minced and dried. The dried material was extracted with ethanol 70%, filtered and evaporated using a rotary evaporator. The resulting crude EBBE was then stored at 4°C.

HSC-3 cell culture

HSC-3 cell culture was performed as previously described⁴ with modification. The HSC-3 cell line was purchased from Sigma-Aldrich Pte. Ltd. (St. Louis, MO, USA). HSC-3 cells were cultured in a complete medium containing Dulbecco's modified Eagle's medium (DMEM) (Sigma-Aldrich), 10% fetal bovine serum (FBS) (PAN-Biotech, Aidenbach, Germany), 50 U/mL penicillin and 50 µg/mL streptomycin (Sigma-Aldrich) in a humidified 5% CO₂, 37°C incubator. HSC-3 cells were then dissociated with trypsin-ethylenediamine tetraacetic acid (EDTA) solution (Sigma-Aldrich) after reaching

80% confluency. Upon reaching the desired cell number, HSC-3 cells were seeded into 24-well plate and 96-well plate for further assays.

Cell viability assay

The number of viable cells were measured quantitatively using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay as previously described²¹ with modification. Briefly, 5×10^3 HSC-3 cells were seeded into 96-well plates and then treated with EBBE (1, 10, or 100 µg/mL), 1 µM Doxorubicin (Dankos Farma, Jakarta, Indonesia) or medium merely for 24 h. After that 100 µL of MTT (Sigma-Aldrich) in culture medium was loaded into each well. After incubating the plates for 4 h, culture medium was discarded, and the formed formazan crystals were dissolved in 100 µL dimethylsulfoxide (DMSO). Results were measured at OD₅₇₀ using a microplate reader (Bio-Rad, Hercules, CA, USA). Each experimental group was measured in six replicates. In addition, the number of untreated cells were counted with hemacytometer and used for interpolating OD₅₇₀ values of HSC-3 cells.

Sub-G1 assay

To investigate the cytotoxic effect of EBBE on HSC-3 cells, the number of apoptotic cells were determined using sub-G1 assay as previously described⁹ with modification. Treated and untreated HSC-3 cells were harvested and suspended in 450 µL hypotonic fluorochrome solution containing 50 µg/mL propidium iodide (Sigma-Aldrich), 0.1% Triton X-100 (Sigma-Aldrich), and 0.1% sodium citrate (Wako, Osaka, Japan). After that, cell suspensions were incubated in the dark for 2 h at room temperature. Fluorescence of individual nuclei was measured using FACSCanto II flow cytometer (Becton Dickinson, Franklin Lakes, NJ, USA) at 100,000 events.

Scratch assay

Scratch assay was performed as previously described²². HSC-3 cells (5×10^4) were seeded into 24-well plates and incubated overnight. Scratch was made on each well with micropipette yellow tips. Then, the wells were washed with phosphate-buffered saline (PBS) (Wako) and treated with/without EBBE or 20 µM cyclopamine tartrate (BioVision, Milpitas, CA, USA) (as positive control) for 24 h. The gap closures were observed and documented under an inverted light microscope (Carl Zeiss, Jena, Germany).

Transwell assay

Transwell assay was performed as previously described²³ with modification. HSC-3 cells (5×10^4) were seeded in the 8- μM pore size chamber insert (Merck, Darmstadt, Germany). After that, 800 μL culture medium with/without EBBE or 20 μM cyclopamine tartrate was loaded into each well of the 24-well plate. HSC-3-containing chamber inserts were then placed on the 24-well plate. After incubating the chamber for 24 h at 37°C, migrated cells in the lower surface of the chamber were fixed. The cells were counted and documented under an inverted light microscope.

Immunoblotting

Expression levels of Sonic hedgehog (SHH) protein were measured using immunoblotting method as previously described²⁴ with modification. HSC-3 cells were lysed and homogenized with Laemmli sample buffer (Bio-Rad). The proteins from each experimental group were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and transferred to a polyvinylidene difluoride (PVDF) sheet. After blocking with 5% skim milk solution, the sheet was incubated with rabbit monoclonal anti-SHH antibody as a primary antibody and horseradish peroxidase (HRP)-conjugated goat anti-rabbit antibody as a secondary antibody (Cell Signaling Technology, Danvers, MA, USA). Immunoblot results were analyzed using ImageJ software (National Institute of Health, Bethesda, MD, USA) to measure the density of each SHH band.

Statistical analysis

Statistical analyses were performed using IBM SPSS Statistics version 26.0 (SPSS IBM, Armonk, NY, USA). Shapiro-Wilk test was used as a normality test. Differences of MTT assay, sub-G1 assay, and scratch assay results between experimental groups were analyzed using Kruskal-Wallis H with Mann-Whitney U *post hoc* analysis. Meanwhile, differences of transwell assay results and SHH band density between experimental groups were analyzed using one-way analysis of variances (ANOVA) with Tukey's honestly significant difference (HSD) *post hoc* analysis. All results were expressed as mean \pm standard deviation (SD). $P < 0.05$ were considered as statistically significant.

Results

EBBE reduced the number of viable HSC-3 cells

The number of viable HSC-3 cells cultured in medium containing extract solvent (sham group) for

24 h was $9,607 \pm 13.81$. Treatment with EBBE showed that the viability of HSC-3 cells was significantly lower compared to the sham group ($P < 0.05$) in a concentration-dependent manner. The number of viable HSC-3 cells treated with 1, 10, and 100 $\mu\text{g/mL}$ EBBE were $9,055 \pm 268.69$, $6,777 \pm 278.04$, and $3,659 \pm 32.86$, respectively. The viability of EBBE-treated HSC-3 cells was significantly higher ($P < 0.05$) than Doxorubicin-treated HSC-3 cells (226 ± 32.34) (Fig. 1). These findings suggested that EBBE could reduce the viability of HSC-3 cells.

EBBE induced apoptosis of HSC-3 cells

Sub-G1 assay results showed that the apoptosis percentage of the sham group was $8.27 \pm 0.14\%$. Apoptotic HSC-3 cells were significantly higher compared to the sham group ($P < 0.05$) in concentration-dependent manner after EBBE treatment. Apoptosis percentages of HSC-3 cells treated with 1, 10, and 100 $\mu\text{g/mL}$ EBBE were $9.60 \pm 0.35\%$, $17.78 \pm 0.32\%$, and $38.64 \pm 0.70\%$, respectively. These percentages were significantly lower ($P < 0.05$) than Doxorubicin-treated HSC-3 cells ($95.95 \pm 2.60\%$) (Fig. 2). Therefore, these results suggested that EBBE treatment could promote apoptosis of HSC-3 cells.

EBBE inhibited HSC-3 cell migration

The gap closure percentage of HSC-3 cells cultured in medium containing extract solvent after 24 h was




Fig. 1 — EBBE decreased the number of viable HSC-3 cells in a concentration-dependent manner. HSC-3 cells were cultured and treated with/without 1 μM Doxorubicin and various concentrations of EBBE for 24 h as indicated in the panel. The number of viable cells were measured with MTT assay as described in Materials and Methods. The data are expressed as mean \pm SD ($n=6$). * $P < 0.05$ compared to the sham group




Fig. 2 — EBBE increased the percentage of apoptotic HSC-3 cells in a concentration-dependent manner. HSC-3 cells were cultured and treated with/without 1 μM Doxorubicin and various concentrations of EBBE for 24 h as indicated. Treated cells were then subjected to sub-G1 assay as described in Materials and Methods. (A) Histogram of flow cytometric results. (B) Percentage of apoptotic HSC-3 cells. The data are expressed as mean \pm SD (n=18). * $P < 0.05$ compared to the sham group

96.11 \pm 4.17%. After EBBE treatment, the migratory activities of HSC-3 cells were significantly lower compared to the sham group ($P < 0.05$) in a concentration-dependent manner. Gap closure percentage of HSC-3 cells treated with 1, 10, and 100 $\mu\text{g/mL}$ EBBE were 89.56 \pm 5.77%, 76.78 \pm 4.87%, and 60.67 \pm 7.62%, respectively. These percentages were significantly higher ($P < 0.05$) than cyclopamine tartrate-treated HSC-3 cells (43.00 \pm 4.82%) (Fig. 3).

Transwell assay also showed similar results. The number of migrated HSC-3 cells in the sham group was 134.44 \pm 11.60. The number of migrated HSC-3 cells was significantly lower than the sham group ($P < 0.05$) in a concentration-dependent manner after EBBE treatment. The number of migrated HSC-3 cells treated with 1, 10, and 100 $\mu\text{g/mL}$ EBBE were 107.44 \pm 15.21, 95.11 \pm 6.01, and 39.22 \pm 3.56, respectively. These numbers were significantly higher ($P < 0.05$) than cyclopamine tartrate-treated HSC-3 cells (17.89 \pm 3.22) (Fig. 4). Overall, these results

indicated that EBBE exhibited an anti-migratory effect towards HSC-3 cells.

EBBE downregulated the expression of SHH

Upon the addition of EBBE, the SHH expression levels in HSC-3 cells were lower in a concentration-dependent manner. Expression level of SHH in HSC-3 cells treated with 1 $\mu\text{g/mL}$ EBBE was significantly higher ($P < 0.05$), while SHH expression levels in HSC-3 cells treated with 10 and 100 $\mu\text{g/mL}$ EBBE were significantly lower ($P < 0.05$) compared to the sham group. SHH expression levels in HSC-3 cells treated with 1 and 10 $\mu\text{g/mL}$ EBBE were significantly higher compared to cyclopamine tartrate-treated HSC-3 cells ($P < 0.05$). However, no significant difference was observed in the SHH expression levels between cyclopamine tartrate and 100 $\mu\text{g/mL}$ EBBE treatment ($P = 0.142$) (Fig. 5). Thus, these findings suggested that EBBE could decrease the expression of SHH.




Fig. 3 — EBBE inhibited migratory activities of HSC-3 cells. HSC-3 cell culture was scratched and treated with/without EBBE or 20 µM cyclopamine tartrate for 24 h as indicated. Gap closure was measured as described in Materials and Methods. (A) Scratched areas of HSC-3 cell culture. White bar: 100 µM. (B) Gap closure percentage of EBBE-treated HSC-3 cell. The data are expressed as mean ± SD (n=9). **P* < 0.05 compared to the sham group




Fig. 4 — EBBE reduced the number of migrated HSC-3 cells. HSC-3 cell culture was treated with/without EBBE or 20 µM cyclopamine tartrate for 24 h as indicated. Migrated cells were measured as described in Materials and Methods. The data are expressed as mean ± SD (n=9). **P* < 0.05 compared to the sham group




Fig. 5 — EBBE reduced the expression of SHH. HSC-3 cell culture was treated with/without EBBE or 20 µM cyclopamine tartrate for 24 h as indicated. HSC-3 cells were lysed, homogenized, and immunoblotted using anti-SHH antibody as described in Materials and Methods. The data are expressed as mean ± SD (n=3). **P* < 0.05 compared to the sham group

Discussion

In the present study, we investigated the potency of EBBE as anticancer agent for OSCC treatment. EBBE showed a cytotoxic effect towards HSC-3 cells by reducing the viability of HSC-3 cells in a concentration-dependent manner. According to the sub-G1 assay results, the decrease in HSC-3 cell viability was caused by apoptosis. These results were in line with previous studies that EBBE was able to induce apoptosis of cancer cells^{18-20,25}. EBBE was also reported to inhibit cell cycle in several cell lines^{19,25}, indicating that EBBE may induce apoptosis of HSC-3 cells through cell cycle arrest since both processes are interconnected²⁶. Based on a molecular docking study, eleutherol, eleutherin, and isoeleutherin could induce apoptosis by activating caspase-8²⁷.

Cell migration is one of the key processes involved in cancer metastasis²⁸. Scratch and transwell assays demonstrated that EBBE restrained the migration of HSC-3 cells in a concentration-dependent manner. This showed the novel activity of EBBE in impairing HSC-3 cell migration. Therefore, EBBE may have anti-metastasis activity towards tongue cancer cells.

Metastasis of tongue cancer is regulated by several proteins, including SHH. One of the SHH signaling pathways that has been reported to regulate cell migration in tongue cancer is SHH/glioma-associated oncogene homologue (SHH/GLI)²⁹. SHH protein binds and inactivates its receptor, Patched 1 (PTCH1). This interaction causes the activation of transcription factors from the GLI family, which play a major role in regulating cancer migration, invasion, and proliferation³⁰.

The present study revealed that EBBE decreased SHH expression levels along with the increase in EBBE concentrations, suggesting that the inhibitory effect of EBBE towards the expression of SHH may be responsible for its anti-migratory property. To the best of our knowledge, this is the first study demonstrating SHH downregulation in EBBE-treated HSC-3 cells. EBBE may inhibit HSC-3 cell migration via inhibition of SHH/GLI signaling pathway. Possibly, all signaling pathways that are regulated by SHH can be inhibited by EBBE. EBBE may also restrain HSC-3 cell migration through another pathway. In SW480 colon cancer cell line, eleutherinoside C and isoeleutherin obtained from *E. bulbosa* bulbs was reported to inhibit Wnt/ β -catenin pathway¹⁶, which has been reported to play an important role in regulating cancer migration³¹.

Therefore, signaling pathways involved in the inhibition of HSC-3 cell migration by EBBE still need to be explored. In addition, signaling components downstream to SHH and target molecules that are affected by EBBE should be identified. Since EBBE also contains various bioactive compounds, further research is needed to identify which compounds are responsible for apoptosis and anti-migratory activity. Thus, more investigations are required to understand the molecular mechanisms underlying EBBE-induced HSC-3 cell apoptosis and confirm which SHH signaling pathway contributes to the inhibition of cell migration by EBBE.

Conclusion

EBBE reduces HSC-3 cell viability through apoptosis and inhibits its migratory activities by **down regulating** SHH expression. Taken together, EBBE could be a potential anti-tongue cancer agent, particularly as an **ant metastatic** agent.

Conflict of interest

All authors declare no conflicts of interest.

References

- Chen YK, Huang HC, Lin LM & Lin CC, Primary oral squamous cell carcinoma: An analysis of 703 cases in southern Taiwan. *Oral Oncol*, 35 (1999) 173.
- Neville BW & Day TA, Oral cancer and precancerous lesions. *CA Cancer J Clin*, 52 (2002) 195.
- Minhas S, Kashif M, Altaf W, Afzal N & Nagi AH, Concomitant-chemoradiotherapy-associated oral lesions in patients with oral squamous-cell carcinoma. *Cancer Biol Med*, 14 (2017) 176.
- Wicaksono BD, Tangkearung E & Sandra F, *Brucea javanica* leaf extract induced apoptosis in human oral squamous cell carcinoma (HSC2) cells by attenuation of mitochondrial membrane permeability. *Indones Biomed J*, 7 (2015) 107.
- Chang WL, Cheng FC, Wang SP, Chou ST & Shih Y, *Cinnamomum cassia* essential oil and its major constituent cinnamaldehyde induced cell cycle arrest and apoptosis in human oral squamous cell carcinoma HSC-3 cells. *Environ Toxicol*, 32 (2017) 456.
- Susianti S, Lesmana R, Salam S, Julaha E, Pratiwi YS, Sylviana N, Goenawan H, Kumiawan A & Supratman U, The effect of nutmeg seed (*M. fragrans*) extracts induces apoptosis in melanoma maligna cell's (B16-F10). *Indones Biomed J*, 13 (2021) 68.
- KameyandaPoonacha** S, Harishkumar M, Radha M, Varadarajan R, Nalilu SK, Shetty SS, Shetty PK, Chandrashekhara RB, Sreenivas MG & Bavabeedu SKB, Insight into oroxylinA-7-O- β -d-glucuronide-enriched *Oroxylum indicum* bark extract in oral cancer HSC-3 cell apoptotic mechanism: Role of mitochondrial microenvironment. *Molecules*, 26 (2021) 7430.
- Ren W, Qiao Z, Wang H, Zhu L & Zhang L, Flavonoids: Promising anticancer agents. *Med Res Rev*, 23 (2003) 519.

- 9 Sandra F & Sidharta MA, Caffeic acid induced apoptosis in MG63 osteosarcoma cells through activation of caspases. *Mol Cell Biomed Sci*, 1(2017)28.
- 10 Ieyama T, Gunawan-Puteri MDPT & Kawabata J, α -Glucosidase inhibitors from the bulb of *Eleutherine americana*. *Food Chem*, 128 (2011) 308.
- 11 Subramaniam K, Suriyamoorthy S, Wahab F, Sharon FB & Rex GR, Antagonistic activity of *Eleutherine palmifolia* Linn. *Asian Pac J Trop Dis*, 2 (2012) S491.
- 12 Kuntorini EM, Dewi M & Misrina M, Anatomical structure and antioxidant activity of red bulb plant (*Eleutherine americana*) on different plant age. *Biodiversitas*, 17 (2016) 229.
- 13 Insanu M, Kusumardiyani S & Hartati R, Recent studies on phytochemicals and pharmacological effects of *Eleutherine americana* Merr. *Procedia Chem*, 13 (2014) 221.
- 14 Chen Z, Huang H, Wang C, Li Y, Ding J, Sankawa U, Noguchi H & Itaka Y, Hongconin, a new naphthalene derivative from hong-cong, the rhizome of *Eleutherine Americana* Merr. et Heyne (Iridaceae). *Chem Pharm Bull*, 34 (1986) 2743.
- 15 Xu J, Qiu F, Duan W, Qu G, Wang N & Yao X, New bioactive constituents from *Eleutherine americana*. *Front Chem China*, 1 (2006) 320.
- 16 Li X, Ohtsuki T, Koyano T, Kowithayakorn T & Ishibashi M, New Wnt/ β -catenin signaling inhibitors isolated from *Eleutherine palmifolia*. *Chem Asian J*, 4(2009) 540.
- 17 Mutiah R, Minggarwati TS, Kristanti RA & Susanti E, Compound identification and anticancer activity of ethyl acetate fraction from bawang sabrang (*Eleutherine palmifolia* (L.) Merr.) on HeLa cervical cancer cell line. *Indones J Cancer Chemoprevent*, 10 (2019) 131.
- 18 Lubis IA, Ichwan M, Mustofa M & Satria D, Anticancer activity of *Eleutherine bulbosa* (Mill.) Urb. extract on WiDr cell line *in vitro*, paper presented to 2nd Public Health International Conference (PHICo 2017), Faculty of Public Health – Universitas Sumatera Utara, Medan, Indonesia, 18-19 December 2017.
- 19 Yuniarti A, Sundhani E & Nurulita NA, The potentiation effect of bawangdayak (*Sisyrinchium palmifolium* L.) extract on T47D cell growth inhibition after 5-fluorouracil treatment. *Pharmaciana*, 8 (2018) 195.
- 20 Kamarudin AA, Sayuti NH, Saad N, Razak NAA & Esa NM, Induction of apoptosis by *Eleutherine bulbosa* (Mill.) Urb. bulb extracted under optimised extraction condition on human retinoblastoma cancer cells (WERI-Rb-1). *J Ethnopharmacol*, 284 (2022) 114770.
- 21 Sandra F, Sudiono J, Trisfilha P & Pratiwi D, Cytotoxicity of *Alpinia galanga* rhizome crude extract on NIH-3T3 cells. *Indones Biomed J*, 9(2017) 23.
- 22 Ariesanti Y, Sandra F, Claresta B & Alvita L, *Coffea canephora* bean extract induces NIH3T3 cell migration. *Indones Biomed J*, 13(2021) 216.
- 23 Oktaviono YH, Al-Farabi MJ, Suastika LOS, Hartono F, Dirgantara Y & Sandra F, Preliminary study: Purple sweet potato extract seems to be superior to increase the migration of impaired endothelial progenitor cells compared to L-Ascorbic acid. *Sci Pharm*, 87 (2019) 16.
- 24 Rizal MI & Sandra F, *Brucea javanica* leaf extract activates caspase-9 and caspase-3 of mitochondrial apoptotic pathway in human oral squamous cell carcinoma. *Indones Biomed J*, 8 (2016) 43.
- 25 Mutiah R, Listiyana A, Suryadinata A, Annisa R, Hakim A, Anggraini W & Susilowati R, Activity of inhibit the cell cycle and induct apoptosis in HeLa cancer cell with combination of sabrang onion (*Eleutherine palmifolia* (L.) Merr) and starfruit mistletoe (*Macrosolen cochinchinensis* (Lour.) Tiegh). *J App Pharm Sci*, 8 (2018) 122.
- 26 Pucci B, Kasten M & Giordano A, Cell cycle and apoptosis. *Neoplasia*, 2(2000) 291.
- 27 Quadros Gomes AR, da Rocha Galucio NC, de Albuquerque KCO, Brigido HPC, Varela ELP, Castro ALG, Vale VV, Bahia MO, Rodriguez Burbano RM, de Molfeta FA, Carneiro LA, Percario S & Dolabela MF, Toxicity evaluation of *Eleutherine plicata* Herb. extracts and possible cell death mechanism. *Toxicol Rep*, 8 (2021) 1480.
- 28 Ellis IR, The migration and invasion of oral squamous carcinoma cells: Matrix, growth factor and signalling involvement. *Cancers*, 13 (2021) 2633.
- 29 Fan HX, Wang S, Zhao H, Liu N, Chen D, Sun M & Zheng JH, Sonic hedgehog signaling may promote invasion and metastasis of oral squamous cell carcinoma by activating MMP-9 and E-cadherin expression. *Med Oncol*, 31 (2014) 41.
- 30 Carballo GB, Honorato JR, de Lopes GPF & Spohr TCLSE, A highlight on Sonic hedgehog pathway. *Cell Commun Signal*, 16 (2018) 11.
- 31 Liang S, Zhang S, Wang P, Yang C, Shang C, Yang J & Wang J, LncRNA, TUG1 regulates the oral squamous cell carcinoma progression possibly *via* interacting with Wnt/ β -catenin signaling. *Gene*, 608 (2017) 49.