



HOME / About the Journal

About the Journal

Journal of Drug Delivery and Therapeutics (JDDT) is a *double-blind peer-reviewed monthly* journal dedicated to the publication of research papers, reviews, mini-reviews, Short communications and case studies. It is an official journal of Society of Pharmaceutical and Medical Professionals ([SPMP](#)). This publication is aimed at a broad, interdisciplinary audience of academic and industrial researchers actively engaged in basic and applied laboratory practice, related to Pharmaceutical Science and Therapeutics.

COPYRIHT NOTICE

This journal has chosen to apply the [Creative Commons Attribution-NonCommercial 4.0 International \(CC BY-NC 4.0\)](#) to all papers to be published. CC BY-NC - Allows users to copy, distribute and transmit an article, adapt the article as long as the author is attributed and the article is not used for commercial purposes.

Authors who wish to publish withis journal agree to the following terms;

1. Authors retain copyright and grant the journal right of first publication with the work simultaneously licensed under a [Creative Commons Attribution-NonCommercial 4.0 International \(CC BY-NC 4.0\)](#). that allows others to share the work with an acknowledgment of the work's authorship and initial publication in this journal.
2. Authors are able to enter into separate, additional contractual arrangements for the non-exclusive distribution of the journal's published version of the work (e.g., post it to an institutional repository or publish it in a book), with an acknowledgment of its initial publication in this journal.
3. Authors are permitted and encouraged to post their work online (e.g., in institutional repositories or on their website) prior to and during the submission process, as it can lead to productive exchanges, as well as earlier and greater citation of published work (See [The Effect of Open Access](#)).

Journal Title: Journal of Drug Delivery and Therapeutics

Focus & Scope: Pharmaceutical Sciences & Medical Sciences

Publisher: Society of Pharmaceutical and Medical Professionals ([SPMP](#))

Publisher Office: 1/7756, Street no. 1, East Gorakh Park, Shahdara, Delhi, India 110032

Country: India

Language: English

Publishing Frequency: Monthly

Editor Name: Ram C Dhakar (**Email:** editor.jddt@gmail.com, **Phone:** +91-9783920207)

Editorial Office: SRG Hospital & Medical College, Jhalawar, Rajasthan, India

Website: jddtonline.info/index.php/jddt

e-ISSN: 2250-1177

Nature of Publication: Online (No Print version)

Year of Starting: 2011

Journal Code:

ISSN: 2250-1177

CODEN(USA-NLM): JDDTAO

NLM ID: 101656626

WorldCat ID: 879704190

CrossRef-DOI prefix: 10.22270

Publication charges (See [Link](#))

Contact: Email- editor.jddt@gmail.com, **Phone-** +91-9783920207

Editor-in-Chief: Ram C Dhakar, SRG Hospital & Medical College, Jhalawar, Rajasthan, India



Online ISSN: 2250-1177

MAKE A SUBMISSION

CATEGORIES

Aims & Scope

Editorial Policies

Publication Ethics

Plagiarism Policy

Retraction Policy

[Open Access Policy](#)

[Processing Charges and Waiver Policy](#)

[Indexing](#)

LATEST PUBLICATIONS

ATOM 1.0

RSS 2.0

RSS 1.0

[Announcements](#) || [Editorial Board](#) || [Indexing](#) || [Contact](#)

The publication is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](#) (CC BY-NC 4.0). [View Legal Code](#)



HOME / Editorial Board

Editorial Board

EDITOR-IN-CHIEF

Dr. Ram C Dhakar, Hospital Pharmacy, SRG Hospital and Medical College Jhalawar, India-326001, India. Email: editor.jddt@gmail.com Phone: +91-9783920207 [Google Scholar](#) [ResearchGate](#) [ORCID](#) [SCOPUS](#) Research fields: Novel Drug delivery, Nanotechnology, Microspheres, TDDS

ASSOCIATE EDITORS

Dr. Aamir Jalal Al-Mosawi, Advisor And Expert Trainer, Baghdad Medical City And The National Training And Development Center, Iraqi Ministry Of Health, Baghdad, Iraq. [ResearchGate](#) [Google Scholar](#) [SCOPUS](#) [ORCID](#) [NCBI-NLM](#) Email: almosawiaj@yahoo.com. Research Fields: Pediatrics, pediatric psychiatry, nephrology, pediatric neurology, genetics

Dr. S K Prajapati, Professor & Head, Institute of Pharmacy, Bundelkhand University, Jhansi (U.P.), India, India. Email: drsunilprajapati@gmail.com [ResearchGate](#) [Website](#) [Google Scholar](#) Research Area: Novel Drug Delivery, Microspheres, Nanoparticles

Dr. Rajesh Ramasamy, Associate Professor, Head of Department, Department of Pathology, Faculty Medicine and Health Science, Universiti Putra Malaysia, Malaysia Phone: +603 8947 2377 Mobile: +6012 393 6444 Fax: +603 8941 2787 Email: rajesh@upm.edu.my [Google Scholar](#) [ResearchGate](#) [Website](#) SCOPUS ID: 26024442700 Research Fields: Stem Cells, Immunology, Diabetes, Medicinal Plant Research, Sidha Medicine

Dr. Syed Mohd Danish Rizvi, Associate Professor, Department of Pharmaceutics, College of Pharmacy, University of Hail, Hail, Saudi Arabia. Email: syedrizvi10@yahoo.com Official Email: sm.danish@uoh.edu.sa | [ORCID ID](#) | [Google Scholar](#) | [Scopus](#) | [Research Gate](#) | [Website](#) | Research fields: Microspheres, Nanoparticles, Cancer Therapy

SECTION EDITORS (Scientific Committee Members)

Dr. Narendar Dudhipala, Department of Pharmaceutics and Drug Delivery, School of Pharmacy, University of Mississippi, USA. Research Interests: Formulation development, preclinical evaluation of small molecules for ocular delivery and oral delivery, Tumor delivery, lipid nanoparticles, lyophilization. Mobile: +1 281 919 5546 Email: narphmreddy@gmail.com ; [View Profile](#) [ResearchGate](#)

Dr. Priyanka Bhatt, Department of Pharmaceutical Sciences, College of Pharmacy, University of South Florida, 12901 Bruce B. Downs Blvd., MDC30, Tampa, FL 33612, Email: priyankabhatter88@yahoo.com psbhatt@health.usf.edu [Google Scholar](#) [ResearchGate](#) [Website](#)

Dr. Siddhartha Pati, Research Associate, Centre of Excellence, Khallikote University, Odisha, India. +91-8895577750/7008251372 Patisiddhartha@gmail.com [ResearchGate](#) [ORCID](#)

Dr. Iqrar A. Ansari, Assistant Professor, Department of Pharmaceutical Chemistry, Prof. Ravindra Nikam College of Pharmacy, Gondur, Dhule, 424002, Maharashtra, India. Mobile No.+91 9284541386, Email:

ansariqrar50@gmail.com, [Google Scholar](#), [Researchgate](#), [LinkedIn](#) Research Fields: [EGFR Inhibitors](#), [CADD](#), [Docking](#), [MD Simulation](#)

Dr. Prakash Doddapattar, Carver College of Medicine, University of Iowa, USA. Research Fields: Molecular Biology · Immunology · Cardiovascular Biology. [ResearchGate](#) [LinkedIn](#) [OICID ID](#) Email: prakashdpbt@gmail.com

Dr. Ashok K Datusalia, Assistant Professor, National Institute of Pharmaceutical Education and Research, Raebareli, Uttar Pradesh, 226002, India Email: ashokdatusalia@gmail.com [Google Scholar](#) [ResearchGate](#) [Website](#) Research Interests: Neuroscience, Pharmacology and Toxicology

Dr. Dinesh Nyavanandi, Formulation Scientist Continuous Pharmaceuticals 25 Olympia Ave, Woburn, Massachusetts, USA. 01801. (P): +1(662) 550-9555 (E): ndinesh624@gmail.com; dnyavanandi@continuuuspharma.com *Research Experience*: Oral drug delivery, Lipid based drug delivery, Strategies for enhancement of solubility and bioavailability, Gastro-retentive Drug delivery, Ocular Drug Delivery, Topical Drug Delivery [LinkedIn](#) [Google Scholar](#) [ORCID](#) [SCOPUS](#)

INTERNATIONAL EDITORIAL MEMBERS

Prof. Dr. rer. Nat. Hesham A. El Enshasy, Director, Institute of Bioproduct Development (IBD), Universiti Teknologi Malaysia (UTM), 81310 UTM, Johor Bahru, Malaysia. Tel: 006-07-5531573 (office), Fax: 006-07-5569706, E-mail: henshasy@ibd.utm.my [ResearchGate](#) [Google Scholar](#)

Dr. Sitansu Sekhar Nanda, Department of Chemistry, Myongji University, Yongin, South Korea, Email: nandasitansusekhar@gmail.com [Google Scholar](#) [ResearchGate](#) [Website](#)

Dr. Kishor Mazumder, Associate Professor at Jashore University of Science and Technology, Chattogram, Bangladesh Email: Mazumderkis2002@Yahoo.Com kmazumder@csu.edu.au kmazumder@just.edu.bd [LinkedIn](#) [Google Scholar](#) [ResearchGate](#) [ORCID ID](#)

Dr. Priya Bharate, Analytical Scientist, Continuous Pharmaceuticals, MA, USA. [Google Scholar](#) [ResearchGate](#) [Website](#) Email: pbharate@continuuuspharma.com Interested fields: Carbohydrates synthesis, Chemical biology, Supramolecular chemistry, Analytical method development and qualifications of drug molecules, Nanomaterials, and organic chemistry

Dr. Ahmed Salih Sahib; Professor, Dept of Pharmacology & Toxicology, College of Pharmacy, University of Kerbala, Iraq, Email: ahmedsalih73@yahoo.com ahmed.alkhezaly@uokerbala.edu.iq Tel : +964-7901585579 [Google Scholar](#) [ResearchGate](#) [Website](#) Interested fields: Pharmacology, Clinical Pharmacology, Pharmacogenomics

Prof. Fahad I. Al-Saikhan, Department of Clinical Pharmacy, Prince Sattam Bin Abdulaziz University, College of Pharmacy, Kingdom of Saudi Arabia Email: f.alsaikhan@psau.edu.sa Telephone: +966115886074 [ResearchGate](#) [Website](#) Interested fields: Primary care/Ambulatory care, health outcome, pharmacogenomics

Dr. Nahlah Elkudssiah Ismail, Professor, Department of Clinical Pharmacy, Malaysian Academy of Pharmacy, Malaysia. researchgate.net orcid.org [LinkedIn](#) [Google Scholar](#) [Official website](#) Email: elkudssiah77@yahoo.com, Research interest: Pharmaceutics (drug delivery; nanotechnology), Pharmaceutical Analysis (method development and validation), Clinical Pharmacy, Pharmacy Education, Pharmacy Practice & Public Health.

Dr. Vijai Kumar Reddy Tangadanchu, Ph. D. Postdoctoral Research Associate, Mallinckrodt Institute of Radiology, Washington University School of Medicine Saint Louis, Missouri-63110, USA E-mail:

tvjajkumarreddy@wustl.edu tvkreddyict@outlook.com Mobile: +1 314-665-8974. [ResearchGate](#)
[Publons](#) [SciProfile](#)

Dr. Solmaz Ghaffari, Department of Pharmaceutics, Faculty of Pharmacy, Islamic Azad University, Tehran, Iran; Email: soligh@yahoo.com [Google Scholar](#) [researchgate.net](#) Tel: +9891252721 Fax: +982122600091

Dr. Mostafa Madmoli, Emergency Medical Technician, Dezful University of Medical Sciences, Dezful, Iran. Email: mostafamadmoli10@yahoo.com Phone: [989370176590] Interested fields: Diabetes, Alzheimer's Disease,

Dr. Vijay Kumar Siripuram, Laboratory of Molecular and Cellular Neuroscience The Rockefeller University, New York, NY, USA. Email: vijay.aleru@gmail.com ; Phone: (+1) 646-238-5620. Interested fields: NMR, FTIR, HPLC, GCMS/LCMS, column chromatography, crystallization [Google Scholar](#)
[Publons](#)

Dr Saket Patel, Research Associate at Memorial Sloan Kettering Cancer Center, Department of Radiology, New York, 10065, USA. Interested fields: Organic chemistry, new molecule design, developing new methods for hyperpolarization via dissolution DNP, and real time metabolism (in vivo/ in vitro). Phone: +1 929-553-4960 Email: saketpatel32@gmail.com patels24@mskcc.org [Google Scholar](#)
[Website](#)

Dr Arun Butreddy, Ph.D., Scientist, Protein Pharmaceutical Development, Biogen Inc, Massachusetts, USA. [ResearchGate](#) [Google Scholar](#) [LinkedIn](#) Email: butreddyaran@gmail.com Mobile: +1 (662)-380-0558

Dr. Madhukiran Parvathaneni, Manager - Global Regulatory Affairs, CRC Pharma, New Jersey, USA. Interested Fields: Regulatory Affairs, Clinical Research, Pharmaceuticals and Medical Devices. Ph: +1 (609)582-6496 Email: madhukiran.parvathaneni@gmail.com [Website](#) [ResearchGate](#) [Google Scholar](#)
[LinkedIn](#)

Praveen Kolimi, Department of Pharmaceutics and Drug Delivery, University of Mississippi, Faser Hall, MA, 38677, USA Mail: praveenkoliminiper@gmail.com [Google Scholar](#) [ResearchGate](#) [LinkedIn](#)

ASSISTANT EDITORS

Dr. Dito Anurogo, M.D., M.Sc., International PhD Program for Cell Therapy and Regeneration Medicine (IPCTRM), College of Medicine, Taipei Medical University Taiwan and Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Makassar, Makassar, Indonesia. Mail: dito.anurogo@med.unismuh.ac.id d151109004@tmu.edu.tw Phone: +62-81224422693. [Google Scholar](#)
[Publons](#) [LinkedIn](#) Research Interest: drugs and delivery, pharmacology, regenerative medicine, oncohematology, neuroscience, molecular biology & medicine, bioinformatics, nanomedicine.

Dr. Jacopo Forte, Fixed-term Researcher (RTD-A) at the Department of Basic Biotechnological Sciences, Catholic University of the Sacred Heart, Rome, Research interests: drug delivery like nanosystems for anticancer drug delivery, vesicular systems based on phospholipids and surfactants suitable for antitubercular drugs, and nanocarriers for targeted delivery to the bladder. [ResearchGate](#) [SciProfiles](#)
[Website](#) [ORCID](#) Email: jacopo.forte@unicatt.it

Dr. Aktsar Roskiana Ahmad, Faculty of Pharmacy, Universitas Muslim Indonesia, Urip Sumoharjo Km. 5, 90231, Indonesia. Email: aktsar.roskiana@umi.ac.id Phone: +6281355184404 [Google Scholar](#)
[ResearchGate](#)

Dr. Sheo Datta Maurya, Sarvodaya College of Pharmacy, Azamgarh, India Email: meetshiv23@gmail.com [Google Scholar](#) [ResearchGate](#) Interested fields: Pharmacy (Novel Drug delivery, Liposomes, Ethosomes,)

Dr. Sandeep K. Singh, Postdoctoral Research Associate, Department of Pharmacy Practice and Science, University of Nebraska Medical Center, Omaha, NE, USA. [Google Scholar](#) sasinghlko@gmail.com

Dr. Mousumi Kar, Professor and Head, Faculty of Pharmacy Medi-Caps University, Indore, INDIA, India. Email: karmousumi@hotmail.com [Website](#) [Google Scholar](#) [ResearchGate](#) [LinkedIn](#) [Website](#) Interested fields: Pharmaceutics, Novel Drug Delivery

Dr. Sachin Goyal, Research Associate, Department of Anesthesiology and Critical Care medicine, School of medicine, University of New Mexico, Albuquerque, NM 87106 USA Email: sgoyal@salud.unm.edu

Mr. Vaibhav Patel, Director (Quality Assurance), Elucida Oncology Inc., 1 Deer Park Drive, Suite E Monmouth Junction, NJ 08852, USA. Website Email: Vpatel0212@gmail.com [Google Scholar](#) Phone No: (551) 208-5652

Mr. Girish Nihalani, Research Fellow of Drug Product Development at Teva Pharmaceuticals Inc. (Actavis) based at Florida, USA. e-mail: girilani679@gmail.com girish9122@gmail.com Contact: +1- 954-647-7740 [LinkedIn](#) [Google Scholar](#) Interested fields: Pharmaceutics, Novel Drug Delivery

Dr. Rajanikant Patel, Ph.D. (Pharmaceutics), Senior Scientist, Product Development Granules Pharmaceuticals Inc. 3701 Concorde Pkwy, Chantilly, Virginia 20151, USA Mobile: +1 571 762 9399 Email: rajnipharmacy@gmail.com [Google Scholar](#) [ResearchGate](#) [ORCID](#)



Online ISSN: 2250-1177

MAKE A SUBMISSION

CATEGORIES

[Aims & Scope](#)

[Editorial Policies](#)

[Publication Ethics](#)

[Palgiarism Policy](#)

[Retraction Policy](#)

[Open Access Policy](#)

[Processing Charges and Waiver Policy](#)

[Indexing](#)

LATEST PUBLICATIONS

[ATOM 1.0](#)

[RSS 2.0](#)

[RSS 1.0](#)

[Announcements](#) || [Editorial Board](#) || [Indexing](#) || [Contact](#)

The publication is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](#) (CC BY-NC 4.0). [View Legal Code](#)



HOME / ARCHIVES / Vol. 13 No. 7 (2023): Volume 13, Issue 7, July 2023

Vol. 13 No. 7 (2023): Volume 13, Issue 7, July 2023

PUBLISHED: 2023-07-15

Research

PREVENTIVE EFFECT OF TOTAL ETHANOLIC EXTRACT OF BUTEA MONOSPERMA ON BONE LOSS IN OSTEOPENIC RATS

 Abstract Display: 656 /  PDF Downloads: 685 /  PDF Downloads: 44

Sonu Khanka, Amardeep Lakra, Ritika Gupta, Kriti Sharma, Rakesh Maurya, Naibedya Chattopadhyay, Divya Singh
6-14

 [VIEW ABSTRACT](#)

 [PDF](#)



<https://doi.org/10.22270/jddt.v13i7.5887>

 [DOWNLOAD PDF](#)

[HTML](#)

FORMULATION AND EVALUATION OF BUCCOADHESIVE TABLET OF VERAPAMIL HYDROCHLORIDE FOR THE TREATMENT OF HYPERTENSION

 Abstract Display: 512 /  PDF Downloads: 608 /  PDF Downloads: 104

Sohan Sukaldas Thipe, Amol V Sawale, Shraddha Dilip Muneshwar, Shiwani A Gulhane, Sakshi S Gahukar
15-21

 [VIEW ABSTRACT](#)

 [PDF](#)



<https://doi.org/10.22270/jddt.v13i7.6122>

 [DOWNLOAD PDF](#)

[HTML](#)

PHENOLIC PROFILE AND ANTIOXIDANT POTENTIAL OF AQUEOUS EXTRACTS OF SELECTED TRADITIONAL ANTI-COUGH PLANTS

 Abstract Display: 723 /  PDF Downloads: 670 /  PDF Downloads: 174

Olachi Lilian Osuagwu, Chidi Uzoma Igwe, Chiamaka Perpetua Nzebude, Onyinyechi Chijioke Njoku, Juliet Chioma Ejiofor
22-29

 [VIEW ABSTRACT](#)

 [PDF](#)



<https://doi.org/10.22270/jddt.v13i7.6124>

 [DOWNLOAD PDF](#)

[HTML](#)

EMERGING PROBLEM EVALUATION OF HYPOCALCEMIA IN SUPER VASMOL HAIR DYE POISONING PATIENTS: CLINICAL STUDY IN RIMS, KADAPA

 Abstract Display: 673 /  PDF Downloads: 512 /  PDF Downloads: 292

K. Leela Prasad Babu, Shaik Kareemulla, M. Ramani, L. Nagamani, B. Abhishek, K. Sravanthi Bai
30-33

 [VIEW ABSTRACT](#)

 [PDF](#)



<https://doi.org/10.22270/jddt.v13i7.6116>

 [DOWNLOAD PDF](#)

[HTML](#)

CAN MORINGA OLEIFERA LEAF ETHYL ACETATE EXTRACT INHIBIT CANDIDA ALBICANS PLANKTONIC CELL GROWTH AND BIOFILM FORMATION IN VITRO?

 Abstract Display: 629 /  PDF Downloads: 767 /  PDF Downloads: 100

Herastuti Sulistyani, Siti Sulastri, Dewi Risnawati, Dewi Agustina
34-37

 [VIEW ABSTRACT](#)

 [PDF](#)



<https://doi.org/10.22270/jddt.v13i7.6146>

 [DOWNLOAD PDF](#)

[HTML](#)

EXPLORING DRUG PRESCRIPTION PATTERNS IN THE NEUROLOGY DEPARTMENT OF A TERTIARY CARE HOSPITAL: A COMPREHENSIVE ANALYSIS

: A COMPREHENSIVE ANALYSIS

 Abstract Display: 679 /  PDF Downloads: 662 /  PDF Downloads: 235

R Nithya, Solanky K Neesha, UDHAYAKUMAR Narmadha, R. R. Redlin, A Ramya, B Manisha
38-45

 [VIEW ABSTRACT](#)

 [PDF](#)



<https://doi.org/10.22270/jddt.v13i7.6139>

 [DOWNLOAD PDF](#)

[HTML](#)

FORMULATION AND EVALUATION OF POLYHERBAL TABLET BY USING NEEM, TULSI, TURMERIC AND GINGER EXTRACT

 Abstract Display: 2373 /  PDF Downloads: 1298 /  PDF Downloads: 2139

Souvik Paul, Tanmoy Dey, Prabin Koirala, Sidhant Tamang, Soumik Bhattacharya, Rajat Das
46-51

 [VIEW ABSTRACT](#)

 [PDF](#)



<https://doi.org/10.22270/jddt.v13i7.5895>

 [DOWNLOAD PDF](#)

[HTML](#)

FACTORS ASSOCIATED WITH PLASMA MALONDIALDEHYDE LEVELS IN PEOPLE OVER 40 YEARS

 Abstract Display: 1069 /  PDF Downloads: 840 /  PDF Downloads: 62

Meiyanti Meiyanti, , Yohana, Eveline Margo, [Juni Chudri](#), Pusparini, Meutia Atika Faradilla
52-56

[VIEW ABSTRACT](#)[PDF](#)<https://doi.org/10.22270/jddt.v13i7.6142>[DOWNLOAD PDF](#)[HTML](#)

ANALYSIS OF THE FACTORS THAT INFLUENCE THE FAMILY IN THE USE OF TRADITIONAL MEDICINE AS SELF MEDICATION IN ILIR TIMUR II DISTRICT

Abstract Display: 603 / [PDF](#) PDF Downloads: 617 / [PDF](#) PDF Downloads: 57Hairun Niza, Sarah Hanifah Az-Zahra Anwar, Wahyudi Wahyudi
57-61[VIEW ABSTRACT](#)[PDF](#)<https://doi.org/10.22270/jddt.v13i7.6149>[DOWNLOAD PDF](#)[HTML](#)

EVALUATION OF THE POTENTIAL FOR MEDICATION ERROR IN OUTPATIENT ELDERLY PATIENTS AT SITI FATIMAH AZ-ZAHRA REGIONAL HOSPITAL

Abstract Display: 661 / [PDF](#) PDF Downloads: 704 / [PDF](#) PDF Downloads: 91Hairun Niza, Sarah Hanifah Az-Zahra Anwar, Sindi Utami
62-64[VIEW ABSTRACT](#)[PDF](#)<https://doi.org/10.22270/jddt.v13i7.6153>[DOWNLOAD PDF](#)[HTML](#)

STRUCTURE-BASED DRUG DESIGN, ADME AND MOLECULAR DOCKING ANALYSES OF ANTI-VIRAL AGENTS AGAINST SARS-COV-2 VIRUS, ZIKA VIRUS AND HEPATITIS C VIRUS

Abstract Display: 711 / [PDF](#) PDF Downloads: 616 / [PDF](#) PDF Downloads: 77Peter Solo, Sangdintuile Zeliang, Muluvelu Lohe, Avünü Neikha, Akumsunep Jamir
65-74[VIEW ABSTRACT](#)[PDF](#)<https://doi.org/10.22270/jddt.v13i7.5909>[DOWNLOAD PDF](#)[HTML](#)

POTENTIAL ACTIVITY OF STEVIA REBAUDIANA BERT. IN INHIBITING CYCLOOXYGENASE AND LIPOOXYGENASE ENZYMES AS ANTI-INFLAMMATORY CANDIDATES: A MOLECULAR DOCKING STUDY AND ADMET PREDICTION

Abstract Display: 511 / [PDF](#) PDF Downloads: 651 / [PDF](#) PDF Downloads: 87Meida Adelia, Arif Setiawansyah, Rina SE Sitindaon
75-86[VIEW ABSTRACT](#)[PDF](#)<https://doi.org/10.22270/jddt.v13i7.6137>[DOWNLOAD PDF](#)[HTML](#)

PHYTOCHEMICAL ANALYSIS OF DIFFERENT EXTRACT OF HYPTIS SUAVEOLENS(L.)POIT

Abstract Display: 703 / [PDF](#) PDF Downloads: 840 / [PDF](#) PDF Downloads: 482

Saurav Gaikwad, Sonali Deore, Akanksha Adsare
87-92

 [VIEW ABSTRACT](#)

 [PDF](#)



<https://doi.org/10.22270/jddt.v13i7.5916>

 [DOWNLOAD PDF](#)

[HTML](#)

MOLECULAR DOCKING ANALYSIS OF PIPERONAL AND ITS ANALOGUES AS PROMISING CANCER THERAPEUTICS BY MODULATING ANGIOGENESIS



Abstract Display: 432 /  PDF Downloads: 672 /  PDF Downloads: 351

Anuradha Venkatraman, Ashwin Prakash Karurkar, Syed Ali Mohamed Yacoob, Sankar Muthumanickam
93-98

 [VIEW ABSTRACT](#)

 [VIEW ABSTRACT](#)

<https://doi.org/10.22270/jddt.v13i7.5917>

 [PDF](#)

 [HTML](#)

 [DOWNLOAD PDF](#)

 [DOWNLOAD HTML](#)

FORMULATION DEVELOPMENT AND EVALUATION OF NEW POLYHERBAL GEL FORMULATIONS FOR THEIR WOUND HEALING ACTIVITY IN RAT



Abstract Display: 1107 /  PDF Downloads: 789 /  PDF Downloads: 132

Yadvendra Dutt Shukla, Karunakar Shukla, Rakesh Kumar Jatav
99-106

 [VIEW ABSTRACT](#)

 [PDF](#)



<https://doi.org/10.22270/jddt.v13i7.5905>

 [DOWNLOAD PDF](#)

[HTML](#)

COMPARISON OF ANTIBIOTIC-RESISTANT PATTERN OF EXTENDED SPECTRUM BETA-LACTAMASE AND CARBAPENEM-RESISTANT ESCHERICHIA COLI ISOLATES FROM CLINICAL AND NON-CLINICAL SOURCES



Abstract Display: 703 /  PDF Downloads: 606 /  PDF Downloads: 56

Itoro Sunday Joseph, Ijeoma Onyinye Okolo, Evangeline Chinyere Udenweze, Chinenye Emelda Nwankwo, Ikemesit Udeme Peter, Ifeoma Precious Ogbonna, Ifeanyichukwu Romanus Iroha
107-118

 [VIEW ABSTRACT](#)

 [PDF](#)



<https://doi.org/10.22270/jddt.v13i7.5918>

 [DOWNLOAD PDF](#)

[HTML](#)

ISOLATION, CHARACTERIZATION AND ANTIFERTILITY ACTIVITY OF ETHANOLIC EXTRACTS OF TABERNAEMONTANA DIVARICATA R.BR. EX ROEM. & SCHULT



Abstract Display: 626 /  PDF Downloads: 645 /  PDF Downloads: 143

Ameena Khatoun, Sourabh Jain, Rajendra S. Bapna
119-126

[VIEW ABSTRACT](#)[PDF](#)<https://doi.org/10.22270/jddt.v13i7.5907>[DOWNLOAD PDF](#)[HTML](#)

FORMULATION AND EVALUATION OF MICRO BEADS OF CLONIDINE FOR THE TREATMENT OF ANXIETY AND HYPERTENSIVE DISORDER

Abstract Display: 461 / [PDF](#) PDF Downloads: 615 / [PDF](#) PDF Downloads: 225Asmita Dwivedi, Napeesha Mansoori, Mahima Dubey, Anjali Namdeo
127-133[VIEW ABSTRACT](#)[PDF](#)<https://doi.org/10.22270/jddt.v13i7.5920>[DOWNLOAD PDF](#)[HTML](#)

Review

A COMPREHENSIVE REVIEW ON EFFERVESCENT TABLETS

Abstract Display: 6406 / [PDF](#) PDF Downloads: 8171 / [PDF](#) PDF Downloads: 1523Kajal Gopinath Vanhere, Deelip Vishram Derle, Sandeep Bhausheb Khatale, Satish Laxman Nangude
141-150[VIEW ABSTRACT](#)[PDF](#)<https://doi.org/10.22270/jddt.v13i7.6120>[DOWNLOAD PDF](#)[HTML](#)

A COMPREHENSIVE OVERVIEW ON PHYTOMEDICINES AS AN UPCOMING/EMERGING CANDIDATE FOR THE MANAGEMENT OF EPILEPSY

Abstract Display: 718 / [PDF](#) PDF Downloads: 587 / [PDF](#) PDF Downloads: 91Priya Kumari, Pravin Kumar, Mahendra Singh Ashawat, Palak Sharma
151-160[VIEW ABSTRACT](#)[PDF](#)<https://doi.org/10.22270/jddt.v13i7.6125>[DOWNLOAD PDF](#)[HTML](#)

A SYSTEMATIC REVIEW ON COMPARISON OF DEEP BRAIN STIMULATION (DBS) AND TRANSCRANIAL MAGNETIC STIMULATION (TMS) IN THE TREATMENT OF REFRACTORY OBSESSIVE COMPULSIVE DISORDER (OCD)

Abstract Display: 610 / [PDF](#) PDF Downloads: 599 / [PDF](#) PDF Downloads: 221K Dhivya, Bibila Samuel
161-171[VIEW ABSTRACT](#)[PDF](#)<https://doi.org/10.22270/jddt.v13i7.6126>[DOWNLOAD PDF](#)[HTML](#)

AN DETAILED OVERVIEW ON MOUTH DISSOLVING FILM



Abstract Display: 1622 / PDF PDF Downloads: 4089 / PDF PDF Downloads: 148

Priyanshi Jain, Ashish Gupta, Gajanan Darwhekar
172-176

VIEW ABSTRACT

PDF



<https://doi.org/10.22270/jddt.v13i7.6121>

DOWNLOAD PDF

HTML

THERAPEUTIC ASPECTS OF BIOLOGICALLY POTENT VANILLIN DERIVATIVES: A CRITICAL REVIEW



Abstract Display: 1535 / PDF PDF Downloads: 1743 / PDF PDF Downloads: 2650

Raju Senthil Kumar, Sundhararajan Naveena, Sekar Praveen, Nagalingam Yogadharshini
177-189

VIEW ABSTRACT

PDF



<https://doi.org/10.22270/jddt.v13i7.6159>

DOWNLOAD PDF

HTML

REVIEW ON GREEN CHEMISTRY



Abstract Display: 1095 / PDF PDF Downloads: 1532 / PDF PDF Downloads: 2017

Tejas Wale, Shrikant Suryavanshi, Vinod Wavare, Pallavi Phalke, Manisha Sharmale
190-193

VIEW ABSTRACT

PDF



<https://doi.org/10.22270/jddt.v13i7.5919>

DOWNLOAD PDF

HTML

SACCOPE TALUM TOMENTOSUM: REVIEW OF ITS BOTANY, MEDICINAL USES, PHARMACOLOGICAL ACTIVITIES AND PHYTOCHEMISTRY



Abstract Display: 572 / PDF PDF Downloads: 538 / PDF PDF Downloads: 98

Prashant Kumar, Karunakar Shukla
194-199

VIEW ABSTRACT

PDF



<https://doi.org/10.22270/jddt.v13i7.5929>

DOWNLOAD PDF

HTML



Online ISSN: 2250-1177

[MAKE A SUBMISSION](#)

CATEGORIES

- [Aims & Scope](#)
- [Editorial Policies](#)
- [Publication Ethics](#)
- [Plagiarism Policy](#)
- [Retraction Policy](#)
- [Open Access Policy](#)
- [Processing Charges and Waiver Policy](#)
- [Indexing](#)

LATEST PUBLICATIONS

[ATOM 1.0](#)

[RSS 2.0](#)

[RSS 1.0](#)

[Announcements](#) || [Editorial Board](#) || [Indexing](#) || [Contact](#)



HOME / ARCHIVES / VOL. 13 NO. 7 (2023): VOLUME 13, ISSUE 7, JULY 2023 / Research

Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years

Meiyanti Meiyanti

Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

<https://orcid.org/0000-0002-1770-5504>

, Yohana

Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

<https://orcid.org/0009-0006-3019-8064>

Eveline Margo

Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

<https://orcid.org/0000-0002-1614-0804>

Juni Chudri

Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

, Pusparini

Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

<https://orcid.org/0000-0002-3371-5825>

Meutia Atika Faradilla

Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

<https://orcid.org/0009-0006-1216-5891>

ABSTRACT

Aging is a progressive process caused by a physiological decline in body functions. It is known that several factors can accelerate aging, namely age, lifestyle, oxidative stress, lipid peroxidation, and mitochondrial damage. Various pollutants such as cigarette smoke, motor vehicle fumes, industrial waste, and exposure to sunlight / ultraviolet light will form free radicals. Malondialdehyde (MDA) level is one of the indicators used to measure oxidative stress. Increased MDA levels are found in various degenerative diseases. This study aims to determine plasma MDA levels in the age group of 40 years and over and factors that can affect plasma MDA levels. This study was a cross-sectional study by including 42 subjects. Data collection was carried out using a questionnaire to obtain subject demographic data and

physical activity, as well as physical examinations complemented by laboratory examinations and MDA levels. The mean age of the subjects was 52.7 years, and most of the 32 (76.2%) subjects were categorized as obese and had high activity. Pearson correlation test obtained body mass index statistically significantly correlated with plasma MDA levels with a value of $p = 0.039$, as well as physical activity statistically significantly correlated with a negative correlation, with a value of $r = -0.26$ and $p = 0.043$. Conclusion Body mass index and activity are significantly correlated with plasma MDA levels.

Keywords: malondialdehyde, stress, oxidative, radicals

Keywords: malondialdehyde, stress, oxidative, radicals

DOI <https://doi.org/10.22270/jddt.v13i7.6142>

AUTHOR BIOGRAPHIES

Meiyanti Meiyanti, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

, Yohana, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Eveline Margo, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Juni Chudri, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

, Pusparini, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Meutia Atika Faradilla, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

REFERENCES

Sharifi-Rad M, Anil Kumar N V., Zucca P, Varoni EM, Dini L, Panzarini E, et al. Lifestyle, Oxidative Stress, and Antioxidants: Back and Forth in the Pathophysiology of Chronic Diseases. *Front Physiol.* 2020; 11(July):1-21. <https://doi.org/10.3389/fphys.2020.00694>

Park MH, Jin HK, Bae J sung. Potential therapeutic target for aging and age-related neurodegenerative diseases: the role of acid sphingomyelinase. *Exp Mol Med [Internet].* 2020; 52(3):380-9. <https://doi.org/10.1038/s12276-020-0399-8>

Całyniuk B, Grochowska-Niedworok E, Walkiewicz K, Kawecka S, Popiołek E, Fatyga E. Malondialdehyde (MDA) - product of lipid peroxidation as marker of homeostasis disorders and aging. *Ann Acad Medicae Silesiensis.* 2016; 70:224-8. <https://doi.org/10.18794/aams/65697>

Savira M, Rusdiana ., Widjaja SS, Syahputra M. Comparison Malondialdehyde (MDA) Level between Obesity Non Metabolic Syndrome and Obesity with Metabolic Syndrome Patients. 2020; (5):644-7.

Yekti R, Bukhari A, Jafar N, Thaha AR. Measurement of malondialdehyde (MDA) as a good indicator of lipid peroxidation. *Int J Allied Med Sci Clin [Internet]*. 2018; 6(4):6-8. Available from: www.ijamscr.com

Denise Mafra ME. Influence of Physical Exercise on Oxidative Stress and Inflammation in Hemodialysis Patients. *J Mol Biomark Diagn*. 2015; 06(03). <https://doi.org/10.4172/2155-9929.1000e120>

Mao C, Yuan JQ, Lv Y Bin, Gao X, Yin ZX, Kraus VB, et al. Associations between superoxide dismutase, malondialdehyde and all-cause mortality in older adults: A community-based cohort study. *BMC Geriatr*. 2019; 19(1):1-9. <https://doi.org/10.1186/s12877-019-1109-z>

Ayala A, Muñoz MF, Argüelles S. Lipid peroxidation: Production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. *Oxid Med Cell Longev*. 2014; 2014. <https://doi.org/10.1155/2014/360438>

Situmorang N, Zulham Z. Malondialdehyde (Mda) (Zat Oksidan Yang Mempercepat Proses Penuaan). *J Keperawatan Dan Fisioter*. 2020; 2(2):117-23. <https://doi.org/10.35451/jkf.v2i2.338>

Hardiany NS, Sucitra S, Paramita R. Profile of malondialdehyde (MDA) and catalase specific activity in plasma of elderly woman. *Heal Sci J Indones*. 2020; 10(2):132-6. <https://doi.org/10.22435/hsji.v12i2.2239>

Hardiany NS, Sadikin M, Siregar N, Wanandi SI. The suppression of manganese superoxide dismutase decreased the survival of human glioblastoma multiforme T98G cells. *Med J Indones*. 2017; 26(1):19-25. <https://doi.org/10.13181/mji.v26i1.1511>

439-445_nova.pdf.

A. Hassan A, Sayyah SG. Oxidative Stress Marker Malondialdehyde and Glutathione Antioxidant in Hypertensive Patients. *Eur J Biomed Res*. 2023; 2(1):31-6. <https://doi.org/10.24018/ejbiomed.2023.2.1.47>

Khoubnasab Jafari M, Ansarin K, Jouyban A. Comments on "use of malondialdehyde as a biomarker for assesing oxidative stress in different disease pathologies: A review." *Iran J Public Health*. 2015; 44(5):714-5.

Shalash M, Badra M, Imbaby S, ElBanna E. Malondialdehyde in Type 2 Diabetics and Association With Cardiovascular Risk Factors. *J Med Res Inst*. 2020; 41(2):21-30. <https://doi.org/10.21608/jmalexu.2020.147116>

Oesterle A, Laufs U, Liao JK. Pleiotropic Effects of Statins on the Cardiovascular System. *Circ Res*. 2017; 120(1):229-43. <https://doi.org/10.1161/CIRCRESAHA.116.308537>

Sekunda KN, Ariosta A, Limantoro C, Setiawan AA. Difference in Profiles of Oxidative Stress Marker (MDA) in STEMI and NSTEMI. *Diponegoro Med J (Jurnal Kedokt Diponegoro)*. 2021; 10(2):145-50. <https://doi.org/10.14710/dmj.v10i2.29673>

Block G, Dietrich M, Norkus EP, Morrow JD, Hudes M, Caan B, et al. Factors associated with oxidative stress in human populations. *Am J Epidemiol*. 2002; 156(3):274-85. <https://doi.org/10.1093/aje/kwf029>

Kubihal C V., Naik HD. Effect of smoking on vitamin C and MDA: a cross sectional comparative study. *Int J Res Med Sci*. 2019; 7(3):746. <https://doi.org/10.18203/2320-6012.ijrms20190511>

Ari Suwondo, Ahmad Mohan, Melyana Nurul Widyawati M. Factors Related to Malondialdehyde (MDA) Levels in Metal Casting Industry Workers. *Preprints*. 2020; 1(October):1-8. <https://doi.org/10.21203/rs.3.rs-99597/v1>

Schöttker B, Brenner H, Jansen EHJM, Gardiner J, Peasey A, Kubínová R, et al. Evidence for the free radical/oxidative stress theory of ageing from the CHANCES consortium: A meta-analysis of individual participant data. *BMC Med*. 2015; 13(1):1-15. <https://doi.org/10.1186/s12916-015-0537-7>

Kander MC, Cui Y, Liu Z. Gender difference in oxidative stress: a new look at the mechanisms for cardiovascular diseases. *J Cell Mol Med.* 2017; 21(5):1024-32. <https://doi.org/10.1111/jcmm.13038>

Bell JR, Bernasochi GB, Varma U, Raaijmakers AJA, Delbridge LMD. Sex and sex hormones in cardiac stress - Mechanistic insights. *J Steroid Biochem Mol Biol.* 2013; 137:124-35. <https://doi.org/10.1016/j.jsbmb.2013.05.015>

P G, Curcio F, Bulli G, Aran L, Della-morte D, Testa G, et al. Oxidative Stress and Diseases. *Oxidative Stress Dis.* 2012; 757-72.

Sabitha K, Venugopal B, Rafi M, V Ramana K. Role of Antioxidant Enzymes in Glucose and Lipid Metabolism in Association with Obesity and Type 2 Diabetes. *Am J Med Sci Med.* 2014; 2(1):21-4. <https://doi.org/10.12691/ajmsm-2-1-5>

Manna P, Jain SK. Obesity, Oxidative Stress, Adipose Tissue Dysfunction, and the Associated Health Risks: Causes and Therapeutic Strategies. *Metab Syndr Relat Disord.* 2015; 13(10):423-44. <https://doi.org/10.1089/met.2015.0095>

Lewis NA, Howatson G, Morton K, Hill J, Pedlar CR. Alterations in Redox Homeostasis in the Elite Endurance Athlete. *Sport Med.* 2015; 45(3):379-409. <https://doi.org/10.1007/s40279-014-0276-5>

 **VIEW ABSTRACT**

 **PDF**

 **DOWNLOAD PDF**

 **HTML**

PUBLISHED

2023-07-15

STATISTICS



Abstract Display: 1069



PDF Downloads: 840



PDF Downloads: 62

HOW TO CITE

Meiyanti M, Yohana , Margo E, Chudri J, Pusparini , Faradilla MA. Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years. *J. Drug Delivery Ther.* [Internet]. 2023 Jul. 15 [cited 2026 Jun. 17];13(7):52-6. Available from: <https://jddtonline.info/index.php/jddt/article/view/6142>

More Citation Formats

ISSUE

[Vol. 13 No. 7 \(2023\): Volume 13, Issue 7, July 2023](#)

SECTION

Research

Authors who publish with this journal agree to the following terms:

- Authors retain copyright and grant the journal right of first publication with the work simultaneously licensed under a [Creative Commons Attribution-NonCommercial 4.0 International \(CC BY-NC 4.0\)](#), that allows others to share the work with an acknowledgment of the work's authorship and initial publication in this journal.
- Authors are able to enter into separate, additional contractual arrangements for the non-exclusive distribution of the journal's published version of the work (e.g., post it to an institutional repository or publish it in a book), with an acknowledgment of its initial publication in this journal.
- Authors are permitted and encouraged to post their work online (e.g., in institutional repositories or on their website) prior to and during the submission process, as it can lead to productive exchanges, as well as earlier and greater citation of published work (See [The Effect of Open Access](#)).

HOW TO CITE

Meiyanti M, Yohana , Margo E, Chudri J, Pusparini , Faradilla MA. Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years. J. Drug Delivery Ther. [Internet]. 2023 Jul. 15 [cited 2026 Jun. 17];13(7):52-6. Available from: <https://jddtonline.info/index.php/jddt/article/view/6142>

MORE CITATION FORMATS



Crossref
Cited-by

0

Scopus[®]

0



Europe
PMC

0

Most read articles by the same author(s)

- Meiyanti Meiyanti, Lyfanni Fiesa Mulia, Mulia, Pusparini, [The Relationship between Antipsychotic Drug Use and Liver Enzyme Levels in People with Mental Disorders](#) , [Journal of Drug Delivery and Therapeutics: Vol. 12 No. 1 \(2022\): Volume 12, Issue 1, Jan-Feb 2022](#)
- Meiyanti Meiyanti, Eveline Margo, Monica Dwi Hartanti, Juni Chudri, Lie Tanu Merijanti, Arleen Devita, [Correlation of Three Anthropometric Parameters with the Percentage of Body Fat and Visceral Fat in the Productive Age Group](#) , [Journal of Drug Delivery and Therapeutics: Vol. 12 No. 2 \(2022\): Volume 12, Issue 2 March-April 2022](#)
- Eveline Margo, , Meiyanti, Fransisca Chondro, Erlani Kartadinata, , Yohana, [Relationship Blood Pressure and Exercise with Vitamin D Levels in Employees](#) , [Journal of Drug Delivery and Therapeutics: Vol. 12 No. 5-S \(2022\): Volume 12, Issue 5-S, Sep-Oct 2022](#)
- Lie Tanu Merijanti, Pusparini Pusparini, Meiyanti Meiyanti, Alvina Alvina, Monica Dwi Hartanti, Muljadi Tjahjadi, [Career Development and Psychopathological Symptoms on Female Workers](#) , [Journal of Drug Delivery and Therapeutics: Vol. 13 No. 4 \(2023\): Volume 13, Issue 4, April 2023](#)
- Yohana Yohana , Meiyanti Meiyanti , Deasyka Yastani , Zakiaturahhmi Zakiaturahhmi , Suweino Tjhia, [Analyzing FOXO3A gene as a marker in adults with hypertension](#) , [Journal of Drug Delivery and Therapeutics: Vol. 16 No. 4 \(2026\): Volume 16, Issue 4, April 2026](#)



Online ISSN: 2250-1177

MAKE A SUBMISSION

CATEGORIES

Aims & Scope

Editorial Policies

Publication Ethics

[Plagiarism Policy](#)

[Retraction Policy](#)

[Open Access Policy](#)

[Processing Charges and Waiver Policy](#)

[Indexing](#)

LATEST PUBLICATIONS

ATOM 1.0

RSS 2.0

RSS 1.0

[Announcements](#) || [Editorial Board](#) || [Indexing](#) || [Contact](#)

The publication is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](#) (CC BY-NC 4.0). [View Legal Code](#)

Parse error: Unclosed '(' on line 53325 in `/home/jddtonline/domains/jddtonline.info/public_html/cache/fc-geolP-all.php` on line **53328**

Available online on 15.07.2023 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

Copyright © 2023 The Author(s). This is an open-access article distributed under the terms of the CC BY-NC 4.0 which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited



Open Access Full Text Article



Check for updates

Research Article

Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years

Meiyanti* , Yohana , Eveline Margo , Juni Chudri, Pusparini , Meutia Atika Faradilla

Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Article Info:

Abstract



Article History:

Received 08 May 2023
Reviewed 11 June 2023
Accepted 24 June 2023
Published 15 July 2023

Cite this article as:

Meiyanti, Yohana, Margo E, Chudri J, Pusparini, Faradilla MA, Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years, *Journal of Drug Delivery and Therapeutics*. 2023; 13(7):52-56

DOI: <http://dx.doi.org/10.22270/jddt.v13i7.6142>

*Address for Correspondence:

Meiyanti, Department of Pharmacology and Medical Pharmacy, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Aging is a progressive process caused by a physiological decline in body functions. It is known that several factors can accelerate aging, namely age, lifestyle, oxidative stress, lipid peroxidation, and mitochondrial damage. Various pollutants such as cigarette smoke, motor vehicle fumes, industrial waste, and exposure to sunlight / ultraviolet light will form free radicals. Malondialdehyde (MDA) level is one of the indicators used to measure oxidative stress. Increased MDA levels are found in

various degenerative diseases. This study aims to determine plasma MDA levels in the age group of 40 years and over and factors that can affect plasma MDA levels. This study was a cross-sectional study by including 42 subjects. Data collection was carried out using a questionnaire to obtain subject demographic data and physical activity, as well as physical examinations complemented by laboratory examinations and MDA levels. The mean age of the subjects was 52.7 years, and most of the 32 (76.2%) subjects were categorized as obese and had high activity. Pearson correlation test obtained body mass index statistically significantly correlated with plasma MDA levels with a value of $p = 0.039$, as well as physical activity statistically significantly correlated with a negative correlation, with a value of $r = -0.26$ and $p = 0.043$. Conclusion Body mass index and activity are significantly correlated with plasma MDA levels.

Keywords: malondialdehyde, stress, oxidative, radicals

INTRODUCTION

Aging is a progressive process caused by a physiological decline in body function. ¹ A decrease in body homeostasis is accompanied by an increased risk of various chronic diseases including diabetes mellitus, cancer, neurodegenerative diseases, and heart disease. It is known that several factors can accelerate aging, namely age, lifestyle, oxidative stress, lipid peroxidation, and mitochondrial damage. Previous studies have shown that changes in sphingomyelin metabolism in the central nervous system and age are associated with neurodegenerative diseases. ² In the US, aging is one of the most important factors in diabetes mellitus and heart disease. In 2015, an estimated 12% of the population had diabetes mellitus and an undetected 24% had diabetes mellitus. Heart disease is the highest contributor to mortality in the population aged 65-84 years. ³ Recent studies have shown that non-diabetic populations whose fasting glucose levels increase in the blood will increase the risk of coronary heart disease. But on the other hand, the data on fasting blood glucose increase is still debated in the population of cardiovascular disease patients who do not have diabetes mellitus. A recent study in North Sumatra showed a significant relationship between MDA levels in obese and non-obese populations. ⁴

Free radicals are formed through two processes, namely

exogenous and endogenous processes. Various pollutants such as cigarette smoke, motor vehicle fumes, industrial waste, and exposure to sunlight/ultraviolet will form free radicals. Illness, stress, and excessive exercise/physical activity are endogenous factors produced by the human body that can form free radicals. The formation of free radicals occurs due to oxidation processes and metabolic processes that take place during exercise or activity, inflammatory processes, consumption of certain foods, and others. Free radicals will begin to activate with molecules in the surrounding cells to obtain electrons to stabilize the condition of one's body. The electrons that have been taken in the cell molecules will trigger the occurrence of free radicals. If this continues for a long time, it can cause oxidative stress which can cause inflammation, and DNA disease can also trigger the disease. Oxidative stress conditions characterized by increased production of Reactive Oxygen Species (ROS) have implications for various diseases such as hypertension, atherosclerosis, diabetes mellitus, stroke, chronic renal failure, heart failure, and other chronic diseases. ⁵⁻⁷

Oxidative stress is an imbalance between oxidants (free radicals) and antioxidants. Free radicals that commonly cause oxidative damage are reactive oxygen species (ROS).⁸ ROS compounds are formed through aerobic metabolism. These compounds can cause extensive damage through catalytic reactions. They are responsible for damage at the cellular level

by oxidizing proteins, lipids, DNA (Deoxyribonucleic acid), and other biomolecules. Various compounds can be used as markers of oxidative stress such as malondialdehyde (MDA), and protein carbonylation. 4-hydroxy-2-nonenal (4HNE), and 8-hydroxy-deoxyguanosine (8OHdG).⁹

Malondialdehyde (MDA) is a result of lipid peroxidation in the body. The concentration of MDA indicates an oxidation process that occurs in the body's cell membranes due to metabolic processes. One of the factors that cause aging is the production of excess reactive oxygen species (ROS). ROS are generated through the process of cell respiration in the mitochondria and the phagocytosis process of foreign body elimination. To stabilize the amount of ROS, the body can use endogenous and exogenous antioxidants. Disruption of the balance between ROS and antioxidants can cause irreversible damage to cell organelle components up to the DNA level.^{10,11}

MDA levels are strongly influenced by many factors such as age, illness, stress, various pollutants, food consumption, antioxidant supplements and excessive physical activity or exercise, and others. The other study found that more than 30% of subjects had high MDA levels, and concluded that there was a significant relationship between eating habits, body mass index, glucose levels with MDA levels, as well as smoking habits significantly related to MDA levels.¹² Recent studies have shown that plasma MDA levels in populations aged 70 years and 60 years are not significantly different but there is an increase in the specific activity of the catalase enzyme that increases with age.¹⁰ In another study, there was a significant difference in plasma MDA levels in the obesity group without metabolic syndrome compared to the obesity group with metabolic syndrome.⁴

There are still variations in MDA data in various groups with different ages and different disease groups, so this study aims to determine plasma MDA levels in the age group 40 years and over and factors that can affect plasma MDA levels.

METHOD

This observational analytic study with a cross-sectional design approach was conducted from November 2022 to February 2023 in Angke village, West Jakarta, Indonesia. The inclusion criteria for this study were: men and women aged ≥ 40 years, willing to participate in the study and sign informed consent. Meanwhile, the exclusion criteria were having a history of malignancy, kidney failure, a history of autoimmune disease, taking drugs related to psychiatric conditions, and having limitations in mobility. This study used the calculation of sample size with the formula $N / 1 + N.\alpha^2$, and obtained the minimum number of samples required was 40 subjects. Demographic data and physical activity were obtained from distributed questionnaires. Blood chemistry examination (complete blood, glucose, total cholesterol, liver enzymes (SGOT and SGPT) using an enzymatic method. Examination of malondialdehyde levels was carried out using the spectrophotometric method. This research has obtained approval from the ethics committee of the Faculty of Medicine, Trisakti University, Number: 178/KER/FK/1X/2022.

RESULTS

A total of 42 subjects participated in this study, most of the subjects were female 27 (64.3%), and the mean age of the subjects was 52.7 years. Anthropometric data of the subjects are as follows body weight 67.53 ± 13.64 and height 156.02 ± 8.73 cm, while for abdominal circumference the average is 93.09 cm. Most of the subjects 59.5% fell into the normotensive category, with an average systolic blood pressure of 141.62 mmHg and an average diastolic blood pressure of 82.45 mmHg. A total of 32 (76.2%) subjects were

categorized as obese with BMI ≥ 25.0 , and most subjects had high activity levels. (Table 1)

Table 1: Distribution of Subject Characteristics (n=42)

Characteristics	n (%)	Mean \pm SD
Gender		
Male	15 (35.7)	
Female	27 (64.3)	
Blood pressure (mmHg)		
Normal (< 140/90)	25 (59.5)	
Hypertension ($\geq 140/90$)	17 (40.5)	
Physical Activity		
Low (< 600 Mets)	11 (26.2)	
High (≥ 600 Mets)	31 (73.8)	
Body Mass Index (kg/m ²)		27.69 \pm 4.90
Not obese (< 25.0)	10 (23.8)	
Obese (≥ 25.0)	32 (76.2)	
Age (years)		52.7 \pm 10.60
Body weight (kg)		67.53 \pm 13.64
Height (cm)		156.02 \pm 8.73
Abdominal circumference (cm)		93.09 \pm 10.54
Blood pressure (mmHg)		
Systole		141.62 \pm 20.77
Diastole		82.45 \pm 12.70

The examination of the subject's plasma malondialdehyde levels obtained a mean of 0.77 with the highest value of 1.03 and the lowest of 0.51. Hemoglobin levels averaged 13.16 g/dL. Fasting blood glucose level was 95.06 ± 41.01 mg/dL and total cholesterol was 179.83 ± 31.85 mg/dL. Laboratory examinations to determine liver function were carried out by measuring SGPT and SGOT levels, obtaining mean values of 24.21 U/L and 26.31 U/L respectively. (Table 2)

Table 2: Laboratory examination results of subjects (n=42)

Variable	Mean \pm SD
Hemoglobin (g/dL)	13.16 \pm 1.56
Fasting blood glucose (mg/dL)	95.06 \pm 41.01
Total cholesterol (mg/dL)	179.83 \pm 31.85
SGPT (U/L)	24.21 \pm 13.50
SGOT (U/L)	26.31 \pm 11.51
Plasma malondialdehyde	0.77 \pm 0.26

Determining the type of correlation test to be used by first conducting a data normality test. The normality test used was Shapiro-Wilk because the number of subjects was less than 50 subjects. The correlation test used to assess the correlation between blood glucose, total cholesterol, SGPT, SGOT levels, age, body mass index and physical activity with plasma MDA levels was the Pearson correlation test. The test results are presented in Table 3. Blood glucose, total cholesterol, SGPT,

SGOT, and age did not have a significant correlation with MDA levels. Body mass index has a significant correlation with plasma MDA ($p = 0.039$), with a positive correlation direction, and the strength of the correlation is statistically weak with a value of $r = 0.396$ (<0.4). The physical activity variable has a significant correlation with plasma MDA levels ($p=0.043$) with a negative correlation direction but with a statistically weak correlation strength ($r=-0.26$).

Table 3: Correlation Test of Age, Gender, Glucose, SGOT, SGPT, BMI, Physical Activity with Plasma MDA Levels (n=42)

	Plasma MDA Level	
	r	p
Age	0.098	0.756
Gender	0.124	0.108
Glucose	0.330	0.835
SGOT	0.104	0.513
SGPT	0.008	0.614
Body Mass Index	0.396	0.039*
Physical activity	-0.260	0.043*

* $p < 0.05$

DISCUSSION

The level of oxidative stress can be measured using Malondialdehyde (MDA) concentration. MDA is the result of lipid peroxidation, which will cause cell damage. Oxidant compounds such as free radicals or non-radical species attack lipids that contain carbon-carbon double bonds, especially unsaturated fatty acids (PUFA). MDA is one of the biomarkers widely used to assess oxidative stress. The mean serum MDA in this study 0.77 was found to be lower than the previous study¹⁰, this is because the age difference of the subjects in this study was younger than in the previous study. In contrast, a study conducted in Iraq on hypertensive patients concluded that the increase in MDA was associated with an increase in age, gender, and duration of disease.¹³

Free radicals such as ROS are the etiology of most chronic diseases, as it is known that active free radicals are the result of metabolic reactions. Damage to body tissue cells occurs because active free electrons remain looking for other electrons to settle, if no other electrons are available then active free electrons will attack cells resulting in cell damage. Although free radicals naturally have benefits for the body, when the level of free radicals exceeds the normal value, it will cause harm to the body.^{11,12}

Measurement of oxidative stress can be done by measuring plasma malondialdehyde (MDA) levels. MDA is a secondary result of the lipid peroxidation process, producing various active compounds that will damage cells. The process of MDA formation through enzymatic and non-enzymatic processes. Currently, MDA is one of the biomarkers used to assess oxidative stress.¹⁴ Some previous researchers concluded that higher serum MDA was found in patients with hyperlipidemia. The condition of increased fat content in the body is in line with increased oxidative stress. Hyperlipidemia is associated with physical changes in cell membranes that facilitate the release of free radicals from the mitochondrial electron transport chain or activation of NADPH oxidase. In patients with diabetes mellitus (DM) there is also an increase in MDA levels, malondialdehyde interacts irreversibly and reversibly with proteins and phospholipids to have a profound effect, especially on the collagen of the cardiovascular system. MDA levels remain within normal limits due to the antioxidant

effect, but in DM, this protective effect is impaired resulting in increased MDA levels.^{1,15}

Increased lipid peroxidation is considered a consequence of oxidative stress that occurs when the dynamic balance between prooxidants and antioxidants is disrupted. The statin class of drugs for hyperlipidemia therapy is proven to be anti-atherogenic, can improve endothelial function, inhibit vascular inflammation, and stabilize atherosclerotic plaques. In addition to statins, non-steroidal anti-inflammatory drugs (NSAIDs) can affect MDA concentration. Previous studies have found that NSAIDs have significant antioxidant properties to scavenge free radicals.^{16,17}

In this study, there was no significant correlation between age and MDA levels. Similar results were also obtained by other researchers, cell damage due to oxidative stress is not related to increasing age but is associated with habits, behavior patterns, and physiological changes due to age. Various habitual patterns such as smoking, fatty food habits, lack of activity, and inflammatory processes can lead to increased oxidative stress. The habit of taking vitamin C (ascorbic acid) has a beneficial impact due to its antioxidant effect so cell damage and DNA damage are lower in this group.¹⁸ MDA levels were significantly higher among all levels of smokers. MDA levels were lower in light smokers compared to heavy or moderate smokers. Higher vitamin C levels were found in the non-smoker group. All body tissues are susceptible to free radical damage, but based on location, the respiratory epithelium is particularly vulnerable to cigarette smoke. Oxygen radicals generated close to the cell membrane oxidize membrane phospholipids which will result in lipid peroxidation with the end products of this reaction being malondialdehyde, ethane, and pentane. this causes an increase in MDA levels in smokers.¹⁹ Another study conducted in Klaten on foundry workers concluded that type of work, body mass index and smoking habits did not affect MDA levels, but length of work and marital status were associated with MDA levels. Long and continuous exposure to metallic materials will result in the accumulation of these materials in the body and will increase the oxidative process of polyunsaturated fats in cell membranes and worsen the level of lipid peroxidase, characterized by an increase in MDA levels as a reflection of the intensity and rate of lipid peroxidase reactions of damage to body cells.²⁰

Hardiany's study also concluded that there was no significant difference in plasma MDA levels in the age population, but increasing age along with an increase in catalase-specific activity.¹⁰ This increase in catalase-specific activity is a protective effect especially for elderly women to overcome high levels of oxidative damage. The role of endogenous antioxidant catalase is to suppress oxidative damage.^{18,21} Oxidative stress plays an important role in the incidence of cardiovascular diseases including coronary artery disease. Oxidative stress is considered an important mechanism for the development of cardiovascular diseases especially atherosclerosis. Hypertension, cardiac hypertrophy, and ischemia-reperfusion injury are three other major cardiovascular conditions for which gender is a risk factor. Gender is also associated with differences in oxidative stress. Under physiological conditions, women appear to be less susceptible to oxidative stress. This may be due to the antioxidant properties of estrogen, sex differences in NADPH-oxidase activity or other mechanisms yet to be determined. Sex hormones including estrogen are globally present with their receptors in the heart so sex hormones can exert effects on the cardiovascular system. Estrogen can improve heart function and reduce injury to the heart. In addition, estrogen has antioxidant effects so that it can function as a cardioprotective.^{22,23}

This study found a significant correlation between body mass index and MDA levels, the higher the BMI value, the higher the MDA levels. Similar results were also obtained in a study in India that concluded that in metabolic syndrome patients with one of the signs is an increase in BMI, there is an increase in MDA levels compared to the control group. In obese subjects, the level of MDA is higher than normal IMT. In a healthy body, there is redox hemostasis that occurs in cells, if an imbalance arises it will cause oxidative stress. Obese patients will have an increased metabolic load and mechanical load on the myocardium, increased body mass, and excessive food intake will cause the formation of lipid peroxidation, and free radicals.²⁴ The increase in some of these things will stimulate antioxidant enzymes in the long run, this will cause antioxidant enzyme reserves to be depleted and unable to overcome and balance the increase in oxidative stress.^{25,26}

These types of training activities are exhausting, long exercise, overtraining syndrome, various types of exercise will induce a significant response to oxidative stress. Exercise activities such as light exercise, low-intensity exercise, and prolonged training will have an impact on improving endogenous antioxidant status. Reactive oxygen species play an important role in cell signaling and in regulating antioxidant gene expression. Exertion results in hyperregulation of nuclear factor kappa B and mitogen-activated protein kinases that activate gene expression of a number of enzymes and proteins with important roles in maintaining intracellular oxidative or antioxidant homeostasis.^{1,6,27} The limitation of this study is that some factors that may affect MDA levels such as consumption of vitamin supplements, diet, and length of illness suffered were not examined in this study.

CONCLUSIONS

Body mass index and activity were significantly correlated with MDA levels, while age, gender, blood glucose levels, and liver enzymes were not significantly correlated with MDA levels.

Conflict of Interest:

The investigators declare there is no conflict of interest in this study.

Acknowledgments

The authors thank all the subjects who participated in this study, as well as the laboratory staff who assisted in the implementation of this study.

REFERENCES

- Sharifi-Rad M, Anil Kumar N V., Zucca P, Varoni EM, Dini L, Panzarini E, et al. Lifestyle, Oxidative Stress, and Antioxidants: Back and Forth in the Pathophysiology of Chronic Diseases. *Front Physiol.* 2020; 11(July):1-21. <https://doi.org/10.3389/fphys.2020.00694>
- Park MH, Jin HK, Bae J sung. Potential therapeutic target for aging and age-related neurodegenerative diseases: the role of acid sphingomyelinase. *Exp Mol Med [Internet].* 2020; 52(3):380-9. <https://doi.org/10.1038/s12276-020-0399-8>
- Całyniuk B, Grochowska-Niedworok E, Walkiewicz K, Kawecka S, Popiołek E, Fatyga E. Malondialdehyde (MDA) - product of lipid peroxidation as marker of homeostasis disorders and aging. *Ann Acad Medicae Silesiensis.* 2016; 70:224-8. <https://doi.org/10.18794/aams/65697>
- Savira M, Rusdiana ., Widjaja SS, Syahputra M. Comparison Malondialdehyde (MDA) Level between Obesity Non Metabolic Syndrome and Obesity with Metabolic Syndrome Patients. 2020; (5):644-7.
- Yekti R, Bukhari A, Jafar N, Thaha AR. Measurement of malondialdehyde (MDA) as a good indicator of lipid peroxidation.

- Int J Allied Med Sci Clin [Internet]. 2018; 6(4):6-8. Available from: www.ijamscr.com
- Denise Mafra ME. Influence of Physical Exercise on Oxidative Stress and Inflammation in Hemodialysis Patients. *J Mol Biomark Diagn.* 2015; 06(03). <https://doi.org/10.4172/2155-9929.1000e120>
 - Mao C, Yuan JQ, Lv Y Bin, Gao X, Yin ZX, Kraus VB, et al. Associations between superoxide dismutase, malondialdehyde and all-cause mortality in older adults: A community-based cohort study. *BMC Geriatr.* 2019; 19(1):1-9. <https://doi.org/10.1186/s12877-019-1109-z>
 - Ayala A, Muñoz MF, Argüelles S. Lipid peroxidation: Production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. *Oxid Med Cell Longev.* 2014; 2014. <https://doi.org/10.1155/2014/360438>
 - Situmorang N, Zulham Z. Malondialdehyde (Mda) (Zat Oksidan Yang Mempercepat Proses Penuaan). *J Keperawatan Dan Fisioter.* 2020; 2(2):117-23. <https://doi.org/10.35451/jkf.v2i2.338>
 - Hardiany NS, Sucitra S, Paramita R. Profile of malondialdehyde (MDA) and catalase specific activity in plasma of elderly woman. *Heal Sci J Indones.* 2020; 10(2):132-6. <https://doi.org/10.22435/hsji.v12i2.2239>
 - Hardiany NS, Sadikin M, Siregar N, Wanandi SI. The suppression of manganese superoxide dismutase decreased the survival of human glioblastoma multiforme T98G cells. *Med J Indones.* 2017; 26(1):19-25. <https://doi.org/10.13181/mji.v26i1.1511>
 - 439-445_nova.pdf.
 - A. Hassan A, Sayyah SG. Oxidative Stress Marker Malondialdehyde and Glutathione Antioxidant in Hypertensive Patients. *Eur J Biomed Res.* 2023; 2(1):31-6. <https://doi.org/10.24018/ejbiomed.2023.2.1.47>
 - Khoubnasab Jafari M, Ansarin K, Jouyban A. Comments on "use of malondialdehyde as a biomarker for assessing oxidative stress in different disease pathologies: A review." *Iran J Public Health.* 2015; 44(5):714-5.
 - Shalash M, Badra M, Imbaby S, ElBanna E. Malondialdehyde in Type 2 Diabetics and Association With Cardiovascular Risk Factors. *J Med Res Inst.* 2020; 41(2):21-30. <https://doi.org/10.21608/jmalexu.2020.147116>
 - Oesterle A, Laufs U, Liao JK. Pleiotropic Effects of Statins on the Cardiovascular System. *Circ Res.* 2017; 120(1):229-43. <https://doi.org/10.1161/CIRCRESAHA.116.308537>
 - Sekunda KN, Ariosta A, Limantoro C, Setiawan AA. Difference in Profiles of Oxidative Stress Marker (MDA) in STEMI and NSTEMI. *Diponegoro Med J (Jurnal Kedokt Diponegoro).* 2021; 10(2):145-50. <https://doi.org/10.14710/dmj.v10i2.29673>
 - Block G, Dietrich M, Norkus EP, Morrow JD, Hudes M, Caan B, et al. Factors associated with oxidative stress in human populations. *Am J Epidemiol.* 2002; 156(3):274-85. <https://doi.org/10.1093/aje/kwf029>
 - Kubihal C V., Naik HD. Effect of smoking on vitamin C and MDA: a cross sectional comparative study. *Int J Res Med Sci.* 2019; 7(3):746. <https://doi.org/10.18203/2320-6012.ijrms20190511>
 - Ari Suwondo, Ahmad Mohan, Melyana Nurul Widyawati M. Factors Related to Malondialdehyde (MDA) Levels in Metal Casting Industry Workers. *Preprints.* 2020; 1(October):1-8. <https://doi.org/10.21203/rs.3.rs-99597/v1>
 - Schöttker B, Brenner H, Jansen EHJM, Gardiner J, Peasey A, Kubínová R, et al. Evidence for the free radical/oxidative stress theory of ageing from the CHANCES consortium: A meta-analysis of individual participant data. *BMC Med.* 2015; 13(1):1-15. <https://doi.org/10.1186/s12916-015-0537-7>
 - Kander MC, Cui Y, Liu Z. Gender difference in oxidative stress: a new look at the mechanisms for cardiovascular diseases. *J Cell Mol Med.* 2017; 21(5):1024-32. <https://doi.org/10.1111/jcmm.13038>
 - Bell JR, Bernasochi GB, Varma U, Raaijmakers AJA, Delbridge LMD. Sex and sex hormones in cardiac stress - Mechanistic insights. J

- Steroid Biochem Mol Biol. 2013; 137:124-35.
<https://doi.org/10.1016/j.jsbmb.2013.05.015>
24. P G, Curcio F, Bulli G, Aran L, Della-morte D, Testa G, et al. Oxidative Stress and Diseases. *Oxidative Stress Dis.* 2012; 757-72.
25. Sabitha K, Venugopal B, Rafi M, V Ramana K. Role of Antioxidant Enzymes in Glucose and Lipid Metabolism in Association with Obesity and Type 2 Diabetes. *Am J Med Sci Med.* 2014; 2(1):21-4.
<https://doi.org/10.12691/ajmsm-2-1-5>
26. Manna P, Jain SK. Obesity, Oxidative Stress, Adipose Tissue Dysfunction, and the Associated Health Risks: Causes and Therapeutic Strategies. *Metab Syndr Relat Disord.* 2015; 13(10):423-44. <https://doi.org/10.1089/met.2015.0095>
27. Lewis NA, Howatson G, Morton K, Hill J, Pedlar CR. Alterations in Redox Homeostasis in the Elite Endurance Athlete. *Sport Med.* 2015; 45(3):379-409. <https://doi.org/10.1007/s40279-014-0276-5>

library FK

Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years

Artikel Jurnal 1

Document Details

Submission ID

trn:oid::3618:143751815

Submission Date

Jun 21, 2026, 12:24 PM GMT+7

Download Date

Jun 21, 2026, 12:54 PM GMT+7

File Name

artikel_factors-associated-with-plasma-malondialdehyde-levels-in-people-over-40-years.pdf

File Size

289.0 KB

7 Pages

4,285 Words

22,824 Characters

19% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

Filtered from the Report

- ▶ Bibliography
- ▶ Quoted Text
- ▶ Small Matches (less than 8 words)

Exclusions

- ▶ 1 Excluded Source
- ▶ 11 Excluded Matches

Match Groups

- 45 Not Cited or Quoted 19%**
Matches with neither in-text citation nor quotation marks
- 0 Missing Quotations 0%**
Matches that are still very similar to source material
- 0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
- 0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 15% Internet sources
- 9% Publications
- 11% Submitted works (Student Papers)

Integrity Flags

0 Integrity Flags for Review

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups

- **45 Not Cited or Quoted 19%**
Matches with neither in-text citation nor quotation marks
- **0 Missing Quotations 0%**
Matches that are still very similar to source material
- **0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
- **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 15% Internet sources
- 9% Publications
- 11% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Internet		
		ejournal3.undip.ac.id	2%
2	Internet		
		www.jddtonline.info	2%
3	Internet		
		onlinelibrary.wiley.com	2%
4	Internet		
		pmc.ncbi.nlm.nih.gov	2%
5	Student papers		
		Trisakti University on 2023-03-29	1%
6	Internet		
		innovpub.org	1%
7	Internet		
		www.azerbaijanmedicaljournal.net	<1%
8	Internet		
		libir.josai.ac.jp	<1%
9	Publication		
		M. Shalash, M. Badra, S. Imbaby, E. ElBanna. "MALONDIALDEHYDE IN TYPE 2 DIAB..."	<1%
10	Student papers		
		Universitas Indonesia on 2025-03-11	<1%

11	Internet	www.pubmedcentral.nih.gov	<1%
12	Internet	jbiomedkes.org	<1%
13	Internet	www.nature.com	<1%
14	Publication	Jancey, Jonine M., Sarah McGann, Robyn Creagh, Krysten D. Blackford, Peter How...	<1%
15	Student papers	Tikrit University on 2016-10-07	<1%
16	Student papers	Mahidol University on 2010-03-23	<1%
17	Publication	Shaikh Ajj Ahmed Makbul, Shamim Irshad Azmi, Mohd. Sayeed Akhtar. "Crude Dr...	<1%
18	Publication	H. Cihan, M. Tunca. "Comparison of Asymmetric Dimethyl Arginine and Oxidative ...	<1%
19	Student papers	Universitas Mataram on 2023-05-18	<1%
20	Internet	www.frontiersin.org	<1%
21	Internet	www.pubtexto.com	<1%
22	Student papers	Fakultas Kedokteran Gigi Universitas Trisakti on 2020-11-28	<1%
23	Student papers	The University of Manchester on 2019-08-16	<1%
24	Internet	link.springer.com	<1%

25

Internet

www.msjonline.org

<1%

Available online on 15.07.2023 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

Copyright © 2023 The Author(s). This is an open-access article distributed under the terms of the CC BY-NC 4.0 which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited



Open Access Full Text Article



Check for updates

Research Article

Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years

Meiyanti* , Yohana , Eveline Margo , Juni Chudri, Pusparini , Meutia Atika Faradilla

Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Article Info:

Abstract



Article History:

Received 08 May 2023
Reviewed 11 June 2023
Accepted 24 June 2023
Published 15 July 2023

Cite this article as:

Meiyanti, Yohana, Margo E, Chudri J, Pusparini, Faradilla MA, Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years, *Journal of Drug Delivery and Therapeutics*. 2023; 13(7):52-56

DOI: <http://dx.doi.org/10.22270/jddt.v13i7.6142>

*Address for Correspondence:

Meiyanti, Department of Pharmacology and Medical Pharmacy, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Aging is a progressive process caused by a physiological decline in body functions. It is known that several factors can accelerate aging, namely age, lifestyle, oxidative stress, lipid peroxidation, and mitochondrial damage. Various pollutants such as cigarette smoke, motor vehicle fumes, industrial waste, and exposure to sunlight / ultraviolet light will form free radicals. Malondialdehyde (MDA) level is one of the indicators used to measure oxidative stress. Increased MDA levels are found in

various degenerative diseases. This study aims to determine plasma MDA levels in the age group of 40 years and over and factors that can affect plasma MDA levels. This study was a cross-sectional study by including 42 subjects. Data collection was carried out using a questionnaire to obtain subject demographic data and physical activity, as well as physical examinations complemented by laboratory examinations and MDA levels. The mean age of the subjects was 52.7 years, and most of the 32 (76.2%) subjects were categorized as obese and had high activity. Pearson correlation test obtained body mass index statistically significantly correlated with plasma MDA levels with a value of $p = 0.039$, as well as physical activity statistically significantly correlated with a negative correlation, with a value of $r = -0.26$ and $p = 0.043$. Conclusion Body mass index and activity are significantly correlated with plasma MDA levels.

Keywords: malondialdehyde, stress, oxidative, radicals

INTRODUCTION

Aging is a progressive process caused by a physiological decline in body function. ¹ A decrease in body homeostasis is accompanied by an increased risk of various chronic diseases including diabetes mellitus, cancer, neurodegenerative diseases, and heart disease. It is known that several factors can accelerate aging, namely age, lifestyle, oxidative stress, lipid peroxidation, and mitochondrial damage. Previous studies have shown that changes in sphingomyelin metabolism in the central nervous system and age are associated with neurodegenerative diseases. ² In the US, aging is one of the most important factors in diabetes mellitus and heart disease. In 2015, an estimated 12% of the population had diabetes mellitus and an undetected 24% had diabetes mellitus. Heart disease is the highest contributor to mortality in the population aged 65-84 years. ³ Recent studies have shown that non-diabetic populations whose fasting glucose levels increase in the blood will increase the risk of coronary heart disease. But on the other hand, the data on fasting blood glucose increase is still debated in the population of cardiovascular disease patients who do not have diabetes mellitus. A recent study in North Sumatra showed a significant relationship between MDA levels in obese and non-obese populations. ⁴

exogenous and endogenous processes. Various pollutants such as cigarette smoke, motor vehicle fumes, industrial waste, and exposure to sunlight/ultraviolet will form free radicals. Illness, stress, and excessive exercise/physical activity are endogenous factors produced by the human body that can form free radicals. The formation of free radicals occurs due to oxidation processes and metabolic processes that take place during exercise or activity, inflammatory processes, consumption of certain foods, and others. Free radicals will begin to activate with molecules in the surrounding cells to obtain electrons to stabilize the condition of one's body. The electrons that have been taken in the cell molecules will trigger the occurrence of free radicals. If this continues for a long time, it can cause oxidative stress which can cause inflammation, and DNA disease can also trigger the disease. Oxidative stress conditions characterized by increased production of Reactive Oxygen Species (ROS) have implications for various diseases such as hypertension, atherosclerosis, diabetes mellitus, stroke, chronic renal failure, heart failure, and other chronic diseases. ⁵⁻⁷

Oxidative stress is an imbalance between oxidants (free radicals) and antioxidants. Free radicals that commonly cause oxidative damage are reactive oxygen species (ROS).⁸ ROS compounds are formed through aerobic metabolism. These compounds can cause extensive damage through catalytic reactions. They are responsible for damage at the cellular level

ISSN: 2250-1177

[52]

CODEN (USA): JDDTAO

by oxidizing proteins, lipids, DNA (Deoxyribonucleic acid), and other biomolecules. Various compounds can be used as markers of oxidative stress such as malondialdehyde (MDA), and protein carbonylation. 4-hydroxy-2-nonenal (4HNE), and 8-hydroxy-deoxyguanosine (8OHdG).⁹

Malondialdehyde (MDA) is a result of lipid peroxidation in the body. The concentration of MDA indicates an oxidation process that occurs in the body's cell membranes due to metabolic processes. One of the factors that cause aging is the production of excess reactive oxygen species (ROS). ROS are generated through the process of cell respiration in the mitochondria and the phagocytosis process of foreign body elimination. To stabilize the amount of ROS, the body can use endogenous and exogenous antioxidants. Disruption of the balance between ROS and antioxidants can cause irreversible damage to cell organelle components up to the DNA level.^{10,11}

MDA levels are strongly influenced by many factors such as age, illness, stress, various pollutants, food consumption, antioxidant supplements and excessive physical activity or exercise, and others. The other study found that more than 30% of subjects had high MDA levels, and concluded that there was a significant relationship between eating habits, body mass index, glucose levels with MDA levels, as well as smoking habits significantly related to MDA levels.¹² Recent studies have shown that plasma MDA levels in populations aged 70 years and 60 years are not significantly different but there is an increase in the specific activity of the catalase enzyme that increases with age.¹⁰ In another study, there was a significant difference in plasma MDA levels in the obesity group without metabolic syndrome compared to the obesity group with metabolic syndrome.⁴

There are still variations in MDA data in various groups with different ages and different disease groups, so this study aims to determine plasma MDA levels in the age group 40 years and over and factors that can affect plasma MDA levels.

METHOD

This observational analytic study with a cross-sectional design approach was conducted from November 2022 to February 2023 in Angke village, West Jakarta, Indonesia. The inclusion criteria for this study were: men and women aged ≥ 40 years, willing to participate in the study and sign informed consent. Meanwhile, the exclusion criteria were having a history of malignancy, kidney failure, a history of autoimmune disease, taking drugs related to psychiatric conditions, and having limitations in mobility. This study used the calculation of sample size with the formula $N / 1 + N.\alpha^2$, and obtained the minimum number of samples required was 40 subjects. Demographic data and physical activity were obtained from distributed questionnaires. Blood chemistry examination (complete blood, glucose, total cholesterol, liver enzymes (SGOT and SGPT) using an enzymatic method. Examination of malondialdehyde levels was carried out using the spectrophotometric method. This research has obtained approval from the ethics committee of the Faculty of Medicine, Trisakti University, Number: 178/KER/FK/1X/2022.

RESULTS

A total of 42 subjects participated in this study, most of the subjects were female 27 (64.3%), and the mean age of the subjects was 52.7 years. Anthropometric data of the subjects are as follows body weight 67.53 ± 13.64 and height 156.02 ± 8.73 cm, while for abdominal circumference the average is 93.09 cm. Most of the subjects 59.5% fell into the normotensive category, with an average systolic blood pressure of 141.62 mmHg and an average diastolic blood pressure of 82.45 mmHg. A total of 32 (76.2%) subjects were

categorized as obese with BMI ≥ 25.0 , and most subjects had high activity levels. (Table 1)

Table 1: Distribution of Subject Characteristics (n=42)

Characteristics	n (%)	Mean \pm SD
Gender		
Male	15 (35.7)	
Female	27 (64.3)	
Blood pressure (mmHg)		
Normal (< 140/90)	25 (59.5)	
Hypertension ($\geq 140/90$)	17 (40.5)	
Physical Activity		
Low (< 600 Mets)	11 (26.2)	
High (≥ 600 Mets)	31 (73.8)	
Body Mass Index (kg/m²)		
Not obese (< 25.0)	10 (23.8)	27.69 \pm 4.90
Obese (≥ 25.0)	32 (76.2)	
Age (years)		52.7 \pm 10.60
Body weight (kg)		67.53 \pm 13.64
Height (cm)		156.02 \pm 8.73
Abdominal circumference (cm)		93.09 \pm 10.54
Blood pressure (mmHg)		
Systole		141.62 \pm 20.77
Diastole		82.45 \pm 12.70

The examination of the subject's plasma malondialdehyde levels obtained a mean of 0.77 with the highest value of 1.03 and the lowest of 0.51. Hemoglobin levels averaged 13.16 g/dL. Fasting blood glucose level was 95.06 ± 41.01 mg/dL and total cholesterol was 179.83 ± 31.85 mg/dL. Laboratory examinations to determine liver function were carried out by measuring SGPT and SGOT levels, obtaining mean values of 24.21 U/L and 26.31 U/L respectively. (Table 2)

Table 2: Laboratory examination results of subjects (n=42)

Variable	Mean \pm SD
Hemoglobin (g/dL)	13.16 \pm 1.56
Fasting blood glucose (mg/dL)	95.06 \pm 41.01
Total cholesterol (mg/dL)	179.83 \pm 31.85
SGPT (U/L)	24.21 \pm 13.50
SGOT (U/L)	26.31 \pm 11.51
Plasma malondialdehyde	0.77 \pm 0.26

Determining the type of correlation test to be used by first conducting a data normality test. The normality test used was Shapiro-Wilk because the number of subjects was less than 50 subjects. The correlation test used to assess the correlation between blood glucose, total cholesterol, SGPT, SGOT levels, age, body mass index and physical activity with plasma MDA levels was the Pearson correlation test. The test results are presented in Table 3. Blood glucose, total cholesterol, SGPT,

SGOT, and age did not have a significant correlation with MDA levels. Body mass index has a significant correlation with plasma MDA ($p = 0.039$), with a positive correlation direction, and the strength of the correlation is statistically weak with a value of $r = 0.396$ (<0.4). The physical activity variable has a significant correlation with plasma MDA levels ($p=0.043$) with a negative correlation direction but with a statistically weak correlation strength ($r=-0.26$).

Table 3: Correlation Test of Age, Gender, Glucose, SGOT, SGPT, BMI, Physical Activity with Plasma MDA Levels (n=42)

	Plasma MDA Level	
	r	p
Age	0.098	0.756
Gender	0.124	0.108
Glucose	0.330	0.835
SGOT	0.104	0.513
SGPT	0.008	0.614
Body Mass Index	0.396	0.039*
Physical activity	-0.260	0.043*

* $p < 0.05$

DISCUSSION

The level of oxidative stress can be measured using Malondialdehyde (MDA) concentration. MDA is the result of lipid peroxidation, which will cause cell damage. Oxidant compounds such as free radicals or non-radical species attack lipids that contain carbon-carbon double bonds, especially unsaturated fatty acids (PUFA). MDA is one of the biomarkers widely used to assess oxidative stress. The mean serum MDA in this study 0.77 was found to be lower than the previous study¹⁰, this is because the age difference of the subjects in this study was younger than in the previous study. In contrast, a study conducted in Iraq on hypertensive patients concluded that the increase in MDA was associated with an increase in age, gender, and duration of disease.¹³

Free radicals such as ROS are the etiology of most chronic diseases, as it is known that active free radicals are the result of metabolic reactions. Damage to body tissue cells occurs because active free electrons remain looking for other electrons to settle, if no other electrons are available then active free electrons will attack cells resulting in cell damage. Although free radicals naturally have benefits for the body, when the level of free radicals exceeds the normal value, it will cause harm to the body.^{11,12}

Measurement of oxidative stress can be done by measuring plasma malondialdehyde (MDA) levels. MDA is a secondary result of the lipid peroxidation process, producing various active compounds that will damage cells. The process of MDA formation through enzymatic and non-enzymatic processes. Currently, MDA is one of the biomarkers used to assess oxidative stress.¹⁴ Some previous researchers concluded that higher serum MDA was found in patients with hyperlipidemia. The condition of increased fat content in the body is in line with increased oxidative stress. Hyperlipidemia is associated with physical changes in cell membranes that facilitate the release of free radicals from the mitochondrial electron transport chain or activation of NADPH oxidase. In patients with diabetes mellitus (DM) there is also an increase in MDA levels, malondialdehyde interacts irreversibly and reversibly with proteins and phospholipids to have a profound effect, especially on the collagen of the cardiovascular system. MDA levels remain within normal limits due to the antioxidant

effect, but in DM, this protective effect is impaired resulting in increased MDA levels.^{1,15}

Increased lipid peroxidation is considered a consequence of oxidative stress that occurs when the dynamic balance between prooxidants and antioxidants is disrupted. The statin class of drugs for hyperlipidemia therapy is proven to be anti-atherogenic, can improve endothelial function, inhibit vascular inflammation, and stabilize atherosclerotic plaques. In addition to statins, non-steroidal anti-inflammatory drugs (NSAIDs) can affect MDA concentration. Previous studies have found that NSAIDs have significant antioxidant properties to scavenge free radicals.^{16,17}

In this study, there was no significant correlation between age and MDA levels. Similar results were also obtained by other researchers, cell damage due to oxidative stress is not related to increasing age but is associated with habits, behavior patterns, and physiological changes due to age. Various habitual patterns such as smoking, fatty food habits, lack of activity, and inflammatory processes can lead to increased oxidative stress. The habit of taking vitamin C (ascorbic acid) has a beneficial impact due to its antioxidant effect so cell damage and DNA damage are lower in this group.¹⁸ MDA levels were significantly higher among all levels of smokers. MDA levels were lower in light smokers compared to heavy or moderate smokers. Higher vitamin C levels were found in the non-smoker group. All body tissues are susceptible to free radical damage, but based on location, the respiratory epithelium is particularly vulnerable to cigarette smoke. Oxygen radicals generated close to the cell membrane oxidize membrane phospholipids which will result in lipid peroxidation with the end products of this reaction being malondialdehyde, ethane, and pentane. this causes an increase in MDA levels in smokers.¹⁹ Another study conducted in Klaten on foundry workers concluded that type of work, body mass index and smoking habits did not affect MDA levels, but length of work and marital status were associated with MDA levels. Long and continuous exposure to metallic materials will result in the accumulation of these materials in the body and will increase the oxidative process of polyunsaturated fats in cell membranes and worsen the level of lipid peroxidase, characterized by an increase in MDA levels as a reflection of the intensity and rate of lipid peroxidase reactions of damage to body cells.²⁰

Hardiany's study also concluded that there was no significant difference in plasma MDA levels in the age population, but increasing age along with an increase in catalase-specific activity.¹⁰ This increase in catalase-specific activity is a protective effect especially for elderly women to overcome high levels of oxidative damage. The role of endogenous antioxidant catalase is to suppress oxidative damage.^{18,21} Oxidative stress plays an important role in the incidence of cardiovascular diseases including coronary artery disease. Oxidative stress is considered an important mechanism for the development of cardiovascular diseases especially atherosclerosis. Hypertension, cardiac hypertrophy, and ischemia-reperfusion injury are three other major cardiovascular conditions for which gender is a risk factor. Gender is also associated with differences in oxidative stress. Under physiological conditions, women appear to be less susceptible to oxidative stress. This may be due to the antioxidant properties of estrogen, sex differences in NADPH-oxidase activity or other mechanisms yet to be determined. Sex hormones including estrogen are globally present with their receptors in the heart so sex hormones can exert effects on the cardiovascular system. Estrogen can improve heart function and reduce injury to the heart. In addition, estrogen has antioxidant effects so that it can function as a cardioprotective.^{22,23}

This study found a significant correlation between body mass index and MDA levels, the higher the BMI value, the higher the MDA levels. Similar results were also obtained in a study in India that concluded that in metabolic syndrome patients with one of the signs is an increase in BMI, there is an increase in MDA levels compared to the control group. In obese subjects, the level of MDA is higher than normal IMT. In a healthy body, there is redox hemostasis that occurs in cells, if an imbalance arises it will cause oxidative stress. Obese patients will have an increased metabolic load and mechanical load on the myocardium, increased body mass, and excessive food intake will cause the formation of lipid peroxidation, and free radicals.²⁴ The increase in some of these things will stimulate antioxidant enzymes in the long run, this will cause antioxidant enzyme reserves to be depleted and unable to overcome and balance the increase in oxidative stress.^{25,26}

These types of training activities are exhausting, long exercise, overtraining syndrome, various types of exercise will induce a significant response to oxidative stress. Exercise activities such as light exercise, low-intensity exercise, and prolonged training will have an impact on improving endogenous antioxidant status. Reactive oxygen species play an important role in cell signaling and in regulating antioxidant gene expression. Exertion results in hyperregulation of nuclear factor kappa B and mitogen-activated protein kinases that activate gene expression of a number of enzymes and proteins with important roles in maintaining intracellular oxidative or antioxidant homeostasis.^{1,6,27} The limitation of this study is that some factors that may affect MDA levels such as consumption of vitamin supplements, diet, and length of illness suffered were not examined in this study.

CONCLUSIONS

Body mass index and activity were significantly correlated with MDA levels, while age, gender, blood glucose levels, and liver enzymes were not significantly correlated with MDA levels.

Conflict of Interest:

The investigators declare there is no conflict of interest in this study.

Acknowledgments

The authors thank all the subjects who participated in this study, as well as the laboratory staff who assisted in the implementation of this study.

REFERENCES

1. Sharifi-Rad M, Anil Kumar N V., Zucca P, Varoni EM, Dini L, Panzarini E, et al. Lifestyle, Oxidative Stress, and Antioxidants: Back and Forth in the Pathophysiology of Chronic Diseases. *Front Physiol.* 2020; 11(July):1-21. <https://doi.org/10.3389/fphys.2020.00694>
2. Park MH, Jin HK, Bae J sung. Potential therapeutic target for aging and age-related neurodegenerative diseases: the role of acid sphingomyelinase. *Exp Mol Med [Internet].* 2020; 52(3):380-9. <https://doi.org/10.1038/s12276-020-0399-8>
3. Całyniuk B, Grochowska-Niedworok E, Walkiewicz K, Kawecka S, Popiołek E, Fatyga E. Malondialdehyde (MDA) - product of lipid peroxidation as marker of homeostasis disorders and aging. *Ann Acad Medicae Silesiensis.* 2016; 70:224-8. <https://doi.org/10.18794/aams/65697>
4. Savira M, Rusdiana ., Widjaja SS, Syahputra M. Comparison Malondialdehyde (MDA) Level between Obesity Non Metabolic Syndrome and Obesity with Metabolic Syndrome Patients. 2020; (5):644-7.
5. Yekti R, Bukhari A, Jafar N, Thaha AR. Measurement of malondialdehyde (MDA) as a good indicator of lipid peroxidation.

- Int J Allied Med Sci Clin [Internet]. 2018; 6(4):6-8. Available from: www.ijamscr.com
6. Denise Mafra ME. Influence of Physical Exercise on Oxidative Stress and Inflammation in Hemodialysis Patients. *J Mol Biomark Diagn.* 2015; 06(03). <https://doi.org/10.4172/2155-9929.1000e120>
 7. Mao C, Yuan JQ, Lv Y Bin, Gao X, Yin ZX, Kraus VB, et al. Associations between superoxide dismutase, malondialdehyde and all-cause mortality in older adults: A community-based cohort study. *BMC Geriatr.* 2019; 19(1):1-9. <https://doi.org/10.1186/s12877-019-1109-z>
 8. Ayala A, Muñoz MF, Argüelles S. Lipid peroxidation: Production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. *Oxid Med Cell Longev.* 2014; 2014. <https://doi.org/10.1155/2014/360438>
 9. Situmorang N, Zulham Z. Malondialdehyde (Mda) (Zat Oksidan Yang Mempercepat Proses Penuaan). *J Keperawatan Dan Fisioter.* 2020; 2(2):117-23. <https://doi.org/10.35451/jkf.v2i2.338>
 10. Hardiany NS, Sucitra S, Paramita R. Profile of malondialdehyde (MDA) and catalase specific activity in plasma of elderly woman. *Heal Sci J Indones.* 2020; 10(2):132-6. <https://doi.org/10.22435/hsj.v12i2.2239>
 11. Hardiany NS, Sadikin M, Siregar N, Wanandi SI. The suppression of manganese superoxide dismutase decreased the survival of human glioblastoma multiforme T98G cells. *Med J Indones.* 2017; 26(1):19-25. <https://doi.org/10.13181/mji.v26i1.1511>
 12. 439-445_nova.pdf.
 13. A. Hassan A, Sayyah SG. Oxidative Stress Marker Malondialdehyde and Glutathione Antioxidant in Hypertensive Patients. *Eur J Biomed Res.* 2023; 2(1):31-6. <https://doi.org/10.24018/ejbiomed.2023.2.1.47>
 14. Khoubnasab Jafari M, Ansarin K, Jouyban A. Comments on "use of malondialdehyde as a biomarker for assessing oxidative stress in different disease pathologies: A review." *Iran J Public Health.* 2015; 44(5):714-5.
 15. Shalash M, Badra M, Imbaby S, ElBanna E. Malondialdehyde in Type 2 Diabetics and Association With Cardiovascular Risk Factors. *J Med Res Inst.* 2020; 41(2):21-30. <https://doi.org/10.21608/jmalex.2020.147116>
 16. Oesterle A, Laufs U, Liao JK. Pleiotropic Effects of Statins on the Cardiovascular System. *Circ Res.* 2017; 120(1):229-43. <https://doi.org/10.1161/CIRCRESAHA.116.308537>
 17. Sekunda KN, Ariosta A, Limantoro C, Setiawan AA. Difference in Profiles of Oxidative Stress Marker (MDA) in STEMI and NSTEMI. *Diponegoro Med J (Jurnal Kedokt Diponegoro).* 2021; 10(2):145-50. <https://doi.org/10.14710/dmj.v10i2.29673>
 18. Block G, Dietrich M, Norkus EP, Morrow JD, Hudes M, Caan B, et al. Factors associated with oxidative stress in human populations. *Am J Epidemiol.* 2002; 156(3):274-85. <https://doi.org/10.1093/aje/kwf029>
 19. Kubihal C V., Naik HD. Effect of smoking on vitamin C and MDA: a cross sectional comparative study. *Int J Res Med Sci.* 2019; 7(3):746. <https://doi.org/10.18203/2320-6012.ijrms20190511>
 20. Ari Suwondo, Ahmad Mohan, Melyana Nurul Widyawati M. Factors Related to Malondialdehyde (MDA) Levels in Metal Casting Industry Workers. *Preprints.* 2020; 1(October):1-8. <https://doi.org/10.21203/rs.3.rs-99597/v1>
 21. Schöttker B, Brenner H, Jansen EHJM, Gardiner J, Peasey A, Kubínová R, et al. Evidence for the free radical/oxidative stress theory of ageing from the CHANCES consortium: A meta-analysis of individual participant data. *BMC Med.* 2015; 13(1):1-15. <https://doi.org/10.1186/s12916-015-0537-7>
 22. Kander MC, Cui Y, Liu Z. Gender difference in oxidative stress: a new look at the mechanisms for cardiovascular diseases. *J Cell Mol Med.* 2017; 21(5):1024-32. <https://doi.org/10.1111/jcmm.13038>
 23. Bell JR, Bernasochi GB, Varma U, Raaijmakers AJA, Delbridge LMD. Sex and sex hormones in cardiac stress - Mechanistic insights. J

- Steroid Biochem Mol Biol. 2013; 137:124-35.
<https://doi.org/10.1016/j.jsbmb.2013.05.015>
24. P G, Curcio F, Bulli G, Aran L, Della-morte D, Testa G, et al. Oxidative Stress and Diseases. Oxidative Stress Dis. 2012; 757-72.
25. Sabitha K, Venugopal B, Rafi M, V Ramana K. Role of Antioxidant Enzymes in Glucose and Lipid Metabolism in Association with Obesity and Type 2 Diabetes. Am J Med Sci Med. 2014; 2(1):21-4.
<https://doi.org/10.12691/ajmsm-2-1-5>
26. Manna P, Jain SK. Obesity, Oxidative Stress, Adipose Tissue Dysfunction, and the Associated Health Risks: Causes and Therapeutic Strategies. Metab Syndr Relat Disord. 2015; 13(10):423-44. <https://doi.org/10.1089/met.2015.0095>
27. Lewis NA, Howatson G, Morton K, Hill J, Pedlar CR. Alterations in Redox Homeostasis in the Elite Endurance Athlete. Sport Med. 2015; 45(3):379-409. <https://doi.org/10.1007/s40279-014-0276-5>

Bukti Bukti Screenshot pada Sinta

The screenshot displays the SINTA Executive interface. On the left is a navigation menu with options: Dashboard, Explore SINTA, Mutation History, List Verificator PT, and My SINTA. The top right shows the user profile for JUNI CHUDRI (Sinta ID: 6727331) with a status of 'Pay' and '3 | Total Records : 27'. The main content area lists three publications:

Publication Title	Author	Journal	Year	Cited
Pengukuran Kadar Kolesterol Total Dan Pelayanan Konsultasi Sindrom Metabolik Di Kelurahan Rawamangun, Jakarta Timur	A Handayani, F Chondro, LM Surjadi, J Chudri	AMMA: Jurnal Pengabdian Masyarakat 2 (5), 515-519, 2023	2023	0
Factors associated with plasma malondialdehyde levels in people over 40 years	M Melyanti, E Margo, J Chudri, MA Faradilla	Journal of Drug Delivery and Therapeutics 13 (7), 52-56, 2023	2023	2
Penyuluhan Protokol 3M Dalam Pencegahan COVID-19 Pada Penduduk Kelurahan Pluit, Kecamatan Penjaringan, Jakarta Utara	J Chudri, F Chondro, A Handayani	AMMA: Jurnal Pengabdian Masyarakat 2 (9: Oktober), 1099-1102, 2023	2023	0