

Indonesian Journal of Pharmaceutical Sciences E-ISSN 2614-6495



J I P I

JURNAL
ILMU KEFARMASIAN
INDONESIA

Volume 22. Issue 2. October 2024

JiPi | Volume 22 | Issue. 2 | Page 179-338 | Jakarta, October 2024

Keterangan: (Gambar Cover yang Vol 22 karena di website tidak ada cover Vol 23 No 1 (2025))



[HOME](#) / [Announcements](#)

NEWS RELEASE

2024-02-27

On Monday, February 26th, 2024, apt. Rahmatul Qodriah, M.Farm., successfully defended his dissertation in the doctoral promotion session, Faculty of Pharmacy, Universitas Pancasila with the title "Metabolomics of Tin leaf extract (Ficus carica L.) from 3 varieties that grow in Indonesia as an anti-metabolic syndrome".



[READ MORE](#)

Indonesian Journal of Pharmaceutical Sciences (JIFI) has collaborated with Indonesian Pharmacist Association (IAI)

2023-11-14

Indonesian Journal of Pharmaceutical Sciences (JIFI) has collaborated with Indonesian Pharmacist Association or Ikatan Apoteker Indonesia (IAI)



[READ MORE](#)

[SUBMIT AN
ARTICLE](#)

Menu

[Editorial Team](#)

[Journal Reviewers](#)

[Peer Review Process](#)

[Focus and Scope](#)

[Publication Ethics](#)

[Online Submission](#)

[Plagiarism Check](#)

[Author Guidelines](#)

[Article Processing Charge](#)

[Open Access Statement](#)

[Copyright](#)

[Indexing & Abstracting](#)

[Letter Statement not
Plagiarism](#)

[Author Statement Letter](#)

In Collaboration with:

[Mobirise](#)



Tools

[turnitin](#)

[MENDELEY](#)

ISSN



1693-1831
p-ISSN

Jifi won 1st place in the Sinta award

📅 2023-11-10

Jifi took first place in the Sinta Award at the Universitas Pancasila anniversary .



[READ MORE >](#)

Journal workshop towards becoming internationally indexed with a reputation in 2023

📅 2023-10-05

Management of the Indonesian pharmaceutical scientific journal (Jifi) took part in a journal workshop towards becoming internationally indexed with a reputation in 2023, DRTPM Ministry of Education and Culture of the Republic of Indonesia.



[READ MORE >](#)

Indonesian Journal of Pharmaceutical Sciences (Jifi) collaborates with PAFI

📅 2023-10-03

The Indonesian Journal of Pharmaceutical Sciences (Jifi) has signed a collaboration with the Management of the Indonesian Association of Pharmacy Experts (PAFI) in journal publication



[READ MORE >](#)



LANGUAGE

English
Bahasa Indonesia

INFORMATION

For Readers
For Authors
For Librarians

CURRENT ISSUE



MAKE A SUBMISSION

Open Journal Systems

395721



Preparation to be an International Journal.

 2023-09-18

Thank you very much for your kind support in submitting your research papers to our journal. Starting from April Vol 21 No 1 (2023), We have been published in English.

READ MORE >

Journal Manajement Training

 2023-09-15

Journal Management Intensive Training to all editorial boards of the Indonesian Journal of Pharmaceutical Sciences (JiFI) and Journal of Natural Product for Degenerative Diseases (JNPDD), Faculty of Pharmacy, Universitas Pancasila. Attended by expert staff of the Institute for Research and Community Service (LPPM) of Universitas Pancasila

READ MORE >

Journal Managemen

 2023-09-15

Journal Management Intensive Training to all editorial boards of the Indonesian Journal of Pharmaceutical Sciences (JiFI) and Journal of Natural Product for Degenerative Diseases (JNPDD), Faculty of Pharmacy, Universitas Pancasila. Attended by expert staff of the Institute for Research and Community Service (LPPM) of Universitas Pancasila

READ MORE >

Editorial Board Meeting " Journal Managemen"

 2023-09-15

Journal Management Intensive Training to all editorial boards of the Indonesian Journal of Pharmaceutical Sciences (JiFI) and Journal of Natural Product for Degenerative Diseases (JNPDD), Faculty of Pharmacy, Universitas Pancasila Attended by.....

READ MORE >

Preparation of accreditation

 2023-09-13

Jurnal Ilmu Kefarmasian Indonesia is applying for accreditation of the Sinta 2 journal. Website improvement, OJS flow, and equalization of perception between editors have been carried out regularly every week

READ MORE >



Faculty of Pharmacy Universitas Pancasila

Editorial Office:

Lenteng Agung St, Srengseng Sawah District,
Jagakarsa Regency, Jakarta Selatan, Special Region of
Jakarta 12640, Indonesia | e-mail:
adminjifi@univpancasila.ac.id

Support Contact

Ahmad Munadi
Phone 082111989075 (WA only)
adminjifi@univpancasila.ac.id



Jurnal Ilmu Kefarmasian Indonesia (ISSN: 1693-1831 (Print) / ISSN: 2614-6495 (Online) is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License](#). Preserved in [LOCKSS](#) through PKP Private LOCKSS Network program.



BRIN
BADAN RISET
DAN INOVASI NASIONAL



powered by OJS | Open Journal Systems

PKP | PUBLIC KNOWLEDGE PROJECT

Jurnal Ilmu Kefarmasian Indonesia

S₂

H-INDEX : 14

H5-INDEX : 14

Terakreditasi **SINTA 2**





[HOME](#) / [ARCHIVES](#) / Vol 23 No 1 (2025): JIFI

DOI: <https://doi.org/10.35814/jifi.v23i1>

PUBLISHED: 2025-04-28

ARTICLES

Assessment on the antioxidant, photoprotective, and antimicrobial properties of *Siraitia grosvenorii*

Figo Figo, Tasya Mariska Candra, Jessica Novia, Marcelia Sugata

1-7



PDF

Statin prescribing pattern and the outcomes for acute coronary syndrome as primary and secondary prevention: a comprehensive review

Pramitha Esha Nirmala Dewi, Farahdina Chairani, Wahyu Himawan

8-17



PDF

Evaluation of drug services based on prescription indicators and patient care according to who at the hospital general practitioner (GP) outpatient clinic in Depok, Indonesia

Nadia Farhanah Syafhan, Hindun Wilda Risni, Azzahra Fahira Salsabila, Raden Jacinda Yasmin Purnama Putri, Marchen Prasetyaningrum

18-28



PDF

Andrographis paniculata Burm. F. in-silico analysis compounds that function as an insulin sensitizer therapy for type 2 diabetes via peroxisome proliferator activated gamma receptors (ppary) receptor activator

Pristiyantoro Pristiyanoro, Siswandono Siswandono, Esti Mumpuni

29-38



PDF

Evaluation of antipsychotic utilization among outpatient schizophrenia patients at the mental health facility in West Java

Entris Sutrisno, Nita Selifiana, Jajang Japar Sodik, Uvika Nurul Fajarrini

39-48



PDF

Characteristic and therapeutic effect of lemongrass leaf encapsulation with a different ratio of chitosan and NaTPP

Florencia Livia Kurniawan, Komariah Komariah, Octarina Octarina, Febry Anggreani, Fauziah Rahma Annisa, Farah Nurlidar

49-57



PDF

[SUBMIT AN
ARTICLE](#)

Menu

[Editorial Team](#)

[Journal Reviewers](#)

[Peer Review Process](#)

[Focus and Scope](#)

[Publication Ethics](#)

[Online Submission](#)

[Plagiarism Check](#)

[Author Guidelines](#)

[Article Processing Charge](#)

[Open Access Statement](#)

[Copyright](#)

[Indexing & Abstracting](#)

[Letter Statement not](#)

[Plagiarism](#)

[Author Statement Letter](#)

In Collaboration with:



Tools



ISSN



p-ISSN



e-ISSN

Formulation of a peel-off gel mask containing mundu fruit extract (*Garcinia dulcis* (roxb.) kurz) for antioxidant and anti-elastase activity

Neneng Siti Silfi Ambarwati, Yesi Desmiaty, Amelia Chandra, Yuslia Noviani, Yulia Triyani, Ratu Mayra Hakim, Reistha Fazlia Amanda, Dion Wahyudi 58-66



Optimization of hand sanitizer gels containing soursop (*Annona muricata* L.) leaf extract using simplex lattice design method

Kosasih Kosasih, Dita Lismawati, Rifael Satrio Adinugroho 67-82



Adverse events among pulmonary drug-resistant tuberculosis patients in Banten, Indonesia

Mita Restinia, Alyya Siddiqi, Wulan Purnama Sari 83-90



Evaluation of the rationality of drugs use inpatient patients with type 2 diabetes mellitus of 2023 at Hospital Y Depok

Sondang Khairani, Reise Manninda, Bitya Aulia Wijayanti 91-96



Anticancer activity of *Stichopus vastus* extract against MCF-7 and HeLa cells from Natuna waters

Mery Sukmiwati, Nurjanah Nurjanah, Andarini Diharmi, Indah Prasetya Putri 97-110



Effectiveness of *Artocarpus heterophyllus* lamk. serum in eliminating dark spot in the guinea pig melasma model due to UV-a exposure

Ida Ayu Dwi Putri Mas Tiari, Ni Putu Kania Mardiyanti Putri, Ni Luh Setia Marliani, Ni Made Elma Agustini Sari, Julia Eva Malelak, Ketut Agus Adrianta 111-122



Natural deep eutectic solvents (NaDES) improves extraction and antioxidant activity of stem bark of *Garcinia cowa* Roxb.

Fahleni Fahleni, Abdul Mun'im, Fadlina Chany Saputri, Silvia Surini 123-130



Tyrosinase inhibitory activity and physicochemical properties of microalgae extract (*Spirulina platensis*)-loaded nanostructured lipid carriers gel

Faizatun Faizatun, Abdullah Abdullah 131-137



Nanostructured lipid carrier (NLC) in topical preparations: a narrative review of components, manufacturing methods, characteristics and activities

Yayan Rizikiyan, Nining Sugihartini, Ichwan Ridwan Rais 138-156



Spray gel film-forming system formulation of *Vernonia amygdalina* as diabetic wound healer


Putri Rachma Novitasari, Yudha Rizky Nuari, Azzahro Aulia Firdausi 157-165



e-ISSN



Visitors



LANGUAGE

English
Bahasa Indonesia

INFORMATION

For Readers
For Authors
For Librarians

CURRENT ISSUE



MAKE A SUBMISSION

Open Journal Systems

395789



The role of pharmacist counseling for patient with type 2 diabetes mellitus and its impact on blood glucose control and quality of life at RSUD Pasar Minggu

Devita Hardi, Yusi Anggriani, Fredrick Dermawan Purba

166-175



Bibliometric analysis on pharmaceutical care publications in Southeast Asia: prospective research trends

Rahmat Riyadi, Sofa Dewi Alfian

176-189



Analysis of pharmaceutical service improvement based IPA (Importance Performance Analysis) at pharmacy Kimia Farma Ijen, Malang

Dian Nurmawati, Tridoso Sapto Agus Priyono, Arifani Siswidasari

190-199



Evaluation of the combination patch of betel leaf extract (Piper sp.) - ultrasonic assisted extraction as a transdermal delivery system on fibroblast cell formation and collagen density

Regina Valencia, Elsa Trinovita, Herlina Eka Shinta, Angeline Novia Toemon, Ysrafil Ysrafil

200-211





[HOME](#) / [Editorial Team](#)

Editor in Chief

[Prof. Dr. apt. Shirly Kumala, M. Biomed](#)

Universitas Pancasila, Faculty of Pharmacy, Jakarta, Indonesia

Editorial Board

[Prof. Dr. Syed Abdul Azeez Basha](#)

Darussalam, Aghapura, Nampally, Hyderabad, India

[Prof. Arshad Mahmood, PhD](#)

Collage of Pharmacy, Al Ain University, Abu Dhabi

[Dr. Erwin Martinez Faller](#)

San Pedro College, Davao City, Philliphine

[Uttam Budhathoki, BPharm, MPharm, PhD](#)

Kathmandu University, Dhulikhel, Kavre, Nepal

[Prof. Dr. Taifo Mahmud, PhD., M.Sc](#)

Oregon State University, Corvallis, United States

[Prof. Dr. Deny Susanti](#)

International Islamic University Malaysia

[apt. Diah Kartika Pratami, M. Farm](#)

Universitas Pancasila, Faculty of Pharmacy, Jakarta, Indonesia

[apt. Safira Nafisa, S. Farm., M. Farm](#)

Universitas Pancasila, Faculty of Pharmacy, Jakarta, Indonesia

[apt. Lusiana Ariani, S. Farm., M. Farm](#)

Universitas Pancasila, Faculty of Pharmacy, Jakarta, Indonesia

[apt. Anarisa Budiati, S.Farm., M. Farm](#)

Universitas Pancasila, Faculty of Pharmacy, Jakarta, Indonesia

[apt. Reise Manninda, S. Farm., M. Farm](#)

Universitas Pancasila, Faculty of Pharmacy, Jakarta, Indonesia

[apt. Rahmatul Qodriah, S. Farm., M. Farm](#)

Universitas Pancasila, Faculty of Pharmacy, Jakarta, Indonesia

[apt. Zainur Rahman Hakim, S. Farm., M. Farm](#)

Universitas Pancasila, Faculty of Pharmacy, Jakarta, Indonesia

[apt. Intan Permata Sari, S. Farm., M. Farm](#)

Universitas Pancasila, Faculty of Pharmacy, Jakarta, Indonesia

[SUBMIT AN
ARTICLE](#)

Menu

[Editorial Team](#)

[Journal Reviewers](#)

[Peer Review Process](#)

[Focus and Scope](#)

[Publication Ethics](#)

[Online Submission](#)

[Plagiarism Check](#)

[Author Guidelines](#)

[Article Processing Charge](#)

[Open Access Statement](#)

[Copyright](#)

[Indexing & Abstracting](#)

[Letter Statement not
Plagiarism](#)

[Author Statement Letter](#)

In Collaboration with:



Tools



ISSN



p-ISSN



e-ISSN

Managing Editor


[Dwi Fajar Saputra S.Sos., MM](#)

Website Management

Ahmad Munadi, S.Kom
Rofiqoh Hadiyah, S.Kom., S.Pd.
Amipriyadi Gerha Permana, SIP

Publisher

[Faculty of Pharmacy, Universitas Pancasila](#)



LANGUAGE

English
Bahasa Indonesia

INFORMATION

For Readers
For Authors
For Librarians

CURRENT ISSUE



MAKE A SUBMISSION

Open Journal Systems

395715





Characteristic and therapeutic effect of lemongrass leaf encapsulation with a different ratio of chitosan and NaTPP

Florencia Livia Kurniawan

Department of Dental Material Science and Technology, Faculty of Dentistry, Universitas Trisakti, West Jakarta, 31657, Indonesia

<http://orcid.org/0009-0007-7107-3142>

Komariah Komariah

Department of Oral Biology, Faculty of Dentistry, Universitas Trisakti, West Jakarta, 31657, Indonesia

Octarina Octarina

Department of Dental Material Science and Technology, Faculty of Dentistry, Universitas Trisakti, West Jakarta, 31657, Indonesia

<http://orcid.org/0000-0002-8764-8365>

Febry Anggreani

Dentistry Study Program, Faculty of Dentistry, Universitas Trisakti, West Jakarta, 31657, Indonesia

Fauziah Rahma Annisa

Dentistry Study Program, Faculty of Dentistry, Universitas Trisakti, West Jakarta, 31657, Indonesia

Farah Nurlidar

Research Center for Radiation Processing Technology, Research Organization for Nuclear Energy, National Research and Innovation Agency, South Jakarta, 12440, Indonesia

DOI: <https://doi.org/10.35814/jifi.v23i1.1570>

ABSTRACT

The lemongrass plant (*Cymbopogon citratus*) in Indonesia has great potential as a natural ingredient for medicines because it contains various compounds such as neral, citral, geranial acetate, flavonoids, and tannins. Previous research has demonstrated various pharmacological activities of kitchen lemongrass leaves, including antibacterial, antifungal, and anti-inflammatory properties. Encapsulation technology in the form of chitosan-NaTPP nanoparticles is used to maintain the activity of kitchen lemongrass leaf compounds. This research aims to determine the physical characteristics of the encapsulation of kitchen lemongrass leaf extract with several comparisons of chitosan and NaTPP by testing PSA (particle

PDF

PUBLISHED

2025-04-30

HOW TO CITE

KURNIAWAN, Florencia Livia et al. Characteristic and therapeutic effect of lemongrass leaf encapsulation with a different ratio of chitosan and NaTPP. JURNAL ILMU KEFARMASIAN INDONESIA, [S.l.], v. 23, n. 1, p. 49-57, apr. 2025. ISSN 2614-6495. Available at: <http://jifi.farmasi.univ.pancasila.ac.id/index.php/jifi/article/view/1570>. Date accessed: 10 July 2025. doi: <https://doi.org/10.35814/jifi.v23i1.1570>.

CITATION FORMATS

[ABNT](#)
[APA](#)
[BibTeX](#)
[CBE](#)
[EndNote - EndNote format \(Macintosh & Windows\)](#)
[MLA](#)
[ProCite - RIS format \(Macintosh & Windows\)](#)
[RefWorks](#)
[Reference Manager - RIS format \(Windows only\)](#)

SUBMIT AN
ARTICLE

Menu

[Editorial Team](#)
[Journal Reviewers](#)
[Peer Review Process](#)
[Focus and Scope](#)
[Publication Ethics](#)
[Online Submission](#)
[Plagiarism Check](#)
[Author Guidelines](#)
[Article Processing Charge](#)
[Open Access Statement](#)
[Copyright](#)
[Indexing & Abstracting](#)
[Letter Statement not Plagiarism](#)
[Author Statement Letter](#)

In Collaboration with:



Tools



ISSN





Characteristic and therapeutic effect of lemongrass leaf encapsulation with a different ratio of chitosan and NaTPP

Florencia Livia Kurniawan¹, Komariah Komariah^{2*}, Octarina Octarina¹, Febry Anggreani JK³, Fauziah Rahma Annisa³, Farah Nurlidar⁴

¹Department of Dental Material Science and Technology, Faculty of Dentistry, Universitas Trisakti, West Jakarta, 31657, Indonesia

²Department of Oral Biology, Faculty of Dentistry, Universitas Trisakti, West Jakarta, 31657, Indonesia

³Dentistry Study Program, Faculty of Dentistry, Universitas Trisakti, West Jakarta, 31657, Indonesia

⁴Research Center for Radiation Processing Technology, Research Organization for Nuclear Energy, National Research and Innovation Agency, South Jakarta, 12440, Indonesia.

*Corresponding Author: komariah@trisakti.ac.id

Received date: 15 March 2025 / Accepted date: 29 April 2025

ABSTRACT: The lemongrass plant (*Cymbopogon citratus*) in Indonesia has great potential as a natural ingredient for medicines because it contains various compounds such as neral, citral, geranial acetate, flavonoids, and tannins. Previous research has demonstrated various pharmacological activities of kitchen lemongrass leaves, including antibacterial, antifungal, and anti-inflammatory properties. Encapsulation technology in the form of chitosan-NaTPP nanoparticles is used to maintain the activity of kitchen lemongrass leaf compounds. This research aims to determine the physical characteristics of the encapsulation of kitchen lemongrass leaf extract with several comparisons of chitosan and NaTPP by testing PSA (particle size analysis test), FTIR (functional group analysis test), viscosity, and stability, including pH and turbidity tests. Encapsulation with a 1:1 ratio of chitosan and NaTPP has a particle size of 524 nm and a PDI value of 0.481, classified as monodisperse (PDI <0.7). The pH value ranges from 6.46 to 7.30, and the viscosity value was 2.134–2.169 cP, which is still within mouthwash standards. The turbidity test showed stable encapsulation results. Therefore, encapsulation with a 1:1 ratio of chitosan and NaTPP is the most optimal choice and has potential for developing the therapeutic effect of mouthwash.

KEYWORDS: Chitosan; encapsulation; lemongrass leaves; mouthwash; NaTPP.

• INTRODUCTION

Indonesia is rich in natural plant sources as raw materials for medicines. One plant that is widely used is the kitchen lemongrass leaves. Many researchers are taking advantage of technological advances and making various efforts to improve the quality and trust in the benefits of natural sources, one of which is kitchen lemongrass leaves [1]. *Cymbopogon citratus*, commonly called kitchen lemongrass, has a larger lower stem, is white, and does not smell too sharp. The part that is often used as a kitchen spice is the young inner part. Lemongrass plays a role as an aromatic spice in cooking and is also used as a complement to chili sauce. These plants can both produce essential oils, but in Indonesia, only citronella is widely processed as an essential oil producer [2]. The results of various studies show that citronella leaves are rich in essential oils and contain neral (31.55%), citral (26.1%), and geranial acetate (2.27%). This ingredient is often used as an aromatherapy oil [3]. This plant also contains active compounds, namely flavonoids, saponins, tannins, phenols, aldehydes, phenolic acids, and terpenoids. Studies also show that lemongrass leaves have various pharmacological properties, such as antibacterial, antifungal, and anti-inflammatory, especially when used as a mouthwash [4].

One way that can be used to protect secondary metabolic activity in lemongrass leaf extract is through encapsulation technology, which is a technique for covering secondary metabolism to protect it from external influences. Encapsulation is divided into micro and nanoparticle forms. Applications of food nanotechnology show an increasing trend. This technology offers advantages in increasing secondary metabolic bioavailability, controlling the release of active compounds, and improving sensory properties [5]. Nanoparticles are solid colloidal particles with a diameter of 1-1000 nm and are materials that can be used as encapsulated active compounds [6]. The active compound of lemongrass leaves has poor stability, so it is encapsulated with horn beetle chitosan to increase its stability, such as antioxidant, anti-cancer, anti-inflammatory, anti-fungal, and antibacterial [7].

How to cite this article: Kurniawan FL, Komariah K, Octarina O, Anggreani F, Annisa FR, Nurlidar F. Characteristic and therapeutic effect of lemongrass leaf encapsulation with a different ratio of chitosan and NaTPP. JIFI. 2025; 23(1): 49-57.

Chitosan is a natural polysaccharide that is non-toxic and easily biodegradable in delivery systems because it has properties as a matrix. Chitosan also forms gels in acidic conditions and has a structure similar to chitin [8]. Chitosan is a compound of poly (N-amino-2-deoxy- β -D-glucopyranose) or glucosamine resulting from the deacetylation of chitin/poly (N-acetyl-2-amino-2-deoxy- β -D-glucopyranose) [9].

Previous studies by Purwandari showed that moringa leaf extract was obtained by repeated ultrasonication of 1x15, 3x15, and 5x15 minutes, frequency 24 kHz, and using 96% alcohol solvent. The encapsulation process was carried out with three variable ratios of the amount of chitosan, Na-TPP added to 10 mL of moringa leaf extract solution, namely 5:1, 2:1 and 1:1. The results of the flavonoid test as quercetin using a UV-Vis spectrophotometer showed that the quercetin content in moringa leaf extract resulting from 5x15 minute ultrasonication was 12.146%, Particle Size Analyzer (PSA) characterization showed encapsulated particle sizes of 170.3 and 209.7 nm with a polydispersion index <0.5, nanoencapsulation of moringa leaf extract has been produced. There has been an increase in the antioxidant activity of moringa leaf extract after encapsulation using chitosan-NaTPP 2:1, namely from 61.4% to 81.2% [9].

Based on the background above, this research was conducted to determine the physical characteristics of chitosan encapsulation of kitchen lemongrass leaf extract with several comparisons of chitosan and NaTPP, which will determine the activity of the encapsulated particles as antibacterial, anticancer, and antioxidant.

MATERIALS AND METHODS

Materials

This study was conducted with lemongrass leaf extract (Bogor, Indonesia), 70% ethanol (Merck, Germany), 1% acetic acid (Merck, Germany), distilled water (Merck, Germany), chitosan *Xylotrupes gideon* (Bogor, Indonesia), NaTPP (Merck, Germany), and Tween 80 (Merck, Germany).

Kitchen lemongrass leaf extraction

Lemongrass leaf simplicia was extracted using the maceration method by soaking samples of kitchen lemongrass leaves in 70% ethanol with a ratio of 1:10 for 3 × 24 hours at a temperature of 27 °C. The solution will be filtered using Whatman filter paper and spun using an evaporator machine at 50-60 °C.

Encapsulation of lemongrass leaf extract

The nano chitosan encapsulation stage was achieved by dissolving 40 mL of chitosan in 1% acetic acid and adding 100 mL of distilled water. Lemongrass leaf extract is dissolved using distilled water. Then, the extract solution was mixed into the chitosan solution. Stirring was carried out at a speed of 2500 rpm for 2 hours. Next, tripolyphosphate was given drop by drop. Finally, 0.1 mL of Tween 80 was added to the solution. In making the encapsulation, a comparison was made between the extract and chitosan, presented in Table 1.

Table 1. Encapsulation comparison.

Group (ratio chitosan : NaTPP)	Chitosan (mg/mL)	NaTPP (mL)	Lemongrass (mL)	Tween 80 (mL)
Group 1 (1 : 1.0)	0.57	0.57	2	0.1
Group 2 (1 : 0.9)	0.57	0.514	2	0.1
Group 3 (1 : 0.8)	0.57	0.468	2	0.1
Group 4 (1 : 0.7)	0.57	0.429	2	0.1

Particle size analysis test

An encapsulated sample of lemongrass leaf extract in liquid form was prepared and dropped into a cuvette to which sufficient distilled water was added as a solvent. Then, it was put into the PSA apparatus, and the temperature was set to 25 °C. The results of the sample particle size distribution will appear on the monitor.

FTIR test

The FTIR (Fourier Transform Infrared Spectroscopy) test was used to analyze the functional groups contained in the sample. The chemical interactions between NaTPP, chitosan, and kitchen lemongrass leaf extract were analyzed using an FTIR spectrophotometer at wave numbers 4000 – 400 cm^{-1} . Analysis was carried out using the Shimadzu Prestige-21 FTIR tool. A total of 1 mL of encapsulated sample for each formulation was frozen using a freeze dryer until it reached a temperature of -80 °C. Then, the sample was put into the tool and analyzed. Next, the results are displayed on the monitor.

Stability test

Turbidity test

Turbidity test by visible observation. This observation aims to see the occurrence of material aggregation within 2 weeks. The measurement results were color and turbidity.

pH determination test

Acid measurements were carried out using a Mettler Toledo pH meter. First, prepare the sample whose acid level you want to measure (place it in a container). Next, the electrode from the pH meter was inserted into a container containing the solution to be tested. When immersed in the solution, the number scale will move randomly. Wait until the number stops and does not change. The results will be visible on the monitor.

Viscosity test

The viscosity test was carried out through measurements using the ViscoQC 100 tool Anton Paar (rotational viscometer). The time is set to 1 minute, the torque is 56%, and the speed is 200 rpm. The results displayed on the monitor screen are recorded. Viscosity measurements were carried out on days 1, 7, and 14.

RESULTS

Particle size analysis test (PSA)

The PSA measurements showed that the encapsulation group with a 1:1 ratio of chitosan and NaTPP had the smallest size of 524.0 nm compared to the other comparison groups. The comparison group for chitosan and NaTPP encapsulation, which has the largest to smallest size, was the ratio of chitosan and NaTPP encapsulation 1:0.7, 1:0.8, 1:0.9, and 1:1, with particle sizes of 1145 nm, 934.2 nm, 614.5 nm, and 524 nm, respectively, as shown in Table 2. Particle measurement results, the ratio of chitosan and NaTPP encapsulation was 1:0.8, 1:0.9, and 1:1, which fall into nano-sized particles, while the encapsulation ratio of chitosan and NaTPP 1:0.7 with a size of more than 1000 nm does not meet the requirements for particle preparation.

Polydispersity Index (Pdl) is an index of the heterogeneity of a collection of particles. The average Pdl value of the encapsulation ratio of chitosan and NaTPP was 1:1, 1:0.9, and 1:0.8, with Pdl values of 0.481, 0.695, and 0.687, respectively. Pdl values from a 1:1 chitosan encapsulation ratio of 1:0.9 and 1:0.8 have a Pdl value of <0.7. According to Chasana et al., particles can be classified into monodisperse groups (Pdl value <0.7) and polydisperse large size distribution (Pdl value >0.7) [10]. A Pdl value <0.7 indicates good particles because it is in the middle range of the polydispersity index. For encapsulation, the ratio of chitosan and NaTPP 1:0.7 has a value of 0.747, which indicates that the sample has a part with a large particle size and is not homogeneous.

Table 2. Particle size and polydispersity index.

Group (ratio chitosan : NaTPP)	Particle size (nm)	Pdl
Group 1 (1 : 1.0)	524.0	0.481
Group 2 (1 : 0.9)	614.5	0.695
Group 3 (1 : 0.8)	934.2	0.687
Group 4 (1 : 0.7)	1145	0.747

FTIR test

The graphical image of the analysis of the FTIR test results showed that several functional groups were identified in the four samples (Figure 1). In the chitosan extract sample, there were absorption peaks in the wave number area 3425 cm^{-1} (O-H stretching vibrations overlap with N-H), 1627.99 cm^{-1} (C=O Amide I stretching vibrations), 1579.77 cm^{-1} (C=O Amide I stretching vibrations). NH_2 bending in Amide II). The four chitosan encapsulation preparations experienced a slight shift in the absorption peak at the amide I carbonyl group, namely around the number area 1642.46 cm^{-1} for the 1:1 formulation, 1640.53 cm^{-1} for the 1:0.9 formulation, 1638.60 cm^{-1} for the 1:0.8 formulation, and 1645.35 cm^{-1} for the 1:0.7 formulation. Furthermore, there is also a shift in N-H bending vibrations in the area of 1566.27 cm^{-1} for the 1:1 formulation, 1564.34 cm^{-1} for the 1:0.9 formulation, 1550.83 cm^{-1} for the 1:0 formulation 8, and 1563.37 cm^{-1} for the 1:0.7 formulation. Besides that, there was an absorption peak of the P-O group in the number area 917.19 cm^{-1} for the 1:1 formulation, 919.12 cm^{-1} for the 1:0.9 formulation, and 921.05 cm^{-1} for the 1:0.8 formulation. Next, the formulation 1:0.7 was located in the number area 922.98 cm^{-1} .




Figure 1. FTIR analysis results.

pH determination

The results of the pH normality test of encapsulated chitosan and NaTPP in trapping lemongrass leaf extract in a ratio of 1:1, 1:0.9, 1:0.8, and 1:0.7, with p value >0.05 . The normality test results were followed by a repeated ANOVA test to see whether there was a difference in pH values on the first day and the seventh day. The parametric test results showed no significant difference in pH values on the first and seventh days ($p > 0.05$). The pH test results can be seen in Table 3.

Encapsulation of chitosan and NaTPP in entrapping lemongrass leaf extract in a ratio of 1:1; 1:0.9; 1:0.8; and 1:0.7 on the first day of pH observation showed that the encapsulation group with a ratio of 1:0.8 was significantly different ($p < 0.05$) from the other encapsulation groups, which were shown by the lowest pH value, namely 5.54 ± 0.17 . On the seventh day, the pH of the 1:1 ratio encapsulation group was not significantly different from the 1:0.7 ratio with a p value of 0.06, but was significantly different from the 1:0.8 and 1:0.9 ratio encapsulation groups ($p < 0.05$). Each formulation has an increase and a decrease in the pH value. Encapsulation 1:1, 1:0.8, and 1:0.7 experienced an increase in pH on the seventh day. Meanwhile, encapsulation 1:0.9 saw a decrease in pH on the seventh day. Changes in pH values can be observed in Table 3.

Table 3. pH determination test.

Group (ratio chitosan : NaTPP)	pH level		pH of commercial mouthwash [11]	Description
	Day 1	Day 7		
Group 1 (1 : 1.0)	6.46 ± 0.14	7.30 ± 0.14	4.1–7.9	Eligible
Group 2 (1 : 0.9)	6.09 ± 0.13	5.47 ± 0.07	4.1–7.9	Eligible
Group 3 (1 : 0.8)	5.54 ± 0.17	5.97 ± 0.07	4.1–7.9	Eligible
Group 4 (1 : 0.7)	6.40 ± 0.17	6.53 ± 0.60	4.1–7.9	Eligible

Turbidity test

Observations on the turbidity test were carried out using visual observation to compare the level of turbidity of each dosage formulation over time. During this observation, the formulation preparations were placed in their respective containers and left to stand for 14 days. Observations were made on day 1, day 7, and day 14.

On the first day of observation, each dosage formulation was light yellow with clear turbidity. On the 7th day of observation, each formulation began to appear cloudy. Furthermore, on the 14th day, all preparations showed a light yellow color, but there was an increase in turbidity, as seen in Table 4.

Table 4. Turbidity test.

Group (ratio chitosan : NaTPP)	1		Day 7		14	
	Color	Turbidity	Color	Turbidity	Color	Turbidity
Group 1 (1 : 1.0)	Yellow	Clear	Yellow	Cloudy	Yellow	Cloudy
Group 2 (1 : 0.9)	Yellow	Clear	Yellow	Cloudy	Yellow	Cloudy
Group 3 (1 : 0.8)	Yellow	Clear	Yellow	Cloudy	Yellow	Cloudy
Group 4 (1 : 0.7)	Yellow	Clear	Yellow	Cloudy	Yellow	Cloudy

Viscosity test

Normality test results of the viscosity of chitosan and NaTPP encapsulation in entrapping kitchen lemongrass leaf extract in a 1:1, 1:0.9, 1:0.8, and 1:0.7 ratio shows a p value <0.05. Statistical testing was carried out non-parametrically using Friedman which showed a p value = 0.42. The viscosity test results did not show any differences between the encapsulation groups, were observed on the first, seventh, and 14th days.

The average results of the viscosity test on the first day between chitosan encapsulation and NaTPP with a ratio of 1:1 and 1:0.7 showed a significant difference with a p value = 0.04. Still, the 1:1 comparison group was not significantly different from the 1:0.8 and 1:0.9 comparison group. For days 7 and 14, there were no significant differences in all groups of the study. The average results of viscosity observations can be seen in Table 5.

On observation from the first day to day 14, the viscosity value of the chitosan encapsulation solution and NaTPP of lemongrass leaf extract 1:1 was relatively stable. The viscosity of the chitosan encapsulation solution and NaTPP of lemongrass leaf extract was 1:0.9 from the beginning of the observation until the 14th day, showing quite good stability. Encapsulation with a ratio of chitosan and NaTPP of lemongrass leaf extract of 1:0.8 showed that the viscosity value from the beginning of the observation until the 14th day saw a decrease in viscosity. Observation of the viscosity value at a ratio of 1:0.7 shows an increase in viscosity from the beginning of the observation to the 14th day. The results of viscosity testing in all comparison groups of chitosan and NaTPP show viscosity values within the standard mouthwash viscosity < 7.25.

Table 5. Viscosity test.

Group (ratio chitosan : NaTPP)	Viscosity level (mPa s)			Viscosity [12]	Description
	Day 1	Day 7	Day 14		
Group 1 (1 : 1.0)	2.169±0.04	2.138±0.01	2.134±0.01	< 7.25	Eligible
Group 2 (1 : 0.9)	2.106±0.01	2.118±0.01	2.118±0.01	< 7.25	Eligible
Group 3 (1 : 0.8)	2.132±0.05	2.120±0.02	2.116±0.02	< 7.25	Eligible
Group 4 (1 : 0.7)	2.081±0.02	2.127±0.01	2.165±0.11	< 7.25	Eligible

The research results showed that all comparison groups of chitosan and NaTPP encapsulation met the viscosity requirements for a mouthwash. A 1:1 ratio of chitosan and NaTPP had the best encapsulation physical characteristics with an even particle size distribution (monodisperse), so it met the requirements as a drug particle in the pharmaceutical field.

• DISCUSSION

Encapsulation is a method used to package a compound within a material wall called a coating. This method offers advantages such as the stability of an active ingredient and increasing the safety of the material; as a result, the release of the compound is easier [13]. Encapsulation can be contained in several forms, such as capsules, emulsions, and particles [14]-[18].

This research used chitosan encapsulation to coat the active compounds of lemongrass leaves obtained from the extraction process using 70% ethanol solvent. An ethanol solvent concentration of 70% showed better antioxidant activity through the DPPH test compared to 50% and 90% ethanol solvent concentrations [19],[20]. However, research by Adiningsih *et al.* showed that compounds with 70% ethanol extract had antibacterial activity that was not much different with compounds extracted using 96% ethanol [21],[22]. This difference can occur because the 70% solvent is able to extract a large number of antioxidant compounds, such as flavonoids and phenolics, from plants. This result is proven by research by Nichitai *et al.*, which states that 70% ethanol extract is able to attract more compounds, but 50% ethanol extract has better antibacterial activity [23]. In addition, research by Rahayu *et al.* states that 96% ethanol extract of roots *Zingiber zerumbet* has the best antibacterial activity against *S. enteritidis* and *S. typhimurium* bacteria compared with concentrations of 45% and 70% [24].

The results of particle measurements in encapsulated preparations showed that the particle size in preparations with a chitosan and NaTPP ratio of 1:1 reached 524.0 nm, smaller than the other comparison groups, namely 1:0.9 (614.5 nm), 1:0.8 (934.2 nm), and 1:0.7 (1145 nm). A particle is said to be nano-sized if the particle has a size between 1 and 1000 nm [6,16]. In addition, of the four preparation ratios, the ratio of chitosan and NaTPP is 1:1, 1:0.9, and 1:0.8 can be classified in the nano size category, while compounds with a ratio of 1:0.7 do not meet the requirements as nanoparticles. In Khoerunnisa's research, the 1:1 ratio variation is a synthesis with optimum characteristics. Variations in the ratio of chitosan: NaTPP are different from research conducted by Dwiki *et al.*, which reported an increase in size with each increase in the ratio of chitosan: NaTPP for bay leaves (*Syzygium polyanthum*), namely 1:1 (284 nm) and 5:1 (410 nm).) and 10:1 (630 nm). This results is different from research conducted by Rizal *et al.*, who reported that there was a parabolic relationship in variations in the chitosan-NaTPP composition of jackfruit leaves (*Artocarpus heterophyllus* Lam.), namely 1:1 (382.6 nm), 2:1 (199.2 nm), and 3:1 (402.2 nm). These results explain that variations and formulations of the chitosan : NaTPP ratio can provide significant differences in particle size [25].

Particle distribution testing can use the Polydispersity Index (Pdl) approach. Particles with a Pdl value <0.7 can be said to be monodisperse, and a Pdl value > 0.7 indicates that they are polydisperse particles [26]. In addition, monodisperse particles can also be considered good because they are located in the middle of the polydispersity index range. The average value of the index Pdl in encapsulation with a ratio of chitosan and NaTPP, namely 1:1, 1:0.9, and 1:0.8, shows a Pdl figure <0.7, respectively 0.481, 0.695, and 0.687. This results shows that the three preparations have monodisperse character. Monodispersion of a nanoparticle means that there is uniformity in the size of the particles formed [27],[28]. This can provide benefits through a greater opportunity to interact with the biological target [29]. Furthermore, encapsulation preparations with a ratio of chitosan and NaTPP 1:0.7 have a value of 0.747, which indicates that the sample is polydisperse, that is, the comparative particles have parts with a wide particle size and have particles with a large size [28].

The presence of active compounds in the formulation preparation can be analyzed to assess the characteristics of the compound's functional groups [30]. In this case, to confirm that the chitosan nanoparticles do not decompose during the encapsulation preparation process, they can be analyzed using an FTIR approach. In the encapsulated preparation, there is an O-H absorption area, which looks steep and wide; this indicates a mixture of O-H from chitosan, NaTPP, and 70% ethanol extract of lemongrass leaves. From the observation process, it was seen that C=O absorption was in the area of 1638.60 – 1645.35 cm⁻¹ for each formulation, where there was a shift from the area of 1627.99 cm⁻¹ of pure chitosan extract. This shift indicates an interaction between chitosan and the 70% ethanol extract of lemongrass leaves. It is a sign that the chitosan compound is still available in each encapsulation formulation [8], [30]-[32]. Apart from that, there is also a shift in the bending N-H absorption area from 1579.77 cm⁻¹ to a number area ranging between 1550.83-1566.27 cm⁻¹ for each encapsulation formulation. This shift indicates that cross-linking has occurred between the chitosan compound, NaTPP, and the 70% ethanol extract of lemongrass leaves [8],[31],[32]. Furthermore, the

absorption peak of the P-O group was obtained in the area of 917.19 - 922.98 cm^{-1} for the encapsulation formulation, which indicates the P-O aliphatic group of NaTPP [30],[33].

When observing the turbidity level for 14 days, it was found that changes occurred in each dosage formulation [34]-[37]. On the 7th day, minimal changes occurred. This is slightly different from research conducted by Syarmalina *et al.*, which reported the stability of the turbidity level in the temu lawak extract nanoparticle formulation, which was observed for 5 days [38]. This change could occur due to several background factors, such as nanoparticle size, size distribution, and concentration [39].

For mouthwash, the viscosity component is necessary to review the comfort of the solution when consumed [33],[40]. This is because the closer the viscosity value of a formulation is to the viscosity of water (1 cP), the easier and more comfortable the formulation is to consume [33],[41]. The smaller the viscosity value, the more dilute the solution; conversely, the greater the viscosity value, the thicker the solution [42]. In this study, the viscosity value of the chitosan encapsulation solution and NaTPP leaf extract on day 14 ranged from 2.116 - 2.165 cP, the highest viscosity value was found by the formulation with a ratio of 1:0.7 and the lowest was owned by the formulation with a ratio of 1:0.8. Formulations with a ratio of 1:1 and 1:0.9 were relatively stable during the first to 14th day of observation. This result is different from the encapsulation formulation with a chitosan and NaTPP ratio of lemongrass leaf extract of 1:0.8, which shows a decrease in viscosity, and a formulation of 1:0.7, which shows an increase in viscosity. This viscosity instability can be caused by the inhomogeneity of the particles in the formulation [33]. Apart from that, the solution is easy to decompose by temperature and light. Generally the viscosity of the liquid will decrease with increasing temperature [34]. In addition, the viscosity values of the four formulations are still below standard (<7.25), so it still meets the standard mouthwash viscosity value requirements.

Thus, the encapsulation ratio 1:1 is the most optimal choice among other ratios. This preparation has a nanoparticle size with a uniform distribution (monodispersion), making it easier for this compound to interact with its biological target. In addition, the stability and good viscosity of this preparation also support these advantages, creating an optimal system for applications in the medical field. Overall, this preparation offers a potential solution for therapeutic development in the context of mouthwash.

CONCLUSION

The physical characteristics of the encapsulation of kitchen lemongrass leaf extract for mouthwash with several comparisons of chitosan and NaTPP have good particle size, pH value, turbidity, and viscosity, as well as cross-linking between the chitosan compound, NaTPP, and 70% ethanol extract of lemongrass leaves, which was confirmed through the FT-IR test.

Acknowledgements: This work was supported by Department of Dental Material Science and Technology and Department of Oral Biology, Faculty of Dentistry, Universitas Trisakti.

Author contributions: Concept - F.L.K., K., O.; Design - F.L.K., K.; Supervision - F.L.K., K.; Resources - F.L.K., K., F.A.J.K.; Materials - F.L.K., K., F.A.J.K.; Data Collection and/or Processing - F.L.K., K., O., F.A.J.K., F.R.A.; Analysis and/or Interpretation - F.L.K., K., F.A.J.K., F.R.A.; Literature Search - F.L.K., K., F.A.J.K.; Writing - F.L.K., K., F.A.J.K.; Critical Reviews - F.L.K., K., O.

Conflict of interest statement: The authors declared no conflict of interest.

REFERENCES

- [1] G. Saputra, Karakteristik nanoenkapsulasi kitosan etanol 70% daun sirih (*Piper betle* Linn) dengan metode gelasi ionik, Naskah Publikasi, vol. 12, no. 1, pp. 579-587, 2016.
- [2] Y. U. Salami, H. Jusuf, and L. N. S. Ayini, Pengaruh perasan daun serai dapur (*Cymbopogon citratus*) terhadap kematian larva *Aedes aegypti*, Jambura Journal of Health Sciences and Research, vol. 2, no. 1, pp. 32-42, 2020.
- [3] B. P. Brügger, L. C. Martínez, A. Plata-Rueda, B. M. de C. Castro, M. A. Soares, C. F. Wilcken, et al., Bioactivity of the *Cymbopogon citratus* (Poaceae) essential oil and its terpenoid constituents on the predatory bug, *Podisus nigripinus* (Heteroptera: Pentatomidae), Scientific Reports, vol. 9, no. 1, p. 1, 2019.

- [4] K. Komariah, W. Poedjiastoeti, R. Amtha, S. Lestari, E. Fibryanto, A. D. P. Roberto, et al., Pemanfaatan limbah serai dapur sebagai obat kumur dan *hand sanitizer* untuk meningkatkan pemberdayaan ekonomi, Abdi Moestopo: Jurnal Pengabdian Pada Masyarakat, vol. 5, no. 1, pp. 74–82, 2022.
- [5] M. Pateiro, B. Gómez, P. E. S. Munekata, F. J. Barba, P. Putnik, D. B. Kovačević, et al., Nanoencapsulation of promising bioactive compounds to improve their absorption, stability, functionality and the appearance of the final food products, *Molecules*, vol. 26, no. 6, pp. 1–26, 2021.
- [6] Y. Juliantoni, W. Hajrin, and W. A. Subaidah, Nanoparticle formula optimization of juwet seeds extract (*Syzygium cumini*) using simplex lattice design method, *Jurnal Biologi Tropis*, vol. 20, no. 3, pp. 416–422, 2020.
- [7] E. T. Webber and M. J. Pashuck, (Macro) molecular self-assembly for hydrogel drug delivery, *Advanced Drug Delivery Reviews*, vol. 172, pp. 275–295, 2021.
- [8] A. Indriani Putri, A. Sundaryono, and I. Nyoman Candra, Karakterisasi nanopartikel kitosan ekstrak daun ubi jalar (*Ipomoea batatas L.*) menggunakan metode gelasi ionik, *Jurnal Pendidikan Dan Ilmu Kimia*, vol. 2, no. 2, pp. 203–207, 2018.
- [9] V. Purwandari, I. Isaeni, R. Rahmi, A. Z. Akbari, and M. Z. Akbari, Formulasi nanoenkapsulasi ekstrak daun kelor (*Moringa oleifera*) /Kitosan-Natrium Tripolifosfat (NaTPP), *Journal of Science and Applicative Technology*, vol. 6, no. 2, p. 77, 2022.
- [10] U. Chasanah, N. Mahmintari, F. Hidayah, F. A. E. Maghfiroh, D. Rahmasari, and R. W. Nugraheni, Thin-layer hydration method to prepare a green tea extract niosomal gel and its antioxidant performance, *European Pharmaceutical Journal*, vol. 68, no. 1, pp. 126–136, 2021.
- [11] B. W. M. Van Swaaij, D. E. Slot, G. A. Van Der Weijden, M. F. Timmerman, and J. M. Ruben, Fluoride, pH value, and titratable acidity of commercially available mouthwashes, *International Dental Journal*, Dec. 2023. [Online]. Available: <https://doi.org/10.1016/j.identj.2023.09.002>
- [12] Departemen Kesehatan Republik Indonesia, Farmakope Indonesia, Edisi VI. Jakarta: Depkes RI, 2020.
- [13] A. Putri, M. S. Rusli, and D. Setyaningsih, Enkapsulasi campuran minyak atsiri sebagai produk sediaan aromaterapi dengan teknik koaservasi kompleks, *Jurnal Teknologi Industri Pertanian*, vol. 30, no. 3, pp. 299–307, 2020.
- [14] I. Kłodová, T. Milota, J. Smetanová, and C. Stathopoulos, Encapsulation: A strategy to deliver therapeutics and bioactive compounds?, *Pharmaceuticals*, vol. 16, no. 3, pp. 2–19, 2023.
- [15] D. O. Z. Hulwah, J. Bobsaid, M. Ramadhani, and Y. Setiawati, Efektivitas mouthwash berbahan dasar ekstrak *Camellia sinensis* dan *Mentha piperita* sebagai antibakteri terhadap *Streptococcus mutans*, *Jurnal Kedokteran Meditek*, vol. 28, no. 1, pp. 30–39, 2022.
- [16] D. Fitri, N. Z. W. Kiromah, and T. C. Widiastuti, Formulasi dan karakterisasi nanopartikel ekstrak etanol daun salam (*Syzygium polyanthum*) pada berbagai variasi komposisi kitosan dengan metode gelasi ionik, *JPSCR: Journal of Pharmaceutical Science and Clinical Research*, vol. 5, no. 1, p. 61, 2020.
- [17] R. Indraswary, R. Amalina, and A. Firmansyah, Effects of nano chitosan mouth spray on the epithelial thickness in the traumatic ulcer healing process (In vivo), *MEDALI Journal*, vol. 4, no. 1, pp. 95–103, 2022.
- [18] G. N. Sumeisey, S. D. Umboh, and T. E. Tallei, Penyalutan bakteri asam laktat menggunakan nanopartikel kitosan, *Pharmakon*, vol. 8, no. 4, pp. 843–850, 2019.
- [19] G. G. Kusumo and D. Dipahayu, The effect of nano extract formulation on antioxidant activity of 70% ethanolic extract of purple sweet potato's leaves (*Ipomoea batatas L.*) antin-3 variety, *Journal Pharmasci (Journal of Pharmacy and Science)*, vol. 7, no. 1, pp. 43–47, 2022.
- [20] M. Kozłowska, I. Ścibisz, J. L. Przybył, A. E. Laudy, E. Majewska, K. Tarnowska, et al, Antioxidant and antibacterial activity of extracts from selected plant material, *Applied Sciences (Switzerland)*, vol. 12, no. 19, p. 9871, 2022.
- [21] W. Adiningsih, R. L. Vifta, R. Yuswantina, Studi PS, dan F. I. Kesehatan, Uji aktivitas antibakteri ekstrak etanol 70% dan ekstrak etanol 96% buah strawberry (*Fragaria x ananassa*) terhadap bakteri *Propionibacterium acnes*, *Generics: Journal of Research in Pharmacy*, vol. 1, no. 1, pp. 1–8, 2021.
- [22] Noviyanti, Pengaruh kepolaran pelarut terhadap aktivitas antioksidan ekstrak etanol daun jambu brazil batu (*Psidium guineense L.*) dengan metode DPPH, *Jurnal Farmako Bahari*, vol. 7, no. 1, pp. 29–35, 2016.
- [23] M. M. Nichitai, A. M. Josceanu, R. D. Isopescu, G. O. Isopencu, E. I. Geana, C. T. Ciucure, et al., Polyphenolics profile effects upon the antioxidant and antimicrobial activity of propolis extracts, *Scientific Reports*, vol. 11, no. 1, 2021.

- [24] I. D. Rahayu, W. Widodo, I. Prihartini, dan A. Winaya, Antibacterial activity of ethanolic extracts from Zingiber zerumbet rhizome against *Salmonella spp.*, Biodiversitas, vol. 20, no. 11, pp. 3322–3327, 2019.
- [25] R. S. Triatmojo, Pembuatan dan karakterisasi nanopartikel ekstrak etanol daun nangka (*Artocarpus Heterophyllus Lam.*) pada berbagai variasi komposisi kitosan-natrium tripolifosfat, 2018.
- [26] R. Ambarwati dan Y. Yulianita, Physical evaluation of transfersome that contains pandan leaves extract (*Pandanus amaryllifolius R.*), Journal of Tropical Pharmacy and Chemistry, vol. 5, no. 4, pp. 369–374, 2021.
- [27] F. Livia, R. Tjandrawinata, C. D. Marpaung, D. Pratiwi, dan Komariah, The effect of horn beetle nano chitosan (*Xylotrupes gideon*) on the surface roughness of glass-ionomer cement, Materials Science Forum, vol. 1069, pp. 161–166, 2022, doi: 10.4028/p-5mk858.
- [28] V. Nurviana, Potensi Antidioksidan Sediaan Nanopartikel Ekstrak Kernel Biji Limus (*Mangifera foetida Lour.*), Jurnal Farmasi Udayana, vol. 3, no. 2, pp. 144–151, 2020.
- [29] A. Masykuroh and H. Puspasari, Aktivitas anti bakteri nano partikel perak (NPP) hasil biosintesis menggunakan ekstrak keladi sarawak *Alocasia macrorrhizos* terhadap *Staphylococcus aureus* dan *Escherichia coli*, BIOMA: Jurnal Biologi Makassar, vol. 7, no. 1, pp. 76–85, 2022.
- [30] I. Antasionasti, I. Jayanto, S. S. Abdullah, and J. P. Siampa, “Karakterisasi nanopartikel ekstrak etanol kayu manis (*Cinnamomum burmannii*) dengan kitosan sodium tripolifosfat sebagai kandidat antioksidan, CHEMISTRY PROGRESS, vol. 13, no. 2, pp. 77–85, 2020.
- [31] A. I. Putri, A. Sundaryono, and I. N. Candra, Karakterisasi nanopartikel kitosan ekstrak daun ubi jalar (*Ipomoea batatas L.*) menggunakan metode gelasi ionik, Jurnal Pendidikan dan Ilmu Kimia, vol. 2, no. 2, pp. 203–207, 2018.
- [32] G. Morales-Olán, S. Luna-Suárez, J. D. D. Figueroa-Cárdenas, M. Corea, and M. Rojas-López, Synthesis and Characterization of Chitosan Particles Loaded with Antioxidants Extracted from Chia (*Salvia hispanica L.*) Seeds, Int. J. Anal. Chem., vol. 2021, 2021.
- [33] Noval, Melviani, Novia, and D. Syahrina, Formulasi dan evaluasi sediaan obat kumur (mouthwash) dari ekstrak etanol tanaman bundung (*Actinoscirpus grossus*) sebagai antiseptik mulut, Jurnal Surya Medika, vol. 6, no. 2, pp. 112–120, 2020.
- [34] Ririn, A. I. Tandjung, and S. Wagola, Formulasi sediaan mouthwash dari sari buah sirih (*Piper betle L.*) varietas siriboah, As-Syifaa, vol. 5, no. 2, pp. 153–161, 2013.
- [35] A. U. Yap, F. Kurniawan, Y. Pragustine, and C. Marpaung, Temporomandibular Disorder and Somatic Symptoms: Relations to Fear of Missing Out and Other Negative Emotional States, Acta Odontol. Scand., vol. 83, pp. 340–347, 2024, doi: 10.2340/aos.v83.40776.
- [36] Gusnawati, Z. Sabara, Munira, and S. Bakhri, Karakterisasi mutu pasta gigi dengan penambahan garam dan Virgin Coconut Oil (VCO) ditinjau dari SNI 12-3524-1995, Jurnal Industri Hasil Perkebunan, vol. 17, no. 1, pp. 41–49, 2022.
- [37] Syarmalina, D. Wirawan, and D. Rahmat, Formulasi nanopartikel ekstrak temu lawak berbasis kitosan sebagai antijerawat, Medical Sains, vol. 3, no. 2, pp. 153–158, 2019.
- [38] F. Livia, R. Tjandrawinata, C. D. Marpaung, D. Pratiwi, and K. Komariah, The Effect of Horn Beetle Nano Chitosan (*Xylotrupes gideon*) on the Hardness of Glass-Ionomer Cement, J. Indones. Dent. Assoc., vol. 5, no. 1, pp. 27–31, 2022, doi: 10.32793/jida.v5i1.719.
- [39] N. Noval, R. Rosyifa, and A. Annisa, Effect of HPMC Concentration Variation as Gelling Agent on Physical Stability of Formulation Gel Ethanol Extract Bundung Plants (*Actinoscirpus Grossus*), in European Alliance for Innovation, 2020.
- [40] F. Handayani, R. Sundu, and R. M. Sari, Formulasi dan Uji Aktivitas Antibakteri Streptococcus mutans dari Sediaan Mouthwash Ekstrak Daun Jambu Biji (*Psidium guajava L.*), Jurnal Sains dan Kesehatan, vol. 1, no. 8, pp. 422–433, 2017.
- [41] D. T. Arya et al., Sintesis nanopartikel zirkonium dioksida (ZrO_2) dengan metode sol gel menggunakan ekstrak nanas sebagai pengkelat untuk aplikasi nanofluida pendingin, PILLAR OF PHYSICS, vol. 8, pp. 25–32, 2016.
- [42] N. Noval, F. Ferlina, and K. B. Indartantri, Formulasi dan evaluasi suspensi ubi cilembu dengan menggunakan perbandingan suspending agent antara NA CMC dan XANTHAN GUM, 2019. [Online]. Available: <https://api.semanticscholar.org/CorpusID:214479893>