

Sources

Title



Enter title

Find sources

Title: [Journal Of Advanced Pharmaceutical Technology And Research](#) x

CiteScore 2023 has been released. [View CiteScore methodology](#) >



Filter refine list

Apply

Clear filters

Display options

Display only Open Access journals

Counts for 4-year timeframe

No minimum selected

Minimum citations

Minimum documents

Citescore highest quartile

Show only titles in top 10 percent

1st quartile

2nd quartile

3rd quartile

4th quartile

Source type

Journals

Book Series

Conference Proceedings

Trade Publications

1 result

[Download Scopus Source List](#) [Learn more about Scopus Source List](#)

All v

[Export to Excel](#)

[Save to source list](#)

View metrics for year: 2023



Source title ↓	CiteScore ↓	Highest percentile ↓	Citations 2020-23 ↓	Documents 2020-23 ↓	% Cited ↓
<input type="checkbox"/> 1 Journal of Advanced Pharmaceutical Technology and Research <i>Open Access</i>	2.0	43% 104/183 Pharmaceutical Science	706	348	53



[Top of page](#)



Source details

Journal of Advanced Pharmaceutical Technology and Research

Open Access ⓘ

Years currently covered by Scopus: from 2010 to 2024

Publisher: Wolters Kluwer Health

ISSN: 2231-4040 E-ISSN: 0976-2094

Subject area: Pharmacology, Toxicology and Pharmaceutics: Pharmaceutical Science

Source type: Journal

CiteScore 2023

2.0 ⓘ

SJR 2023

0.279 ⓘ

SNIP 2023

0.473 ⓘ

[View all documents >](#)

[Set document alert](#)

[Save to source list](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

CiteScore 2023 ▾

$$2.0 = \frac{706 \text{ Citations } 2020 - 2023}{348 \text{ Documents } 2020 - 2023}$$

Calculated on 05 May, 2024

CiteScoreTracker 2024 ⓘ

$$2.2 = \frac{738 \text{ Citations to date}}{330 \text{ Documents to date}}$$

Last updated on 08 July, 2024 • Updated monthly

CiteScore rank 2023 ⓘ

Category	Rank	Percentile
Pharmacology, Toxicology and Pharmaceutics	#104/183	43rd
Pharmaceutical Science		

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site ↗](#)

About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

ELSEVIER

[Terms and conditions ↗](#) [Privacy policy ↗](#)

All content on this site: Copyright © 2024 Elsevier B.V. ↗, its licensors, and contributors. All rights are reserved, including those for text and data mining, AI training, and similar technologies. For all open access content, the Creative Commons licensing terms apply. We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies ↗.



ISSN 2231-4040

Journal of Advanced Pharmaceutical Technology & Research

An Official Publication of Society of Pharmaceutical Education & Research

Vol 2 / Issue 4 / Oct-Dec 2011

www.japtr.org

JAPTR



Medknow
Publications

Editorial Board : Journal of Advanced Pharmaceutical Technology & Research

Advertisement

Editorial Board

Editor-in-Chief

Dr. Upendra Nagaich
M.Pharm., Ph.D., FSPER
Professor & Research Coordinator,
Amity Institute of Pharmacy, Amity University,
Noida (U.P.) India
& National Secretary, Society of Pharmaceutical Education & Research (SPER), India

Executive Editors

Dr. Paul Wan Sia Heng
Department of Pharmacy,
National University of Singapore,
Singapore

Dr. Jaya Pandey
School of Studies in Pharmaceutical Sciences,
Jiwaji University,
Gwalior (M.P.) India

Associate Editor

Dr. Vandana Chaudhary
School of Pharmacy,
Pharmacy Australia Centre of Excellence,
The University of Queensland,
Brisbane, Australia

Consultant Editor

Prof. (Dr.) Laszlo Endrenyi
Professor Emeritus
Department of Pharmacology & Toxicology
University of Toronto, Toronto, Canada

Section Editors (Alphabetical Order)

Dr. Ikoni Ogaji
Dept. of Pharmaceutics & Pharmaceutical Technology,
Faculty of Pharmaceutical Sciences,
University of Jos, Nigeria

Dr. Ke Li
J. David Gladstone Institutes,
San Francisco, California, USA

Editorial Board Members (In Alphabetical Order)

Prof. (Dr.) A. N. Kalia
Vice-President, Society of Pharmacognosy,
Professor & Head
I.S.F. College of Pharmacy,
Moga (Punjab), India

Dr. Abdul Wahab
Department of Pharmacy
Kohat University of Science & Technology
Kohat, KPK, Pakistan

Prof. (Dr.) Arun Dharmarajan
Professor, School of Biomedical Sciences,
Faculty of Health Sciences,
Curtin University,
Perth, Australia

Prof. (Dr.) Ashoke Kumar Ghosh
Vice-Chancellor, IFTM University,
Moradabad (Uttar Pradesh), India

Prof. (Dr.) B. Mishra
Professor & Head
Department of Pharmaceutics, Indian Institute of Technology,
Banaras Hindu University,
Varanasi (Uttar Pradesh), India

Prof. B. G. Shivananda
Secretary, APTI
Bangalore (Karnataka), India

Prof. (Dr.) Bharat B. Aggarwal
Professor of Cancer Medicine & Chief, Cytokine Research Laboratory
The University of Texas
Texas, USA

Dr. Bhaswat Sundar Chakraborty
Senior VP & Chairman, R&D Core Committee
Cadila Pharmaceuticals Ltd.
Ahmedabad (Gujarat), India

Prof. (Dr.) C. K. Kokate
National Advisor, Society of Pharmacognosy, Vice-Chancellor, KLE University,
Belgaum (Karnataka), India

Prof. (Dr.) Cedric Pearce
Adjunct Professor
University of North Carolina, CEO, Mycosynthetix Inc.
North Carolina, USA
[Associate Editor, Journal of Natural Products]

Prof. (Dr.) Ciddi Veeresham
Professor, University College of Pharmaceutical Sciences,
Kakatiya University,
Warangal (Andhra Pradesh), India

Dr. Dâmaris Silveira

Universidade de Brasilia
Brasilia, Brazil

Prof. (Dr.) Dilipkumar Pal
Professor, Dept. of Pharmaceutical Sciences,
Guru Ghasidas Vishwavidyalaya,
Bilaspur (Chhattisgarh), India

Prof. (Dr.) Divakar Goli
Associate Secretary, IPA
Secretary, Education Division, IPA

Professor & Principal
Acharya & B.M. Reddy College of Pharmacy,
Bangalore (Karnataka), India

Prof. (Dr.) G. K. Dash
Professor, Universiti Kuala Lumpur
Perak Darul Ridzuan, Malaysia

Prof. (Dr.) H. S. Buttar
Adjunct Professor,
School of Medicine,
University of Ottawa,
Ottawa, Canada

Kaushik Desai
Gen. Secretary, IPA
Immediate Vice President, IPA
Mumbai (Maharashtra), India

Prof. (Dr.) Mainul Haque
Professor, Universiti Sultan Zainal Abidin
Kuala Terengganu, Terengganu, Malaysia

Dr. Muhammad Akhlaq
University of Auckland
Auckland, New Zealand

Prof. (Dr.) N. Udupa
Professor & Research Director [Health Sciences]
Manipal University,
Manipal (Karnataka), India

Prof. (Dr.) Pornsak Sriamornsak
Professor, Department of Pharmaceutical Technology &
Pharmaceutical Biopolymer Group (PBiG),
Faculty of Pharmacy, Silpakorn University,
Nakhon Pathom, Thailand

Dr. Rakesh Kumar Sharma
Scientist 'F', Director,
Defence Food Research Laboratory (DFRL),
Mysuru (Karnataka), India

Prof. (Dr.) Roop K. Khar
Former Professor, Department of Pharmaceutics,
Jamia Hamdard (Hamdard University)

New Delhi, India

Prof. (Dr.) Roopa Karki
Professor & Head Department of Pharmaceutics,
Acharya & BM Reddy College of Pharmacy,
Bangalore (Karnataka), India
Prof. (Dr.) S. D. Tonpay
Former Professor & Head, Govt. Gajra Raja Medical College,
Gwalior (Madhya Pradesh), India

Prof. (Dr.) S. H. Ansari
Professor & Head,
Department of Pharmacognosy & Phytochemistry,
Jamia Hamdard (Hamdard University)
New Delhi, India

Prof. (Dr.) S. H. Mishra
Professor Emeritus, Faculty of Pharmacy,
The M. S. University of Baroda,
Vadodara (Gujarat), India

Prof. (Dr.) Satish Sardana
Immediate Past President, APTI (Haryana)
Director,
Amity Institute of Pharmacy, Amity University,
Manesar (Haryana), India

Prof. (Dr.) T. V. Narayana
President, IPA, Director
Vikas Institute of Pharmaceutical Sciences
Rajahmundry (Andhra Pradesh), India

Prof. (Dr.) V. K. Dixit
President, Society of Pharmacognosy,
Former Professor, Department of Pharmaceutical Sciences,
Dr. H. S. Gour University,
Sagar (M.P.), India

Prof. (Dr.) Y. K. Agrawal
Director, Institute of Research & Development,
Gujarat Forensic Sciences University,
Gandhinagar (Gujarat), India

Advertisement

[^Back to Top](#)



Never Miss an Issue

Get new journal Tables of Contents sent right to your email inbox

[Get New Issue Alerts](#)

Browse Journal Content

- [Register on the website](#)
- [Get eTOC Alerts](#)

Customer Service

- Contact us at:
 - Support: [Submit a Service Request](#)
 - TEL: 800-638-3030 (within the USA)
301-223-2300 (outside of the USA)

[Manage Cookie Preferences](#)

- [Privacy Policy](#)
- [Legal Disclaimer](#)
- [Terms of Use](#)
- [Open Access Policy](#)
- [Your California Privacy Choices](#)



- Copyright © 2024
- Journal of Advanced Pharmaceutical Technology & Research | Published by Wolters Kluwer - [Medknow](#)

Advertisement

[Close](#)

Current Issue : Journal of Advanced Pharmaceutical Technology & Research

[Previous Issue](#)



Jul-Sep 2024 - Volume 15 - Issue 3

pp: 135-241

[Table of Contents Outline](#)

[eTOC Alerts](#)

[Contributor Index](#)

Advertisement

Review Article

[Gene polymorphism impact on opioid analgesic usage](#)

Widjaja, Sry Suryani; Ichwan, Muhammad; Chowbay, Balram; More

Journal of Advanced Pharmaceutical Technology & Research. 15(3):135-138, Jul-Sep 2024.

- [Abstract](#)
-
-
- [Permissions](#)
- [Open](#)

[Bioadhesive polymer in antifungal drug delivery for therapeutic treatment of candidiasis](#)

Roslan, Muhamad Naquib Faisal; Naharudin, Idanawati; Musa, Nafisah; More

Journal of Advanced Pharmaceutical Technology & Research. 15(3):139-143, Jul-Sep 2024.

- [Abstract](#)
- [Favorite](#)
- [PDF](#)
- [Permissions](#)

- [Open](#)

[Table of Contents Outline](#) | [Back to Top](#)

Original Article

[Botanical and pharmacognostic investigation of *Strobilanthes kalimantanensis*](#)

Prabowo, Wisnu Cahyo; Kuncoro, Hadi; Irawan, Budi; More

Journal of Advanced Pharmaceutical Technology & Research. 15(3):144-149, Jul-Sep 2024.

- [Abstract](#)
- [Favorite](#)
- [PDF](#)
- [Permissions](#)

- [Open](#)

[Genetic analysis: therapeutic drug monitoring of metformin and glimepiride on diabetic patients' plasma including genetic polymorphism](#)

Ibrahim, Areen; Odeh, Mohanad; Mallah, Eyad; More

Journal of Advanced Pharmaceutical Technology & Research. 15(3):150-155, Jul-Sep 2024.

- [Abstract](#)
- [Favorite](#)
- [PDF](#)
- [Permissions](#)

- [Open](#)

[Inhibitory effects against zoonotic bacteria by *Oroxylum indicum* ointment and effects to dog wound](#)

Sithisarn, Patchima; Rojsanga, Piyanuch; Ruenraroengsak, Pakatip; More

Journal of Advanced Pharmaceutical Technology & Research. 15(3):156-160, Jul-Sep 2024.

- [Abstract](#)
- [Favorite](#)
- [PDF](#)
- [Permissions](#)
- [Open](#)

[**Comparing monotherapy with tadalafil or tamsulosin in men with benign prostatic hyperplasia: A case-control study**](#)

Fawzi, Hayder Adnan; Akram, Saif Nabeel; Talib, Aya Fawzi; More

Journal of Advanced Pharmaceutical Technology & Research. 15(3):161-165, Jul-Sep 2024.

- [Abstract](#)
- [Favorite](#)
- [PDF](#)
- [Permissions](#)
- [Open](#)

[**Exploring the antioxidant potential of *Moringa oleifera* leaf extracts mitigating doxorubicin-induced cardiotoxicity in male rats**](#)

Obeid, Ruaa Ali; Mohammed, Rusul Ahmed; Kaskoos, Raad A.; More

Journal of Advanced Pharmaceutical Technology & Research. 15(3):166-170, Jul-Sep 2024.

- [Abstract](#)
- [Favorite](#)
- [PDF](#)
- [Permissions](#)
- [Open](#)

[**Potential antioxidant and antiradical agents from *Allium ascalonicum*: Superoxide dismutase and density functional theory *in silico* studies**](#)

Ajiati, Dwipa; Sumiarsa, Dadan; Amin, Meiny Faudah; More

Journal of Advanced Pharmaceutical Technology & Research. 15(3):171-176, Jul-Sep 2024.

- [Abstract](#)
- [Favorite](#)
- [PDF](#)
- [Permissions](#)
- [Open](#)

[**Potential target and mechanism exploration from \$\alpha\$ -mangostin against triple-negative breast cancer: An *in silico* study**](#)

Rivaldo, Rafly Mochamad; Chandra, Paulus

Journal of Advanced Pharmaceutical Technology & Research. 15(3):177-184, Jul-Sep 2024.

- [Abstract](#)
- [Favorite](#)
- [PDF](#)
- [Permissions](#)

- [Open](#)

[Evaluation of the methanol extract of BaiYangJie: Toxicology and protective effect against acute kidney injury](#)

Li, Jiaxin; Zhou, Lingjuan; Wang, Jinhui; More

Journal of Advanced Pharmaceutical Technology & Research. 15(3):185-193, Jul-Sep 2024.

- [Abstract](#)
- [Favorite](#)
- [PDF](#)
- [Permissions](#)

- [Open](#)

[Effect of hesperidin in the prevention of aluminum chloride-induced testicular dysfunction in rats](#)

Parasuraman, Subramani; Huey, Loh Yee; Jee, Yap Shen

Journal of Advanced Pharmaceutical Technology & Research. 15(3):194-199, Jul-Sep 2024.

- [Abstract](#)
- [Favorite](#)
- [PDF](#)
- [Permissions](#)

- [Open](#)

[Role of *Moringa oleifera* irrigation solution on the cell metabolism change of *Streptococcus mutans*](#)

Soraya, Cut; Batubara, Fitri Yunita; Nasroen, Saskia L.; More

Journal of Advanced Pharmaceutical Technology & Research. 15(3):200-207, Jul-Sep 2024.

- [Abstract](#)
- [Favorite](#)
- [PDF](#)
- [Permissions](#)

- [Open](#)

[Nephroprotective effect of sugarcane \(*Saccharum officinarum* L.\) leaves ethanol extract on gentamicin-induced nephrotoxicity in rats](#)

Dewi, Ika Puspita; Aldiana, Merinda; Viadina, Zildjian Adela; More

Journal of Advanced Pharmaceutical Technology & Research. 15(3):208-213, Jul-Sep 2024.

- [Abstract](#)
-
-
- [Permissions](#)

- [Open](#)

[New method to induce neurotrophin gene expression in human adipose-derived stem cells *in vitro*](#)

Altememy, Dhiya; Kashani, Maryam Haji Ghasem; Fateme, Amirahmadi; More

Journal of Advanced Pharmaceutical Technology & Research. 15(3):214-219, Jul-Sep 2024.

- [Abstract](#)
-
-
- [Permissions](#)

- [Open](#)

[Physical characteristics of quercetin pulmospheres using combination of alginate-carrageenan: Effect of polymer concentration](#)

Hariyadi, Dewi Melani; Fawwaz, Lubby Razan; Fattah, Abdul; More

Journal of Advanced Pharmaceutical Technology & Research. 15(3):220-224, Jul-Sep 2024.

- [Abstract](#)
-
-
- [Permissions](#)

- [Open](#)

[Genetic polymorphism and immunological evaluation of PD-1 in Iraqi patients with acute myeloid leukemia](#)

Mahmood, Aseel S.

Journal of Advanced Pharmaceutical Technology & Research. 15(3):225-230, Jul-Sep 2024.

- [Abstract](#)
-
-
- [Permissions](#)

- [Open](#)

[Enhancing antioxidant properties of lime juice powder through polyelectrolyte microparticles of chitosan-alginate: Formulation, characterization and stability](#)

[study](#)

Rahmiati, Nur; Sari, Retno; Wahyuni, Tutik Sri; More

Journal of Advanced Pharmaceutical Technology & Research. 15(3):231-236, Jul-Sep 2024.

- [Abstract](#)
- [Favorite](#)
- [PDF](#)
- [Permissions](#)

- [Open](#)

[Table of Contents Outline](#) | [Back to Top](#)

Short Communication

[Glucomannan is a promising isoniazid's enhancer that inducing macrophage phagocytosis](#)

Novita, Bernadette Dian; Wasiyastuti, Widya; Tjahjono, Yudy; More

Journal of Advanced Pharmaceutical Technology & Research. 15(3):237-241, Jul-Sep 2024.

- [Abstract](#)
- [Favorite](#)
- [PDF](#)
- [Permissions](#)

- [Open](#)

[Table of Contents Outline](#) | [Back to Top](#)

Advertisement

^Back to Top



Never Miss an Issue

Get new journal Tables of Contents sent right to your email inbox

Browse Journal Content

- [Register on the website](#)
- [Get eTOC Alerts](#)

Customer Service

- Contact us at:
 - Support:
[Submit a Service Request](#)
 - TEL:
800-638-3030 (within the USA)
301-223-2300 (outside of the USA)

-

- [Privacy Policy](#)
- [Legal Disclaimer](#)
- [Terms of Use](#)
- [Open Access Policy](#)
- [Your California Privacy Choices](#)



- Copyright © 2024
- Journal of Advanced Pharmaceutical Technology & Research | Published by Wolters Kluwer - [Medknow](#)

Advertisement

Potential antioxidant and antiradical agents from *Allium ascalonicum*: Superoxide dismutase and density functional theory *in silico* studies

Dwipa Ajiati, Dadan Sumiarsa,
Meiny Faudah Amin¹, Dikdik
Kurnia

Department of Chemistry, Faculty of
Mathematics and Natural Science,
Universitas Padjadjaran, Sumedang,
¹Department of Dental Conservation,
Faculty of Dentistry, Universitas
Trisakti, Jakarta, Indonesia

J. Adv. Pharm. Technol. Res.

ABSTRACT

Antioxidants are compounds that can inhibit excessive free radical reactions in the body. Excessive free radicals can cause system imbalances in the body which can trigger oxidative stress and cause serious illness. The limitations of antioxidants in the body can be overcome by consuming safe natural additional antioxidants that can be obtained from natural products. Isolating compounds of *Allium ascalonicum* leaves as antioxidant and antiradical agents in inhibiting excessive free radicals by *in vitro* and *in silico*. The extracted compounds were purified by column chromatography. The compounds obtained were then characterized using ultraviolet, infrared, NMR, and mass spectrometry. Determination of antioxidant activity was carried out by *in vitro* using 2,2-diphenyl-1-picrylhydrazyl (DPPH) and the non-enzymatic superoxide dismutase (SOD) methods. The *in silico* study used the density functional theory (DFT) calculation method with global descriptive parameters (GDP), donor acceptor map (DAM), and frontier molecular orbitals (FMO) analysis. Three compounds have been isolated, of which compound **1** is a new compound. In the DPPH method, compound **1** has more strong antioxidant activity than others, as well as in the non-enzymatic SOD method. Whereas, in the DFT calculation shows that compound **1** has the best reactivity and stability between other compounds and was categorized as the best antiradical. Compound **1** has the highest antioxidant activity compared to the other compounds by *in vitro* both the DPPH and non-enzymatic SOD methods. *In silico*, compound **1** has the potential as the best antiradical.

Key words: *Allium ascalonicum*, antioxidant, density functional theory, oxidative stress, the nonenzymatic superoxide dismutase

Address for correspondence:

Prof. Dikdik Kurnia,
Department of Chemistry, Faculty of Mathematics and Natural
Sciences, Universitas Padjadjaran, Sumedang, Indonesia.
E-mail: dikdik.kurnia@unpad.ac.id

Submitted: 03-Dec-2023

Revised: 02-May-2024

Accepted: 06-May-2024

Published: 22-Jul-2024

Access this article online

Quick Response Code:



Website:

<https://journals.lww.com/JAPTR>

DOI:

10.4103/japtr.japtr_525_23

INTRODUCTION

Shallot (*Allium ascalonicum* L.) is a plant which is widely cultivated and used as a spice and believed to be a traditional medicine.^[1] According to previous studies, *A. ascalonicum* bulbs contain several important compounds that have potential as natural antioxidant compounds.^[2]

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Ajiati D, Sumiarsa D, Amin MF, Kurnia D. Potential antioxidant and antiradical agents from *Allium ascalonicum*: Superoxide dismutase and density functional theory *in silico* studies. *J Adv Pharm Technol Res* 2024;15:171-6.

Antioxidants are compounds that can inhibit the occurrence of free radical reactions in the body.^[3] It is important to protect the body from free radicals that can harm and prevent the body from oxidative stress that causes several serious diseases such as premature aging and cancer.^[4] In addition, it is also caused by excessive production of reactive oxygen species.^[5] Natural antioxidants can be obtained from fruits and vegetables because they contain important compounds which have an antioxidant activity.^[6]

The antioxidant activity of a compound is influenced by its structure. The more polar the compound, the greater its ability is as an antioxidant. The structure which contains many hydroxyl groups bound to aromatic rings or conjugated carbon double bonds that can easily donate hydrogen or electrons to stabilize radical atoms because it has good resonance ability.^[7] In other studies, antioxidant activity assay is mostly conducted using the 2,2-diphenyl-1-picrylhydrazyl (DPPH) method, which is considered more effective, easy, and cheap.^[8] DPPH is a stable radical compound which has reactive nitrogen species.^[9] Another method used in this study is non-enzymatic superoxide dismutase (SOD) or known as mimic SOD because it mimics the workings of the antioxidant testing method using the SOD enzyme. In this method, the source of superoxide anion comes from riboflavin photoreduction and the role of SOD enzyme is replaced by secondary metabolite compounds.^[10]

In recent years, many studies on antioxidants have used computational studies or better known as *in silico* studies to predict the antioxidant ability of a molecule because it is considered more effective and certainly more affordable. One method that is widely used to predict antioxidant reactivity and activity is density functional theory (DFT).^[11,12] This method involves electron transfer ability and electron affinity in determining antioxidant ability.^[13] This study uses quantitative and qualitative approaches involving laboratory research. In addition to a new compound, the use of mimic SOD (mSOD) method *in vitro* and DFT *in silico* is a novelty of this study.

MATERIALS AND METHODS

Materials

The leaves of *A. ascalonicum* were obtained from Demak, Indonesia, which were then determined (No. 25/LBM/IT/12/2021) at the Taxonomy Laboratory, Padjadjaran University, Indonesia. Several organic solvents such as *n*-hexane, ethyl acetate (EtOAc), and methanol (MeOH) as well as several other chemicals, namely, Silica Gel 60 (0.2–0.5 mm) and (0.063–0.200 mm), Silica ODS RP-18 (0.040–0.063 mm), KLT Gel 60 F₂₅₄ plate and ODS RP-18 F₂₅₄ S plate were used for extraction, fractionation and isolation processes. In the process of identifying compound spots with KLT, 10% H₂SO₄ in ethanol (v/v)

spotting solution was used, which was then monitored under UV light at λ 254 and 365 nm. *In vitro* research, antioxidant activity test with DPPH and non-enzymatic SOD methods using several chemicals such as DPPH, methanol, riboflavin, phosphate buffer pH 7.4, TEMED, distilled water, and NBT.^[10,14] Materials for *in silico* study only require 3D-conformers of compounds 1–3 downloaded from PubChem in sdf format.

Extraction and isolation of *Allium ascalonicum* leaves

A. ascalonicum leaves were extracted using methanol. The methanol extract obtained was concentrated with a rotary evaporator at \pm 50°C. The extract was partitioned using column chromatography (Silica G 60) with *n*-hexane, EtOAc, and MeOH to obtain fractions of each solvent. EtOAc and MeOH fractions were selected to be purified using Silica G 60 column chromatography (*n*-hexane-EtOAc) and ODS RP-18 (H₂O-MeOH) to obtain pure compounds 1 and 2. Characterization of pure compounds was carried out using ¹H-NMR (JEOL, 500 MHz), ultraviolet-visible, infrared, and mass spectrometry.

Antioxidant activity assays using the 2,2-diphenyl-1-picrylhydrazyl method

This method was carried out twice, in the preliminary test and the pure compounds. Solvent used is methanol and DPPH as a radical with variation of sample concentration. The absorbance measurement was conducted using a microplate reader at 510 nm after being left in a dark room for \pm 30 min. Percentage inhibition was calculated using the absorbance value of measurement result using $(A_{\text{control}} - A_{\text{sample}}) / A_{\text{control}} \times 100\%$. Whereas, the IC₅₀ value was calculated using $Y = ax + b$, which is the result of the regression equation.^[15]

Antioxidant activity assays using the mSOD method

Initially, we prepared sample solutions of compounds with varying concentrations. The concentration variation was done by microdilution. Then, the sample solution, MeOH, working solution A (aquabides, phosphate buffer pH 7.4, NBT, TEMED, Riboflavin) and B (aquabides, phosphate buffer pH 7.4, NBT, TEMED) were put into 96-well and then irradiated for 10–15 min in a closed box. Absorbance measurement was performed using a microplate reader at λ 550 nm to obtain percentage inhibition and IC₅₀ value.

Study of antioxidant activity with density functional theory calculation by *in silico*

In this antioxidant activity study by *in silico*, we used DFT calculation with basis sets B3 LYP/6-31G (2d, 2p) using MarvinSketch, Gaussian 09 W, and GaussView 5.0. The calculation was carried out under gas phase conditions.^[16] We used three analyses consisting of global descriptive parameters (GDP), donor acceptor map (DAM), and frontier molecular orbitals (FMO). The three analyses involved calculations and DAM diagrams as follows [Figure 1].

RESULTS

Compounds **1** and **2** were obtained from the EtOAc fraction of the methanol extract of *A. ascalonicum* leaves. Meanwhile, compound **3** was obtained from the methanol fraction. Based on the characterization data, it shows that compound **1** belongs to the aromatic compound group and compounds **2–3** belongs to the steroid group.

2-((2-hydroxyphenanthrene-1-yl)oxy)-2-oxoacetic acid (**1**) is a white crystal-shaped compound and has the molecular formula of $C_{16}H_{10}O_5$; IR, ν_{max} 1094 (C-O), 1494 (C=C), 1683, 1701 (C=O) and 3453 cm^{-1} (OH); MS, m/z 282 (M-H)⁺; It has the carbon sp^2 at the shift δ_c 167.7 ppm which is a characteristic of the ester or carboxylic acid group. Then, there are several characteristics of alkene and aromatic carbon at δ_c 121.6–135.6 ppm.^[17] According to the UV data of compound **1** at λ_{max} 255 nm is the absorption band of the benzenoid indicating that compound **1** has a benzene ring. Based on the shift δ_c , compound **1** has eight methine (CH) and eight sp^2 quaternary (Cq) carbons.^[18] β -sitosterol (**2**), the white crystal-shaped compound, has the molecular formula of $C_{29}H_{50}O$; IR, ν_{max} 1050 (C-O), 1381 (*gem*-dimethyl), 1465 (C=C), 2937 (C-H) and 3429 cm^{-1} (OH); MS, m/z 414 (M+H)⁺; These data provide information that compound **2** has six methyl, eleven methylene, nine methine and three quaternary (Cq) carbons.^[19] Sitosterol-3-*O*-glucoside (**3**) is a white powder-shaped compound with the molecular

formula of $C_{35}H_{60}O_6$; IR, ν_{max} 1021 (C-O), 1367 (*gem*-dimethyl), 1639 (C=C), 2932 (C-H) and 3400 (OH); MS, m/z 576 (M-H)⁺.^[20]

Based on ^1H and ^{13}C -NMR analysis, compounds **2** and **3** have almost the same chemical shift. However, compound **3** has chemical shift that was the characteristics of glucoside at δ_c 61.5 (C-26), 73.9 (C-28), 77.2 (C-29, C-29a), 77.3 (C-30), and 101.2 ppm (C-31).^[21] According to ^{13}C -NMR, compound **1** has carboxyl group at δ_c 167.7 ppm (C-12, C-12a).^[22] Structures of compounds **1–3** are shown in Figure 2.

DISCUSSION

Antioxidant activity assays of *Allium ascalonicum* leave compounds by *in vitro*

Table 1 shows that compound **1** has the best antioxidant activity among the other compounds in DPPH and non-enzymatic SOD methods with IC_{50} of 76 and 40 $\mu\text{g/mL}$, respectively. The IC_{50} value of compound **1** indicates that the compound has strong antioxidant activity. Meanwhile, compounds **2–3** showed weak antioxidant activity in both methods. In this study, we used quercetin as a positive control in both methods because quercetin is known to have a strong antioxidant ability to inhibit free radicals.^[14]

Based on Table 1, compound **1** has good activity because it is influenced by its structure. There are several influential groups from compound **1** that can affect antioxidant activity, namely the aromatic benzene ring, carbonyl groups, and phenol hydroxyl groups attached to it.^[23] Compound **1** could donate its electrons to stabilize radical atoms to become more stable.^[24] Meanwhile, compounds **2–3** do not have an influential group to donate its electrons. Although compound **3** has some hydroxyl groups on the




Figure 1: Donor acceptor map analysis diagram used to determine antioxidant activity *in silico*

Table 1: Data of antioxidant activity of compounds^[1–3]

Compounds	IC_{50} ($\mu\text{g/mL}$)	
	DPPH	SOD nonenzymatic
1	76	40
2	3273	3837
3	2500	10,000
Quercetin	3.8	5

DPPH: 2-diphenyl-1-picrylhydrazyl, SOD: Superoxide dismutase, IC: Inhibitory concentration




Figure 2: Compounds **1–3** were obtained from *Allium ascalonicum* leaves

glucoside substituents in its structure, it cannot conjugate and delocalize electrons when donating or transferring electrons.^[25] Those functional groups have a relationship with the polarity of compounds and effect on antioxidant activity.^[26] Figure 2 also shows that compound 1 is more polar than compounds 2 and 3. The more polar the compound, the easier it is to break hydrogen bonds and donate electrons.^[27]

Antioxidant activity assay of *Allium ascalonicum* leave compounds by *in silico*

In the DFT calculation in this study, we used three analysis methods to determine and validate the results of antioxidant activity *in vitro*, namely, GDP, DAM, and FMO.

Table 2: The result of global descriptive parameter calculation

Parameters (eV)	Compounds			
	1	2	3	Quercetin
I	10.85	7.62	7.51	9.43
A	1.12	-1.93	1.63	1.75
η	4.86	4.77	4.57	3.83
S	0.10	0.10	0.11	0.13
χ	5.99	2.84	2.94	5.59
μ	-5.99	-2.84	-2.94	-5.59
ω	17.96	4.05	4.32	15.64

Table 3: The calculation results of compounds 1–3 using donor acceptor map analysis

Compounds	Ω^- (eV)	ω^+ (eV)	Ra	Rd
1	7.29	1.30	0.62	2.08
2	2.86	0.02	0.01	0.82
3	2.98	0.04	0.02	0.85
Quercetin	7.35	1.75	0.84	2.10
Natrium	3.50	0.57	0.27	1.00
Flourin	10.68	2.09	1.00	3.05

Ra: Electron acceptor, Rd: Electron donor

Global descriptive parameters

GDP calculation was obtained from single point energy value of each charge (neutral, anion, and cation) using formulas that were mentioned above (method section). Those parameters are used to predict reactivities and stability of compounds.^[28] They consist of ionization potential (I), electron affinity (A), hardness (η), softness (S), electronegativity (χ), chemical potential (μ), and electrophilicity index (ω).^[29] Table 2 shows that compound 1 is more stable than compounds 2–3 as indicated by the higher hardness values than the others. The higher the hardness value, the more stable the compound. Stability can also be seen based on a more negative chemical potential value. Reactivity and stability greatly affect antioxidant activity as they are related to electron transfer in inhibiting radicals.^[30]

Donor acceptor map

Determination of antioxidant activity based on DAM analysis is inseparable from the previous analysis, namely GDP. I and A values as in Table 2 were used to




Figure 3: The results of compounds 1–3 were obtained using donor acceptor map diagram analysis




Figure 4: Frontier molecular orbitals analysis of compounds 1–3 involved highest occupied molecular orbital-lowest unfilled molecular orbital energy. HOMO: Highest occupied molecular orbital, LUMO: Lowest unfilled molecular orbital

calculate electron donor (ω^-) and electron acceptor (ω^+) to obtain electron donor (Rd) and electron acceptor (Ra) index.^[29,31] We determined these values using the formulas mentioned above. In this analysis, fluorine and sodium were also involved to obtain the Ra and Rd values. Fluorine (F) represents an electron acceptor and sodium (Na) a good electron donor.^[31] The Ra and Rd values that have been obtained as in Table 3, then plotted on the DAM diagram. Figure 3 shows that compound **1** is categorized as the best antiradical that acts as a good acceptor and donor based on the DAM diagram [Figure 1]. Meanwhile, compounds **2–3** were categorized as a good antiradical with poor acceptor, good donor, and good antioxidant. When a molecule acts as a good electron donor and acceptor, it has high reactivity, so that hydrogen bonds can be broken more easily and donate electrons or hydrogen.^[32]

Frontier molecular orbital

The third analysis is FMO which involves the highest occupied molecular orbital energy (HOMO)–the lowest unfilled molecular orbital energy (LUMO). In this analysis, the reactivity of the compound can be seen from the energy gap between the HOMO-LUMO energies of the molecule. The lower the energy gap, the more reactive the molecule is.^[32] HOMO-LUMO visualization can also shed light on orbital distribution and electron density. In addition, this analysis can show the active sites or radical attack sites in the molecule, qualitatively.^[33] As shown in Figure 4, of the three compounds, compound **1** has a lower energy gap than the other two compounds and even lower than the positive control. This shows that compound **1** has the highest reactivity with an energy gap of 0.11 eV.^[13] Figure 2 shows that radicals can attack compound **1** at the aromatic ring and phenol hydroxyl group. Meanwhile, in compounds **2** and **3**, radicals can attack the cyclohexanol site. This can be seen from the orbital distribution in the HOMO-LUMO analysis as shown in Figure 4.

CONCLUSION

In this study, we have successfully isolated three compounds and one of them is a new compound. Based on the *in vitro* antioxidant activity test, compound **1** has strong antioxidant activity on DPPH and non-enzymatic SOD method. Based on *in silico* study, in GDP analysis, compound **1** has the highest reactivity and stability compared to other compounds. This is also reinforced by the results of FMO analysis which can be seen from the energy gap value. In DAM analysis, compound **1** is categorized as the best antiradical among the other. Thus, the antioxidant activity of compounds using *in vitro* methods does not correlate to certain analyses in *in silico* studies.

Acknowledgement

The authors are grateful to the Academic Leadership Grant (ALG) Dikdik Kurnia, Indonesia (1439/UN6.3.1/

PT.00/2024, March 18, 2024) and also grateful to Universitas Padjadjaran for supporting all research facilities.

Financial support and sponsorship

The Academic Leadership Grant (ALG) Dikdik Kurnia, Indonesia (1439/UN6.3.1/PT.00/2024, March 18, 2024), Universitas Padjadjaran, Indonesia.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Thuy NM, Tuyen NT, Cuong NP, Huyen LT, Phuong NP, Nguyen LT, *et al.* Identification and extraction method of quercetin from flesh and skin of shallot (*Allium ascalonicum*) cultivated in Soc Trang Province, Vietnam. *Food Res* 2020;4:358-65.
2. Deswati DA, Dhina MA, Mubaroq SR. IOP Conference Series: Materials Science and Engineering. Bristol, England: Institute of Physics Publishing; 2018.
3. Sandhiutami NM, Rahayu L. Uji toksisitas akut, aktivitas antioksidan *in vitro* dan efek rebusan bunga kamboja merah (*Plumeria rubra* L.) terhadap kadar malondialdehid. *J Ilmu Kefarmasian Indones* 2014;12:43-9.
4. Batinić Haberle I, Rebouças JS, Spasojević I. Superoxide dismutase mimics: Chemistry, pharmacology, and therapeutic potential. *Antioxid Redox Signal* 2010;13:877-918.
5. Gospodaryov D, Lushchak V. Oxidative Stress: Cause and Consequence of Diseases. *Oxid Stress Dis* 2012;2:14-38. [doi: 10.5772/38093].
6. Rahal A, Kumar A, Singh V, Yadav B, Tiwari R, Chakraborty S, *et al.* Oxidative stress, prooxidants, and antioxidants: The interplay. *Biomed Res Int* 2014;2014:761264.
7. Ahmadi SM, Farhoosh R, Sharif A, Rezaie M. Structure-antioxidant activity relationships of luteolin and catechin. *J Food Sci* 2020;85:298-305.
8. Romulo A. The Principle of Some *in vitro* Antioxidant Activity Methods: Review. *IOP Conf Ser Earth Environ Sci* 2020;426:1-7. [doi: 10.1088/1755-1315/426/1/012177].
9. Tirezitis G, Bartosz G. Determination of antiradical and antioxidant activity: Basic principles and new insights. *Acta Biochim Pol* 2010;57:139-42.
10. Deawati Y, Onggo D, Mulyani I, Hastiawan I, Kurnia D. Activity of Superoxide Dismutase Mimic of [Mn(salen)OAc] Complex Compound Non-enzymatically *In Vitro* Through Riboflavin Photoreduction. *Molekul* 2017;12:61.
11. Dupont S, Fleurat Lessard P, Cruz RG, Lafarge C, Grangeteau C, Yahou F, *et al.* Antioxidant properties of ergosterol and its role in yeast resistance to oxidation. *Antioxidants (Basel)* 2021;10:1024.
12. Chong DP. Density functional theory study of allopurinol. *Can J Chem* 2013;91:637-41.
13. Farrokhnia M. Density functional theory studies on the antioxidant mechanism and electronic properties of some bioactive marine meroterpenoids: Sargahydroquinonic acid and sargachromanol. *ACS Omega* 2020;5:20382-90.
14. Ozgen S, Kilinc OK, Selamoğlu Z. Antioxidant Activity of Quercetin: A Mechanistic Review. *Turk J Agric Food Sci Technol* 2016;4:1134.
15. Rivero Cruz JF, Granados Pineda J, Pedraza Chaverri J, Pérez Rojas JM, Kumar Passari A, Diaz Ruiz G, *et al.* Phytochemical constituents, antioxidant, cytotoxic, and antimicrobial activities of the ethanolic extract of Mexican brown propolis. *Antioxidants (Basel)* 2020;9:70.

16. Soliva R, Luque FJ, Orozco M. Reliability of MEP and MEP-derived properties computed from DFT methods for molecules containing P, S and CL. *Theor Chem Acc* 1997;98:42-9.
17. Rakhmatullin IZ, Efimov SV, Klochkov AV, Gnezdilov OI, Varfolomeev MA, Klochkov VV. NMR chemical shifts of carbon atoms and characteristic shift ranges in the oil sample. *Pet Res* 2022;7:269-74.
18. Rawat DS, Kumar D. Nature of electronic transitions and factors affecting it. *Org Spectro* 2017;12:1-21.
19. Khan N, Sagar Hossain MN, Mofiz Uddin Khan Lecturer M, Mofiz Uddin Khan N. Scopoletin and β -sitosterol glucoside from roots of *Ipomoea digitata*. *J Pharmacogn Phytochem* 2015;4:5-7.
20. Klawun C, Wilkins CL, Chem J. Optimization of functional group prediction from infrared spectra using neural networks. *Inf Comput Sci* 1996;36:69-81.
21. Khatun M, Billah M, Quader MA. Sterols and Sterol Glucoside from *Phyllanthus* Species. *Dhaka Univ J Sci* 2012;60:5-10.
22. Zhang X, Macmillan DW. Alcohols as Latent Coupling Fragments for Metallaphotoredox Catalysis: sp³-sp² Cross-Coupling of Oxalates with Aryl Halides *J Am Chem Soc* 2016. p. 1-5.
23. Muzolf-Panek M, Gliszczynska-Świgło A, Szymusiak H, Tyrakowska B. The influence of stereochemistry on the antioxidant properties of catechin epimers. *Eur Food Res Technol* 2012;235:1001-9.
24. Heim KE, Tagliaferro AR, Bobilya DJ. Flavonoid antioxidants: Chemistry, metabolism and structure-activity relationships. *J Nutr Biochem* 2002;13:572-84.
25. Godlewska Żyłkiewicz B, Świsłocka R, Kalinowska M, Golonko A, Świdorski G, Arciszewska Ż, et al. Biologically active compounds of plants: Structure-related antioxidant, microbiological and cytotoxic activity of selected carboxylic acids. *Materials (Basel)* 2020;13:4454.
26. Saptarini NM, Wardati Y. Effect of extraction methods on antioxidant activity of papery skin extracts and fractions of Maja Cipanas onion (*Allium cepa* L. var. *ascalonicum*). *ScientificWorldJournal* 2020;2020:1-6.
27. Rice Evans C. Flavonoid antioxidants. *Curr Med Chem* 2001;8:797-807.
28. Safna Hussan KP, Shahin Thayyil M, Rajan VK, Muraleedharan K. DFT studies on global parameters, antioxidant mechanism and molecular docking of amlodipine besylate. *Comput Biol Chem* 2019;80:46-53.
29. Pottachola S, Kaniyantavida A, Karuvathodiyil M, Density Functional Theory – Recent Advances, New Perspectives and Applications. Croatia: IntechOpen; 2012. p. 1-18.
30. Xue Y, Zheng Y, An L, Dou Y, Liu Y. Density functional theory study of the structure-antioxidant activity of polyphenolic deoxybenzoins. *Food Chem* 2014;151:198-206.
31. Hernandez DA, Rodriguez Zavala JG, Tenorio FJ. DFT study of antioxidant molecules from traditional Japanese and Chinese teas: comparing allylic and phenolic antiradical activity. *Struct Chem* 2020;31:359-69.
32. Rajan VK, Ragi C, Muraleedharan K. A computational exploration into the structure, antioxidant capacity, toxicity and drug-like activity of the anthocyanidin “petunidin”. *Heliyon* 2019;5:e02115.
33. Mahmoudi S, Dehkordi MM, Asgarshamsi MH. Density functional theory studies of the antioxidants-a review. *J Mol Model* 2021;27:271.

Potential antioxidant and antiradical agents from *Allium ascalonicum*: Superoxide dismutase and density functional theory in silico studies

by Meiny F Amin FKG

Submission date: 27-Sep-2024 02:07PM (UTC+0700)

Submission ID: 2427490523

File name: potential_antioxidant_and_antiradical_agents_from.8.pdf (1.32M)

Word count: 4145

Character count: 22352

Potential antioxidant and antiradical agents from *Allium ascalonicum*: Superoxide dismutase and density functional theory *in silico* studies

Dwipa Ajiati, Dadan Sumiarsa,
Meiny Faudah Amin¹, Dikdik
Kurnia

Department of Chemistry, Faculty of
Mathematics and Natural Science,
Universitas Padjadjaran, Sumedang,
¹Department of Dental Conservation,
Faculty of Dentistry, Universitas
Trisakti, Jakarta, Indonesia

J. Adv. Pharm. Technol. Res.

ABSTRACT

Antioxidants are compounds that can inhibit excessive free radical reactions in the body. Excessive free radicals can cause system imbalances in the body which can trigger oxidative stress and cause serious illness. The limitations of antioxidants in the body can be overcome by consuming safe natural additional antioxidants that can be obtained from natural products. Isolating compounds of *Allium ascalonicum* leaves as antioxidant and antiradical agents in inhibiting excessive free radicals by *in vitro* and *in silico*. The extracted compounds were purified by column chromatography. The compounds obtained were then characterized using ultraviolet, infrared, NMR, and mass spectrometry. Determination of antioxidant activity was carried out by *in vitro* using 2,2-diphenyl-1-picrylhydrazyl (DPPH) and the non-enzymatic superoxide dismutase (SOD) methods. The *in silico* study used the density functional theory (DFT) calculation method with global descriptive parameters (GDP), donor acceptor map (DAM), and frontier molecular orbitals (FMO) analysis. Three compounds have been isolated, of which compound 1 is a new compound. In the DPPH method, compound 1 has more strong antioxidant activity than others, as well as in the non-enzymatic SOD method. Whereas, in the DFT calculation shows that compound 1 has the best reactivity and stability between other compounds and was categorized as the best antiradical. Compound 1 has the highest antioxidant activity compared to the other compounds by *in vitro* both the DPPH and non-enzymatic SOD methods. *In silico*, compound 1 has the potential as the best antiradical.

Key words: *Allium ascalonicum*, antioxidant, density functional theory, oxidative stress, the nonenzymatic superoxide dismutase

Address for correspondence:

Prof. Dikdik Kurnia,
Department of Chemistry, Faculty of Mathematics and Natural
Sciences, Universitas Padjadjaran, Sumedang, Indonesia.
E-mail: dikdik.kurnia@unpad.ac.id

Submitted: 03-Dec-2023

Revised: 02-May-2024

Accepted: 06-May-2024

Published: 22-Jul-2024

Access this article online

Quick Response Code:



Website:

<https://journals.lww.com/JAPTR>

DOI:

10.4103/japtr.japtr_525_23

INTRODUCTION

Shallot (*Allium ascalonicum* L.) is a plant which is widely cultivated and used as a spice and believed to be a traditional medicine.^[1] According to previous studies, *A. ascalonicum* bulbs contain several important compounds that have potential as natural antioxidant compounds.^[2]

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Ajiati D, Sumiarsa D, Amin MF, Kurnia D. Potential antioxidant and antiradical agents from *Allium ascalonicum*: Superoxide dismutase and density functional theory *in silico* studies. *J Adv Pharm Technol Res* 2024;15:171-6.

Antioxidants are compounds that can inhibit the occurrence of free radical reactions in the body.^[3] It is important to protect the body from free radicals that can harm and prevent the body from oxidative stress that causes several serious diseases such as premature aging and cancer.^[4] In addition, it is also caused by excessive production of reactive oxygen species.^[5] Natural antioxidants can be obtained from fruits and vegetables because they contain important compounds which have an antioxidant activity.^[6]

The antioxidant activity of a compound is influenced by its structure. The more polar the compound, the greater its ability is as an antioxidant. The structure which contains many hydroxyl groups bound to aromatic rings or conjugated carbon double bonds that can easily donate hydrogen or electrons to stabilize radical atoms because it has good resonance ability.^[7] In other studies, antioxidant activity assay is mostly conducted using the 2,2-diphenyl-1-picrylhydrazyl (DPPH) method, which is considered more effective, easy, and cheap.^[8] DPPH is a stable radical compound which has reactive nitrogen species.^[9] Another method used in this study is non-enzymatic superoxide dismutase (SOD) or known as mimic SOD because it mimics the workings of the antioxidant testing method using the SOD enzyme. In this method, the source of superoxide anion comes from riboflavin photoreduction and the role of SOD enzyme is replaced by secondary metabolite compounds.^[10]

In recent years, many studies on antioxidants have used computational studies or better known as *in silico* studies to predict the antioxidant ability of a molecule because it is considered more effective and certainly more affordable. One method that is widely used to predict antioxidant reactivity and activity is density functional theory (DFT).^[11,12] This method involves electron transfer ability and electron affinity in determining antioxidant ability.^[13] This study uses quantitative and qualitative approaches involving laboratory research. In addition to a new compound, the use of mimic SOD (mSOD) method *in vitro* and DFT *in silico* is a novelty of this study.

13 MATERIALS AND METHODS

Materials

The leaves of *A. ascalonicum* were obtained from Demak, Indonesia, which were then determined (No. 25/LBM/IT/12/2021) at the Taxonomy Laboratory, Padjadjaran University, Indonesia. Several organic solvents such as *n*-hexane, ethyl acetate (EtOAc), and methanol (MeOH) as well as several other chemicals, namely, Silica Gel 60 (0.2–0.5 mm) and (0.063–0.200 mm), Silica ODS RP-18 (0.040–0.063 mm), KLT Gel 60 F₂₅₄ plate and ODS RP-18 F₂₅₄ S plate were used for extraction, fractionation and isolation processes. In the process of identifying compound spots with KLT, 10% H₂SO₄ in ethanol (v/v)

spotting solution was used, which was then monitored under UV light at λ 254 and 365 nm. *In vitro* research, antioxidant activity test with DPPH and non-enzymatic SOD methods using several chemicals such as DPPH, methanol, riboflavin, phosphate buffer pH 7.4, TEMED, distilled water, and NBT.^[10,14] Materials for *in silico* study only require 3D-conformers of compounds 1–3 downloaded from PubChem in sdf format.

Extraction and isolation of *Allium ascalonicum* leaves

A. ascalonicum leaves were extracted using methanol. The methanol extract obtained was concentrated with a rotary evaporator at \pm 50°C. The extract was partitioned using column chromatography (Silica G 60) with *n*-hexane, EtOAc, and MeOH to obtain fractions of each solvent. EtOAc and MeOH fractions were selected to be purified using Silica G 60 column chromatography (*n*-hexane-EtOAc) and ODS RP-18 (H₂O-MeOH) to obtain pure compounds 1 and 2. Characterization of pure compounds was carried out using ¹H-NMR (JEOL, 500 MHz), ultraviolet-visible, infrared, and mass spectrometry.

5 Antioxidant activity assays using the 2,2-diphenyl-1-picrylhydrazyl method

This method was carried out twice, in the preliminary test and the pure compounds. Solvent used is methanol and DPPH as a radical with variation of sample concentration. The absorbance measurement was conducted using a microplate reader at 510 nm after being left in a dark room for \pm 30 min. Percentage inhibition was calculated using the absorbance value of measurement result using $(A_{\text{control}} - A_{\text{sample}}) / A_{\text{control}} \times 100\%$. Whereas, the IC₅₀ value was calculated using $Y = ax + b$, which is the result of the regression equation.^[15]

Antioxidant activity assays using the mSOD method

Initially, we prepared sample solutions of compounds with varying concentrations. The concentration variation was done by microdilution. Then, the sample solution, MeOH, working solution A (aquabides, phosphate buffer pH 7.4, NBT, TEMED, Riboflavin) and B (aquabides, phosphate buffer pH 7.4, NBT, TEMED) were put into 96-well and then irradiated for 10–15 min in a closed box. Absorbance measurement was performed using a microplate reader at λ 550 nm to obtain percentage inhibition and IC₅₀ value.

Study of antioxidant activity with density functional theory calculation by *in silico*

In this antioxidant activity study by *in silico*, we used DFT calculation with basis sets B3 LYP/6-31G (2d, 2p) using MarvinSketch, Gaussian 09 W, and GaussView 5.0. The calculation was carried out under gas phase conditions.^[16] We used three analyses consisting of global descriptive parameters (GDP), donor acceptor map (DAM), and frontier molecular orbitals (FMO). The three analyses involved calculations and DAM diagrams as follows [Figure 1].

RESULTS

Compounds **1** and **2** were obtained from the EtOAc fraction of the methanol extract of *A. ascalonicum* leaves. Meanwhile, compound **3** was obtained from the methanol fraction. Based on the characterization data, it shows that compound **1** belongs to the aromatic compound group and compounds **2–3** belongs to the steroid group.

2-((2-hydroxyphenanthrene-1-yl) oxy)-2-oxoacetic acid (**1**) is a white crystal-shaped compound and has the molecular formula of $C_{16}H_{10}O_5$; IR, ν_{\max} 1094 (C-O), 1494 (C=C), 1683, 1701 (C=O) and 3453 cm^{-1} (OH); MS, m/z 282 (M-H)⁺; It has the carbon sp^2 at the shift δ_c 167.7 ppm which is a characteristic of the ester or carboxylic acid group. Then, there are several characteristics of alkene and aromatic carbon at δ_c 121.6–135.6 ppm.^[17] According to the UV data of compound **1** at λ_{\max} 255 nm is the absorption band of the benzenoid indicating that compound **1** has a benzene ring. Based on the shift δ_c , compound **1** has eight methine (CH) and eight sp^2 quaternary (Cq) carbons.^[18] β -sitosterol (**2**), the white crystal-shaped compound, has the molecular formula of $C_{29}H_{50}O$; IR, ν_{\max} 1050 (C-O), 1381 (*gem*-dimethyl), 1465 (C=C), 2937 (C-H) and 3429 cm^{-1} (OH); MS, m/z 414 (M+H)⁺; These data provide information that compound **2** has six methyl, eleven methylene, nine methine and three quaternary (Cq) carbons.^[19] Sitosterol-3-O-glucoside (**3**) is a white powder-shaped compound with the molecular

formula of $C_{35}H_{60}O_6$; IR, ν_{\max} 1021 (C-O), 1367 (*gem*-dimethyl), 1639 (C=C), 2932 (C-H) and 3400 (OH); MS, m/z 576 (M-H)⁺.^[20]

Based on ^1H and ^{13}C -NMR analysis, compounds **2** and **3** have almost the same chemical shift. However, compound **3** has chemical shift that was the characteristics of glucoside at δ_c 61.5 (C-26), 73.9 (C-28), 77.2 (C-29, C-29a), 77.3 (C-30), and 101.2 ppm (C-31).^[21] According to ^{13}C -NMR, compound **1** has carboxyl group at δ_c 167.7 ppm (C-12, C-12a).^[22] Structures of compounds **1–3** are shown in Figure 2.

DISCUSSION

Antioxidant activity assays of *Allium ascalonicum* leave compounds by *in vitro*

Table 1 shows that compound **1** has the best antioxidant activity among the other compounds in DPPH and non-enzymatic SOD methods with IC_{50} of 76 and 40 $\mu\text{g/mL}$, respectively. The IC_{50} value of compound **1** indicates that the compound has strong antioxidant activity. Meanwhile, compounds **2–3** showed weak antioxidant activity in both methods. In this study, we used quercetin as a positive control in both methods because quercetin is known to have a strong antioxidant ability to inhibit free radicals.^[14]

Based on Table 1, compound **1** has good activity because it is influenced by its structure. There are several influential groups from compound **1** that can affect antioxidant activity, namely the aromatic benzene ring, carbonyl groups, and phenol hydroxyl groups attached to it.^[23] Compound **1** could donate its electrons to stabilize radical atoms to become more stable.^[24] Meanwhile, compounds **2–3** do not have an influential group to donate its electrons. Although compound **3** has some hydroxyl groups on the




Figure 1: Donor-acceptor map analysis diagram used to determine antioxidant activity *in silico*

Table 1: Data of antioxidant activity of compounds^[1-3]

Compounds	IC_{50} ($\mu\text{g/mL}$)	
	DPPH	SOD nonenzymatic
1	76	40
2	3273	3837
3	2500	10,000
Quercetin	3.8	5

DPPH: 2-diphenyl-1-picrylhydrazyl, SOD: Superoxide dismutase, IC: Inhibitory concentration




Figure 2: Compounds **1–3** were obtained from *Allium ascalonicum* leaves

glucoside substituents in its structure, it cannot conjugate and delocalize electrons when donating or transferring electrons.^[25] Those functional groups have a relationship with the polarity of compounds and effect on antioxidant activity.^[26] Figure 2 also shows that compound 1 is more polar than compounds 2 and 3. The more polar the compound, the easier it is to break hydrogen bonds and donate electrons.^[27]

Antioxidant activity assay of *Allium ascalonicum* leave compounds by *in silico*

In the DFT calculation in this study, we used three analysis methods to determine and validate the results of antioxidant activity *in vitro*, namely, GDP, DAM, and FMO.

Table 2: The result of global descriptive parameter calculation

Parameters (eV)	Compounds			
	1	2	3	Quercetin
I	10.85	7.62	7.51	9.43
A	1.12	-1.93	1.63	1.75
η	4.86	4.77	4.57	3.83
S	0.10	0.10	0.11	0.13
χ	5.99	2.84	2.94	5.59
μ	-5.99	-2.84	-2.94	-5.59
ω	17.96	4.05	4.32	15.64

Table 3: The calculation results of compounds 1–3 using donor acceptor map analysis

Compounds	Ω^- (eV)	ω^+ (eV)	Ra	Rd
1	7.29	1.30	0.62	2.08
2	2.86	0.02	0.01	0.82
3	2.98	0.04	0.02	0.85
Quercetin	7.35	1.75	0.84	2.10
Natrium	3.50	0.57	0.27	1.00
Flourin	10.68	2.09	1.00	3.05

Ra: Electron acceptor, Rd: Electron donor

Global descriptive parameters

GDP calculation was obtained from single point energy value of each charge (neutral, anion, and cation) using formulas that were mentioned above (method section). Those parameters are used to predict reactivities and stability of compounds.^[28] They consist of ionization potential (I), electron affinity (A), hardness (η), softness (S), electronegativity (χ), chemical potential (μ), and electrophilicity index (ω).^[29] Table 2 shows that compound 1 is more stable than compounds 2–3 as indicated by the higher hardness values than the others. The higher the hardness value, the more stable the compound. Stability can also be seen based on a more negative chemical potential value. Reactivity and stability greatly affect antioxidant activity as they are related to electron transfer in inhibiting radicals.^[30]

Donor acceptor map

Determination of antioxidant activity based on DAM analysis is inseparable from the previous analysis, namely GDP. I and A values as in Table 2 were used to




Figure 3: The results of compounds 1–3 were obtained using donor acceptor map diagram analysis




Figure 4: Frontier molecular orbitals analysis of compounds 1–3 involved highest occupied molecular orbital-lowest unfilled molecular orbital energy. HOMO: Highest occupied molecular orbital, LUMO: Lowest unfilled molecular orbital

calculate electron donor (ω^-) and electron acceptor (ω^+) to obtain electron donor (Rd) and electron acceptor (Ra) index.^[29,31] We determined these values using the formulas mentioned above. In this analysis, florin and sodium were also involved to obtain the Ra and Rd values. Florin (F) represents an electron acceptor and sodium (Na) a good electron donor.^[31] The Ra and Rd values that have been obtained as in Table 3, then plotted on the DAM diagram. Figure 3 shows that compound 1 is categorized as the best antiradical that acts as a good acceptor and donor based on the DAM diagram [Figure 1]. Meanwhile, compounds 2–3 were categorized as a good antiradical with poor acceptor, good donor, and good antioxidant. When a molecule acts as a good electron donor and acceptor, it has high reactivity, so that hydrogen bonds can be broken more easily and donate electrons or hydrogen.^[32]

Frontier molecular orbital

The third analysis is FMO which involves the highest occupied molecular orbital energy (HOMO)—the lowest unfilled molecular orbital energy (LUMO). In this analysis, the reactivity of the compound can be seen from the energy gap between the HOMO-LUMO energies of the molecule. The lower the energy gap, the more reactive the molecule is.^[32] HOMO-LUMO visualization can also shed light on orbital distribution and electron density. In addition, this analysis can show the active sites or radical attack sites in the molecule, qualitatively.^[33] As shown in Figure 4, of the three compounds, compound 1 has a lower energy gap than the other two compounds and even lower than the positive control. This shows that compound 1 has the highest reactivity with an energy gap of 0.11 eV.^[13] Figure 2 shows that radicals can attack compound 1 at the aromatic ring and phenol hydroxyl group. Meanwhile, in compounds 2 and 3, radicals can attack the cyclohexanol site. This can be seen from the orbital distribution in the HOMO-LUMO analysis as shown in Figure 4.

CONCLUSION

In this study, we have successfully isolated three compounds and one of them is a new compound. Based on the *in vitro* antioxidant activity test, compound 1 has strong antioxidant activity on DPPH and non-enzymatic SOD method. Based on *in silico* study, in GDP analysis, compound 1 has the highest reactivity and stability compared to other compounds. This is also reinforced by the results of FMO analysis which can be seen from the energy gap value. In DAM analysis, compound 1 is categorized as the best antiradical among the other. Thus, the antioxidant activity of compounds using *in vitro* methods does not correlate to certain analyses in *in silico* studies.

Acknowledgement

The authors are grateful to the Academic Leadership Grant (ALG) Dikdik Kurnia, Indonesia (1439/UN6.3.1/

PT.00/2024, March 18, 2024) and also grateful to Universitas Padjadjaran for supporting all research facilities.

Financial support and sponsorship

The Academic Leadership Grant (ALG) Dikdik Kurnia, Indonesia (1439/UN6.3.1/PT.00/2024, March 18, 2024), Universitas Padjadjaran, Indonesia.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Thuy NM, Tuyen NT, Cuong NP, Huyen LT, Phuong NP, Nguyen LT, et al. Identification and extraction method of quercetin from flesh and skin of shallot (*Allium ascalonicum*) cultivated in Soc Trang Province, Vietnam. *Food Res* 2020;4:358-65.
- Deswati DA, Dhina MA, Mubaroq SR. IOP Conference Series: Materials Science and Engineering. Bristol, England: Institute of Physics Publishing; 2018.
- Sandhiutami NM, Rahayu L. Uji toksisitas akut, aktivitas antioksidan *in vitro* dan efek rebusan bunga kamboja merah (*Plumeria rubra* L.) terhadap kadar malondialdehid. *J Ilmu Kefarmasian Indones* 2014;12:43-9.
- Batinić Haberle I, Rebouças JS, Spasojević I. Superoxide dismutase mimics: Chemistry, pharmacology, and therapeutic potential. *Antioxid Redox Signal* 2010;13:877-918.
- Gospodaryov D, Lushchak V. Oxidative Stress: Cause and Consequence of Diseases. *Oxid Stress Dis* 2012;2:14-38. [doi: 10.5772/38093].
- Rahal A, Kumar A, Singh V, Yadav B, Tiwari R, Chakraborty S, et al. Oxidative stress, prooxidants, and antioxidants: The interplay. *Biomed Res Int* 2014;2014:761264.
- Ahmadi SM, Farhoosh R, Sharif A, Rezaei M. Structure-antioxidant activity relationships of luteolin and catechin. *J Food Sci* 2020;85:298-305.
- Romulo A. The Principle of Some *in vitro* Antioxidant Activity Methods: Review. *IOP Conf Ser Earth Environ Sci* 2020;426:1-7. [doi: 10.1088/1755-1315/426/1/012177].
- Tirzitis G, Bartosz G. Determination of antiradical and antioxidant activity: Basic principles and new insights. *Acta Biochim Pol* 2010;57:139-42.
- Deawati Y, Onggo D, Mulyani I, Hastiawan I, Kurnia D. Activity of Superoxide Dismutase Mimic of [Mn(salen)OAc] Complex Compound Non-enzymatically *In Vitro* Through Riboflavin Photoreduction. *Molekul* 2017;12:61.
- Dupont S, Fleurat Lessard P, Cruz RG, Lafarge C, Grangeteau C, Yahou F, et al. Antioxidant properties of ergosterol and its role in yeast resistance to oxidation. *Antioxidants (Basel)* 2021;10:1024.
- Chong DP. Density functional theory study of allopurinol. *Can J Chem* 2013;91:637-41.
- Farrokhnia M. Density functional theory studies on the antioxidant mechanism and electronic properties of some bioactive marine meroterpenoids: Sargahydroquinonic acid and sargachromanol. *ACS Omega* 2020;5:20382-90.
- Ozgen S, Kilinc OK, Selamoğlu Z. Antioxidant Activity of Quercetin: A Mechanistic Review. *Turk J Agric Food Sci Technol* 2016;4:1134.
- Rivero Cruz JF, Granados Pineda J, Pedraza Chaverri J, Pérez Rojas JM, Kumar Passari A, Diaz Ruiz G, et al. Phytochemical constituents, antioxidant, cytotoxic, and antimicrobial activities of the ethanolic extract of Mexican brown propolis. *Antioxidants (Basel)* 2020;9:70.

16. Soliva R, Luque FJ, Orozco M. Reliability of MEP and MEP-derived properties computed from DFT methods for molecules containing P, S and Cl. *Theor Chem Acc* 1997;98:42-9.
17. Rakhmatullin IZ, Efimov SV, Klochkov AV, Gnezdilov OI, Varfolomeev MA, Klochkov VV. NMR chemical shifts of carbon atoms and characteristic shift ranges in the oil sample. *Pet Res* 2022;7:269-74.
18. Rawat DS, Kumar D. Nature of electronic transitions and factors affecting it. *Org Spectro* 2017;12:1-21.
19. Khan N, Sagar Hossain MN, Mofiz Uddin Khan Lecturer M, Mofiz Uddin Khan N. Scopoletin and β -sitosterol glucoside from roots of *Ipomoea digitata*. *J Pharmacogn Phytochem* 2015;4:5-7.
20. Klawun C, Wilkins CL, Chem J. Optimization of functional group prediction from infrared spectra using neural networks. *Inf Comput Sci* 1996;36:69-81.
21. Khatun M, Billah M, Quader MA. Sterols and Sterol Glucoside from *Phyllanthus* Species. *Dhaka Univ J Sci* 2012;60:5-10.
22. Zhang X, Macmillan DW. Alcohols as Latent Coupling Fragments for Metallaphotoredox Catalysis: sp³-sp² Cross-Coupling of Oxalates with Aryl Halides *J Am Chem Soc* 2016. p. 1-5.
23. Muzolf-Panek M, Gliszczynska-Swigło A, Szymusiak H, Tyrakowska B. The influence of stereochemistry on the antioxidant properties of catechin epimers. *Eur Food Res Technol* 2012;235:1001-9.
24. Heim KE, Tagliaferro AR, Bobilya DJ. Flavonoid antioxidants: Chemistry, metabolism and structure-activity relationships. *J Nutr Biochem* 2002;13:572-84.
25. Godlewska Żyłkiewicz B, Świsłocka R, Kalinowska M, Golonko A, Świdorski G, Arciszewska Ż, et al. Biologically active compounds of plants: Structure-related antioxidant, microbiological and cytotoxic activity of selected carboxylic acids. *Materials (Basel)* 2020;13:4454.
26. Saptarini NM, Wardati Y. Effect of extraction methods on antioxidant activity of papery skin extracts and fractions of Maja Cipanas onion (*Allium cepa* L. var. *ascalonicum*). *ScientificWorldJournal* 2020;2020:1-6.
27. Rice Evans C. Flavonoid antioxidants. *Curr Med Chem* 2001;8:797-807.
28. Safna Hussan KP, Shahin Thayyil M, Rajan VK, Muraleedharan K. DFT studies on global parameters, antioxidant mechanism and molecular docking of amlodipine besylate. *Comput Biol Chem* 2019;80:46-53.
29. Pottachola S, Kaniyantavida A, Karuvathodiyil M, Density Functional Theory – Recent Advances, New Perspectives and Applications. Croatia: IntechOpen; 2012. p. 1-18.
30. Xue Y, Zheng Y, An L, Dou Y, Liu Y. Density functional theory study of the structure-antioxidant activity of polyphenolic deoxybenzoins. *Food Chem* 2014;151:198-206.
31. Hernandez DA, Rodriguez Zavala JG, Tenorio FJ. DFT study of antioxidant molecules from traditional Japanese and Chinese teas: comparing allylic and phenolic antiradical activity. *Struct Chem* 2020;31:359-69.
32. Rajan VK, Ragi C, Muraleedharan K. A computational exploration into the structure, antioxidant capacity, toxicity and drug-like activity of the anthocyanidin “petunidin”. *Heliyon* 2019;5:e02115.
33. Mahmoudi S, Dehkordi MM, Asgarshamsi MH. Density functional theory studies of the antioxidants-a review. *J Mol Model* 2021;27:271.

Potential antioxidant and antiradical agents from *Allium ascalonicum*: Superoxide dismutase and density functional theory in silico studies

ORIGINALITY REPORT

14%

SIMILARITY INDEX

9%

INTERNET SOURCES

13%

PUBLICATIONS

8%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to University of Newcastle Student Paper	2%
2	www.srpc.ca Internet Source	2%
3	Riza Andriani, Toto Subroto, Safri Ishmayana, Dikdik Kurnia. "Enhancement Methods of Antioxidant Capacity in Rice Bran: A Review", <i>Foods</i> , 2022 Publication	2%
4	www.japtr.org Internet Source	1%
5	<i>Methods in Molecular Biology</i> , 2010. Publication	1%
6	Venkatachalam Vijayalakshmi, Narayanasamy Nivetha, Arumugam Thangamani. "Synthesis, molecular docking, anti-cancer activity, and in-silico ADME analysis of novel spiroacenaphthylene pyrrolizidine	1%

derivatives", Journal of Molecular Structure,
2022

Publication

7	Zenika Febian Ramadhanty, Dikdik Kurnia, Boima Situmeang, Mieke Hemiawati, Nur Asmah. "Antibacterial and Antioxidant Superoxide Anion Radical Inhibitors from Myrmecodia pendans: An In silico Study", The Natural Products Journal, 2023 Publication	1 %
8	Submitted to Universitas Airlangga Student Paper	1 %
9	www.eurekaselect.com Internet Source	1 %
10	www.researchgate.net Internet Source	1 %
11	Safna Hussan K. P, Thekkekara D. Babu, Pareeth C. M, Greena Joshy, Deepu Mathew, M. Shahin Thayyil. "Antioxidant activity of erlotinib and gefitinib: theoretical and experimental insights", Free Radical Research, 2022 Publication	1 %
12	repository.qu.edu.iq Internet Source	1 %
13	vibdoc.com Internet Source	1 %

14

Houssem Boulebd. "The role of benzylic-allylic hydrogen atoms on the antiradical activity of prenylated natural chalcones: A Thermodynamic and Kinetic Study", Journal of Biomolecular Structure and Dynamics, 2020

Publication

<1 %

Exclude quotes On

Exclude matches < 15 words

Exclude bibliography On