

Bukti cover-turnitin journal health of science

Journal of Health Sciences

Register Login

Home Latest articles Current issue Archives For authors About News

Search

About the Journal

The Journal of Health Sciences is an international, open access, peer-reviewed, and evidence-based scientific journal published by the University of Sarajevo, Faculty of Health Studies (UNSA-FHS: www.unsa.ba/en - www.fhs.unsa.ba/)

The journal's mission is to promote excellence in nursing and a range of disciplines and specialties of [allied health professions](#). It welcomes submissions from international academic and health professionals community. The journal publishes evidence-based articles with solid and sound methodology, clinical application, description of best clinical practices, and discussion of relevant professional issues or perspectives. Articles can be submitted in the form of research articles, reviews, case reports, and letters to the editor or commentaries.

The journal's priorities are papers in the fields of nursing, physical therapy, medical laboratory science, environmental health, and medical imaging and radiologic technologies. Relevant articles from other disciplines of [allied health professions](#) may be considered for publication.

It follows publishing standards set by the International Committee of Medical Journal Editors (ICMJE), Committee on Publication Ethics (COPE), and the World Association of Medical Editors (WAME).

This journal is indexed or abstracted by [SCOPUS](#), [EBSCO](#), [DOAJ](#), [Scilis](#), [Google Scholar](#), [Dimensions](#), [Researcher](#), [Scimago](#), [PROQUEST](#), [ScienceOpen](#), [MIAR](#), [Bielefeld Academic Search Engine \(BASE\)](#), [Index Copernicus](#), [CAB Abstracts](#), [ULRICH'S](#), [Academic Journal Database](#), [Elektronische Zeitschriftenbibliothek](#), [Geneva Foundation for Medical Education and Research](#), and other libraries.



Current issue

Archives

Submit a manuscript

Announcements

December 2023 issue published

Journal of Health Sciences

Register Login

Home Latest articles Current issue Archives For authors About News

Search

Home / Editorial Team

Editorial Team

Editor-in-chief

Arzija Pasalić, Faculty of Health Studies, University of Sarajevo, Sarajevo, *Bosnia and Herzegovina*

Managing Editor

Adnan Šehić, Faculty of Health Studies, University of Sarajevo, Sarajevo, *Bosnia and Herzegovina*

Statistical Editor

Anes Jogunčić, Faculty of Health Studies, University of Sarajevo; Public Health Institute of Canton Sarajevo, Sarajevo, *Bosnia and Herzegovina*

Editorial Board

Adnan Beganović, Department for medical physics, Clinical Center of Sarajevo University, Sarajevo, *Bosnia and Herzegovina*
Aida Pilav, Faculty of Health Studies, University of Sarajevo, Sarajevo, *Bosnia and Herzegovina*
Amra Mačak Hadžomerović, Faculty of Health Studies, University of Sarajevo, Sarajevo, *Bosnia and Herzegovina*
Amila Jaganjac, Faculty of Health Studies, University of Sarajevo, Sarajevo, *Bosnia and Herzegovina*
Andrej Starc, Faculty of Health Sciences, University of Ljubljana, Ljubljana, *Slovenia*



Current issue

Archives

Submit a manuscript

Announcements

December 2023 issue published

2023 December 30
The December 2023 issue of the journal of Health Sciences is

Bakir Katana, Faculty of Health Studies, University of Sarajevo, Sarajevo, *Bosnia and Herzegovina*
Borut Poljsak, Laboratory of Oxidative Stress Research, University of Ljubljana, Ljubljana, *Slovenia*
Darija Vukić-Lušić, Faculty of Medicine, University of Rijeka, Rijeka, *Croatia*
Fuad Julardžija, Faculty of Health Studies, University of Sarajevo, Sarajevo, *Bosnia and Herzegovina*
Isabelle Richard, Department of Physical Medicine and Rehabilitation, CHU of Angers Rue des Capucins, Angers, *France*
Jasmina Mahmutović, Faculty of Health Studies, University of Sarajevo, Sarajevo, *Bosnia and Herzegovina*
Jonathan McNulty, Graduate Taught Studies, School of Medicine, University College Dublin, Dublin, *Ireland*
Luca Testarelli, Department of Oral Sciences and Maxillofacial Surgery, Sapienza University of Rome, *Italy*
Maja Cigrovski Berković, Faculty of Kinesiology, University of Zagreb, Zagreb, *Croatia*
Maja Miškulin, Faculty of Medicine, University of Osijek, Osijek, *Croatia*
Majda Pajnikhar, Institute for Nursing Care, Faculty of Health Sciences, University of Maribor, Maribor, *Slovenia*
Martina Lovrić Benčić, Clinical Hospital Center Zagreb Rebro, Zagreb, *Croatia*
Mirna Habuda Stanić, Faculty of Food Technology, "Josip Juraj Strossmayer" University of Osijek, *Croatia*
Muhammed Ajanović, Faculty of Dentistry, University of Sarajevo, Sarajevo, *Bosnia and Herzegovina*
Nachiappan Chockalingam, Faculty of Health Sciences, Staffordshire University, Stoke-on-Trent, *United Kingdom*
Naris Poljskić, Institute for Genetic Engineering and Biotechnology, Sarajevo, *Bosnia and Herzegovina*
Nejc Mekš, Faculty of Health Sciences, University of Ljubljana, Ljubljana, *Slovenia*
Rijad Konjhodžić, Faculty of Health Studies, University of Sarajevo, Sarajevo, *Bosnia and Herzegovina*
Samir Bojičić, Faculty of Health Studies, University of Sarajevo, Sarajevo, *Bosnia and Herzegovina*
Selma Jakupović, Faculty of Dentistry, University of Sarajevo, Sarajevo, *Bosnia and Herzegovina*
Senka Mesihović-Dinarević, Academy of Sciences and Arts of *Bosnia and Herzegovina*
Theodoros B. Grivas, Department of Orthopedic and Traumatology, "Tzaneio" General Hospital, Piraeus, *Greece*
Valentina Colonnello, Department of Experimental, Diagnostic and Specialty Medicine, University of Bologna, Bologna, *Italy*
Vedran Đido, Faculty of Health Studies, University of Sarajevo, Sarajevo, *Bosnia and Herzegovina*
Zarema Obradović, Faculty of Health Studies, University of Sarajevo, Sarajevo, *Bosnia and Herzegovina*
Zhaohui Su, Center on Smart and Connected Health Technologies, Mays Cancer Center, School of Nursing, UT Health San Antonio, San Antonio, Texas, *United States of America*

Editorial associate

Amna Pezo, Faculty of Health Studies, University of Sarajevo, Sarajevo, *Bosnia and Herzegovina*

published, it contains research articles in radiology technology, laboratory engineering, mental health, and more.

September 2023 issue published

2023 October 6

The September 2023 issue of the Journal of Health Sciences is published. It contains research articles in physical therapy, midwifery, laboratory engineering, nursing, and more.

[Show all announcements ...](#)

Newsletter

Stay up to date with our occasional newsletter

[Subscribe](#)

Latest publications

[Current issue](#)
[Archives](#)
[Submit a manuscript](#)

Journal of Health Sciences

[Home](#) [Latest articles](#) [Current issue](#) [Archives](#) [For authors](#) [About](#) [News](#)

[Search](#)

[Home](#) / [Archives](#) / Vol. 13 No. 2 (2023)

Vol. 13 No. 2 (2023)

Published: 30.09.2023

Research articles

Table 1. Studies from the Faculty of Health Studies comparing the prevalence of stress and anxiety among students during the COVID-19 pandemic.

Study	Year	N	Stress (%)	Anxiety (%)
Alma Gavranić-Giamoć et al.	2020	100	100	100
Alma Gavranić-Giamoć et al.	2021	100	100	100
Alma Gavranić-Giamoć et al.	2022	100	100	100
Alma Gavranić-Giamoć et al.	2023	100	100	100
Alma Gavranić-Giamoć et al.	2024	100	100	100
Alma Gavranić-Giamoć et al.	2025	100	100	100
Alma Gavranić-Giamoć et al.	2026	100	100	100
Alma Gavranić-Giamoć et al.	2027	100	100	100
Alma Gavranić-Giamoć et al.	2028	100	100	100
Alma Gavranić-Giamoć et al.	2029	100	100	100

© 2023 Published online by John Wiley & Sons, Ltd. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Association between stress and self-reported bruxism among students from University of Sarajevo during the COVID-19 pandemic

Alma Gavranić-Giamoć, Zinejda Šabić, Selma Alić-Drina, Sanela Strujić-Porović, Selma Jakupović, Alma Kamber, Emir Berhamović, Lejla Berhamović, Amela Džonlagić, Bakir Katana, Lejla Kazazić

[PDF](#)

51-58



[Current issue](#)

[Archives](#)

[Submit a manuscript](#)

TABLE 2. Analysis of body weight, body length and body mass index of the respondents.

Groupings	n	AG	22	Mean	Stdev
Weight					
Body weight (kg) (p=0.03, p<0.001)					
Group 1	30	87.54	-0.36	10.05	144.35
Group 2	30	86.38	-0.62	10.02	135.58
Total	60	86.92	-0.39	10.03	140.50
Height (cm) (p=0.228, p<0.001)					
Group 1	30	173.03	0.19	6.02	150.56
Group 2	30	173.22	0.09	6.00	152.59
Total	60	173.08	0.14	6.01	149.50
HR (p=0.2, p=0.001)					
Group 1	30	27.89	0.47	21.48	80.45
Group 2	30	27.76	0.78	21.29	76.21
Total	60	27.82	0.61	21.38	80.81

© 2000 Blackwell Science Ltd, *Journal of Internal Medicine* 247: 399–406

© 2004 Blackwell Publishing Ltd, *Journal of Internal Medicine* 255: 103–110

Year	N	%
Never help		
Other	21	66.9
Black	4	11.9
Hispanic	22	68.9
Multiethnic of Two Races	2	3.7
Total	55	100
Never help		
Other	1	1.7
Black	22	58.8
Hispanic	20	43.1
Total	43	100

© 2002 Blackwell Science Ltd *Journal of Internal Medicine* 252: 105–112

[illegible]

1978) suggest that the observed increase in the proportion of *Chironomus tentans* in the sediments of the Tiber River may be due to the fact that this species is more resistant to pollution than other species of the genus *Chironomus* (Bertoni 1978).

Comparative analysis of different physical therapy programs in the treatment of people with knee osteoarthritis

Bakir Katana, Amra Maćak Hadžibierović, Eldad Kaljić, Adnan Mujezinović, Sead Kojić, Bojana Stanišić, Jasmina Bajrović, Marcella Niková, Ademir Spahić, Samir Boyčik

PDF

The effect of the application of Kinesio Tape on pain relief in musculoskeletal disorders

Dinko Remić, Samir Bojčić, Amra Maćak Hadžomerović, Amila Jaganjac, Eldad Kaljić, Dženan Pleho, Bakir Katana 77-83

PDF

Inflammatory and endothelial dysfunction markers in females with metabolic syndrome aged 50-75 years

Ruspanini, Raditya Wratsangka, Merianti Lie Tanu, Melivanti, Rismawati Yaswir 114-90

PDF

Announcements

December 2023 issue
published

2023 December 30
The December 2023 issue of the Journal of Health Sciences is published. It contains research articles in radiology technology, laboratory engineering, mental health, and more.


September 2023 issue
published

2023 October 6
The September 2023 issue of the Journal of Health Sciences is published. It contains research articles in physical therapy, midwifery, laboratory engineering, nursing, and more.

[Show all announcements](#)

Newsletter


Stay up to date with our occasional newsletter



Quality of life of families with children with intellectual and developmental disabilities: Family health domain

Anyka Yukićević, Emira Švraka, Amra Mačak Hadžomerović, Naim Salkić


91-97



R517817449 FTO gene variation associated with familial disease burden rather than individual risk for breast cancer

Berisa Hasanbegović, Belmina Sarić Medić, Emir Sokolović, Nikola Tomašević, Lejla Pojskić

88-104



Knowledge, attitude, and preventive practices of leptospirosis affected populations in South Andaman, India: A cross-sectional study

Amreen Fatema, Manjunatha Ramu, Kannan Thiruvengadam, Iitoo Pulikkottil, Sunithi, Paluru Vijayachari

105-112



TABLE 1: Correlation between age, sex, and inflammatory markers in COVID-19 patients with mild and moderate disease. (P < 0.05 is significant)

Parameter	Age	Sex	WBC	CRP	ESR	IL-6	IL-10	IL-17
Age	1.00	0.12	0.05	0.01	0.01	0.01	0.01	0.01
Sex	0.12	1.00	0.01	0.01	0.01	0.01	0.01	0.01
WBC	0.05	0.01	1.00	0.01	0.01	0.01	0.01	0.01
CRP	0.01	0.01	0.01	1.00	0.01	0.01	0.01	0.01
ESR	0.01	0.01	0.01	0.01	1.00	0.01	0.01	0.01
IL-6	0.01	0.01	0.01	0.01	0.01	1.00	0.01	0.01
IL-10	0.01	0.01	0.01	0.01	0.01	0.01	1.00	0.01
IL-17	0.01	0.01	0.01	0.01	0.01	0.01	0.01	1.00

Systemic inflammatory indexes (NLR, DNL, PLR and SII) role in predicting COVID-19 progression

Berina Hasanefendić, Emir Šehinčehajić, Armina Dedić, Suzana Tihvić-Kapidžić, Ermin Begović, Sanela Hajro, Belma Gazibera, Ahmed Velić, Melina Drijo

113-117



Journal of Health Sciences



← Show this widget in your own website

Just copy the code below and paste within your html code:

`<a href="https://www.scimaç`

newsletter

[Subscribe](#)

Latest publications





RESEARCH ARTICLE

Open Access

Inflammatory and endothelial dysfunction markers in females with metabolic syndrome aged 50–75 years

Pusparini^{1*}, Raditya Wratsangka², Lie Tanu Merijanti³, Meiyanti⁴, Rismawati Yaswir⁵

¹Department of Clinical Pathology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia, ²Department of Obstetrics Gynecology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia, ³Department of Occupational Medicine, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia, ⁴Department of Pharmacology and Pharmacy, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia, ⁵Department of Clinical Pathology, Faculty of Medicine, Andalas University, Padang, Indonesia

ABSTRACT

Introduction: The prevalence of metabolic syndrome (MetS) in Indonesia is 39% and it is higher in females than in males (46% vs. 28%). Features of this syndrome comprise low-grade chronic inflammation and endothelial dysfunction from cardiometabolic abnormalities. High-sensitive C-reactive protein (HsCRP) is a biomarker of cardiovascular disease, whereas ferritin and pentraxin-3 (PTX-3) are markers of chronic inflammation. One of the markers of endothelial dysfunction is an intercellular cell adhesion molecule-1 (ICAM-1). Studies on MetS in women aged 50–70 years and using several markers of inflammation and endothelial dysfunction are rare. The study aimed to determine differences in HsCRP, ferritin, PTX-3, and ICAM-1 level between females with and without MetS aged 50–75 years.

Methods: For this cross-sectional study, 160 females aged 50–75 years from South Jakarta were recruited, divided into MetS and non-MetS (control) groups, 80 participants each. Participants that met the inclusion criteria were assessed for vital signs and metabolic parameters (e.g., lipid profile, fasting blood glucose level, and waist circumference [WC]). The groups were compared using an independent *t*-test for inflammatory (i.e., HsCRP, PTX-3, and ferritin) and endothelial dysfunction (ICAM-1) ($p < 0.05$) biomarkers.

Results: The mean age of the MetS and control groups was 59.6 ± 5.6 and 59.2 ± 5.5 years. MetS criteria included WC of ≥ 80 cm (95.2%), increased blood pressure level (90.6%); and high-density lipoprotein cholesterol of < 50 mg/dL (76.2%). Significant differences were found between the MetS and control groups for ferritin ($p = 0.027$), PTX-3 ($p = 0.045$), and ICAM-1 ($p = 0.004$) but not for HsCRP ($p = 0.136$).

Conclusion: Elevated ferritin, PTX-3 level, and ICAM-1 level are significantly associated with an increased MetS risk in females aged 50–75 years; however, the same is not true for HsCRP.

Keywords: Ferritin; high-sensitive C-reactive protein; intercellular cell adhesion molecule-1; inflammation; metabolic syndrome; pentraxin-3

INTRODUCTION

Currently, metabolic syndrome (MetS) is developing into a global epidemic, with approximately 12–37% of Asians and 12–26% of Europeans suffering from MetS (1). The incidence of MetS in Indonesia is higher in females than in males. The prevalence of MetS in adults aged 45–65 years is 39%, namely 28% in males and 46% in females (2). MetS is a group of symptoms comprising abdominal obesity, hypertension, dyslipidemia, hyperglycemia, and prothrombotic and pro-inflammatory conditions (3).

The higher prevalence of abdominal obesity in Indonesian females, accounting for 57.5% versus 16.8% in males,

explains the higher MetS prevalence in Indonesian females than in males. Individuals with MetS are at an increased risk of morbidity and mortality from cardiovascular disease. MetS is a risk factor for the occurrence of type 2 diabetes mellitus (T2DM), cardiovascular disease, and stroke (2). The pathophysiology of cardiovascular disease in MetS probably involves risk factors such as insulin resistance, chronic low-grade systemic inflammation, genetics, dietary status, and individual habits (3,4).

Abdominal fat is an active endocrine organ that generates cytokines that affect other body tissues (4). Abdominal fat releases several inflammatory markers, such as C-reactive protein (CRP), and acute-phase proteins, such as pentraxin-3 (PTX-3) (3). The level of plasma PTX-3 was inversely related to MetS in the Korean population (5). Another acute-phase protein is ferritin (6,7). Ferritin is not only a marker of iron reserves in the body but may also be a marker of inflammation (6). Abnormal cardiometabolic

Corresponding author: Pusparini, Department of Clinical Pathology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia.
E-mail: pusparini@trisakti.ac.id

Submitted: 27 October 2022/Accepted: 12 May 2023

DOI: <https://doi.org/10.17532/jhsci.2023.2003>



components contribute to the occurrence of endothelial dysfunction followed by blood vessel inflammation, resulting in atherosclerosis and cardiovascular disease (8). Intercellular cell adhesion molecule-1 (ICAM-1) is an endothelial dysfunction marker (8,9). Many studies have reported the relationship between MetS and inflammatory markers, but MetS occurrence varies by population, ethnicity, and sex (4,8,10). Studies on MetS conducted in females aged 50–75 years and using several inflammation (high-sensitive CRP [HsCRP], PTX-3, and ferritin) and endothelial dysfunction (ICAM-1) markers are rare (4,6,8–10). This study aimed to determine differences in the levels of inflammatory and endothelial dysfunction biomarkers between females with and without MetS aged 50–75 years.

METHODS

This cross-sectional study was conducted from January to June 2021 in the catchment area of *Puskesmas* in Mampang District, South Jakarta. The investigators enrolled 80 females aged 50–75 years with MetS and 80 non-MetS females (controls). Inclusion criteria for the MetS group were the presence of three MetS criteria, i.e., waist circumference (WC) of ≥ 80 cm, systolic blood pressure (SBP) level of ≥ 130 mmHg or diastolic blood pressure (DBP) level of ≥ 85 mmHg or taking antihypertensive medications; fasting blood glucose (FBG) level of ≥ 100 mg/dL; high-density lipoprotein (HDL) cholesterol level of < 50 mg/dL; triglyceride level of ≥ 150 mmHg; no active infection in 2 weeks before the study, not smoking, not consuming alcohol, and providing written informed consent. Exclusion criteria for the MetS and control groups in this study comprised the administration of corticosteroids, oral anticoagulants, and hormone replacement therapy; history of malignancies; or abnormal renal function. The MetS criteria employed in this study followed the revised National Cholesterol Education Programme's Adult Treatment Panel III criteria for Asians (1).

The participants were interviewed by five individuals (having at least senior high school or equivalent education) with knowledge of the aims of the study and trained in the filling of the questionnaire (including demographic characteristics, concomitant disease, smoking status, alcohol drinking status, concomitant medication usage, and anthropometrics) (Table 1). The filling-in of the questionnaire was completed in 2 weeks. Following physical examination (vital signs); anthropometry; and laboratory bloodwork for lipid profile, FBG, and creatinine, participants were divided into MetS and control groups. The investigators determined the blood pressure level, body weight, height, WC, and hip circumference of participants through physical examination. Venous blood was drawn (following a 12-h fast) using a 10-mL plain vacutainer for FBG and a complete lipid profile. An aliquot of the blood sample was used to obtain serum, which was stored at -70°C for determination of PTX-3, ICAM-1, HsCRP, and ferritin levels in participants. Since differences in intraday repeatability of laboratory procedures may cause spurious differences in the results between the MetS and control groups, the concentrations of PTX-3, ICAM-1, HsCRP, and ferritin were simultaneously determined, to ensure that differences in PTX-3, ICAM-1, HsCRP, and ferritin levels were not

due to laboratory errors but were true differences. The biomarkers were assessed using the following techniques: PTX-3 through enzyme-linked immunosorbent assay (ELISA) using Quantikine® ELISA Human PTX-3/TSG-14 reagents (R&D Systems, Inc., Minneapolis, USA) and Biorad model 680 microplate reader; ICAM-1 via ELISA using Quantikine® ELISA Human ICAM-1 reagent (R&D Systems, Inc., Minneapolis, USA) and Biorad model 680 microplate reader; HsCRP using immunoturbidimetry, Architect 6K26-30 reagents, and Architect c-8000 analyzer; and ferritin via chemiluminescent microparticle immunoassay using Architect 7K59 reagents and Architect i-2000 analyzer. This study was approved by the Ethics Committee, Faculty of Medicine, Universitas Trisakti (approval no. 165/KER/FK/XII/2020). Independent t-test was used to determine differences in biomarker levels between the MetS and control groups, and $p < 0.05$ were considered to indicate statistical significance.

RESULTS

The characteristics of participants with MetS and controls are summarized in Table 2. The SBP, DBP, WC, and hip circumference of participants were considerably different between the two groups (p -values of 0.000, 0.016, 0.009, and 0.006, respectively). The biochemical characteristics of participants are listed in Table 3. The MetS group showed significantly increased triglyceride and FBG and reduced HDL cholesterol levels compared with the control group. PTX-3, ferritin, and ICAM-1 level (p -values of 0.045, 0.027, and 0.004, respectively) were significantly higher in the MetS than in the control groups, whereas HsCRP levels did not indicate any significant differences. In the MetS group, the five MetS parameters presented in the order of their frequency were: WC of ≥ 80 cm, increased blood pressure level, HDL cholesterol level of < 50 mg/dL, increased FBG level, and triglyceride level of ≥ 150 mg/dL.

DISCUSSION

In the present study, out of the five MetS-contributing elements, the most frequently encountered in the participants with MetS were: WC of ≥ 80 cm (97.5%) and increased blood pressure (91.3%), followed by HDL cholesterol concentration of < 50 mg/dL (80%). The study by Sigit et al. (2) showed that the predominant MetS components in the Indonesian population were hypertension (61.3%; 56.2% in males vs. 64.6% in females) and hyperglycemia (51%; 51% in males vs. 50.9% in females), and hypertension (61.7%; 70% in males vs. 55.2% in females) and abdominal obesity (40%; 34.6% in males vs. 44.2% in females) in the Dutch population. The prevalence of abdominal obesity was approximately 57.5% in Indonesian females and 16.8% in Indonesian males (2). The aforementioned study highlighted that MetS in Indonesia was more common in females (46.2%) than in males (28%) (2). The differences between the study by Sigit et al. (2) and the current study may have occurred because participants in the current study were all females with a mean age of 59.6 ± 5.6 years in the MetS group and 59.2 ± 5.5 years in the control group, whereas Sigit et al. (2) included male and female participants, with a mean

TABLE 1. Questionnaire used during the enrollment process (English translation)

General Information	
Name:	
Address:	
Telephone/Cellphone number:	
Interviewer:	
Date of interview:	
Circle option number according to your answer to items No. 1–10 (Please include photocopy of your National Identity Card)	
1. Age (years, months):	
2. Date of birth (DD/MM/YY):	
3. Educational background	
1. None 2) Elementary School 3) Junior High School 4) Senior High School 5) Academy/Bachelor 6) Did not complete elementary school	
4. Current employment status	
1. Unemployed 2) Trader 3) Entrepreneur 4) Other	
5. Are you in menopause ?	
1. Yes 2) No	
6. If in menopause, give duration of menopause.....year (s);.....month (s)	
7. Did you have any infection in the past 2 weeks?	
1. Yes 2) No	
8. Do you have any of the following diseases or a past history of these diseases?	
a. Hypertension	1) Yes 2) No
b. Diabetes Mellitus	1) Yes 2) No
c. Kidney disease	1) Yes 2) No
d. Stroke	1) Yes 2) No
e. Cancer	1) Yes 2) No
f. Other	1) Yes 2) No
If yes, state the diagnosis	
9. Do you take any of the following drugs?	
a. Corticosteroids	1) Yes 2) No
b. Hormonal replacement therapy	1) Yes 2) No
c. Antihypertensive drugs	1) Yes 2) No
d. Antidiabetic drugs	1) Yes 2) No
e. Antihyperlipidemic drugs	1) Yes 2) No
f. Oral anticoagulants	1) Yes 2) No
10. Do you smoke?	1) Yes 2) No
If yes, how many cigarettes per day	
11. Do you consume alcoholic drinks?	1) Yes 2) No
If yes, how many glasses per day	

age of 52.2 ± 5.8 years in the Indonesian population (2). The prevalence of MetS increases with age, as reported in a Turkish study: 11.9% (age 20–29 years), 23.4% (age 30–39 years), 26.9% (age 40–49 years), 40.4% (age 50–59 years), 45.3% (age 60–69 years), and 26.3% (age >70 years) (11). Mehndiratta et al. (12) conducted a study on 200 women aged 45–60 years receiving ambulatory treatment in an obstetrics and gynecology department at Amritsar, Punjab, India. The study results indicated that 29% of women were found to have newly onset MetS, with an incidence of 16% in the premenopausal group and 42% in the postmenopausal group (12). MetS incidence is higher in postmenopausal women than in premenopausal women, because following menopause, the estrogen-to-testosterone ratio decreases, dilutes the vasorelaxant effects of estrogen on the vessel wall and promotes the production of vasoconstrictor factors such as endothelin which lead to metabolic culminating into MetS (12).

TABLE 2. Characteristics of study participants

Parameter	X $\pm$ SD/n (%) MetS n=80	X $\pm$ SD/n (%) Control n=80	p-value
Age (years)	59.6 $\pm$ 5.6	59.2 $\pm$ 5.5	0.256
Weight (kg)	60.4 $\pm$ 10.2	55.3 $\pm$ 8.9	0.173
Height (cm)	152.5 $\pm$ 5.7	152.3 $\pm$ 5.3	0.532
BMI (kg/m ²)	25.9 $\pm$ 4.6	21.5 $\pm$ 3.5	0.076
Pulse rate	84.9 $\pm$ 14.2	80.6 $\pm$ 11.9	0.060
Blood pressure level			
Systolic (mmHg)	150.5 $\pm$ 23.0	127.6 $\pm$ 23.7	0.000*
Diastolic (mmHg)	88.4 $\pm$ 12.1	79.5 $\pm$ 10.7	0.016*
Waist circumference (cm)	91.7 $\pm$ 7.2	77.8 $\pm$ 8.0	0.009*
Hip circumference (cm)	99.2 $\pm$ 7.8	90.4 $\pm$ 6.5	0.006*
Education			
None	3 (3.8)	2 (2.5)	0.785
Elementary school	30 (37.5)	27 (33.7)	
Junior high school	21 (26.3)	20 (25)	
Senior high school	19 (23.7)	26 (32.5)	
Academy/Bachelor	5 (6.2)	3 (3.8)	
Did not complete elementary school	2 (2.5)	2 (2.5)	
Employment			
Unemployed	58 (72.4)	57 (71.3)	0.482
Trader	5 (6.3)	6 (7.5)	
Entrepreneur	5 (6.3)	3 (3.7)	
Other	12 (15)	14 (17.5)	
Menopause			
Yes	72 (90)	73 (91.3)	0.348
No	8 (10)	7 (8.7)	
Duration of menopause (years)	10.11 $\pm$ 2.3	10.7 $\pm$ 2.1	0.753
History of diabetes mellitus	16 (20)	0 (0)	
History of hypertension	47 (58.8)	0 (0)	
Medications			
Antihypertensive	32 (40)	0 (0)	
Antidiabetic	9 (11.3)	0 (0)	
Antihyperlipidemic	12 (15)	0 (0)	

* $p < 0.05$ =significant difference (independent *t*-test); Chi-square test. BMI: Body mass index

TABLE 3. Biochemical and inflammatory markers

Parameter	X $\pm$ SD MetS n=80	X $\pm$ SD Non-MetS n=80	p-value
Biochemical			
Total cholesterol (mg/dL)	215.8 $\pm$ 15.9	208.4 $\pm$ 13.7	0.286
Triglycerides (mg/dL)	145.8 $\pm$ 10.5	94.3 $\pm$ 12.8	0.000*
HDL cholesterol (mg/dL)	48.8 $\pm$ 9.8	58.9 $\pm$ 10.5	0.000*
LDL cholesterol (mg/dL)	139.9 $\pm$ 18.7	132.2 $\pm$ 10.7	0.198
Fasting blood glucose (mg/dL)	124.7 $\pm$ 13.5	88.6 $\pm$ 9.3	0.000*
Creatinine (mg/dL)	0.77 $\pm$ 0.2	0.79 $\pm$ 0.3	0.686
eLFG (mL/min)	85.5 $\pm$ 17.4	85.7 $\pm$ 17.9	0.931
Inflammatory			
Pentraxin-3 (ng/dL)	0.81 $\pm$ 0.20	0.72 $\pm$ 0.19	0.045*
HsCRP (mg/L)	3.1 $\pm$ 0.6	2.6 $\pm$ 0.3	0.136
Ferritin (μ g/L)	165.9 $\pm$ 14.7	125.7 $\pm$ 8.7	0.027*
ICAM-1 (ng/mL)	356.6 $\pm$ 9.9	316.1 $\pm$ 9.5	0.004*

* $p < 0.05$ significant difference (independent *t*-test). HDL: High-density lipoprotein, LDL: Low-density lipoprotein, HsCRP: High-sensitive C-reactive protein, ICAM-1: Intercellular cell adhesion molecule-1

This study demonstrated no significant differences were found in educational level (Table 2; $p = 0.785$) and

employment status (Table 2; $p = 0.482$) between the MetS and control group. Most of the subjects in the two groups had an educational level of senior high school and below (MetS 91.3% vs. non-MetS 93.7%) and were not anymore employed (MetS 72.4% vs. non-MetS 71.3%). In this study, no distinction was made between senior high school and tertiary education. Research in China on 3175 respondents aged ≥ 45 years showed that MetS prevalence was quite high in the rural population with low income and low education (13). Stephens et al. reported that a higher education level correlates to significantly better metabolic health when compared to a lower education level and carries significantly less risk for WC, SBP, glucose and glycosylated hemoglobin, triglycerides, HDL cholesterol, and MetS (all $p < 0.05$), except for DBP, basal insulin, uric acid, low-density lipoprotein cholesterol, and total cholesterol (14).

Low-grade chronic inflammation is identified as the leading factor for the occurrence of MetS and subsequent clinical symptoms (15,16). Among the four biomarkers assessed in this study, ferritin, ICAM-1, and PTX-3, but not HsCRP, showed significant differences between the MetS and control groups (Table 3). The threshold values of HsCRP (<1 , $1-3$, and $3-10$) indicate low, moderate, and high risk of cardiovascular disease, respectively (17). The MetS group had higher mean HsCRP levels than the control group based on these criteria. The mean HsCRP level in the MetS group was 3.1 ± 0.4 mg/L, indicating a high risk, whereas that in the control group was 2.6 ± 0.3 mg/L, indicating a moderate risk, although no significant differences were observed between the two groups. Carrero et al. (17) studied myocardial infarction (MI) survivors (17,464 participants aged >18 years) undergoing HsCRP testing for >30 days after MI and revealed that most patients with MI had increased HsCRP levels. HsCRP may be used as a prognostic factor for determining the occurrence of cardiovascular disease (17). Yilmaz et al. (18) analyzed 100 females aged 20–45 years with obesity and revealed that HsCRP, tumor necrosis factor (TNF)- α , and malondialdehyde may be used as diagnostic markers of MetS in females with obesity. Furthermore, WC is correlated with HsCRP ($r = 0.315$; $p = 0.003$) (18). The current study revealed a correlation between WC and HsCRP, both in the MetS ($r = 0.353$, $p = 0.002$; data not shown) and control ($r = 0.560$, $p = 0.003$) groups. A WC of >80 cm in females indicates the occurrence of visceral fat accumulation. Visceral fat is an endocrine organ that can secrete pro-inflammatory substances, such as leptin, adiponectin, resistin, TNF- α , and interleukin-6, which play a significant role in the occurrence of MetS (18). The study of Hong et al. in China revealed that after a follow-up period of 5 years, HsCRP was linked to MetS, but only in females. The reason why females are at higher risk for MetS than males remains unclear, but physiologically, females may have more visceral fat than males, which is associated with MetS risk (16). Four estrogens have been identified in humans: estrone, estradiol, estriol, and estetrol. Estradiol is the predominant estrogen in women during reproduction. Post-menopausal estrogen production in the ovaries is reduced therefore circulatory estrogen concentrations are also drastically reduced. After menopause, estradiol is replaced by estrone, synthesized in adipose tissue from adrenal dehydroepiandrosterone (19). The current study

revealed that the HsCRP level did not significantly differ between the MetS and control groups, probably because the body mass index (BMI) of both groups was not significantly different ($p = 0.076$). A study in China revealed that BMI was correlated with HsCRP (10). In addition, 90% of our participants were in menopause, both in the MetS and control groups. Estrogen has pleiotropic effects and plays a crucial role in preserving endothelial functions (20). Estrogen increases the bioavailability of nitric oxide by promoting its synthesis and exerts direct and indirect antioxidant and anti-inflammatory effects. The decreased estrogen level in menopause causes endothelial dysfunction followed by vascular inflammation (20). Furthermore, several drugs used in the treatment of metabolic risk factors, such as statins, nicotinic acid, fibrates, angiotensin-converting enzyme inhibitors, and thiazolidinediones, have been reported to decrease CRP levels (21). Among the MetS group in the present study, 40% used anti-hypertensive drugs, 11.3% used anti-diabetics, and 15% used anti-hyperlipidemic drugs (Table 2). Silva et al. (22) reported that different anti-hypertensive drugs have different effects on inflammatory markers and markers of endothelial function. Although angiotensin-converting enzyme inhibitors and angiotensin receptor blockers do beneficially affect endothelial function and inflammation, first and second-generation beta-blockers have no such effect (22). A systematic review on the effect of statin therapy on inflammatory markers showed that statin therapy has lipid-lowering and anti-inflammatory effects (23). Forty-one participants with a BMI of $25-39.9$ kg/m² and impaired glucose tolerance test were randomized to take simvastatin or metformin for 16 weeks. The study demonstrated a significant reduction in CRP and interleukin-6 concentration at the completion of the study (24).

PTX-3 is both an inflammatory marker and an acute-phase protein (25-27). Obesity and MetS are linked to chronic inflammation (27). This study revealed that PTX-3 levels were significantly higher in the MetS group than in the control group ($p = 0.045$) (Table 3). This is consistent with the study by Karakas et al., indicating that the MetS group has higher PTX-3 levels than the control group (26). Moreover, Karakas et al. found that the PTX-3 levels in the group with 5 MetS components (0.90 ± 0.06 ng/dL) were higher than in the groups with 4 MetS components (0.63 ± 0.12 ng/dL) and 3 MetS components (0.58 ± 0.11 ng/dL) (26). In our study, the mean PTX-3 levels in the MetS and control groups were 0.81 ± 0.20 and 0.72 ± 0.19 ng/dL, respectively. The mean PTX-3 level in the MetS group was higher than that reported by Karakas et al. (0.58 ± 0.11 ng/dL) with the same number of components of MetS criteria (26). The participants in our study were older than those in the study by Karakas et al. (approximately 49 years of age). Accordingly, our study participants might have had MetS for a longer period than those in the study of Karakas et al. Nevertheless, our study is in line with the study of Karakas et al. that PTX-3 level is correlated with MetS severity. Kizilgul et al. (3) stated that a number of study results related to PTX-3 level in MetS were contradictory. For instance, a study conducted on 1749 healthy Japanese participants (818 men and 931 women) in the age range of 37–87 years indicated that PTX-3 levels were lower in participants with MetS than in the non-MetS group and

that the PTX-3 levels were negatively correlated with triglyceride levels and BMI (28). Zanetti et al. (29) reported that plasma PTX-3 levels were higher in the MetS group with subclinical atherosclerosis than in the control group. Ogawa et al. (30) reported low PTX-3 and high CRP levels in males with obesity and/or MetS compared to the control group. Ogawa et al. (30) study indicate that the reduction in plasma PTX-3 may accelerate chronic inflammation and atherosclerosis in the obese and/or MetS population because of the tissue protective and antiprotective roles of PTX-3 (30). Kizilgul et al. (3) compared PTX-3 levels in patients with MetS, in patients with rheumatoid arthritis, and with healthy controls. Their results do not demonstrate an increase in the levels of PTX-3 as chronic inflammatory markers in patients with MetS (3). The symptoms of MetS are specific for each age group and population and therefore each group or population should evolve its own strategy for MetS prevention (2,25).

PTX-3, a marker of cardiovascular risk, is produced by various cells, such as adipocytes, macrophages, dendritic cells, fibroblasts, and vascular endothelial cells, and is released in response to inflammatory stimuli (26,27). The current study revealed that PTX-3 was negatively correlated with HDL cholesterol ($r = -0.253$ and $p = 0.028$) in the MetS group, whereas no significant correlation was noted in the control group (data not shown). This indicates that HDL cholesterol plays a role in inhibiting PTX-3. This is consistent with the study of Kardas et al., which showed that children with HDL cholesterol levels of <40 mg/dL have higher PTX-3 levels than those with normal HDL cholesterol levels ($p < 0.05$) (27). A study on overweight children and children with obesity indicated that the mean PTX-3 level was significantly higher in the overweight (10.23 ± 4.42 ng/mL) and obese (11.20 ± 4.12 ng/mL) groups compared to that in the control group (7.93 ± 4.35 ng/mL). In addition, PTX-3 level was linked to intimal and epicardial fat thickness in overweight children. PTX-3 may be an early marker of cardiovascular risk in overweight children (31).

Ferritin is an essential protein regulating iron homeostasis in the body. Serum ferritin is used as a biomarker to determine the status of iron reserves in the body. Excessive iron accumulation may lead to different pathological conditions, such as hemochromatosis and hemosiderosis (7). In addition, serum ferritin is an acute-phase protein associated with insulin resistance. Iron accumulation interferes with pancreatic insulin synthesis and secretion and affects hepatic insulin-extracting capacity, resulting in peripheral hyperinsulinemia and impaired insulin secretion (32). Our study results revealed significant differences in serum ferritin levels between the MetS and control groups ($p = 0.027$) (Table 3). The mean ferritin level in the MetS group was significantly higher than that in the control group. Chand et al. revealed comparable results; the serum ferritin levels were higher in the MetS group than in the control group ($p < 0.0001$) (32). Our study findings are consistent with those of Shim et al. who revealed that serum ferritin was correlated with insulin resistance and abdominal obesity, both in males and females (7). Additionally, increased serum iron levels linked to an increased risk of MetS were reported in the Vietnamese population (33). Overweight individuals with high serum ferritin levels are at an increased

risk of diabetes than normal individuals (10). Tofano et al. revealed that MetS components such as hyperglycemia, hypertriglyceridemia, increased blood pressure level, and increased WC were positively correlated with hyperferritinemia, whereas HDL cholesterol was negatively correlated with hyperferritinemia (6). An increased level of serum ferritin was linked to an increased risk of MetS in Chinese males but not in females (34). A Korean study revealed the same result both in males and females. The pathophysiology underlying the association between ferritin and MetS remains unknown. A possible acceptable explanation is that the increased body iron level is linked to insulin resistance, which plays an important role in the occurrence of MetS (7). Iron is a transition metal that can become highly reactive by catalyzing the formation of free radicals associated with oxidative damage to the body. Increased oxidative stress is linked to intracellular signaling and gene expression abnormalities and can result in pathological conditions, such as insulin resistance (7).

Our study results revealed significant differences in ICAM-1 level between the two groups, with a higher mean ICAM-1 level in the MetS group than in the control group ($p = 0.004$). This indicates that ICAM-1 in the MetS group is expressed to a greater extent than that in the control group. ICAM-1 is influenced by genetic polymorphisms, as indicated in the studies by Kent et al. and Zhang et al. (35,36). ICAM-1 is an adhesion molecule that plays an important role in the initiation of inflammatory processes in the vascular endothelium. ICAM-1 is usually expressed at lower levels on the surface of arterial endothelial cells. ICAM-1 expression increases when the endothelium is stimulated due to physical or chemical injury. The interaction between adhesion molecules and leukocytes influences the onset of leukocyte migration into the arterial intima. ICAM-1 plays a role in immune cell recruitment during atherosclerotic plaque formation (35). Kent et al. revealed the correlation between ICAM-1 and MetS components such as obesity, insulin resistance, and HDL cholesterol (35). Lee et al. reported that the increase in MetS score resulted in a 1.26-fold increase in the levels of endothelial dysfunction markers (such as ICAM-1, but not vascular cell adhesion molecule-1) (8).

Our study was limited by the lack of data on the duration of MetS in the participants; moreover, the recruited participants were limited to one district, such that the results cannot be generalized to the Indonesian population. Further prospective multicenter studies are needed to validate our results.

CONCLUSION

Ferritin, PTX-3, and ICAM-1 level were significantly different between the MetS and control groups. Elevated ferritin, PTX-3, and ICAM-1 level are greatly linked to increased MetS risk in females aged 50–75 years, but the same is not true for HsCRP.

ACKNOWLEDGMENTS

The investigators wish to extend their heartfelt gratitude to the respondents of this study who agreed to participate and the *Puskesmas* of Mampang Prapatan District and the

Health Service Office of South Jakarta for their cooperation in the conduct of this study.

AUTHORS' CONTRIBUTIONS

Conceptualization: Pr. Data curation: Pr, RW, LTM, Me. Formal analysis: Pr, RW. Methodology: Pr, Me. Writing—original draft: Pr. Writing—review, and editing: RW, LTM, Me, RY.

STATEMENT OF INTERESTS

The authors declare that there is no conflict of interest.

FUNDING

Universitas Trisakti, Jakarta, Indonesia.

REFERENCES

- Ranasinghe P, Mathangasinghe Y, Jayawardena R, Hills AP, Misra A. Prevalence and trends of metabolic syndrome among adults in the Asia-Pacific region: A systematic review. *BMC Public Health* 2017;17(1):101. <https://doi.org/10.1186/s12889-017-4041-1>
- Sigit FS, Tahapary DL, Trompet S, Sartono E, van Dijk KW, Rosendaal FR, et al. The prevalence of metabolic syndrome and its association with body fat distribution in middle-aged individuals from Indonesia and the Netherlands: A cross-sectional analysis of two population-based studies. *Diabetol Metab Syndr* 2020;12(1):2. <https://doi.org/10.1186/s13098-019-0503-1>
- Kizilgul M, Uzunlulu M, Karancakaya M, Semerci A, Isbilen B, Isman FK. Pentraxin 3 levels as a marker of chronic inflammation in patients with metabolic syndrome. *J Endocrinol Metab* 2012;2(6):220-7. <https://doi.org/10.4021/jem142w>
- Altieri PI, Marcial J, Banchs HL, Escobales N, Crespo M. The metabolic syndrome in Hispanics—the role of inflammation. *Glob J Obes Diabetes Metab Syndr* 2015;2(1):12-7. <https://doi.org/10.17352/2455-8583.000009>
- Lee R, Ahn HR, Shin MH, Kim HN, Lee YH, Choi SW, et al. Association of plasma pentraxin-3 level with lipid levels and cardiovascular risk factors in people with no history of lipid-lowering medication: The Dong-gu study. *J Atheroscler Thromb* 2019;26(8):738-45. <https://doi.org/10.5551/jat.47167>
- Tofano RJ, Salzedas LM, Chagas EF, Detregiachi CR, Guiguer EL, Araujo AC, et al. Association of metabolic syndrome and hyperferritinemia in patients at cardiovascular risk. *Diabetes Metab Syndr Obes* 2020;13:3239-48. <https://doi.org/10.2147/DMSO.S271050>
- Shim YS, Kang MJ, Oh YJ, Baek JW, Yang S, Hwang IT. Association of serum ferritin with insulin resistance, abdominal obesity, and metabolic syndrome in Korean adolescent and adults: The Korean National Health and Nutrition Examination Survey, 2008 to 2011. *Medicine (Baltimore)* 2017;96(8):e6179. <https://doi.org/10.1097/md.00000000000006179>
- Lee HA, Choi EJ, Park B, Lee H, Hong YS, Kim HS, et al. The association between metabolic components and markers of inflammatory and endothelial dysfunction in adolescents, based on the Ewha Birth and Growth Cohort Study. *PLoS One* 2020;15(5):e0233469. <https://doi.org/10.1371/journal.pone.0233469>
- Takashi Y, Koga M, Matsuzawa Y, Saito J, Omura M, Nishikawa T. Circulating pentraxin 3 is positively associated with chronic hyperglycemia but negatively associated with plasma aldosterone concentration. *PLoS One* 2018;13(5):e0196526. <https://doi.org/10.1371/journal.pone.0196526>
- Zhou B, Liu S, Yuan G. Combined associations of serum ferritin and body size phenotypes with cardiovascular risk profiles: A Chinese population-based study. *Front Public Health* 2021;9:550011. <https://doi.org/10.3389/fpubh.2021.550011>
- Gündogan K, Bayram F, Capak M, Tanriverdi F, Karaman A, Ozturk A, et al. Prevalence of metabolic syndrome in the Mediterranean region of Turkey: Evaluation of hypertension, diabetes mellitus, obesity, and dyslipidemia. *Metab Syndr Relat Disord* 2009;7(5):427-34. <https://doi.org/10.1089/met.2008.0068>
- Mehndiratta N, Sharma S, Sharma RK, Grover S. A prospective study on the incidence of metabolic syndrome in premenopausal and postmenopausal women. *J Midlife Health* 2020;11(1):17-21. https://doi.org/10.4103/jmh.jmh_57_19
- Bao J, Wang L, Hu P, Liu J, Tu J, Wang J, et al. Burden of metabolic syndrome among a low-income population in China: A population-based cross-sectional study. *Diabetes Metab Syndr Obes* 2022;15:2713-23. <https://doi.org/10.2147/DMSO.S377490>
- Stephens CR, Easton JF, Robles-Cabrera A, Fossion R, de la Cruz L, Martínez-Tapia R, et al. The impact of education and age on metabolic disorders. *Front Public Health* 2020;8:180. <https://doi.org/10.3389/fpubh.2020.00180>
- Farhadnejad H, Parastouei K, Rostami H, Mirmiran P, Azizi F. Dietary and lifestyle inflammatory scores are associated with increased risk of metabolic syndrome in Iranian adults. *Diabetol Metab Syndr* 2021;13(1):30. <https://doi.org/10.1186/s13098-021-00648-1>
- Hong GB, Gao PC, Chen YY, Xia Y, Ke XS, Shao XF, et al. High-sensitivity C-reactive protein leads to increased incident metabolic syndrome in women but not in men: A five-year follow-up study in a Chinese population. *Diabetes Metab Syndr Obes* 2020;13:581-90. <https://doi.org/10.2147/dms0.s241774>
- Carrero JJ, Franko MA, Obergfell A, Gabrielsen A, Jernberg T. HsCRP level and the risk of death or recurrent cardiovascular events in patients with myocardial infarction: A healthcare-based study. *J Am Heart Assoc* 2019;8(11):e012638. <https://doi.org/10.1161/JAHA.119.012638>
- Yilmaz SK, Eskici G, Mertoğlu C, Ayaz A. Adipokines, inflammation, oxidative stress: Critical components in obese women with metabolic syndrome. *Prog Nutr* 2021;23:1-12. <https://doi.org/10.23751/pn.v23i1.9072>
- Kurylowicz A. Estrogens in adipose tissue physiology and obesity-related dysfunction. *Biomedicines* 2023;11(3):690. <https://doi.org/10.3390/biomedicines11030690>
- Somani YB, Pawelczyk JA, De Souza MJ, Kris-Etherton PM, Proctor DN. Aging women and their endothelium: Probing the relative role of estrogen on vasodilator function. *Am J Physiol Heart Circ Physiol* 2019;317(2):H395-404. <https://doi.org/10.1152/ajpheart.00430.2018>
- Grundey SM, Cleeman JI, Daniels SR, Donato KA, Eckel RH, Franklin BA, et al. Diagnosis and management of the metabolic syndrome: An American Heart Association/National Heart, Lung, and Blood Institute scientific statement. *Circulation* 2005;112(17):2735-52. <https://doi.org/10.1161/circulationaha.105.169404>
- Silva IV, De Figueiredo RC, Rios DR. Effect of different classes of antihypertensive drugs on endothelial function and inflammation. *Int J Mol Sci* 2019;20(14):3458. <https://doi.org/10.3390/ijms20143458>
- Proute MC, Kothur N, Georgiou P, Serhiyenia T, Shi W, Kerolos ME, et al. The effect of statin therapy on inflammatory biomarkers: A systematic review. *Cureus* 2021;13(9):e18273. <https://doi.org/10.7759/cureus.18273>
- Bulcão C, Ribeiro-Filho FF, Sañudo A, Ferreira SG. Effects of simvastatin and metformin on inflammation and insulin resistance in individuals with mild metabolic syndrome. *Am J Cardiovasc Drugs* 2007;7:219-24. <https://doi.org/10.2165/00129784-200707030-00007>
- Liu S, Qu X, Liu F, Wang C. Pentraxin 3 as a prognostic biomarker in patients with systemic inflammation or infection. *Mediators Inflamm* 2014;2014:421429. <https://doi.org/10.1155/2014/421429>
- Karakas MF, Buyukkaya E, Kurt M, Motor S, Akcay AB, Karakas E, et al. Serum pentraxin-3 levels are associated with the severity of metabolic syndrome. *Med Princ Pract* 2013;22(3):274-9. <https://doi.org/10.1159/000343904>
- Kardas F, Akin L, Kurtoglu S, Kendirci M, Kardas Z. Plasma pentraxin 3 as a biomarker of metabolic syndrome. *Indian J Pediatr* 2015;82(1):35-8. <https://doi.org/10.1007/s12098-014-1542-0>
- Yamasaki K, Kurimura M, Kasai T, Sagara M, Kodama T, Inoue K. Determination of physiological plasma pentraxin 3 (PTX3) levels in healthy populations. *Clin Chem Lab Med* 2009;47(4):471-7. <https://doi.org/10.1515/CCLM.2009.110>
- Zanetti M, Bosutti A, Ferreira C, Vinci P, Biolo G, Fonda M, et al. Circulating pentraxin 3 levels are higher in metabolic syndrome with subclinical atherosclerosis: Evidence for association with atherogenic lipid profile. *Clin Exp Med* 2009;9(3):243-8. <https://doi.org/10.1007/s10238-009-0039-z>
- Ogawa T, Kawano Y, Imamura T, Kawakita K, Sagara M, Matsuo T, et al. Reciprocal contribution of pentraxin 3 and C-reactive protein to obesity and metabolic syndrome. *Obesity (Silver Spring)* 2010;18(9):1871-4. <https://doi.org/10.1038/oby.2009.507>
- Derisoglu P, Elmas B. Pentraxin 3 as a marker for cardiovascular disease risk in overweight and obese children. *Acta Cardiol Sin* 2021;37(2):177-83. [https://doi.org/10.6515/acs.202103_37\(2\).20201006a](https://doi.org/10.6515/acs.202103_37(2).20201006a)
- Chand MR, Agarwal N, Nishanth D, Jhuma S. Association of serum ferritin levels with

- metabolic syndrome in India: A cross-sectional study. *Maedica (Bucur)* 2021;16:48-53.
<https://doi.org/10.26574/maedica.2020.16.1.48>
33. Tran TN, Tran HD, Tran-Huu TT, Tran DM, Tran QN. A cross-sectional study of serum ferritin levels in Vietnamese adults with metabolic syndrome. *Diabetes Metab Syndr Obes* 2022;15:1517-23.
<https://doi.org/10.2147/dmso.S360689>
 34. Han LL, Wang YX, Li J, Zhan XL, Bian C, Wang H, et al. Gender differences in associations of serum ferritin and diabetes, metabolic syndrome, and obesity in the China Health and Nutrition survey. *Mol Nutr Food Res* 2014;58(11):2189-95.
<https://doi.org/10.1002/mnfr.201400088>
 35. Kent JW Jr., Comuzzie AG, Mahaney MC, Almasy L, Rainwater DL, VandeBerg JL, et al. Intercellular adhesion molecule-1 concentration is genetically correlated with insulin resistance, obesity, and HDL concentration in Mexican Americans. *Diabetes* 2004;53(10):2691-5.
<https://doi.org/10.2337/diabetes.53.10.2691>
 36. Zhang X, Seman NA, Falhammar H, Brismar K, Gu HF. Genetic and biological effects of ICAM-1 E469K polymorphism in diabetic kidney disease. *J Diabetes Res* 2020;2020:8305460.
<https://doi.org/10.1155/2020/8305460>

Inflammatory and endothelial dysfunction markers in females with metabolic syndrome aged 50–75 years

by Pusparini Pusparini

Submission date: 16-Jun-2023 09:46AM (UTC+0700)

Submission ID: 2117020656

File name: 2003-Article_Text-12196-1-10-20230614-1.pdf (253.95K)

Word count: 6589

Character count: 34339



RESEARCH ARTICLE

Open Access

57

Inflammatory and endothelial dysfunction markers in females with metabolic syndrome aged 50–75 years

Pusparini^{1*}, Raditya Wratsangka², Lie Tanu Merijanti³, Meiyanti⁴, Rismawati Yaswir⁵

¹Department of Clinical Pathology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia, ²Department of Obstetrics Gynecology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia, ³Department of Occupational Medicine, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia, ⁴Department of Pharmacology and Pharmacy, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia, ⁵Department of Clinical Pathology, Faculty of Medicine, Andalas University, Padang, Indonesia

ABSTRACT

Introduction: The prevalence of metabolic syndrome (MetS) in Indonesia is 39% and it is higher in females than in males (46% vs. 28%). Features of the syndrome comprise low-grade chronic inflammation and endothelial dysfunction from cardiometabolic abnormalities. High-sensitive C-reactive protein (HsCRP) is a biomarker of cardiovascular disease, whereas ferritin and pentraxin-3 (PTX-3) are markers of chronic inflammation. One of the markers of endothelial dysfunction is an intercellular cell adhesion molecule-1 (ICAM-1). Studies on MetS in women aged 50–70 years and using several markers of inflammation and endothelial dysfunction are rare. The study aimed to determine differences in HsCRP, ferritin, PTX-3, and ICAM-1 level between females with and without MetS aged 50–75 years.

Methods: For this cross-sectional study, 160 females aged 50–75 years from South Jakarta were recruited, divided into MetS and non-MetS (control) groups, 80 participants each. Participants that met the inclusion criteria were assessed for vital signs and metabolic parameters (e.g., lipid profile, fasting blood glucose level, and waist circumference [WC]). The groups were compared using an independent t-test for inflammatory (i.e., HsCRP, PTX-3, and ferritin) and endothelial function (ICAM-1) ($p < 0.05$) biomarkers.

Results: The mean age of the MetS and control groups was 59.6 ± 5.6 and 59.2 ± 5.5 years. MetS criteria included WC of ≥ 80 cm (95.2%), increased blood pressure level (90.6%); and high-density lipoprotein cholesterol of < 50 mg/dL (76.2%). Significant differences were found between the MetS and control groups for ferritin ($p = 0.027$), PTX-3 ($p = 0.045$), and ICAM-1 ($p = 0.004$) but not for HsCRP ($p = 0.136$).

Conclusion: Elevated ferritin, PTX-3 level, and ICAM-1 level are significantly associated with an increased Mets risk in females aged 50–75 years; however, the same is not true for HsCRP.

Keywords: Ferritin; high-sensitive C-reactive protein; intercellular cell adhesion molecule-1; inflammation; metabolic syndrome; pentraxin-3

INTRODUCTION

Currently, metabolic syndrome (MetS) is developing into a global epidemic, with approximately 12–37% of Asians and 12–26% of Europeans suffering from MetS (1). The incidence of MetS in Indonesia is higher in females than in males. The prevalence of MetS in adults aged 45–65 years is 39%, namely 28% in males and 46% in females (2). MetS is a group of symptoms comprising abdominal obesity, hypertension, dyslipidemia, hyperglycemia, and prothrombotic and proinflammatory conditions (3).

The higher prevalence of abdominal obesity in Indonesian females, accounting for 57.5% versus 16.8% in males,

explains the higher MetS prevalence in Indonesian females than in males. Individuals with MetS are at an increased risk of morbidity and mortality from cardiovascular disease. MetS is a risk factor for the occurrence of type 2 diabetes mellitus (T2DM), cardiovascular disease, and stroke (2). The pathophysiology of cardiovascular disease in MetS probably involves risk factors such as insulin resistance, chronic low-grade systemic inflammation, genetics, dietary status, and individual habits (3,4).

Abdominal fat is an active endocrine organ that generates cytokines that affect other body tissues (4). Abdominal fat releases several inflammatory markers, such as C-reactive protein (CRP), and acute-phase proteins, such as pentraxin-3 (PTX-3) (3). The level of plasma PTX-3 was inversely related to MetS in the Korean population (5). Another acute-phase protein is ferritin (6,7). Ferritin is not only a marker of iron reserves in the body but may also be a marker of inflammation (6). Abnormal cardiometabolic

Corresponding author: Pusparini, Department of Clinical Pathology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia.
E-mail: pusparini@trisakti.ac.id

Submitted: 27 October 2022/Accepted: 12 May 2023

DOI: <https://doi.org/10.17532/jhsci.2023.2003>



UNIVERSITY OF SARAJEVO
FACULTY OF HEALTH STUDIES

© 2023 Pusparini, et al.; licensee University of Sarajevo - Faculty of Health Studies. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

components contribute to the occurrence of endothelial dysfunction followed by blood vessel inflammation, ⁴³ultimately in atherosclerosis and cardiovascular disease (8). Intercellular cell adhesion molecule-1 (ICAM-1) is an endothelial dysfunction marker (8,9). Many studies have reported the relationship between MetS and inflammatory markers, but MetS occurrence varies by population, ethnicity, and sex (4,8,10). Studies on MetS conducted in females aged 50–75 years and using several inflammation (high-sensitive CRP [HsCRP], PTX-3, and ferritin) and endothelial dysfunction (ICAM-1) markers are rare (4,6,8–10). This study aimed to determine differences in the levels of inflammatory and endothelial dysfunction biomarkers between females with and without MetS aged 50–75 years.

METHODS

This cross-sectional study was conducted from January to June 2021 in the catchment area of Puskesmas in Mampang District, South Jakarta. The investigators enrolled 80 females aged 50–75 years with MetS and 80 non-MetS females (controls). Inclusion criteria for the MetS group were the presence of three ²¹MetS criteria, i.e., waist circumference (WC) of ≥ 80 cm, systolic blood pressure (SBP) level of ≥ 130 mmHg or diastolic blood pressure (DBP) level of ≥ 85 mmHg or taking antihypertensive ¹⁴medications; fasting blood glucose (FBG) level of ≥ 100 mg/dL; high-density lipoprotein (HDL) cholesterol level of < 50 mg/dL; triglyceride level of ≥ 150 mg/dL; no active infection in 2 weeks before the study; not smoking, not consuming alcohol, and providing written informed consent. Exclusion criteria for the MetS and control groups in this study comprised the administration of corticosteroids, oral anticoagulants, and hormone replacement therapy; history of malignancies ²⁴abnormal renal function. The MetS criteria employed in this study followed the revised National Cholesterol Education Programme's Adult Treatment Panel III criteria for Asians (1).

The ⁶¹participants were interviewed by five individuals (having at least senior high school or equivalent education) with knowledge of the aims of the study and trained in the filling of the questionnaire (including demographic characteristics, concomitant disease, smoking status, alcohol drinking status, concomitant medication usage, and anthropometrics) (Table 1). The filling-in of the questionnaire was completed in 2 weeks. Following physical examination (vital signs); anthropometry; and laboratory bloodwork for lipid profile, FBG, and creatinine, participants were divided into MetS and control groups. The ¹¹investigators determined the blood pressure level, body weight, height, WC, and hip circumference of participants through physical examination. Venous blood was drawn (following a 12-h fast) using a 10-mL plain vacutainer for FBG and a complete lipid profile. An aliquot of the blood sample was used to obtain serum, which was stored at -70°C for determination of PTX-3, ICAM-1, HsCRP, and ferritin levels in participants. Since differences in intraday repeatability of laboratory procedures may cause spurious differences in the results between the MetS and control groups, the concentrations of PTX-3, ICAM-1, HsCRP, and ferritin were simultaneously determined, to ensure that differences in PTX-3, ICAM-1, HsCRP, and ferritin levels were not

due to laboratory errors but were true differences. The biomarkers were ¹⁹assayed using the following techniques: PTX-3 through enzyme-linked immunosorbent assay (ELISA) using Quantikine® ELISA Human PTX-3/TSG-14 reagents (R&D Systems, Inc., Minneapolis, USA) and Biorad model 680 microplate reader; ICAM-1 via ELISA using Quantikine® ELISA Human ICAM-1 reagent (R&D Systems, Inc., Minneapolis, USA) and Biorad model 680 microplate reader; HsCRP using immunoturbidimetry, Architect 6K26-30 reagents, and Architect c-8000 analyzer; and ferritin via chemiluminescent microparticle immunoassay using Architect 7K59 reagents and Architect i-2000 analyzer. This study was approved by the Ethics Committee, Faculty of Medicine, Universitas Tr ⁵⁹ti (approval no. 165/KER/FK/XII/2020). Independent t-test was used to ²⁵determine differences in biomarker levels between the MetS and control groups, and $p < 0.05$ were considered to indicate statistical significance.

RESULTS

The characteristics of participants with MetS and controls are summarized in Table 2. The SBP, DBP, WC, and hip circumference of participants were considerably different between the two groups (p -values of 0.00 ¹³1.016, 0.009, and 0.006, respectively). The biochemical characteristics of participants are listed in Table 3. The MetS group showed significantly increased triglyceride and FBG and reduced HDL cholesterol levels compared with ³⁴control group. PTX-3, ferritin, and ICAM-1 ⁶level (p -values of 0.045, 0.027, and 0.004, respectively) were significantly higher in the MetS than in the control groups, whereas HsCRP levels did not indicate any significant differences. In the MetS group, the five MetS parameters presented in the order of their frequency ³³were: WC of ≥ 80 cm, increased blood pressure level, HDL cholesterol level of < 50 mg/dL, increased FBG level, and triglyceride level of ≥ 150 mg/dL.

DISCUSSION

In the present study, out of the five MetS-contributing elements, the most frequently encountered in the participants with MetS were: WC of ≥ 80 cm (97.5%) and increased blood pressure (91.3%), followed by ⁴⁸L cholesterol concentration of < 50 mg/dL (80%). The study by Sigit et al. (2) showed that the predominant MetS components in the Indonesian population were hypertension (61.3%; 56.2% in males vs. 64.6% in females) and hyperglycemia (51%; 51% in males vs. 50.9% in females), and hypertension (61.7%; 70% in males vs. 55.2% in females) ²³abdominal obesity (40%; 34.6% in males vs. 44.2% in females) in the Dutch population. The prevalence of abdominal obesity was approximately 57.5% in Indonesian females and 16.8% in Indonesian males (2). The aforementioned study highlighted that MetS in Indonesia was more common in females (46.2%) than in males (28%) (2). The differences between the study by Sigit et al. (2) and the current study may have occurred ⁵⁶because participants in the current study were all females with a mean age of 59.6 ± 5.6 years in the MetS group and 59.2 ± 5.5 years in the control group, whereas Sigit et al. (2) included male and female participants, with a mean

TABLE 1. Questionnaire used during the enrollment process (English translation)

General Information	
Name:	
Address:	
Telephone/Cellphone number:	
Interviewer:	
Date of interview:	
Circle option number according to your answer to items No. 1–10 (Please include photocopy of your National Identity Card)	
1. Age (years, months):	
2. Date of birth (DD/MM/YY):	
3. Educational background	
1. None 2) Elementary School 3) Junior High School 4) Senior High School 5) Academy/Bachelor 6) Did not complete elementary school	
4. Current employment status	
1. Unemployed 2) Trader 3) Entrepreneur 4) Other	
5. Are you in menopause?	
1. Yes 2) No	
6. If in menopause, give duration of menopause.....year (s);.....month (s)	
7. Did you have any infection in the past 2 weeks?	
1. Yes 2) No	
8. Do you have any of the following diseases or a past history of these diseases?	
a. Hypertension	1) Yes 2) No
b. Diabetes Mellitus	1) Yes 2) No
c. Kidney disease	1) Yes 2) No
d. Stroke	1) Yes 2) No
e. Cancer	1) Yes 2) No
f. Other	1) Yes 2) No
If yes, state the diagnosis	
9. Do you take any of the following drugs?	
a. Corticosteroids	1) Yes 2) No
b. Hormonal replacement therapy	1) Yes 2) No
c. Antihypertensive drugs	1) Yes 2) No
d. Antidiabetic drugs	1) Yes 2) No
e. Antihyperlipidemic drugs	1) Yes 2) No
f. Oral anticoagulants	1) Yes 2) No
10. Do you smoke?	1) Yes 2) No
If yes, how many cigarettes per day	
11. Do you consume alcoholic drinks?	1) Yes 2) No
If yes, how many glasses per day	

age of 52.2 ± 5.8 years in the Indonesian population (2). The prevalence of MetS increased with age, as reported in a Turkish study: 11.9% (age 20–29 years), 23.4% (age 30–39 years), 26.9% (age 40–49 years), 40.4% (age 50–59 years), 45.3% (age 60–69 years), and 26.3% (age >70 years) (11). Mehndiratta et al. (12) conducted a study on 200 women aged 45–60 years receiving ambulatory treatment in an obstetrics and gynecology department at Gurnagar, Punjab, India. The study results indicated that 29% of women were found to have newly onset MetS, with an incidence of 16% in the premenopausal group and 2% in the postmenopausal group (12). MetS incidence is higher in postmenopausal women than in premenopausal women, because following menopause, the estrogen-to-testosterone ratio decreases, dilutes the vasorelaxant effects of estrogen on the vessel wall and promotes the production of vasoconstrictor factors such as endothelin which lead to metabolic culminating into MetS (12).

TABLE 2. Characteristics of study participants

Parameter	X $\pm$ SD/n (%) MetS n=80	X $\pm$ SD/n (%) Control n=80	p-value
Age (years)	59.6 $\pm$ 5.6	59.2 $\pm$ 5.5	0.256
Weight (kg)	60.4 $\pm$ 10.2	55.3 $\pm$ 8.9	0.173
Height (cm)	152.5 $\pm$ 5.7	152.3 $\pm$ 5.3	0.532
BMI (kg/m ²)	25.9 $\pm$ 4.6	21.5 $\pm$ 3.5	0.076
Pulse rate	84.9 $\pm$ 14.2	80.6 $\pm$ 11.9	0.060
Blood pressure level			
Systolic (mmHg)	150.5 $\pm$ 23.0	127.6 $\pm$ 23.7	0.000*
Diastolic (mmHg)	88.4 $\pm$ 12.1	79.5 $\pm$ 10.7	0.016*
Waist circumference (cm)	91.7 $\pm$ 7.2	77.8 $\pm$ 8.0	0.009*
Hip circumference (cm)	99.2 $\pm$ 7.8	90.4 $\pm$ 6.5	0.006*
Education			
Elementary school	3 (3.8)	2 (2.5)	
Junior high school	30 (37.5)	27 (33.7)	
Senior high school	21 (26.3)	20 (25)	0.785
Academy/Bachelor	19 (23.7)	26 (32.5)	
Did not complete elementary school	5 (6.2)	3 (3.8)	
Employment			
Unemployed	2 (2.5)	2 (2.5)	
Trader	58 (72.4)	57 (71.3)	
Entrepreneur	5 (6.3)	6 (7.5)	0.482
Other	5 (6.3)	3 (3.7)	
Menopause			
Yes	12 (15)	14 (17.5)	
No	72 (90)	73 (91.3)	0.348
Duration of menopause (years)	8 (10)	7 (8.7)	
History of diabetes mellitus	10.11 $\pm$ 2.3	10.7 $\pm$ 2.1	0.753
History of hypertension	16 (20)	0 (0)	
Medications	47 (58.8)	0 (0)	
Antihypertensive	32 (40)	0 (0)	
Antidiabetic	9 (11.3)	0 (0)	
Antihyperlipidemic	12 (15)	0 (0)	

*p<0.05=significant difference (independent t-test); Chi-square test.

BMI: Body mass index

TABLE 3. Biochemical and inflammatory markers

Parameter	X $\pm$ SD MetS n=80	X $\pm$ SD Non-MetS n=80	p-value
Biochemical			
Total cholesterol (mg/dL)	215.8 $\pm$ 15.9	208.4 $\pm$ 13.7	0.286
Triglycerides (mg/dL)	145.8 $\pm$ 10.5	94.3 $\pm$ 12.8	0.000*
HDL cholesterol (mg/dL)	48.8 $\pm$ 9.8	58.9 $\pm$ 10.5	0.000*
LDL cholesterol (mg/dL)	139.9 $\pm$ 18.7	132.2 $\pm$ 10.7	0.198
55 mg blood glucose (mg/dL)	124.7 $\pm$ 13.5	88.6 $\pm$ 9.3	0.000*
Creatinine (mg/dL)	0.77 $\pm$ 0.2	0.79 $\pm$ 0.3	0.686
eLFG (mL/min)	85.5 $\pm$ 17.4	85.7 $\pm$ 17.9	0.931
Inflammatory			
Pentraxin-3 (ng/dL)	0.81 $\pm$ 0.20	0.72 $\pm$ 0.19	0.045*
HsCRP (mg/L)	3.1 $\pm$ 0.6	2.6 $\pm$ 0.3	0.136
Ferritin (μ g/L)	165.9 $\pm$ 14.7	125.7 $\pm$ 8.7	0.027*
60 ICAM-1 (ng/mL)	356.6 $\pm$ 9.9	316.8 $\pm$ 9.5	0.004*

*p<0.05 significant difference (independent t-test). HDL: High-density lipoprotein, LDL: low-density lipoprotein, HsCRP: High-sensitive C-reactive protein, ICAM-1: Intercellular cell adhesion molecule-1

This study demonstrated no significant differences were found in educational level (Table 2; $p = 0.785$) and

employment status (Table 2; $p = 0.482$) between the MetS and control group. Most of the subjects in the two groups had an educational level of senior high school and below (MetS 91.3% vs. non-MetS 93.7%) and were not anymore employed (MetS 72.4% vs. non-MetS 71.3%). In this study, no distinction was made between senior high school and tertiary education. Research in China on 3175 subjects aged ≥ 45 years showed that MetS prevalence was quite high in the rural population with low income and low education (13). Stephens et al. reported that a higher education level correlates to significantly better metabolic health when compared to a lower education level and carries significantly less risk for WC, SBP, glucose and glycosylated hemoglobin, triglycerides, HDL cholesterol, and MetS (all $p < 0.05$), except for DBP, basal insulin, uric acid, low-density lipoprotein cholesterol, and total cholesterol (14).

Low-grade chronic inflammation is identified as the leading factor for the occurrence of MetS and subsequent clinical symptoms (15,16). Among the four biomarkers assessed in this study, ferritin, ICAM-1, and PTX-3, but not HsCRP, showed significant differences between the MetS and control groups (Table 3). The threshold values of HsCRP (<1 , $1-3$, and $3-10$) indicate low, moderate, and high risk of cardiovascular disease, respectively (17). The MetS group had higher mean HsCRP levels than the control group based on these criteria. The mean HsCRP level in the MetS group was 3.1 ± 0.4 mg/L, indicating a high risk, whereas that in the control group was 2.6 ± 0.3 mg/L, indicating a moderate risk, although no significant differences were observed between the two groups. Carrero et al. (17) studied myocardial infarction (MI) survivors (17,464 participants aged >18 years) undergoing HsCRP testing for >30 days after MI and revealed that most patients with MI had increased HsCRP levels. HsCRP may be used as a prognostic factor for determining the occurrence of cardiovascular disease (17). Yilmaz et al. (18) analyzed 100 females aged 20–45 years with obesity and revealed that HsCRP, tumor necrosis factor (TNF)- α , and malondialdehyde may be used as diagnostic markers of MetS in females with obesity. Furthermore, WC is correlated with HsCRP ($r = 0.315$; $p = 0.003$) (18). The current study revealed correlation between WC and HsCRP, both in the MetS ($r = 0.353$, $p = 0.002$; data not shown) and control ($r = 0.560$, $p = 0.003$) groups. A WC of >80 cm in females indicates the occurrence of visceral fat accumulation. Visceral fat is an endocrine organ that can secrete pro-inflammatory substances, such as leptin, adiponectin, resistin, TNF- α , and interleukin-6, which play a significant role in the occurrence of MetS (18). The study of Hong et al. in China revealed that after a follow-up period of 5 years, HsCRP was linked to MetS, but only in females. The reason why females are at higher risk for MetS than males remains unclear, but physiologically, females may have more visceral fat than males, which is associated with MetS risk (16). Four estrogens have been identified in humans: estrone, estradiol, estriol, and estretrol. Estradiol is the predominant estrogen in women during reproduction. Post-menopausal estrogen production in the ovaries is reduced therefore circulating estrogen concentrations are also drastically reduced. After menopause, estradiol is replaced by estrone, synthesized in adipose tissue from adrenal dehydroepiandrosterone (19). The current study

revealed that the HsCRP level did not significantly differ between the MetS and control groups, probably because the body mass index (BMI) of both groups was not significantly different ($p = 0.076$). A study in China revealed that BMI was correlated with HsCRP (10). In addition, 90% of our participants were in menopause, both in the MetS and control groups. Estrogen has pleiotropic effects and plays a crucial role in preserving endothelial functions (20). Estrogen increases the bioavailability of nitric oxide by promoting its synthesis and exerts direct and indirect antioxidant and anti-inflammatory effects. The decreased estrogen level in menopause causes endothelial dysfunction followed by vascular inflammation (20). Furthermore, several drugs used in the treatment of metabolic risk factors, such as statins, nicotinic acid, fibrates, angiotensin-converting enzyme inhibitors, and thiazolidinediones, have been reported to decrease CRP levels (21). Among the MetS group in the present study, 40% used anti-hypertensive drugs, 11.3% used anti-diabetics, and 15% used anti-hyperlipidemic drugs (Table 2). Silva et al. (22) reported that different anti-hypertensive drugs have different effects on inflammatory markers and markers of endothelial function. Although angiotensin-converting enzyme inhibitors and angiotensin receptor blockers do beneficially affect endothelial function and inflammation, first and second-generation β -blockers have no such effect (22). A systematic review on the effect of statin therapy on inflammatory markers showed that statin therapy has lipid-lowering and anti-inflammatory effects (23). Forty-one participants with a BMI of $25-39.9$ kg/m² and impaired glucose tolerance test were randomized to take simvastatin or metformin for 16 weeks. The study demonstrated a significant reduction in CRP and interleukin-6 concentration at the completion of the study (24).

PTX-3 is both an inflammatory marker and an acute-phase protein (25-27). Obesity and MetS are linked to chronic inflammation (27). This study revealed that PTX-3 levels were significantly higher in the MetS group than in the control group ($p = 0.045$) (Table 3). This is consistent with the study by Karakas et al., indicating that the MetS group has higher PTX-3 levels than the control group (26). Moreover, Karakas et al. found that the PTX-3 levels in the group with 5 MetS components (0.90 ± 0.06 ng/dL) were higher than in the groups with 4 MetS components (0.63 ± 0.12 ng/dL) and 3 MetS components (0.58 ± 0.11 ng/dL) (26). In our study, the mean PTX-3 levels in the MetS and control groups were 0.81 ± 0.20 and 0.72 ± 0.19 ng/dL, respectively. The mean PTX-3 level in the MetS group was higher than that reported by Karakas et al. (0.58 ± 0.11 ng/dL) with the same number of components of MetS criteria (26). The participants in our study were older than those in the study by Karakas et al. (approximately 49 years of age). Accordingly, our study participants might have had MetS for a longer period than those in the study of Karakas et al. Nevertheless, our study is in line with the study of Karakas et al. that PTX-3 level is correlated with MetS severity. Kizilgul et al. (3) stated that a number of study results related to PTX-3 level in MetS were contradictory. For instance, a study conducted on 1749 healthy Japanese participants (818 men and 931 women) in the age range of 37–87 years indicated that PTX-3 levels were lower in participants with MetS than in the non-MetS group and

that the PTX-3 levels were negatively correlated with triglyceride levels and HDL (28). Zanetti et al. (29) reported that plasma PTX-3 levels were higher in the MetS group with subclinical atherosclerosis than in the control group. Ogawa et al. (30) reported low PTX-3 and high CRP levels in males with obesity and/or MetS compared to the control group. Ogawa et al. (30) study indicate that the reduction in plasma PTX-3 may accelerate chronic inflammation and atherosclerosis in the obese and/or MetS population because of the tissue protective and antiprotective roles of PTX-3 (30). Kizilgul et al. (3) compared PTX-3 levels in patients with MetS, in patients with rheumatoid arthritis, and with healthy controls. Their results do not demonstrate an increase in the levels of PTX-3 as chronic inflammatory markers in patients with MetS (3). The symptoms of MetS are specific for each age group and population and therefore each group or population should evolve its own strategy for MetS prevention (2,25).

PTX-3, a marker of cardiovascular risk, is produced by various cells, such as adipocytes, macrophages, dendritic cells, fibroblasts, and vascular endothelial cells, and is released in response to inflammatory stimuli (26,27). The current study revealed that PTX-3 was negatively correlated with HDL cholesterol ($r = -0.253$ and $p = 0.028$) in the MetS group, whereas no significant correlation was noted in the control group (data not shown). This indicates that HDL cholesterol plays a role in inhibiting PTX-3. This is consistent with the study of Kardas et al., which showed that children with HDL cholesterol levels of <40 mg/dL have higher PTX-3 levels than those with normal HDL cholesterol levels ($p < 0.05$) (27). A study on overweight children and children with obesity indicated that the mean PTX-3 level was significantly higher in the overweight (10.23 ± 4.42 ng/mL) and obese (11.20 ± 4.12 ng/mL) groups compared to that in the control group (7.93 ± 4.35 ng/mL). In addition, PTX-3 level was linked to intima and epicardial fat thickness in overweight children. PTX-3 may be an early marker of cardiovascular risk in overweight children (31).

Ferritin is an essential protein regulating iron homeostasis in the body. Serum ferritin is used as a biomarker to determine the status of iron reserves in the body. Excessive iron accumulation may lead to different pathological conditions, such as hemochromatosis and hemosiderosis (7). In addition, serum ferritin is an acute-phase protein associated with insulin resistance. Iron accumulation interferes with pancreatic insulin synthesis and secretion and affects hepatic insulin-extracting capacity, resulting in peripheral hyperinsulinemia and impaired insulin secretion (32). Our study results revealed significant differences in serum ferritin levels between the MetS and control groups ($p = 0.027$) (table 3). The mean ferritin level in the MetS group was significantly higher than that in the control group. Cui et al. revealed comparable results; the serum ferritin levels were higher in the MetS group than in the control group ($p < 0.0001$) (42). Our study findings are consistent with those of Shim et al. who revealed that serum ferritin was correlated with insulin resistance and abdominal obesity, both in males and females (7). Additionally, increased serum iron levels linked to an increased risk of MetS were reported in the Vietnamese population (33). Overweight individuals with high serum ferritin levels are at an increased

risk of diabetes than normal individuals (10). Tofano et al. revealed that MetS components such as hyperglycemia, hypertriglyceridemia, increased blood pressure level, and increased WC were positively correlated with hyperferritinemia, whereas HDL cholesterol was negatively correlated with hyperferritinemia (6). An increased level of serum ferritin was linked to an increased risk of MetS in Chinese males but not in females (34). A Korean study revealed the same result both in males and females. The pathophysiology underlying the association between ferritin and MetS remains unknown. A possible acceptable explanation is that the increased body iron level is linked to insulin resistance, which plays an important role in the occurrence of MetS (7). Iron is a transition metal that can become highly reactive by catalyzing the formation of free radicals associated with oxidative damage to the body. Increased oxidative stress is linked to intracellular signaling and gene expression abnormalities and can result in pathological conditions, such as insulin resistance (7).

Our study results revealed significant differences in ICAM-1 level between the two groups, with a higher mean ICAM-1 level in the MetS group than in the control group ($p = 0.004$). This indicates that ICAM-1 in the MetS group is expressed to a greater extent than that in the control group. ICAM-1 is influenced by genetic polymorphisms, as indicated in the studies by Kent et al. and Zhang et al. (35,36). ICAM-1 is an adhesion molecule that plays an important role in the initiation of inflammatory processes in the vascular endothelium. ICAM-1 is usually expressed at lower levels on the surface of arterial endothelial cells. ICAM-1 expression increases when the endothelium is stimulated due to physical or chemical injury. The interaction between adhesion molecules and leukocytes influences the onset of leukocyte migration into the arterial intima. ICAM-1 plays a role in immune cell recruitment during atherosclerotic plaque formation (35). Kent et al. revealed the correlation between ICAM-1 and MetS components such as obesity, insulin resistance, and HDL cholesterol (35). Lee et al. reported that the increase in MetS score resulted in a 1.26-fold increase in the levels of endothelial dysfunction markers (such as ICAM-1, but not vascular cell adhesion molecule-1) (8).

Our study was limited by the lack of data on the duration of MetS in the participants; moreover, the recruited participants were limited to one district, such that the results cannot be generalized to the Indonesian population. Further prospective multicenter studies are needed to validate our results.

CONCLUSION

Ferritin, PTX-3, and ICAM-1 level were significantly different between the MetS and control groups. Elevated ferritin, PTX-3, and ICAM-1 level are greatly linked to increased MetS risk in females aged 50–75 years, but the same is not true for HsCRP.

ACKNOWLEDGMENTS

The investigators wish to extend their heartfelt gratitude to the respondents of this study who agreed to participate and the Puskesmas of Mampang Prapatan District and the

Health Service Office of South Jakarta for their cooperation in the conduct of this study.

AUTHORS' CONTRIBUTIONS

Conceptualization: Pr. Data curation: Pr, RW, LTM, Me. Formal analysis: Pr, RW. Methodology: Pr, Me. Writing—original draft: Pr. Writing—review, and editing: RW, LTM, Me, RY.

STATEMENT OF INTERESTS

The authors declare that there is no conflict of interest.

FUNDING

Universitas Trisakti, Jakarta, Indonesia.

REFERENCES

- Ranasinghe P, Mathangasinghe Y, Jayawardena R, Hills AP, Misra A. Prevalence and trends of metabolic syndrome among adults in the Asia-Pacific region: A systematic review. *BMC Public Health* 2017;17(1):101. <https://doi.org/10.1186/s12889-017-4041-1>
- Sigst F, Tahapary DL, Trompet S, Sartono E, van Dijk KW, Rosendaal FR, et al. The prevalence of metabolic syndrome and its association with body fat distribution in middle-aged individuals from Indonesia and the Netherlands: A cross-sectional analysis of two population-based studies. *Diabetol Metab Syndr* 2020;12(1):2. <https://doi.org/10.1186/s13098-019-0503-1>
- Kizilgul M, Uzunlulu M, Karancakaya M, Semerci A, Isbilen B, Isman FK. Pentraxin 3 levels as a marker of chronic inflammation in patients with metabolic syndrome. *J Endocrinol Metab* 2012;2(6):220-7. <https://doi.org/10.4021/jem142w>
- Altieri PI, Marcial J, Banchs HL, Escobales N, Crespo M. The metabolic syndrome in Hispanics-the role of inflammation. *Glob J Obes Diabetes Metab Syndr* 2015;2(1):12-7. <https://doi.org/10.17352/2455-8583.000009>
- Lee R, Ahn HR, Shin MH, Kim HN, Lee YH, Choi SW, et al. Association of plasma pentraxin-3 level with lipid levels and cardiovascular risk factors in people with no history of lipid-lowering medication: The Dong-gu study. *J Atheroscler Thromb* 2019;26(8):738-45. <https://doi.org/10.5551/jat.47167>
- Tofano RJ, Salzedas LM, Chagas EF, Detregiachi CR, Guiguer EL, Araujo AC, et al. Association of metabolic syndrome and hyperferritinemia in patients at cardiovascular risk. *Diabetes Metab Syndr* 2020;13:323-48. <https://doi.org/10.2147/DMSO.S271050>
- Shim YS, Kang MJ, Oh YJ, Baek JW, Yang S, Hwang IT. Association of serum ferritin with insulin resistance, abdominal obesity, and metabolic syndrome in Korean adolescents and adults: The Korean National Health and Nutrition Examination Survey, 2008 to 2011. *Medicine (Baltimore)* 2017;96(8):e6179. <https://doi.org/10.1097/md.00000000000006179>
- Lee HA, Choi EJ, Park B, Lee H, Hong YS, Kim HS, et al. The association between metabolic components and markers of inflammatory and endothelial dysfunction in adolescents, based on the Ewha Birth and Growth Cohort Study. *PLoS One* 2020;15(5):e0233469. <https://doi.org/10.1371/journal.pone.0233469>
- Takashi Y, Koga M, Matsuzawa Y, Saito J, Omura M, Nishikawa T. Circulating pentraxin 3 is positively associated with chronic hyperglycemia but negatively associated with plasma aldosterone concentration. *PLoS One* 2018;13(5):e0196526. <https://doi.org/10.1371/journal.pone.0196526>
- Zhou B, Liu S, Yuan G. Combined associations of serum ferritin and body size phenotypes with cardiovascular risk profiles: A Chinese population-based study. *Front Public Health* 2021;9:550011. <https://doi.org/10.3389/fpubh.2021.550011>
- Gündogan K, Bayram F, Capak M, Tanrıverdi F, Karaman A, Öztürk A, et al. Prevalence of metabolic syndrome in the Mediterranean region of Turkey: Evaluation of hypertension, diabetes mellitus, obesity, and dyslipidemia. *Metab Syndr Relat Disord* 2009;7(5):427-34. <https://doi.org/10.1089/met.2008.0068>
- Mehndiratta N, Shama S, Sharma RK, Grover S. A prospective study on the incidence of metabolic syndrome in premenopausal and postmenopausal women. *J Midlife Health* 2020;11(1):17-21. https://doi.org/10.4103/jmh.jmh_57_19
- Bao J, Wang L, Hu P, Liu J, Tu J, Wang J, et al. Burden of metabolic syndrome among a low-income population in China: A population-based cross-sectional study. *Diabetes Metab Syndr* 2022;15:2713-23. <https://doi.org/10.2147/DMSO.S377490>
- Stephens CR, Easton JF, Robles-Cabrera A, Fossion R, de la Cruz L, Martínez-Tapia R, et al. The impact of education and age on metabolic disorders. *Front Public Health* 2020;8:180. <https://doi.org/10.3389/fpubh.2020.00180>
- Farhadnejad H, Parastouei K, Rostami H, Mirmiran P, Azizi F. Dietary and lifestyle inflammatory scores are associated with increased risk of metabolic syndrome in Iranian adults. *Diabetol Metab Syndr* 2021;13(1):30. <https://doi.org/10.1186/s13098-021-00648-1>
- Hong GB, Gao PC, Chen YY, Xia Y, Ke XS, Shao XF, et al. High-sensitivity C-reactive protein leads to increased incident metabolic syndrome in women but not in men: A five-year follow-up study in a Chinese population. *Diabetes Metab Syndr* 2020;13:581-90. <https://doi.org/10.2147/dmsd.s241774>
- Carrero JJ, Franko MA, Obergfell A, Gabrielsen A, Jernberg T. HsCRP level and the risk of death or recurrent cardiovascular events in patients with myocardial infarction: A healthcare-based study. *J Am Heart Assoc* 2019;8(11):e012638. <https://doi.org/10.1161/JAHA.119.012638>
- Yilmaz SK, Eskici G, Mertoglu C, Ayaz A. Adipokines, inflammation, oxidative stress: Critical components in obese women with metabolic syndrome. *Prog Nutr* 2021;23:1-12. <https://doi.org/10.23751/pn.v23i1.9072>
- Kurylowicz A. Estrogens in adipose tissue physiology and obesity-related dysfunction. *Biomedicines* 2023;11(3):690. <https://doi.org/10.3390/biomedicines11030690>
- Somani YB, Pawelczyk JA, De Souza MJ, Kris-Etherton PM, Proctor DN. Aging women and their endothelium: Probing the relative role of estrogen on vasodilator function. *Am J Physiol Heart Circ Physiol* 2019;317(2):H395-404. <https://doi.org/10.1152/ajpheart.00430.2018>
- Grundey SM, Cleeman JI, Daniels SR, Donato KA, Eckel RH, Franklin BA, et al. Diagnosis and management of the metabolic syndrome: An American Heart Association/National Heart, Lung, and Blood Institute scientific statement. *Circulation* 2005;112(17):2735-52. <https://doi.org/10.1161/circulationaha.105.169404>
- Silva IV, De Figueiredo RC, Rios DR. Effect of different classes of antihypertensive drugs on endothelial function and inflammation. *Int J Mol Sci* 2019;20(14):3458. <https://doi.org/10.3390/ijms20143458>
- Proute MC, Kothur N, Georgiou P, Serhiyenia T, Shi W, Kerolos ME, et al. The effect of statin therapy on inflammatory biomarkers: A systematic review. *Cureus* 2021;13(9):e18273. <https://doi.org/10.7759/cureus.18273>
- Bulcão C, Ribeiro-Filho FF, Sañudo A, Ferreira SG. Effects of simvastatin and metformin on inflammation and insulin resistance in individuals with mild metabolic syndrome. *Am J Cardiovasc Drugs* 2007;7:219-24. <https://doi.org/10.2165/00129784-200707030-00007>
- Liu S, Qu X, Liu F, Wang C. Pentraxin 3 as a prognostic biomarker in patients with systemic inflammation or infection. *Mediators Inflamm* 2014;2014:421429. <https://doi.org/10.1155/2014/421429>
- Karakas MF, Buyukkaya E, Kurt M, Motor S, Akcay AB, Karakas E, et al. Serum pentraxin-3 levels are associated with the severity of metabolic syndrome. *Med Princ Pract* 2013;22(3):274-9. <https://doi.org/10.1159/000343904>
- Kardas F, Akın L, Kurtoglu S, Kendirci M, Kardas Z. Plasma pentraxin 3 as a biomarker of metabolic syndrome. *Indian J Pediatr* 2015;82(1):35-8. <https://doi.org/10.1007/s12098-014-1542-0>
- Yamasaki K, Kurimura M, Kasai T, Sagara M, Kodama T, Inoue K. Determination of physiological plasma pentraxin 3 (PTX3) levels in healthy populations. *Clin Chem Lab Med* 2009;47(4):471-7. <https://doi.org/10.1515/CCLM.2009.110>
- Zanetti M, Bosutti A, Ferreira C, Vinci P, Biolo G, Fonda M, et al. Circulating pentraxin 3 levels are higher in metabolic syndrome with subclinical atherosclerosis: Evidence for association with atherogenic lipid profile. *Clin Exp Med* 2009;9(3):243-8. <https://doi.org/10.1007/s10238-009-0039-z>
- Ogawa T, Kawano Y, Imamura T, Kawakita K, Sagara M, Matsuo T, et al. Reciprocal contribution of pentraxin 3 and C-reactive protein to obesity and metabolic syndrome. *Obesity (Silver Spring)* 2010;18(9):1871-4. <https://doi.org/10.1038/oby.2009.507>
- Devissoglu P, Elmas B. Pentraxin 3 as a marker for cardiovascular disease risk in overweight and obese children. *Acta Cardiol Sin* 2021;37(2):177-83. [https://doi.org/10.6515/acs.202103_37\(2\).20201006a](https://doi.org/10.6515/acs.202103_37(2).20201006a)
- Chand MR, Agarwal N, Nishanth D, Jhuma S. Association of serum ferritin levels with

- metabolic syndrome in India: A cross-sectional study. *Maedica (Bucur)* 2021;16:48-53.
<https://doi.org/10.26574/maedica.2020.16.1.48>
33. Tran TN, Tran HD, Tran-Huu TT, Tran DM, Tran QN. A cross-sectional study of serum ferritin levels in Vietnamese adults with metabolic syndrome. *Diabetes Metab Syndr Obes* 2022;15:1517-23.
<https://doi.org/10.2147/dms0.S360689>
34. Han LL, Wang YX, Li J, Zhan XL, Bian C, Wang H, et al. Gender differences in associations of serum ferritin and diabetes, metabolic syndrome, and obesity in the China Health and Nutrition survey. *Mol Nutr Food Res* 2014;58(11):2189-95.
<https://doi.org/10.1002/mnfr.201400088>
35. Kent JW Jr., Comuzzie AG, Mahaney MC, Almasy L, Rainwater DL, VandeBerg JL, et al. Intercellular adhesion molecule-1 concentration is genetically correlated with insulin resistance, obesity, and HDL concentration in Mexican Americans. *Diabetes* 2004;53(10):2691-5.
<https://doi.org/10.2337/diabetes.53.10.2691>
36. Zhang X, Seman NA, Falhammar H, Brismar K, Gu HF. Genetic and biological effects of ICAM-1 E469K polymorphism in diabetic kidney disease. *J Diabetes Res* 2020;2020:8305460.
<https://doi.org/10.1155/2020/8305460>

Inflammatory and endothelial dysfunction markers in females with metabolic syndrome aged 50–75 years

ORIGINALITY REPORT

20%
SIMILARITY INDEX

15%
INTERNET SOURCES

12%
PUBLICATIONS

4%
STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|-----|
| 1 | Submitted to American University of Beirut
Student Paper | 1 % |
| 2 | jmidlifehealth.org
Internet Source | 1 % |
| 3 | Nuha A. Alkhattabi, Maha J. Balgoon, Afnan A. Alraddadi, Maryam H. Al-Zahrani, Rana A. Alghamdi. "Association of Apelin gene rs3115757 variation and expression with metabolic syndrome among female students in King Abdulaziz University", Gene Reports, 2021
Publication | 1 % |
| 4 | www.tsoc.org.tw
Internet Source | 1 % |
| 5 | www.tkd-online.org
Internet Source | 1 % |
| 6 | jmhg.springeropen.com
Internet Source | 1 % |

7	J. W. Kent, A. G. Comuzzie, M. C. Mahaney, L. Almasy, D. L. Rainwater, J. L. VandeBerg, J. W. MacCluer, J. Blangero. "Intercellular Adhesion Molecule-1 Concentration Is Genetically Correlated With Insulin Resistance, Obesity, and HDL Concentration in Mexican Americans", Diabetes, 2004 Publication	1 %
8	link.springer.com Internet Source	1 %
9	worldwidescience.org Internet Source	<1 %
10	www.endocrino.org.br Internet Source	<1 %
11	www.nature.com Internet Source	<1 %
12	www.dovepress.com Internet Source	<1 %
13	drum.lib.umd.edu Internet Source	<1 %
14	www.jstage.jst.go.jp Internet Source	<1 %
15	dmsjournal.biomedcentral.com Internet Source	<1 %

- | | | |
|----|--|------|
| 16 | Sevilay Vural, Levent Albayrak. "Can calcitonin gene-related peptide (CGRP) and pentraxin-3 (PTX-3) be useful in diagnosing acute migraine attack?", Journal of Receptors and Signal Transduction, 2022
Publication | <1 % |
| 17 | pubmed.ncbi.nlm.nih.gov
Internet Source | <1 % |
| 18 | Submitted to Fakultas Kedokteran Gigi Universitas Trisakti
Student Paper | <1 % |
| 19 | Submitted to Mississippi State Board for Community & Junior Colleges
Student Paper | <1 % |
| 20 | Parimita Routray, Belen Torondel, Thomas Clasen, Wolf-Peter Schmidt. "Women's role in sanitation decision making in rural coastal Odisha, India", PLOS ONE, 2017
Publication | <1 % |
| 21 | Tawfiq Almadi, Ian Cathers, Chin Moi Chow. "Associations among work-related stress, cortisol, inflammation, and metabolic syndrome", Psychophysiology, 2013
Publication | <1 % |
| 22 | jurnal.unismuhpalu.ac.id
Internet Source | <1 % |

23	"Insulin Resistance", Springer Science and Business Media LLC, 2020 Publication	<1 %
24	Submitted to The Robert Gordon University Student Paper	<1 %
25	www.revistas.usp.br Internet Source	<1 %
26	Submitted to University of Duhok Student Paper	<1 %
27	jpma.org.pk Internet Source	<1 %
28	www.scitepress.org Internet Source	<1 %
29	real.mtak.hu Internet Source	<1 %
30	Alyssa B. Schultz. "Analysis of the Association between Metabolic Syndrome and Disease in a Workplace Population over Time", Value in Health, 03/2010 Publication	<1 %
31	Giuseppe Danilo Norata, Patrizia Marchesi, Vivek Krishna Pulakazhi Venu, Fabio Pasqualini et al. "Deficiency of the Long Pentraxin PTX3 Promotes Vascular Inflammation and Atherosclerosis", Circulation, 2009	<1 %

32

Jie Bao, Lifeng Wang, Peng Hu, Jie Liu, Jun Tu, Jinghua Wang, Jidong Li, Xianjia Ning. "Burden of Metabolic Syndrome Among a Low-Income Population in China: A Population-Based Cross-Sectional Study", Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy, 2022

Publication

<1 %

33

S. R. Srinivasan, M. G. Frontini, J. Xu, G. S. Berenson. "Utility of Childhood Non-High-Density Lipoprotein Cholesterol Levels in Predicting Adult Dyslipidemia and Other Cardiovascular Risks: The Bogalusa Heart Study", PEDIATRICS, 2006

Publication

<1 %

34

benthamscience.com

Internet Source

<1 %

35

lib.ui.ac.id

Internet Source

<1 %

36

natap.org

Internet Source

<1 %

37

topsecretapiaccess.dovepress.com

Internet Source

<1 %

38

Ayumi Nakamura, Shin-ichiro Miura, Yuhei Shiga, Kenji Norimatsu, Yuiko Miyase, Yasunori Suematsu, Ryoko Mitsutake, Keijiro

<1 %

Saku. "Is pentraxin 3 a biomarker, a player, or both in the context of coronary atherosclerosis and metabolic factors?", Heart and Vessels, 2014

Publication

39

Fikret Akyurek, Fatma Tunccez Akyurek. "Investigation of pregnancy - associated plasma protein A and neopterin levels in Behçet's patients", Dermatologic Therapy, 2020

Publication

40

Israel S. Ribeiro, Ítalo S. Pereira, Denisar P. Santos, Danilo N. Lopes et al. "Association between body composition and inflammation: A central role of IL-17 and IL-10 in diabetic and hypertensive elderly women", Experimental Gerontology, 2019

Publication

41

Lee, Mi Kyung, Sang Hui Chu, Duk Chul Lee, Ki Yong An, Ji-Hye Park, Dong Il Kim, Jiyoung Kim, Sunghyun Hong, Jee Aee Im, Ji Won Lee, and Justin Y. Jeon. "The association between chemerin and homeostasis assessment of insulin resistance at baseline and after weight reduction via lifestyle modifications in young obese adults", Clinica Chimica Acta, 2013.

Publication

<1 %

<1 %

<1 %

- | | | |
|----|---|------|
| 42 | Yangting Zhao, Xiaoyu Lv, Chongyang Chen, Kai Li, Yawen Wang, Jingfang Liu. "The association between triglyceride-glucose index and hyperferritinemia in patients with type 2 diabetes mellitus", Hormones, 2023
Publication | <1 % |
| 43 | researchers.mq.edu.au
Internet Source | <1 % |
| 44 | rossstores.com
Internet Source | <1 % |
| 45 | www.eje-online.org
Internet Source | <1 % |
| 46 | www.ijscia.com
Internet Source | <1 % |
| 47 | Ehab K. El Melegy, Eman A. Badr, Ahmed M. ElKersh, Wassam H. EL Shafey, Walaa A. Fareed. "Pentraxin 3 genotyping in relation to serum levels of pentraxin 3 in patients with acute ST-segment elevation myocardial infarction", Clinical Trials and Regulatory Science in Cardiology, 2016
Publication | <1 % |
| 48 | J. - F. Yang, X. - F. Xiong, Y. Xiao, L. Wei et al. "The single nucleotide polymorphism rs11643718 in is associated with the development of diabetic kidney disease in | <1 % |

Chinese people with type 2 diabetes ", Diabetic Medicine, 2020

Publication

49

Liang-Yin Ke, Hua-Chen Chan, Hsiu-Chuan Chan, Franklin Chikodi Udo Kalu et al.
"Electronegative Low-density Lipoprotein L5 Induces Adipose Tissue Inflammation Associated with Metabolic Syndrome", The Journal of Clinical Endocrinology & Metabolism, 2017

Publication

<1 %

50

Thallapaneni Sasikala, Suchitra M Manohar, Aparna RR Bitla, S Sarala, Suresh Vaikkakara.
"Intercellular adhesion molecule-1 is a surrogate biomarker for subclinical atherosclerosis in Type 2 diabetes mellitus", Biomarkers in Medicine, 2021

Publication

<1 %

51

bmcgastroenterol.biomedcentral.com

Internet Source

<1 %

52

edepot.wur.nl

Internet Source

<1 %

53

ijmrr.medresearch.in

Internet Source

<1 %

54

journal.fk.unpad.ac.id

Internet Source

<1 %

services.rmh.med.sa

55

Internet Source

<1 %

56

turkishjcrd.com

Internet Source

<1 %

57

wjgnet.com

Internet Source

<1 %

58

www.aafp.org

Internet Source

<1 %

59

www.repository.unipr.it

Internet Source

<1 %

60

www.reproductive-health-journal.com

Internet Source

<1 %

61

www.unchina.org

Internet Source

<1 %

62

A Zlibut, IC Bocsan, RM Pop, SC Vesa, K Bheecarry, R Revnic, B Cojan-Minzat, S Lupu, AD Buzoianu, L Agoston-Coldea. "Role of pentraxin-3 in risk assessment of patients with metabolic syndrome", Physiology International, 2019

Publication

<1 %

63

F Gündoğan, A Bayram, M Kalkan, İ Özcan. "Plasma levels of endothelial cell-specific molecule-1 and pentraxin-3 in idiopathic sudden sensorineural hearing loss", The Journal of Laryngology & Otology, 2018

<1 %

64

Asiyeh Sadat Zahedi, Mahdi Akbarzadeh, Bahareh Sedaghati-Khayat, Atefeh Seyedhamzehzadeh, Maryam S. Daneshpour. "GCKR common functional polymorphisms are associated with metabolic syndrome and its components: a 10-year retrospective cohort study in Iranian adults", *Diabetology & Metabolic Syndrome*, 2021

Publication

<1 %

65

Caroline Bulc??o. "Effects of Simvastatin and Metformin on Inflammation and Insulin Resistance in Individuals with Mild Metabolic Syndrome", *American Journal of Cardiovascular Drugs*, 2007

Publication

<1 %

66

Hamideh Ghazizadeh, Majid Rezaei, Amir Avan, Mohammad Fazilati et al. "Association between serum cell adhesion molecules with hs-CRP, uric acid and VEGF genetic polymorphisms in subjects with metabolic syndrome", *Molecular Biology Reports*, 2019

Publication

<1 %

67

Mahdi Akbarzadeh, Parisa Riahi, Azra Ramezankhani, Saeid Rasekhi Dehkordi et al. "Parental Transmission Plays the Major Role in High Aggregation of Type 2 Diabetes in

<1 %

Iranian Families: Tehran Lipid and Glucose Study", Canadian Journal of Diabetes, 2021

Publication

68

Natsuki Ohnishi, Yuhei Shiga, Kohei Tashiro, Yuto Kawahira et al. "Association between major adverse cardiovascular events and pentraxin-3 in patients who have undergone coronary computed tomography angiography: from the FU-CCTA registry", Heart and Vessels, 2022

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Inflammatory and endothelial dysfunction markers in females with metabolic syndrome aged 50–75 years

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7