



Vol.6 No.3 | Desember 2023

# JURNAL BIOMEDIKA DAN KESEHATAN

Publikasi dari Fakultas Kedokteran Universitas Trisakti

## Editorial

JN.1 COVID-19: Variant of interest  
*Husnun Amalia, Yasmine Mashabi*

## Original Article

Vitamin D Level with Dry Eye Syndrome in the Employees  
*Eveline Margo, Meiyanti Meiyanti, Erlani Kartadinata et al*

Relationship between Magnesium Intake and Handgrip Strength in Elderly  
*Muhammad Iqbal Arigi Putra, Elly Herwana*

Pulmonary Function Test and its Correlation with Exhaled Carbon Monoxide and Smoking Habits in Ojek Drivers  
*Rita Khairani, Mustika Anggiane Putri, Dyah Ayu Woro Setyaningrum*

The Association of Laboratory Parameters with COVID-19 Severity in Pregnancy  
*Rully Ayu Nirmalasari, Lily Marlany Surjadi, Laksmi Maharani et al*

The Relation between Sleep Quality and Screen Time with Student Learning Concentration  
*Ratu Aurany Desnissa, Nuryani Sidarta*

The Risk Factors Associated with Handgrip Strength and Endurance in Adolescence  
*Ni Putu Laksmi Martini, Dian Mediana*

Ojek Driver's Knowledge of Basic Life Support in Daily After Training  
*Diani Harahap, Antin Trilaksmi, Lira Panduwati et al*

Relationship Between Fatigue and Burnout Syndrome Among Residents in Anesthesiology and Intensive Therapy Department  
*Feni Venawati, Andriamuri Prima Lubis, Christmas Gideon Bangun et al*

## Case Report

A Scarce Case of Sarcomatoid Bladder Carcinoma and Bladder Stone in a 58-Year-Old Man: A Case Report  
*Vincent Vincent, Mikha Mikha, Aries Alpendri*

Metachronous Multifocal Osteosarcoma After 5-month Therapy: Metastasis or Other Primary Lesion?  
*Astien Astien, Nany Hairunisa, Nugroho Abikusno et al*

## Review Article

Medical and Ethical Studies Cannabis/Marijuana Use Controversy in Indonesia: A Literature Review  
*Ervin Dyah Ayu Masita Dewi, Jefman Efendi Marzuki H. Y, Rico Mulyono*

The Risk of Latent Tuberculosis Reactivation in COVID-19 Therapy  
*Ghina Mutiara Abas, Fithriyah Sjatha, Yeva Rosana*

Chronic Kidney Disease : is plant-based diet effective ?  
*Wawan Kurniawan, Nany Hairunisa, Gerie Amarendra*

# Dewan Redaksi



## **Ketua Penyunting (Editor-in-Chief)**

Dr. dr. Husnun Amalia, Sp.M  
Departemen Ilmu Penyakit Mata, Fakultas Kedokteran  
Universitas Trisakti, Indonesia

## **Wakil Ketua Penyunting (Deputy Editor-in-Chief)**

Dr. Drs. ML. Edy Parwanto, M.Biomed  
Departemen Biologi Kedokteran, Fakultas Kedokteran  
Universitas Trisakti, Indonesia

## **Penyunting Ahli (Associate Editor)**

dr. Nany Hairunisa, MCHSc  
Departemen Ilmu Kedokteran Kerja, Fakultas Kedokteran  
Universitas Trisakti, Indonesia

## **Dewan Penyunting (Editorial Boards)**

Prof. Dr. dr. Adi Hidayat, MS (Indonesia)  
Dr. dr. Elly Herwana, M.Biomed (Indonesia)  
Dr. dr. Yenny, Sp.FK (Indonesia)  
dr. Laksmi Maharani, Sp.OG (Indonesia)  
dr. Monica Dwi Hartanti, M.Biomed, PhD (Indonesia)  
Dr. dr. Raditya Wratsangka, Sp.O.G, Subsp. Obginsos (Indonesia)  
Dr. Siti Sugih Hartiningsih, S.Si, M.Kes (Indonesia)  
dr. Dito Anurogo, M.Sc (Indonesia)

## **Editor Produksi**

Afton Muhandis, S.I.Kom

## **Alamat Korespondensi**

Fakultas Kedokteran Universitas Trisakti  
Jalan Kyai Tapa Np. 260 (Kampus B) Grogol, Jakarta 11440  
Telp. 021-5672731 ext. 2502 | Fax. 021-5660706  
[www.jbiomedkes.org](http://www.jbiomedkes.org) | E-mail: [jbiomedkes@trisakti.ac.id](mailto:jbiomedkes@trisakti.ac.id)

## **Penerbit**

Fakultas Kedokteran Universitas Trisakti

# Petunjuk Penulisan

## Format penyusunan manuskrip

Manuskrip diketik pada kertas berukuran A4 (210 x 297 mm) dengan batas tepi 254 mm (*margin Normal*), huruf diketik dengan tipe huruf (*font*) *Times New Roman*, besar huruf (*font size*) 12 point dengan menggunakan spasi rangkap 2 (*double space*). Setiap bagian dari manuskrip dimulai pada halaman baru dengan urutan sebagai berikut: halaman judul, abstrak dan kata kunci (*keywords*), teks keseluruhan, ucapan terima kasih, daftar pustaka, tabel dan gambar (setiap tabel dan gambar pada halaman terpisah). Nomor halaman dicantumkan secara berurutan dimulai dari halaman judul pada sudut sebelah kanan bawah. Manuskrip sebaiknya ditulis maksimal 16 halaman.

## Halaman judul

Halaman judul mencakup: a) judul manuskrip yang dibuat sesingkat mungkin, spesifik informatif dan ringkasan judul tidak lebih dari 40 karakter (hitung huruf dan spasi) yang dicantumkan dibawah judul, b) nama penulis disusun berurutan dengan nama mahasiswa sebagai pengarang pertama, diikuti oleh Pembimbing sebagai pengarang kedua. Nama penulis ditulis lengkap tanpa gelar dan dicantumkan seperti aslinya, tidak dibalik seperti pada daftar pustaka dan sitasi, c) alamat setiap penulis, nama departemen dan lembaga afiliasi penulis, d) nama dan alamat penulis untuk korespondensi serta nomor telepon, nomor faksimili, alamat email. Judul penelitian dibuat jelas, singkat, spesifik, informatif, dan sesuai dengan topik manuskrip. Jumlah kata tidak lebih dari 12 kata agar mudah dan cepat dipahami pembaca.

## Abstrak dan kata kunci

Abstrak berjumlah 200-250 kata ditulis dalam bahasa Indonesia dan Inggris. Abstrak berisikan latar belakang termasuk tujuan penelitian, metode, hasil, dan kesimpulan. Kata kunci dicantumkan di bawah abstrak pada halaman yang sama sebanyak 4-6 kata. Bagian abstrak merupakan ringkasan dari isi makalah yang dibuat secara singkat, informatif, dengan menekankan pada aspek baru dan penting dari penelitian.

## Teks

Teks makalah manuskrip dibagi dalam beberapa bagian dengan judul sebagai berikut: **Pendahuluan, Metode, Hasil, Pembahasan, Kesimpulan dan saran.**

## Pendahuluan

a. Latar belakang merupakan bagian yang menjelaskan alasan mengapa masalah ini penting untuk diteliti. Bagian ini memuat penjelasan mengapa masalah itu dipandang menarik, penting, dan perlu diteliti untuk mencari pemecahannya. Penjelasan dapat diperoleh dari penelusuran pustaka yang berkaitan erat dengan

masalah yang diteliti.

b. Keaslian penelitian dikemukakan dengan menunjukkan bahwa masalah yang dihadapi belum pernah dipecahkan oleh peneliti terdahulu atau dinyatakan dengan tegas perbedaan penelitian ini dengan penelitian terdahulu.

c. Tujuan penelitian yang menjelaskan hasil yang akan dicapai.

## Metode

Metode penelitian berisi uraian terpadu dan sistematis mengenai bagaimana penelitian akan dilaksanakan. Metode terdiri dari :

a. Desain

b. Populasi / sampel (subjek) penelitian

Diuraikan kriteria inklusi dan eksklusi subjek penelitian, cara pemilihan sampel (subjek penelitian) secara random atau non-random, serta besar sampel yang akan di pilih. Teknik pemilihan sampel harus dijelaskan secara rinci. Bila perlu dibuat alur pemilihan sampel.

c. Bahan dan alat serta pengukuran

Bahan dan alat yang harus disajikan pada laporan terbatas pada bahan (materi) dan alat utama yang diperlukan untuk penelitian dan harus disebutkan spesifikasinya. Prosedur pengukuran perlu dijelaskan sesuai dengan tahapan yang dilakukan.

d. Alur kerja penelitian

Jalannya penelitian perlu dijelaskan mengenai jenis pendekatan yang dipakai untuk mendapatkan data, melalui pendekatan laboratorium, klinik, komunitas, observasi, dll.

e. Analisis data

Perlu dijelaskan jenis teknik statistik yang digunakan untuk menjawab masalah dan mencapai tujuan penelitian. Data yang diperoleh dapat dianalisis menggunakan teknik statistik secara parametrik dan non-parametrik.

## Hasil

Suatu hasil penelitian hendaknya disajikan dengan jelas, logis, runut, sehingga mudah untuk dimengerti. Hasil penelitian sebaiknya ditampilkan selain dalam bentuk narasi dapat pula berupa gambar, tabel, foto, dan grafik sehingga memudahkan untuk dipahami. Hasil dan interpretasi analisis statistik dituliskan secara jelas dalam uraian hasil penelitian.

Pada tahap awal disajikan distribusi karakteristik subjek penelitian, yang biasanya dibuat pada sebuah tabel. Kemudian disajikan temuan penting yang diperoleh, kalau cukup banyak sebaiknya pada sebuah tabel. Bila terbatas misalkan hanya satu atau dua temuan cukup dalam bentuk narasi/teks.

Tabel, bagan/gambar, grafik dibuat dengan jelas, diberi nomor urut serta keterangan yang jelas. Keterangan

tabel diletakkan di atas tabel dan keterangan gambar diletakkan di bawah gambar. Maksimal tabel dan gambar 5. Semua tabel, grafik dan gambar diberi nomor dan keterangan yang jelas. Setiap tabel dianalisis dan diinterpretasi secara sistematis, dan hasilnya ditulis di bawah tabel tersebut. Perhitungan statistik detail tidak perlu ditulis dalam bagian hasil ini. Bila perhitungan statistik dianggap perlu ditulis, maka sebaiknya diletakkan dalam lampiran saja.

### **Pembahasan**

Langkah awal harus diuraikan temuan penting yang diperoleh dari penelitian sesuai dengan tujuan penelitian. Kemudian bandingkan hasil penelitian yang diperoleh dengan hasil-hasil penelitian sebelumnya. Perlu dijelaskan kesesuaian dan ketidaksesuaian hasil penelitian yang didapat terhadap kerangka teori atau hasil penelitian lain yang telah dilakukan sebelumnya. Selanjutnya menggunakan teori-teori yang ada uraikan mekanisme terjadinya hasil penelitian tersebut. Bagian pembahasan juga menjelaskan mengenai kelemahan dan kelebihan penelitian yang telah dilakukan. Uraikan implikasi dari hasil penelitian yang diperoleh.

### **Kesimpulan**

Kesimpulan hendaknya dibuat dalam bentuk narasi dan menguraikan secara singkat, jelas, padat menurut urutan yang sistematis. Bagian ini memuat tentang hasil penelitian yang telah diperoleh untuk menjawab tujuan penelitian. Saran menguraikan perlunya dilakukan penelitian lebih lanjut untuk memperbaiki kelemahan/keterbatasan dari penelitian yang telah dilakukan.

### **Ucapan terima kasih**

Ditujukan kepada pihak-pihak yang memberikan bantuan dana dan dukungan antara lain dukungan dari bagian dan lembaga, para profesional yang memberikan kontribusi dalam penyusunan makalah, dan untuk penguji I maupun penguji II. Pembimbing tidak perlu dicantumkan pada Ucapan Terima Kasih karena sudah dicantumkan sebagai penulis.

### **Daftar Referensi**

Daftar referensi/rujukan hanya mencatumkan rujukan yang telah digunakan dan ditulis menurut sistem Vancouver.

### **Online Submission**

Penulis dapat mengirim naskah manuskrip melalui *online submission* di *website* Jurnal Biomedika dan Kesehatan.

Langkah online submission:

1. Pastikan naskah yang akan diunggah sudah mengikuti semua petunjuk penulisan
2. Lakukan pendaftaran author di :  
<https://jbiomedkes.org/index.php/jbk/user/register>
3. Setelah terdaftar silakan unggah naskah manuskrip dan isi form yang terdapat di dalam website, dan ikuti langkah selanjutnya.

### **Daftar Cek Pengiriman Naskah Manuskrip**

- ☐ Naskah manuskrip belum pernah dipublikasikan sebelumnya, juga tidak dalam pengajuan ke jurnal lain.
- ☐ File manuskrip harus berformat OpenOffice, Ms. Word atau RTF dokumen, *font* 12, *Times New Roman*, *double spacing*.
- ☐ Halaman judul harus memuat jelas judul, nama lengkap penulis tanpa gelar, departemen penulis, universitas, alamat lengkap, nomor telepon dan email.
- ☐ Pelaporan data manuskrip dari penelitian yang melibatkan manusia dan hewan memerlukan persetujuan formal (kaji etik) oleh dewan peninjau atau komisi etik institusi yang bersangkutan.
- ☐ Daftar rujukan memuat semua rujukan yang terdapat di dalam manuskrip dan ditulis sesuai urutan pengutipannya menggunakan sistem Vancouver.

# Daftar Isi



*Jurnal Biomedika dan Kesehatan - Vol. 5 No. 3 December 2022*

|                                                                                                                    |         |
|--------------------------------------------------------------------------------------------------------------------|---------|
| Editorial                                                                                                          | ▼       |
| <b>JN.1 COVID-19: Variant of interest</b>                                                                          |         |
| Husnun Amalia, Yasmine Mashabi                                                                                     | 261-263 |
| <a href="#">PDF</a>                                                                                                |         |
| <hr/>                                                                                                              |         |
| Original Article                                                                                                   | ▼       |
| <b>Vitamin D Level with Dry Eye Syndrome in the Employees</b>                                                      |         |
| Eveline Margo, Meiyanti Meiyanti, Erlani Kartadinata, Noviani Kartadinata                                          | 264-274 |
| <a href="#">PDF</a>                                                                                                |         |
| <b>Relationship between Magnesium Intake and Handgrip Strength in Elderly</b>                                      |         |
| Muhammad Iqbal Arigi Putra, Elly Herwana                                                                           | 275-282 |
| <a href="#">PDF</a>                                                                                                |         |
| <b>Pulmonary Function Test and its Correlation with Exhaled Carbon Monoxide and Smoking Habits in Ojek Drivers</b> |         |
| Rita Khairani, Mustika Anggiane Putri, Dyah Ayu Woró Setyaningrum                                                  | 283-290 |
| <a href="#">PDF</a>                                                                                                |         |
| <b>The Association of Laboratory Parameters with COVID-19 Severity in Pregnancy</b>                                |         |
| Rully Ayu Nirmalasari, Lily Marliany Surjadi, Laksmi Maharani, Yuyun Lisnawati, Dhea Mangun                        | 291-299 |
| <a href="#">PDF</a>                                                                                                |         |
| <b>The Relation between Sleep Quality and Screen Time with Student Learning Concentration</b>                      |         |
| Ratu Aurany Desnissa, Nuryani Sidarta                                                                              | 300-309 |
| <a href="#">PDF</a>                                                                                                |         |
| <b>The Risk Factors Associated with Handgrip Strength and Endurance in Adolescence</b>                             |         |
| Ni Putu Laksmi Martini, Dian Mediana                                                                               | 310-319 |
| <a href="#">PDF</a>                                                                                                |         |
| <b>Ojek Driver's Knowledge of Basic Life Support in Daily After Training</b>                                       |         |
| Diani Nazma, Antin Trilaksmi, Lira Panduwati, M Ridho Devantoro                                                    | 320-329 |
| <a href="#">PDF</a>                                                                                                |         |

### Relationship Between Fatigue and Burnout Syndrome Among Residents in Anesthesiology and Intensive Therapy Department

Feni Venawati, Andriamuri Prima Lubis, Christmas Gideon Bangun, Yuki Yunanda

330-339



PDF

### Case Report



### A Scarce Case of Sarcomatoid Bladder Carcinoma and Bladder Stone in a 58-Year-Old Man: A Case Report

Vincent Vincent, Mikha Mikha, Aries Alpendri

340-346



PDF

### Metachronous Multifocal Osteosarcoma After 5-month Therapy: Metastasis or Other Primary Lesion?

Astien Astien, Nany Hairunisa, Nugroho Abikusno, Revalita Abikusno, Nadifa Agil, Febrina T

347-353



PDF

### Review Article



### Medical and Ethical Studies Cannabis/Marijuana Use Controversy in Indonesia: A Literature Review

Ervin Dyah Ayu Masita Dewi, Jefman Efendi Marzuki H. Y, Rico Mulyono

354-362



PDF

### The Risk of Latent Tuberculosis Reactivation in COVID-19 Therapy

Ghina Mutiara Abas, Fithriyah Sjatha, Yeva Rosana

363-372



PDF

### Chronic Kidney Disease : is plant-based diet effective ?

Wawan Kurniawan, Nany Hairunisa, Gerie Amarendra

373-382



PDF

ORIGINAL ARTICLE

## Pulmonary Function Test and its Correlation with Exhaled Carbon Monoxide and Smoking Habits in Ojek Drivers

### Korelasi Karbon Monoksida dan Derajat Rokok dengan Fungsi Paru pada Pengemudi Ojek Online

Rita Khairani<sup>1</sup> , Mustika Anggiane Putri<sup>2</sup>, Dyah Ayu Woro Setyaningrum<sup>3</sup>

<sup>1</sup>Department of Internal Medicine, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

<sup>2</sup>Department of Physiology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

<sup>3</sup>Department of Anatomical Pathology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

 rita.khairani@trisakti.ac.id

 <https://doi.org/10.18051/jbk.283-290>

#### ABSTRACT

##### Background

Cigarette smoke is the main source of exposure to carbon monoxide (CO) besides air pollution. When cigarette smoke is inhaled, carbon monoxide is absorbed through the lungs, enters the bloodstream, and then binds to hemoglobin to form carboxyhemoglobin (COHb), whose levels in the blood can be measured as a marker of cigarette smoke absorption. Both smoking habits and CO levels can affect lung function, after smoking, blood CO levels will increase and lung function will decrease significantly.

##### Methods

This study used an analytic observational method with a cross-sectional design with a total sample of 99 respondents. Data collection in September-December 2022 using the Brinkman Index questionnaire, *Fagerstrom Test for Nicotine Dependence* (FTND) questionnaire, exhaled CO levels, and spirometry examination.

##### Results

47.5% exhaled CO levels on 0-6 ppm or in a green zone, 77.8% of subjects smoked, 57.6% FTND score was 0-2, 51.5% of subjects had restricted lung function, and 79.5% subjects with obstruction. Exhaled CO levels were strongly correlated with the Brinkman index ( $r=0.654$ ,  $p=0.000$ ), and moderately significant correlation with the FTND score ( $r=0.544$ ,  $p=0.000$ ). There is no correlation between exhaled CO levels and degree of smoking with lung function.

##### Conclusions

Exhaled CO levels have a significant correlation with smoking and nicotine dependence but do not correlate with pulmonary function test.

**Keywords:** Exhaled carbon monoxide level, smoking, pulmonary function test, nicotine dependence

## ABSTRAK

### Latar Belakang

Asap rokok merupakan sumber utama pajanan terhadap karbon monoksida (CO) selain polusi udara. Saat asap rokok terinhalasi, karbon monoksida akan diabsorpsi melalui paru, masuk ke dalam aliran darah kemudian akan berikatan dengan hemoglobin untuk membentuk karboksi – hemoglobin (COHb) yang kadarnya dalam darah dapat diukur sebagai marker absorpsi asap rokok. Baik rokok atau CO dapat mempengaruhi fungsi paru, setelah merokok maka kadar CO dalam darah akan meningkat dan fungsi paru akan menurun secara bermakna.

### Metode

Penelitian ini menggunakan metode observasional analitik dengan desain potong lintang dengan besar sampel 99 responden. Pengambilan data pada bulan September-Desember 2022 dengan menggunakan kuesioner Indeks Brinkman, *Fagerstrom Test for Nicotine Dependence* (FTND), pengukuran CO ekspirasi dan pemeriksaan spirometri

### Hasil

Sebanyak 47.5% responden memiliki kadar CO ekspirasi sebesar 0-6 ppm yang masih berada pada zona hijau, 77.8% responden merokok, 57.6% memiliki skor FTND 0-2, sebanyak 51.5% mengalami gangguan restriksi dan 79.5% responden mengalami gangguan obstruksi saluran napas. Kadar CO ekspirasi berkorelasi erat dengan indeks Brinkman ( $r=0.654$ ,  $p=0.000$ ), dan berkorelasi sedang bermakna dengan skor FTND ( $r=0.544$ ,  $p=0.000$ ). Tidak ada korelasi kadar CO ekspirasi, indeks Brinkman dan ketergantungan terhadap nikotin dengan fungsi paru.

### Kesimpulan

kadar CO ekspirasi berkorelasi erat dengan rokok dan ketergantungan nikotin tetapi tidak berkorelasi dengan fungsi paru.

**Kata Kunci:** karbon monoksida ekspirasi, rokok, uji fungsi paru, ketergantungan nikotin

## INTRODUCTION

The most common source of air pollution in urban areas comes from motorized vehicles. The development of traffic volume in urban areas has reached 15% every year in the last 10 years. Transportation in big cities is the largest source of air pollution, where 70% of air pollution in urban areas is caused by motorized vehicle activities, especially motorbikes, which reach 30%.<sup>1</sup> Parameters of air pollution from motorized vehicles such as carbon monoxide (CO), nitrogen oxides (NOx), methane (CH<sub>4</sub>), sulfur dioxide (SO<sub>2</sub>) and particulate matter can affect global warming.<sup>1,2</sup>

Apart from air pollution, cigarette smoke is also the main source of exposure to carbon monoxide (CO).<sup>3</sup> Examination of exhaled air CO levels can be used as a biomarker for the degree of smoking. Smoking is a habit that is often found throughout the world, although it is generally known that cigarettes can cause health problems.<sup>4</sup> The percentage of smokers aged 15-19 years has increased in 2020, namely 10.61%, an increase from 10.54% in 2019. More than half (52.1%) of smokers in Indonesia smoked for the first time at the age of 15-19 years, according to Basic Health Research data.<sup>5</sup>

Cigarettes cause damage to almost all organs of the body, when cigarette smoke is inhaled, carbon monoxide will be absorbed through the lungs, enter the bloodstream, and then bind with hemoglobin to form carboxy-hemoglobin (COHb) whose levels in the blood can be measured as a marker for cigarette smoke absorption.<sup>6</sup> Carbon monoxide will remain in the blood for 24 hours after inhalation of cigarette smoke depending on several factors such as gender, physical activity, and respiratory rate.<sup>7</sup> In the blood, CO will then re-enter the alveoli because there is a

concentration gradient in the alveoli, so the levels of CO contained in exhaled air can be measured using a portable CO meter.<sup>8</sup>

Endogenous carbon monoxide is produced primarily by systemic heme oxygenation and is eliminated through respiration. In healthy populations who do not smoke, end-tidal concentrations are in the range of 1-3 ppm.<sup>9</sup> Carbon monoxide is one of the most widely distributed pollutants in the air. CO concentrations are released into the air in the highest quantities among other air pollutants every year except for carbon dioxide (CO<sub>2</sub>) concentrations.<sup>3</sup> In areas with high populations the ratio of mixed CO levels can reach up to 10 ppm. The most important biological characteristic of CO is its ability to bind to hemoglobin to form carboxyhemoglobin (HbCO) which is 200 times more stable than oxyhemoglobin (HbO<sub>2</sub>) which can have fatal consequences because it can disrupt muscle metabolism and intracellular enzyme function due to the stable CO bond.<sup>4,10</sup>

Measurement of expiratory monoxide levels has been used to increase the validity of investigations of the degree of smoking because carbon monoxide measurements are easy to use, non-invasive, and affordable.<sup>11</sup> This method can provide a direct assessment of the degree of smoking and is a potential biomarker for assessing air pollution exposure.<sup>3,4</sup> Environmental exposures other than cigarette smoke are important factors that can differentiate exhaled CO results in passive smokers and non-smokers.<sup>11,12</sup>

Subjectively, a person's level of addiction to cigarettes is assessed using the Fagerstrom Test for Nicotine Dependence questionnaire, a questionnaire that was introduced in 1978.<sup>13,14</sup> This questionnaire contains a series of questions regarding how much a person cannot give up smoking. Objectively, the degree of smoking can be assessed by examining exhaled carbon monoxide levels between smokers and non-smokers and there is a strong correlation between exhaled carbon monoxide and the severity of smoking.<sup>13,15</sup>

Both cigarettes and CO can affect lung function. In research conducted by Yalcin et al., who conducted experimental research, after smoking, CO levels will increase and lung function will decrease significantly.<sup>16</sup> Ejazi et al.'s research also found that expiratory CO levels were significantly negatively correlated with the first-second forced expiratory volume (VEP<sub>1</sub>) in patients with chronic obstructive pulmonary disease (COPD).<sup>17</sup> Different research results were obtained by Salepci et al because there was no relationship between lung function and exhaled CO levels.<sup>12</sup> This study aims to assess the relationship between the degree of smoking and exhaled carbon monoxide levels with lung function in online motorcycle taxi drivers.<sup>18</sup>

## METHODS

This research uses an analytical observational method with a cross-sectional design. Data was collected at the Faculty of Medicine Campus, Universitas Trisakti in September-December 2022. Respondents were selected using consecutive non-random sampling with the required sample size of 94 respondents and rounded up to 100 respondents.

The research population was online motorcycle taxi drivers who met the inclusion criteria (online motorcycle taxi drivers in DKI Jakarta and surrounding areas, male and female, aged 17 years – 60 years, negative antigen swab results, willing to be respondents, and able to carry out spirometry and expiratory CO procedures well). Respondents were excluded if they had a history

of chronic lung disease and the spirometry examination was unacceptable or had no reproducibility). Respondents were then interviewed, filled out questionnaires, spirometry, and exhaled CO examinations.

Interview data includes identity and sociodemographic data, and filling out a questionnaire about smoking habits using the Brinkman index by calculating the number of cigarettes smoked per day and the number of years of smoking. The degree of nicotine dependence uses the Fagerstrom Test for Nicotine Dependence (FTND), with categories light (0-2), moderate (3-4), and severe (5-7). CO levels are checked using the piCO Smokerlyzer device, by holding the respondent's breath for 15 seconds, then exhaling it slowly until it runs out and the results will be visible on the measuring instrument. Light levels/green zone (0-6 ppm), medium/yellow zone (7-10 ppm) and heavy/red zone ( $\geq 11$ ). Lung function examination used a Minato AS-507 spirometer, measuring vital capacity (CV) and first-second expiratory volume (VEP1) which was carried out three times, and the highest value was taken and compared with the predicted value. Lung function test results are then categorized as normal if CV and VEP1 are  $\sim 80\%$ , mild restriction (CV 60-79.9%), moderate restriction (CV 30-59.8%), and severe restriction (CV  $< 30\%$ ), while mild obstruction (VEP1 60% -79.9%), moderate obstruction (VEP1 30-59.9%) and severe obstruction (VEP1  $< 30\%$ ). Statistical test analysis uses the Spearman correlation test using SPSS version 25. This research has passed ethical review by the ethics commission of the Faculty of Medicine, Universitas Trisakti, with number 166/KER/FK/VIII/2022.

## RESULTS

Univariate analysis was used to determine the frequency distribution of subject characteristics in the form of gender, age, serum 25(OH)D levels, Schirmer test, TBUT, and OSDI scores.

Table 1. Data on subject characteristics (N=57)

| Variable             |    | Frequency (%) | Mean $\pm$ SD    |
|----------------------|----|---------------|------------------|
| Gender               |    |               |                  |
| Men                  | 26 | (41.3%)       |                  |
| Women                | 31 | (67.7%)       |                  |
| Age (year)           |    |               |                  |
| 25 – 40              | 24 | (42.1%)       | 42.0 $\pm$ 8.7   |
| > 40                 | 33 | (57.9%)       |                  |
| Serum level 25 (OH)D |    |               |                  |
| $\geq 30$ ng/ml      | 11 | (19.3%)       | 22.0 $\pm$ 11.64 |
| $< 30$ ng/ml         | 46 | (80.7%)       |                  |
| Schirmer test        |    |               |                  |
| >10 mm               | 32 | (56.1%)       | 13.13 $\pm$ 8.91 |
| $\leq 10$ mm         | 25 | (43.9%)       |                  |
| TBUT                 |    |               |                  |
| $\geq 10$ seconds    | 22 | (38.6%)       | 9.08 $\pm$ 4.56  |
| <10 seconds          | 35 | (61.4%)       |                  |
| OSDI score           |    |               |                  |
| 0-12                 | 34 | (59.6%)       | 22 $\pm$ 11.64   |
| >13                  | 23 | (40.4%)       |                  |

Based on characteristic data obtained from 57 research subjects, 67.7% were dominated by women with an average age of  $42.0 \pm 8.7$ . Judging from the serum 25(OH)D levels, it was found that 80.7% had vitamin D deficiency. For the DES examination with the Schirmer test, 56.1% were normal, while for the TBUT examination, 61.4% showed disruption of the tear film and for OSDI the score was 59.6. % normal.

Table 2. Analysis of Respondent Characteristics using the Schirmer test and TBUT

| Variable            | Schirmer test |           | p value | TBUT        |             | p value            |
|---------------------|---------------|-----------|---------|-------------|-------------|--------------------|
|                     | >10mm         | ≤10mm     |         | >10 seconds | ≤10 seconds |                    |
| Gender              |               |           |         |             |             |                    |
| Men                 | 5(19.2%)      | 21(80.8%) | 0.174   | 7(26.9%)    | 19(73.1%)   | 0.860 <sup>€</sup> |
| Women               | 11(35.5%)     | 0(64.5%)  |         | 9(29.0%)    | 22(71.0%)   |                    |
| Age (year)          |               |           |         |             |             |                    |
| 25 – 40             | 18(69.2%)     | 8(30.8%)  | 0.783   | 10(41.7%)   | 14(58.3%)   | 0.044 <sup>€</sup> |
| >40                 | 12(38.7%)     | 19(61.3%) |         | 12(36.4%)   | 21(63.6%)   |                    |
| Serum level 25(OH)D |               |           |         |             |             |                    |
| ≥30ng/ml            | (27.3%)       | 8(72.7%)  | 0.948   | 6(54.5%)    | 5(45.5%)    | 0.030 <sup>€</sup> |
| <30ng/ml            | 13(28.3%)     | 33(71.7%) |         | 36(78.3%)   |             |                    |

€ = Chi-square test(p<0.05)

Based on Table 2 above, it can be concluded that there is no significant relationship between gender (p=0.174), age (p=0.783) vitamin D levels (p=0.948), and the Schirmer test. However, the TBUT examination showed a significant relationship with age (p=0.044) and vitamin D levels (p=0.030), whereas the relationship between gender and the TBUT examination was not found to be significant (p=0.860).

Table 3 Bivariate Analysis of Respondent Characteristics with OSDI scores

| Variable            | OSDI Score |           | p value <sup>€</sup> |
|---------------------|------------|-----------|----------------------|
|                     | 0 – 12     | > 13      |                      |
| Gender              |            |           |                      |
| Men                 | 15(57.7%)  | 11(42.3%) | 0.783                |
| Women               | 19(61.3%)  | 12(38.7%) |                      |
| Age (year)          |            |           |                      |
| 25 – 40             | 15(62.5%)  | 9(37.5%)  | 0.708                |
| >40                 | 19(57.6%)  | 14(42.4%) |                      |
| Serum level 25(OH)D |            |           |                      |
| ≥30ng/dl            | 5(45.5%)   | 6(54.5%)  | 0.285                |
| <30ng/dl            | 29(63.0%)  | 17(37.0%) |                      |

€ = Chi-square test (p<0.05)

Based on Table 3 above, it can be concluded that there is no significant relationship between gender (p=0.783), age (p=0.708), and vitamin D levels (p=0.285) with the OSDI questionnaire.

## DISCUSSION

This study aimed to assess the correlation between CO levels and the degree of smoking and lung function in online motorcycle taxi drivers. The median CO level in this study was 7 ppm with the highest value being 36 ppm almost half of the respondents (47.5%) were in the green zone and the remaining 37.3% were in the red zone. Respondents who smoked in this study amounted

to 75.7% with a median Brinkman index of 80 and a maximum value of 936. In this study, the average CO level was found to be 12.16 ppm in respondents who smoked, while in those who did not smoke it was 2.76 ppm. Smokers' exhaled CO levels in this study were higher than in similar studies, such as Maga et al (smokers 8.25 ppm vs non-smokers 3.26 pm), but lower than Warsaw et al (smokers 14.4 ppm vs non-smokers 5.1 ppm).<sup>3</sup> In many studies CO levels Expiration in smokers is significantly higher than in non-smokers, but the cut-off point between the two is not yet clear.<sup>7,19</sup> Research in Korea by Kim et al concluded that a cut-point value of 5 ppm has a sensitivity of 80-98.3% with a specificity of 100% in smoking cessation programs for stated that the respondent had stopped completely. If a smoker stops smoking for a long period, his exhaled CO levels can decrease to the same as those of a non-smoker.<sup>19</sup> Research by Cropsey et al even created a cut point of 3 ppm to differentiate smokers from non-smokers, which is considered accurate for all races and does not differentiate between genders. 20 Differences in exhaled CO levels, especially in smokers, can be influenced by many things, such as age, distribution of respondents who are heavy smokers, intensity of exposure to cigarette smoke, depth of inhalation, type of cigarette, last time smoked, or the influence of other factors such as air and environmental pollution.<sup>19, 21</sup> In addition to the cut point for smokers and nonsmokers, some researchers recommend a limit of 10 ppm as a threshold for dangerous CO levels.<sup>19</sup>

The pulmonary function test in this study uses CV and VEP1 parameters to assess whether there is restriction or obstruction. Of the respondents, the results showed that more than half (51.5%) of the respondents experienced restrictive disorders with mild restrictions at 42.4% and 78.8% of respondents experienced obstructive disorders. The results of the spirometry examination were normally distributed with the mean CV and VEP1 between smokers and non-smokers not significantly different, the CV values (79.35% vs 78% with  $p = 0.732$ ) and VEP1 (55.31% VC 66.96% with  $p = 0.630$ ). Similar research results were obtained by Saiphoklang et al who did not obtain a significant difference in VEP1 values between smokers and non-smokers ( $p=0.444$ ).<sup>21</sup>

In the correlation test, it was found that CO levels were significantly positively correlated with the degree of smoking, whether assessed from the Brinkman index or the FTND questionnaire. Expired CO levels have been widely used to assess indicators of the severity of a person's smoking habit.<sup>22</sup> Many studies have been carried out to assess the relationship between exhaled CO levels and smoking habits, the severity of smoking is related to higher expiratory CO levels.<sup>23</sup>

Cigarettes contain thousands of chemicals that are dangerous to humans. Free radicals, nicotine, and CO are believed to have the most dangerous effects. Cigarettes increase CO levels in the blood which will cause various cardiovascular diseases because CO replaces oxygen in the blood to form carboxyhemoglobin /COHb bonds. Measurement of expiratory CO and CoHb levels is a non-invasive and objective measurement to assess the degree of smoking and has been widely used in various studies.<sup>23</sup>

The results of this study did not show a correlation between lung function and exhaled CO levels or the degree of smoking. In experimental tests, there was a significant difference in lung function (KVP and VEP1) before and after smoking shisha ( $p<0.05$ ). Measurements of lung function and expiratory CO were carried out after the respondents were given treatment to smoke shisha, there was a significant increase in CO and a decrease in lung function before and

after the intervention.<sup>9</sup> Soeroso et al, research also found a weak correlation between CO and peak expiratory flow, the higher the CO levels. Expiration will reduce peak expiratory flow ( $r = -0.26$  and  $p = 0.106$ ).<sup>23</sup> Similar results to researchers obtained by Salepci et al, expiratory CO levels were also not found to have a significant correlation with force expiratory flow/ FEF 25-75% ( $r = -0.05$ ,  $p = 0.527$ ).<sup>18</sup> This lack of correlation could be caused by several factors including no data on how many cigarettes were smoked in the last 12 hours and how long the respondent had not smoked before the examination.

## CONCLUSION

This research found that the heavier the degree of smoking and the higher the nicotine dependence, the higher the exhaled CO levels. There was no correlation between KVP and VEP1 with exhaled CO levels and degree of smoking.

## ACKNOWLEDGEMENT

None

## AUTHORS CONTRIBUTION

RK: Concept, research design, analysis, interpretation of results, and preparation of published articles; RK, MA, DAWS: Data collection, reviewed the article and approved publication.

## FUNDING

All sources of funds used in research and access to this journal use internal grants from Universitas Trisakti.

## CONFLICT OF INTEREST

Competing interests: No relevant disclosures

## REFERENCES

1. Ismiyati I, Marlita D, Saidah D. Pencemaran Udara Akibat Emisi Gas Buang Kendaraan Bermotor. *J Manaj Transp Logistik*. 2014;1(3):241. DOI: 10.54324/j.mtl.v1i3.23
2. Valavanidis A. Air Pollution as a Major Risk Factor of Global Burden of Disease Study. Global estimates for indoor-outdoor particulate matter, ozone nitrogen oxides and sulphur dioxide. *Sci Rev*. 2021:1–34.
3. Gregorczyk-Maga I, Maga M, Wachsmann A, Janik MK, Chrzastek-Janik O, Bajkowski M, et al. Air pollution may affect the assessment of smoking habits by exhaled carbon monoxide measurements. *Environ Res*. 2019;172:258–65. DOI:10.1016/j.envres.2019.01.063
4. Lawin H, Ayi Fanou L, Hinson V, Wanjiku J, Ukwaja NK, Gordon SB, et al. Exhaled carbon monoxide: a non-invasive biomarker of short-term exposure to outdoor air pollution. *BMC Public Health*. 2017;17(1):4–8. DOI: 10.1186/s12889-017-4243-6
5. Balitbang Kemenkes RI. Riset Kesehatan Dasar. RISKESDAS 2013. Available from: [https://repository.badankebijakan.kemkes.go.id/id/eprint/4467/1/Laporan\\_riskesdas\\_2013\\_final.pdf](https://repository.badankebijakan.kemkes.go.id/id/eprint/4467/1/Laporan_riskesdas_2013_final.pdf)
6. Vasthare R, Kumar S, Arron L. Carbon monoxide breath analyzers and its role in tobacco cessation: A narrative review of literature. *J Int Oral Heal*. 2018;10(2):71–6.
7. Pan KT, Leonardi GS, Ucci M, et al. Can exhaled carbon monoxide be used as a marker of exposure? A cross-sectional study in young adults. *Int J Environ Res Public Health*. 2021;18(22). 11893. DOI:10.3390/ijerph.182211893

8. Ghorbani R, Blomberg A, Schmidt FM. Impact of breath sampling on exhaled carbon monoxide. *J Breath Res.* 2020;14(4). 047105
9. Wiratmoko MR, Adhalia CK. Gambaran Kadar Kabon Monoksida Udara Ekspirasi pada Pengguna Shisha dan Faktor yang Mempengaruhi. *J Respir Indones.* 2019;39(1):37–43.
10. Ryter SW, Choi AMK. Carbon monoxide in exhaled breath testing and therapeutics. *J Breath Res.* 2013;7(1):1–23. DOI: 10.1088/1752-7155/7/1/017111
11. Sandberg A, Sköld CM, Grunewald J, et al. Assessing recent smoking status by measuring exhaled carbon monoxide levels. *PLoS One.* 2011;6(12). DOI: 10.1371/journal.pone.0028864
12. Moscato U, Poscia A, Gargaruti R, et al. Normal values of exhaled carbon monoxide in healthy subjects: Comparison between two methods of assessment. *BMC Pulm Med.* 2014;14(1):1–7.
13. Chen Z, Yu X, Lv D, et al. Do nicotine dependence influencing and non-influencing behaviors have an association with high nicotine dependence in smokers? *Tob Induc Dis.* 2021;19:1–6. DOI: 18332/tid/142866
14. Nurhidayati IR, Syafnita E. Hubungan antara Ketergantungan Nikotin berdasarkan Fagerstrom Test for Nicotine Dependence (FTND )dengan Gangguan Fungsi Kognitif yang Dinilai dengan MoCA-Ina. *Maj Sainstekes.* 2019;5(1):53–7.
15. Põld M, Pärna K. Nicotine dependence and factors related to smoking cessation among physicians in Estonia. *Int J Environ Res Public Health.* 2020;17(9). DOI: 10.3390/ijerph17093217
16. Yalcin FK, Er M, Hasanoglu HC, et al. Deteriorations of pulmonary function, elevated carbon monoxide levels and increased oxidative stress amongst water-pipe smokers. *Int J Occup Med Environ Health.* 2017;30(5):731–42. DOI:10.13075/ijom.1896.00912
17. Ejazi MA, Shameem M Bhargava R, Ahmad Z Akhtar J, et al. . Correlation of exhaled carbon monoxide level with disease severity in chronic obstruction pulmonary disease. *Lung India.* 2018;35(5):401-6. DOI:10.4103/lungindia.lungindia\_11\_18.
18. Salepci BM, Fida A, TorunDoes the Finding of Small Airway Obstruction in Pulmonary Function Tests Effect End-Expiratory Carbonmonoxide Level? *Eurasian J Pulmonol.* 2015;17(3):154–8. DOI:105152/ejpp.2015.76476
19. Kim SS, Kim S, Gona PN. Determining Optimal Cutoffs for Exhaled Carbon Monoxide and Salivary Cotinine to Identify Smokers among Korean Americans in a Smoking Cessation Clinical Trial. *J Smok Cessat.* 2021;15:6678237 DOI:10.1155/2021/6678237
20. Cropsey KL, Trent LR, Clark CB, et al. How low should you go? Determining the optimal cutoff for exhaled carbon monoxide to confirm smoking abstinence when using cotinine as reference. *Nicotine Tob Res.* 2014;16(10):1348–55. DOI:10.1093/ntr/ntu085
21. Saiphoklang N, Poachanukoon O, Soorapan S. Smoking characteristics and lung functions among university athletes. *Sci Rep* 2020;10(1):1–6. DOI:10.1038/s41598-020-77248-y
22. Herath P, Wimalasekera SW, Amarasekara TD, et al. Adverse effects of cigarette smoking on exhaled breath carbon monoxide, blood carboxyhemoglobin, and hematological parameters amongst Sri Lankan adult tobacco smokers: A descriptive study. *Popul Med.* 2021;3(Cvd):1–10. DOI:10.18332/popmed/143076
23. Soeroso N, Intan TK, Ichwan M, et al. The Relationship between Exhaled Carbon Monoxide Test and Peak Expiratory Flow Rate in Smokers and Non-smokers. *ICOSTEEERR* 2018:635–8. DOI:10.5220/0010081306350638



This work is licensed under a Creative Commons Attribution Non-Commercial 4.0 International License

JURNAL BIOMEDIKA DAN KESEHATAN  
(JOURNAL OF BIOMEDIKA AND HEALTH)  
Vol. 6 No. 3 (2023) pp.283-290 e-ISSN:  
2621-5470 283 ORIGINAL ARTICLE  
Pulmonary Function Test and its  
Correlation with Exhaled Carbon  
Monoxide and Smoking Habits in Ojek  
Drivers

*By Rita Khairani*

## ORIGINAL ARTICLE

# Pulmonary Function Test and its Correlation with Exhaled Carbon Monoxide and Smoking Habits in Ojek Drivers

Korelasi Karbon Monoksida dan Derajat Rokok dengan Fungsi Paru pada Pengemudi Ojek Online

Rita Khairani<sup>1</sup>, Mustika Anggiane Putri<sup>2</sup>, Dyah Ayu Woro Setyaningrum<sup>3</sup>

<sup>1</sup>Department of Internal Medicine, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

<sup>2</sup>Department of Physiology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

<sup>3</sup>Department of Anatomical Pathology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

✉ rita.khairani@trisakti.ac.id

 <https://doi.org/10.18051/jbk.283-290>

## ABSTRACT

### Background

Cigarette smoke is the main source of exposure to carbon monoxide (CO) besides air pollution. When cigarette smoke is inhaled, carbon monoxide is absorbed through the lungs, enters the bloodstream, and then binds to hemoglobin to form carboxyhemoglobin (COHb), whose levels in the blood can be measured as a marker of cigarette smoke absorption. Both smoking habits and CO levels can affect lung function, after smoking, blood CO levels will increase and lung function will decrease significantly.

### Methods

This study used an analytic observational method with a cross-sectional design with a total sample of 99 respondents. Data collection in September-December 2022 using the Brinkman Index questionnaire, Fagerstrom Test for Nicotine Dependence (FTND) questionnaire, exhaled CO levels, and spirometry examination.

### Results

47.5% exhaled CO levels on 0-6 ppm or in a green zone, 77.8% of subjects smoked, 57.6% FTND score was 0-2, 51.5% of subjects had restricted lung function, and 79.5% subjects with obstruction. Exhaled CO levels were strongly correlated with the Brinkman index ( $r=0.654$ ,  $p=0.000$ ), and moderately significant correlation with the FTND score ( $r=0.544$ ,  $p=0.000$ ). There is no correlation between exhaled CO levels and degree of smoking with lung function.

### Conclusions

Exhaled CO levels have a significant correlation with smoking and nicotine dependence but do not correlate with pulmonary function test.

**Keywords:** Exhaled carbon monoxide level, smoking, pulmonary function test, nicotine dependence

## ABSTRAK

### Latar Belakang

Asap rokok merupakan sumber utama pajanan terhadap karbon monoksida (CO) selain polusi udara. Saat asap rokok terinhalasi, karbon monoksida akan diabsorpsi melalui paru, masuk ke dalam aliran darah kemudian akan berikatan dengan hemoglobin untuk membentuk karboksi – hemoglobin (COHb) yang kadarnya dalam darah dapat diukur sebagai marker absorpsi asap rokok. Baik rokok atau CO dapat mempengaruhi fungsi paru, setelah merokok maka kadar CO dalam darah akan meningkat dan fungsi paru akan menurun secara bermakna.

### Metode

Penelitian ini menggunakan metode observasional analitik dengan desain potong lintang dengan besar sampel 99 responden. Pengambilan data pada bulan September-Desember 2022 dengan menggunakan kuesioner Indeks Brinkman, *Fagerstrom Test for Nicotine Dependence* (FTND), pengukuran CO ekspirasi dan pemeriksaan spirometri

### Hasil

Sebanyak 47.5% responden memiliki kadar CO ekspirasi sebesar 0-6 ppm yang masih berada pada zona hijau, 77.8% responden merokok, 57.6% memiliki skor FTND 0-2, sebanyak 51.5% mengalami gangguan restriksi dan 79.5% responden mengalami gangguan obstruksi saluran napas. Kadar CO ekspirasi berkorelasi erat dengan indeks Brinkman ( $r=0.654$ ,  $p=0.000$ ), dan berkorelasi sedang bermakna dengan skor FTND ( $r=0.544$ ,  $p=0.000$ ). Tidak ada korelasi kadar CO ekspirasi, indeks Brinkman dan ketergantungan terhadap nikotin dengan fungsi paru.

### Kesimpulan

kadar CO ekspirasi berkorelasi erat dengan rokok dan ketergantungan nikotin tetapi tidak berkorelasi dengan fungsi paru.

**Kata Kunci:** karbon monoksida ekspirasi, rokok, uji fungsi paru, ketergantungan nikotin

## INTRODUCTION

The most common source of air pollution in urban areas comes from motorized vehicles. The development of traffic volume in urban areas has reached 15% every year in the last 10 years. Transportation in big cities is the largest source of air pollution, where 70% of air pollution in urban areas is caused by motorized vehicle activities, especially motorbikes, which reach 30%.<sup>1</sup> Parameters of air pollution from motorized vehicles such as carbon monoxide (CO), nitrogen oxides (NOx), methane (CH<sub>4</sub>), sulfur dioxide (SO<sub>2</sub>) and particulate matter can affect global warming.<sup>1,2</sup>

Apart from air pollution, cigarette smoke is also the main source of exposure to carbon monoxide (CO).<sup>3</sup> Examination of exhaled air CO levels can be used as a biomarker for the degree of smoking. Smoking is a habit that is often found throughout the world, although it is generally known that cigarettes can cause health problems.<sup>4</sup> The percentage of smokers aged 15-19 years has increased in 2020, namely 10.61%, an increase from 10.54% in 2019. More than half (52.1%) of smokers in Indonesia smoked for the first time at the age of 15-19 years, according to Basic Health Research data.<sup>5</sup>

Cigarettes cause damage to almost all organs of the body, when cigarette smoke is inhaled, carbon monoxide will be absorbed through the lungs, enter the bloodstream, and then bind with hemoglobin to form carboxy-hemoglobin (COHb) whose levels in the blood can be measured as a marker for cigarette smoke absorption.<sup>6</sup> Carbon monoxide will remain in the blood for 24 hours after inhalation of cigarette smoke depending on several factors such as gender, physical activity, and respiratory rate.<sup>7</sup> In the blood, CO will then re-enter the alveoli because there is a

concentration gradient in the alveoli, so the levels of CO contained in exhaled air can be measured using a portable CO meter.<sup>8</sup>

<sup>14</sup> Endogenous carbon monoxide is produced primarily by systemic heme oxygenation and is eliminated through respiration. In healthy populations who do not smoke, end-tidal concentrations are in the range of 1-3 ppm.<sup>9</sup> Carbon monoxide is one of the most widely distributed pollutants in the air. CO concentrations are released into the air in the highest quantities among other air pollutants every year except for carbon dioxide (CO<sub>2</sub>) concentrations.<sup>3</sup> <sup>8</sup> areas with high populations the ratio of mixed CO levels can reach up to 10 ppm. The most important biological <sup>8</sup> characteristic of CO is its ability to bind to hemoglobin to form carboxyhemoglobin (HbCO) which is 200 times more stable than oxyhemoglobin (HbO<sub>2</sub>) which can have fatal consequences because it can disrupt muscle metabolism and intracellular enzyme function due to the stable CO bond.<sup>4,10</sup>

Measurement of expiratory monoxide levels has been used to increase the validity of investigations of the degree of smoking because carbon monoxide measurements are easy to use, non-invasive, and affordable.<sup>11</sup> This method can provide a direct assessment of the degree of smoking and is a potential biomarker for assessing air pollution exposure.<sup>3,4</sup> Environmental exposures other than cigarette smoke are important factors that can differentiate exhaled CO results in passive smokers and non-smokers.<sup>11,12</sup>

Subjectively, a person's level of addiction to cigarettes is <sup>25</sup> assessed using the Fagerstrom Test for Nicotine Dependence questionnaire, a questionnaire that was introduced in 1978.<sup>13,14</sup> This questionnaire contains a series of questions regarding how much a person <sup>10</sup> cannot give up smoking. Objectively, the degree of smoking can be assessed by examining exhaled <sup>12</sup> carbon monoxide levels between smokers and non-smokers and there is a strong correlation between exhaled carbon monoxide and the severity of smoking.<sup>13,15</sup>

Both cigarettes and CO can affect lung function. In research conducted by Yalcin et al., who conducted experimental research, after smoking, CO levels will increase and lung function will decrease significantly.<sup>16</sup> Ejazi et al.'s <sup>5</sup> research also found that expiratory CO levels were significantly negatively correlated with the first-second forced expiratory volume (VEP1) in patients. chronic obstructive pulmonary disease (COPD).<sup>17</sup> Different research results were obtained by Salepci et al because <sup>16</sup> there was no relationship between lung function and exhaled CO levels.<sup>12</sup> This study aims to assess the relationship between the degree of smoking and exhaled carbon monoxide levels with lung function in online motorcycle taxi drivers.<sup>18</sup>

## METHODS

<sup>17</sup> This research uses an analytical observational method with a cross-sectional design. Data was collected at the Faculty of Medicine Campus, Universitas Trisakti in September-December 2022. <sup>26</sup> Respondents were selected using consecutive non-random sampling with the required sample size of 94 respondents and rounded up to 100 respondents.

<sup>22</sup> The research population was online motorcycle taxi drivers who met the inclusion criteria (online motorcycle taxi drivers in DKI Jakarta and surrounding areas, male and female, aged 17 years – 60 years, negative antigen swab results, willing to be respondents, and able to carry out spirometry and expiratory CO procedures well). Respondents were excluded if they had a history

of chronic lung disease and the spirometry examination was unacceptable or had no reproducibility). Respondents were then interviewed, filled out questionnaires, spirometry, and exhaled CO examinations.

Interview data includes identity and sociodemographic data, and filling out a questionnaire about smoking habits using the Brinkman index by calculating the number of cigarettes smoked per day and the number of years of smoking. The degree of nicotine dependence uses the Fagerstrom Test for Nicotine Dependence (FTND), with categories light (0-2), moderate (3-4), and severe (5-7). CO levels are checked using the piCO Smokerlyzer device, by holding the respondent's breath for 15 seconds, then exhaling it slowly until it runs out and the results will be visible on the measuring instrument. Light levels/green zone (0-6 ppm), medium/yellow zone (7-10 ppm) and heavy/red zone ( $\geq 11$ ). Lung function examination used a Minato AS-507 spirometer, measuring vital capacity (CV) and first-second expiratory volume (VEP1) which was carried out three times, and the highest value was taken and compared with the predicted value. Lung function test results are then categorized as normal if CV and VEP1 are  $\sim 80\%$ , mild restriction (CV 60-79.9%), moderate restriction (CV 30-59.8%), and severe restriction (CV  $< 30\%$ ), while mild obstruction (VEP1 60% -79.9%), moderate obstruction (VEP1 30-59.9%) and severe obstruction (VEP1  $< 30\%$ ). Statistical test analysis uses the Spearman correlation test using SPSS version 25. This research has passed ethical review by the ethics commission of the Faculty of Medicine, Universitas Trisakti, with number 166/KER/FK/VIII/2022.

## RESULTS

Univariate analysis was used to determine the frequency distribution of subject characteristics in the form of gender, age, serum 25(OH)D levels, Schirmer test, TBUT, and OSDI scores.

Table 1. Data on subject characteristics (N=57)

| Variable             |    | Frequency (%) | Mean $\pm$ SD    |
|----------------------|----|---------------|------------------|
| Gender               |    |               |                  |
| Men                  | 26 | (41.3%)       |                  |
| Women                | 31 | (67.7%)       |                  |
| Age (year)           |    |               |                  |
| 25 – 40              | 24 | (42.1%)       | 42.0 $\pm$ 8.7   |
| > 40                 | 33 | (57.9%)       |                  |
| Serum level 25 (OH)D |    |               |                  |
| $\geq 30$ ng/ml      | 11 | (19.3%)       | 22.0 $\pm$ 11.64 |
| $< 30$ ng/ml         | 46 | (80.7%)       |                  |
| Schirmer test        |    |               |                  |
| >10 mm               | 32 | (56.1%)       | 13.13 $\pm$ 8.91 |
| $\leq 10$ mm         | 25 | (43.9%)       |                  |
| TBUT                 |    |               |                  |
| $\geq 10$ seconds    | 22 | (38.6%)       | 9.08 $\pm$ 4.56  |
| $< 10$ seconds       | 35 | (61.4%)       |                  |
| OSDI score           |    |               |                  |
| 0-12                 | 34 | (59.6%)       | 22 $\pm$ 11.64   |
| >13                  | 23 | (40.4%)       |                  |

Based on characteristic data obtained from 57 research subjects, 67.7% were dominated by women with an average age of  $42.0 \pm 8.7$ . Judging from the serum 25(OH)D levels, it was found that 80.7% had vitamin D deficiency. For the DES examination with the Schirmer test, 56.1% were normal, while for the TBUT examination, 61.4% showed disruption of the tear film and for OSDI the score was 59.6. % normal.

Table 2. Analysis of Respondent Characteristics using the Schirmer test and TBUT

| Variable            | Schirmer test |           | p value | TBUT        |             | p value            |
|---------------------|---------------|-----------|---------|-------------|-------------|--------------------|
|                     | >10mm         | ≤10mm     |         | >10 seconds | ≤10 seconds |                    |
| Gender              |               |           |         |             |             |                    |
| Men                 | 5(19.2%)      | 21(80.8%) | 0.174   | 7(26.9%)    | 19(73.1%)   | 0.860 <sup>€</sup> |
| Women               | 11(35.5%)     | 0(64.5%)  |         | 9(29.0%)    | 22(71.0%)   |                    |
| Age (year)          |               |           |         |             |             |                    |
| 25 – 40             | 18(69.2%)     | 8(30.8%)  | 0.783   | 10(41.7%)   | 14(58.3%)   | 0.044 <sup>€</sup> |
| >40                 | 12(38.7%)     | 19(61.3%) |         | 12(36.4%)   | 21(63.6%)   |                    |
| Serum level 25(OH)D |               |           |         |             |             |                    |
| ≥30ng/ml            | (27.3%)       | 8(72.7%)  | 0.948   | 6(54.5%)    | 5(45.5%)    | 0.030 <sup>€</sup> |
| <30ng/ml            | 13(28.3%)     | 33(71.7%) |         |             | 36(78.3%)   |                    |

€ = Chi-square test(p<0.05)

Based on Table 2 above, it can be concluded that there is no significant relationship between gender (p=0.174), age (p=0.783) vitamin D levels (p=0.948), and the Schirmer test. However, the TBUT examination showed a significant relationship with age (p=0.044) and vitamin D levels (p=0.030), whereas the relationship between gender and the TBUT examination was not found to be significant (p=0.860).

Table 3 Bivariate Analysis of Respondent Characteristics with OSDI scores

| Variable            | OSDI Score |           | p value <sup>€</sup> |
|---------------------|------------|-----------|----------------------|
|                     | 0 – 12     | > 13      |                      |
| Gender              |            |           |                      |
| Men                 | 15(57.7%)  | 11(42.3%) | 0.783                |
| Women               | 19(61.3%)  | 12(38.7%) |                      |
| Age (year)          |            |           |                      |
| 25 – 40             | 15(62.5%)  | 9(37.5%)  | 0.708                |
| >40                 | 19(57.6%)  | 14(42.4%) |                      |
| Serum level 25(OH)D |            |           |                      |
| ≥30ng/dl            | 5(45.5%)   | 6(54.5%)  | 0.285                |
| <30ng/dl            | 29(63.0%)  | 17(37.0%) |                      |

€ = Chi-square test (p<0.05)

Based on Table 3 above, it can be concluded that there is no significant relationship between gender (p=0.783), age (p=0.708), and vitamin D levels (p=0.285) with the OSDI questionnaire.

## DISCUSSION

This study aimed to assess the correlation between CO levels and the degree of smoking and lung function in online motorcycle taxi drivers. The median CO level in this study was 7 ppm with the highest value being 36 ppm almost half of the respondents (47.5%) were in the green zone and the remaining 37.3% were in the red zone. Respondents who smoked in this study amounted

to 75.7% with a median Brinkman index of 80 and a maximum value of 936. In this study, the average CO level was found to be 12.16 ppm in respondents who smoked, while in those who did not smoke it was 2.76 ppm. Smokers' exhaled CO levels in this study were higher than in similar studies, such as Maga et al (smokers 8.25 ppm vs non-smokers 3.26 ppm), but lower than Warsaw et al (smokers 14.4 ppm vs non-smokers 5.1 ppm).<sup>23</sup> In many studies CO levels Expiration in smokers is significantly higher than in non-smokers, but the cut-off point between the two is not yet clear.<sup>7,19</sup> Research in Korea by Kim et al concluded that a cut-point value of 5 ppm has a sensitivity of 80-98.3% with a specificity of 100% in smoking cessation programs for stated that the respondent had stopped completely. If a smoker stops smoking for a long period, his exhaled CO levels can decrease to the same as those of a non-smoker.<sup>10</sup> Research by Cropsey et al even created a cut point of 3 ppm to differentiate smokers from non-smokers, which is considered accurate for all races and does not differentiate between genders.<sup>20</sup> Differences in exhaled CO levels, especially in smokers, can be influenced by many things, such as age, distribution of respondents who are heavy smokers, intensity of exposure to cigarette smoke, depth of inhalation, type of cigarette, last time smoked, or the influence of other factors such as air and environmental pollution.<sup>19, 21</sup> In addition to the cut point for smokers and nonsmokers, some researchers recommend a limit of 10 ppm as a threshold for dangerous CO levels.<sup>19</sup>

The pulmonary function test in this study uses CV and VEP1 parameters to assess whether there is restriction or obstruction. Of the respondents, the results showed that more than half (51.5%) of the respondents experienced restrictive disorders with mild restrictions at 42.4% and 78.8% of respondents experienced obstructive disorders. The results of the spirometry examination were normally distributed with the mean CV and VEP1 between smokers and non-smokers not significantly different, the CV values (79.35% vs 78% with  $p = 0.732$ ) and VEP1 (55.31% VC 66.96% with  $p = 0.630$ ). Similar research results were obtained by Saiphoklang et al who did not obtain a significant difference in VEP1 values between smokers and non-smokers ( $p=0.444$ ).<sup>21</sup>

In the correlation test, it was found that CO levels were significantly positively correlated with the degree of smoking, whether assessed from the Brinkman index or the FTND questionnaire. Expired CO levels have been widely used to assess indicators of the severity of a person's smoking habit.<sup>22</sup> Many studies have been carried out to assess the relationship between exhaled CO levels and smoking habits, the severity of smoking is related to higher expiratory CO levels.<sup>23</sup>

Cigarettes contain thousands of chemicals that are dangerous to humans. Free radicals, nicotine, and CO are believed to have the most dangerous effects. Cigarettes increase CO levels in the blood which will cause various cardiovascular diseases because CO replaces oxygen in the blood to form carboxyhemoglobin /COHb bonds. Measurement of expiratory CO and COHb levels is a non-invasive and objective measurement to assess the degree of smoking and has been widely used in various studies.<sup>23</sup>

The results of this study did not show a correlation between lung function and exhaled CO levels or the degree of smoking. In experimental tests, there was a significant difference in lung function (KVP and VEP1) before and after smoking shisha ( $p<0.05$ ). Measurements of lung function and expiratory CO were carried out after the respondents were given treatment to smoke shisha, there was a significant increase in CO and a decrease in lung function before and

after the intervention.<sup>9</sup> Soeroso et al, research also found a weak correlation between CO and peak expiratory flow, the higher the CO levels. Expiration will reduce peak expiratory flow ( $r = -0.26$  and  $p = 0.106$ ).<sup>23</sup> Similar results to researchers obtained by Salepci et al, expiratory CO levels were also not found to have a significant correlation with force expiratory flow/ FEF 25-75% ( $r = -0.05$ ,  $p = 0.527$ ).<sup>18</sup> This lack of correlation could be caused by several factors including no data on how many cigarettes were smoked in the last 12 hours and how long the respondent had not smoked before the examination.

## CONCLUSION

This research found that the heavier the degree of smoking and the higher the nicotine dependence, the higher the exhaled CO levels. There was no correlation between KVP and VEP1 with exhaled CO levels and degree of smoking.

## ACKNOWLEDGEMENT

None

## AUTHORS CONTRIBUTION

RK: Concept, research design, analysis, interpretation of results, and preparation of published articles; RK, MA, DAWs: Data collection, reviewed the article and approved publication.

## FUNDING

All sources of funds used in research and access to this journal use internal grants from Universitas Trisakti.

## CONFLICT OF INTEREST

Competing interests: No relevant disclosures

## REFERENCES

1. Ismiyati I, Marlita D, Saidah D. Pencemaran Udara Akibat Emisi Gas Buang Kendaraan Bermotor. *J Manaj Transp Logistik*. 2014;1(3):241. DOI: 10.54324/j.mtl.v1i3.23
2. Valavanidis A. Air Pollution as a Major Risk Factor of Global Burden of Disease Study. Global estimates for indoor-outdoor particulate matter, ozone nitrogen oxides and sulphur dioxide. *Sci Rev*. 2021;1:34.
3. Gregorczyk-Maga I, Maga M, Wachsmann A, Janik MK, Chrzastek-Janik O, Bajkowski M, et al. Air pollution may affect the assessment of smoking habits by exhaled carbon monoxide measurements. *Environ Res*. 2019;172:258–65. DOI :10.1016/j.envres.2019.01.063
4. Lawin H, Ayi Fanou L, Hinson V, Wanjiku J, Ukwaja NK, Gordon SB, et al. Exhaled carbon monoxide: a non-invasive biomarker of short-term exposure to outdoor air pollution. *BMC Public Health*. 2017;17(1):4–8. DOI: 10.1186/s12889-017-4243-6
5. Balitbang Kemenkes RI. Riset Kesehatan Dasar. RISKESDAS 2013. Available from: [https://repository.badankebijakan.kemkes.go.id/id/eprint/4467/1/Laporan\\_riskesdas\\_2013\\_final.pdf](https://repository.badankebijakan.kemkes.go.id/id/eprint/4467/1/Laporan_riskesdas_2013_final.pdf)
6. Vasthare R, Kumar S, Arron L. Carbon monoxide breath analyzers and its role in tobacco cessation: A narrative review of literature. *J Int Oral Heal*. 2018;10(2):71–6.
7. Pan KT, Leonardi GS, Ucci M, et al. Can exhaled carbon monoxide be used as a marker of exposure? A cross-sectional study in young adults. *Int J Environ Res Public Health*. 2021;18(22). 11893. DOI:10.3390/ijerph.182211893

8. Ghorbani R, Blomberg A, Schmidt FM. Impact of breath sampling on exhaled carbon monoxide. *J Breath Res.* 2020;14(4). 047105
9. Wiratmoko MR, Adhalia CK. Gambaran Kadar Kabon Monoksida Udara Ekspirasi pada Pengguna Shisha dan Faktor yang Mempengaruhi. *J Respir Indones.* 2019;39(1):37–43.
10. Ryter SW, Choi AMK. Carbon monoxide in exhaled breath testing and therapeutics. *J Breath Res.* 2013;7(1):1–23. DOI: 10.1088/1752-7155/7/1/017111
11. Sandberg A, Sköld CM, Grunewald J, et al. Assessing recent smoking status by measuring exhaled carbon monoxide levels. *PLoS One.* 2011;6(12). DOI: 10.1371/journal.pone.0028864
12. Moscato U, Poscia A, Gargaruti R, et al. Normal values of exhaled carbon monoxide in healthy subjects: Comparison between two methods of assessment. *BMC Pulm Med.* 2014;14(1):1–7.
13. Chen Z, Yu X, Lv D, et al. Do nicotine dependence influencing and non-influencing behaviors have an association with high nicotine dependence in smokers? *Tob Induc Dis.* 2021;19:1–6. DOI: 18332/tid/142866
14. Nurhidayati IR, Syafnita E. Hubungan antara Ketergantungan Nikotin berdasarkan Fagerstrom Test for Nicotine Dependence (FTND) dengan Gangguan Fungsi Kognitif yang Dinilai dengan MoCA-Ina. *Maj Sainstekes.* 2019;5(1):53–7.
15. Põld M, Pärna K. Nicotine dependence and factors related to smoking cessation among physicians in Estonia. *Int J Environ Res Public Health.* 2020;17(9). DOI: 10.3390/ijerph17093217
16. Yalcin FK, Er M, Hasanoglu HC, et al. Deteriorations of pulmonary function, elevated carbon monoxide levels and increased oxidative stress amongst water-pipe smokers. *Int J Occup Med Environ Health.* 2017;30(5):731–42. DOI:10.13075/ijomeh.1896.00912
17. Ejazi MA, Shameem M Bhargava R, Ahmad Z Akhtar J, et al. . Correlation of exhaled carbon monoxide level with disease severity in chronic obstruction pulmonary disease. *Lung India.* 2018;35(5):401-6. DOI:10.4103/lungindia.lungindia\_11\_18.
18. Salepci BM, Fida A, TorunDoes the Finding of Small Airway Obstruction in Pulmonary Function Tests Effect End-Expiratory Carbonmonoxide Level? *Eurasian J Pulmonol.* 2015;17(3):154–8. DOI:105152/ejip.2015.76476
19. Kim SS, Kim S, Gona PN. Determining Optimal Cutoffs for Exhaled Carbon Monoxide and Salivary Cotinine to Identify Smokers among Korean Americans in a Smoking Cessation Clinical Trial. *J Smok Cessat.* 2021;15:6678237 DOI:10.1155/2021/6678237
20. Cropsey KL, Trent LR, Clark CB, et al. How low should you go? Determining the optimal cutoff for exhaled carbon monoxide to confirm smoking abstinence when using cotinine as reference. *Nicotine Tob Res.* 2014;16(10):1348–55. DOI:10.1093/ntr/ntu085
21. Saiphoklang N, Poachanukoon O, Soorapan S. Smoking characteristics and lung functions among university athletes. *Sci Rep* 2020;10(1):1–6. DOI:10.1038/s41598-020-77248-y
22. Herath P, Wimalasekera SW, Amarasekara TD, et al. Adverse effects of cigarette smoking on exhaled breath carbon monoxide, blood carboxyhemoglobin, and hematological parameters amongst Sri Lankan adult tobacco smokers: A descriptive study. *Popul Med.* 2021;3(Cvd):1–10. DOI:10.18332/popmed/143076
23. Soeroso N, Intan TK, Ichwan M, et al. The Relationship between Exhaled Carbon Monoxide Test and Peak Expiratory Flow Rate in Smokers and Non-smokers. *ICOSTEEERR* 2018:635–8. DOI:10.5220/0010081306350638



This work is licensed under a Creative Commons Attribution Non-Commercial 4.0 International License

JURNAL BIOMEDIKA DAN KESEHATAN (JOURNAL OF BIOMEDIKA AND HEALTH) Vol. 6 No. 3 (2023) pp.283-290 e-ISSN: 2621-5470 283 ORIGINAL ARTICLE Pulmonary Function Test and its Correlation with Exhaled Carbon Monoxide and Smoking Habits in Ojek Drivers

---

ORIGINALITY REPORT

---

17%

SIMILARITY INDEX

---

PRIMARY SOURCES

---

- |   |                                                                                                                                                                                                    |                |
|---|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| 1 | <a href="https://repositori.usu.ac.id">repositori.usu.ac.id</a><br>Internet                                                                                                                        | 103 words — 3% |
| 2 | <a href="https://jamb.s.poltekkes-mataram.ac.id">jamb.s.poltekkes-mataram.ac.id</a><br>Internet                                                                                                    | 43 words — 1%  |
| 3 | <a href="http://www.populationmedicine.eu">www.populationmedicine.eu</a><br>Internet                                                                                                               | 37 words — 1%  |
| 4 | <a href="http://www.scitepress.org">www.scitepress.org</a><br>Internet                                                                                                                             | 23 words — 1%  |
| 5 | Basar, M.M.. "Female Sexual and Hormonal Status in Patients with Bronchial Asthma: Relationship with Respiratory Function Tests and Psychological and Somatic Status", Urology, 200703<br>Crossref | 21 words — 1%  |
| 6 | <a href="https://jurnal.payungnegeri.ac.id">jurnal.payungnegeri.ac.id</a><br>Internet                                                                                                              | 21 words — 1%  |
| 7 | <a href="http://www.lib.okayama-u.ac.jp">www.lib.okayama-u.ac.jp</a><br>Internet                                                                                                                   | 21 words — 1%  |

|    |                                                                                                                                                                                                                                                                                                                                     |                 |
|----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------|
| 8  | Susanti, Diah, Endah Lutfiana, Haniffudin Nurdiansah, Hariyati Purwaningsih, Lukman Noerochiem, and George Endri Kusuma. "Preparation of CO Gas Sensor from ZnO Material Synthesized via Thermo-Oxidation Process", Modern Applied Science, 2015.<br>Crossref                                                                       | 20 words — 1%   |
| 9  | mail2.isainsmedis.id<br>Internet                                                                                                                                                                                                                                                                                                    | 20 words — 1%   |
| 10 | discovery.ucl.ac.uk<br>Internet                                                                                                                                                                                                                                                                                                     | 19 words — 1%   |
| 11 | Kiki Rizky Mariani, Kartini Kartini. "Derajat merokok berhubungan dengan kadar hemoglobin pada pria usia 30-40 tahun", Jurnal Biomedika dan Kesehatan, 2018<br>Crossref                                                                                                                                                             | 17 words — 1%   |
| 12 | SALEPÇİ, Banu Musaffa, FİDAN, Ali, PARMAKSIZ, Elif Torun, COŞKUN, Esmâ, KIRAL, Nesrin, ÇAĞLAYAN, Benan, CÖMERT, Sevda Şener and DOĞAN, Coşkun. "Does the Finding of Small Airway Obstruction in Pulmonary Function Tests Effect End-Expiratory Carbonmonoxide Level?", Türkiye Solunum Araştırmaları Derneği, 2015.<br>Publications | 17 words — 1%   |
| 13 | apps.who.int<br>Internet                                                                                                                                                                                                                                                                                                            | 16 words — 1%   |
| 14 | www.frontiersin.org<br>Internet                                                                                                                                                                                                                                                                                                     | 13 words — < 1% |
| 15 | midwifery.iocspublisher.org<br>Internet                                                                                                                                                                                                                                                                                             | 12 words — < 1% |
| 16 | www.sciencegate.app<br>Internet                                                                                                                                                                                                                                                                                                     | 12 words — < 1% |

- 
- 17 [repository.trisakti.ac.id](https://repository.trisakti.ac.id) 11 words — < 1%  
Internet
- 
- 18 [www.offshoresource.com](http://www.offshoresource.com) 11 words — < 1%  
Internet
- 
- 19 D Dewanto, I M Muliatna, W Warju. "Increasing community participation in the green technology program through design and application of alternative mufflers", IOP Conference Series: Materials Science and Engineering, 2021 10 words — < 1%  
Crossref
- 
- 20 [coek.info](http://coek.info) 10 words — < 1%  
Internet
- 
- 21 [jurnalrespirologi.org](http://jurnalrespirologi.org) 10 words — < 1%  
Internet
- 
- 22 Dina Lusiana Setyowati, Swandari Paramita, Riza Hayati Ifroh, Tanti Asrianti, Efi Fitriani, Wahnadita Rahman. "Work readiness during COVID-19 among taxibike online drivers in Samarinda, Indonesia", International Journal of Public Health Science (IJPHS), 2021 9 words — < 1%  
Crossref
- 
- 23 J Obermeier, P Trefz, K Wex, B Sabel, J K Schubert, W Miekisch. "Electrochemical sensor system for breath analysis of aldehydes, CO and NO", Journal of Breath Research, 2015 9 words — < 1%  
Crossref
- 
- 24 [ejournal.almaata.ac.id](http://ejournal.almaata.ac.id) 9 words — < 1%  
Internet
- 
- 25 [njcmindia.org](http://njcmindia.org)  
Internet

9 words — < 1%

26 Sari Riastiningsih, Nuryani Sidarta. "Tingkat depresi dan jenis kelamin berhubungan dengan perilaku anti sosial pada pelajar SMA", Jurnal Biomedika dan Kesehatan, 2018

Crossref

8 words — < 1%

27 Vinod K Ramani, Manohar Mhaske, Radheshyam Naik. "Assessment of Carbon Monoxide in Exhaled Breath using the Smokerlyzer Handheld Machine: A Cross-Sectional Study", Tobacco Use Insights, 2023

Crossref

8 words — < 1%

28 [www.dovepress.com](http://www.dovepress.com)

Internet

8 words — < 1%

29 [www.researchgate.net](http://www.researchgate.net)

Internet

8 words — < 1%

30 İlknur Aykurt Karlıbel, Seyhan Dülger, Meliha Kasapoğlu Aksoy, Muhammet Güzelsoy et al. "Effect of cigarette smoking on sexual functions, psychological factors, and disease activity in male patients with ankylosing spondylitis", The Aging Male, 2018

Crossref

6 words — < 1%

EXCLUDE QUOTES ON

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES

EXCLUDE MATCHES

OFF

OFF

☆ **Dr. dr. Husnun Amalia, Sp.M**

Inbox -...trisakti.ac.id

July 3, 2023 at 9:11 PM

SD

[JBK] Submission Acknowledgement

To: Rita Khairani



Siri found new contact info Dr. dr. Husnun Amalia, Sp. M jbiomedkes@trisakti.ac.id

[add...](#) (x)

Rita Khairani:

Thank you for submitting the manuscript, "KORELASI KARBON MONOKSIDA DAN DERAJAT ROKOK DENGAN FUNGSI PARU PADA PENGEMUDI OJEK ONLINE" to Jurnal Biomedika dan Kesehatan. With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal web site:

Submission URL: <https://jbiomedkes.org/index.php/jbk/authorDashboard/submission/396>

Username: rkhairani

If you have any questions, please contact me. Thank you for considering this journal as a venue for your work.

Dr. dr. Husnun Amalia, Sp.M

---

[Jurnal Biomedika dan Kesehatan](#)

Jl. Kyai Tapa No. 260 Kampus B, Grogol, DKI Jakarta 11440

Telp. (021) 5672731 | E-mail: [jbiomedkes@trisakti.ac.id](mailto:jbiomedkes@trisakti.ac.id)

**Husnun Amalia**

Inbox -...trisakti.ac.id

December 25, 2023 at 11:37 AM

HA

[JBK] Editor Decision

To: Rita Khairani

Rita Khairani:

We have reached a decision regarding your submission to Jurnal Biomedika dan Kesehatan, "KORELASI KARBON MONOKSIDA DAN DERAJAT ROKOK DENGAN FUNGSI PARU PADA PENGEMUDI OJEK ONLINE".

Our decision is: Revisions Required

---

[Jurnal Biomedika dan Kesehatan](#)

Jl. Kyai Tapa No. 260 Kampus B, Grogol, DKI Jakarta 11440

Telp. (021) 5672731 | E-mail: [jbiomedkes@trisakti.ac.id](mailto:jbiomedkes@trisakti.ac.id)

**Husnun Amalia**

Inbox -...trisakti.ac.id

December 25, 2023 at 11:38 AM

HA

[JBK] New notification from Jurnal Biomedika dan Kesehatan

To: Rita Khairani,

Reply-To: Dr. dr. Husnun Amalia, Sp.M

---

You have a new notification from Jurnal Biomedika dan Kesehatan:

You have been added to a discussion titled "Revision required" regarding the submission "KORELASI KARBON MONOKSIDA DAN DERAJAT ROKOK DENGAN FUNGSI PARU PADA PENGEMUDI OJEK ONLINE".

Link: <https://jbiomedkes.org/index.php/jbk/authorDashboard/submission/396>

Dr. dr. Husnun Amalia, Sp.M

---

[Jurnal Biomedika dan Kesehatan](#)

Jl. Kyai Tapa No. 260 Kampus B, Grogol, DKI Jakarta 11440

Telp. (021) 5672731 | E-mail: [jbiomedkes@trisakti.ac.id](mailto:jbiomedkes@trisakti.ac.id)