

The background of the cover features a stylized illustration of laboratory glassware. It includes two round-bottom flasks on the left and right, each containing a purple liquid. In the center, there is a piece of equipment that looks like a pipette or a burette. To the right of the central equipment, there are two test tubes, also containing purple liquid. The entire scene is set against a solid purple background.

**Biomedical Journal of
Indonesia**

BJI

**Published by Faculty of Medicine
Universitas Sriwijaya**

Biomedical Journal of Indonesia


Published by Fakultas Kedokteran Universitas Sriwijaya

Online ISSN : 2685-0184 | Print ISSN : 2407-7097

[🏠](#) > Editorial Team

Editorial Team

Editor in Chief

- **Zen Hafy**, Department of Histology, Faculty of Medicine, Universitas Sriwijaya, Palembang, Indonesia  [Scopus'](#)

Editors

- **Noriyuki Koibuchi**, Department of Integrative Physiology, Gunma University, Maebashi, Japan  [Scopus'](#)
- **Ella Amalia**, Department of Microbiology, Faculty of Medicine, Universitas Sriwijaya, Palembang, Indonesia  [Scopus'](#)
- **Veny Larasati**, Department of Histology, Faculty of Medicine, Universitas Sriwijaya, Palembang, Indonesia  [Scopus'](#)
- **Dwi Handayani**, Department of Parasitology, Faculty of Medicine, Universitas Sriwijaya, Palembang, Indonesia  [Scopus'](#)
- **Syarinta Adenina**, Department of Pharmacology, Faculty of Medicine, Universitas Sriwijaya, Palembang, Indonesia  [Scopus'](#)
- **Rizki A. Nawawi**, Department of Clinical Microbiology, Faculty of Medicine, Universitas Sriwijaya, Palembang, Indonesia 
- **Muhammad Ichwan**, Department of Pharmacology and Therapeutic, Faculty of Medicine, Universitas Sumatera Utara, Medan, Indonesia  [Scopus'](#)
- **Radiana Dhewayani Antarianto**, Stem Cell and Tissue Engineering Research Cluster Indonesian Medical Education and Research Institute (IMERI), Universitas Indonesia, Jakarta, Indonesia  [Scopus'](#)
- **Andri Rezano**, Andrology Study Program, Faculty of Medicine, Universitas Padjajaran, Bandung, Indonesia  [Scopus'](#)

E-ISSN



E-ISSN : **2685-0184**

Additional Menu

[Focus and Scope](#)

[Reviewers](#)

[Peer Review](#)

[Copyright Notice](#)

[Publication Fees](#)

[Plagiarism Policy](#)

[Publication Ethics](#)

[Online Submission](#)

[Author Guidelines](#)

Indexing



Journal Template



**Original Article
Template**

DOC

Template



Literature Review
Template

Visitor Statistics

Old Site:



New (Current) Site:



0004364 [View Stats](#)



Reference Manager



Biomedical Journal of Indonesia

— Biomedical Journal of Indonesia is licensed under a Creative Commons Attribution
ShareAlike 4.0 International License.

Biomedical Journal of Indonesia

Published by Fakultas Kedokteran Universitas Sriwijaya

Online ISSN : 2685-0184 | Print ISSN : 2407-7097

🏠 > Archive > Vol. 9 No. 1 (2023): Vol 9, No 1, 2023



Vol. 9 No. 1 (2023): Vol 9, No 1, 2023

Published: 📅 01-02-2023

📄 Articles

Cytotoxic Activity Test Of Ethanolic Extract Of Berenuk Fruit (Crescentia Cujete L.) On Artemia Salina Leach Shrimp Larva Using Brine Shrimp Lethality Test (BSLT) Method

Dona Suzana, Isnani Handayanti

📄 PDF

👁 304

📄 1-6



Relationship Between The Location Of Benign Vocal Cord Lesion And The Degree Of Voice Handicap Index-30

Lisa Apri Yanti, Irfannuddin Irfannuddin, Nilam Kusuma Anggraeni

 PDF

 177

 7-11

The Characteristics of Deviated Nasal Septum Patients and Computed Tomography Findings at Dr. Mohammad Hoesin General Hospital Palembang

Ermalinda Kurnia, Fiona Widayari, Puspa Zuleika, Ahmad Hifni, M Ricky Al Hafiz Sazili

 PDF

 460

 12-18

Correlation Between Hemoglobin Level And Functional Capacity In Young Adult Population

Fiona Rista Utami, Patwa Amani

 PDF

 188

 19-22

Percentage Of Body Fat Related With Functional Capacity In Young Adults

Delia Ainnaya, Mustika Anggiane Putri

 PDF

 322

 23-26

E-ISSN



E-ISSN : 2685-0184

Additional Menu

Focus and Scope

Reviewers

Peer Review



[Copyright Notice](#)

[Publication Fees](#)

[Plagiarism Policy](#)

[Publication Ethics](#)

[Online Submission](#)

[Author Guidelines](#)

Indexing



Journal Template



**Original Article
Template**



**Case Report
Template**



**Literature Review
Template**

Visitor Statistics

Old Site:



New (Current) Site:



0004364 [View Stats](#)





Reference Manager



Biomedical Journal of Indonesia is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-sa/4.0/).





Percentage Of Body Fat Related With Functional Capacity In Young Adults

Delia Ainnaya¹, Mustika Anggiane Putri^{2*}

¹Undergraduate Student of Medical Program Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

²Department of Physiology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

ARTICLE INFO

Keywords:

Young Adults

Bioelectrical impedance analysis (BIA)

Body fat

Functional capacity

VO2max

*Corresponding author:

Mustika Anggiane Putri

E-mail address:

inge.mustika@trisakti.ac.id

All authors have reviewed and approved the final version of the manuscript.

<https://doi.org/10.32539/BJI.v9i1.159>

ABSTRACT

Introduction. The percentage of body fat is a component of body composition in addition to bone mass, muscle mass, and body water content related to obesity. Diseases caused by an increase in body fat will have an impact on decreasing the body's functional capacity, disability, and death. Maximum Oxygen Volume (VO2 max) refers to the maximum amount of oxygen that humans can use during physical activity. It is usually used to test aerobic endurance or cardiovascular fitness so that VO2 max can describe a person's functional capacity. Percentage of body fat can affect VO2 max. Based on data from the Sports Development Index (SDI) in Indonesia, 43.9% of young adults have a poor VO2 max. This study aims to determine whether or not there is a relationship between body fat percentage and functional capacity in young adults. **Methods.** The study used a cross-sectional method and was followed by 92 respondents throughout October and November 2022. Bioelectrical Impedance Analysis (BIA) is used to measure body fat percentage. The 6-minute Walk Test is used to measure VO2 max. **Results.** The results of the study showed that the bivariate analysis of the Fisher's Exact test obtained a significant relationship between body fat percentage (measured by BIA) and body functional capacity (VO2 max) ($p=0.028$). **Conclusion.** There was a significant relationship between body fat percentage through BIA measurements and functional capacity in young adults.

1. Introduction

Functional capacity is a person's ability to take in maximum oxygen also called VO2 max. Maximum Oxygen Volume (VO2 max) refers to the maximum amount of oxygen that humans can use during physical activity. VO2 max is the result of calculating the cardiac output and oxygen in the veins and arteries which describes a person's aerobic endurance to carry out activities.¹ Functional capacity can be measured by measuring VO2 max. Factors that can affect VO2 max include age, gender, lung & heart function, exercise, genetics, abdominal circumference, body fat percentage, and a person's nutritional status (determined by BMI).² In Indonesia, 43.9% of young adults have a poor VO2 max and 37.4% have a very poor VO2 max.³ VO2 max is one of the important factors in cardiovascular health. Age is related to a decrease in VO2 max, one of the reasons is that there is a decrease in the level of physical activity with age so aerobic endurance decreases. Storen et al said that there was a decrease in VO2 max of 1% per year. there will be an increased risk of decreasing VO2 max levels during old age.⁴⁻⁶ The

composition of the human body consists of extracellular fluid, bone, fat, and muscle cells. The combination of these four components at a perfect level will achieve a perfect balance of body composition.⁷ Fat is one of the components of the body that is very influential on health. Obesity is one of the effects of increasing high levels of fat in the body.⁸⁻¹⁰ An increase in body fat will cause disease that can reduce functional capacity, and disability, and even increase the risk of mortality. Body fat needs to be assessed, this is an effort to prevent the occurrence of metabolic syndrome due to obesity.¹¹ Bioelectrical Impedance Analysis (BIA) is a common and widely used method of measuring body fat percentage. Measurements using BIA are often carried out because they are easy and quite accurate.¹²

Based on a study conducted by Schnurr TM et al stated that there was a relationship ($p=0.0034$) between the percentage of body fat and VO2 max.¹³ Meanwhile, Demirkan E et al stated that there was no relationship ($p>0.05$) statistically between the percentage of body fat with VO2 max.¹⁴ Previous

studies have had different conclusions, so researchers are interested in conducting further research to find out whether or not there is a relationship between body fat percentage and functional capacity in young adults.

2. Methods

This research is a cross-sectional study conducted in October - November 2022. This study was followed by 92 respondents aged 18 - 25 years. Respondents who used walking aids had a history of asthma, chronic lung, and heart disease, anemia, and were on long-term glucocorticoid treatment were excluded.

Functional Capacity

The functional capacity of the body is assessed by measuring the respondent's VO₂ max. A 6-minute walking test (6MWT) was performed to assess VO₂ max. The procedure for the 6-minute walk test is as follows: The walking track was a meter wide and 15 meters long, marked by a cone at the end where the subject made a U-turn. The midline was observed every 50 cm to determine the final walking position quickly. The test procedure allows the subject to walk continuously on the track for 6 minutes. Subjects are instructed to stop walking when the time is up and put a mark where the subject stops. These results are then used to calculate the VO₂ max prediction with the following formula:¹⁵

$$\text{VO}_2 \text{ Max} = 0.053 (\text{distance}) + 0.022 (\text{age}) + 0.032 (\text{Body height}) - 0.164 (\text{Body Weight}) - 2.228 (\text{sex}^*) - 2.287$$

Notes: VO₂ max in kg/b.w/min, Distance in meters, Age in years, Body height in cm, Bodyweight in kg, Sex: male = 0 and Female = 1

Interpretation :

1. Poor: <15 ml/ kg/b.w/min
2. Average: 15,1 – 30 ml/ kg/b.w/min
3. Good: 30,1 – 37,5 ml/ kg/b.w/min
4. Excellent : > 37,6 – 45 ml/ kg/b.w/min

Body Fat Percentage

In this study, the percentage of body fat was measured using Bioelectrical Impedance Analysis (BIA): a digital scale (brand: KOVA Body Scale Smart). Subjects were asked to stand on the scales in an upright position to measure total fat.

Interpretation

1.Fair:

Male:11 - 14%

Female: 16 - 19%

2.Normal:

Male: 15 - 18%

Female: 20% - 25%

3.Overweight:

Male: 19% - 24%

Female: 26% - 29%

4.Obese:

Male: 25 - >25%

Female: 30 - >30%

Data analysis used in this study is univariate and bivariate analysis. Univariate analysis was used to analyze the variables studied by explaining the characteristics of each variable. The results of the data are expressed in percentages and presented in tabular form. Bivariate analysis is used to determine the relationship between body fat percentage and functional capacity. The collected data were processed in Microsoft Excel and then analyzed using the Fisher's Exact test through the Statistical Software Package for Social Sciences version 27 (SPSS v.27) program. This research has received ethical approval from the Health Research Ethics Committee, Faculty of Medicine, Universitas Trisakti (No:78//KER-FK/VII/2022).

3. Results

Characteristics of Research Respondents

Based on the results of the univariate analysis provided in Table 1, it was observed that the majority of participants, 57.6% or 53 individuals, were female. Moreover, a significant proportion of the participants, 66.3%, had a fair-normal range of body fat percentage. In terms of functional capacity, it was found that the VO₂max results categorized 89.1% of the participants as being in the poor-average category. Based on the study findings (Table 2), it has been established that the respondents' body fat percentage is 22.16% on average. Likewise, the average VO₂ max level, which is 21.61 ml/kg/b.w/min, has also been determined.

Body Fat Percentage Relationship with Functional Capacity

Based on the data presented in Table 3, it can be concluded that 95.1% of the respondents had a fair to normal body fat percentage and poor-average functional capacities. Meanwhile, the remaining 4.9% of the respondents had good - excellent functional capacities. Furthermore, out of the respondents with overweight - obese body fat percentage, 77.4% had poor - average functional capacity, while 22.6% had good-excellent functional capacity.

The results of the Fisher's Exact test indicate a significant relationship between body fat percentage and functional capacity in young adults, with a probability value of 0.028 ($p=0.028 < 0.05$). This suggests that body fat percentage may have a significant impact on the functional capacity of young adults.

Table 1. Characteristics of research respondents (n = 92)

Characteristics	Frequency (n)	Percentage (%)
Gender		
Male	39	42,4
Female	53	57,6
Body Fat Percentage (BIA)		
Fair - Normal	61	66,3
Overweight - Obese	31	33,7
Functional Capacity (VO2max)		
Poor - Average	82	89,1
Good - Excellent	10	10,9

Table 2. Distribution of body fat percentage and VO2max based on gender

Gender		Body Fat Percentage (BIA) (%)	VO2max (Functional Capacity) (ml/ kg/b.w/min)
Male	Mean	17,325	24,179
	N	39	39
	St. Deviation	4,77597	5,41039
Female	Mean	25,718	19,713
	N	53	53
	St. Deviation	4,578	4,44222
Total	Mean	22,160	21,606
	N	92	92
	St. Deviation	6,236	5,33147

Table 3. Relationship of body Fat Percentage (BIA) with Functional Capacity (VO2max)

Characteristics	Functional capacity (VO ₂ max) (ml/ kg/b.w/min)		Total n(%)	p-value
	Poor - Average	Good - Excellent		
	n(%)	n(%)		
Body Fat Percentage				
Fair - Normal	58 (95,1)	3 (4,9)	61 (66,3)	.028*
Overweight - Obese	24 (77,4)	7 (22,6)	31 (33,7)	

4. Discussion

Based on the findings of this study, it appears that women tend to have higher body fat percentages than men. This is due to the fact that women have increased fat stores in certain areas of the body, such as the upper arms, breasts, and hips. In addition, it was observed that women generally have lower VO₂ max levels compared to men. This is because their total volume of circulating red blood cells tends to decrease, leading to a lower maximum capacity to deliver oxygen to muscles. It should also be noted that age can also affect VO₂ max levels. Starting at age 20, there is a gradual decrease that continues until age 55, where there is a significant drop of 27% compared to age 25.

Moreover, it was discovered that the majority of participants in the study, specifically 66.3% or 61 out of 92, had a fair-normal body fat percentage. Young adults were also observed to fall into this category, most likely due to their balanced energy source from food intake and physical activity. Urban

young adults were found to engage in moderate physical activity, which could contribute to their fair-normal body fat percentage.

The average VO₂ max in young adults is 21.61 ml/kg/min which is included in the average category. Most of the young adults aged 18-25 years are students and workers. they often spend their time sitting in front of gadgets to study and work so they rarely do regular exercise.¹⁶

In this study, 82 people (89.1%) had a poor-average VO₂ max. the average VO₂ max in adolescents is 21.61 ml/kg/min which is included in the average category. Most of the young adults aged 18-25 years are students and workers. they often spend their time sitting in front of gadgets to study and work so they rarely do regular exercise.¹⁶

In this study, it was found that there was a relationship between the percentage of body fat (BIA) and functional capacity with a value of $p = 0.028$ ($p < 0.05$), the higher the percentage of body fat, the lower a person's VO₂ max. VO₂ max is not

only influenced by body fat percentage but also by physical activity which is one of the important factors and greatly affects a person's functional capacity.¹⁷ The results of a previous study conducted by Huldani et al said that there was a significant relationship between the percentage of body fat and VO₂ max with a value of $p=0.013$, where the percentage of body fat in the overweight-obese category had a lower VO₂ max value than people with a normal body. In obese people, it will be more difficult for them to move causing tissue fat to increase. This fatty tissue can ultimately inhibit the functional capacity of the body's cardiorespiratory system.²

5. Conclusion

There was a significant relationship between body fat percentage through BIA measurements and functional capacity in young adults.

6. Acknowledgements

The authors would like to thank all the subjects for their participation. The publication of this manuscript is supported by the Medical Faculty, Universitas Trisakti.

7. References

- Herrera-santelices A, Argüello-florencio G, Westphal G, Junior NN. Effects of Supervised Physical Exercise as Prehabilitation on Body Composition, Functional Capacity and Quality of Life in Bariatric Surgery Candidates: A Systematic Review and. *J Clin Med* 2022;11:1–15.
- Huldani, Asnawati, Auliadina D, Amilia F, NUarti N, Jayanti R. Abdominal Circumference, Body Fat Percent, and VO₂ Max in Pilgrims of Hulu Sungai Tengah Regency. *J Phys Conf Ser* 2019.
- Gantarialdha N. Hubungan Indeks Massa Tubuh Terhadap Ketahanan Kardiorepirasi Dinyatakan Dalam VO₂max. *J Med Utama* 2021;02:1162–8.
- Storen O, Helgerud J, Saebo M, Stoa EM, et al. High-Intensity Interval Training. *Am Coll Sport Med* 2016:78–85.
- Koons NJ, Suresh MR, Schlotman TE, Convertino VA. Interrelationship Between Sex, Age, Blood Volume, and V̇. *Aerosp Med Hum Perform* 2019;90:362–8.
- Myers J, Kaminsky LA, Lima R, Christle JW, Ashley E, Arena R. A Reference Equation for Normal Standards for VO₂ Max: Analysis from the Fitness Registry and the Importance of Exercise National Database (FRIEND Registry). *Prog Cardiovasc Dis* 2017;60.
- Uçar MK, Uçar Z, Köksal F, Daldal N. Estimation of body fat percentage using hybrid machine learning algorithms. *Meas J Int Meas Confed* 2021;167.
- Ribeiro SML, Kehayias JJ. Sarcopenia and the analysis of body composition. *Adv Nutr* 2014;5.
- Pulgarón ER. Childhood Obesity: A Review of Increased Risk for Physical and Psychological Comorbidities. *Clin Ther* 2013;35.
- Chung S, Jeong YG. Effects of the craniocervical flexion and isometric neck exercise compared in patients with chronic neck pain: A randomized controlled trial. *Physiother Theory Pract* 2018;34.
- Danielsen KK, Svendsen M, Mæhlum S, Sundgot-Borgen J. Changes in body composition, cardiovascular disease risk factors, and eating behavior after an intensive lifestyle intervention with high volume of physical activity in severely obese subjects: A prospective clinical controlled trial. *J Obes* 2013;2013.
- Lubis L, Zhe Yiin NN, Luftimas DE. Differentiation of body fat composition between skinfold caliper and bioelectrical impedance analysis methods among professors. *Bali Anat J* 2018;1. <https://doi.org/10.36675/baj.v1i1.6>.
- Schnurr TM, Gjesing AP, Sandholt CH, Jonsson A, Mahendran Y, Have CT, et al. Genetic correlation between body fat percentage and cardiorespiratory fitness suggests common genetic etiology. *PLoS One* 2016;11.
- Demirkan E, Can S, Arslan E. The Relationship between Body Composition and Aerobic Fitness in Boys and Girls Distance Runners. *Int J Sport Sci* 2016;6.
- Ilmul M, Amha Y, Putri MA, Lutfie SH. Effect of School-based Aerobic Exercise Model on Health Fitness Medical Students. *Malaysian J Med Heal Sci* 2022;18:62–6.
- Noordia A. Hubungan Persentase Lemak Tubuh terhadap Kemampuan VO₂max Pada Atlet Gulat Kota Surabaya. *J Kesehat Olahraga Vol* 2015;3.
- Khairani R, Adriani D, and Amani P. Obesity is the most influential risk factor of cardiopulmonary endurance in older women. *Universa Medicina*. 2021; 40(3): 254-262.