

BAGHDAD JOURNAL OF BIOCHEMISTRY AND APPLIED BIOLOGICAL SCIENCES

Volume 7, Issue 2, Year 2026



**Serological Biomarkers
in Forensic Sciences:
Applications and
Challenges**

**Advanced Biochemistry
Applied Biological Sciences
ISSN 2706-9915 (Online)**



Open Access

www.bjbabs.org

Editorial Board

Editor-in-Chief




Omar F. Abdul-Rasheed, Ph.D.

Affiliation: Department of Chemistry and Biochemistry, College of Medicine, Al-Nahrain University, Baghdad, Iraq

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed

Co-Editor-in-Chief




Mohammad Amjad Kamal, Ph.D.

Affiliation: Princess Nourah Bint Abdulrahman University, Riyadh, Saudi Arabia

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed


Associate Editors



Mahmoud Bahmani, Ph.D.

Affiliation: Biotechnology and Medicinal Plants Research Center, Ilam University of Medical Sciences, Ilam, Iran


Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Emad A. Yousif, Ph.D.

Affiliation: Department of Chemistry, College of Science, Al-Nahrain University, Baghdad, Iraq


Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Gamal A. El-Hiti, Ph.D.

Affiliation: Department of Optometry, College of Applied Medical Sciences, King Saud University, Riyadh, Saudi Arabia


Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Rayah S. Baban, Ph.D.

Affiliation: Department of Chemistry and Biochemistry, College of Medicine, Al-Nahrain University, Baghdad, Iraq

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Davoodbasha MubarakAli, Ph.D.

Affiliation: B S Abdur Rahman Crescent Institute of Science and Technology, Chennai, India


Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Jorddy N. Cruz, Ph.D.

Affiliation: Federal University of Amapá, Macapá, Brazil

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Salleh, Wan Mohd Nuzul Hakimi Wan, PhD

Affiliation: Department of Chemistry, Faculty of Science and Mathematics, Universiti Pendidikan Sultan Idris, Tanjung Malim, Malaysia

Email | ORCID | Google Scholar | Web of Science | Scopus | PubMed | ResearchGate



Jalila Ben Salah-ABBES, PhD

Affiliation: High Institute of Biotechnology of Monastir (ISBM), University of Monastir, Tunisia

Email | ORCID | Google Scholar | Web of Science | Scopus | PubMed | ResearchGate



Hayder A. Al-Aubaidy, M.D., Ph.D.

Affiliation: School of Life Science, La Trobe University, Melbourne, Australia

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Hamid Zahednasab, Ph.D.

Affiliation: Institute of Biochemistry and Biophysics, University of Tehran, Tehran, Iran

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Mahmoud H. Hadwan, Ph.D.

Affiliation: Department of Chemistry, College of Science, University of Babylon, Hillah, Iraq

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Muhammed T. Abdel Ghafar, Ph.D.

Affiliation: Department of Clinical Pathology, Faculty of Medicine, Tanta University, Tanta, Egypt

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Zaid M. Al-Obaidi, Ph.D.

Affiliation: Department of Pharmaceuticals and Pharmaceutical Chemistry, College of Pharmacy, University of Alkafel, Karbala, Iraq

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Murad A. AlDamen, Ph.D.

Affiliation: University of Jordan, School of Science, Amman, Jordan


Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Wassan Nori Mohammed, Ph.D.

Affiliation: Department of Obstetrics and Gynecology, College of Medicine, Al Mustansiriyah University, Baghdad, Iraq

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed | ResearchGate



Nazmul Haque, Ph.D.

Affiliation: School of Medicine and Dentistry, University of Central Lancashire, Burnley, United Kingdom

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Nahidh Al-Jaberi, Ph.D.

Affiliation: School of Medicine and Dentistry, University of Central Lancashire, Burnley, United Kingdom

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed | LinkedIn



JEBALI JAMEL, Ph.D.

Affiliation: Professor of Biochemistry and Environmental Sciences, Higher Institute of Biotechnology of Monastir, Monastir, Tunisia

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Jiban Shrestha, Ph.D.

Affiliation: National Plant Breeding and Genetics Research Centre, Khumaltar, Lalitpur, Nepal

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Omer Q. B. Al-Lela, Ph.D.

Affiliation: Department of Pharmacy, Alnoor University, Mosul, Ninevah, Iraq

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Mahara Valverde, Ph.D.

Affiliation: Department of Medicine and Environmental Toxicology, National Autonomous University of Mexico, Ciudad de México, Mexico

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Alireza Heidari, Ph.D.

Affiliation: California South University, Irvine, United States


Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed | Biography



Carlos Augusto C. de Vasconcelos, Ph.D.

Affiliation: Department of Nutrition, Health Sciences Center, Federal University of Pernambuco, Recife, 50670-901, Brazil


Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Khalid Tobal, Ph.D.

Affiliation: Manara University, Iraq


Email | ORCID | Google Scholar | ResearchGate | Scopus | PubMed | LinkedIn



Sam S. Al-Dalali, Ph.D.

Affiliation: Ibb University, Ibb, Yemen


Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Fulgence N. Mpenda, Ph.D.

Affiliation: Department of Molecular Biology and Biotechnology, University of Dar es Salaam, P.O. Box 35179, Dar es Salaam, Tanzania

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Adem Keskin, Ph.D.

Affiliation: Department of Medical Biochemistry, Institute of Health Sciences, Aydin Adnan Menderes University, Aydin, Turkey

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Mohammed Al-Mashhadani, Ph.D.

Affiliation: Department of Chemistry, College of Science, Al-Nahrain University, Baghdad, Iraq

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Mohammed El-Mezgueldi, Ph.D.

Affiliation: King's College London, London, United Kingdom

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Khawla Abdalkarim Kasar, Ph.D.

Affiliation: Department of Pathological Analysis, College of Science, Al-Nahrain University, Baghdad, Iraq


Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed



Hafsan H., Ph.D.

Affiliation: Alauddin State Islamic University of Makassar, Indonesia


Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed | Biography



William Richard Webb, Ph.D.

Affiliation: Research and Innovation Hub, Innovation Aesthetics, London, UK

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed | Biography




Amany A. Ghazy, Ph.D.

Affiliation: Department of Medicine, College of Medicine, Al-Jouf University, Sakakah, Saudi Arabia

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed

Typesetting and Language Editor




Scott Bryant

Affiliation: The Dougmar Publishing Group Inc, Hamilton, Ontario, Canada

Email

Technical Support



Montadher Issam Abdulkareem, MSC..

Affiliation: College of Information Engineering, Al-Nahrain University, Baghdad, Iraq

Email

Former Editor-in-Chief (2020-2022)




Rayah S. Baban, Ph.D.

Affiliation: Department of Chemistry and Biochemistry, College of Medicine, Al-Nahrain University, Baghdad, Iraq

Email | ORCID | Google Scholar | ResearcherID | Scopus | PubMed

Diversity of the Editorial Board in Geographical Distribution



All members of the Editorial Board have identified their affiliated institutions or organizations, along with the corresponding country or geographic region. Al-Nahrain University remains neutral with regard to any jurisdictional claims.

Journal's information

Publisher: College of Medicine, Al-Nahrain University

Peer-review process: Double-blind

Publishing frequency: Quarterly

Launching year: 2020

Abbreviation: Baghdad J Biochem Appl Biol Sci

E-ISSN: 2706-9915

P-ISSN: 2706-9907

DOI Prefix: 10.47419


Make a Submission

Indexing and Abstracting



Academic Scientific Journals

Membership



Digital Archiving



Download Center

Cover Letter Form

Manuscript Template File

Conflict of Interest Form

Publishing Agreement Form

Graphical Abstract

visitors



Click on the visitor counter above to view more details about visitor statistics and the time spent on the journal website, as well as learn more about the journal's audience.

Keywords



Announcements



Latest publications



Browse

Categories

Biochemistry

Genetics and Molecular Biology

Clinical Biochemistry


Translational Medicine

Agricultural and Biological Sciences


Molecular Biotechnology

Pharmacology, Toxicology and

Pharmaceuticals



Vol. 7 No. 2 (2026): April-June



Published: 10-04-2026

Narrative reviews



Antibiotic Resistance and the Role of Nanoparticles as Antimicrobial/Adjuvant Agents

Abeer Hassan Kadhum, Hasan A. Aal Owaif

88-96




Serological Biomarkers in Forensic Science

A Review

Dunya Abdullah Mohammed, Omar A Mahmoud, Sahar M. Ibrahim, Farah Badri Abed, Muhammed A. Hameed, Shahrzad H. Muhi

110-124



The Hepatoprotective Potential of Alpha-Lipoic Acid

Mechanisms and Translational Evidence

Huda Ayad Al-Mosawi, Ausama Ayob Jaccob, Muhsin S.G. Al-Mozie'i

133-148

Research articles




Ruta graveolens L. Leaves as a Potential Source of Natural Antioxidants and Phytochemicals

A Study from Msallata, Libya

Khaled Muftah Elsherif, Marwa A. Sulaiman, Adel Miltan

155-163



Immunomodulatory Role of Serum Interleukin-37 and Its Association with Biochemical Indicators of Liver Injury in Patients with Chronic HBV and HCV Infections

Rana Talib Mohsen, Sally Hassan

164-168



Remifentanyl vs Fentanyl impact on Ejection Fraction and Troponin in off pump cardiac surgery

Zhiman Ibrahim, Haidar Nassir Mohammad, Majeed Hussein Mustafa

169-176



Evaluation of the Role of Uterine Artery Color Doppler Ultrasound in the Assessment of Low-Risk Pregnancies and in Predicting their Outcome

Bushra Jarallah

149-154



Prevalence and Risk Factors of Pediculosis in Different Primary School in Zakho City, Iraq

Araz Ramadhan Issa, Avan Amr Ali, Wijdan M. Mero

82-87



Impact of Active Helicobacter Pylori Infection on Disability and Relapses in Multiple Sclerosis

Sahar A. Taha, Iman Nozad Mahmood, Mufeed Akram Taha, Esraa Abdulkareem Mohammed

97-101



A Study of Lactate Dehydrogenase (LDH) Levels and Other Biomarkers in Breast Cancer Patients Undergoing Chemotherapy

Implications for Prognosis and Monitoring

Abeer Erfan, Emad Yousif, Ahmed Alshanon, Muna Bufaroosha, Husnun Amalia, Yudhisman Imran, Nany Hairunisa, Raghdha Alsayed

102-109



Evaluation of Serum 25-Hydroxyvitamin D, Ferritin, and CA 15-3 Levels in Women with Breast Cancer

An Iraqi Comparative Study

Marwa Adel Hussein, Ibrahim Ramzi Hadi, Mais Adnan Al_Ward, Mithat Güllü

125-132


Journal's information

Publisher: College of Medicine, Al-Nahrain University**Peer-review process:** Double-blind**Publishing frequency:** Quarterly**Launching year:** 2020**Abbreviation:** Baghdad J Biochem Appl Biol Sci**E-ISSN:** 2706-9915**P-ISSN:** 2706-9907**DOI Prefix:** 10.47419[Make a Submission](#)

Indexing and Abstracting



Membership



Digital Archiving



Download Center


[Cover Letter Form](#)[Manuscript Template File](#)[Conflict of Interest Form](#)[Publishing Agreement Form](#)[Graphical Abstract](#)

visitors



Click on the visitor counter above to view more details about visitor statistics and the time spent on the journal website, as well as learn more about the journal's audience.

Keywords




Browse Issues

First issue
Vol. 1 No. 01 (2020)
October-December (2020)**Previous issue**
Vol. 7 No. 1 (2026)
January-March**Next issue**
Vol. 7 No. 3 (2026)
July-September**Last issue**
Vol. 7 No. 3 (2026)
July-September[All issues](#)

Announcements




Latest publications



Browse

Categories

[Biochemistry](#)
[Genetics and Molecular Biology](#)
[Clinical Biochemistry](#)
[Translational Medicine](#)
[Agricultural and Biological Sciences](#)
[Molecular Biotechnology](#)
[Pharmacology; Toxicology and Pharmaceutics](#)



A Study of Lactate Dehydrogenase (LDH) Levels and Other Biomarkers in Breast Cancer Patients Undergoing Chemotherapy: Implications for Prognosis and Monitoring

Abeer Erfan¹, Emad Yousif^{2*}, Ahmed Alshanon³, Muna Bufaroosha⁴, Husnun Amalia⁵,
Yudhisman Imran⁶, Nany Hairunisa⁷, and Raghda Alsayed²

¹Department of Biology, College of Science, Al-Nahrain University, Baghdad 64021, Iraq

²Department of Chemistry, College of Science, Al-Nahrain University, Baghdad 64021, Iraq

³Center of Biotechnology Researches, University of Al-Nahrain, Baghdad, Iraq

⁴Department of Chemistry, College of Science, United Arab Emirates University, P.O. Box 15551, Al-Ain 1818, UAE

⁵Department of Ophthalmology, Universitas Trisakti, Indonesia

⁶Neurology Department, Faculty of Medicine, Universitas Trisakti, 11440 Jakarta, Indonesia

⁷Department of Occupational Medicine, Universitas Trisakti, Indonesia

Article Info.

Keywords:

Breast cancer,
Chemotherapy response,
Iraqi patients,
Lactate dehydrogenase (LDH),
Prognostic biomarker

Abstract

Background: Breast cancer, after lung cancer, is the second leading cause of cancer-related deaths worldwide, making early diagnosis and continuous treatment crucial. This study evaluated the levels of serum lactate dehydrogenase (LDH) enzyme and other biochemical parameters in post-operative chemotherapy in Iraqi women diagnosed with breast cancer. This study highlights the potential use of LDH as a biomarker for monitoring treatment response, rather than being limited to its traditional diagnostic role.

Objective: To assess the prognostic value of serum LDH along with other biochemical markers, such as alkaline phosphatase (ALP), alanine aminotransferase (ALT), aspartate aminotransferase (AST), electrolytes (Na^+ , K^+ , and Cl^-), kidney functions (urea and creatinine), and prolactin hormone.

Methods: A cross-sectional study was conducted on 150 serum samples collected from 100 female patients diagnosed with breast cancer (after surgery and during chemotherapy) and 50 age-matched healthy women, aged 26–60 years, as a control group. The samples were collected at Al-Amal National Hospital for Cancer Management, Medical City, Baghdad, Iraq, from July 2021 to October 2021. All biochemical parameters (LDH, ALP, ALT, AST, Na^+ , K^+ , Cl^- , urea, creatinine, and prolactin) were measured in the hospital laboratory using standard automated biochemical analyzers and commercially available diagnostic kits.

Results: LDH activity was significantly elevated in breast cancer patients, highlighting its potential as an early prognostic marker for progression or recurrence of disease. Renal function tests (urea and creatinine) and liver enzymes (ALT and AST) were significantly higher in patients, compared to controls ($P < 0.05$). Electrolytes, specifically K^+ and Cl^- , were elevated ($P < 0.05$), while Na^+ showed no significant difference ($P > 0.05$). Levels of ALP were not significantly different between patients and controls ($P > 0.05$). Level of prolactin hormone was also significantly higher in patients ($P < 0.05$) than in controls. These findings underscore LDH as a central prognostic marker, with other biochemical parameters providing supportive information on patient status.

Conclusion: Elevated serum LDH levels in breast cancer patients may serve as an early prognostic marker for progression or recurrence of disease, offering a practical tool for monitoring response to treatment.

Received: 08.20.2025

Accepted: 10.04.2025

Published online: 04.10.2026

Published: 04.10.2026

*Corresponding author: Emad Yousif: emad.yousif@nahrainuniv.edu.iq

How to cite this article: Erfan A et al. A Study of Lactate Dehydrogenase (LDH) Levels and Other Biomarkers in Breast Cancer Patients Undergoing Chemotherapy: Implications for Prognosis and Monitoring. Baghdad Journal of Biochemistry and Applied Biological Sciences, 2026, Vol. 7, No. 2, 102–109. <https://doi.org/10.47419/bjbabs.v7i2.451>

License: Distributed under the terms of The Creative Commons Attribution 4.0 International License (CC BY 4.0), which Permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are properly cited.

Copyright: © 2026 the Authors CC BY license (<http://creativecommons.org/licenses/by/4.0/>).



Introduction

Cancer is characterized by uncontrolled cell growth leading to the formation of tumors, which may be benign—localized and slow-growing—or malignant, with the ability to invade and spread [1–4]. Breast cancer is among the most common malignant tumors worldwide, with over two million new cases reported annually [5,6]. Several risk factors, such as obesity, physical inactivity, and alcohol consumption, increase its incidence, while early detection remains critical for improving outcomes [7–9]. Patients may present with symptoms such as nipple changes, tissue heterogeneity, or unexplained weight loss [8,10]. Breast cancer can be invasive, spreading to surrounding tissues, or non-invasive, restricted to ducts or lobules, with treatment varying accordingly [11].

Breast cancer is characterized by a long doubling time of cancer cells (100–300 days), which provides a valuable opportunity for early detection and treatment. The classification of breast cancer is based on tumor size, lymph node involvement, and the presence of metastasis. Stage 0: the tumor is confined to the site of origin. Stage 1: the tumor spreads into fatty tissue of the breast. Stage 2: The tumor increases in size and may extend further within the breast. Stage 3: The tumor involves nearby lymph nodes and the chest wall. Stage 4 (metastatic stage): The tumor spreads to distant organs, such as bones, liver, lungs, and brain [13,14].

Early diagnosis, adjuvant chemotherapy, hormonal therapy, and radiotherapy play a major role in the treatment of breast cancer and in reducing disease-related risks [15]. Both estrogen (ER) and progesterone (PR) receptors are considered predictive factors in breast cancer. Radical mastectomy was applied in the early 20th century but proved less efficient compared to less severe surgical approaches. Chemotherapy, endocrine therapy, and anti-human epidermal growth factor receptor 2 (HER2)-targeted therapy have demonstrated high efficacy in reducing breast cancer risks and improving the survival [16]. Breast cancer is a heterogeneous disease, and in cases of small tumors, locoregional treatment may be sufficient, whereas cancer that has spread to distant sites requires additional therapy. The main goals of treatment are surgical resection, sampling, and removal of axillary lymph nodes, followed by radiation [17].




Figure (1): Difference between normal cells lining the milk duct, ductal carcinoma *in situ* (DCIS), and invasive cancer [12].

In order to better understand disease progression and treatment response, biomarker analysis has become an essential tool. Histological grade, obtained via core needle biopsy (CNB), axillary status, and tumor size are important parameters in breast cancer. Estrogen receptor alpha (ER- α) is a key predictive factor for advanced-stage patients responding to hormone therapy, which is nontoxic and suitable for long-term use [18]. Cancer antigen CA15.3 is elevated pre-surgically and serves as an important marker for evaluating response after chemotherapy [19].

Among emerging biomarkers, serum lactate dehydrogenase (LDH) has gained attention because of its strong association with progression and early detection of breast cancer [20]. Serum LDH catalyzes the reversible conversion of pyruvate to lactate, playing a crucial role in anaerobic metabolism and maintaining levels of nicotinamide adenine dinucleotide (NAD⁺) for glycolysis [21]. It is found in the heart and muscle tissues and exhibits high activity and stability, making it a central enzyme in the glycolytic pathway and a potential therapeutic target [22]. Elevated serum LDH levels are observed in a variety of disorders, such as cardiac, hepatic, skeletal muscle, and renal diseases as well as leukemia [23–25]. In cancer, high serum LDH levels reflect altered metabolism that supports tumor growth and metastasis, establishing its value as a diagnostic and prognostic biomarker [23,26].

Lactate dehydrogenase is an enzyme that catalyzes the reversible conversion of pyruvate to lactate, playing a key role in maintaining levels of NAD⁺ for glycolysis. In cancer cells, LDH contributes to the “Warburg effect,” a metabolic shift where glucose is preferentially metabolized to lactate even in the presence of oxygen, supporting rapid tumor growth and survival [27–31]. Mitochondrial dysfunction in tumor cells leads to increased membrane permeability and the release of intracellular enzymes, including LDH, which correlates with abnormal apoptotic signaling [32,33]. Elevated serum LDH levels in serum thus reflect altered cellular metabolism and progression of




Figure (2): The reaction catalyzed by lactate dehydrogenase (LDH).

tumor, highlighting its value as a diagnostic and prognostic biomarker in breast cancer.

2. Materials and Methods

2.1. Study subjects

This cross-sectional study included 150 Iraqi women, aged 25–60 years, who were divided into two groups. The first group, the control group, included 50 women with no history of cancer or any other chronic disease. The second group included 100 women diagnosed with breast cancer, and were enrolled at the beginning of their chemotherapy and before undergoing radiotherapy. Diagnosis was confirmed through medical record review. Patients with chronic diseases, such as diabetes mellitus, hypertension, ischemic heart disease, and thyroid disorders, as well as smokers or alcohol consumers, were excluded from the study. Verbal informed consent was obtained from all participants prior to sample collection, and the study was approved by the Ethics Committee of the Al Amal National Cancer Hospital, Medical City, Baghdad, Iraq.

2.2. Specimen collection and preparation

Sample collection was approved by the management of Al Amal National Cancer Hospital, Baghdad, Iraq, under the direction of Dr. Mussab Kadhim Al-Aboodi. Blood samples (5 mL) were collected from each participant and placed in gel tubes. Samples from breast cancer patients were collected at different time points relative to chemotherapy, such as prior to starting of chemotherapy, after the first session, and after two to three sessions. Tubes were kept at room temperature until coagulation, and then centrifuged at 3,000 rpm for 15 min. The resulting sera were transferred into Eppendorf tubes and stored at low temperatures until analysis.

All biochemical analyses, such as LDH, alanine aminotransferase (ALT), aspartate aminotransferase (AST), alkaline phosphatase (ALP), prolactin (PRL), electrolytes, and kidney function tests, were performed using automated analyzers available in the hospital laboratory. Data distribution was tested using Shapiro–Wilk and Kolmogorov–Smirnov tests, and multifactorial analysis was used to control for potential confounding variables, such as age, disease stage, and treatment type. Statistical significance was set at $P < 0.05$.

2.3. Statistical analysis

All statistical analyses were performed using GraphPad Prism version 8.0.2. Data distribution was first tested for normality using Shapiro–Wilk and Kolmogorov–Smirnov tests. Continuous variables were presented as mean $\pm$ standard deviation (SD). The unpaired t -test was used to compare normally distributed data between the study groups. Multifactorial analysis was applied to control potential confounding variables, viz. age, disease stage, and treatment type. Differences with $P < 0.05$ were considered statistically significant [34].

3. Results and Discussion

3.1. Evaluation of serum lactate dehydrogenase

Lactate dehydrogenase enzyme plays an important role in glycolysis by converting pyruvate to lactate at the end of this process. The release of high levels of serum LDH into the blood is related to several diseases, such as cancers causes destruction the tissue [35]. The serum LDH levels in postoperative patients of breast cancer and controls are shown in Table 1 and Figure 5, which explained that the serum LDH levels in cancer patients were significantly high (352.28 ± 67.4) with $P < 0.0001$, compared to controls (120 ± 27.34). The LDH activity in breast cancer patients was higher, compared to that in controls, and this could be due to increased cell growth and cell migration compared to normal cells. Additionally, cancer cells have high energy demands and rely on aerobic glycolysis (Warburg effect), mitochondrial dysfunction, increased cellular permeability, and chemotherapy-induced inflammation, all of which contribute to elevated LDH levels in the serum. The cancer cells require excess of energy so the LDH can be considered an alternative to meet metabolic and aerobic glycolytic needs of cancer cells. In breast cancer patients, higher levels of LDH activity can give a prognostic indicator for the early warning of metastasis or recurrence [36,37].

3.2. Evaluation of serum urea & creatinine levels

The urea cycle reactions are important for removing free ammonia by hepatic detoxification and expelling it through the urethra; hence, urea levels are particularly important in patients exposed to diseases or treatments that can impair kidney function [38]. In this work, both urea and creatinine levels were evaluated in breast cancer patients, and the results shown in Table 2 and Figure 6 illustrate significantly higher levels of urea in breast cancer patients (41.76 ± 5.76), compared to controls (20.2 ± 3.61 , $P < 0.0001$).




Figure (3): Activity levels of serum lactate dehydrogenase (LDH) in breast cancer patients and healthy controls.

Table (1): Serum LDH levels in breast cancer patients and healthy controls.

Group	LDH (mean $\pm$ SD)	P value
Controls	120 $\pm$ 27.34	<0.0001
Patients	352.28 $\pm$ 67.4	

Creatinine levels were also elevated (0.822 ± 0.16 in cancer patients vs. 0.506 ± 0.1 in controls, $P < 0.0001$).

The increase in urea levels in breast cancer patients may be attributed not only to tissue destruction and nucleic acid turnover but also to chemotherapy-induced renal stress, inflammation, and altered metabolism, which affect nitrogen balance. The levels of creatinine in serum play an important role in evaluating glomerular filtration rate and kidney function efficiency. Chemotherapeutic drugs can impair kidney function directly by damaging renal cells or indirectly through vascular and tubular effects, contributing to elevated serum urea and creatinine levels in cancer patients [39–41].

3.3. Evaluation of liver enzymes in serum (AST, ALT, and ALP)

Both serum AST and serum ALT, in addition to (ALP), were evaluated in breast cancer patients and healthy controls, and the results are shown in Table 3 and Figure 7. ALP is a nonspecific enzyme involved in hydrolyzing phosphate in alkaline medium, with activity reflecting presence of isoenzymes in the bone, liver, kidney, and intestinal lining [42]. ALT and AST are liver-related enzymes, and their serum levels increase in various diseases, including liver damage, viral hepatitis, myocardial infarction, bile duct obstruction, acute pancreatitis, and anemia [43].

The levels of ALP in cancer patients (113.64 ± 49.50 , $P = 0.0630$) did not show a statistically significant difference, compared to controls (82.1 ± 22.3), suggesting that early-stage breast cancer or non-hepatic factors may not markedly affect ALP levels. Increase in ALP is generally observed in advanced stages of breast cancer and may serve to distinguish between stages or indicate metastasis [44,45].

The ALT activity was significantly higher in breast cancer patients (61.68 ± 30.5), compared to controls (18.1 ± 4.4 , $P < 0.0001$), and AST was also elevated (39.52 ± 22.5 vs. 20.6 ± 5.6 , $P = 0.0140$). Both ALT and AST catalyze the conversion of alanine to pyruvate and aspartate to oxaloacetate, respectively, and elevated levels reflect liver stress or damage, which may be secondary to tumor invasion, chemotherapy effects, or systemic inflammation. Previous studies report higher ALT and AST in malignant versus benign breast cancer patients, supporting their use as indicators of compromise of liver and renal functioning in progression of cancer [46].

3.4. Evaluation of serum electrolyte levels (Na^+ , K^+ , and Cl^-)

Kidney dysfunction is widely common in patients with malignant cancer because chemotherapy and other treatments can impair renal function, cause electrolyte disturbances, and induce immune deficiency [47]. However, many factors affect electrolyte levels, so identifying the causes is not straightforward [48].

Electrolyte levels were assessed in breast cancer patients and compared to healthy controls (Table 4 and Figure 8). Sodium levels did not differ significantly in

cancer patients (142.7 ± 6.5 , $P = 0.0663$), compared to controls (138.62 ± 3.4), suggesting that homeostatic mechanisms may maintain serum sodium despite renal stress or chemotherapy.

On the other hand, chloride levels were significantly higher in cancer patients (113.1 ± 4.5 , $P < 0.0001$), compared to controls (103.28 ± 2.9). Potassium, which plays a key role in muscle excitability and nerve functioning, was also elevated in cancer patients (4.36 ± 0.55 , $P = 0.0212$), compared to controls (3.92 ± 0.25). This may result from the release of intracellular contents from damaged tumor cells, chemotherapy-induced renal impairment, or altered hormonal regulation associated with cancer.

Serum electrolyte imbalance is clinically important for monitoring breast cancer patients, helping to assess renal injury, nephrotoxicity, and the overall prognosis. They




Figure (4): Serum levels of prolactin (PRL) in breast cancer patients and healthy controls.

Table (2): Serum levels of urea and creatinine in breast cancer patients and healthy controls.

Group	Urea (mean $\pm$ SD)	P value	Creatinine (mean $\pm$ SD)	P value
Controls	20.2 $\pm$ 3.61	<0.0001	0.506 $\pm$ 0.1	<0.0001
Patients	41.76 $\pm$ 5.76		0.822 $\pm$ 0.16	

Table (3): Serum levels of ALP, AST, and ALT in breast cancer patients and healthy controls.

Parameters	Mean $\pm$ SD		
	Controls	Patients	P value
ALP (U/L)	82.1 $\pm$ 22.3	113.64 $\pm$ 49.50	0.0630
Serum AST (U/L)	20.6 $\pm$ 5.6	39.52 $\pm$ 22.5	0.0140
Serum ALT (U/L)	18.1 $\pm$ 4.4	61.68 $\pm$ 30.5	<0.0001

Table (4): Serum levels of electrolytes in breast cancer patients and healthy controls.

Parameters	Mean $\pm$ SD		
	Controls	Patients	P value
Na^+	138.62 $\pm$ 3.4	142.7 $\pm$ 6.5	0.0663
K^+	3.92 $\pm$ 0.25	4.36 $\pm$ 0.55	0.0212
Cl^-	103.28 $\pm$ 2.9	113.1 $\pm$ 4.5	<0.0001

may also reflect systemic effects of tumor metabolism and treatment-related stress [49,50].

3.5. Evaluation of serum prolactin levels

Imbalance in hormone levels is associated with the development of various diseases. Recently, studies have focused on serum PRL in the prognosis of breast cancer. Many studies have shown a possible relationship between breast cancer and PRL hormone. PRL has cytokine-like properties; its receptor binds to the cell surface and cytoplasm, activating the Jak2/Stat5 signaling pathway [51,52].

Table 5 and Figure 9 show that breast cancer patients had significantly higher PRL levels (32.45 ± 7.0 , $P < 0.0001$), compared to controls (7.186 ± 2.5). Elevated PRL may increase the risk of breast cancer, potentially by promoting cell division, reducing apoptosis, and contributing

Table (5): Serum levels of prolactin (PRL) in breast cancer patients and healthy controls.

Group	Prolactin (mean ± SD)	P value
Controls	7.186 ± 2.5	<0.0001
Patients	32.45 ± 7.0	




Figure (5): (A) Serum levels of urea in breast cancer patients and healthy controls; (B) Serum levels of creatinine in breast cancer patients and healthy controls.




Figure (6): (A) Serum levels of ALP in breast cancer patients and healthy controls; (B) Serum levels of ALT in breast cancer patients and healthy controls; (C) Serum levels of AST in breast cancer patients and healthy controls.




Figure (7): (A) Serum levels of Na⁺ in breast cancer patients and healthy control; (B) Serum levels of Cl⁻ in breast cancer patients and healthy controls; (C) Serum levels of K⁺ in breast cancer patients and healthy controls.

to resistance to cytotoxic drugs, although the exact mechanism remains unclear [53].

4. Conclusions

This study evaluates the levels of serum LDH in patients with breast cancer at different stages of chemotherapy, and demonstrates that LDH activity in cancer patients is higher, compared to that in controls. This can be due to increased cell growth and migration of cancer cells, compared to normal cells. Cancer cells require excess energy, so serum LDH serves as an alternative to meet the metabolic and aerobic glycolytic needs of cancer cells. In breast cancer patients, the higher levels of LDH activity can serve as a prognostic indicator for early warning of metastasis or recurrence. Additionally, it can be used for monitoring cancer progression and response to treatment.

Finally, this study demonstrates that elevated serum LDH levels are significantly associated with breast cancer and its progression. Serum LDH can serve as a valuable biomarker for prognosis, early detection of metastasis, and monitoring of therapeutic response, highlighting its potential role in clinical management of breast cancer patients.

Acknowledgments

The authors appreciated the efforts and assistance of the management of Al Amal National Cancer Hospital under the supervision of the hospital director, Dr. Mussab Kadhim Al-Aboodi, and the staff who facilitated the samples and collection process.

Conflict of Interest

The authors declared no conflict of interest.

Author Contributions (CRediT Taxonomy)

Contributor Role	Degree of Contribution		
	Lead	Equal	Supporting
Conceptualization		X	
Data curation		X	
Formal analysis		X	
Funding acquisition		X	
Investigation		X	
Methodology		X	
Project administration		X	
Resources		X	
Software		X	
Supervision		X	
Validation		X	
Visualization		X	
Writing-original draft		X	
Writing-review & editing		X	

References

[1] Sinha, T., Tumors: benign and malignant. *Cancer Ther Oncol Int J*, 10(3): 1–3, 2018. <https://doi.org/10.19080/CTOIJ.2018.10.555790>.

[2] Gomes, C.C., Recurrent driver mutations in benign tumors. *Mutat Res Rev Mutat Res*, 789(108412), 2022. <https://doi.org/10.1016/j.mrrev.2022.108412>.

[3] Patel A., Benign vs malignant tumors. *Am Med Ass JAMA Oncol*, 6(9): 1488, 2020. <https://doi.org/10.1001/jamaoncol.2020.2592>.

[4] Lee, Y.; Jee, W.H.; Whang, Y.S.; Jung, C.K.; Chung, Y.G.; Lee S.Y.; Benign versus malignant soft-tissue tumors: differentiation with 3t magnetic resonance image textural analysis including diffusion-weighted imaging. *Investig Magn Reson Imaging*, 25(1): 118–128, 2021. <https://doi.org/10.13104/imri.2021.25.2.118>.

[5] Liu, Y.; Li, H.; Zhu, Z.; Chen, C.; Zhang, X.; Jin, G.; Li, H., RSDCNet: an efficient and lightweight deep learning model for benign and malignant pathology detection in breast cancer. *Digital Health*, 11: 1–11, 2025. <https://doi.org/10.1177/20552076251336286>.

[6] Satpathi, S.; Gaurkar, S.S.; Potdukhe, A.; Wanjari, M.B., Unveiling the role of hormonal imbalance in breast cancer development: a comprehensive review. *Cureus*, 15(7): e41737, 2023. <https://doi.org/10.7759/cureus.41737>.

[7] Giaquinto, A.N.; Sung, H.; Miller, K.D.; Kramer, J.L.; Newman, L.A.; Minihan, A.; et al., Breast Cancer Statistics, 2022. *CA Cancer J Clin*, 72(1): 524–541, 2022. <https://doi.org/10.3322/caac.21754>.

[8] Lina, L., Signs and symptoms of breast cancer. *Eur J Clin Oncol*, 3(2): 1, 2021.

[9] Prusty, R.K.; Begum, S.; Patil, A.; Naik, D.; Pimple, S.; Mishra, G., Knowledge of symptoms and risk factors of breast cancer among women: a community based study in a low socio-economic area of Mumbai, India. *BMC Women's Health*, 20(106): 1–12, 2020. <https://doi.org/10.1186/s12905-020-00967-x>.

[10] Ikhuoria, E.B.; Bach, C., Introduction to breast carcinogenesis symptoms, risks factors, treatment and management. *Eur J Eng Res Sci*, 3(7): 58–66, 2018. <https://doi.org/10.24018/ejeng.2018.3.7.745>.

[11] Hessari, P.M.; Montazeri, A., Health-related quality of life in breast cancer patients: review of reviews from 2008 to 2018. *Health Qual Life Outcomes*, 18(1): 1–25, 2020. <https://doi.org/10.1186/s12955-020-01591-x>.

[12] Lips, E.H.; Kumar, T.; Megalios, A.; Visser, L.; Sheinman, M.; Fortunato, A.; et al., Genomic analysis defines clonal relationships of ductal carcinoma *in situ* and recurrent invasive breast cancer. *Nat Genet*, 54(1): 850–860, 2022. <https://doi.org/10.1038/s41588-022-01082-3>.

[13] Nargis, H.F.; Nawaz, H.; Ditta, A.; Mahmood, T.; Majeed, M.I.; Rashid, N.; et al., Raman Spectroscopy of Blood Plasma Samples from breast Cancer Patients at Different Stages. *Technological University, Dublin, UK*; 2019; pp. 1–24. <https://doi.org/10.1016/j.saa.2019.117210>.

[14] Alwan, N.A.; Kerr, D.; Al-Okati, D.; Pezella, F.; Tawfeeq, F.N., Comparative study on the clinicopathological profiles of breast cancer among Iraqi and British patients. *Open Public Health J*, 11(1): 177–191, 2018. <https://doi.org/10.2174/1874944501811010177>.

[15] Hassan, M.S.U.; Ansari, J.; Spooner, D.; Hussian, S.A., Chemotherapy for breast cancer (review). *Oncol Rep*, 24(1): 1121–1131, 2010. https://doi.org/10.3892/or_00000963.

[16] Anampa, J.; Makower, D.; Sparano, J.A., Progress in adjuvant chemotherapy for breast cancer: an overview. *BMC*

- Med, 13(195): 1–13, 2015. <https://doi.org/10.1186/s12916-015-0439-8>.
- [17] Waks, A.G.; Winer, E.P., Breast cancer treatment a review. *JAMA*, 321(3): 288–300, 2019. <https://doi.org/10.1001/jama.2018.19323>.
 - [18] Colomer, R.; Lopez, I.A.; Albanell, J.; Caballero, T.G.; Ciruelos, E.; Cortes, J.; et al, Biomarkers in breast cancer: a consensus statement by the Spanish Society of Medical Oncology and the Spanish Society of Pathology. *Clin Transl Oncol*, 20: 815–826, 2018 (Springer). <https://doi.org/10.1007/s12094-017-1800-5>.
 - [19] Patani, N.; Martin, L.A.; Dowsett, M., Biomarkers for the clinical management of breast cancer: international perspective. *Int J Cancer*, 133(1): 1–13, 2013. <https://doi.org/10.1002/ijc.27997>.
 - [20] Adnan, S.F.; Al-Abady, Z.N., Investigation the role of lactate dehydrogenase, caspase and the oxidative stress levels in breast cancer patients. *J Biomed Biochem*, 3(4), 2024. <https://doi.org/10.57238/jbb.2024.7432.1129>.
 - [21] Khan, A.A.; Allemaillem, K.S.; Alhumaydhi, F.A., Gowder, S.; Rahmani A.H., The biochemical and clinical perspectives of lactate dehydrogenase: an enzyme of active metabolism, endocrine, metabolic & immune disorders. *Drug Targets*, 20(1): 1–14, 2020. <https://doi.org/10.2174/1871530320666191230141110>.
 - [22] Chaari, A.; Al-Ali, D.; Roach, J., Biochemistry course-based undergraduate research experience: purification, characterization, and identification of an unknown lactate dehydrogenase isoenzyme. *Biochem Mol Biol Educ*, 48(1): 369–380, 2020. <https://doi.org/10.1002/bmb.21363>.
 - [23] Kumar, P.; Nagarajan, A.; Uchil, P.D, Analysis of Cell Viability by the Lactate Dehydrogenase Assay. Cold Spring Harbor Laboratory Press, Woodbury, NY; 2018; pp. 465–468. <https://doi.org/10.1101/pdb.prot095497>.
 - [24] Klein, R.; Nagy, O.; Tothovam, C.; Frederika, C., Clinical and diagnostic significance of lactate dehydrogenase and its isoenzymes in animals. *Vet Med Int*, 2020(5346483): 1–11, 2020. <https://doi.org/10.1155/2020/5346483>.
 - [25] Cai, X.; Wang, T.; Xie, L., Lactate dehydrogenase is associated with flow mediated dilation in hypertensive patients. *Sci Rep*, 13(768): 1–9, 2023. <https://doi.org/10.1038/s41598-023-27614-3>.
 - [26] Valvona, C.J.; Fillmore, H.L.; Nunn, P.B.; Pilkington, G.J., The regulation and function of lactate dehydrogenase A: therapeutic potential in brain tumor. *Brain Pathol*, 1(26): 3–17, 2016. <https://doi.org/10.1111/bpa.12299>.
 - [27] Farhana, A.; Lappin, S.L., Biochemistry, lactate dehydrogenase. StatPearls [Internet], Treasure Island, FL. StatPearls Publishing; 2026; pp. 1–8.
 - [28] Forkasiewicz, A.; Dorociak, M.; Stach, K.; Szelachowski, P.; Tabola, R.; Augoff, K., The usefulness of lactate dehydrogenase measurements in current oncological practice. *Cell Mol Biol Lett*, 25(35): 1–14, 2020. <https://doi.org/10.1186/s11658-020-00228-7>.
 - [29] Zhou, Y.; Niu, W.; Luo, Y.; Li, H.; Xie, Y.; Wang, H.; et al., p53/Lactate dehydrogenase A axis negatively regulates aerobic glycolysis and tumor progression in breast cancer expressing wild-type p53. *Cancer Sci*, 110(1): 939–949, 2019. <https://doi.org/10.1111/cas.13928>.
 - [30] Zhou, Y.; Qi, M.; Yang M., Current status and future perspectives of lactate dehydrogenase detection and medical implications: a review. *Biosensors*, 12(1145): 1–17, 2022. <https://doi.org/10.3390/bios12121145>.
 - [31] Fukushi, A.; Kim, H.D.; Chang, Y.C.; Kim, C.H., Revisited metabolic control and reprogramming cancers by means of the Warburg effect in tumor cells. *Int J Mol Sci*, 23(1): 1–21, 2022. <https://doi.org/10.3390/ijms231710037>.
 - [32] Jurisic, V.; Radenkovic, S.; Konjevic, G., The actual role of LDH as tumor marker, biochemical and clinical aspects, advances in cancer biomarkers. *Adv Exp Med Biol*, 867(1): 115–124, 2015. https://doi.org/10.1007/978-94-017-7215-0_8.
 - [33] Zaib, S.; Hayyat, A.; Ali, N.; Gul, A.; Naveed, M.; Khan, I., Role of mitochondrial membrane potential and lactate dehydrogenase A in apoptosis. *Anti-Cancer Agents Med Chem*, 22(11): 2048–2062, 2022. <https://doi.org/10.2174/1871520621666211126090906>.
 - [34] Glover, T.; Mitchell, K., An Introduction to Biostatistics. Waveland Press, Long Grove, IL; 2008.
 - [35] Liu, D.; Wang, D.; Wu, C.; Zhang, L.; Mei, Q.; Hu, G.; et al., Prognostic significance of serum lactate dehydrogenase in patients with breast cancer: a meta-analysis. *Cancer Manag Res*, 11(1): 3611–3619, 2019. <https://doi.org/10.2147/CMAR.S199260>.
 - [36] Kasht, G.J.; Abed, B.A.; Farhan, L.O.; Salman, I.N.; Dawood, A.S., A comparative study to determine LDH enzyme levels in serum samples of women with breast cancer and women with breast cancer and type 2 diabetes mellitus. *J Med Chem Sci*, 6(1): 883–890, 2023. <https://doi.org/10.26655/JMCHEMSCI.2023.4.20>.
 - [37] Basnyat, A.S.; Jha, A.; Pathak, R; Shrestha, B.G., Study of serum lactate dehydrogenase and gamma-glutamyl transpeptidase in breast cancer patients receiving chemotherapy. *J Trop Life Sci*, 7(2): 128–132, 2017. <https://doi.org/10.11594/jtls.07.02.07>.
 - [38] Attallah, B.M.; Al-Ani, W.Y.; Alboaisa, N.S., Total antioxidant, renal function test in Iraqi women with breast cancer. *World Bull Public Health*, 16(1): 158–161, 2022.
 - [39] Kebede, T.; Melak, T.; Ibn-Sina, A.; Fasil, A., Assessment of serum uric acid, urea, and glucose levels and associated factors among breast cancer patients attending a tertiary hospital in Bahirdar, Ethiopia: a comparative cross-sectional study. *Ethiop J Health Sci*, 32(6): 1183–1192, 2022. <https://doi.org/10.4314/ejhs.v32i6.16>.
 - [40] Prabhu, S.N.; Mukhopadhyay, S.C.; Gooneratne, C.; Davidson, A.S.; Liu, G., Interdigital sensing system for detection of levels of creatinine from the samples. *Springer Nat*, 36(1): 267–309, 2021. https://doi.org/10.1007/978-3-030-62684-6_11.
 - [41] Parker, A.L.; Toulabi, L.; Oike, T.; Kanke, Y.; Patel, D.; Tada T.; et al., Creatine riboside is a cancer cell-derived metabolite associated with arginine auxotrophy. *J Clin Invest*, 132(14): 1–17, 2022. <https://doi.org/10.1172/JCI157410>.
 - [42] Jiang, C.; Hu, F.; Xia, X.; Guo X., Prognostic value of alkaline phosphatase and bone-specific alkaline phosphatase in breast cancer: a systematic review and meta-analysis. *Int J Biol Markers*, 38(1): 25–36, 2023. <https://doi.org/10.1177/03936155231154662>.
 - [43] AL-Mashhadani, Z.I.; Mukhlis, A.J.; AL-Faraji, A.S., Estimation of ALP, GPT and GOT activities in Iraqi patients female with breast cancer. *Ibn Al-Haitham J Pure Appl Sci*, 25(1): 1–10, 2012.
 - [44] Long, Z.Q.; Hua, X.; Zhang, W.; Lv, S.; Deng, J.; Guo, L.; et al., Prognostic impact of the pretreatment albumin to alkaline phosphatase ratio for nonmetastatic breast cancer patients. *Cancer Manag Res*, 11(1): 4809–4814, 2019. <https://doi.org/10.2147/CMAR.S200759>.
 - [45] Singh, A.K.; Pandey, A.; Tewari, M.; Kumar, R.; Sharma, A.; Singh, K.A.; et al., Advanced stage of breast cancer hoist alkaline phosphatase activity: risk factor for females in India. *Biotech*, 3(1): 517–520, 2013. <https://doi.org/10.1007/s13205-012-0113-1>.
 - [46] Yadav, C.P., Kaushal, V.; Beniwal P., Evaluation of serum biochemical profile of breast cancer patients. *Int J Med Res Health Sci*, 5(7): 1–7, 2016.

- [47] Abdulaziz, O.; Almeshmadi, M., Dysregulation of serum electrolytes is detected in untreated cancer patients in Taif city. *Ann Cancer Res Ther*, 29(2): 126–130, 2021. <https://doi.org/10.4993/acrt.29.126>.
- [48] Alem, A.; Edae, C.K.; Wabalo, E.K.; Tareke, A.A.; Bedanie, A.A.; Reta, W.; et al., Factors influencing the occurrence of electrolyte disorders in cancer patients. *SAGE Open Med*, 9(1): 1–7, 2021. <https://doi.org/10.1177/20503121211052861>.
- [49] Yadav, A.S.; Khodke, P.V., Status of serum electrolytes in cancer patients. *Int J Basic Appl Med Sci*, 5(1): 208–211, 2015.
- [50] Kumari, R.; Siddiqui, A.A.; Zubair, T.; Fatima, S.; Shaheer, A.; Imtiaz, S.; et al., Levels of electrolyte in cancer patients; a prospective study focusing on the variations before and after therapy. *Palliat Med Care*, 5(4): 1–4, 2018. <https://doi.org/10.15226/2374-8362/5/4/00168>.
- [51] Ali, J.K.; Hassan, S.H.; Merzah, M.A., Prolactin serum levels and breast cancer: relationships with hematological factors among cases in Karbala province, Iraq. *Med J Babylon*, 15(2): 178–181, 2018.
- [52] Kolomiiets, O.; Yazykov, O.; Piddubnyi, A.; Lyndin, M.; Lukavenko, I.; Andryushchenko, V.; et al, The expression of prolactin receptors in benign breast tumors is not associated with serum prolactin level. *J Clin Med*, 10(1): 1–10. <https://doi.org/10.3390/jcm10245866>

A Study of Lactate Dehydrogenase (LDH) Levels and Other Biomarkers in Breast Cancer Patients Undergoing Chemotherapy/ Implications for Prognosis and Monitoring

by Nany Hairunisa

A Study of Lactate Dehydrogenase (LDH) Levels and Other Biomarkers in Breast Cancer Patients Undergoing Chemotherapy/ Implications for Prognosis and Monitoring

ORIGINALITY REPORT

11%
SIMILARITY INDEX

8%
INTERNET SOURCES

10%
PUBLICATIONS

5%
STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

2%
★ downloads.hindawi.com
Internet Source

Exclude quotes On
Exclude bibliography On

Exclude matches < 15 words



A Study of Lactate Dehydrogenase (LDH) Levels and Other Biomarkers in Breast Cancer Patients Undergoing Chemotherapy: Implications for Prognosis and Monitoring

Abeer Erfan¹, Emad Yousif^{2*}, Ahmed Alshanon³, Muna Bufaroosha⁴, Husnun Amalia⁵,
Yudhisman Imran⁶, Nany Hairunisa⁷, and Raghda Alsayed²

¹Department of Biology, College of Science, Al-Nahrain University, Baghdad 64021, Iraq

²Department of Chemistry, College of Science, Al-Nahrain University, Baghdad 64021, Iraq

³Center of Biotechnology Researches, University of Al-Nahrain, Baghdad, Iraq

⁴Department of Chemistry, College of Science, United Arab Emirates University, P.O. Box 15551, Al-Ain 1818, UAE

⁵Department of Ophthalmology, Universitas Trisakti, Indonesia

⁶Neurology Department, Faculty of Medicine, Universitas Trisakti, 11440 Jakarta, Indonesia

⁷Department of Occupational Medicine, Universitas Trisakti, Indonesia

Article Info.

Keywords:

Breast cancer,
Chemotherapy response,
Iraqi patients,
Lactate dehydrogenase (LDH),
Prognostic biomarker

Abstract

Background: Breast cancer, after lung cancer, is the second leading cause of cancer-related deaths worldwide, making early diagnosis and continuous treatment crucial. This study evaluated the levels of serum lactate dehydrogenase (LDH) enzyme and other biochemical parameters in post-operative chemotherapy in Iraqi women diagnosed with breast cancer. This study highlights the potential use of LDH as a biomarker for monitoring treatment response, rather than being limited to its traditional diagnostic role.


Objective: To assess the prognostic value of serum LDH along with other biochemical markers, such as alkaline phosphatase (ALP), alanine aminotransferase (ALT), aspartate aminotransferase (AST), electrolytes (Na⁺, K⁺, and Cl⁻), kidney functions (urea and creatinine), and prolactin hormone.

Methods: A cross-sectional study was conducted on 150 serum samples collected from 100 female patients diagnosed with breast cancer (after surgery and during chemotherapy) and 50 age-matched healthy women, aged 26–60 years, as a control group. The samples were collected at Al-Amal National Hospital for Cancer Management, Medical City, Baghdad, Iraq, from July 2021 to October 2021. All biochemical parameters (LDH, ALP, ALT, AST, Na⁺, K⁺, Cl⁻, urea, creatinine, and prolactin) were measured in the hospital laboratory using standard automated biochemical analyzers and commercially available diagnostic kits.

Results: LDH activity was significantly elevated in breast cancer patients, highlighting its potential as an early prognostic marker for progression or recurrence of disease. Renal function tests (urea and creatinine) and liver enzymes (ALT and AST) were significantly higher in patients, compared to controls ($P < 0.05$). Electrolytes, specifically K⁺ and Cl⁻, were elevated ($P < 0.05$), while Na⁺ showed no significant difference ($P > 0.05$). Levels of ALP were not significantly different between patients and controls ($P > 0.05$). Level of prolactin hormone was also significantly higher in patients ($P < 0.05$) than in controls. These findings underscore LDH as a central prognostic marker, with other biochemical parameters providing supportive information on patient status.

Conclusion: Elevated serum LDH levels in breast cancer patients may serve as an early prognostic marker for progression or recurrence of disease, offering a practical tool for monitoring response to treatment.

Received: 08.20.2025
Accepted: 10.04.2025
Published online: 04.10.2026
Published: 04.10.2026



*Corresponding author: Emad Yousif, email: yousif@nahrainu.edu.iq

How to cite this article: Erfan A et al. A Study of Lactate Dehydrogenase (LDH) Levels and Other Biomarkers in Breast Cancer Patients Undergoing Chemotherapy: Implications for Prognosis and Monitoring. Baghdad Journal of Biochemistry and Applied Biological Sciences, 2026, Vol. 7, No. 2, 102–109. <https://doi.org/10.47419/bjabs.v7i2.451>

License: Distributed under the terms of The Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are properly cited.

Copyright: © 2026 the Authors CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

Cancer is characterized by uncontrolled cell growth leading to the formation of tumors, which may be benign—localized and slow-growing—or malignant, with the ability to invade and spread [1-4]. Breast cancer is among the most common malignant tumors worldwide, with over two million new cases reported annually [5,6]. Several risk factors, such as obesity, physical inactivity, and alcohol consumption, increase its incidence, while early detection remains critical for improving outcomes [7-9]. Patients may present with symptoms such as nipple changes, tissue heterogeneity, or unexplained weight loss [8,10]. Breast cancer can be invasive, spreading to surrounding tissues, or non-invasive, restricted to ducts or lobules, with treatment varying accordingly [11].

Breast cancer is characterized by a long doubling time of cancer cells (100-300 days), which provides a valuable opportunity for early detection and treatment. The classification of breast cancer is based on tumor size, lymph node involvement, and the presence of metastasis. Stage 0: the tumor is confined to the site of origin. Stage 1: the tumor spreads into fatty tissue of the breast. Stage 2: The tumor increases in size and may extend further within the breast. Stage 3: The tumor involves nearby lymph nodes and the chest wall. Stage 4 (metastatic stage): The tumor spreads to distant organs, such as bones, liver, lungs, and brain [13,14].

Early diagnosis, adjuvant chemotherapy, hormonal therapy, and radiotherapy play a major role in the treatment of breast cancer and in reducing disease-related risks [15]. Both estrogen (ER) and progesterone (PR) receptors are considered predictive factors in breast cancer. Radical mastectomy was applied in the early 20th century but proved less efficient compared to less severe surgical approaches. Chemotherapy, endocrine therapy, and anti-human epidermal growth factor receptor 2 (HER2)-targeted therapy have demonstrated high efficacy in reducing breast cancer risks and improving the survival [16]. Breast cancer is a heterogeneous disease, and in cases of small tumors, locoregional treatment may be sufficient, whereas cancer that has spread to distant sites requires additional therapy. The main goals of treatment are surgical resection, sampling, and removal of axillary lymph nodes, followed by radiation [17].




Figure (1): Difference between normal cells lining the milk duct, ductal carcinoma *in situ* (DCIS), and invasive cancer [12].

In order to better understand disease progression and treatment response, biomarker analysis has become an essential tool. Histological grade, obtained via core needle biopsy (CNB), axillary status, and tumor size are important parameters in breast cancer. Estrogen receptor alpha (ER- α) is a key predictive factor for advanced-stage patients responding to hormone therapy, which is nontoxic and suitable for long-term use [18]. Cancer antigen CA15.3 is elevated pre-surgically and serves as an important marker for evaluating response after chemotherapy [19].

Among emerging biomarkers, serum lactate dehydrogenase (LDH) has gained attention because of its strong association with progression and early detection of breast cancer [20]. Serum LDH catalyzes the reversible conversion of pyruvate to lactate, playing a crucial role in anaerobic metabolism and maintaining levels of nicotinamide adenine dinucleotide (NAD⁺) for glycolysis [21]. It is found in the heart and muscle tissues and exhibits high activity and stability, making it a central enzyme in the glycolytic pathway and a potential therapeutic target [22]. Elevated serum LDH levels are observed in a variety of disorders, such as cardiac, hepatic, skeletal muscle, and renal diseases as well as leukemia [23-25]. In cancer, high serum LDH levels reflect altered metabolism that supports tumor growth and metastasis, establishing its value as a diagnostic and prognostic biomarker [23,26].

Lactate dehydrogenase is an enzyme that catalyzes the reversible conversion of pyruvate to lactate, playing a key role in maintaining levels of NAD⁺ for glycolysis. In cancer cells, LDH contributes to the "Warburg effect," a metabolic shift where glucose is preferentially metabolized to lactate even in the presence of oxygen, supporting rapid tumor growth and survival [27-31]. Mitochondrial dysfunction in tumor cells leads to increased membrane permeability and the release of intracellular enzymes, including LDH, which correlates with abnormal apoptotic signaling [32,33]. Elevated serum LDH levels in serum thus reflect altered cellular metabolism and progression of




Figure (2): The reaction catalyzed by lactate dehydrogenase (LDH).

tumor, highlighting its value as a diagnostic and prognostic biomarker in breast cancer.

2. Materials and Methods

2.1. Study subjects

This cross-sectional study included 150 Iraqi women, aged 25–60 years, who were divided into two groups. The first group, the control group, included 50 women with no history of cancer or any other chronic disease. The second group included 100 women diagnosed with breast cancer, and were enrolled at the beginning of their chemotherapy and before undergoing radiotherapy. Diagnosis was confirmed through medical record review. Patients with chronic diseases, such as diabetes mellitus, hypertension, ischemic heart disease, and thyroid disorders, as well as smokers or alcohol consumers, were excluded from the study. Verbal informed consent was obtained from all participants prior to sample collection, and the study was approved by the Ethics Committee of the Al Amal National Cancer Hospital, Medical City, Baghdad, Iraq.

2.2. Specimen collection and preparation

Sample collection was approved by the management of Al Amal National Cancer Hospital, Baghdad, Iraq, under the direction of Dr. Mussab Kadhim Al-Aboodi. Blood samples (5 mL) were collected from each participant and placed in gel tubes. Samples from breast cancer patients were collected at different time points relative to chemotherapy, such as prior to starting of chemotherapy, after the first session, and after two to three sessions. Tubes were kept at room temperature until coagulation, and then centrifuged at 3,000 rpm for 15 min. The resulting sera were transferred into Eppendorf tubes and stored at low temperatures until analysis.

All biochemical analyses, such as LDH, alanine aminotransferase (ALT), aspartate aminotransferase (AST), alkaline phosphatase (ALP), prolactin (PRL), electrolytes, and kidney function tests, were performed using automated analyzers available in the hospital laboratory. Data distribution was tested using Shapiro-Wilk and Kolmogorov-Smirnov tests, and multifactorial analysis was used to control for potential confounding variables, such as age, disease stage, and treatment type. Statistical significance was set at $P < 0.05$.

2.3. Statistical analysis

All statistical analyses were performed using GraphPad Prism version 8.0.2. Data distribution was first tested for normality using Shapiro-Wilk and Kolmogorov-Smirnov tests. Continuous variables were presented as mean $\pm$ standard deviation (SD). The unpaired t-test was used to compare normally distributed data between the study groups. Multifactorial analysis was applied to control potential confounding variables, viz. age, disease stage, and treatment type. Differences with $P < 0.05$ were considered statistically significant [34].

3. Results and Discussion

3.1. Evaluation of serum lactate dehydrogenase

Lactate dehydrogenase enzyme plays an important role in glycolysis by converting pyruvate to lactate at the end of this process. The release of high levels of serum LDH into the blood is related to several diseases, such as cancers causes destruction the tissue [35]. The serum LDH levels in postoperative patients of breast cancer and controls are shown in Table 1 and Figure 5, which explained that the serum LDH levels in cancer patients were significantly high (352.28 ± 67.4) with $P < 0.0001$, compared to controls (120 ± 27.34). The LDH activity in breast cancer patients was higher, compared to that in controls, and this could be due to increased cell growth and cell migration compared to normal cells. Additionally, cancer cells have high energy demands and rely on aerobic glycolysis (Warburg effect), mitochondrial dysfunction, increased cellular permeability, and chemotherapy-induced inflammation, all of which contribute to elevated LDH levels in the serum. The cancer cells require excess of energy so the LDH can be considered an alternative to meet metabolic and aerobic glycolytic needs of cancer cells. In breast cancer patients, higher levels of LDH activity can give a prognostic indicator for the early warning of metastasis or recurrence [36,37].

3.2. Evaluation of serum urea & creatinine levels

The urea cycle reactions are important for removing free ammonia by hepatic detoxification and expelling it through the urethra; hence, urea levels are particularly important in patients exposed to diseases or treatments that can impair kidney function [38]. In this work, both urea and creatinine levels were evaluated in breast cancer patients, and the results shown in Table 2 and Figure 6 illustrate significantly higher levels of urea in breast cancer patients (41.76 ± 5.76), compared to controls (20.2 ± 3.61 , $P < 0.0001$).




Figure (3): Activity levels of serum lactate dehydrogenase (LDH) in breast cancer patients and healthy controls.

Table (1): Serum LDH levels in breast cancer patients and healthy controls.

Group	LDH (mean $\pm$ SD)	P value
Controls	120 $\pm$ 27.34	<0.0001
Patients	352.28 $\pm$ 67.4	

Creatinine levels were also elevated (0.822 ± 0.16 in cancer patients vs. 0.506 ± 0.1 in controls, $P < 0.0001$).

The increase in urea levels in breast cancer patients may be attributed not only to tissue destruction and nucleic acid turnover but also to chemotherapy-induced renal stress, inflammation, and altered metabolism, which affect nitrogen balance. The levels of creatinine in serum play an important role in evaluating glomerular filtration rate and kidney function efficiency. Chemotherapeutic drugs can impair kidney function directly by damaging renal cells or indirectly through vascular and tubular effects, contributing to elevated serum urea and creatinine levels in cancer patients [39–41].

3.3. Evaluation of liver enzymes in serum (AST, ALT, and ALP)

Both serum AST and serum ALT, in addition to (ALP), were evaluated in breast cancer patients and healthy controls, and the results are shown in Table 3 and Figure 7. ALP is a nonspecific enzyme involved in hydrolyzing phosphate in alkaline medium, with activity reflecting presence of isoenzymes in the bone, liver, kidney, and intestinal lining [42]. ALT and AST are liver-related enzymes, and their serum levels increase in various diseases, including liver damage, viral hepatitis, myocardial infarction, bile duct obstruction, acute pancreatitis, and anemia [43].

The levels of ALP in cancer patients (113.64 ± 49.50 , $P = 0.0630$) did not show a statistically significant difference, compared to controls (82.1 ± 22.3), suggesting that early-stage breast cancer or non-hepatic factors may not markedly affect ALP levels. Increase in ALP is generally observed in advanced stages of breast cancer and may serve to distinguish between stages or indicate metastasis [44,45].

The ALT activity was significantly higher in breast cancer patients (61.68 ± 30.5), compared to controls (18.1 ± 4.4 , $P < 0.0001$), and AST was also elevated (39.52 ± 22.5 vs. 20.6 ± 5.6 , $P = 0.0140$). Both ALT and AST catalyze the conversion of alanine to pyruvate and aspartate to oxaloacetate, respectively, and elevated levels reflect liver stress or damage, which may be secondary to tumor invasion, chemotherapy effects, or systemic inflammation. Previous studies report higher ALT and AST in malignant versus benign breast cancer patients, supporting their use as indicators of compromise of liver and renal functioning in progression of cancer [46].

3.4. Evaluation of serum electrolyte levels (Na^+ , K^+ , and Cl^-)

Kidney dysfunction is widely common in patients with malignant cancer because chemotherapy and other treatments can impair renal function, cause electrolyte disturbances, and induce immune deficiency [47]. However, many factors affect electrolyte levels, so identifying the causes is not straightforward [48].

Electrolyte levels were assessed in breast cancer patients and compared to healthy controls (Table 4 and Figure 8). Sodium levels did not differ significantly in

cancer patients (142.7 ± 6.5 , $P = 0.0663$), compared to controls (138.62 ± 3.4), suggesting that homeostatic mechanisms may maintain serum sodium despite renal stress or chemotherapy.

On the other hand, chloride levels were significantly higher in cancer patients (113.1 ± 4.5 , $P < 0.0001$), compared to controls (103.28 ± 2.9). Potassium, which plays a key role in muscle excitability and nerve functioning, was also elevated in cancer patients (4.36 ± 0.55 , $P = 0.0212$), compared to controls (3.92 ± 0.25). This may result from the release of intracellular contents from damaged tumor cells, chemotherapy-induced renal impairment, or altered hormonal regulation associated with cancer.

Serum electrolyte imbalance is clinically important for monitoring breast cancer patients, helping to assess renal injury, nephrotoxicity, and the overall prognosis. They




Figure (4): Serum levels of prolactin (PRL) in breast cancer patients and healthy controls.

Table (2): Serum levels of urea and creatinine in breast cancer patients and healthy controls.

Group	Urea (mean $\pm$ SD)	P value	Creatinine (mean $\pm$ SD)	P value
Controls	20.2 $\pm$ 3.61	<0.0001	0.506 $\pm$ 0.1	<0.0001
Patients	41.76 $\pm$ 5.76		0.822 $\pm$ 0.16	

Table (3): Serum levels of ALP, AST, and ALT in breast cancer patients and healthy controls.

Parameters	Mean $\pm$ SD		
	Controls	Patients	P value
ALP (U/L)	82.1 $\pm$ 22.3	113.64 $\pm$ 49.50	0.0630
Serum AST (U/L)	20.6 $\pm$ 5.6	39.52 $\pm$ 22.5	0.0140
Serum ALT (U/L)	18.1 $\pm$ 4.4	61.68 $\pm$ 30.5	<0.0001

Table (4): Serum levels of electrolytes in breast cancer patients and healthy controls.

Parameters	Mean $\pm$ SD		
	Controls	Patients	P value
Na^+	138.62 $\pm$ 3.4	142.7 $\pm$ 6.5	0.0663
K^+	3.92 $\pm$ 0.25	4.36 $\pm$ 0.55	0.0212
Cl^-	103.28 $\pm$ 2.9	113.1 $\pm$ 4.5	<0.0001

may also reflect systemic effects of tumor metabolism and treatment-related stress [49,50].

3.5. Evaluation of serum prolactin levels

Imbalance in hormone levels is associated with the development of various diseases. Recently, studies have focused on serum PRL in the prognosis of breast cancer. Many studies have shown a possible relationship between breast cancer and PRL hormone. PRL has cytokine-like properties; its receptor binds to the cell surface and cytoplasm, activating the Jak2/Stat5 signaling pathway [51,52].

Table 5 and Figure 9 show that breast cancer patients had significantly higher PRL levels (32.45 ± 7.0 , $P < 0.0001$), compared to controls (7.186 ± 2.5). Elevated PRL may increase the risk of breast cancer, potentially by promoting cell division, reducing apoptosis, and contributing

Table (5): Serum levels of prolactin (PRL) in breast cancer patients and healthy controls.

Group	Prolactin (mean $\pm$ SD)	P value
Controls	7.186 ± 2.5	<0.0001
Patients	32.45 ± 7.0	




Figure (5): (A) Serum levels of urea in breast cancer patients and healthy controls; (B) Serum levels of creatinine in breast cancer patients and healthy controls.




Figure (6): (A) Serum levels of ALP in breast cancer patients and healthy controls; (B) Serum levels of ALT in breast cancer patients and healthy controls; (C) Serum levels of AST in breast cancer patients and healthy controls.




Figure (7): (A) Serum levels of Na⁺ in breast cancer patients and healthy control; (B) Serum levels of Cl⁻ in breast cancer patients and healthy controls; (C) Serum levels of K⁺ in breast cancer patients and healthy controls.

to resistance to cytotoxic drugs, although the exact mechanism remains unclear [53].

4. Conclusions

This study evaluates the levels of serum LDH in patients with breast cancer at different stages of chemotherapy, and demonstrates that LDH activity in cancer patients is higher, compared to that in controls. This can be due to increased cell growth and migration of cancer cells, compared to normal cells. Cancer cells require excess energy, so serum LDH serves as an alternative to meet the metabolic and aerobic glycolytic needs of cancer cells. In breast cancer patients, the higher levels of LDH activity can serve as a prognostic indicator for early warning of metastasis or recurrence. Additionally, it can be used for monitoring cancer progression and response to treatment.

Finally, this study demonstrates that elevated serum LDH levels are significantly associated with breast cancer and its progression. Serum LDH can serve as a valuable biomarker for prognosis, early detection of metastasis, and monitoring of therapeutic response, highlighting its potential role in clinical management of breast cancer patients.

Acknowledgments

The authors appreciated the efforts and assistance of the management of Al Amal National Cancer Hospital under the supervision of the hospital director, Dr. Mussab Kadhim Al-Aboudi, and the staff who facilitated the samples and collection process.

Conflict of Interest

The authors declared no conflict of interest.

Author Contributions (CRediT Taxonomy)

Contributor Role	Degree of Contribution		
	Lead	Equal	Supporting
Conceptualization		X	
Data curation		X	
Formal analysis		X	
Funding acquisition		X	
Investigation		X	
Methodology		X	
Project administration		X	
Resources		X	
Software		X	
Supervision		X	
Validation		X	
Visualization		X	
Writing-original draft		X	
Writing-review & editing		X	

References

- [1] Sinha, T., Tumors: benign and malignant. *Cancer Ther Oncol Int J*, 10(3): 1-3, 2018. <https://doi.org/10.19080/CTOIJ.2018.10.555790>.
- [2] Gomes, C.C., Recurrent driver mutations in benign tumors. *Mutat Res Rev Mutat Res*, 789(108412), 2022. <https://doi.org/10.1016/j.mrrrev.2022.108412>.
- [3] Patel, A., Benign vs malignant tumors. *Am Med Ass JAMA Oncol*, 6(9): 1488, 2020. <https://doi.org/10.1001/jamaoncol.2020.2592>.
- [4] Lee, Y.; Jee, W.H.; Whang, Y.S.; Jung, C.K.; Chung, Y.G.; Lee S.Y., Benign versus malignant soft-tissue tumors: differentiation with 3t magnetic resonance image textural analysis including diffusion-weighted imaging. *Investig Magn Reson Imaging*, 25(1): 118-128, 2021. <https://doi.org/10.13104/imri.2021.25.2.118>.
- [5] Liu, Y.; Li, H.; Zhu, Z.; Chen, C.; Zhang, X.; Jin, G.; Li, H., RSDNet: an efficient and lightweight deep learning model for benign and malignant pathology detection in breast cancer. *Digital Health*, 11: 1-11, 2025. <https://doi.org/10.1177/20552076251336286>.
- [6] Satpathi, S.; Gaurkar, S.S.; Potdukhe, A.; Wanjari, M.B., Unveiling the role of hormonal imbalance in breast cancer development: a comprehensive review. *Cureus*, 15(7): e41737, 2023. <https://doi.org/10.7759/cureus.41737>.
- [7] Giaquinto, A.N.; Sung, H.; Miller, K.D.; Kramer, J.L.; Newman, L.A.; Minihan, A.; et al., Breast Cancer Statistics, 2022. *CA Cancer J Clin*, 72(1): 524-541, 2022. <https://doi.org/10.3322/caac.21754>.
- [8] Lina, L., Signs and symptoms of breast cancer. *Eur J Clin Oncol*, 3(2): 1, 2021.
- [9] Prusty, R.K.; Begum, S.; Patil, A.; Naik, D.; Pimple, S.; Mishra, G., Knowledge of symptoms and risk factors of breast cancer among women: a community based study in a low socio-economic area of Mumbai, India. *BMC Women's Health*, 20(106): 1-12, 2020. <https://doi.org/10.1186/s12905-020-00967-x>.
- [10] Ikhuoria, E.B.; Bach, C., Introduction to breast carcinogenesis symptoms, risks factors, treatment and management. *Eur J Eng Res Sci*, 3(7): 58-66, 2018. <https://doi.org/10.24018/ejeng.2018.3.7.745>.
- [11] Hessari, P.M.; Montazeri, A., Health-related quality of life in breast cancer patients: review of reviews from 2008 to 2018. *Health Qual Life Outcomes*, 18(1): 1-25, 2020. <https://doi.org/10.1186/s12955-020-01591-x>.
- [12] Lips, E.H.; Kumar, T.; Megalios, A.; Visser, L.; Sheinman, M.; Fortunato, A.; et al., Genomic analysis defines clonal relationships of ductal carcinoma in situ and recurrent invasive breast cancer. *Nat Genet*, 54(1): 850-860, 2022. <https://doi.org/10.1038/s41588-022-01082-3>.
- [13] Nargis, H.F.; Nawaz, H.; Ditta, A.; Mahmood, T.; Majeed, M.I.; Rashid, N.; et al., Raman Spectroscopy of Blood Plasma Samples from breast Cancer Patients at Different Stages. *Technological University, Dublin, UK*, 2019; pp. 1-24. <https://doi.org/10.1016/j.saa.2019.117210>.
- [14] Alwan, N.A.; Kerr, D.; Al-Okati, D.; Pezella, F.; Tawfeeq, F.N., Comparative study on the clinicopathological profiles of breast cancer among Iraqi and British patients. *Open Public Health J*, 11(1): 177-191, 2018. <https://doi.org/10.2174/1874944501811010177>.
- [15] Hassan, M.S.U.; Ansari, J.; Spooner, D.; Hussain, S.A., Chemotherapy for breast cancer (review). *Oncol Rep*, 24(1): 1121-1131, 2010. <https://doi.org/10.3892/or.00000963>.
- [16] Anampa, J.; Makower, D.; Sparano, J.A., Progress in adjuvant chemotherapy for breast cancer: an overview. *BMC*

- Med. 13(195): 1-13, 2015. <https://doi.org/10.1186/s12916-015-0439-8>.
- [17] Waks, A.G.; Winer, E.P. Breast cancer treatment a review. *JAMA*. 321(3): 288-300, 2019. <https://doi.org/10.1001/jama.2018.19323>.
- [18] Colomer, R.; Lopez, I.A.; Albanell, J.; Caballero, T.G.; Cluvelas, E.; Cortes, J.; et al. Biomarkers in breast cancer: a consensus statement by the Spanish Society of Medical Oncology and the Spanish Society of Pathology. *Clin Transl Oncol*. 20: 815-826, 2018 (Springer). <https://doi.org/10.1007/s12094-017-1800-5>.
- [19] Patani, N.; Martin, L.A.; Dowsett, M. Biomarkers for the clinical management of breast cancer: international perspective. *Int J Cancer*. 133(1): 1-13, 2013. <http://doi.org/10.1002/ijc.27997>.
- [20] Adnan, S.F.; Al-Abady, Z.N., Investigation the role of lactate dehydrogenase, caspase and the oxidative stress levels in breast cancer patients. *J Biomed Biochem*. 3(4), 2024. <https://doi.org/10.57238/jbb.2024.7432.1120>.
- [21] Khan, A.A.; Allemaleem, K.S.; Alhumaydhi, F.A.; Gowder, S.; Rahmani A.H., The biochemical and clinical perspectives of lactate dehydrogenase: an enzyme of active metabolism, endocrine, metabolic & immune disorders. *Drug Targets*. 20(1): 1-14, 2020. <https://doi.org/10.2174/1871530320666191230141110>.
- [22] Chaari, A.; Al-Ali, D.; Roach, J., Biochemistry course-based undergraduate research experience: purification, characterization, and identification of an unknown lactate dehydrogenase isoenzyme. *Biochem Mol Biol Educ*. 48(1): 369-380, 2020. <https://doi.org/10.1002/bmb.21363>.
- [23] Kumar, P.; Nagarajan, A.; Uchil, P.D. Analysis of Cell Viability by the Lactate Dehydrogenase Assay. Cold Spring Harbor Laboratory Press, Woodbury, NY; 2018; pp. 465-468. <https://doi.org/10.1101/pdb.prot095497>.
- [24] Klein, R.; Nagy, O.; Tothovam, C.; Frederika, C., Clinical and diagnostic significance of lactate dehydrogenase and its isoenzymes in animals. *Vet Med Int*. 2020(5346483): 1-11, 2020. <https://doi.org/10.1155/2020/5346483>.
- [25] Cai, X.; Wang, T.; Xie, L., Lactate dehydrogenase is associated with flow mediated dilation in hypertensive patients. *Sci Rep*. 13(768): 1-9, 2023. <https://doi.org/10.1038/s41598-023-27614-3>.
- [26] Valvona, C.J.; Fillmore, H.L.; Nunn, P.B.; Pilkington, G.J., The regulation and function of lactate dehydrogenase A: therapeutic potential in brain tumor. *Brain Pathol*. 1(26): 3-17, 2016. <https://doi.org/10.1111/bpa.12299>.
- [27] Farhana, A.; Lappin, S.L., Biochemistry, lactate dehydrogenase. StatPearls [Internet]. Treasure Island, FL: StatPearls Publishing; 2026; pp. 1-8.
- [28] Forkasiewicz, A.; Doruciak, M.; Stach, K.; Szelachowski, P.; Tabola, R.; Augoff, K., The usefulness of lactate dehydrogenase measurements in current oncological practice. *Cell Mol Biol Lett*. 25(35): 1-14, 2020. <https://doi.org/10.1186/s11658-020-00228-7>.
- [29] Zhou, Y.; Niu, W.; Luo, Y.; Li, H.; Xie, Y.; Wang, H.; et al., p53/Lactate dehydrogenase A axis negatively regulates aerobic glycolysis and tumor progression in breast cancer expressing wild-type p53. *Cancer Sci*. 110(1): 939-949, 2019. <https://doi.org/10.1111/cas.13928>.
- [30] Zhou, Y.; Qi, M.; Yang M., Current status and future perspectives of lactate dehydrogenase detection and medical implications: a review. *Biosensors*. 12(1145): 1-17, 2022. <https://doi.org/10.3390/bios12121145>.
- [31] Fukushi, A.; Kim, H.D.; Chang, Y.C.; Kim, C.H., Revisited metabolic control and reprogramming cancers by means of the Warburg effect in tumor cells. *Int J Mol Sci*. 23(1): 1-21, 2022. <https://doi.org/10.3390/ijms231710037>.
- [32] Jurisic, V.; Radenkovic, S.; Konjevic, G., The actual role of LDH as tumor marker, biochemical and clinical aspects, advances in cancer biomarkers. *Adv Exp Med Biol*. B67(1): 115-124, 2015. https://doi.org/10.1007/978-94-017-7215-0_8.
- [33] Zaib, S.; Hayyat, A.; Ali, N.; Gul, A.; Naveed, M.; Khan, I., Role of mitochondrial membrane potential and lactate dehydrogenase A in apoptosis. *Anti-Cancer Agents Med Chem*. 22(11): 2048-2062, 2022. <https://doi.org/10.2174/1871520621666211126090906>.
- [34] Glover, T.; Mitchell, K., An Introduction to Biostatistics, Waveland Press, Long Grove, IL; 2008.
- [35] Liu, D.; Wang, D.; Wu, C.; Zhang, L.; Mei, Q.; Hu, G.; et al., Prognostic significance of serum lactate dehydrogenase in patients with breast cancer: a meta-analysis. *Cancer Manag Res*. 11(1): 3611-3619, 2019. <https://doi.org/10.2147/CMAR.S199260>.
- [36] Kasht, G.J.; Abed, B.A.; Farhan, L.O.; Salman, I.N.; Dawood, A.S., A comparative study to determine LDH enzyme levels in serum samples of women with breast cancer and women with breast cancer and type 2 diabetes mellitus. *J Med Chem Sci*. 6(1): 883-890, 2023. <https://doi.org/10.26655/JMCHMSCI.2023.4.20>.
- [37] Basnyat, A.S.; Jha, A.; Pathak, R.; Shrestha, B.G., Study of serum lactate dehydrogenase and gamma-glutamyl transpeptidase in breast cancer patients receiving chemotherapy. *J Trop Life Sci*. 7(2): 128-132, 2017. <https://doi.org/10.11594/jtla.07.02.07>.
- [38] Attallah, B.M.; Al-Ani, W.Y.; Alhoaisa, N.S., Total antioxidant, renal function test in Iraqi women with breast cancer. *World Bull Public Health*. 16(1): 158-161, 2022.
- [39] Kebede, T.; Melak, T.; Ibn-Sina, A.; Fasil, A., Assessment of serum uric acid, urea, and glucose levels and associated factors among breast cancer patients attending a tertiary hospital in Bahirdar, Ethiopia: a comparative cross-sectional study. *Ethiop J Health Sci*. 32(6): 1183-1192, 2022. <https://doi.org/10.4314/ejhs.v32i6.16>.
- [40] Prabhu, S.N.; Mukhopadhyay, S.C.; Gooneratne, C.; Davidson, A.S.; Liu, G., Interdigital sensing system for detection of levels of creatinine from the samples. *Springer Nat*. 36(1): 267-309, 2021. https://doi.org/10.1007/978-3-030-62684-6_11.
- [41] Parker, A.L.; Toulabi, L.; Oike, T.; Kanke, Y.; Patel, D.; Tada T.; et al., Creatine riboside is a cancer cell-derived metabolite associated with arginine auxotrophy. *J Clin Invest*. 132(14): 1-17, 2022. <https://doi.org/10.1172/JCI157410>.
- [42] Jiang, C.; Hu, F.; Xia, X.; Guo X., Prognostic value of alkaline phosphatase and bone-specific alkaline phosphatase in breast cancer: a systematic review and meta-analysis. *Int J Biol Markers*. 38(1): 25-36, 2023. <https://doi.org/10.1177/03936155231154662>.
- [43] Al-Mashhadani, Z.I.; Mukhlis, A.J.; Al-Faraji, A.S., Estimation of ALP, GPT and GOT activities in Iraqi patients female with breast cancer. *Ibn Al-Haitham J Pure Appl Sci*. 25(1): 1-10, 2012.
- [44] Long, Z.Q.; Hua, X.; Zhang, W.; Lv, S.; Deng, J.; Guo, L.; et al., Prognostic impact of the pretreatment albumin to alkaline phosphatase ratio for nonmetastatic breast cancer patients. *Cancer Manag Res*. 11(1): 4809-4814, 2019. <https://doi.org/10.2147/CMAR.S200759>.
- [45] Singh, A.K.; Pandey, A.; Tewari, M.; Kumar, R.; Sharma, A.; Singh, K.A.; et al., Advanced stage of breast cancer host alkaline phosphatase activity: risk factor for females in India. *Biotech*. 3(1): 517-520, 2013. <https://doi.org/10.1007/s11205-012-0113-1>.
- [46] Yadav, C.P.; Kaushal, V.; Beniwal P., Evaluation of serum biochemical profile of breast cancer patients. *Int J Med Res Health Sci*. 5(7): 1-7, 2016.

- [47] Abdulaziz, O.; Almelhadi, M., Dysregulation of serum electrolytes is detected in untreated cancer patients in Taif city. *Ann Cancer Res Ther*, 29(2): 126-130, 2021. <https://doi.org/10.4993/acrt.29.126>.
- [48] Alem, A.; Edae, C.K.; Wabalo, E.K.; Tareke, A.A.; Bedanie, A.A.; Reta, W.; et al., Factors influencing the occurrence of electrolyte disorders in cancer patients. *SAGE Open Med*, 9(1): 1-7, 2021. <https://doi.org/10.1177/20503121211052861>.
- [49] Yadav, A.S.; Khodke, P.V., Status of serum electrolytes in cancer patients. *Int J Basic Appl Med Sci*, 5(1): 208-211, 2015.
- [50] Kumari, R.; Siddiqui, A.A.; Zubair, T.; Fatima, S.; Shaheer, A.; Imtiaz, S.; et al., Levels of electrolyte in cancer patients; a prospective study focusing on the variations before and after therapy. *Palliat Med Care*, 5(4): 1-4, 2018. <https://doi.org/10.15226/2374-8362/5/4/00168>.
- [51] Ali, J.K.; Hassan, S.H.; Merzah, M.A., Prolactin serum levels and breast cancer; relationships with hematological factors among cases in Karbala province, Iraq. *Med J Babylon*, 15(2): 178-181, 2018.
- [52] Kolomiets, O.; Yazykov, O.; Piddubnyi, A.; Lyndin, M.; Lukavenko, I.; Andryushchenko, V.; et al, The expression of prolactin receptors in benign breast tumors is not associated with serum prolactin level. *J Clin Med*, 10(1): 1-10. <https://doi.org/10.3390/jcm10245866>

A Study of Lactate Dehydrogenase (LDH) Levels and Other Biomarkers in Breast Cancer Patients Undergoing Chemotherapy/ Implications for Prognosis and Monitoring

GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/100

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8