

## SURAT TUGAS

Nomor: 4121b/Usakti/FK/01.B/IX/2024

- Dasar : 1. Pedoman Evaluasi, Kinerja dan Rekam Jejak Dosen Dalam Pelaksanaan Tridharma Perguruan tinggi Fakultas Kedokteran Universitas Trisakti.  
2. Rencana Strategis Pelaksanaan Tridharma Perguruan Tinggi Fakultas Kedokteran Universitas Trisakti dalam kegiatan penelitian.

Dekan Fakultas Kedokteran Universitas Trisakti


### MENUGASKAN



- Kepada : dr. Noviani Prasetyaningsih, Sp.M  
NIK/Usakti : 2791/Usakti  
NIDN / NIDK : 0313116301  
Untuk : melaksanakan kegiatan penelitian  
Waktu : Semester Ganjil dan Genap Tahun Akademik 2024/2025  
Tempat : Fakultas Kedokteran Universitas Trisakti.

Demikian surat tugas ini untuk dilaksanakan dengan seksama dan penuh tanggung jawab.

Apabila dikemudian hari terdapat kekeliruan atau kesalahan dalam surat tugas ini, akan diubah dan diperbaiki sebagaimana mestinya.

Jakarta, 2 September 2024

Dekan  
  
Dr. dr. Yenny, Sp.FK  
NIK: 2613/ Usakti

WD I	KTU
	

## Editorial

Screening T4 and TSH in Early Detection of Congenital Hypothyroidism in Newborns:  
What's the Dilemma?  
*Yasmine Mashabi*

## Original Article

CD68 Expression on Macrophages as Anti-Inflammatory Effect of  
Tamarillo (*Solanum betaceum* Cav.) Fruit Peel Ethanol Extract (Study on Carrageenan-  
Induced Buccal Mucosa of Rats)  
*Jelita Febrilia Bindaputri, Janti Sudiono*

Relationship Between Blood Lead (Pb) Levels and Hypertension in Motorcycle Taxi  
Drivers  
*Julian Chendrasari, Indah Widya Lestari, Florinda Ilona et al*

The association between Stress Levels and Social Support in Mothers Regarding  
Exclusive Breastfeeding in Samarinda  
*Ratih Wirapuspita Wisnuwardani, Nurul Afiah, Siti A'isyah et al*

The Effect of Sleep Patterns to Incident of Hypertension: A Case Control Study of  
Fishermen on the Puger Coast, Jember District  
*Nur Fitri Widya Astuti, Nazilatul Wahyuni Munawaroh*

Patient Characteristics Correlation with Cost of Hospitalisation in Ischemic Stroke  
Geriatric Patient  
*Dhanang Prawira Nugraha, Martanty Aditya*

The Relationship of Flat Foot to Agility in Children Aged 7-10 Years  
*Nuryani Sidarta, Ririn Afyora*

The Correlation between Laboratory Metabolic Profile and Blood Pressure  
*Diana Samara, Magdalena Wartono, Adrianus Kosasih*

Relationship Between Pterygium and Dry Eye Syndrome Among Delivery Motorbike  
Drivers  
*Erlani Kartadinata, Husnun Amalia, Anggraeni Adiwardhani et al*

Relationship between Blood Magnesium Level, Physical Fitness and Stress Level in  
Online Driver  
*Fransisca Chondro, Eveline Margo, Astri Handayani et al*

## Case Report

Painless Placental Abruption with 80% Retroplacental Bleeding: Case Report  
*Atut Cichih Mayasari, Nugroho Abikusno, Laksmi Maharani et al*

The Challenges in Treating Obesity Patients with Major Depressive Disorder (MDD)  
Treatment: a Case Report  
*Erita Istriana, Verawati Sudarma*

## Review Article

Estradiol Towards Sepsis  
*Agustina Br. Haloho, Ramzi Amin, Mgs. Irsan Saleh et al*

Review: an Overview of Neurodegenerative Diseases: Huntington, Alzheimer, and  
Parkinson  
*Ina Karlina, Eka Fitri Siti Andriyani, Arini Dian Pratiwi et al*

Role Of Exercise Intensity in Skeletal Muscle Hypertrophy  
*Nur Ayu Virginia Irawati, Nova Sylviana, Leonardo Lubis*

# Dewan Redaksi



## **Ketua Penyunting (Editor-in-Chief)**

Dr. dr. Husnun Amalia, Sp.M  
Departemen Ilmu Penyakit Mata, Fakultas Kedokteran  
Universitas Trisakti, Indonesia

## **Wakil Ketua Penyunting (Deputy Editor-in-Chief)**

Dr. Drs. ML. Edy Parwanto, M.Biomed  
Departemen Biologi Kedokteran, Fakultas Kedokteran  
Universitas Trisakti, Indonesia

## **Penyunting Ahli (Associate Editor)**

dr. Nany Hairunisa, MCHSc  
Departemen Ilmu Kedokteran Kerja, Fakultas Kedokteran  
Universitas Trisakti, Indonesia

## **Dewan Penyunting (Editorial Boards)**

Prof. Dr. dr. Adi Hidayat, MS (Indonesia)  
Dr. dr. Yenny, Sp.FK (Indonesia)  
dr. Laksmi Maharani, Sp.OG (Indonesia)  
dr. Monica Dwi Hartanti, M.Biomed, PhD (Indonesia)  
Dr. dr. Raditya Wratsangka, Sp.O.G, Subsp. Obginsos (Indonesia)  
Dr. Siti Sugih Hartiningsih, S.Si, M.Kes (Indonesia)  
dr. Dito Anurogo, M.Sc (Indonesia)  
Prof. Dr. Emad Yousif (Irak)

## **Editor Produksi**

Afton Muhandis, S.I.Kom

## **Alamat Korespondensi**

Fakultas Kedokteran Universitas Trisakti  
Jalan Kyai Tapa Np. 260 (Kampus B) Grogol, Jakarta 11440  
Telp. 021-5672731 ext. 2502 | Fax. 021-5660706  
[www.jbiomedkes.org](http://www.jbiomedkes.org) | E-mail: [jbiomedkes@trisakti.ac.id](mailto:jbiomedkes@trisakti.ac.id)

## **Penerbit**

Fakultas Kedokteran Universitas Trisakti

# Petunjuk Penulisan

## Format penyusunan manuskrip

Manuskrip diketik pada kertas berukuran A4 (210 x 297 mm) dengan batas tepi 254 mm (*margin Normal*), huruf diketik dengan tipe huruf (*font*) *Times New Roman*, besar huruf (*font size*) 12 point dengan menggunakan spasi rangkap 2 (*double space*). Setiap bagian dari manuskrip dimulai pada halaman baru dengan urutan sebagai berikut: halaman judul, abstrak dan kata kunci (*keywords*), teks keseluruhan, ucapan terima kasih, daftar pustaka, tabel dan gambar (setiap tabel dan gambar pada halaman terpisah). Nomor halaman dicantumkan secara berurutan dimulai dari halaman judul pada sudut sebelah kanan bawah. Manuskrip sebaiknya ditulis maksimal 16 halaman.

## Halaman judul

Halaman judul mencakup: a) judul manuskrip yang dibuat sesingkat mungkin, spesifik informatif dan ringkasan judul tidak lebih dari 40 karakter (hitung huruf dan spasi) yang dicantumkan dibawah judul, b) nama penulis disusun berurutan dengan nama mahasiswa sebagai pengarang pertama, diikuti oleh Pembimbing sebagai pengarang kedua. Nama penulis ditulis lengkap tanpa gelar dan dicantumkan seperti aslinya, tidak dibalik seperti pada daftar pustaka dan sitasi, c) alamat setiap penulis, nama departemen dan lembaga afiliasi penulis, d) nama dan alamat penulis untuk korespondensi serta nomor telepon, nomor faksimili, alamat email. Judul penelitian dibuat jelas, singkat, spesifik, informatif, dan sesuai dengan topik manuskrip. Jumlah kata tidak lebih dari 12 kata agar mudah dan cepat dipahami pembaca.

## Abstrak dan kata kunci

Abstrak berjumlah 200-250 kata ditulis dalam bahasa Indonesia dan Inggris. Abstrak berisikan latar belakang termasuk tujuan penelitian, metode, hasil, dan kesimpulan. Kata kunci dicantumkan di bawah abstrak pada halaman yang sama sebanyak 4-6 kata. Bagian abstrak merupakan ringkasan dari isi makalah yang dibuat secara singkat, informatif, dengan menekankan pada aspek baru dan penting dari penelitian.

## Teks

Teks makalah manuskrip dibagi dalam beberapa bagian dengan judul sebagai berikut: ***Pendahuluan, Metode, Hasil, Pembahasan, Kesimpulan dan saran.***

## Pendahuluan

a. Latar belakang merupakan bagian yang menjelaskan alasan mengapa masalah ini penting untuk diteliti. Bagian ini memuat penjelasan mengapa masalah itu dipandang menarik, penting, dan perlu diteliti untuk mencari pemecahannya. Penjelasan dapat diperoleh dari penelusuran pustaka yang berkaitan erat dengan

masalah yang diteliti.

b. Keaslian penelitian dikemukakan dengan menunjukkan bahwa masalah yang dihadapi belum pernah dipecahkan oleh peneliti terdahulu atau dinyatakan dengan tegas perbedaan penelitian ini dengan penelitian terdahulu.

c. Tujuan penelitian yang menjelaskan hasil yang akan dicapai.

## Metode

Metode penelitian berisi uraian terpadu dan sistematis mengenai bagaimana penelitian akan dilaksanakan. Metode terdiri dari :

a. Desain

b. Populasi / sampel (subjek) penelitian

Diuraikan kriteria inklusi dan eksklusi subjek penelitian, cara pemilihan sampel (subjek penelitian) secara random atau non-random, serta besar sampel yang akan di pilih. Teknik pemilihan sampel harus dijelaskan secara rinci. Bila perlu dibuat alur pemilihan sampel.

c. Bahan dan alat serta pengukuran

Bahan dan alat yang harus disajikan pada laporan terbatas pada bahan (materi) dan alat utama yang diperlukan untuk penelitian dan harus disebutkan spesifikasinya. Prosedur pengukuran perlu dijelaskan sesuai dengan tahapan yang dilakukan.

d. Alur kerja penelitian

Jalannya penelitian perlu dijelaskan mengenai jenis pendekatan yang dipakai untuk mendapatkan data, melalui pendekatan laboratorium, klinik, komunitas, observasi, dll.

e. Analisis data

Perlu dijelaskan jenis teknik statistik yang digunakan untuk menjawab masalah dan mencapai tujuan penelitian. Data yang diperoleh dapat dianalisis menggunakan teknik statistik secara parametrik dan non-parametrik.

## Hasil

Suatu hasil penelitian hendaknya disajikan dengan jelas, logis, runut, sehingga mudah untuk dimengerti. Hasil penelitian sebaiknya ditampilkan selain dalam bentuk narasi dapat pula berupa gambar, tabel, foto, dan grafik sehingga memudahkan untuk dipahami. Hasil dan interpretasi analisis statistik dituliskan secara jelas dalam uraian hasil penelitian.

Pada tahap awal disajikan distribusi karakteristik subjek penelitian, yang biasanya dibuat pada sebuah tabel. Kemudian disajikan temuan penting yang diperoleh, kalau cukup banyak sebaiknya pada sebuah tabel. Bila terbatas misalkan hanya satu atau dua temuan cukup dalam bentuk narasi/teks.

Tabel, bagan/gambar, grafik dibuat dengan jelas, diberi nomor urut serta keterangan yang jelas. Keterangan

tabel diletakkan di atas tabel dan keterangan gambar diletakkan di bawah gambar. Maksimal tabel dan gambar 5. Semua tabel, grafik dan gambar diberi nomor dan keterangan yang jelas. Setiap tabel dianalisis dan diinterpretasi secara sistematis, dan hasilnya ditulis di bawah tabel tersebut. Perhitungan statistik detail tidak perlu ditulis dalam bagian hasil ini. Bila perhitungan statistik dianggap perlu ditulis, maka sebaiknya diletakkan dalam lampiran saja.

### **Pembahasan**

Langkah awal harus diuraikan temuan penting yang diperoleh dari penelitian sesuai dengan tujuan penelitian. Kemudian bandingkan hasil penelitian yang diperoleh dengan hasil-hasil penelitian sebelumnya. Perlu dijelaskan kesesuaian dan ketidaksesuaian hasil penelitian yang didapat terhadap kerangka teori atau hasil penelitian lain yang telah dilakukan sebelumnya. Selanjutnya menggunakan teori-teori yang ada uraikan mekanisme terjadinya hasil penelitian tersebut. Bagian pembahasan juga menjelaskan mengenai kelemahan dan kelebihan penelitian yang telah dilakukan. Uraikan implikasi dari hasil penelitian yang diperoleh.

### **Kesimpulan**

Kesimpulan hendaknya dibuat dalam bentuk narasi dan menguraikan secara singkat, jelas, padat menurut urutan yang sistematis. Bagian ini memuat tentang hasil penelitian yang telah diperoleh untuk menjawab tujuan penelitian. Saran menguraikan perlunya dilakukan penelitian lebih lanjut untuk memperbaiki kelemahan/keterbatasan dari penelitian yang telah dilakukan.

### **Ucapan terima kasih**

Ditujukan kepada pihak-pihak yang memberikan bantuan dana dan dukungan antara lain dukungan dari bagian dan lembaga, para professional yang memberikan kontribusi dalam penyusunan makalah, dan untuk penguji I maupun penguji II. Pembimbing tidak perlu dicantumkan pada Ucapan Terima Kasih karena sudah dicantumkan sebagai penulis.

### **Daftar Referensi**

Daftar referensi/rujukan hanya mencantumkan rujukan yang telah digunakan dan ditulis menurut sistem Vancouver.

### **Online Submission**

Penulis dapat mengirim naskah manuskrip melalui *online submission* di *website* Jurnal Biomedika dan Kesehatan.

Langkah online submission:

1. Pastikan naskah yang akan diunggah sudah mengikuti semua petunjuk penulisan
2. Lakukan pendaftaran author di : <https://jbiomedkes.org/index.php/jbk/user/register>
3. Setelah terdaftar silakan unggah naskah manuskrip dan isi form yang terdapat di dalam website, dan ikuti langkah selanjutnya.

### **Daftar Cek Pengiriman Naskah Manuskrip**

- Naskah manuskrip belum pernah dipublikasikan sebelumnya, juga tidak dalam pengajuan ke jurnal lain.
- File manuskrip harus berformat OpenOffice, Ms. Word atau RTF dokumen, *font* 12, *Times New Roman*, *double spacing*.
- Halaman judul harus memuat jelas judul, nama lengkap penulis tanpa gelar, departemen penulis, universitas, alamat lengkap, nomor telepon dan email.
- Pelaporan data manuskrip dari penelitian yang melibatkan manusia dan hewan memerlukan persetujuan formal (kaji etik) oleh dewan peninjau atau komisi etik institusi yang bersangkutan.
- Daftar rujukan memuat semua rujukan yang terdapat di dalam manuskrip dan ditulis sesuai urutan pengutipannya menggunakan sistem Vancouver.

# Daftar Isi



*Jurnal Biomedika dan Kesehatan - Vol. 7 No. 1 Maret 2024*

## **Editorial**

- Screening T4 and TSH in Early Detection of Congenital Hypothyroidism in Newborns: What's the Dilemma?** 1  
*Yasmine Mashabi*

## **Original Article**

- CD68 Expression on Macrophages as Anti-Inflammatory Effect of Tamarillo (*Solanum betaceum* Cav.) Fruit Peel Ethanol Extract (Study on Carrageenan-Induced Buccal Mucosa of Rats)** 6  
*Jelita Febrilia Bindaputri, Janti Sudiono*

- Relationship Between Blood Lead (Pb) Levels and Hypertension in Motorcycle Taxi Drivers** 17  
*Julian Chendrasari, Indah Widya Lestari, Florinda Ilona et al*

- The association between Stress Levels and Social Support in Mothers Regarding Exclusive Breastfeeding in Samarinda** 24  
*Ratih Wirapuspita Wisnuwardani, Nurul Afiah, Siti A'isyah et al*

- The Effect of Sleep Patterns to Incident of Hypertension: A Case-Control Study of Fishermen on the Puger Coast, Jember District** 34  
*Nur Fitri Widya Astuti, Nazilatul Wahyuni Munawaroh*

- Patient Characteristics Correlation with Cost of Hospitalisation in Ischemic Stroke Geriatric Patient** 44  
*Dhanang Prawira Nugraha, Martanty Aditya*

- The Relationship of Flat Foot to Agility in Children Aged 7-10 Years** 52  
*Nuryani Sidarta, Ririn Afyora*

- The Correlation between Laboratory Metabolic Profile and Blood Pressure** 61  
*Diana Samara, Magdalena Wartono, Adrianus Kosasih*

- Relationship Between Pterygium and Dry Eye Syndrome Among Delivery Motorbike Drivers** 71  
*Erlani Kartadinata, Husnun Amalia, Anggraeni Adiwardhani et al*

- Relationship between Blood Magnesium Level, Physical Fitness and Stress Level in Online Driver** 82  
*Fransisca Chondro, Eveline Margo, Astri Handayani et al*

<b>Case Report</b>	
<b>Painless Placental Abruption with 80% Retroplacental Bleeding: Case Report</b>	91
<i>Atut Cicih Mayasari, Nugroho Abikusno, Laksmi Maharani et al</i>	
<b>The Challenges in Treating Obesity Patients with Major Depressive Disorder (MDD) Treatment: a Case Report</b>	98
<i>Erita Istriana, Verawati Sudarma</i>	
<b>Review Article</b>	
<b>Estradiol Towards Sepsis</b>	
<i>Agustina Br. Haloho, Ramzi Amin, Mgs. Irsan Saleh et al</i>	104
<b>Review: an Overview of Neurodegenerative Diseases: Huntington, Alzheimer, and Parkinson</b>	113
<i>Ina Karlina, Eka Fitri Siti Andriyani, Arini Dian Pratiwi et al</i>	
<b>Role Of Exercise Intensity in Skeletal Muscle Hypertrophy</b>	124
<i>Nur Ayu Virginia Irawati, Nova Sylviana, Leonardo Lubis</i>	

## ORIGINAL ARTICLE

# Relationship Between Pterygium and Dry Eye Syndrome Among Delivery Motorbike Drivers


## Pterigium berhubungan dengan Sindroma Mata Kering pada Pengemudi Ojek Online


Erlani Kartadinata<sup>1</sup> , Husnun Amalia<sup>1</sup>, Anggraeni Adiwardhani<sup>1</sup>, Riani Witjaksana<sup>1</sup>, Noviani Prasetyaningsih<sup>1</sup>, Eveline Margo<sup>2</sup>, Yohana<sup>3</sup>

<sup>1</sup>Departement Ophthalmology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

<sup>2</sup>Departement Physiology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

<sup>3</sup>Departement Biochemistry, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

 [erlani.kartadinata@trisakti.ac.id](mailto:erlani.kartadinata@trisakti.ac.id)

 <https://doi.org/10.18051/JBiomedKes.2024.v7.71-81>

### ABSTRACT

#### Background

Dry eye syndrome (DES) is an eye disease caused by many factors involving the ocular surface, with characteristic damage to tear layer homeostasis accompanied by ocular symptoms due to tear layer instability, hyperosmolarity, damage and inflammation of the ocular surface, and neurosensory abnormalities. Delivery motorbike drivers are often on the streets where the role of environmental factors will be able to pose a risk of experiencing DES. This study aims to assess risk factors associated with the incidence of dry eye syndrome in delivery motorbike drivers. The benefit of this research is to improve public health, especially in preventive behavior against factors associated with dry eye syndrome.

#### Methods

This study used observational analytical methods with a cross-sectional research design. The selection of samples by consequential non-random sampling amounted to 124 with the criteria of inclusion of delivery motorbike drivers, exclusion of not using artificial tears eye drops, and not use of contact lenses. Data were collected by filling out a DES risk factor questionnaire and a Tear Break Up Time (TBUT) examination to assess DES. The research data were analyzed using the Chi-Square test.

#### Results

This study's results show that subjects with dry eye syndrome are dominated by adults aged 19–44 and male sex. Subjects who experienced DES 76.6%, adult age 78.2%, male 80.6%, no pterygium 66.1%, working period 2-5 years 62.9%, length of work  $\geq$ 8 hours 94.4%, always using a closed helmet 83.9%, not using protective glasses 39.5%, coffee consumption 71.8%, active smokers 66.1%, using gadgets >6 hours 91.9%. There is no significant relationship between subjects' characteristics and DES ( $p > 0.05$ ). The incidence of DES in pterygium showed significant results ( $p = 0.001$ ).

#### Conclusions

There is a significant relationship between the incidence of pterygium and DES.

**Keywords:** Dry Eye Syndrome; Pterygium; Risk Factor; Tear Break Up Time (TBUT)

## ABSTRAK

### Latar Belakang

Sindroma mata kering (DES) adalah penyakit mata yang disebabkan oleh banyak faktor yang melibatkan permukaan okular, dengan karakteristik rusaknya homeostasis lapisan air mata disertai dengan gejala okular diakibatkan ketidak-stabilan lapisan air mata, hiperosmolaritas, kerusakan dan inflamasi pada permukaan okular, serta abnormalitas neurosensoris. Pengemudi ojek online sering berada di jalanan di mana peran faktor lingkungan akan dapat menimbulkan risiko mengalami DES. Penelitian ini bertujuan untuk menilai faktor risiko yang terkait dengan kejadian sindrom mata kering pada pengemudi sepeda motor pengiriman. Manfaat dari penelitian ini adalah untuk meningkatkan kesehatan masyarakat, terutama dalam perilaku preventif terhadap faktor-faktor yang terkait dengan sindrom mata kering.

### Metode

Penelitian ini menggunakan metode analisis observasional dengan desain penelitian cross sectional. Pemilihan sampel secara konsekuensial non-random sampling berjumlah 124 dengan kriteria inklusi pengemudi ojek online, eksklusi tidak menggunakan tetes mata air mata buatan dan tidak menggunakan lensa kontak. Data dikumpulkan melalui pengisian kuesioner faktor risiko DES dan pemeriksaan Tear Break Up Time (TBUT) untuk menilai DES. Data penelitian dianalisis menggunakan uji Chi-Square.

### Hasil

Hasil penelitian ini menunjukkan bahwa subjek yang memiliki sindrom mata kering didominasi oleh orang dewasa pada usia 19-44 tahun dan berjenis kelamin laki-laki. Subjek yang mengalami DES 76.6%, dewasa usia 78.2%, laki-laki 80.6%, tidak ada pterygium 66.1%, masa kerja 2-5 tahun 62.9%, lama kerja  $\geq$  8 jam 94.4%, selalu menggunakan helm tertutup 83.9%, tidak menggunakan kacamata pelindung 39.5%, konsumsi kopi 71.8%, perokok aktif 66.1%, menggunakan gadget  $>6$  jam 91.9%. Tidak ada hubungan yang bermakna antara karakteristik subjek dan DES ( $p > 0.05$ ). Kejadian DES pada pterygium menunjukkan hasil yang bermakna ( $p = 0.001$ ).

### Kesimpulan

Ada hubungan yang bermakna antara kejadian pterygium dan DES.

**Kata Kunci:** Sindroma Mata Kering; Pterygium; Faktor Risiko; Tear Break Up Time (TBUT)

## INTRODUCTION

Dry eye syndrome (DES) is an eye disease caused by many factors involving the ocular surface, characterized by damage to tear film homeostasis accompanied by ocular symptoms due to tear film instability, hyperosmolarity, damage, and inflammation on the ocular surface, as well as neurosensory abnormalities.<sup>1-3</sup>

Symptoms of DES vary from mild irritation, burning sensation, itching, red eyes, gritty feeling in the eyes, sore eyes, tired eyes, and in severe cases can cause vision problems.<sup>1,4,5</sup> DES is the most common eye disease. In daily practice, with a prevalence of between 5-50% of the entire world population.<sup>2</sup> The prevalence of DES in Southeast Asia is around 20-52.4%.<sup>3</sup> The prevalence of dry eyes in Indonesia in 2017 was 26.5%.<sup>4</sup>

Several risk factors that are closely related to the incidence of DES include age, gender, race, connective tissue disorders, Sjogren's syndrome, estrogen replacement therapy, androgen hormone deficiency, use of drugs (antihistamines, antidepressants, anxiolytics, and isotretinoin), use of computers, use of contact lenses, environmental conditions (pollution, low humidity).<sup>1,2</sup> Probable risk factors include diabetes, rosacea, viral infections, thyroid disease, psychiatric conditions, pterygium, low fat intake, refractive surgery, and allergic conjunctivitis.<sup>2</sup>

Dry eyes are very common in older people, around 5-30% of the elderly population, and more often affect women than men.<sup>1</sup> The disparity in prevalence according to age ranges from 8.4% for those aged less than 60 years, 5% for those aged 70-79 years. and 20% over 80 years of age.<sup>1</sup>

Complications of DES that do not receive therapy or do not respond well to therapy can result in loss of visual acuity.<sup>1,2</sup> In severe cases of DES it can affect the patient's quality of life physically, socially, psychologically, and work productivity.<sup>4</sup>

Online motorcycle taxi driving is a relatively new profession in Indonesia and this profession helps community activities this profession is very popular and the number of online motorcycle taxi drivers is increasing. This job requires online drivers to always be on the road where environmental factors can pose a risk of experiencing DES.

Research conducted by Zsafidda and Kamellisha found that motorbike taxi workers in the city of Malang who had been exposed to smoke for more than 10 years experienced severe DES.<sup>6</sup> Currently there is not much research on online drivers, especially in the field of eye health, as well as the negative impacts that can arise. As a result of this work, researchers are interested in conducting research to assess the factors related to the incidence of DES in online motorcycle taxi drivers, for prevention efforts, especially in the field of eye health.

## METHODS

This research uses an analytical observational method with a cross-sectional design. This research was carried out at FK USAKTI, Jl. Kyai Tapa- Grogol, West Jakarta, in September 2022. The population in this study was online motorcycle taxi drivers, totaling 124 respondents. Sample selection used a non-random consecutive sampling technique. The inclusion criteria are online motorcycle taxi drivers, male and female. Exclusion criteria were using contact lenses and using artificial tears and eye drops.

Data was collected by filling out a questionnaire containing the respondent's identity and risk factors for dry eye syndrome. Then an anterior eye examination and Tear Break Up Time (TBUT) examination were carried out. TBUT examination shows whether there is a disturbance in the stability of the tear film. The examination is carried out using fluorescent strips. The inspection was carried out using a cobalt blue filter spotlight. Then the patient is asked to blink 3 times, then measure the time interval required after the last eye blink until a black spot appears on the cornea using a stopwatch. The measurement result is the time it takes for the tear film to break apart when the eye blinks. Normal TBUT results are 15-30 seconds. If  $TBUT \leq 10$  seconds indicates a disruption in tear film stability.

The statistical tests used were the Chi-square test and the Fisher Exact test with a significance level of  $p < 0.05$ . Data were analyzed univariately and bivariately using the SPSS (Statistical Package for Social Science) program version 23.0. The results are presented in the form of a frequency table of respondent characteristics and risk factors associated with dry eye syndrome.

## RESULTS

Table 1. Characteristics and Risk Factors for Dry Eye Syndrome

Variable	Respondent (n)	Percentage (%)
<b>Age</b>		
Adult (19-44 y.o)	97	78.2
Pre-elderly (45-59 y.o)	26	21.0
Elderly (> 60 y.o)	1	0.8
<b>Gender</b>		
Man	100	80.6
Woman	24	19.4
<b>Dry Eye</b>		
Yes	95	76.6
No	29	23.4
<b>Pterygium</b>		
Yes	42	33.9
No	82	66.1
<b>Years of service</b>		
< 2 years	1	0.8
2-5 2-5 Years	78	62.9
> 5 years	45	36.3
<b>Length of working</b>		
< 8 hours	7	5.6
> 8 hours	117	94.4
<b>Closed helmet</b>		
No	6	4.8
Rarely	14	11.3
Always	104	83.9
<b>Protective glasses</b>		
No	49	39.5
Rarely	36	29.0
Always	39	31.5
<b>Coffee Consumption</b>		
Yes	89	71.8
No	35	28.2
<b>Active smoker</b>		
Yes	82	66.1
No	42	33.9
<b>Gadgets/day</b>		
≤ 6 hours	10	8.1
> 6 hours	114	91.9

In table 1, most respondents were male (80.6%), mature age (78.2%), incidence of dry eye syndrome (76.6%), no pterygium (66.1%), working period 2-5 years (62.9%), working time ≥ 8 hours (94.4%), always using a closed helmet (83.9%), not using protective glasses (39.5%), consuming coffee (71.8%), active smoker (66.1%), using gadgets > 6 hours (91.9%).

Table 2. Relationship between respondent characteristics and risk factors for dry eye syndrome

Variable	Dry eye syndrome				p
	Yes		No		
	n	%	n	%	
<b>Age</b>					0.462 <sup>€</sup>
Adults (19-44 years)	72	74.23	25	25.77	
Pre-elderly (25-59 years)	22	84.62	4	15.38	
Elderly (> 60 years old)	1	100.00	0	0.00	
<b>Gender</b>					0.456 <sup>€</sup>
Man	78	78.00	22	22.00	
Woman	17	70.83	7	29.17	
<b>Pterygium</b>					0.001 <sup>b*</sup>
Yes	39	92.86	3	7.14	
No	56	68.29	26	31.71	
<b>Years of service</b>					0.440 <sup>€</sup>
< 2 years	1	100.00	0	0.00	
2-5 years	57	73.08	21	26.92	
> 5 years	37	82.22	8	17.78	
<b>Working hours/day</b>					0.482 <sup>b</sup>
< 8 hours	6	85.71	1	14.29	
> 8 hours	89	76.07	28	23.93	
<b>Closed helmet</b>					0.087 <sup>€</sup>
No	6	100.00	0	0.00	
Rarely	8	57.14	6	42.86	
Always	81	77.88	23	22.12	
<b>Protective glasses</b>					0.087 <sup>€</sup>
No	39	79.59	10	20.41	
Rarely	23	63.89	13	36.11	
Always	33	84.62	6	15.38	
<b>Coffee Consumption</b>					0.072 <sup>€</sup>
Yes	72	80.90	17	19.10	
No	23	65.71	12	34.29	
<b>Active smoker</b>					0.414 <sup>€</sup>
Yes	61	74.39	21	25.61	
No	34	80.95	8	19.05	
<b>Gadgets/day</b>					0.115 <sup>b</sup>
< 6 hours	10	100.00	0	0.00	
> 6 hours	85	74.56	29	25.44	

<sup>€</sup> = chi-square test

<sup>b</sup> = Fisher Exact Test

\* = p < 0.05

Table 2 shows that more adult respondents aged 19-44 years experienced dry eye syndrome at 74.2% (72 respondents). In the statistical test results, the value of p=0.087 (p>0.05) was obtained, which means there is no significant relationship between age and dry eye syndrome. In the gender variable, the results showed that men experienced more dry eye syndrome, namely 78% (78 respondents). The statistical test results obtained a value of p=0.456 (p>0.05), which means there

is no significant relationship between gender and dry eye syndrome. Respondents who did not have pterygium experienced dry eye syndrome at 68.3% (39 respondents). The statistical test results obtained a value of  $p=0.001$  ( $p<0.05$ ), which means there is a significant relationship between pterygium and dry eye syndrome. Respondents with 2-5 years of work experienced the most dry eye syndrome, namely 73.1% (57 respondents). The statistical test results showed that  $p=0.440$  ( $p>0.05$ ), which means that there was no significant relationship between work experience and dry eye syndrome. Working hours  $\geq 8$  hours per day most frequently experienced dry eye syndrome at 76.1% (89%). The statistical test results showed that  $p=0.482$  ( $p>0.05$ ) which means there is no significant relationship between working hours  $\geq 8$  hours per day and dry eye syndrome. Respondents who always used closed helmets experienced dry eye syndrome of 77.9% (81 respondents). The statistical test results obtained  $p=0.087$  ( $p>0.05$ ), which means there is no significant relationship between using a closed helmet and dry eye syndrome. Respondents who did not use protective glasses experienced dry eyes of 79.6% (39 respondents). The statistical test results obtained  $p=0.087$  ( $p>0.05$ ), which means there is no significant relationship between using protective glasses and dry eye syndrome. 80.9% of respondents who consumed coffee had dry eye syndrome (72 respondents). The statistical test results obtained  $p=0.072$  ( $p>0.05$ ), which means there is no significant relationship between coffee consumption and dry eye syndrome. Active smoker respondents who had dry eye syndrome were 74.4% (61 respondents). The statistical test results showed  $p=0.414$  ( $p>0.05$ ), which means there is no significant relationship between active smoking and dry eye syndrome. Respondents who used gadgets  $>6$  hours per day had dry eye syndrome of 74.6% (85 respondents). The statistical test results obtained  $p=0.115$  ( $p>0.05$ ), which means there is no significant relationship between using gadgets  $>6$  hours per day and dry eye syndrome.

## DISCUSSION

The results in Table 1 show that online motorcycle taxi drivers are dominated by adult men (19-44 years). This is because adult men mostly use two-wheeled motorized vehicles and are breadwinners to meet family needs.

The results in Table 1 show that 76.6% of motorbike taxi drivers experienced dry eye syndrome. This is because in this study there were risk factors that caused dry eye syndrome, including 94.4% having a history of working more than 8 hours, and 71.8% consuming coffee, 66.1% are active smokers, and 91.9% use gadgets for more than 6 hours.

Table 2 shows that more adult respondents aged 19-44 years experienced dry eye syndrome at 74.2% (72 respondents). These results are in accordance with research conducted by Zong JY, et al. which states that there is a change in tear production with increasing age and dry eye syndrome is mostly in the younger generation (18-49 years).<sup>7</sup> In this study, 78.2% were aged 19-44 years, where at this age the use of mobile phones is increasing, especially in this era. modern today. This causes the risk of dry eye syndrome to increase at a young age. In the statistical test results, the value of  $p=0.087$  ( $p>0.05$ ) was obtained, which means there is no significant relationship between age and dry eye syndrome. This is in line with research conducted by Paulsen AJ, et al. which states that there is no significant relationship between age and dry eye syndrome with a value of  $p = 0.06$ .<sup>8</sup> Older respondents are said to be more at risk of experiencing a decrease in tear production because they have a decrease in corneal sensitivity which is related to the mechanism of dry eye. With

increasing age, apart from decreasing tear secretion, there is also an increase in tear evaporation such as atrophy of the meibomian glands. The distribution of dry eye syndrome in the age range may vary, this is because the causes of dry eye syndrome are multifactorial.<sup>9</sup>

In the gender variable, the results showed that men experienced more dry eye syndrome, namely 78% (78 respondents). The statistical test results obtained a value of  $p=0.456$  ( $p>0.05$ ), which means there is no significant relationship between gender and dry eye syndrome. These results are in line with research conducted by Gierow JP, et al. which stated that no significant relationship was found between gender and dry eye syndrome with a  $p\text{-value} > 0.05$ .<sup>9</sup> In several epidemiological studies, the prevalence of dry eye syndrome was higher in women. This is due to low estrogen in menopausal women, whereas in this study it was dominated by men at 80.6%.

Respondents who did not have pterygium experienced dry eye syndrome at 68.3% (39 respondents). The statistical test results obtained a value of  $p=0.001$  ( $p<0.05$ ), which means there is a significant relationship between pterygium and dry eye syndrome. These results are in accordance with research conducted by Manhas A, et al. which states that there is a significant relationship between pterygium and dry eye syndrome with a value of  $p<0.001$ .<sup>10</sup> Research conducted by Gupta AK also states that there is a significant relationship between pterygium and dry eye syndrome with  $p<0.05$ .<sup>11</sup> Pterygium is a condition where the tissue under the conjunctiva experiences degeneration in the form of proliferation of fibrovascular tissue that invades the cornea. The causes of pterygium include environmental factors such as ultraviolet (sun exposure) and chronic inflammation (exposure to dust and wind).<sup>12,13</sup> Tear film abnormalities are also one of the factors causing pterygium. Disorders of the stability of the tear film (tear film) are generally caused by two factors, namely chronic inflammation of the ocular surface and changes in tear dynamics.<sup>12</sup> Dry cornea and conjunctiva trigger the growth of new tissue. Pterygium causes the surface of the eyeball to become abnormal, resulting in tear film dysfunction and dry eye syndrome.<sup>10</sup> Tear film dysfunction is a precursor to pterygium or whether pterygium causes tear film dysfunction is still unclear.<sup>11,14</sup> The results of research conducted by Thatee S in 2023 stated that pterygium causes dry eye syndrome where the higher the stage/degree of the pterygium, the more severe the degree of dry eye syndrome.<sup>15</sup>

Respondents with 2-5 years of work experience experienced the most dry eye syndrome, namely 73.1% (57 respondents). The statistical test results showed that  $p=0.440$  ( $p>0.05$ ), which means that there was no significant relationship between work experience and dry eye syndrome. The results of research conducted by Paudel N, et al. stated that drivers who worked for 5 years experienced dry eye syndrome due to air pollution and there was no significant difference between the length of work of 5-10 years and 10 years.<sup>16</sup> The impact of air pollution on the surface of the eyeball occurs after a cumulative process of many years.

Working hours  $\geq 8$  hours per day most frequently experienced dry eye syndrome at 76.1% (89%). In the statistical test results, it was found that  $p=0.482$  ( $p>0.05$ ) which means there is no significant relationship between working hours  $\geq 8$  hours per day and dry eye syndrome. Respondents who always used closed helmets experienced dry eye syndrome of 77.9% (81 respondents). The statistical test results obtained  $p=0.087$  ( $p>0.05$ ), which means there is no significant relationship between using a closed helmet and dry eye syndrome. Respondents who did not use protective glasses experienced dry eyes of 79.6% (39 respondents). The statistical test results obtained

$p=0.087$  ( $p>0.05$ ), which means there is no significant relationship between using protective glasses and dry eye syndrome. Working as a driver will involve environmental exposure such as pollution, high temperatures, sun, and wind. The longer you drive, the greater the risk of exposure to environments that cause dry eye syndrome. The use of helmets and protective glasses aims to reduce exposure to this environment. Research conducted by Liang K, et al. states that exposure to accumulated air pollution is significantly related to dry eye syndrome.<sup>17</sup> The many factors that cause dry eye syndrome cause dry eye syndrome to be found in respondents who always wear helmets or protective glasses, for example, because they have had pterygium for a long time, the weather is heat, and other influencing factors.

80.9% of respondents who consumed coffee had dry eye syndrome (72 respondents). The statistical test results obtained  $p=0.072$  ( $p>0.05$ ), which means there is no significant relationship between coffee consumption and dry eye syndrome. The caffeine contained in coffee can affect tear production because it has the effect of increasing the activation of sympathetic innervation in the lacrimal gland which causes a decrease in tear secretion.<sup>18</sup> Research conducted by Juddy AU, et al. shows that there is a significant relationship between caffeine and a decrease in tear production.<sup>18</sup> Likewise, research conducted by Fayola PA shows that there is a significant relationship between coffee and dry eye syndrome with  $p = 0.000$ .<sup>19</sup> Meanwhile, research conducted by Magno M. S, et al, in 2023, states that there is no significant relationship between caffeine and dry eye syndrome.<sup>20</sup> Caffeine reaches its maximum concentration in 20-120 minutes in plasma and is eliminated from the body after 4 hours.<sup>21</sup> Caffeine stimulates the parasympathetic by increasing acetylcholine which causes activation of water secretion, proteins, electrolytes and stimulation of tear secretion. The difference is in research conducted by Judy AU, et al. It uses experimental methods in collecting data. Research subjects were asked to consume caffeinated coffee and then had their tear production measured via the Schirmer test, before and after consuming coffee.

Active smoker respondents who had dry eye syndrome were 74.4% (61 respondents). The statistical test results showed  $p=0.414$  ( $p>0.05$ ), which means there is no significant relationship between active smoking and dry eye syndrome. Cigarette smoke irritates the surface of the eyeball, if it occurs continuously it will activate T cells which cause local inflammation and activation of cytokines which block sensory signals from the lacrimal gland resulting in a decrease in tear secretion.<sup>22</sup> Cigarette smoke on the surface of the eyeball causes disruption of the lipid layer. tear film which results in increased evaporation from the tear film.<sup>23</sup> Research conducted by Ibrahim AM, stated that there was a significant relationship with increasing degrees of dry eye syndrome in smokers compared to non-smokers with  $p=0.01$ .<sup>24</sup> This is different from research conducted by Attri S, who stated that there was no significant relationship between smoking, diabetes mellitus, hypertension, and dry eye syndrome.<sup>25</sup> Likewise, research conducted by Tariq MA, et al. stated that smoking was not indicated as a risk factor for dry eye syndrome.<sup>26</sup> In this study, although many respondents were smokers, more detailed and accurate data was still needed such as how to smoke, type of cigarette, duration of smoking, comorbidities, and other influencing factors.

Respondents who used gadgets >6 hours per day had dry eye syndrome of 74.6% (85 respondents). The statistical test results obtained  $p=0.115$  ( $p>0.05$ ), which means there is no significant relationship between using gadgets >6 hours per day and dry eye syndrome. Using a digital screen for more than 6 hours can increase the risk of dry eye syndrome.<sup>27</sup> Staring at a digital

screen for a long period can reduce the frequency of blinking which fails in the distribution of tears to lubricate the surface of the eye, so that tear evaporation will increase.<sup>27</sup> In research conducted by Mohtaseb ZA, et al. shows a significant relationship between the use of digital screens and a decrease in the blink reflex and the risk of developing dry eye syndrome.<sup>28</sup> Research conducted by Tawonkasiwattanakum P, et al. stated that there was no significant relationship between gadget use and dry eye syndrome in university students and staff.<sup>29</sup> In this study, the online driver profession will often use mobile phones in carrying out their work. The length of time you use a mobile phone depends on the length of work and the drivers' activities while working. If drivers often take breaks without using their mobile phones for a while or take short naps in their free time while working, this will reduce the risk of dry eye syndrome. Another thing that influences the results of this study is that there are many risk factors for dry eye syndrome, such as hot weather, cold rooms using air conditioners, wind, and other factors when using a mobile phone.

This research has the limitation of being carried out subjectively using a multifactorial questionnaire of risk factors for dry eye and the causes of dry eye syndrome, so a more detailed questionnaire is needed for each risk factor and adding other tests to confirm the diagnosis of dry eye syndrome and increasing the number of respondents studied more.

This research implies that it is hoped that preventive and educational measures regarding dry eye syndrome and pterygium can be carried out among online motorcycle taxi drivers so that the productivity of online motorcycle taxi drivers is optimal.

## CONCLUSION

The conclusion from the analysis of this research data is that there is a relationship between the incidence of pterygium and dry eye syndrome in online motorcycle taxi drivers.

## ACKNOWLEDGEMENT

Researchers would like to thank the Occupational Medicine Study Center, Faculty of Medicine, Universitas Trisakti, and the Jabodetabek Ojol Community Association who were willing to help and contribute to this research.

## AUTHORS CONTRIBUTION

Research concept and design: EK, data collection: EK, HA, AA, RW, NP, analysis and interpretation of results: EK, HA, manuscript preparation: EK, EM, Y, review of final manuscript results: EK, EM

## FUNDING

Research funding is covered by Universitas Trisakti.

## CONFLICT OF INTEREST

Researchers have no conflict of interest regarding the results of this research.

## REFERENCES

1. Sharma A, Hindman HB. Aging: A Predisposition to Dry Eyes. *J Ophthalmol.* 2014;2014:1-8. doi:10.1155/2014/781683
2. Craig JP, Nelson JD, Azar DT, et al. TFOS DEWS II Report Executive Summary. *Ocul Surf.* 2017;15(4):802-12. doi: <http://dx.doi.org/10.1016/j.jtos.2017.08.003>
3. Tsubota K, Yokoi N, Shimazaki J, et al. New Perspectives on Dry Eye Definition and Diagnosis: A Consensus Report by the Asia Dry Eye Society. *Ocul Surf.* 2017; 5(1):65-76. doi:<https://doi.org/10.1016/j.jtos.2016.09.003>
4. Pedoman Nasional Pelayanan Kedokteran Dry Eye. PERDAMI. Available at: <https://perdami.or.id/wp-content/uploads/2022/03/PNPK-Dry-Eye-Final.pdf>
5. Gurnani B, Kaur K. Current Approach in Surgical Management of Dry Eyes - Dry Eye Review II. *TNOA J Ophthalmic Sci Res.* 2021;59:241-9. doi:10.4103/tjosr.tjosr\_56\_21
6. Mahardika ZA, Amani KF. Pengaruh Usia dan Lama Paparan Asap Kendaraan Bermotor terhadap Derajat Dry Eye Syndrome pada Pekerja Ojek. *Jurnal Ilmiah STIKES Kendal.* 2021;11(2):401-10.
7. Zhong YJ, Lee YC, Hsieh CJ, et al. Association between Dry Eye Disease, Air Pollution and Weather Changes in Taiwan. *Int J Environ Res Public Health.* 2018;15. doi:10.3390/ijerph15102269
8. Paulsen AJ, Cruickshanks KJ, Fischer ME, et al. Dry Eye in the Beaver Dam Offspring Study: Prevalence, Risk Factors, and Health-Related Quality of Life. *Am J Ophthalmol.* 2014;23(1):1-7. doi: 10.1016/j.ajo.2013.12.023.Dry.
9. Gierow JP, Kacz L. Effect of Age and Gender on Dry Eye according to Tests and Symptoms. *ARVO J.* 2018;59(9).
10. Manhas A, Gupta D, Gupta A, et al. Clinical Correlation between Dry eye and Pterygium: A Study Done at Government Medical College Jammu, Jammu and Kashmir, North India. *Int J Res Med Sci.* 2017;5(7):3087-3094. doi: <http://dx.doi.org/10.18203/2320-6012.ijrms20172992>
11. Gupta AK, Nathwani Y. Correlation between Pterygium and Dry Eye. *Kerala J Ophthalmol.* 2019;31:217-20. doi: 10.4103/kjo.kjo\_65\_19
12. Onkar A, Pandey DJ, Bist HK, et al. Tear and Pterygium: A Clinico-Pathological Study of Conjunctiva for Tear Film Anomaly in Pterygium. *J Eye Cataract Surg.* 2017;3(24):1-8. doi: 10.21767/2471-8300.100024.
13. Panigrahi S, Jyothi K. A Study on Dry Eye among Patients with Pterygium : A Tertiary Care Experience. *Journal of Cardiovascular Disease Research.* 2022;13(4):409-17.
14. Kiyat P, Karti O. Evaluation of Dry Eye in Eyes with Unilateral Pterygium. *Eur Eye Res.* 2023;3(1):12-5. doi: 10.14744/eer.2023.74946
15. Thatte S, Singh SO, Agrawal D, et al. Correlation of the Different Grades of Pterygium with Severity of Dry Eye Disease. *J Opht Res Rev Rep.* 2023;4(1):1-5. doi: [doi.org/10.47363/JORRR/2023\(4\)139](https://doi.org/10.47363/JORRR/2023(4)139).
16. Paudel N, Adhikari S, Manandhar S, et al. Ocular Surface Symptoms among Individuals Exposed to Ambient Levels of Traffic Derived Air Pollution. *F1000Research.* 2018;6:2167. doi: <https://doi.org/10.12688/f1000research.13483.2>.
17. Liang K, Gui SY, Qiao JC, et al. Association between Air Pollution Exposure and Daily Outpatient Visits for Dry Eye Disease: A Time-Series Study in Urumqi, China. *Atmosphere.* 2023;14(90). doi:<https://doi.org/10.3390/atmos14010090>.
18. Juddy AU, Esenwah EC, Ikoro NC, et al. Effect of Caffeinated Coffee on Tear Production. *Int J Res.* 2014;1(9):1264-8.
19. Fayola Ap, Kartadinata E. Relationship between Coffee Consumption and Dry Eye Syndrom in Adulthood. *J Biomedika dan Kesehatan.* 2023;6(1):5-12.
20. Magno MT, Utheim TP, Morthen MK, et al. The Relationship Between Caffeine Intake and Dry Eye Disease. *Cornea.* 2023;42:186-93. doi:10.1097/ICO.0000000000002979.

21. Murari S, Ho A, Hayes J, et al. Effects of caffeine intake on visual performance of the eye among normal healthy adults. *Coll Optom.* 2018;842. Available at: <https://commons.pacificu.edu/opt/842>.
22. Pritasari AMS, Faidah SN, Zulaikhah ST. Smoking as Risk Factors to Dry Eye Syndrome. *KEMAS.* 2019;15(1):1-5. doi:<https://doi.org/10.15294/kemas.v15i1.8611>
23. Najmee NABA, Nasir MNABA, Muhammad N, et al. Smoking Behaviour and Dry Eye Symptoms Among Uitm Puncak Alam Students During Covid 19. *Mal J Med Health Sci.* 2022;18(15):293-300. doi:[10.47836/mjmhs18.s15.41](https://doi.org/10.47836/mjmhs18.s15.41).
24. Ibrahim AM, Awara AM, Eldsouky M, et al. Ocular Surface Changes among Smokers. *J Adv Med Med Res.* 2023;15(35):28-35. doi: [10.9734/JAMMR/2023/v35i155066](https://doi.org/10.9734/JAMMR/2023/v35i155066).
25. Attri S, Dwivedi J, Mithal S, et al. Dry eye- Study of Prevalence, Associated Risk Factors and Frequency of Symptoms in Meerut District. *J. Evolution Med. Dent. Sci.* 2019;8(45):3382-3386. doi: [10.14260/jemds/2019/734](https://doi.org/10.14260/jemds/2019/734)
26. Tariq MA, Amin A, Ahmed B, et al. Association of Dry Eye Disease with Smoking: A Systematic Review and Meta analysis. *Indian J Ophthalmol.* 2022;70:1892-904. doi: [10.4103/ijo.IJO\\_2193\\_21](https://doi.org/10.4103/ijo.IJO_2193_21).
27. Thatte S, Choudhary R. The Prevalence of Dry Eye in Young Individuals Exposed to Visual Display Terminal. *Clin Ophthalmol J.* 2020;1(1):1004.
28. Mohtaseb ZA, Schachter S, Lee BS, et al. The Relationship Between Dry Eye Disease and Digital Screen Use. *Clin Ophthalmol J.* 2021;15:3811-20. doi:<https://doi.org/10.2147/OPTH.S321591>
29. Tawonkasiwattanakun P, Kosaisavee V, Suwannapong N, et al. Symptomatic Dry Eye Prevalence and Related Factors in an Open University, Bangkok, Thailand. *THJPH.* 2021; 51(3): 195-203.



This work is licensed under a Creative Commons Attribution Non-Commercial 4.0 International License

---

# Relationship Between Pterygium and Dry Eye Syndrome Among Delivery Motorbike Drivers

*by Anggraeni Adiwardhani FK*

---

**Submission date:** 21-Jun-2026 07:51PM (UTC+0700)

**Submission ID:** 2986919430

**File name:** 9\_395\_OA\_71-81\_1.pdf (323.67K)

**Word count:** 5381

**Character count:** 26891

**ORIGINAL ARTICLE**

## Relationship Between Pterygium and Dry Eye Syndrome Among Delivery Motorbike Drivers

Pterygium berhubungan dengan Sindroma Mata Kering pada Pengemudi Ojek Online

Erlani Kartadinata<sup>1</sup>, Husnun Amalia<sup>1</sup>, Anggraeni Adiwardhani<sup>1</sup>, Riani Witjaksana<sup>1</sup>, Noviani Prasetyaningsih<sup>1</sup>, Eveline Margo<sup>2</sup>, Yohana<sup>3</sup>

<sup>1</sup>Departement Ophthalmology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

<sup>2</sup>Departement Physiology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

<sup>3</sup>Departement Biochemistry, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

✉ [erlani.kartadinata@trisakti.ac.id](mailto:erlani.kartadinata@trisakti.ac.id)

<https://doi.org/10.18051/JBiomedKes.2024.v7.71-81>

### ABSTRACT

#### Background

Dry eye syndrome (DES) is an eye disease caused by many factors involving the ocular surface, with characteristic damage to tear layer homeostasis accompanied by ocular symptoms due to tear layer instability, hyperosmolarity, damage and inflammation of the ocular surface, and neurosensory abnormalities. Delivery motorbike drivers are often on the streets where the role of environmental factors will be able to pose a risk of experiencing DES. This study aims to assess risk factors associated with the incidence of dry eye syndrome in delivery motorbike drivers. The benefit of this research is to improve public health, especially in preventive behavior against factors associated with dry eye syndrome.

#### Methods

This study used observational analytical methods with a cross-sectional research design. The selection of samples by consequential non-random sampling amounted to 124 with the criteria of inclusion of delivery motorbike drivers, exclusion of not using artificial tears eye drops, and not use of contact lenses. Data were collected by filling out a DES risk factor questionnaire and a Tear Break Up Time (TBUT) examination to assess DES. The research data were analyzed using the Chi-Square test.

#### Results

This study's results show that subjects with dry eye syndrome are dominated by adults aged 19–44 and male sex. Subjects who experienced DES 76.6%, adult age 78.2%, male 80.6%, no pterygium 66.1%, working period 2-5 years 62.9%, length of work ≥8 hours 94.4%, always using a closed helmet 83.9%, not using protective glasses 39.5%, coffee consumption 71.8%, active smokers 66.1%, using gadgets >6 hours 91.9%. There is no significant relationship between subjects' characteristics and DES ( $p > 0.05$ ). The incidence of DES in pterygium showed significant results ( $p = 0.001$ ).

#### Conclusions

There is a significant relationship between the incidence of pterygium and DES.

**Keywords:** Dry Eye Syndrome; Pterygium; Risk Factor; Tear Break Up Time (TBUT)

## ABSTRAK

### Latar Belakang

Sindroma mata kering (DES) <sup>5</sup> adalah karakteristik rusaknya homeostasis lapisan air mata disertai dengan gejala okular diakibatkan ketidak-stabilan lapisan air mata, hiperosmolaritas, kerusakan dan inflamasi pada permukaan okular, serta abnormalitas neurosensoris. Pengemudi ojek online sering <sup>20</sup> ada di jalanan di mana peran faktor lingkungan akan dapat menimbulkan risiko mengalami DES. Penelitian ini bertujuan untuk menilai faktor risiko yang terkait dengan kejadian sindrom mata kering pada pengemudi sepeda motor pengiriman. Manfaat dari penelitian ini adalah untuk meningkatkan kesehatan masyarakat, terutama dalam perilaku preventif terhadap faktor-faktor yang terkait dengan <sup>35</sup> sindrom mata kering.

### Metode

Penelitian ini menggunakan metode analisis observasional dengan desain penelitian cross sectional. Pemilihan sampel secara konsekuensial non-random sampling berjumlah 124 dengan kriteria inklusi pengemudi ojek online, eksklusif tidak menggunakan tetes mata air mata buatan dan tidak menggunakan lensa kontak. Data dikumpulkan melalui pengisian kuesioner faktor risiko DES dan pemeriksaan Tear Break Up Time (TBUT) untuk menilai DES. <sup>34</sup> Data penelitian dianalisis menggunakan uji Chi-Square.

### Hasil

Hasil penelitian ini menunjukkan bahwa subjek yang memiliki sindrom mata kering didominasi oleh orang dewasa pada usia 19-44 tahun dan berjenis kelamin laki-laki. Subjek yang mengalami DES 76.6%, dewasa usia 78.2%, laki-laki 80.6%, tidak ada pterygium 66.1%, masa kerja 2-5 tahun 62.9%, lama kerja  $\geq$  8 jam 94.4%, selalu menggunakan helm tertutup 83.9%, tidak menggunakan kacamata pelindung 39.5%, konsumsi kopi 71.8%, perokok aktif 66.1%, menggunakan gadget >6 jam 91.9%. Tidak ada hubungan yang bermakna antara karakteristik subjek dan DES ( $p>0.05$ ). Kejadian DES pada pterygium menunjukkan hasil yang bermakna ( $p=0.001$ ).

### Kesimpulan

Ada hubungan yang bermakna antara kejadian pterygium dan DES.

<sup>38</sup> **Kata Kunci:** Sindroma Mata Kering; Pterygium; Faktor Risiko; Tear Break Up Time (TBUT)

## INTRODUCTION

<sup>10</sup> Dry eye syndrome (DES) is an eye disease caused by many factors involving the ocular surface, characterized by damage to tear film homeostasis accompanied by ocular symptoms due to tear film instability, hyperosmolarity, damage, and inflammation on the ocular surface, as well as neurosensory abnormalities.<sup>1,3</sup>

Symptoms of DES vary from mild irritation, burning sensation, itching, red eyes, gritty feeling in the eyes, sore eyes, tired eyes, and in severe cases can cause vision problems.<sup>1,4,5</sup> DES is the most common eye disease. in daily practice, with a prevalence of between 5-50% of the entire world population.<sup>2</sup> The prevalence of DES in Southeast Asia is around 20-52.4%.<sup>3</sup> The prevalence of dry eyes in Indonesia in 2017 was 26.5%.<sup>4</sup>

Several risk factors that are closely related to the incidence of DES include age, gender, race, connective tissue disorders, Sjogren's syndrome, estrogen replacement therapy, androgen hormone deficiency, use of drugs (antihistamines, antidepressants, anxiolytics, and isotretinoin), use of computers, use of contact lenses, environmental conditions (pollution, low humidity).<sup>1,2</sup> Probable risk factors include diabetes, rosacea, viral infections, thyroid disease, psychiatric conditions, pterygium, low fat intake, refractive surgery, and allergic conjunctivitis.<sup>2</sup>

Dry eyes are very common in older people, around 5-30% of the elderly population, and more often affect women than men.<sup>1</sup> The disparity in prevalence according to age ranges from 8.4% for those aged less than 60 years, 5% for those aged 70-79 years, and 20% over 80 years of age.<sup>1</sup>

Complications of DES that do not receive therapy or do not respond well to therapy can result in loss of visual acuity.<sup>1,2</sup> In severe cases of DES it can affect the patient's quality of life physically, socially, psychologically, and work productivity.<sup>4</sup>

Online motorcycle taxi driving is a relatively new profession in Indonesia and this profession helps community activities this profession is very popular and the number of online motorcycle taxi drivers is increasing. This job requires online drivers to always be on the road where environmental factors can pose a risk of experiencing DES.

Research conducted by Zsafidda and Kamellisha found that motorbike taxi workers in the city of Malang who had been exposed to smoke for more than 10 years experienced severe DES.<sup>6</sup> Currently there is not much research on online drivers, especially in the field of eye health, as well as the negative impacts that can arise. As a result of this work, researchers are interested in conducting research to assess the factors related to the incidence of DES in online motorcycle taxi drivers, for prevention efforts, especially in the field of eye health.

## METHODS

This research uses an analytical observational method with a cross-sectional design. This research was carried out at FK USAKTI, Jl. Kyai Tapa- Grogol, West Jakarta, in September 2022. The population in this study was online motorcycle taxi drivers, totaling 124 respondents. Sample selection used a non-random consecutive sampling technique. The inclusion criteria are online motorcycle taxi drivers, male and female. Exclusion criteria were using contact lenses and using artificial tears and eye drops.

Data was collected by filling out a questionnaire containing the respondent's identity and risk factors for dry eye syndrome. Then an anterior eye examination and Tear Break Up Time (TBUT) examination were carried out. TBUT examination shows whether there is a disturbance in the stability of the tear film. The examination is carried out using fluorescent strips. The inspection was carried out using a cobalt blue filter spotlight. Then the patient is asked to blink 3 times, then measure the time interval required after the last eye blink until a black spot appears on the cornea using a stopwatch. The measurement result is the time it takes for the tear film to break apart when the eye blinks. Normal TBUT results are 15-30 seconds. If TBUT  $\leq$  10 seconds indicates a disruption in tear film stability.

The statistical tests used were the Chi-square test and the Fisher Exact test with a significance level of  $p < 0.05$ . Data were analyzed univariately and bivariately using the SPSS (Statistical Package for Social Science) program version 23.0. The results are presented in the form of a frequency table of respondent characteristics and risk factors associated with dry eye syndrome.

## RESULTS

Table 1. Characteristics and Risk Factors for Dry Eye Syndrome

Variable	Respondent (n)	Percentage (%)
<b>Age</b>		
Adult (19-44 y.o)	97	78.2
Pre-elderly (45-59 y.o)	26	21.0
Elderly (> 60 y.o)	1	0.8
<b>Gender</b>		
Man	100	80.6
Woman	24	19.4
<b>Dry Eye</b>		
Yes	95	76.6
No	29	23.4
<b>Pterygium</b>		
Yes	42	33.9
No	82	66.1
<b>Years of service</b>		
< 2 years	1	0.8
2-5 Years	78	62.9
> 5 years	45	36.3
<b>Length of working</b>		
< 8 hours	7	5.6
> 8 hours	117	94.4
<b>Closed helmet</b>		
No	6	4.8
Rarely	14	11.3
Always	104	83.9
<b>Protective glasses</b>		
No	49	39.5
Rarely	36	29.0
Always	39	31.5
<b>Coffee Consumption</b>		
Yes	89	71.8
No	35	28.2
<b>Active smoker</b>		
Yes	82	66.1
No	42	33.9
<b>Gadgets/day</b>		
< 6 hours	10	8.1
> 6 hours	114	91.9

In table 1, most respondents were male (80.6%), mature age (78.2%), incidence of dry eye syndrome (76.6%), no pterygium (66.1%), working period 2-5 years (62.9%), working time  $\geq$  8 hours (94.4%), always using a closed helmet (83.9%), not using protective glasses (39.5%), consuming coffee (71.8%), active smoker (66.1%), using gadgets > 6 hours (91.9%).

Table 2. Relationship between respondent characteristics and risk factors for dry eye syndrome <sup>27</sup>

Variable	Dry eye syndrome				p
	Yes		No		
	n	%	n	%	
<b>Age</b>					0.462 <sup>c</sup>
Adults (19-44 years)	72	74.23	25	25.77	
Pre-elderly (25-59 years)	22	84.62	4	15.38	
Elderly (> 60 years old)	1	100.00	0	0.00	
<b>Gender</b>					0.456 <sup>c</sup>
Man	78	78.00	22	22.00	
Woman	17	70.83	7	29.17	
<b>Pterygium</b>					0.001 <sup>b*</sup>
Yes	39	92.86	3	7.14	
No	56	68.29	26	31.71	
<b>Years of service</b>					0.440 <sup>c</sup>
< 2 years	1	100.00	0	0.00	
2-5 years	57	73.08	21	26.92	
> 5 years	37	82.22	8	17.78	
<b>Working hours/day</b>					0.482 <sup>b</sup>
< 8 hours	6	85.71	1	14.29	
> 8 hours	89	76.07	28	23.93	
<b>Closed helmet</b>					0.087 <sup>c</sup>
No	6	100.00	0	0.00	
Rarely	8	57.14	6	42.86	
Always	81	77.88	23	22.12	
<b>Protective glasses</b>					0.087 <sup>c</sup>
No	39	79.59	10	20.41	
Rarely	23	63.89	13	36.11	
Always	33	84.62	6	15.38	
<b>Coffee Consumption</b>					0.072 <sup>c</sup>
Yes	72	80.90	17	19.10	
No	23	65.71	12	34.29	
<b>Active smoker</b>					0.414 <sup>c</sup>
Yes	61	74.39	21	25.61	
No	34	80.95	8	19.05	
<b>Gadgets/day</b>					0.115 <sup>b</sup>
< 6 hours	10	100.00	0	0.00	
> 6 hours	85	74.56	29	25.44	

<sup>c</sup> = chi-square test<sup>b</sup> = Fisher Exact Test

\* = p &lt; 0.05

Table 2 shows that more adult respondents aged 19-44 years experienced dry eye syndrome at 74.2% (72 respondents). In the statistical test results, the value of p=0.087 (p>0.05) was obtained, which means there is no significant relationship between age and dry eye syndrome. In the gender variable, the results showed that men experienced more dry eye syndrome, namely 78% (78 respondents). The statistical test results obtained a value of p=0.456 (p>0.05), which means there

is no significant relationship between gender and dry eye syndrome. Respondents who did not have pterygium experienced dry eye syndrome at 68.3% (39 respondents). The statistical test results obtained a value of  $p=0.001$  ( $p<0.05$ ), which means there is a significant relationship between pterygium and dry eye syndrome. Respondents with 2-5 years of work experienced the most dry eye syndrome, namely 73.1% (52 respondents). The statistical test results showed that  $p=0.440$  ( $p>0.05$ ), which means that there was no significant relationship between work experience and dry eye syndrome. Working hours  $\geq 8$  hours per day most frequently experienced dry eye syndrome at 76.1% (89%). The statistical test results showed that  $p=0.482$  ( $p>0.05$ ) which means there is no significant relationship between working hours  $\geq 8$  hours per day and dry eye syndrome. Respondents who always used closed helmets experienced dry eye syndrome of 77.9% (81 respondents). The statistical test results obtained  $p=0.087$  ( $p>0.05$ ), which means there is no significant relationship between using a closed helmet and dry eye syndrome. Respondents who did not use protective glasses experienced dry eyes of 79.6% (39 respondents). The statistical test results obtained  $p=0.087$  ( $p>0.05$ ), which means there is no significant relationship between using protective glasses and dry eye syndrome. 80.9% of respondents who consumed coffee had dry eye syndrome (72 respondents). The statistical test results obtained  $p=0.072$  ( $p>0.05$ ), which means there is no significant relationship between coffee consumption and dry eye syndrome. Active smoker respondents who had dry eye syndrome were 74.4% (61 respondents). The statistical test results showed  $p=0.414$  ( $p>0.05$ ), which means there is no significant relationship between active smoking and dry eye syndrome. Respondents who used gadgets  $>6$  hours per day had dry eye syndrome of 74.6% (85 respondents). The statistical test results obtained  $p=0.115$  ( $p>0.05$ ), which means there is no significant relationship between using gadgets  $>6$  hours per day and dry eye syndrome.

## DISCUSSION

The results in Table 1 show that online motorcycle taxi drivers are dominated by adult men (19-44 years). This is because adult men mostly use two-wheeled motorized vehicles and are breadwinners to meet family needs.

The results in Table 1 show that 76.6% of motorbike taxi drivers experienced dry eye syndrome. This is because in this study there were risk factors that caused dry eye syndrome, including 94.4% having a history of working more than 8 hours, and 71.8% consuming coffee, 66.1% are active smokers, and 91.9% use gadgets for more than 6 hours.

Table 2 shows that more adult respondents aged 19-44 years experienced dry eye syndrome at 74.2% (72 respondents). These results are in accordance with research conducted by Zong JY, *et al.* which states that there is a change in tear production with increasing age and dry eye syndrome is mostly in the younger generation (18-49 years).<sup>7</sup> In this study, 78.2% were aged 19-44 years, where at this age the use of mobile phones is increasing, especially in this era. modern today. This causes the risk of dry eye syndrome to increase at a young age. In the statistical test results, the value of  $p=0.087$  ( $p>0.05$ ) was obtained, which means there is no significant relationship between age and dry eye syndrome. This is in line with research conducted by Paulsen AJ, *et al.* which states that there is no significant relationship between age and dry eye syndrome with a value of  $p = 0.06$ .<sup>8</sup> Older respondents are said to be more at risk of experiencing a decrease in tear production because they have a decrease in corneal sensitivity which is related to the mechanism of dry eye. With

increasing age, apart from decreasing tear secretion, there is also an increase in tear evaporation such as atrophy of the meibomian glands. The distribution of dry eye syndrome in the age range may vary, this is because the causes of dry eye syndrome are multifactorial.<sup>9</sup>

In the gender variable, the results showed that men experienced more dry eye syndrome, namely 78% (78 respondents). The statistical test results obtained a value of  $p=0.456$  ( $p>0.05$ ), which means there is no significant relationship between gender and dry eye syndrome. These results are in line with research conducted by Gierow JP, et al. which stated that no significant relationship was found between gender and dry eye syndrome with a  $p$ -value  $> 0.05$ .<sup>9</sup> In several epidemiological studies, the prevalence of dry eye syndrome was higher in women. This is due to low estrogen in menopausal women, whereas in this study it was dominated by men at 80.6%.

Respondents who did not have pterygium experienced dry eye syndrome at 68.3% (39 respondents). The statistical test results obtained a value of  $p=0.001$  ( $p<0.05$ ), which means there is a significant relationship between pterygium and dry eye syndrome. These results are in accordance with research conducted by Manhas A, et al. which states that there is a significant relationship between pterygium and dry eye syndrome with a value of  $p<0.001$ .<sup>10</sup> Research conducted by Gupta AK also states that there is a significant relationship between pterygium and dry eye syndrome with  $p<0.05$ .<sup>11</sup> Pterygium is a condition where the tissue under the conjunctiva experiences degeneration in the form of proliferation of fibrovascular tissue that invades the cornea. The causes of pterygium include environmental factors such as ultraviolet (sun exposure) and chronic inflammation (exposure to dust and wind).<sup>12,13</sup> Tear film abnormalities are also one of the factors causing pterygium. Disorders of the stability of the tear film (tear film) are generally caused by two factors, namely chronic inflammation of the ocular surface and changes in tear dynamics.<sup>12</sup> Dry cornea and conjunctiva trigger the growth of new tissue. Pterygium causes the surface of the eyeball to become abnormal, resulting in tear film dysfunction and dry eye syndrome.<sup>10</sup> Tear film dysfunction is a precursor to pterygium or whether pterygium causes tear film dysfunction is still unclear.<sup>14</sup> The results of research conducted by Thatee S in 2023 stated that pterygium causes dry eye syndrome where the higher the stage/degree of the pterygium, the more severe the degree of dry eye syndrome.<sup>15</sup>

Respondents with 2-5 years of work experienced the most dry eye syndrome, namely 73.4% (57 respondents). The statistical test results showed that  $p=0.440$  ( $p>0.05$ ), which means that there was no significant relationship between work experience and dry eye syndrome. The results of research conducted by Paudel N, et al. stated that drivers who worked for 5 years experienced dry eye syndrome due to air pollution and there was no significant difference between the length of work of 5-10 years and 10 years.<sup>16</sup> The impact of air pollution on the surface of the eyeball occurs after a cumulative process of many years.

Working hours  $\geq 8$  hours per day most frequently experienced dry eye syndrome at 76.1% (89%). In the statistical test results, it was found that  $p=0.482$  ( $p>0.05$ ) which means there is no significant relationship between working hours  $\geq 8$  hours per day and dry eye syndrome. Respondents who always used closed helmets experienced dry eye syndrome of 77.9% (81 respondents). The statistical test results obtained  $p=0.087$  ( $p>0.05$ ), which means there is no significant relationship between using a closed helmet and dry eye syndrome. Respondents who did not use protective glasses experienced dry eyes of 79.6% (39 respondents). The statistical test results obtained

$p=0.087$  ( $p>0.05$ ), which means there is no significant relationship between using protective glasses and dry eye syndrome. Working as a driver will involve environmental exposure such as pollution, high temperatures, sun, and wind. The longer you drive, the greater the risk of exposure to environments that cause dry eye syndrome. The use of helmets and protective glasses aims to reduce exposure to this environment. Research conducted by Liang K, et al. states that exposure to accumulated air pollution is significantly related to dry eye syndrome.<sup>17</sup> The many factors that cause dry eye syndrome cause dry eye syndrome to be found in respondents who always wear helmets or protective glasses, for example, because they have had pterygium for a long time, the weather is heat, and other influencing factors.

80.9% of respondents who consumed coffee had dry eye syndrome (72 respondents). The statistical test results obtained  $p=0.072$  ( $p>0.05$ ), which means there is no significant relationship between coffee consumption and dry eye syndrome. The caffeine contained in coffee can affect tear production because it has the effect of increasing the activation of sympathetic innervation in the lacrimal gland which causes a decrease in tear secretion.<sup>18</sup> Research conducted by Juddy AU, et al. shows that there is a significant relationship between caffeine and a decrease in tear production.<sup>18</sup> Likewise, research conducted by Fayola PA shows that there is a significant relationship between coffee and dry eye syndrome with  $p=0.000$ .<sup>19</sup> Meanwhile, research conducted by Magno M. S, et al, in 2023, states that there is no significant relationship between caffeine and dry eye syndrome.<sup>20</sup> Caffeine reaches its maximum concentration in 20-120 minutes in plasma and is eliminated from the body after 4 hours.<sup>21</sup> Caffeine stimulates the parasympathetic by increasing acetylcholine which causes activation of water secretion, proteins, electrolytes and stimulation of tear secretion. The difference is in research conducted by Judy AU, et al. It uses experimental methods in collecting data. Research subjects were asked to consume caffeinated coffee and then had their tear production measured via the Schirmer test, before and after consuming coffee.

Active smoker respondents who had dry eye syndrome were 74.4% (61 respondents). The statistical test results showed  $p=0.414$  ( $p>0.05$ ), which means there is no significant relationship between active smoking and dry eye syndrome. Cigarette smoke irritates the surface of the eyeball, if it occurs continuously it will activate T cells which cause local inflammation and activation of cytokines which block sensory signals from the lacrimal gland resulting in a decrease in tear secretion.<sup>22</sup> Cigarette smoke on the surface of the eyeball causes disruption of the lipid layer. tear film which results in increased evaporation from the tear film.<sup>23</sup> Research conducted by Ibrahim AM, stated that there was a significant relationship with increasing degrees of dry eye syndrome in smokers compared to non-smokers with  $p=0.01$ .<sup>24</sup> This is different from research conducted by Attri S, who stated that there was no significant relationship between smoking, diabetes mellitus, hypertension, and dry eye syndrome.<sup>25</sup> Likewise, research conducted by Tariq MA, et al. stated that smoking was not indicated as a risk factor for dry eye syndrome.<sup>26</sup> In this study, although many respondents were smokers, more detailed and accurate data was still needed such as how to smoke, type of cigarette, duration of smoking, comorbidities, and other influencing factors.

Respondents who used gadgets >6 hours per day had dry eye syndrome of 74.6% (85 respondents). The statistical test results obtained  $p=0.115$  ( $p>0.05$ ), which means there is no significant relationship between using gadgets >6 hours per day and dry eye syndrome. Using a digital screen for more than 6 hours can increase the risk of dry eye syndrome.<sup>27</sup> Staring at a digital

screen for a long period can reduce the frequency of blinking which fails in the distribution of tears to lubricate the surface of the eye, so that tear evaporation will increase.<sup>27</sup> In research conducted by Mohtaseb ZA, et al. shows a significant relationship between the use of digital screens and a decrease in the blink reflex and the risk of developing dry eye syndrome.<sup>28</sup> Research conducted by Tawonkasiwattanakum P, et al. stated that there was no significant relationship between gadget use and dry eye syndrome in university students and staff.<sup>29</sup> In this study, the online driver profession will often use mobile phones in carrying out their work. The length of time you use a mobile phone depends on the length of work and the drivers' activities while working. If drivers often take breaks without using their mobile phones for a while or take short naps in their free time while working, this will reduce the risk of dry eye syndrome. Another thing that influences the results of this study is that there are many risk factors for dry eye syndrome, such as hot weather, cold rooms using air conditioners, wind, and other factors when using a mobile phone.

This research has the limitation of being carried out subjectively using a multifactorial questionnaire of risk factors for dry eye and the causes of dry eye syndrome, so a more detailed questionnaire is needed for each risk factor and adding other tests to confirm the diagnosis of dry eye syndrome and increasing the number of respondents studied more.

This research implies that it is hoped that preventive and educational measures regarding dry eye syndrome and pterygium can be carried out among online motorcycle taxi drivers so that the productivity of online motorcycle taxi drivers is optimal.

## CONCLUSION

The conclusion from the analysis of this research data is that there is a relationship between the incidence of pterygium and dry eye syndrome in online motorcycle taxi drivers.

## ACKNOWLEDGEMENT

Researchers would like to thank the Occupational Medicine Study Center, Faculty of Medicine, Universitas Trisakti, and the Jabodetabek Ojol Community Association who were willing to help and contribute to this research.

## AUTHORS CONTRIBUTION

Research concept and design: EK, data collection: EK, HA, AA, RW, NP, analysis and interpretation of results: EK, HA, manuscript preparation: EK, EM, Y, review of final manuscript results: EK, EM

28

## FUNDING

Research funding is covered by Universitas Trisakti.

## CONFLICT OF INTEREST

Researchers have no conflict of interest regarding the results of this research.

## REFERENCES

1. Sharma A, Hindman HB. Aging: A Predisposition to Dry Eyes. *J Ophthalmol.* 2014;2014:1-8. doi:10.1155/2014/781683
2. Craig JP, Nelson JD, Azar DT, et al. TFOS DEWS II Report Executive Summary. *Ocul Surf.* 2017;15(4):802-12. doi: <http://dx.doi.org/10.1016/j.jtos.2017.08.003>
3. Tsubota K, Yokoi N, Shimazaki J, et al. New Perspectives on Dry Eye Definition and Diagnosis: A Consensus Report by the Asia Dry Eye Society. *Ocul Surf.* 2017; 5(1):65-76. doi:<https://doi.org/10.1016/j.jtos.2016.09.003>
4. Pedoman Nasional Pelayanan Kedokteran Dry Eye. PERDAMI. Available at: <https://perdami.or.id/wp-content/uploads/2022/03/PNPK-Dry-Eye-Final.pdf>
5. Gumani B, Kaur K. Current Approach in Surgical Management of Dry Eyes - Dry Eye Review II. *TNOA J Ophthalmic Sci Res.* 2021;59:241-9. doi:10.4103/tjosr.tjosr\_56\_21
6. Mahardika ZA, Amani KF. Pengaruh Usia dan Lama Paparan Asap Kendaraan Bermotor terhadap Derajat Dry Eye Syndrome pada Pekerja Ojek. *Jurnal Ilmiah STIKES Kendal.* 2021;11(2):401-10.
7. Zhong YJ, Lee YC, Hsieh CJ, et al. Association between Dry Eye Disease, Air Pollution and Weather Changes in Taiwan. *Int J Environ Res Public Health.* 2018;15. doi:10.3390/ijerph15102269
8. Paulsen AJ, Cruickshanks KJ, Fischer ME, et al. Dry Eye in the Beaver Dam Offspring Study: Prevalence, Risk Factors, and Health-Related Quality of Life. *Am J Ophthalmol.* 2014;23(1):1-7. doi: 10.1016/j.ajo.2013.12.023.Dry.
9. Gierow JP, Kacz L. Effect of Age and Gender on Dry Eye according to Tests and Symptoms. *ARVO J.* 2018;59(9).
10. Manhas A, Gupta D, Gupta A, et al. Clinical Correlation between Dry eye and Pterygium: A Study Done at Government Medical College Jammu, Jammu and Kashmir, North India. *Int J Res Med Sci.* 2017;5(7):3087-3094. doi: <http://dx.doi.org/10.18203/2320-6012.ijrms20172992>
11. Gupta AK, Nathwani Y. Correlation between Pterygium and Dry Eye. *Kerala J Ophthalmol.* 2019;31:217-20. doi: 10.4103/kjo.kjo\_65\_19
12. Onkar A, Pandey DJ, Bist HK, et al. Tear and Pterygium: A Clinico-Pathological Study of Conjunctiva for Tear Film Anomaly in Pterygium. *J Eye Cataract Surg.* 2017;3(24):1-8. doi: 10.21767/2471-8300.100024.
13. Panigrahi S, Jyothi K. A Study on Dry Eye among Patients with Pterygium : A Tertiary Care Experience. *Journal of Cardiovascular Disease Research.* 2022;13(4):409-17.
14. Kiyat P, Karti O. Evaluation of Dry Eye in Eyes with Unilateral Pterygium. *Eur Eye Res.* 2023;3(1):12-5. doi: 10.14744/eer.2023.74946
15. Thatte S, Singh SO, Agrawal D, et al. Correlation of the Different Grades of Pterygium with Severity of Dry Eye Disease. *J Opht Res Rev Rep.* 2023;4(1):1-5. doi: [doi.org/10.47363/JORRR/2023\(4\)139](https://doi.org/10.47363/JORRR/2023(4)139).
16. Paudel N, Adhikari S, Manandhar S, et al. Ocular Surface Symptoms among Individuals Exposed to Ambient Levels of Traffic Derived Air Pollution. *F1000Research.* 2018;6:2167. doi: <https://doi.org/10.12688/f1000research.13483.2>.
17. Liang K, Gui SY, Qiao JC, et al. Association between Air Pollution Exposure and Daily Outpatient Visits for Dry Eye Disease: A Time-Series Study in Urumqi, China. *Atmosphere.* 2023;14(90). doi:<https://doi.org/10.3390/atmos14010090>.
18. Juddy AU, Esenwah EC, Ikoru NC, et al. Effect of Caffeinated Coffee on Tear Production. *Int J Res.* 2014;1(9):1264-8.
19. Fayola Ap, Kartadinata E. Relationship between Coffee Consumption and Dry Eye Syndrom in Adulthood. *J Biomedika dan Kesehatan.* 2023;6(1):5-12.
20. Magno MT, Utheim TP, Morthen MK, et al. The Relationship Between Caffeine Intake and Dry Eye Disease. *Cornea.* 2023;42:186-93. doi:10.1097/ICO.0000000000002979.

21. Murari S, Ho A, Hayes J, et al. Effects of caffeine intake on visual performance of the eye among normal healthy adults. *Coll Optom.* 2018;842. Available at: <https://commons.pacificu.edu/opt/842>.
22. Pritasari AMS, Faidah SN, Zulaikhah ST. Smoking as Risk Factors to Dry Eye Syndrome. *KEMAS.* 2019;15(1):1-5. doi:<https://doi.org/10.15294/kemas.v15i1.8611>
23. Najmee NABA, Nasir MNABA, Muhammad N, et al. Smoking Behaviour and Dry Eye Symptoms Among Uitm Puncak Alam Students During Covid 19. *Mal J Med Health Sci.* 2022;18(15):293-300. doi:10.47836/mjmhs18.s15.41.
24. Ibrahim AM, Awara AM, Eldsouky M, et al. Ocular Surface Changes among Smokers. *J Adv Med Med Res.* 2023;15(35):28-35. doi: 10.9734/JAMMR/2023/v35i155066.
25. Attri S, Dwivedi J, Mithal S, et al. Dry eye- Study of Prevalence, Associated Risk Factors and Frequency of Symptoms in Meerut District. *J. Evolution Med. Dent. Sci.* 2019;8(45):3382-3386. doi: 10.14260/jjemds/2019/734
26. Tariq MA, Amin A, Ahmed B, et al. Association of Dry Eye Disease with Smoking: A Systematic Review and Meta analysis. *Indian J Ophthalmol.* 2022;70:1892-904. doi: 10.4103/ijoo.IJO\_2193\_21.
27. Thatte S, Choudhary R. The Prevalence of Dry Eye in Young Individuals Exposed to Visual Display Terminal. *Clin Ophthalmol J.* 2020;1(1):1004.
28. Mohtaseb ZA, Schachter S, Lee BS, et al. The Relationship Between Dry Eye Disease and Digital Screen Use. *Clin Ophthalmol J.* 2021;15:3811-20. doi:<https://doi.org/10.2147/OPTH.S321591>
29. Tawonkasiwattanakun P, Kosaisavee V, Suwannapong N, et al. Symptomatic Dry Eye Prevalence and Related Factors in an Open University, Bangkok, Thailand. *THJPH.* 2021; 51(3): 195-203.



This work is licensed under a Creative Commons Attribution Non-Commercial 4.0 International License

# Relationship Between Pterygium and Dry Eye Syndrome Among Delivery Motorbike Drivers

## ORIGINALITY REPORT

19%

SIMILARITY INDEX

14%

INTERNET SOURCES

12%

PUBLICATIONS

3%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="https://publications.aston.ac.uk">publications.aston.ac.uk</a> Internet Source	1%
2	Rita Khairani, Mustika Anggiane Putri, Dyah Ayu Woro Setyaningrum. "Korelasi Karbon Monoksida dan Derajat Rokok dengan Fungsi Paru pada Pengemudi Ojek Online", Jurnal Biomedika dan Kesehatan, 2023 Publication	1%
3	Julian Chendrasari, Indah Widya Lestari, Florinda Lestari, Reza Lestari. "Hubungan Kadar Timbal (Pb) Darah Dengan Hipertensi Pada Pengemudi Ojek", Jurnal Biomedika dan Kesehatan, 2023 Publication	1%
4	<a href="https://journal.unnes.ac.id">journal.unnes.ac.id</a> Internet Source	1%
5	<a href="https://perdami.or.id">perdami.or.id</a> Internet Source	1%
6	Ayu Oktaviana, Elly Herwana. "Demographic Factors and Total Muscle Mass are Associated with Handgrip Strength in Selected Indonesian Adults", 2021 IEEE International Conference on Health, Instrumentation & Measurement, and Natural Sciences (InHeNce), 2021 Publication	1%

7	<a href="http://midwifery.iocspublisher.org">midwifery.iocspublisher.org</a> Internet Source	1 %
8	<a href="http://pubmed.ncbi.nlm.nih.gov">pubmed.ncbi.nlm.nih.gov</a> Internet Source	1 %
9	Submitted to Deakin University Student Paper	1 %
10	<a href="http://www.mdpi.com">www.mdpi.com</a> Internet Source	1 %
11	<a href="http://ojs.unpkediri.ac.id">ojs.unpkediri.ac.id</a> Internet Source	1 %
12	<a href="http://pmc.ncbi.nlm.nih.gov">pmc.ncbi.nlm.nih.gov</a> Internet Source	1 %
13	Joekly Wahidan Muharaam, Diana Samara. "Fenomena Gejala Pilek paling Sering pada Pasien COVID-19", Jurnal Biomedika dan Kesehatan, 2023 Publication	<1 %
14	<a href="http://doaj.org">doaj.org</a> Internet Source	<1 %
15	<a href="http://jurnal.globalhealthsciencegroup.com">jurnal.globalhealthsciencegroup.com</a> Internet Source	<1 %
16	<a href="http://scholar.archive.org">scholar.archive.org</a> Internet Source	<1 %
17	Yasinta Saraswati Hakim, Setyoko Setyoko, Yanurita Tursinawati. "Hubungan antara Indeks Massa Tubuh dan Lingkar Pinggang dengan Kejadian Hipertensi Sistolik Terisolasi pada Lansia", Jurnal Biomedika dan Kesehatan, 2022 Publication	<1 %
18	<a href="http://medic.upm.edu.my">medic.upm.edu.my</a> Internet Source	<1 %

19	Submitted to King Saud bin Abdulaziz University for Health Sciences Student Paper	<1 %
20	repository.umy.ac.id Internet Source	<1 %
21	Sunita Sabarwal, Priyanka ., Bruttendu Moharana, Rituka Gupta, Bhavana Sharma. "Prevalence, pattern and associated risk factors of Dry eye disease: Results of a prospective patient database from a tertiary eye care centre in Central India", Springer Science and Business Media LLC, 2024 Publication	<1 %
22	Tanushree V, Venkate Gowda H T. "CORELATION OF DRY EYE STATUS WITH SEVERITY OF DIABETIC RETINOPATHY", Journal of Evolution of Medical and Dental Sciences, 2014 Publication	<1 %
23	e-journal.unair.ac.id Internet Source	<1 %
24	www.dovepress.com Internet Source	<1 %
25	pure.rug.nl Internet Source	<1 %
26	www.medrxiv.org Internet Source	<1 %
27	Masoud Reza Manaviat, Maryam Rashidi, Mohammad Afkhami-Ardekani, Mohammad Reza Shoja. "Prevalence of dry eye syndrome and diabetic retinopathy in type 2 diabetic patients", BMC Ophthalmology, 2008 Publication	<1 %

28 Wahdaniar Wahdaniar, Imran Pashar, Miladiarsi Miladiarsi. "Pengaruh Pencucian Luka menggunakan Infus Daun Sirih Merah (Piper Crocatum) 40% terhadap Proses Penyembuhan Ulkus Diabetik", Jurnal Biomedika dan Kesehatan, 2022

Publication

<1 %

29 Yu-Ting Jeng, Shu-Yi Lin, Hsiao-Yun Hu, Oscar K. Lee, Li-Lin Kuo. "Osteoporosis and dry eye syndrome: A previously unappreciated association that may alert active prevention of fall", PLOS ONE, 2018

Publication

<1 %

30 E P Akbar, F Kurniadi, C Destria. "Adaptation of Ojek Online's Behavior to Open Space Settings in Pontianak City", IOP Conference Series: Earth and Environmental Science, 2020

Publication

<1 %

31 Essentials in Ophthalmology, 2015.

Publication

<1 %

32 Monica Alves, Penny Asbell, Murat Dogru, Giuseppe Giannaccare et al. "TFOS Lifestyle Report: Impact of environmental conditions on the ocular surface", The Ocular Surface, 2023

Publication

<1 %

33 [www.researchgate.net](http://www.researchgate.net)

Internet Source

<1 %

34 [www.frontiersin.org](http://www.frontiersin.org)

Internet Source

<1 %

35 [repository.unair.ac.id](http://repository.unair.ac.id)

Internet Source

<1 %

36

[vbn.aau.dk](http://vbn.aau.dk)

Internet Source

<1 %

37

[www.researchsquare.com](http://www.researchsquare.com)

Internet Source

<1 %

38

Shashank Dokania, Parth Aphale, Neeta Mishra. "Efficacy of homoeopathic preparation of Euphrasia officinalis as Eye Drops in management of dry eye syndrome: An Experimental Study", International Journal of High Dilution Research - ISSN 1982-6206, 2025

Publication

<1 %

39

[healthdocbox.com](http://healthdocbox.com)

Internet Source

<1 %

40

[www.coursehero.com](http://www.coursehero.com)

Internet Source

<1 %

Exclude quotes Off

Exclude matches < 10 words

Exclude bibliography On