

Malaysian Journal of Soil Science

COUNTRY

Malaysia



Universities and research institutions in Malaysia



Media Ranking in Malaysia

SUBJECT AREA AND CATEGORY


Agricultural and Biological Sciences
- Agronomy and Crop Science
- Soil Science

PUBLISHER

Malaysian Society of Soil Science

H-INDEX

14



PUBLICATION TYPE

Journals

ISSN


13947990



Documents cited by public policy (Overton)



Documents related to SDGs (UN)



Malaysian Journal of Soil Science

Q2

Agronomy and Crop Science

best quartile

SJR 2023

← Show this widget in your own website


Just copy the code below and paste within your html code:

SCImago Graphica

Explore, visually communicate and make sense of data with our **new data visualization**



Cover



Malaysian Journal of Soil Science

Editorial Board of MJSS

Chief Editor



Dr. Daljit Singh Karam Singh
Universiti Putra Malaysia

Managing Editor



Dr. Arina Shairah Abdul Sukor
Universiti Putra Malaysia

Associate Editor



Prof. Dr. Shamshuddin Jusop
Universiti Putra Malaysia



Prof. Dr. Che Fauziah Ishak
Universiti Putra Malaysia



Assoc. Prof. Dr. Hasmah Mohidin
Universiti Teknologi MARA (UITM)



Dr. Samsuri Abd Wahid
Universiti Putra Malaysia



Dr. Rosazlin Abdullah
Universiti Malaya



Dr. Jeyanny Vijayanathan
Forest Research Institute Malaysia
(FRIM)

International Associate Editors



Prof. Dr. Prakash B. Nagabovanalli
Department of Soil Science &
Agricultural Chemistry,
University of Agricultural Sciences,
Bangalore, India



Prof. Dr. Ibrahim Ortas
Department of Soil Science,
Faculty of Agriculture,
Cukurova University, Turkiye



Prof. Dr. Hseu Zeng-Yei
Department of Agricultural Chemistry,
National Taiwan University Taipei,
Taiwan



Prof. Dr. Udaya W.A. Vitharana
Department of Soil Science,
Faculty of Agriculture,
University of Peradeniya, Sri Lanka



Assoc. Prof. Dr. Mami Irei
Department of International
Agricultural Development,
Faculty of International Agriculture
and Food Studies,
Tokyo University of Agriculture, Japan

Daftar Isi

[Home](#) [Publication](#) [Journal \(MJSS\)](#) [Directory of Members](#) [Quick links](#)



Malaysian Society of Soil Science

Home / MJSS / Vol. 28

The Journal
Editorial Board
Author Instructions
Publication Fees
MJSS Submission

Vol. 28 | 2024

Vol. 27 | 2023

Vol. 26 | 2022

Vol. 25 | 2021

Vol. 24 | 2020

Vol. 23 | 2019

Vol. 22 | 2018

Vol. 21 | 2017

Vol. 20 | 2016

Vol. 19 | 2015

Vol. 18 | 2014

Vol. 17 | 2013

Vol. 16 | 2012

Vol. 15 | 2011

Vol. 14 | 2010

Vol. 13 | 2009

Vol. 12 | 2008

Vol. 11 | 2007

Vol. 10 | 2006

Vol. 09 | 2005

Vol. 08 | 2004

Vol. 07 | 2003

Vol. 06 | 2002

Vol. 05 | 2001

Vol. 04 | 2000

Vol. 03 | 1999

Vol. 02 | 1998

Vol. 01 | 1997

Malaysian Journal Of Soil Science

Vol. 28 | December 2024

Mitigating Nitrogen Leaching in Mineral Soils using Pineapple Leaf Biochar

Pages 1-11

Jos, Syahira; Mohidin, Hasmah; Ahmed, Osumanu H; Kassim, Nur Qursyna B; Mahdian, Suraiya; & Rosli, Nurmaliena
▼ Abstract | [Full Text](#)

Keywords: pineapple leaf biochar, soil amendment, nitrogen retention

Assessing Soil Bulk Density, Plasticity Index, Porosity, and Degree of Saturation through Electrical Resistivity using Correlation Analysis

Pages 12-25

Muhammad Burhan Memon; Zheng Yang; Waqar Hussain Qazi; Shafi Muhammad Pathan; and Saleem Raza Chalgrri
▼ Abstract | [Full Text](#)

Keywords: Geophysical technique, soil properties, correlation model, non-invasive method, borehole sampling

Effect of Co-application of Vetiver Grass Biochar and NPK Fertilizer on the Growth of Oil Palm (*Elaeis guineensis* Jacq.) Seedlings and Soil Chemical Properties

Pages 26-37

Aleladeh, Hemam; Idris, Juferi; Mohidin, Hasmah; Omar, Latifah; Man, Sulaiman; Munir, Siraj; & Jong, Vivien
▼ Abstract | [Full Text](#)

Keywords: Biochar, NPK, oil palm, growth, soil chemical properties

Analysis of Soil Quality through Aerial Biomass Contribution of Three Forest Species in Relict High Andean Forests of Peru

Pages 38-52

Ysaías Zanabria Cáceres; Betty Cordova Torres; Gelly Clemente Archi; Rosario Zanabria Mallqui; Lucia Enriquez Pinedo; Dennis Ccopi Trucios; & Kevin Ortega Quispe
▼ Abstract | [Full Text](#)

Keywords: Soil quality, relict forests, nutrient recycling, biomass, coverage

Efficiency of Indigenous Soil Rhizobia on Growth Performance of the Common Bean (*Phaseolus vulgaris* L.)

Pages 53-62

Mulugeta Mekonnen; Ameha Kebede; Meseret Chimdessa Egilgu; and Manikandan Muthuswamy
▼ Abstract | [Full Text](#)

Keywords: indigenous rhizobia, common bean, symbiotic efficiency

Optimizing the Retention and Leaching of Potassium of Tropical Mineral Acid Soils with Application of Charcoal and Sago Bark Ash

Pages 63-78

Puvan Paramisaperam; Osamenu Haruna Ahmed; Latifah Omar; Prisca Divra Johan; Nur Hideyah Hamidi; Huck Ywih Ch'ng; and Adiza Alhassan Musah
▼ Abstract | [Full Text](#)

Keywords: Soil amendments, potassium availability, soil fertility, cation exchange capacity, sustainable agriculture

Effect of Mango Tree Age on Soil Fertility, Plant Nutrient Contents and Their Relationship

Pages 79-91

Kamarudin, K.N.; Abdul Rahman, M.H.; Abd Rashid, N.F.; Mohamad, M.; Mohamad, M.; Khairun, N.M.; Shahidin, N.M.; and Roslan, N.
▼ Abstract | [Full Text](#)

Keywords: Harumanis variety, linear relationship, MA128, *Mangifera indica*, soil and plant nutrient contents

Assessment of Plant Height and Trunk Diameter of Oil Palm as a Sole Function of Soil Textural Grains (Sand, Clay, and Silt)

Pages 92-104

Idris Sani; Teh Christopher Boon Sung; Zaibon Syaharudin; and Sim Choon Cheak
▼ Abstract | [Full Text](#)

Keywords: Plant height, trunk diameter, clay, sand, silt

Effects of Local Resource-based Ameliorant Residues on Chemical Properties of Inceptisols and Green Bean Production (*Phaseolus vulgaris* L.)

Pages 105-116

Herdiyanti, Herdiyanti; Maulana, A.; Fathi, A.N.M.; Monikasari, M.; Kasim, M.; Ryswaldi, R.; Habazar, T.; Noor, M.; Lita, R.P.; & Refdl, C.W.

▼ Abstract | [Full Text](#)

Keywords: ameliorant formulations, green beans, intercropping, local resources, residue

Mitigating Rapid Release of Nitrate from Rejected Sago Starch-coated Urea and Uncoated Urea

Pages 117-133

Rajan Kavitha; Omar Latifah; Osumanu Haruna Ahmed; Primus Walter Charles; and Kasim Susilawati

▼ Abstract | [Full Text](#)

Keywords: coated urea, urea hydrolysis, natural polymer, nitrogen loss

The Relative Importance of Stand and Soil Properties Parameters on Soil Organic Matter Content of Acacia Hybrid Forests in the South Central Coast Region of Vietnam

Pages 134-146

Cuong, L.V.; Quy, N.V.; Hung, B.M.; Chau, M.H.; and Doan, P.V.T.D.

▼ Abstract | [Full Text](#)

Keywords: Factors affecting, stand structural characteristics, soil physico-chemical properties, plantation forests, stand age

Effects of Peat in Reducing the Salinity of Spent Mushroom Waste as Growing Medium

Pages 147-152

Nurhidayah; Abdul Rahman; Salwa Adani; and Nur Qursyria Bull Kassim

▼ Abstract | [Full Text](#)

Keywords: spent mushroom waste, salinity, peat soil, spinach

Adsorption Behaviour and Microstructural Study of Non-Polar Amino Acids in Natural Soil Amended with Pesticide

Pages 153-164

Ankita Nagar; Shagufra Jabin; and Jyoti Chawla

▼ Abstract | [Full Text](#)

Keywords: Adsorption, mobility, plant nutrition, scanning electron microscopy, Soil thin layer chromatography, X-ray diffraction

Distribution Of Manganese Fractions in Soil under Long-term Tea Cultivation at Thai Nguyen Province, Vietnam

Pages 165-171

Hoang Huu Chien; Dang Van Minh; Nguyen Huy Trung; Chu Van Trung; Nguyen Quang Thi; Nguyen Duy Hai; Sota Tanaka; and Kozo Iwasaki

▼ Abstract | [Full Text](#)

Keywords: *Camellia sinensis*, manganese fractions, soil acidification, tea gardens, Thai Nguyen

NORMs Spatial Distribution and Radiological Risk Assessment in the Soil of The Kesang River Basin, Malaysia

Pages 172-185

Ismail, Nurul IzzatiAffi; Jaafar, Mohd Zuli; Hashim, Mohammad Mu'az; Nik Ariffin, Nik Azlin; Zaharullil, Nur Atiqah; Wan Mohd Zain; & Wan Zuraida

▼ Abstract | [Full Text](#)

Keywords: River, soil, NORMs, radiological hazard, principal component analysis

Spatial Variability and Distribution of Soil Nutrients at Harumanis Mango Farm During Vegetative and Flowering Stages

Pages 186-198

Shahidin, N.M.; Karim, M.A.A.; Pauzi, S.A.; Rashid N.F.A.; Kamarudin, K.N.; and Chuah, T.S.

▼ Abstract | [Full Text](#)

Keywords: Harumanis mango, phenological stages, soil nutrients distribution, spatial variability.

The Potential of Biochar's Properties in Ameliorating Soil Productivity of Acidic Ferralsols

Pages 199-211

Walelign Demisie, Mingkui Zhang

▼ Abstract | [Full Text](#)

Keywords: Acidic ferralsol, biochar, surface properties, nutritional properties, soil productivity

The Role of Microclimate, Vegetation Variety and Land Use in the Formation of Humic Substances

Pages 212-229

Mindari, W; Chakim, MG; Sasongko, PE; Aditya, HF; Karam, DS; and Masri, IN

▼ Abstract | [Full Text](#)

Keywords: Organic-C, humic substances, land use, microclimate, vegetation

Variations in Soil Phosphorus Levels in *Acacia* Hybrid Plantations Across Different Ages in Southern Vietnam

Pages 230-243

Cuong, L.V.; Karam D.S.; Hung, B.M.; Chau, M.H.; and Quy, N.V.

▼ Abstract | [Full Text](#)

Keywords: *Acacia* hybrid plantation, soil features, stand characteristics, structural equation modeling, stand age

Maize Yield Pot Response Toward Different Rates of Indigenous Microorganism Biocompost Integrated with Rice Husk Biochar Cultivated on Gajah Mati Series Soil

Pages 244-254

Lim, M.R.Z.M.R.; Muzamal, Z.; & Kamarudin, K.N.

▼ Abstract | [Full Text](#)

Keywords: IMO, problematic soil, production, sweet corn, *Zea mays* L.

Determination of Available Phosphorus In Soil Using ICP-OES And UV-VIS Spectrophotometer: A Comparison

Pages 255-267

Mohd Kamaruddin Jaffar, Nur Syafiqah Mohamad Sa'adan; & Lee Suan Chua

▼ Abstract | [Full Text](#)

Keywords: soils, available phosphorus, ICP-OES, UV-Vis, productivity, linearity

Model of Artificial Sprinkling to Study the Influence of Slope Steepness and Rainfall Rate on Soil Erosion

Pages 268-275

Djalilova Gulnora Tulkunovna & Nurullaev Azamkhon

▼ Abstract | [Full Text](#)

Keywords: erosion, soil, washout, slope steepness, rainfall

Soil Quality Assessment and Climatic Condition of Selected Forest Reserves in Terengganu, Malaysia

Pages 276-292

M.H.M. Ibrahim; N. Yunus; A. Abdu; D.S. Karam; K.S. Rajoo; R. Rani; Z. Ibrahim; A.R.A. Wahid; M.A.A. Abdullah; D. Zulperi; & M.I.H. Johari

▼ Abstract | [Full Text](#)

Keywords: soil conservation, tropical forest, sustainable forest management

Assessing the Effectiveness of Buffer Zones in Logged Malaysia Borneo Tropical Forests using Tropical Soil Quality Index (TSQI)

Pages 293-304

Bodos, V.A.; Rajoo, K.S.; Karam, D.S.; Arifin, A.; & Irie, M.

▼ Abstract | [Full Text](#)

Keywords: Tropical forest management, soil quality, buffer zones, logging impacts, Tropical Soil Quality Index (TSQI)

Peat Soil Verification and Assessment for Independent Smallholders RSPO Certification in Pontian and Batu Pahat, Johor: A Case Study

Pages 305-311

Ata, Asad; Adam, Salwa; Faizal, Mohamad Amirul Shafiq; Ajman, Nadiatul Nazleen; Ramachandran, Vasagi; & Boll Kassim, Nur Qursyna

▼ Abstract | [Full Text](#)

Keywords: oil palm, peat soil, RSPO, smallholders, sustainability

Preliminary Study on Carbofuran-Degrading Bacteria Isolated from Agricultural Soil

Pages 312-220

Jafarudin, N. & Ariffin, F.

▼ Abstract | [Full Text](#)

Keywords: Biodegradation, agricultural site, carbofuran, soil bacteria

Comparison of Five Phosphorus (P) and Potassium (K) Extraction Methods in Tomato Cultivation (*Solanum lycopersicum*) in Andisols and Inceptisols

Pages 332-344

Dermawan, Rahmansyah; Susila, Anas D; Purwono, Purwono; Nugroho, Budi; & Faried, Muhammad

▼ Abstract | [Full Text](#)

Keywords: extraction methods, fertilization, nutrient availability, tomato

Potassium, Zinc, Gypsum and Organic Manure Influence Electrical Conductivity, Biomass Yield and Nutrient Accumulation in Maize Irrigated with Brackish Water in Coastal Saline Soil

Pages 345-358

Mishra, Sancharita; Alomar, Rahaf; Barala, Jatiprasad; Patel, Sonali; Panda, Debadarshini; Pattnaik, Manoranjan; & Jena, Dinabandhu

▼ Abstract | [Full Text](#)

Keywords: brackish water, potassium, gypsum, zinc, maize

Potassium, Zinc, Gypsum and Organic Manure Influence Electrical Conductivity, Biomass Yield and Nutrient Accumulation in Maize Irrigated with Brackish Water in Coastal Saline Soil

Pages 345-358

Mishra, Sancharita; Alomar, Rahaf; Barala, Jatiprasad; Patel, Sonali; Panda, Debadarshini; Pattnaik, Manoranjan; & Jena, Dinabandhu

▼ Abstract | [Full Text](#)

Keywords: brackish water, potassium, gypsum, zinc, maize

***Gluconacetobacter diazotrophicus* Bacteria combined Nitrogen Fertilization Promotes Rice Yield and Soil Quality in The Paddy Field in The Vietnamese Mekong Delta Region**

Pages 359-368

Vu Van Long; Vo Trung Duong; Le Bich Tuyen; Tran Van Dung; Nguyen Khoi Nghia; Tran Sy Nam; Nguyen Quoc Viet; Le Van Cuong; & Huynh Van Tien

▼ Abstract | [Full Text](#)

Keywords: Chlorophyll, *Gluconacetobacter diazotrophicus*, grow-promoting bacteria, seed inoculation, soil fertility

Systematic Review of Emerging Trends in Soil-Based Probiotic

Pages 369-381

Khairul, S.R.; Leong, S.S.; Korel, F.; Lingoh, A.D.; & Toh, Seng Chiew

▼ Abstract | [Full Text](#)

Keywords: Pathogen, plant, soil-based probiotics, probiotics

Soil Chemical Characteristics And Soil Fertility Status In Coffee Agroforestry In The Upper Ciliwung Watershed

Pages 382-387

Fitri, R.; Simangunsong, N.I.; Danniswari, D.; and Taki, H.M.

▼ Abstract | [Full Text](#)

Keywords: agroforestry, coffee, soil fertility, watershed

Calcium Has Been a Neglected Nutrient in Oil Palm Cultivation

Pages 388-399

Shamshuddin, J.; Fauziah, C.I.; & Syed Omar, S.R.

▼ Abstract | [Full Text](#)

Keywords: Acidic soil, aluminium toxicity, Ca-mineral, highly weathered soil, oil palm

Assessing The Role Of Land Use In Enhancing AMF Growth, Phosphorus Solubilization, And Cowpea (*Vigna unguiculata*) Productivity

Pages 400-411

Nwokocha Amarachi G.; Idris Sani; Joseph Chibueze M.; & Fagbola O.

▼ Abstract | [Full Text](#)

Keywords: Rhizosphere, Arbuscular Mycorrhizal, Phosphate solubilization, Cowpea, Land use

Assessment of Retention Factors and Morphological Studies of Clayey Sand Soil in Conjunction with Polyvinylpyrrolidone

Pages 412-423

Tewatia Nisha; Jabin Shagufta; & Kumar Loveneesh

▼ Abstract | [Full Text](#)

Keywords: Polyvinylpyrrolidone, scanning electron microscopy, soil-amino acid interaction, soil-thin layer chromatography, X-ray diffraction

Effect of 2-chloroethylphosphonic Acid Concentration on Growth Performance and Nutrient Status of Sweet Corn at Vegetative Stage Grown on Rasau Soil

Pages 424-436

Abbas Hartinee; Wahab Zakaria; Katimon Ayob; and Abdul Rahman Zaharah

▼ Abstract | [Full Text](#)

Keywords: ethephon, sweet corn, root growth, nutrient uptake, crop productivity

Effects of Soil Types and Fertilizers on Growth Performance of RevoTropix Paulownia

Pages 437-446

Nur Bazilah Ismail; Geoffery James Gerusu; Keeren Sundara Rajoo; Daljit Singh Karam; & Dzarifah Mohamed Zulperi

▼ Abstract | [Full Text](#)

Keywords: Ultisols, Spodosols, RevoTropix Paulownia, Sarawak

Relationship of Rice Straw at Varying Water Depth Conditions on Methane (CH₄) Emissions

Pages 447-463

Mohd Ridzuan, Rabeatuladawiah; Abdul Sukor, Arina Shalrah; Ishak, Che Fauziah; & Sulaiman, Muhammad Firdaus

▼ Abstract | [Full Text](#)

Keywords: Methane emission, burnt rice straw, unburnt rice straw, continuous flooding system, simple meta-analysis



Soil Chemical Characteristics And Soil Fertility Status In Coffee Agroforestry In The Upper Ciliwung Watershed

Fitri, R.¹; Simangunsong, N.I.¹; Danniswari, D.^{1*} and Taki, H.M.²

¹Department of Landscape Architecture, Faculty of Landscape Architecture and Environmental Technology, Universitas Trisakti, Indonesia

²Department of Urban and Regional Planning, Faculty of Landscape Architecture and Environmental Technology, Universitas Trisakti, Indonesia

*Corresponding author: dibyanti@trisakti.ac.id

ABSTRACT

The conversion of forest land to coffee agroforestry in the Upper Ciliwung Watershed may result in accelerated leaching and impoverishment of soil quality and decreased litterfall. This study aims to analyse the status of soil chemical fertility in coffee agroforestry in the Upper Ciliwung Watershed. This research used the field survey method with a qualitative data analysis approach. The results showed that the status of soil fertility in the aspect of soil chemistry is in the medium - very high category. For the cation exchange capacity (CEC) parameter, the values were 36.02 cmol/kg (high) in Cibulao, 25.45 cmol/kg (high) in Cikoneng, 24.46 cmol/kg (medium) in Rawa Gede, and 38.85 cmol/kg (high) in Cisuren. Furthermore, the P₂O₅ parameter values were 187.8 mg/100g (very high) in Cibulao, 38.6 mg/100g (medium) in Cikoneng, 39.2 mg/100g (medium) in Rawa Gede, and 22.6 mg/100g (medium) in Cisuren. K₂O was 610.6 mg/100g (very high) in Cibulao, 87.5 mg/100g (very high) in Cikoneng, 306.9 mg/100g (very high) in Rawa Gede, and 31.1 mg/100g (medium) in Cisuren, and C-organic was 6.00% (very high) in Cibulao, 7.59% (very high) in Cikoneng, 6.39% (very high) in Rawa Gede and 17.33% (very high) in Cisuren. The findings of this study contribute to understanding the factors maintaining good soil health and fertility under coffee agroforestry in the Upper Ciliwung Watershed, while also providing a foundation for sustainable land management and ensure a productive coffee agroforestry.

Keywords: agroforestry, coffee, soil fertility, watershed

INTRODUCTION

The Ciliwung watershed is one of the priority watersheds to be restored because it supports the downstream area of Jakarta. The upstream area of the Ciliwung watershed has many coffee agroforestry practices, especially in Cibulao, Rawa Gede, Cikoneng, and Cisuren (Fitri *et al.*, 2023; Fitri *et al.*, 2024). Land use in the Upper Ciliwung Watershed has the potential for land degradation and decreased soil fertility. Soil as a medium for plant growth, soil is able to provide organic matter, a place for plant roots to develop, a habitat for soil flora and fauna and soil is also a place to provide nutrients and water for plants (Arsyad 2006). The need to know the status of soil fertility and the ability of soil to provide nutrients to support plant growth (Yadav *et al.*, 2023). Soil fertility is the ability of soil to receive, store and distribute energy for plant growth and development. The fertility of a soil is also influenced by soil properties and slope (Jamaluddin *et al.*, 2022; Rahmayanti, 2017).

Analysis of soil chemical properties can indicate the level of soil fertility in supporting plant growth (Arévalo-Gardini *et al.*, 2015). It is important to assess the chemical properties and soil fertility status of land used for coffee agroforestry activities to identify constraints in the area (Martunis *et al.*, 2017). The problem with coffee agroforestry is that it requires good soil management in the form of applying sufficient fertilizer to meet the nutrient needs of plants (Erlansyah *et al.*, 2022; Jawang, 2021; Suleman *et al.*, 2016). Diversification of land use, intensive land utilization and inadequate land use planning lead to degradation of soil properties and low soil fertility (Aji dan Arifin., 2024; Kharal *et al.*, 2018). It takes a long time to improve

soil quality degraded by intensive land use (Veldkamp et al. 2020; Sena *et al.*, 2021). Improper management of coffee agroforestry can lead to soil degradation and erosion (Kurniawan *et al.*, 2024). Furthermore, agroforestry land in steep slope area would have more degraded soil chemical properties compared to land in gentle slope (Setyastika *et al.*, 2022). Research on the assessment of soil fertility status is important in coffee agroforestry in order to know the ability of the soil to provide nutrients for the growth of coffee plants. An existing study explored the soil physical characteristics of dry agricultural lands in the Upper Ciliwung Watershed (Fitri *et al.*, 2023). To the best of authors' knowledge, there is no information on the soil chemical characteristics of coffee agroforestry lands in Upper Ciliwung Watershed. Therefore, this study was conducted to analyze the status of soil chemical characteristics and soil fertility in coffee agroforestry in the Upper Ciliwung Watershed.

MATERIALS AND METHODS

This research was conducted in the Upper Ciliwung Watershed in West Java Province, the research was conducted from May to September 2023. The research was conducted using a survey method at the time of soil sampling, and the analysis of soil chemical properties was carried out in the laboratory of the Department of Soil Science, Soil and Land Resources, Faculty of Agricultural Sciences, IPB University. Soil chemical properties analyzed included cation exchange capacity, total P and K content, and organic C content. Soil samples were taken from four locations in the Upper Ciliwung Watershed, namely Cibulao, Rawa Gede, Cikoneng and Cisuren. Soil samples were collected at depths of 0 to 30 cm and 30 to 60 cm, then composited depending on depth and analyzed in the laboratory. Determination of soil fertility status based on "Technical Guidelines for Soil Fertility Assessment PPT (1995)".

RESULTS AND DISCUSSION

Soil chemical characteristics in coffee agroforestry in the Upper Ciliwung Watershed based on the results of the soil chemical properties test, the highest soil CEC value in Cisuren was 38.85 me/100g, P₂O₅ parameter value obtained 187.8 mg/100g (very high) in Cibulao, K₂O value of 610.6 mg/100g (very high) in Cibulao, 87.5 mg/100g (very high) in Cikoneng, 306.9 mg/100g (very high) in Rawa Gede. C-organic values in all coffee agroforestry locations were obtained very high at 6.00% in Cibulao, 7.59% (very high) in Cikoneng, 6.39% (very high) in Rawa Gede and 17.33% (very high) in Cisuren. The results of the analysis of soil chemical properties are presented in **Table 1**.

Table 1: Soil Chemical Properties in Coffee Agroforestry in the Upper Ciliwung Watershed

Location	CEC (cmol/kg)	P ₂ O ₅ (mg/100g)	K ₂ O (mg/100g)	C-organic (%)
Cibulao	36.02	187.8	610.6	6.00
Cikoneng	25.45	39.2	87.5	7.59
Rawa Gede	24.46	38.6	306.9	6.38
Cisuren	38.85	22.5	31.1	17.33

Cation exchange capacity (CEC)

One of the soil chemical properties that is closely related to the availability of nutrients for plants is cation exchange capacity, the value of cation exchange capacity is an indicator of soil fertility. Cation exchange capacity on coffee agroforestry land in the Upper Ciliwung Watershed based on the results of laboratory analysis obtained CEC values that vary in all locations. The highest CEC value is found in coffee agroforestry in Cisuren which is 38.85 cmol/kg which is included in the high class. The lowest CEC value was found in coffee agroforestry in Rawa Gede at 24.46 cmol/kg.

Cation exchange capacity is the ability of soil colloids to absorb and exchange cations. A study found that the CEC level of rubber agroforestry land in North Sumatera is low (Muhdi *et al.*, 2023). Despite having the same land use as agroforestry land, the study area in Upper Ciliwung Watershed, West Java, has higher CEC value than in North Sumatera. The predominant soil type in Upper Ciliwung Hulu Watershed, West Java, is Inceptisol (latosol) (Fitri, 2020), and latosol soil type tend to be clayey or have high clay content (Roehrs *et al.*, 2020). Meanwhile, the predominant soil type in North Sumatera is Litosol (Bappeda Sumatera Utara, 2013), and Litosol type is usually sandy (Sari, 2021).

The CEC value is strongly influenced by the type of clay, clay content, soil texture, and organic matter content contained in the soil. (Putri *et al.*, 2019; Yunanto *et al.*, 2022). Soil with higher clay content tend to have higher CEC (Sufardi *et al.*, 2020). This shows that CEC is not influenced by the agroforestry type of land use but rather influenced by soil type and clay content.

P₂O₅

P₂O₅ parameters based on the results of laboratory analysis contained in coffee agroforestry in the class obtained very high to moderate. A significantly high P₂O₅ values was found in Cibulao at 187.8 mg/100g, while other locations show medium values ranging from 22.6 mg/100g to 39.2 mg/100g, and the lowest was in Cisuren. Phosphorus level often become the limiting factor for many crop productions because P quantity is generally small in natural ecosystems (Pradhan *et al.*, 2020). The high value of P₂O₅ is likely by intensive fertilizer application and long planting of coffee agroforestry while at the Cisuren location the low P₂O₅ value is caused by the washing process.

High concentrations of P elements are generally found on land with gentle slopes, this is due to the leaching process from steep slopes that are deposited on gentle slopes (Jakšić *et al.*, 2021). The type of agroforestry significantly influences the concentration of P₂O₅ in soil, and robusta coffee agroforestry tends to have the highest P₂O₅ level compared to other agroforestry systems, including arabica coffee, cinnamon, rubber, chili, and ginger (Rahmawati, 2001). This is aligned with the findings of this study because Cibulao, the location with highest P₂O₅, is the only location planted with robusta coffee, while other locations are planted with arabica coffee (Fitri *et al.*, 2024).

K₂O

The results of K₂O data analysis based on laboratory tests obtained at the research location are very high and medium. The K₂O value in three coffee agroforestry locations in the Upper Ciliwung Watershed is very high including Cibulao at 610.6 mg/100g (very high), in Cikoneng at 87.5 mg/100g (very high), Rawa Gede at 306.9 mg/100g (very high), and K₂O value of 31.1 mg/100g (medium) in Cisuren.

The total K content in soil is influenced by several factors, such as soil colloid type, wet-dry conditions, soil pH and weathering rate (Suarjana *et al.*, 2015). Existing studies revealed that, in agroforestry systems, higher K₂O tends to be found near tree rows or under the native/shade tree canopies (Fahad *et al.*, 2022; Gota *et al.*, 2024). The sampling point location during the data collection might influence the K₂O values. Other than that, the high K concentration in the study area is likely due to fertilization and agricultural land processing by coffee farmers.

C-Organic

C-organic values in coffee agroforestry in the Upper Ciliwung Watershed ranged from 6.00% -17.33%. The highest soil C-organic value was found in coffee agroforestry in Cisuren with a value of 17.33%. Agroforestry systems tend to have higher C-organic content compared to

monoculture cropping systems (Setyastika *et al.*, 2022). Compared to crop monocultures, agroforestry increased C-organic by 40% (Muchane *et al.*, 2020). High C-organic is thought to be influenced by the vegetation that grows on it, the greater number of trees will also increase the C-organic value (Gunawan *et al.*, 2018; Bahnemiri *et al.*, 2019).

According to a previous study (Fitri *et al.*, 2024), the dominant shade tree in coffee agroforestry of Cisuren is *Pinus merkusii* (pine). Pine trees produce high litter, leaving a thick layer of litter under the tree stands, especially when they are old (Imanuddin *et al.*, 2020). C-organic content is high because there are many dead and fallen plant debris (litterfall) in various stages of decomposition, and accumulate on the forest floor, thus affecting the high C-organic content of the coffee agroforestry land.

Soil fertility status

Soil fertility is an important factor determining crop productivity, the condition of the physical, chemical and biological properties of the soil greatly affects the availability of nutrients for plant growth. Soil chemical fertility status includes soil CEC, total K content, organic matter content, and available P. The results of soil analysis are then linked to the criteria for assessing the fertility status of soil chemical properties, indicating that the status of soil chemical fertility in coffee agroforestry in the Upper Ciliwung Watershed is classified as high. (Table 2).

Table 2: Results of Soil Fertility Status Analysis on Coffee Agroforestry in the Upper Ciliwung Watershed

Location	CEC (cmol/kg)	P ₂ O ₅ (mg/100g)	K ₂ O (mg/100g)	C-organic (%)	Fertility status
Cibulao	36.02	187.8	610.6	6.00	High
Cikoneng	25.45	39.2	87.5	7.59	High
Rawa Gede	24.46	38.6	306.9	6.38	High
Cisuren	38.85	22.5	31.1	17.33	High

The resulting high soil fertility status of coffee agroforestry in the Upper Ciliwung Watershed indicates a good balance of soil nutrients. High soil fertility status is thought to be due to plant roots in coffee agroforestry being able to effectively absorb nutrients needed for vegetative growth and generative development of plants. Agroforestry practices can also improve soil fertility through enrichment of organic matter, tree vegetation, and soil microbial activity. (Dollinger and Jose, 2018). The level of soil fertility is high in coffee agroforestry in the Upper Ciliwung Watershed. This is likely to the combination of trees, legumes and annual crops so as to increase the amount of litter that falls from the trees. Complex agroforestry systems with a high number of tree species and individuals can improve soil physical properties (Bahuguna *et al.*, 2018; Bahnemiri *et al.*, 2019). Nutrient sources are enriched from a balanced biomass input and output system. The largest biomass production from plant debris left in agroforestry systems contributes to nutrient and organic carbon stocks. In general, agroforestry improves soil health and fertility (Muchane *et al.*, 2020).

Agroforestry can revitalize degraded lands by improving nutrient recycling and soil fertility. Trees in these systems draw nutrients from deeper soil layers, increase N fixation, and produce litters that enrich the soil as they decompose (Sileshi *et al.*, 2020). In coffee production, higher CEC and C-organic contribute to larger beans, meanwhile higher available P and K contribute to smaller beans (Yadessa *et al.*, 2020). To gain a deeper understanding of how soil fertility impacts coffee production, further research about the relationship between soil health and coffee yields is encouraged.

CONCLUSION

The quality of soil chemical properties at the Cibulao, Rawa Gede, Cikoneng and Cisuren research sites is different. Despite the differences, overall soil fertility status is highly fertile. The vegetation composition, plant canopy density and the amount of litter influence the chemical properties of soil including cation exchange capacity, total K content, organic matter content, and available P. The results emphasize the importance of site-specific soil management practices in coffee agroforestry within the Upper Ciliwung Watershed. Soil fertility can be enhanced through strategies that promote vegetation diversity, canopy cover, and litter retention. A long-term study to monitor changes in soil chemical properties and fertility under different management practices is encouraged to deepen our understanding of soil dynamics in coffee agroforestry.

ACKNOWLEDGEMENTS

The authors are grateful to the Ministry of Education, Culture, Research and Technology for financial support through the BIMA Grant. The authors would like to thank the Institute for Research and Community Service of Trisakti University for technical support in this research.

REFERENCES

- Aji, A.B. & Arifin, M.M. 2024. Status kesuburan tanah sebagai rekomendasi perbaikan lahan pada berbagai tingkat kemiringan lereng di Kecamatan Wonosalam Kabupaten Jombang. *Agroteknika* 7(1): 1-10.
- Arévalo-Gardini, E., Canto, M., Alegre, J., Loli, O., Julca, A. & Baligar, V. 2015. Changes in soil physical and chemical properties in long term improved natural and traditional agroforestry management systems of Cacao genotypes in Peruvian Amazon. *PLoS ONE* 10: e0132147.
- Arsyad. 2006. Konservasi Tanah dan Air. Bogor (ID): IPB Press.
- Bahnemiri, A.K., Abkenar, K.T., Kooch, Y. & Salehi, A. 2019. Evaluating ecological potential of forest stands based on soil quality indices in Hyrcanian beech forest stands, Northern Iran (Case study: Korkoroud forests in Noshahr). *Journal of Forest Science* 65(10): 397–407.
- Bahuguna, H.S., Chaturvedi, R.K. & Rajwar, G.S. 2018. Carbon sequestration potential of the forest soils of district Tehri Garhwal, Uttarakhand, India. *Tropical Ecology* 59(4): 659–678.
- Dollinger, J. & Jose, S. 2018. Agroforestry for soil health. *Agroforestry System Journals* 92: 213-219.
- Erlansyah, T.Z.F., Azis, M.A. & Dude, S. 2022. Karakteristik sifat kimia dan status kesuburan tanah pada Agrowisata Asmara Garden di Kecamatan Bulango Timur Kabupaten Bone Bolango. *Jurnal Lahan Pertanian Tropis* 1(2): 17-22.
- Fitri, R., Simangunsong, N.I., Danniswari, D. & Taki, H.M. 2024. Mapping the distribution of coffee Agroforestry in the Upper Ciliwung Watershed, West Java, Indonesia. *IOP Conference Series: Earth and Environmental Science* 1290: 012032.
- Fitri, R., Simangunsong, N.I., Danniswari, D. & Taki, H.M. 2024. Coffee agroforestry for soil erosion control in the upstream of Ciliwung Watershed, West Java, Indonesia. *EnvironmentAsia* 17(2).
- Gunawan, Wijayanto, N. & Budi, S.W. 2018. Karakteristik sifat kimia tanah dan status kesuburan tanah pada agroforestri tanaman sayuran berbasis *Eucalyptus* sp. *Jurnal Silvikultur Tropika* 10(2): 63-69.
- Jakšić, S., Ninkov, J., Milić, S., Vasin, J., Živanov, M., Jakšić, D. & Komlen, V. 2021. Influence of slope gradient and aspect on soil organic carbon content in the region of Niš, Serbia. *Sustainability* 13(15): 8332.
- Jamaluddin, A.S., Ibrahim, Z., Karam, D.S., Rajoo, K.S., Jusop, S., Gandaseca, S. & Abdu, A. 2022. Soil fertility status of rehabilitated forest soil in Bintulu, Sarawak after 30 Years of planting. *Malaysian Journal of Soil Science* 26: 133-150.
- Jawang, U.P. 2021. Penilaian status kesuburan dan pengelolaan tanah sawah tadah hujan di Desa Umu Pabal Selatan, Kecamatan Umu Ratu Nggay Barat. *Jurnal Ilmu Pertanian Indonesia* 26(3): 421–427.
- Kharal, S., Khanal, B.R. & Panday, D. 2018. Assessment of soil fertility under different land use systems in Dhading District of Nepal. *Soil Systems* 2(4): 57.

- Martunis, L., Sufardi & Muyassir. 2017. Karakteristik kimia tanah dan status kesuburan tanah beberapa jenis tanah dilahan kering Kabupaten Aceh Besar, Provinsi Aceh (Indonesia). *Jurnal Agrotan* 3(1): 77-90.
- PPT. 1995. Petunjuk Teknis Evaluasi Kesuburan Tanah. Laporan Teknis Pusat Penelitian Tanah No.14. Versi 1,0.1. REP II Project. Bogor: CSAR.
- Putri, O.H., Utami, S.R. & Kurniawan, S. 2019. Sifat kimia tanah pada berbagai penggunaan lahan di UB Forest. *Jurnal Tanah Dan Sumberdaya Lahan* 6(1): 1075–1081.
- Rahmayanti, F.D. 2017. Pengaruh kelas kemiringan dan posisi lereng terhadap kandungan Fe tanah sebagai indikator kualitas lingkungan dan kesuburan tanah pada Alfisol di Desa Gunungsari Kabupaten Tasikmalaya. *AGRISIA - Jurnal Ilmu-Ilmu Pertanian* 9(2): 17–27.
- Sena, K.L., Yeager, K.M., Barton, C.D., Lhotka, J.M., Bond, W.E. & Schindler, K.J. 2021. Development of mine soils in a chronosequence of forestry-reclaimed sites in Eastern Kentucky. *Minerals* 11: 422.
- Suarjana, I.W., Supadma, A.A.N. & Arthagama, I.D.M. 2015. Kajian status kesuburan tanah sawah untuk menentukan anjuran pemupukan berimbang spesifik lokasi tanaman padi di Kecamatan Manggis. *E-Jurnal Agroekoteknologi Tropika* 4(4): 314-323.
- Suleman, S., Rajamuddin, U.A. & Isrun. 2016. Penilaian kualitas tanah pada beberapa tipe penggunaan lahan di Kecamatan Sigi Biromaru Kabupaten Sigi. *Agrotekbis: Jurnal Ilmu Pertanian* 4(6): 712–718.
- Veldkamp, E., Schmidt, M., Powers, J.S. & Corre, M.D. 2020. Deforestation and reforestation impacts on soils in the tropics. *Nature Reviews Earth & Environment* 1-16.
- Yadav, A., Yadav, K. & Abd-Elsalam, K.A. 2023. Nanofertilizers: Types, delivery and advantages in agricultural sustainability. *Agrochemicals* 2(2): 296–336.
- Yunanto, T., Amanah, F., Wulansari, A.R. & Wisnu, N.P. 2022. Effect of soil properties on plant growth and diversity at various ages of coal mine reclamation in Indonesia. *Biodiversitas* 23(1): 459-468.

Soil Chemical Characteristics And Soil Fertility Status

by Silia Yuslim FALTL

Submission date: 13-Feb-2025 11:15AM (UTC+0700)

Submission ID: 2545291998

File name: V28_32.pdf (307.25K)

Word count: 3347

Character count: 17725



Soil Chemical Characteristics And Soil Fertility Status In Coffee Agroforestry In The Upper Ciliwung Watershed

Fitri, R.¹; Simangunsong, N.I.¹; Danniswari, D.^{1*} and Taki, H.M.²

¹Department of Landscape Architecture, Faculty of Landscape Architecture and Environmental Technology, Universitas Trisakti, Indonesia

²Department of Urban and Regional Planning, Faculty of Landscape Architecture and Environmental Technology, Universitas Trisakti, Indonesia

*Corresponding author: dibyanti@trisakti.ac.id

ABSTRACT

The conversion of forest land to coffee agroforestry in the Upper Ciliwung Watershed may result in accelerated leaching and impoverishment of soil quality and decreased litterfall. This study aims to analyse the status of soil chemical fertility in coffee agroforestry in the Upper Ciliwung Watershed. This research used the field survey method with a qualitative data analysis approach. The results showed that the status of soil fertility in the aspect of soil chemistry is in the medium - very high category. For the cation exchange capacity (CEC) parameter, the values were 36.02 cmol/kg (high) in Cibulao, 25.45 cmol/kg (high) in Cikoneng, 24.46 cmol/kg (medium) in Rawa Gede, and 38.85 cmol/kg (high) in Cisuren. Furthermore, the P₂O₅ parameter values were 187.8 mg/100g (very high) in Cibulao, 38.6 mg/100g (medium) in Cikoneng, 39.2 mg/100g (medium) in Rawa Gede, and 22.6 mg/100g (medium) in Cisuren. K₂O was 610.6 mg/100g (very high) in Cibulao, 87.5 mg/100g (very high) in Cikoneng, 306.9 mg/100g (very high) in Rawa Gede, and 31.1 mg/100g (medium) in Cisuren, and C-organic was 6.00% (very high) in Cibulao, 7.59% (very high) in Cikoneng, 6.39% (very high) in Rawa Gede and 17.33% (very high) in Cisuren. The findings of this study contribute to understanding the factors maintaining good soil health and fertility under coffee agroforestry in the Upper Ciliwung Watershed, while also providing a foundation for sustainable land management and ensure a productive coffee agroforestry.

Keywords: agroforestry, coffee, soil fertility, watershed

INTRODUCTION

The Ciliwung watershed is one of the priority watersheds to be restored because it supports the downstream area of Jakarta. The upstream area of the Ciliwung watershed has many coffee agroforestry practices, especially in Cibulao, Rawa Gede, Cikoneng, and Cisuren (Fitri *et al.*, 2023; Fitri *et al.*, 2024). Land use in the Upper Ciliwung Watershed has the potential for land degradation and decreased soil fertility. Soil as a medium for plant growth, soil is able to provide organic matter, a place for plant roots to develop, a habitat for soil flora and fauna and soil is also a place to provide nutrients and water for plants (Arsyad 2006). The need to know the status of soil fertility and the ability of soil to provide nutrients to support plant growth (Yadav *et al.*, 2023). Soil fertility is the ability of soil to receive, store and distribute energy for plant growth and development. The fertility of a soil is also influenced by soil properties and slope (Jamaluddin *et al.*, 2022; Rahmayanti, 2017).

Analysis of soil chemical properties can indicate the level of soil fertility in supporting plant growth (Arévalo-Gardini *et al.*, 2015). It is important to assess the chemical properties and soil fertility status of land used for coffee agroforestry activities to identify constraints in the area (Martunis *et al.*, 2017). The problem with coffee agroforestry is that it requires good soil management in the form of applying sufficient fertilizer to meet the nutrient needs of plants (Erlansyah *et al.*, 2022; Jawang, 2021; Suleman *et al.*, 2016). Diversification of land use, intensive land utilization and inadequate land use planning lead to degradation of soil properties and low soil fertility (Aji dan Arifin., 2024; Kharal *et al.*, 2018). It takes a long time to improve

soil quality degraded by intensive land use (Veldkamp et al. 2020; Sena et al., 2021). Improper management of coffee agroforestry can lead to soil degradation and erosion (Kurniawan et al., 2024). Furthermore, agroforestry land in steep slope area would have more degraded soil chemical properties compared to land in gentle slope (Setyastika et al., 2022). Research on the assessment of soil fertility status is important in coffee agroforestry in order to know the ability of the soil to provide nutrients for the growth of coffee plants. An existing study explored the soil physical characteristics of dry agricultural lands in the Upper Ciliwung Watershed (Fitri et al., 2023). To the best of authors' knowledge, there is no information on the soil chemical characteristics of coffee agroforestry lands in Upper Ciliwung Watershed. Therefore, this study was conducted to analyze the status of soil chemical characteristics and soil fertility in coffee agroforestry in the Upper Ciliwung Watershed.

1 MATERIALS AND METHODS

This research was conducted in the Upper Ciliwung Watershed in West Java Province, the research was conducted from May to September 2023. The research was conducted using a survey method at the time of soil sampling, and the analysis of soil chemical properties was carried out in the laboratory of the Department of Soil Science, Soil and Land Resources, Faculty of Agricultural Sciences, IPB University. Soil chemical properties analyzed included cation exchange capacity, total P and K content, and organic C content. Soil samples were taken from four locations in the Upper Ciliwung Watershed, namely Cibulao, Rawa Gede, Cikoneng and Cisuren. Soil samples were collected at depths of 0 to 30 cm and 30 to 60 cm, then composited depending on depth and analyzed in the laboratory. Determination of soil fertility status based on "Technical Guidelines for Soil Fertility Assessment PPT (1995)".

1.1 RESULTS AND DISCUSSION

Soil chemical characteristics in coffee agroforestry in the Upper Ciliwung Watershed based on the results of the soil chemical properties test, the highest soil CEC value in Cisuren was 38.85 me/100g, P₂O₅ parameter value obtained 187.8 mg/100g (very high) in Cibulao, K₂O value of 610.6 mg/100g (very high) in Cibulao, 87.5 mg/100g (very high) in Cikoneng, 306.9 mg/100g (very high) in Rawa Gede. C-organic values in all coffee agroforestry locations were obtained very high at 6.00% in Cibulao, 7.59% (very high) in Cikoneng, 6.39% (very high) in Rawa Gede and 17.33% (very high) in Cisuren. The results of the analysis of soil chemical properties are presented in Table 1.

Table 1: Soil Chemical Properties in Coffee Agroforestry in the Upper Ciliwung Watershed

Location	CEC (cmol/kg)	P ₂ O ₅ (mg/100g)	K ₂ O (mg/100g)	C-organic (%)
Cibulao	36.02	187.8	610.6	6.00
Cikoneng	25.45	39.2	87.5	7.59
Rawa Gede	24.46	38.6	306.9	6.38
Cisuren	38.85	22.5	31.1	17.33

Cation exchange capacity (CEC)

One of the soil chemical properties that is closely related to the availability of nutrients for plants is cation exchange capacity, the value of cation exchange capacity is an indicator of soil fertility. Cation exchange capacity on coffee agroforestry land in the Upper Ciliwung Watershed based on the results of laboratory analysis obtained CEC values that vary in all locations. The highest CEC value is found in coffee agroforestry in Cisuren which is 38.85 cmol/kg which is included in the high class. The lowest CEC value was found in coffee agroforestry in Rawa Gede at 24.46 cmol/kg.

Cation exchange capacity is the ability of soil colloids to absorb and exchange cations. A study found that the CEC level of rubber agroforestry land in North Sumatera is low (Muhdi *et al.*, 2023). Despite having the same land use as agroforestry land, the study area in Upper Ciliwung Watershed, West Java, has higher CEC value than in North Sumatera. The predominant soil type in Upper Ciliwung Hulu Watershed, West Java, is Inceptisol (latosol) (Fitri, 2020), and latosol soil type tend to be clayey or have high clay content (Roehrs *et al.*, 2020). Meanwhile, the predominant soil type in North Sumatera is Litosol (Bappeda Sumatera Utara, 2013), and Litosol type is usually sandy (Sari, 2021).

The CEC value is strongly influenced by the type of clay, clay content, soil texture, and organic matter content contained in the soil. (Putri *et al.*, 2019; Yunanto *et al.*, 2022). Soil with higher clay content tend to have higher CEC (Sufardi *et al.*, 2020). This shows that CEC is not influenced by the agroforestry type of land use but rather influenced by soil type and clay content.

P₂O₅

P₂O₅ parameters based on the results of laboratory analysis contained in coffee agroforestry in the class obtained very high to moderate. A significantly high P₂O₅ values was found in Cibulao at 187.8 mg/100g, while other locations show medium values ranging from 22.6 mg/100g to 39.2 mg/100g, and the lowest was in Cisuren. Phosphorus level often become the limiting factor for many crop productions because P quantity is generally small in natural ecosystems (Pradhan *et al.*, 2020). The high value of P₂O₅ is likely by intensive fertilizer application and long planting of coffee agroforestry while at the Cisuren location the low P₂O₅ value is caused by the washing process.

High concentrations of P elements are generally found on land with gentle slopes, this is due to the leaching process from steep slopes that are deposited on gentle slopes (Jakšić *et al.*, 2021). The type of agroforestry significantly influences the concentration of P₂O₅ in soil, and robusta coffee agroforestry tends to have the highest P₂O₅ level compared to other agroforestry systems, including arabica coffee, cinnamon, rubber, chili, and ginger (Rahmawati, 2001). This is aligned with the findings of this study because Cibulao, the location with highest P₂O₅, is the only location planted with robusta coffee, while other locations are planted with arabica coffee (Fitri *et al.*, 2024).

K₂O

The results of K₂O data analysis based on laboratory tests obtained at the research location are very high and medium. The K₂O value in three coffee agroforestry locations in the Upper Ciliwung Watershed is very high including Cibulao at 610.6 mg/100g (very high), in Cikoneng at 87.5 mg/100g (very high), Rawa Gede at 306.9 mg/100g (very high), and K₂O value of 31.1 mg/100g (medium) in Cisuren.

The total K content in soil is influenced by several factors, such as soil colloid type, wet-dry conditions, soil pH and weathering rate (Suarjana *et al.*, 2015). Existing studies revealed that, in agroforestry systems, higher K₂O tends to be found near tree rows or under the native/shade tree canopies (Fahad *et al.*, 2022; Gota *et al.*, 2024). The sampling point location during the data collection might influence the K₂O values. Other than that, the high K concentration in the study area is likely due to fertilization and agricultural land processing by coffee farmers.

C-Organic

C-organic values in coffee agroforestry in the Upper Ciliwung Watershed ranged from 6.00% -17.33%. The highest soil C-organic value was found in coffee agroforestry in Cisuren with a value of 17.33%. Agroforestry systems tend to have higher C-organic content compared to

monoculture cropping systems (Setyastika *et al.*, 2022). Compared to crop monocultures, agroforestry increased C-organic by 40% (Muchane *et al.*, 2020). High C-organic is thought to be influenced by the vegetation that grows on it, the greater number of trees will also increase the C-organic value (Gunawan *et al.*, 2018; Bahnemiri *et al.*, 2019).

According to a previous study (Fitri *et al.*, 2024), the dominant shade tree in coffee agroforestry of Cisuren is *Pinus merkusii* (pine). Pine trees produce high litter, leaving a thick layer of litter under the tree stands, especially when they are old (Imanuddin *et al.*, 2020). C-organic content is high because there are many dead and fallen plant debris (litterfall) in various stages of decomposition, and accumulate on the forest floor, thus affecting the high C-organic content of the coffee agroforestry land.

Soil fertility status

Soil fertility is an important factor determining crop productivity, the condition of the physical, chemical and biological properties of the soil greatly affects the availability of nutrients for plant growth. Soil chemical fertility status includes soil CEC, total K content, organic matter content, and available P. The results of soil analysis are then linked to the criteria for assessing the fertility status of soil chemical properties, indicating that the status of soil chemical fertility in coffee agroforestry in the Upper Ciliwung Watershed is classified as high. (Table 2).

Table 2: Results of Soil Fertility Status Analysis on Coffee Agroforestry in the Upper Ciliwung Watershed

Location	CEC (cmol/kg)	P ₂ O ₅ (mg/100g)	K ₂ O (mg/100g)	C-organic (%)	Fertility status
Cibulao	36.02	187.8	610.6	6.00	High
Cikoneng	25.45	39.2	87.5	7.59	High
Rawa Gede	24.46	38.6	306.9	6.38	High
Cisuren	38.85	22.5	31.1	17.33	High

The resulting high soil fertility status of coffee agroforestry in the Upper Ciliwung Watershed indicates a good balance of soil nutrients. High soil fertility status is thought to be due to plant roots in coffee agroforestry being able to effectively absorb nutrients needed for vegetative growth and generative development of plants. Agroforestry practices can also improve soil fertility through enrichment of organic matter, tree vegetation, and soil microbial activity. (Dollinger and Jose, 2018). The level of soil fertility is high in coffee agroforestry in the Upper Ciliwung Watershed. This is likely to the combination of trees, legumes and annual crops so as to increase the amount of litter that falls from the trees. Complex agroforestry systems with a high number of tree species and individuals can improve soil physical properties (Bahuguna *et al.*, 2018; Bahnemiri *et al.*, 2019). Nutrient sources are enriched from a balanced biomass input and output system. The largest biomass production from plant debris left in agroforestry systems contributes to nutrient and organic carbon stocks. In general, agroforestry improves soil health and fertility (Muchane *et al.*, 2020).

Agroforestry can revitalize degraded lands by improving nutrient recycling and soil fertility. Trees in these systems draw nutrients from deeper soil layers, increase N fixation, and produce litters that enrich the soil as they decompose (Sileshi *et al.*, 2020). In coffee production, higher CEC and C-organic contribute to larger beans, meanwhile higher available P and K contribute to smaller beans (Yadessa *et al.*, 2020). To gain a deeper understanding of how soil fertility impacts coffee production, further research about the relationship between soil health and coffee yields is encouraged.

CONCLUSION

The quality of soil chemical properties at the Cibulao, Rawa Gede, Cikongeng and Cisuren research sites is different. Despite the differences, overall soil fertility status is highly fertile. The vegetation composition, plant canopy density and the amount of litter influence the chemical properties of soil including cation exchange capacity, total K content, organic matter content, and available P. The results emphasize the importance of site-specific soil management practices in coffee agroforestry within the Upper Ciliwung Watershed. Soil fertility can be enhanced through strategies that promote vegetation diversity, canopy cover, and litter retention. A long-term study to monitor changes in soil chemical properties and fertility under different management practices is encouraged to deepen our understanding of soil dynamics in coffee agroforestry.

ACKNOWLEDGEMENTS

The authors are grateful to the Ministry of Education, Culture, Research and Technology for financial support through the BIMA Grant. The authors would like to thank the Institute for Research and Community Service of Trisakti University for technical support in this research.

REFERENCES

- Aji, A.B. & Arifin, M.M. 2024. Status kesuburan tanah sebagai rekomendasi perbaikan lahan pada berbagai tingkat kemiringan lereng di Kecamatan Wonosalam Kabupaten Jombang. *Agroteknika* 7(1): 1-10.
- Arévalo-Gardini, E., Canto, M., Alegre, J., Loli, O., Julca, A. & Baligar, V. 2015. Changes in soil physical and chemical properties in long term improved natural and traditional agroforestry management systems of Cacao genotypes in Peruvian Amazon. *PLoS ONE* 10: e0132147.
- Arsyad. 2006. *Konservasi Tanah dan Air*. Bogor (ID): IPB Press.
- Bahnemiri, A.K., Abkenar, K.T., Kooch, Y. & Salehi, A. 2019. Evaluating ecological potential of forest stands based on soil quality indices in Hyrcanian beech forest stands, Northern Iran (Case study: Korkoroud forests in Noshahr). *Journal of Forest Science* 65(10): 397-407.
- Bahuguna, H.S., Chaturvedi, R.K. & Rajwar, G.S. 2018. Carbon sequestration potential of the forest soils of district Tehri Garhwal, Uttarakhand, India. *Tropical Ecology* 59(4): 659-678.
- Dollinger, J. & Jose, S. 2018. Agroforestry for soil health. *Agroforestry System Journals* 92: 213-219.
- Erlansyah, T.Z.F., Azis, M.A. & Dude, S. 2022. Karakteristik sifat kimia dan status kesuburan tanah pada Agrowisata Asmara Garden di Kecamatan Bulango Timur Kabupaten Bone Bolango. *Jurnal Lahan Pertanian Tropis* 1(2): 17-22.
- Fitri, R., Simangunsong, N.I., Danniswari, D. & Taki, H.M. 2024. Mapping the distribution of coffee Agroforestry in the Upper Ciliwung Watershed, West Java, Indonesia. *IOP Conference Series: Earth and Environmental Science* 1290: 012032.
- Fitri, R., Simangunsong, N.I., Danniswari, D. & Taki, H.M. 2024. Coffee agroforestry for soil erosion control in the upstream of Ciliwung Watershed, West Java, Indonesia. *EnvironmentAsia* 17(2).
- Gunawan, Wijayanto, N. & Budi, S.W. 2018. Karakteristik sifat kimia tanah dan status kesuburan tanah pada agroforestri tanaman sayuran berbasis *Eucalyptus* sp. *Jurnal Silvikultur Tropika* 10(2): 63-69.
- Jakšić, S., Ninkov, J., Milić, S., Vasin, J., Živanov, M., Jakšić, D. & Komlen, V. 2021. Influence of slope gradient and aspect on soil organic carbon content in the region of Niš, Serbia. *Sustainability* 13(15): 8332.
- Jamaluddin, A.S., Ibrahim, Z., Karam, D.S., Rajoo, K.S., Jusop, S., Gandaseca, S. & Abdu, A. 2022. Soil fertility status of rehabilitated forest soil in Bintulu, Sarawak after 30 Years of planting. *Malaysian Journal of Soil Science* 26: 133-150.
- Jawang, U.P. 2021. Penilaian status kesuburan dan pengelolaan tanah sawah tadah hujan di Desa Uumbu Pabal Selatan, Kecamatan Uumbu Ratu Nggay Barat. *Jurnal Ilmu Pertanian Indonesia* 26(3): 421-427.
- Kharal, S., Khanal, B.R. & Panday, D. 2018. Assessment of soil fertility under different land use systems in Dhading District of Nepal. *Soil Systems* 2(4): 57.

- Martunis, L., Sufardi & Muyassir. 2017. Karakteristik kimia tanah dan status kesuburan tanah beberapa jenis tanah dilahan kering Kabupaten Aceh Besar, Provinsi Aceh (Indonesia). *Jurnal Agrotan* 3(1): 77-90.
- PPT. 1995. Petunjuk Teknis Evaluasi Kesuburan Tanah. Laporan Teknis Pusat Penelitian Tanah No.14. Versi 1.0.1. REP II Project. Bogor: CSAR.
- Putri, O.H., Utami, S.R. & Kurniawan, S. 2019. Sifat kimia tanah pada berbagai penggunaan lahan di UB Forest. *Jurnal Tanah Dan Sumberdaya Lahan* 6(1): 1075–1081.
- Rahmayanti, F.D. 2017. Pengaruh kelas kemiringan dan posisi lereng terhadap kandungan Fe tanah sebagai indikator kualitas lingkungan dan kesuburan tanah pada Alfisol di Desa Gunungsari Kabupaten Tasikmalaya. *AGRISIA - Jurnal Ilmu-Ilmu Pertanian* 9(2): 17–27.
- Sena, K.L., Yeager, K.M., Barton, C.D., Lhotka, J.M., Bond, W.E. & Schindler, K.J. 2021. Development of mine soils in a chronosequence of forestry-reclaimed sites in Eastern Kentucky. *Minerals* 11: 422.
- Suarjana, I.W., Supadma, A.A.N. & Arthagama, I.D.M. 2015. Kajian status kesuburan tanah sawah untuk menentukan anjuran pemupukan berimbang spesifik lokasi tanaman padi di Kecamatan Manggis. *E-Jurnal Agroekoteknologi Tropika* 4(4): 314-323.
- Suleman, S., Rajamuddin, U.A. & Isrun. 2016. Penilaian kualitas tanah pada beberapa tipe penggunaan lahan di Kecamatan Sigi Biromaru Kabupaten Sigi. *Agrotekbis: Jurnal Ilmu Pertanian* 4(6): 712–718.
- Veldkamp, E., Schmidt, M., Powers, J.S. & Corre, M.D. 2020. Deforestation and reforestation impacts on soils in the tropics. *Nature Reviews Earth & Environment* 1-16.
- Yadav, A., Yadav, K. & Abd-Elsalam, K.A. 2023. Nanofertilizers: Types, delivery and advantages in agricultural sustainability. *Agrochemicals* 2(2): 296–336.
- Yunanto, T., Amanah, F., Wulansari, A.R. & Wisnu, N.P. 2022. Effect of soil properties on plant growth and diversity at various ages of coal mine reclamation in Indonesia. *Biodiversitas* 23(1): 459-468.

Soil Chemical Characteristics And Soil Fertility Status

ORIGINALITY REPORT

18%

SIMILARITY INDEX

10%

INTERNET SOURCES

14%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- 1 R Fitri, N I Simangunsong, D Danniswari, H M Taki. "Mapping the Distribution of Coffee Agroforestry in the Upper Ciliwung Watershed, West Java, Indonesia", IOP Conference Series: Earth and Environmental Science, 2024
Publication 6%
- 2 www.smujo.id
Internet Source 3%
- 3 Sulakhudin, Sukirno, A M Abdillah. "Monitoring soil fertility to mitigate soil degradation in reclamation land after bauxite mining", IOP Conference Series: Earth and Environmental Science, 2024
Publication 2%
- 4 msss.com.my
Internet Source 2%
- 5 R Mardhatillah, P Pamoengkas, Darwo. " The Growth of Kapur Tanduk (Burck.) on Different Levels of Canopy Opening and Fertilization ", IOP Conference Series: Earth and Environmental Science, 2019
Publication 1%
- 6 Tuti Mutia, Sumarmi Sumarmi, Ravines Rohit Prasad. "SOIL FERTILITY STATUS IN MANDALA COSTUMARY FOREST BASED ON TOPOGRAPHY LEVELS", GEOGRAPHY : Jurnal 1%

Kajian, Penelitian dan Pengembangan Pendidikan, 2023

Publication

7	journal.ipb.ac.id Internet Source	1%
8	ojs.untika.ac.id Internet Source	1%
9	Submitted to University of Nottingham Student Paper	1%

Exclude quotes On

Exclude matches < 20 words

Exclude bibliography On

Soil Chemical Characteristics And Soil Fertility Status

GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/100

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
