


COVER

<https://www.midwifery.iocspublisher.org/index.php/midwifery/issue/view/33>

Vol. 14 No.3 (2026) : Health Sciences and related fields



DAFTAR ISI

midwifery.iocspublisher.org/index.php/midwifery/issue/view/33

Home / Archives / Vol. 14 No. 3 (2026): Health Sciences and related fields

Vol. 14 No. 3 (2026): Health Sciences and related fields

Published: 2026-07-22

Articles

Combination of slow deep breathing with rose aromatherapy and progressive muscle relaxation on blood pressure among hypertensive patients in Manado City

Faradilla Miftah Suranata, Rahmat Hidayat Djali, Neifa Fitra Takahapis, Bunga Pakaya

519-521

PDF

The effect of giving papaya fruit on the sufficiency of breast milk in post-part mothers

Fenny Fernando, Mella Pebriana, Dewi Fransisca, Ramah Hayu

561-570

PDF

Effectiveness of nutritional counseling and folic acid supplementation on hemoglobin levels of pregnant women in the working area of the Hajimena Health Center, Lampung

Nur Wachmah Mawadda, Hanny Citra Purnama, Nurul Amalia, Qorinah Estiningtyas Sakilah Adhoni, Chyntia Haryana

618-625

PDF

Sigmoid colon perforation complicating neonatal rectovestibular fistula: A case report

Pertogi Napitupulu, Nadifa Agil, Revalita Wahan, Nathalia Ningrum

663-672

PDF

Nutrition care in pediatric patients with severe diabetic ketoacidosis, hyperglycemia and suspected type 1 diabetes

Wahyu Dwi Suzanti, Sukmawati Sukmawati, Sutanti Rauf, Musyawarah Thahir

530-540

PDF

Identifying anemia determinant factors in adolescent girls to improve productivity: A systematic review

Ni Putu Wahyu Laksmi Natha Devi, Luh Seri Anji, Komang Ayu Karlikasari

597-609

PDF

Factors associated with the intention of students at universitas muhammadiyah surakarta to quit smoking

Nova Deslita Agusetyani, Tanjung Anisari Indah Kusumaningrum

643-653

PDF

The effect of classical instrumental music (Kenny G-Style) on reducing anxiety in pregnant women during labor: A literature review

Rut Yohana Girsang, Dime'aarij Agustia, Novita Adeline Sipayung, Dinda Dian Meldita, Ria Fazellia Gullom

622-629

PDF



Editorial Team

Peer Review Process

Reviewer

Focus & Scope

Author Guidelines

Online Submission

Publication Ethics

Copyright Notice

Policy of Screening for Plagiarism

Indexing

Journal Fee

ISSN

p-ISSN: 2088-7689 | E-ISSN: 2721-9453

TEMPLATE





EDITORIAL BOARD

<https://www.midwifery.iocspublisher.org/index.php/midwifery/about/editorialTeam>

Editorial Team

Editor in Chief

- Dr. dr. D. K. Keliat, Spa, ([Scopus Profil](#)), Medan, Universitas Sumatera Utara, Inedonesia

Associate Editors

- Dr. Leila Mohammadinia, Ph.D., MS, BS, ([Scopus Profil](#)) School of Management and Information Sciences, Shiraz University of Medical Sciences, Shiraz, Iran
- Dr. Malakeh Zuhdi Malak, Ph.D., ([Scopus Profil](#)) Associate Professor in Community Health Nursing, Al-Zaytoonah University of Jordan, Amman
- Dr. Satyanarayan Labani, ([Scopus Profil](#)) National Institute of Cancer Prevention & Research, Noida, India
- Joko Gunawan, S.Kep.Ns MN, Ph.D., ([Scopus Profil](#)) Chulalongkrong University, Thailand,
- Prof. Aytac GUDER, PhD., ([Scopus Profil](#)) Department Of Medical Services and Techniques, Vocational High School of Health, Giresun University, Turkey
- Prof. Eileen Savage, ([Scopus Profil](#)) School of Nursing and Midwifery, University College Cork, Ireland
- Prof. Francesco Carelli, FD, MD, MSc., ([Scopus Profil](#)) Professor at University of Milan, Rome and Bari for Undergraduate Students and Specializing Doctors and Masters, Italy
- Prof. Magda Moawad Mohamad Mohsen, PhD., ([Scopus Profil](#)) Professor of Community Health Nursing, Faculty of Nursing, Menoufia University, Egypt
- Prof. Urmila Bhardwaj, ([Scopus Profil](#)) Principal, Rufaida College of Nursing, Jamia Hamdard, New Delhi
- Prof. Yau Sui Yu, PhD., ([Scopus Profil](#)) Associate Professor, School of Nursing and Health Studies, Open University of Hong Kong, Hong Kong
- Prof. Denise Fassett, BHS.RN.MN.PHD, ([Scopus Profil](#)) Faculty Of Health, Departement Nursing and Midfiwifery, University of Tasmania, Australia
- Prof. John Christian Fox, MD, RDMS, FAIUM, ([Scopus Profil](#)) School Of Medicine, University Of California, Irvine, US
- dr.Leo Simanjuntak, SpOG, ([Scopus Profil](#)), Medan, Universitas Sumatera Utara, Inedonesia
- Prof Dr H Aslim Sihotang, SPM-KVR, ([Scopus Profil](#)), Medan, Universitas Sumatera Utara, Inedonesia
- Naety, SST., M.Keb, ([Google Scholar Profil](#)) Medan, Universitas Darma Agung, Indonesia
- Efi Satriana Silalahi, SST, MKM, ([Google Scholar Profil](#)), STIKE Mitra Sejati, Medan, Indonesia
- Ns. Sumandar, S.Kep, M.Kes, ([Google Scholar Profil](#)) STIKes Al-Insyirah Pekanbaru, Riau, Indonesia
- Riny Apriani S.Kep., Ns., M.Kep., ([Google Scholar Profil](#)) STIKES Binalita Sudama Medan, Sumatera Utara, Indonesia

ARTIKEL TERAKREDITASI SINTA 3

<https://sinta.kemdiktisaintek.go.id/journals/profile/6857>

SERTIFIKAT

Direktorat Jendral Pendidikan Tinggi, Riset dan Teknologi
Kementerian Pendidikan, Kebudayaan, Riset dan Teknologi Republik Indonesia





Kutipan dari Keputusan Direktorat Jendral Pendidikan Tinggi, Riset, dan Teknologi
Kementerian Pendidikan, Kebudayaan, Riset dan Teknologi Republik Indonesia

Nomor: 152/E/KPT/2023

Peringkat Akreditasi Jurnal Ilmiah Periode II Tahun 2023

Nama Jurnal Ilmiah:
Science Midwifery

E-ISSN: 27219453
Institute Of Computer Science
Ditetapkan Sebagai Jurnal Ilmiah:

TERAKREDITASI PERINGKAT 3

Akreditasi Berlaku selama 5 (lima) Tahun, yaitu:
Volume 11 Nomor 1 Tahun 2023 sampai Volume 15 Nomor 6 Tahun 2027
Jakarta, 25 September 2023

Plt. Direktur Jendral Pendidikan Tinggi, Riset, dan Teknologi

Prof. Ir. Nizam, M.Sc., DIC, Ph.D., IPU, ASEAN Eng
NIP. 196107061987101001



BUKTI KOMUNIKASI DAN SUBMISSION

Tab Baru

iocspublisher.org/index.php/midwifery/authorDashboard/submission/2426#

Round 1

Round 1 Status
All reviewers have responded and a decision is needed.

Review Discussions [Add discussion](#)

Name	From	Last Reply	Replies	Closed
Accepted submission with minor revision	midwifery 2026-08-04 04:28 AM	partogi 2026-08-04 03:34 PM	1	☐
▶ Manuscript after revision as sugested	partogi 2026-08-04 06:38 AM	-	0	☐
▶ Manuscript after revision as sugested	partogi 2026-08-04 03:43 PM	-	0	☐

Pencarian

[Submission](#) [Review](#) [Copyediting](#) [Production](#)

Round 1

Round 1 Status
All reviewers have responded and a decision is needed.

Review Discussions [Add discussion](#)

Name	From	Last Reply	Replies	Closed
Accepted submission with minor revision	midwifery 2026-08-04 04:28 AM	partogi 2026-08-04 03:34 PM	1	☐
▶ Manuscript after revision as sugested	partogi 2026-08-04 06:38 AM	-	0	☐

Round 1

Round 1 Status

All reviewers have responded and a decision is needed.

Review Discussions

[Add discussion](#)

Name	From	Last Reply	Replies	Closed
Accepted submission with minor revision	midwifery 2026-08-04 04:28 AM	partogi 2026-08-04 03:34 PM	1	☐
▶ Manuscript after revision as sugested	partogi 2026-08-04 06:38 AM	-	0	☐
▶ Manuscript after revision as sugested	partogi 2026-08-04 03:43 PM	-	0	☐



Pencarian



ASUS Vivobook

Workflow

Publication

Submission

Review

Copyediting

Production

Copyediting Discussions

Add discussion

Name	From	Last Reply	Replies	Closed
No Items				

Q Search

Copyedited

		August 18, 2026	Article Text
15655	Partogi Napitupulu.doc		

Gmail

Search mail

Active

2 of 1,836

Compose

Inbox 923

Starred

Snoozed

Drafts 17

Categories

Purchases

More

Labels +

←

📅

🕒

🗑️

✉️

🕒

🔄

📁

🗨️

⋮

[midwifery] Editor Decision

External

Inbox x

Juliana Batubara

<midwiferyofficial@gmail.com>

to me

6:01 PM (1 hour ago)

☆

😊

↩️

⋮

Partogi Napitupulu:

The editing of your submission, "275 Sigmoid Colon Perforation Complicating Neonatal Rectovestibular Fistula: A Case Report," is complete. We are now sending it to production.

Submission URL: <https://midwifery.iospublisher.org/index.php/midwifery/authorDashboard/submission/2426>

Science Midwifery

↩️ Reply

➡️ Forward

😊

🗨️ Share in chat

New

2026-08-07 01:41 AM

IOCSOURCE



Institute of
Computer
Science

LETTER OF ACCEPTANCE

ISSN: 2086-7689 (Print) | 2721-9453 (Online)

Secret ID: 2426

Dear Author(s),

Partogi Napitupulu, Nadifa Agil, Revalita Wahab, Nathalia Ningrum

Warm Greetings!

It's a great pleasure to inform you that, after the peer review process, your article entitled "**275 Sigmoid Colon Perforation Complicating Neonatal Rectovestibular Fistula: A Case Report**" has been **ACCEPTED** and considered for publication in **Science Midwifery** in Volume 14 Issue 3, August (2026).

Thank you for Submitting your work to this journal. We hope you Submit your articles in future.

Warm regards,



Pencarian



ARTIKEL

<https://midwifery.iocspublisher.org/index.php/midwifery/article/view/2253>

Home / Archives / Vol. 14 No. 3 (2026): Health Sciences and related fields / Articles	
Sigmoid colon perforation complicating neonatal rectovestibular fistula: A case report	
Partogi Napitupulu Universitas Trisakti, Indonesia	 PDF
Nadifa Agil Universitas Trisakti, Indonesia	Published 2026-08-18
Revalita Wahab Universitas Trisakti, Indonesia	How to Cite Napitupulu, P. (2026) "Sigmoid colon perforation complicating neonatal rectovestibular fistula: A case report", <i>Science Midwifery</i> , 14(3), pp. 663–672. doi: 10.35335/midwifery.v14i3.2426.
Nathalia Ningrum Universitas Trisakti, Indonesia	More Citation Formats ▼
DOI: https://doi.org/10.35335/midwifery.v14i3.2426	Issue Vol. 14 No. 3 (2026): Health Sciences and related fields
Keywords: Anorectal Malformation , Bowel Obstruction, Pneumoperitonuem, Rectovestibular Fistula, Sigmoid Colon Perforation	Section Articles
Abstract Anorectal malformation (ARM) with rectovestibular fistula may allow small amounts of meconium to pass despite ongoing distal bowel obstruction. We report a three-day-old term female neonate with an imperforate anus, progressive abdominal distension, bilious vomiting, and minimal meconium passage through a rectovestibular fistula. Abdominal radiography showed marked bowel dilatation and pneumoperitoneum. Emergency laparotomy revealed sigmoid colon perforation with diffuse fecal peritonitis. The perforation was repaired, followed by a double-barrel sigmoid colostomy. The infant recovered well and was discharged on postoperative day eight. At three-month follow-up, the wound was completely healed and the colostomy remained healthy and functional. This case illustrates a rare but serious complication of rectovestibular fistula, in which limited meconium passage may coexist with significant bowel obstruction and subsequent perforation. These findings emphasize that thorough	License Copyright (c) 2026 Partogi Napitupulu, Nadifa Agil, Revalita Wahab, Nathalia Ningrum  This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License .

Partogi Napitupulu FK

Sigmoid colon perforation complicating neonatal rectovestibular fistula A case report

 Artikel Jurnal 1

Document Details

Submission ID

trn:oid:::3618:146687734

Submission Date

Jul 28, 2026, 2:58 PM GMT+7

Download Date

Jul 28, 2026, 3:22 PM GMT+7

File Name

Sigmoid colon perforation complicating neonatal rectovestibular fistula A case report.docx

File Size

3.5 MB

11 Pages

3,986 Words

25,885 Characters

17% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

Filtered from the Report

- Bibliography
- Quoted Text

Match Groups

-  **33 Not Cited or Quoted 14%**
Matches with neither in-text citation nor quotation marks
-  **9 Missing Quotations 3%**
Matches that are still very similar to source material
-  **0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 2%  Internet sources
- 2%  Publications
- 15%  Submitted works (Student Papers)

Integrity Flags

0 Integrity Flags for Review

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups

- 33 Not Cited or Quoted 14%**
Matches with neither in-text citation nor quotation marks
- 9 Missing Quotations 3%**
Matches that are still very similar to source material
- 0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
- 0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 2% Internet sources
- 2% Publications
- 15% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Student papers	Trisakti University on 2026-07-21	14%
2	Internet	www.jneonataisurg.com	<1%
3	Publication	Ecem Koyun, Derya Suluhan. "The relationship between the caregiver burden and...	<1%
4	Publication	Phanthila Sitthikarnkha, Sirapoom Niamsanit, Leelawadee Techasatian, Suchaorn...	<1%
5	Student papers	University of Worcester on 2024-04-03	<1%
6	Internet	files01.core.ac.uk	<1%
7	Internet	www.dovepress.com	<1%
8	Student papers	iHeed on 2026-06-08	<1%

Abstract

Anorectal malformation (ARM) with rectovestibular fistula may allow small amounts of meconium to pass despite ongoing distal bowel obstruction. We report a three-day-old term female neonate with an imperforate anus, progressive abdominal distension, bilious vomiting, and minimal meconium passage through a rectovestibular fistula. Abdominal radiography showed marked bowel dilatation and pneumoperitoneum. Emergency laparotomy revealed sigmoid colon perforation with diffuse fecal peritonitis. The perforation was repaired, followed by a double-barrel sigmoid colostomy. The infant recovered well and was discharged on postoperative day eight. At three-month follow-up, the wound was completely healed and the colostomy remained healthy and functional. This case illustrates a rare but serious complication of rectovestibular fistula, in which limited meconium passage may coexist with significant bowel obstruction and subsequent perforation.

Keywords: anorectal malformation; bowel obstruction; pneumoperitoneum; rectovestibular fistula; sigmoid colon perforation

Introduction

Anorectal malformations (ARMs) encompass a heterogeneous group of congenital abnormalities involving the distal rectum and anus, with variable fistulous communications to the urinary or genital tract and frequent associations with other congenital anomalies (Amerstorfer et al., 2022; Hageman et al., 2024). Because the anatomical presentation of ARM varies considerably, accurate identification of the malformation during the neonatal period is important for determining appropriate diagnostic evaluation and subsequent surgical management (Amerstorfer et al., 2022). Careful inspection of the anus and perineum should therefore be an integral component of routine neonatal examination, particularly for identifying abnormalities in anal position, caliber, or fistulous openings (Amerstorfer et al., 2022). Although uncommon, ARM occurs worldwide and shows geographical variation in reported birth prevalence (Kancherla et al., 2023). A large multicountry surveillance study involving 9,438 cases among nearly 29 million births estimated a pooled ARM prevalence of 3.26 per 10,000 total births (Kancherla et al., 2023). Population-based data from Southeast Asia remain comparatively limited; however, a nationwide analysis using Thailand's National Health Security Office database reported an ARM prevalence of 2.57 per 10,000 births between 2017 and 2020 (Sirichamratsakul et al., 2023).

Among female patients with ARM, rectovestibular fistula represents an important anatomical phenotype in which the rectum opens into the vestibule anterior to the expected anal position (Hageman et al., 2024). Data from the European ARM-Net registry, which included 2,619 patients from 34 centers, showed that vestibular fistula accounted for 31.8% of ARM phenotypes among female patients (Hageman et al., 2024). Although a rectovestibular fistula permits passage of meconium and may provide some degree of distal bowel decompression, visible meconium through an abnormal perineal opening should not be interpreted as evidence of normal anorectal anatomy or necessarily adequate intestinal emptying (Long et al., 2024). This distinction is clinically relevant because delayed recognition of ARM continues to occur despite routine neonatal assessment (Kruger et al., 2019; Long et al., 2024). In a prospective population-based cohort of 174 infants with ARM in the United Kingdom and Ireland, 39 infants (22%) were diagnosed late, while female sex, a visible perineal opening, and visible meconium on the perineum were associated with an increased likelihood of late diagnosis (Long et al., 2024). These findings emphasize the importance of careful anatomical assessment

and continued clinical observation when meconium is passed through an abnormal perineal opening (Amerstorfer et al., 2022; Long et al., 2024).

Gastrointestinal perforation is an uncommon but potentially life-threatening complication of ARM, particularly when diagnosis or effective intestinal decompression is delayed (Kruger et al., 2019; Singh et al., 2023). Delayed recognition of ARM may be accompanied by progressive bowel distension and obstructive manifestations, with intestinal perforation documented among affected neonates (Kruger et al., 2019). Although colonic perforation is rarely encountered in ARM, neonatal cases have been documented, indicating that this complication should be considered when abdominal deterioration occurs in an affected infant (King et al., 2016; Singh et al., 2023). More specifically, neonatal sigmoid colon perforation has been reported in association with low ARM, indicating that severe proximal colonic complications may occur even in anatomically lower forms of the malformation (Parelkar et al., 2016; Singh et al., 2023). The pathophysiology of perforation is likely multifactorial, with inadequate distal decompression, progressive bowel distension, and increased intraluminal pressure proposed as contributing mechanisms (Parelkar et al., 2016; Singh et al., 2023). Therefore, the mechanism responsible for perforation in an individual neonate should be interpreted by correlating the clinical course with radiological and intraoperative findings rather than being attributed to distal obstruction alone (Parelkar et al., 2016; Singh et al., 2023).

Early recognition of an evolving intra-abdominal complication requires integration of serial clinical assessment and appropriate imaging rather than reliance solely on the apparent patency of a distal fistula (Riccabona et al., 2017; Long et al., 2024). Progressive abdominal distension, vomiting or feeding intolerance, and clinical deterioration in a neonate with ARM warrant reassessment for possible intestinal obstruction or perforation (Kruger et al., 2019; Singh et al., 2023). Radiological evaluation complements physical examination by helping define anorectal anatomy, identify associated abnormalities, and provide information relevant to subsequent surgical decision-making (Riccabona et al., 2017). Despite previous reports of sigmoid colon perforation associated with low ARM, published evidence specifically describing sigmoid colon perforation complicating neonatal rectovestibular fistula remains limited (Parelkar et al., 2016; Singh et al., 2023). This represents a clinically relevant knowledge gap because distal meconium passage through a rectovestibular fistula may coexist with significant proximal bowel pathology and therefore should not be considered sufficient evidence of effective colonic decompression (Long et al., 2024; Singh et al., 2023). The present case describes sigmoid colon perforation complicating neonatal rectovestibular fistula, with particular emphasis on the temporal relationship among clinical deterioration, radiological findings, and intraoperative anatomy. The clinical value of this report lies in highlighting that continued meconium passage through a rectovestibular fistula does not necessarily exclude a serious proximal intestinal complication and that integrated clinical, radiological, and surgical assessment may facilitate timely recognition and appropriate management of this rare presentation (Long et al., 2024; Singh et al., 2023).

RESEARCH METHOD

This study was designed as a retrospective descriptive case report based on a review of the medical records. Clinical information was retrospectively collected from the patient's medical records, including perinatal history, clinical presentation, physical examination findings, laboratory investigations, radiological findings, operative findings, postoperative management, and clinical outcomes. The clinical course was reconstructed chronologically from the initial presentation through diagnostic evaluation, surgical intervention, and postoperative follow-up,

with particular attention to the relationship between clinical deterioration, radiological evidence of gastrointestinal perforation, and intraoperative findings. (Chu et al., 2021)

RESULTS AND DISCUSSIONS

A 3-day-old female neonate, born at term and weighing 3,000 g, was referred to our emergency department because of an absent anal opening that had been recognized since birth. She developed progressive abdominal distension accompanied by recurrent bilious vomiting. Her parents initially reported that she had not passed meconium after delivery. However, further questioning revealed intermittent passage of small amounts of meconium through a tiny opening in the vestibule, located just below the vaginal introitus.

The infant was delivered vaginally at term following an uncomplicated pregnancy. The delivery was uneventful, with no reported birth trauma. The mother had no significant illness during pregnancy, and there was no known family history of congenital anomalies, anorectal malformations, or hereditary gastrointestinal disorders.

At presentation, the infant was hemodynamically stable. Her abdomen was markedly distended but remained soft, with no overlying erythema or edema. Careful examination of the perineum confirmed the absence of a normal anal opening. A small opening was identified within the vestibule, from which only a minimal amount of meconium could be expressed. The location of this opening and the passage of meconium through it were consistent with an anorectal malformation with a rectovestibular fistula.

Laboratory evaluation showed marked leukocytosis ($25.98 \times 10^3/\mu\text{L}$) with neutrophil predominance and an elevated C-reactive protein level of 6.4 mg/L, suggesting an ongoing inflammatory response. Prothrombin time was mildly prolonged, whereas serum electrolyte levels were within normal limits.

Plain abdomen foto showed marked gaseous dilatation of multiple bowel loops, consistent with distal intestinal obstruction. Free intraperitoneal air was also evident, indicating pneumoperitoneum and raising strong suspicion of gastrointestinal perforation (Figure 1). No significant abnormality was observed on the chest radiograph.

In view of the intestinal obstruction and pneumoperitoneum, the infant was initially stabilized with gastric decompression and supportive therapy. Emergency exploratory laparotomy was subsequently performed.

A transverse supraumbilical incision was made, and extensive fecal contamination was encountered upon entering the peritoneal cavity. Exploration revealed a perforation of the sigmoid colon with diffuse fecal peritonitis. The perforation was primarily repaired with a two-layer closure. A double-barrel sigmoid colostomy was then created to divert the fecal stream and protect the repaired bowel segment.

After surgery, the infant was transferred to the neonatal intensive care unit for close observation and postoperative care. She was managed with bowel rest, nasogastric decompression, broad-spectrum intravenous antibiotics, intravenous fluids, nutritional support, and regular assessment of colostomy function.

Her postoperative recovery was uneventful. Abdominal distension gradually subsided, and subsequent abdominal radiography demonstrated resolution of the bowel dilatation with no residual or recurrent pneumoperitoneum (Figure 2). Enteral feeding was gradually introduced on postoperative day 4 after bowel function had recovered and was well tolerated.

1
1
1
The infant continued to improve and was discharged on postoperative day 8 in good general condition. At the time of discharge, her abdomen was soft and non-distended, the surgical wound was healing without evidence of infection, and the double-barrel sigmoid colostomy was viable and functioning well (Figure 3).

1
1
At the 3-month follow-up, the patient remained clinically well. The laparotomy incision had completely healed, and the abdomen remained non-distended. The colostomy was viable and functioning normally, with healthy-appearing mucosa and no clinically apparent stoma-related complications (Figure 4).



Figure 1. plain abdominal radiograph obtained at presentation showing marked gaseous dilatation of multiple bowel loops, consistent with distal intestinal obstruction. Free intraperitoneal air (pneumoperitoneum) is also present, raising suspicion of gastrointestinal perforation.



1
1
Figure 2. Postoperative supine abdominal radiograph showing resolution of the previously observed bowel dilatation and pneumoperitoneum. An intra-abdominal drainage catheter is visible, with its tip projecting over the left mid-abdomen.



Figure 3. Clinical appearance on postoperative day 8. Abdominal distension had resolved, and the transverse laparotomy incision was healing well without clinical signs of surgical-site infection. The double-barrel sigmoid colostomy was viable and functioning appropriately.



Figure 4. Clinical appearance at the 3-month follow-up showing complete healing of the laparotomy incision and a non-distended abdomen. The double-barrel sigmoid colostomy remained viable, with healthy-appearing mucosa and no clinically apparent stoma-related complications.

RESULTS AND DISCUSSIONS

he findings of the present case support evidence that visible meconium passage through a perineal fistula does not necessarily exclude clinically significant anorectal obstruction. In this

Commented [MOU1]: The article sections consist of Introduction, Research Method, Results and Discussions, and Conclusion. Please adjust accordingly

neonate, only small amounts of meconium intermittently passed through the rectovestibular fistula despite progressive abdominal distension, bilious vomiting, and marked bowel dilatation, indicating inadequate distal decompression. Visible meconium on the perineum has been associated with delayed diagnosis, and obstructive manifestations are reported more frequently among patients whose ARM is recognized late (Davidson et al., 2024; Long et al., 2024). Rectovestibular fistula is also a common ARM phenotype in female patients and may present with an abnormal opening in the vestibule through which stool or meconium passes (Hageman et al., 2024). In contrast to the complicated presentation in our patient, selected patients with uncomplicated rectovestibular fistula can undergo definitive repair without preliminary fecal diversion with satisfactory outcomes (Amanollahi & Ketabchian, 2016; Aziz et al., 2017). These contrasting findings suggest that the clinical significance of a rectovestibular fistula depends not only on its anatomical presence but also on its functional adequacy for fecal drainage, the severity of obstruction, the patient's clinical condition, and timing of diagnosis (Xiao et al., 2026). Therefore, the distinctive feature of the present case was the coexistence of limited meconium passage through a rectovestibular fistula with progressive evidence of functionally significant distal obstruction.

The clinical course suggests a plausible pathophysiological progression from inadequate distal drainage to progressive bowel distension and subsequent perforation. A small or functionally inadequate fistulous opening may allow some meconium to pass while remaining insufficient for effective decompression, and persistent abdominal distension or defecation difficulty despite conservative management may necessitate fecal diversion in selected patients (Xiao et al., 2026). In our patient, marked gaseous dilatation of multiple bowel loops on admission radiography provided objective evidence of substantial distal intestinal obstruction. Persistent obstruction may increase intraluminal pressure and bowel-wall stress, providing a plausible mechanism by which an obstructed colon becomes susceptible to perforation; nevertheless, this mechanism cannot be directly proven from the present case. Pneumoperitoneum on admission indicated gastrointestinal perforation, which was subsequently confirmed at laparotomy as sigmoid colon perforation with diffuse fecal peritonitis. Although colonic perforation is rare in ARM, sigmoid perforation has been reported in neonates with low ARM and represents a potentially life-threatening complication requiring prompt surgical treatment (Singh et al., 2023). Delayed ARM diagnosis has also been associated with serious complications, including intestinal perforation, reinforcing the importance of early recognition of distal obstruction (Kruger et al., 2019). Accordingly, the sigmoid perforation in our patient should be interpreted as a severe complication occurring in the setting of unresolved distal obstruction rather than as a proven direct consequence of the rectovestibular fistula itself.

Delayed recognition provides an important clinical context for understanding the progression observed in this case. A substantial proportion of neonates with ARM may still experience delayed diagnosis, particularly when an abnormal perineal opening or visible meconium creates the impression that distal bowel drainage is adequate (Davidson et al., 2024; Long et al., 2024). Delayed presentation has also been reported in settings where early neonatal recognition and access to specialized pediatric surgical care are limited (Bhavsar et al., 2021). Importantly, the available evidence demonstrates an association rather than a direct causal relationship between delayed recognition and perforation; therefore, diagnostic delay should be regarded as a potential contributor that prolongs unresolved obstruction in susceptible neonates rather than as an independent cause of intestinal perforation (Kruger et al., 2019; Davidson et al., 2024). In the present case, intermittent passage of a small amount of meconium

8 may have contributed to initial false reassurance despite the absence of a normal anal opening. This observation emphasizes that the presence of meconium on the perineum should not be interpreted as evidence of normal anorectal anatomy or adequate intestinal decompression (Davidson et al., 2024; Long et al., 2024).

1 The surgical strategy in this patient was determined primarily by the acute complication rather than by the rectovestibular fistula alone. In uncomplicated rectovestibular fistula, both primary and staged reconstruction have been described, and treatment should be individualized according to anatomy, clinical condition, associated abnormalities, and surgical expertise (Amanollahi & Ketabchian, 2016). In our patient, sigmoid perforation with diffuse fecal peritonitis shifted the immediate therapeutic priority from definitive anorectal reconstruction to source control, repair of the perforation, and fecal diversion. Primary two-layer closure of the sigmoid perforation followed by a double-barrel sigmoid colostomy provided diversion of the fecal stream while allowing definitive anorectal reconstruction to be deferred until clinical recovery. Staged repair with colostomy remains an accepted strategy for selected patients with rectovestibular fistula, particularly when clinical circumstances make immediate definitive reconstruction inappropriate (Xiao et al., 2026). Large registry data further demonstrate that temporary stomas remain part of ARM management, although stoma-related complications require appropriate surveillance (Hageman et al., 2024). The resolution of abdominal distension and pneumoperitoneum, successful initiation of enteral feeding, discharge on postoperative day eight, and a viable functioning colostomy at three months demonstrated a favorable short-term outcome. Nevertheless, longer follow-up remains essential because constipation, soiling, fecal accidents, and the need for bowel management may persist after reconstruction of rectovestibular fistula (van der Steeg et al., 2022; Hakalmaz & Tekant, 2023).

The major clinical and educational implication of this case is that meconium passage should never substitute for direct assessment of anorectal anatomy in a newborn. In female neonates, careful inspection is essential because rectovestibular fistula and other ARM phenotypes may present with abnormal perineal openings through which meconium is visible (Hageman et al., 2024). Visible meconium may paradoxically contribute to delayed diagnosis when it is incorrectly assumed to originate from a normally positioned anus (Davidson et al., 2024; Long et al., 2024). Therefore, physicians, midwives, nurses, and other healthcare professionals responsible for newborn assessment should systematically inspect the perineum, confirm the presence and normal position of the anus, examine the urethral and genital openings, and identify any additional perineal opening. When meconium is observed, its exact anatomical source should be verified rather than assumed. An absent anal opening, meconium emerging through an abnormal opening, progressive abdominal distension, bilious vomiting, or inadequate stool evacuation should raise suspicion of ARM with intestinal obstruction and prompt early pediatric surgical referral. The importance of careful postnatal examination is further reinforced by the continuing difficulty of reliably detecting ARM prenatally (Huijgen et al., 2024). Improving neonatal examination and training of healthcare professionals performing newborn assessment may therefore facilitate earlier recognition and reduce preventable diagnostic delay (Kruger et al., 2019; Long et al., 2024).

This case is strengthened by the concordance between clinical presentation, direct perineal findings, laboratory evidence of inflammation, radiographic demonstration of bowel obstruction and pneumoperitoneum, and operative confirmation of sigmoid perforation with

1

diffuse fecal peritonitis. Nevertheless, as a single case report, it cannot establish the incidence of perforation or prove a causal relationship between delayed recognition, inadequate fistulous drainage, and sigmoid perforation. The proposed pathophysiological mechanism remains inferential because fistula caliber, intraluminal pressure, local bowel perfusion, and histopathological changes at the perforation site were not objectively assessed. Furthermore, the three-month follow-up is insufficient to determine outcomes after definitive anorectal reconstruction, particularly constipation, continence, urinary function, and long-term quality of life, which remain important components of ARM follow-up (van der Steeg et al., 2022; Hakalmaz & Tekant, 2023). Despite these limitations, the present case demonstrates an important clinical principle: limited passage of meconium does not confirm normal anorectal anatomy and should not be considered sufficient evidence of adequate distal bowel decompression. Systematic perineal examination, confirmation of the actual route of meconium passage, recognition of obstructive warning signs, and timely referral may reduce diagnostic delay and provide an opportunity for intervention before severe complications develop (Kruger et al., 2019; Davidson et al., 2024). Therefore, systematic perineal examination of every newborn should remain an essential component of routine neonatal assessment, particularly at primary and maternity care facilities where the first opportunity to identify ARM commonly occurs.

7

CONCLUSION

This case emphasizes the importance of systematic perineal examination in every newborn for the early recognition of anorectal malformations. Healthcare professionals should verify normal anorectal anatomy and the actual route of meconium passage rather than relying solely on a history of stool passage. This report reinforces the need to educate healthcare practitioners, particularly those in primary and maternity care facilities, to perform careful perineal examination as part of routine newborn assessment, enabling timely diagnosis and referral and potentially preventing progression to severe complications.

1


ACKNOWLEDGEMENTS

The authors would like to express their sincere gratitude to RS Hermina, Kabupaten Bogor, Jawa Barat, and to the Departments of Pediatric Surgery, Neonatology, and Medical Records for their assistance in patient management and data collection. Special appreciation is extended to all healthcare professionals involved in the multidisciplinary care of this patient, and to the patient's family for their consent to the publication of clinical information and images.

References

1. Amerstorfer, E. E., Schmiedeke, E., Samuk, I., Sloots, C. E. J., van Rooij, I. A. L. M., Jenetzky, E., Midrio, P., & ARM-Net Consortium. (2022). Clinical differentiation between a normal anus, anterior anus, congenital anal stenosis, and perineal fistula: Definitions and consequences—The ARM-Net Consortium consensus. *Children*, 9(6), 831. <https://doi.org/10.3390/children9060831>. (Amerstorfer et al., 2022)
2. Amanollahi, O., & Ketabchian, S. (2016). One-stage vs. three-stage repair in anorectal malformation with rectovestibular fistula. *African Journal of Paediatric Surgery*, 13, 20–25. <https://doi.org/10.4103/0189-6725.181702>
3. Aziz, D. A., Velayutham, R., Osman, M., Latiff, Z., Lim, F., & Mohd Nor, M. (2017). Anorectal anomaly with rectovestibular fistula: A historical comparison of neonatal anterior sagittal anorectoplasty without covering colostomy and postoperative anal dilatation to the classical three-stage posterior sagittal anorectoplasty. *Open Access Surgery*, 10, 33–44. DOI: 10.2147/OAS.S137904
4. Bhavsar, R., Ray, S., Verma, M., Agarwal, S., & Nundy, S. (2021). Single stage repair of anorectal malformation with rectovestibular fistula in adult. *Annals of Medicine and Surgery*, 72. DOI: 10.1016/j.amsu.2021.103057
5. Chanchlani, R., & Budhwani, K. S. (2023). A study of the clinical profile and management of children with anorectal malformations. *Cureus*, 15. DOI: 10.7759/cureus.36772
6. Chu, H., Kim, Y., & Min, B. (2021). A guide to writing a case report according to CARE guideline. *Journal of Korean Medical Society of Soft Tissue*, 5(1), 41–45. DOI 10.54461/jkmst.2021.5.1.41
7. Davidson, J. R., Zaparackaite, I., Holbrook, C., & Thakkar, H. (2024). Late diagnosis of anorectal malformation: How good is good enough? *Pediatric Surgery International*, 40. DOI: 10.1007/s00383-024-05765-2
8. Hageman, I. C., Midrio, P., van der Steeg, H. J. J., et al. (2024). The European Anorectal Malformation Network (ARM-Net) patient registry: 10-year review of clinical and surgical characteristics. *British Journal of Surgery*, 111. <https://doi.org/10.1093/bjs/znae019>. (Hageman et al., 2024)
9. Hakalmaz, A. E., & Tekant, G. (2023). Anorectal malformations and late-term problems. *Turkish Archives of Pediatrics*, 58, 572–579. DOI: 10.5152/turkarchpediatr.2023.23090
10. Huijgen, D., Versteegh, H., Wijnen, R., Galjaard, S., Peters, N., & Sloots, C. (2024). Prenatal ultrasound in the diagnosis of anorectal malformations: Correlating prenatal signs with postnatal outcomes. *Prenatal Diagnosis*, 45, 35–43. DOI: 10.1002/pd.6723
11. Kancherla, V., et al. (2023). Prevalence and mortality among children with anorectal malformation: A multi-country analysis. *Birth Defects Research*. <https://doi.org/10.1002/bdr2.2129>. (Kancherla et al., 2023)
12. King, S. K., Cooksey, R., Atkinson, J., McLeod, E., & Levitt, M. A. (2016). Colonic perforation in a neonate with an anorectal malformation. *ANZ Journal of Surgery*, 86. DOI 10.1111/ans.12704

13. Kruger, P., Teague, W. J., Khanal, R., Hutson, J. M., & King, S. K. (2019). Delayed diagnosis of anorectal malformations in neonates. *ANZ Journal of Surgery*, 89. <https://doi.org/10.1111/ans.15374>. (Kruger et al., 2019)
14. Lawal, T. A. (2019). Overview of anorectal malformations in Africa. *Frontiers in Surgery*, 6, 7. DOI: 10.3389/fsurg.2019.00007
15. Long, A. M., Davidson, J. R., Tyraskis, A., Knight, M., & De Coppi, P. (2024). A population-based cohort study on diagnosis and early management of anorectal malformation in the UK and Ireland. *Journal of Pediatric Surgery*. <https://doi.org/10.1016/j.jpedsurg.2024.03.009>. (Long et al., 2024)
16. Parelkar, S., Kapadnis, S., Sanghvi, B., Joshi, P., Mundada, D., & Oak, S. (2016). Neonatal sigmoid colon perforation: A rare occurrence in low anorectal malformation and review of the literature. *Pediatrics & Neonatology*, 57(3), 232–235. <https://doi.org/10.1016/j.pedneo.2013.07.011>. (Parelkar et al., 2016)
17. Riccabona, M., Lobo, M. L., Ording-Müller, L. S., et al. (2017). European Society of Paediatric Radiology abdominal imaging task force recommendations in paediatric uro-radiology, part IX: Imaging in anorectal and cloacal malformation, imaging in childhood ovarian torsion, and efforts in standardising paediatric uro-radiology terminology. *Pediatric Radiology*, 47, 1369–1380. <https://doi.org/10.1007/s00247-017-3837-6>. (Riccabona et al., 2017)
18. Singh, R., Suman, B., Sinha, A., & Kumar, B. (2023). Neonatal sigmoid colon perforation in low anorectal malformation. *Journal of Indian Association of Pediatric Surgeons*, 28, 342–344. https://doi.org/10.4103/jiaps.jiaps_36_23. (Singh et al., 2023)
19. Sohail, H., Nawaz, T., Mirza, M., Ara, J., Nawaz, M., Arif, A., & Ahmed, M. (2022). Late presentation of anorectal malformations—A dilemma in modern world: Cross-sectional analysis from a tertiary care hospital, Pakistan. *Pakistan Armed Forces Medical Journal*. DOI: 10.51253/pafmj.v72isuppl-2.4426.
20. Sirichamratsakul, K., Laochareonsuk, W., Surachat, K., & Sangkhathat, S. (2023). Population-based prevalence study of common congenital malformations of the alimentary tract and abdominal wall in Thailand: A study using data from the National Health Security Office. *World Journal of Pediatric Surgery*, 6, e000540. <https://doi.org/10.1136/wjps-2022-000540>. (Sirichamratsakul et al., 2023)
21. van der Steeg, H. V. D., van Rooij, I. A. L. M., Iacobelli, B. D., et al. (2022). Bowel function and associated risk factors at preschool and early childhood age in children with anorectal malformation type rectovestibular fistula: An ARM-Net Consortium study. *Journal of Pediatric Surgery*. DOI: 10.1016/j.jpedsurg.2022.02.015
22. Xiao, X., Feng, W., Wang, Z., Chen, P., Hou, J., & Wang, Y. (2026). Colostomy in patients with anorectal malformation with rectoperineal fistula or rectovestibular fistula: A single-center experience. *BMC Gastroenterology*, 26. DOI: 10.1186/s12876-026-04734-4



Sigmoid colon perforation complicating neonatal rectovestibular fistula: A case report

Partogi Napitupulu¹, Nadifa Agil², Revalita Wahab³, Nathalia Ningrum⁴

^{1,3}Department of Radiology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

²Department of Surgery, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

⁴Department of Pediatrics, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

ARTICLE INFO

Article history:

Received Jul 21, 2026

Revised Jul 30, 2026

Accepted Aug 7, 2026

Keywords:

Anorectal Malformation
Bowel Obstruction
Pneumoperitonuem
Rectovestibular Fistula
Sigmoid Colon Perforation

ABSTRACT

Anorectal malformation (ARM) with rectovestibular fistula may allow small amounts of meconium to pass despite ongoing distal bowel obstruction. We report a three-day-old term female neonate with an imperforate anus, progressive abdominal distension, bilious vomiting, and minimal meconium passage through a rectovestibular fistula. Abdominal radiography showed marked bowel dilatation and pneumoperitoneum. Emergency laparotomy revealed sigmoid colon perforation with diffuse fecal peritonitis. The perforation was repaired, followed by a double-barrel sigmoid colostomy. The infant recovered well and was discharged on postoperative day eight. At three-month follow-up, the wound was completely healed and the colostomy remained healthy and functional. This case illustrates a rare but serious complication of rectovestibular fistula, in which limited meconium passage may coexist with significant bowel obstruction and subsequent perforation. These findings emphasize that thorough perineal inspection, high vigilance for obstruction despite minor meconium passage, and timely surgical intervention are essential to prevent bowel perforation in neonates with anorectal malformations.

This is an open access article under the [CC BY-NC](#) license.



Corresponding Author:

Partogi Napitupulu,
Faculty of Medicine,
Universitas Trisakti,
Jl. Kyai Tapa No. 1, Grogol, Jakarta Barat, 11440, Jakarta, Indonesia
Email: partoginapitupulu@trisakti.ac.id

INTRODUCTION

Anorectal malformations (ARMs) encompass a heterogeneous group of congenital abnormalities involving the distal rectum and anus, with variable fistulous communications to the urinary or genital tract and frequent associations with other congenital anomalies (Amerstorfer et al., 2022; Hageman et al., 2024). Because the anatomical presentation of ARM varies considerably, accurate identification of the malformation during the neonatal period is important for determining appropriate diagnostic evaluation and subsequent surgical management (Amerstorfer et al., 2022). Careful inspection of the anus and perineum should therefore be an integral component of routine neonatal examination, particularly for identifying abnormalities in anal position, caliber, or fistulous openings (Amerstorfer et al., 2022). Although uncommon, ARM occurs worldwide and

shows geographical variation in reported birth prevalence (Kancherla et al., 2023). A large multicountry surveillance study involving 9,438 cases among nearly 29 million births estimated a pooled ARM prevalence of 3.26 per 10,000 total births (Kancherla et al., 2023). Population-based data from Southeast Asia remain comparatively limited; however, a nationwide analysis using Thailand's National Health Security Office database reported an ARM prevalence of 2.57 per 10,000 births between 2017 and 2020 (Sirichamratsakul et al., 2023).

Among female patients with ARM, rectovestibular fistula represents an important anatomical phenotype in which the rectum opens into the vestibule anterior to the expected anal position (Hageman et al., 2024). Data from the European ARM-Net registry, which included 2,619 patients from 34 centers, showed that vestibular fistula accounted for 31.8% of ARM phenotypes among female patients (Hageman et al., 2024). Although a rectovestibular fistula permits passage of meconium and may provide some degree of distal bowel decompression, visible meconium through an abnormal perineal opening should not be interpreted as evidence of normal anorectal anatomy or necessarily adequate intestinal emptying (Long et al., 2024). This distinction is clinically relevant because delayed recognition of ARM continues to occur despite routine neonatal assessment (Kruger et al., 2019; Long et al., 2024). In a prospective population-based cohort of 174 infants with ARM in the United Kingdom and Ireland, 39 infants (22%) were diagnosed late, while female sex, a visible perineal opening, and visible meconium on the perineum were associated with an increased likelihood of late diagnosis (Long et al., 2024). These findings emphasize the importance of careful anatomical assessment and continued clinical observation when meconium is passed through an abnormal perineal opening (Amerstorfer et al., 2022; Long et al., 2024).

Recognizing signs of bowel obstruction in a neonate with a rectovestibular fistula carries substantial clinical urgency, because diagnostic delay is consistently linked to a higher risk of intestinal perforation, sepsis, and mortality, particularly when a patent fistula creates a false impression of adequate decompression (Oyania et al., 2024; Piperidis et al., 2025). International reviews emphasize that ARM is a clinical diagnosis that should be actively excluded within the first 24-72 hours of life through a structured perineal examination performed by trained birth attendants, including midwives, rather than inferred from spontaneous meconium passage alone (Murthi et al., 2023; Oyania et al., 2024). Every hour of unrecognized obstruction allows further bowel distension and wall stress to accumulate, so the interval between birth and a confirmatory perineal examination represents a critical window for preventing progression to perforation and peritonitis (Piperidis et al., 2025). However, substantial knowledge gaps persist regarding the clinical trajectory, warning signs, and monitoring protocols for sigmoid colon perforation in neonates with rectovestibular fistulas, particularly in detecting early intestinal wall stress before perforation occurs.

Gastrointestinal perforation is an uncommon but potentially life-threatening complication of ARM, particularly when diagnosis or effective intestinal decompression is delayed (Kruger et al., 2019; Singh et al., 2023). Delayed recognition of ARM may be accompanied by progressive bowel distension and obstructive manifestations, with intestinal perforation documented among affected neonates (Kruger et al., 2019). Although colonic perforation is rarely encountered in ARM, neonatal cases have been documented, indicating that this complication should be considered when abdominal deterioration occurs in an affected infant (King et al., 2016; Singh et al., 2023). More specifically, neonatal sigmoid colon perforation has been reported in association with low ARM, indicating that severe proximal colonic complications may occur even in anatomically lower forms of the malformation (Parelkar et al., 2016; Singh et al., 2023). The pathophysiology of perforation is likely multifactorial, with inadequate distal decompression, progressive bowel distension, and increased intraluminal pressure proposed as contributing mechanisms (Parelkar et al., 2016; Singh et al., 2023). Therefore, the mechanism responsible for perforation in an individual neonate should be interpreted by correlating the clinical course with radiological and

intraoperative findings rather than being attributed to distal obstruction alone (Parelkar et al., 2016; Singh et al., 2023).

Early recognition of an evolving intra-abdominal complication requires integration of serial clinical assessment and appropriate imaging rather than reliance solely on the apparent patency of a distal fistula (Riccabona et al., 2017; Long et al., 2024). Progressive abdominal distension, vomiting or feeding intolerance, and clinical deterioration in a neonate with ARM warrant reassessment for possible intestinal obstruction or perforation (Kruger et al., 2019; Singh et al., 2023). Radiological evaluation complements physical examination by helping define anorectal anatomy, identify associated abnormalities, and provide information relevant to subsequent surgical decision-making (Riccabona et al., 2017). Despite previous reports of sigmoid perforation associated with low ARM, published evidence specifically describing sigmoid colon perforation complicating neonatal rectovestibular fistula remains limited (Parelkar et al., 2016; Singh et al., 2023). This represents a clinically relevant knowledge gap because distal meconium passage through a rectovestibular fistula may coexist with significant proximal bowel pathology and therefore should not be considered sufficient evidence of effective colonic decompression (Long et al., 2024; Singh et al., 2023). The present case describes sigmoid colon perforation complicating neonatal rectovestibular fistula, with particular emphasis on the temporal relationship among clinical deterioration, radiological findings, and intraoperative anatomy. The clinical value of this report lies in highlighting that continued meconium passage through a rectovestibular fistula does not necessarily exclude a serious proximal intestinal complication and that integrated clinical, radiological, and surgical assessment may facilitate timely recognition and appropriate management of this rare presentation (Long et al., 2024; Singh et al., 2023).

RESEARCH METHOD

This study was designed as a retrospective descriptive case report based on a review of the medical records (Chu et al., 2021). Clinical information was retrospectively collected from the patient's medical records, including admission records, laboratory results, radiological reports, operative notes, and postoperative records. including perinatal history, clinical presentation, physical examination findings, laboratory investigations, radiological findings, operative findings, postoperative management, and clinical outcomes. Clinical, radiological, and intraoperative findings were correlated according to their chronological sequence and concordance, with emphasis on the consistency between progressive clinical signs of bowel obstruction, radiological evidence of bowel dilatation and pneumoperitoneum, and operative confirmation of sigmoid colon perforation) The case analysis primarily focused on the clinical progression of bowel obstruction and its correlation with radiological and intraoperative findings.

RESULTS AND DISCUSSIONS

A 3-day-old female neonate, born at term and weighing 3,000 g, was referred to our emergency department because of an absent anal opening that had been recognized since birth. She developed progressive abdominal distension accompanied by recurrent bilious vomiting. Her parents initially reported that she had not passed meconium after delivery. However, further questioning revealed intermittent passage of small amounts of meconium through a tiny opening in the vestibule, located just below the vaginal introitus. The infant was delivered vaginally at term following an uncomplicated pregnancy. The delivery was uneventful, with no reported birth trauma. The mother had no significant illness during pregnancy, and there was no known family history of congenital anomalies, anorectal malformations, or hereditary gastrointestinal disorders.

At presentation, the infant was hemodynamically stable. Her abdomen was markedly distended but remained soft, with no overlying erythema or edema. Careful examination of the perineum confirmed the absence of a normal anal opening. A small opening was identified within

the vestibule, from which only a minimal amount of meconium could be expressed. The location of this opening and the passage of meconium through it were consistent with an anorectal malformation with a rectovestibular fistula.

Laboratory evaluation showed marked leukocytosis ($25.98 \times 10^3/\mu\text{L}$) with neutrophil predominance and an elevated C-reactive protein level of 6.4 mg/L, suggesting an ongoing inflammatory response. Prothrombin time was mildly prolonged, whereas serum electrolyte levels were within normal limits. Plain abdomen foto showed marked gaseous dilatation of multiple bowel loops, consistent with distal intestinal obstruction. Free intraperitoneal air was also evident, indicating pneumoperitoneum and raising strong suspicion of gastrointestinal perforation (Figure 1). In neonates with anorectal malformations, finding pneumoperitoneum indicates severe bowel perforation, signaling an critical need for immediate emergency surgical intervention. In view of the intestinal obstruction and pneumoperitoneum, the infant was initially stabilized with gastric decompression and supportive therapy. Emergency exploratory laparotomy was subsequently performed.

A transverse supraumbilical incision was made, and extensive fecal contamination was encountered upon entering the peritoneal cavity. Exploration revealed a perforation of the sigmoid colon with diffuse fecal peritonitis. The perforation was primarily repaired with a two-layer closure. A double-barrel sigmoid colostomy was then created to divert the fecal stream and protect the repaired bowel segment. After surgery, the infant was transferred to the neonatal intensive care unit for close observation and postoperative care. She was managed with bowel rest, nasogastric decompression, broad-spectrum intravenous antibiotics, intravenous fluids, nutritional support, and regular assessment of colostomy function.

Her postoperative recovery was uneventful. Abdominal distension gradually subsided, and subsequent abdominal radiography demonstrated resolution of the bowel dilatation with no residual or recurrent pneumoperitoneum (Figure 2). Enteral feeding was gradually introduced on postoperative day 4 after bowel function had recovered and was well tolerated.

The infant continued to improve and was discharged on postoperative day 8 in good general condition. At the time of discharge, her abdomen was soft and non-distended, the surgical wound was healing without evidence of infection, and the double-barrel sigmoid colostomy was viable and functioning well (Figure 3). At the 3-month follow-up, the patient remained clinically well. The laparotomy incision had completely healed, and the abdomen remained non-distended. The colostomy was viable and functioning normally, with healthy-appearing mucosa and no clinically apparent stoma-related complications (Figure 4).



Figure 1. Plain abdominal foto obtained at presentation, showing marked gaseous dilatation of multiple bowel loops, consistent with distal intestinal obstruction. Free intraperitoneal air (pneumoperitoneum) is also present, raising suspicion of gastrointestinal perforation



Figure 2. Postoperative supine abdominal radiograph showing resolution of the previously observed bowel dilatation and pneumoperitoneum. An intra-abdominal drainage catheter is visible, with its tip projecting over the left mid-abdomen



Figure 3. Clinical appearance on postoperative day 8. Abdominal distension had resolved, and the transverse laparotomy incision was healing well without clinical signs of surgical-site infection. The double-barrel sigmoid colostomy was viable and functioning appropriately



Figure 4. Clinical appearance at the 3-month follow-up showing complete healing of the laparotomy incision and a non-distended abdomen. The double-barrel sigmoid colostomy remained viable, with healthy-appearing mucosa and no clinically apparent stoma-related complications

The findings of the present case support evidence that visible meconium passage through a perineal fistula does not necessarily exclude clinically significant anorectal obstruction. In this neonate, only small amounts of meconium intermittently passed through the rectovestibular fistula despite progressive abdominal distension, bilious vomiting, and marked bowel dilatation, indicating inadequate distal decompression. Visible meconium on the perineum has been associated with delayed diagnosis, and obstructive manifestations are reported more frequently among patients whose ARM is recognized late (Davidson et al., 2024; Long et al., 2024). Rectovestibular fistula is also a common ARM phenotype in female patients and may present with an abnormal opening in the vestibule through which stool or meconium passes (Hageman et al., 2024). In contrast to the complicated presentation in our patient, selected patients with uncomplicated rectovestibular fistula can undergo definitive repair without preliminary fecal diversion with satisfactory outcomes (Abdul Aziz et al., 2017; Amanollahi & Ketabchian, 2016). These contrasting findings suggest that the clinical significance of a rectovestibular fistula depends not only on its anatomical presence but also on its functional adequacy for fecal drainage, the severity of obstruction, the patient's clinical condition, and timing of diagnosis (Xiao et al., 2026). Therefore, the distinctive feature of the present case was the coexistence of limited meconium passage through a rectovestibular fistula with progressive evidence of functionally significant distal obstruction.

The clinical course suggests a plausible pathophysiological progression from inadequate distal drainage to progressive bowel distension and subsequent perforation. A small or functionally inadequate fistulous opening may allow some meconium to pass while remaining insufficient for effective decompression, and persistent abdominal distension or defecation difficulty despite conservative management may necessitate fecal diversion in selected patients (Xiao et al., 2026). In our patient, marked gaseous dilatation of multiple bowel loops on admission radiography provided objective evidence of substantial distal intestinal obstruction. Persistent obstruction may increase intraluminal pressure and bowel-wall stress, providing a plausible mechanism by which an obstructed colon becomes susceptible to perforation; nevertheless, this mechanism cannot be directly proven from the present case. Pneumoperitoneum on admission indicated gastrointestinal perforation, which was subsequently confirmed at laparotomy as sigmoid colon perforation with diffuse fecal peritonitis. Although colonic perforation is rare in ARM, sigmoid perforation has been reported in neonates with low ARM and represents a potentially life-threatening complication requiring prompt surgical treatment (Singh et al., 2023). Delayed ARM diagnosis has also been associated with serious complications, including intestinal perforation, reinforcing the importance of early recognition of distal obstruction (Kruger et al., 2019). Accordingly, the sigmoid perforation in our patient should be interpreted as a severe complication occurring in the setting of unresolved distal obstruction rather than as a proven direct consequence of the rectovestibular fistula itself.

Delayed recognition provides an important clinical context for understanding the progression observed in this case. A substantial proportion of neonates with ARM may still experience delayed diagnosis, particularly when an abnormal perineal opening or visible meconium creates the impression that distal bowel drainage is adequate (Davidson et al., 2024; Long et al., 2024). Delayed presentation has also been reported in settings where early neonatal recognition and access to specialized pediatric surgical care are limited (Bhavsar et al., 2021). Importantly, the available evidence demonstrates an association rather than a direct causal relationship between delayed recognition and perforation; therefore, diagnostic delay should be regarded as a potential contributor that prolongs unresolved obstruction in susceptible neonates rather than as an independent cause of intestinal perforation (Kruger et al., 2019; Davidson et al., 2024). In the present case, intermittent passage of a small amount of meconium may have contributed to initial false reassurance despite the absence of a normal anal opening. This observation emphasizes that the presence of meconium on the perineum should not be interpreted

as evidence of normal anorectal anatomy or adequate intestinal decompression (Davidson et al., 2024; Long et al., 2024).

Several concrete clinical lessons can be drawn from this case to help prevent delayed diagnosis in similar presentations. First, any meconium observed at an abnormal perineal site should prompt a complete anorectal examination rather than being accepted as reassurance of normal bowel patency, since delayed diagnosis is disproportionately common precisely among neonates who pass some meconium through an ectopic opening (Piperidis et al., 2025). Second, progressive abdominal distension, bilious vomiting, or feeding intolerance occurring after birth in an infant with a known or suspected perineal fistula should trigger urgent radiological evaluation for obstruction or perforation rather than a period of watchful waiting (Oyania et al., 2024). Third, structured, checklist-based newborn examination performed by trained birth attendants, together with explicit documentation of the anatomical source of any meconium passage, has been associated with earlier detection of ARM in national screening programs and should be reinforced through ongoing training of midwives and other first-line examiners (Murthi et al., 2023). Applying these lessons systematically may not eliminate delayed diagnosis, but it can shorten the interval between birth and definitive recognition of obstruction, potentially preventing progression to perforation as occurred in the present case (Murthi et al., 2023; Piperidis et al., 2025).

The surgical strategy in this patient was determined primarily by the acute complication rather than by the rectovestibular fistula alone. In uncomplicated rectovestibular fistula, both primary and staged reconstruction have been described, and treatment should be individualized according to anatomy, clinical condition, associated abnormalities, and surgical expertise (Amanollahi & Ketabchian, 2016). In our patient, sigmoid perforation with diffuse fecal peritonitis shifted the immediate therapeutic priority from definitive anorectal reconstruction to source control, repair of the perforation, and fecal diversion. Primary two-layer closure of the sigmoid perforation followed by a double-barrel sigmoid colostomy provided diversion of the fecal stream while allowing definitive anorectal reconstruction to be deferred until clinical recovery. Staged repair with colostomy remains an accepted strategy for selected patients with rectovestibular fistula, particularly when clinical circumstances make immediate definitive reconstruction inappropriate (Xiao et al., 2026). Large registry data further demonstrate that temporary stomas remain part of ARM management, although stoma-related complications require appropriate surveillance (Hageman et al., 2024). The resolution of abdominal distension and pneumoperitoneum, successful initiation of enteral feeding, discharge on postoperative day eight, and a viable functioning colostomy at three months demonstrated a favorable short-term outcome. Nevertheless, longer follow-up remains essential because constipation, soiling, fecal accidents, and the need for bowel management may persist after reconstruction of rectovestibular fistula (Hakalmaz & Topuzlu Tekant, 2023; van der Steeg et al., 2022).

The major clinical and educational implication of this case is that meconium passage should never substitute for direct assessment of anorectal anatomy in a newborn. In female neonates, careful inspection is essential because rectovestibular fistula and other ARM phenotypes may present with abnormal perineal openings through which meconium is visible (Hageman et al., 2024). Visible meconium may paradoxically contribute to delayed diagnosis when it is incorrectly assumed to originate from a normally positioned anus (Davidson et al., 2024; Long et al., 2024). Therefore, physicians, midwives, nurses, and other healthcare professionals responsible for newborn assessment should systematically inspect the perineum, confirm the presence and normal position of the anus, examine the urethral and genital openings, and identify any additional perineal opening. When meconium is observed, its exact anatomical source should be verified rather than assumed. An absent anal opening, meconium emerging through an abnormal opening, progressive abdominal distension, bilious vomiting, or inadequate stool evacuation should raise suspicion of ARM with intestinal obstruction and prompt early pediatric surgical referral. The importance of careful postnatal examination is further reinforced by the continuing difficulty of

reliably detecting ARM prenatally (Huijgen et al., 2025) neonatal examination and training of healthcare professionals performing newborn assessment may therefore facilitate earlier recognition and reduce preventable diagnostic delay (Kruger et al., 2019; Long et al., 2024).

This case is strengthened by the concordance between clinical presentation, direct perineal findings, laboratory evidence of inflammation, radiographic demonstration of bowel obstruction and pneumoperitoneum, and operative confirmation of sigmoid perforation with diffuse fecal peritonitis. Nevertheless, as a single case report, it cannot establish the incidence of perforation or prove a causal relationship between delayed recognition, inadequate fistulous drainage, and sigmoid perforation. The proposed pathophysiological mechanism remains inferential because fistula caliber, intraluminal pressure, local bowel perfusion, and histopathological changes at the perforation site were not objectively assessed. Furthermore, the three-month follow-up is insufficient to determine outcomes after definitive anorectal reconstruction, particularly constipation, continence, urinary function, and long-term quality of life, which remain important components of ARM follow-up (van der Steeg et al., 2022; Hakalmaz & Tekant, 2023). Despite these limitations, the present case demonstrates an important clinical principle: limited passage of meconium does not confirm normal anorectal anatomy and should not be considered sufficient evidence of adequate distal bowel decompression. Systematic perineal examination, confirmation of the actual route of meconium passage, recognition of obstructive warning signs, and timely referral may reduce diagnostic delay and provide an opportunity for intervention before severe complications develop (Kruger et al., 2019; Davidson et al., 2024). Therefore, systematic perineal examination of every newborn should remain an essential component of routine neonatal assessment, particularly at primary and maternity care facilities where the first opportunity to identify ARM commonly occurs.

CONCLUSION

This case demonstrates that fistula patency must not be equated with adequate bowel decompression, as visible meconium can coexist with life-threatening colonic perforation. To improve early detection in primary care settings, four recommendations are proposed: (1) routine, structured perineal inspection before discharge; (2) explicit documentation of the meconium source; (3) low threshold for pediatric surgical referral upon detecting abnormal perineal openings or obstructive symptoms; and (4) continuous training for birth attendants on screening pathways. Future research should focus on prospective multicenter studies to identify specific risk factors for colonic perforation in neonates with anorectal malformations and evaluate standardized management protocols to strengthen current evidence.

ACKNOWLEDGEMENTS

The authors would like to express their sincere gratitude to RS Hermina, Kabupaten Bogor, Jawa Barat, and to the Departments of Pediatric Surgery, Neonatology, and Medical Records for their assistance in patient management and data collection. Special appreciation is extended to all healthcare professionals involved in the multidisciplinary care of this patient, and to the patient's family for their consent to the publication of clinical information and images.

References

- Abdul Aziz, D. A., Velayutham, R., Osman, M., Abdul Latiff, Z., Lim, F., & Mohd Nor, M. (2017). Anorectal anomaly with rectovestibular fistula: a historical comparison of neonatal anterior sagittal anorectoplasty without covering colostomy and postoperative anal dilatation to the classical three-stage posterior sagittal anorectoplasty. *Open Access Surgery, Volume 10*, 33–44. <https://doi.org/10.2147/OAS.S137904>
- Amanollahi, O., & Ketabchian, S. (2016). One-stage vs. three-stage repair in anorectal malformation with rectovestibular fistula. *African Journal of Paediatric Surgery : AJPS*, 13(1), 20–25.

- <https://doi.org/10.4103/0189-6725.181702>
- Amerstorfer, E. E., Schmiedeke, E., Samuk, I., Sloots, C. E. J., van Rooij, I. A. L. M., Jenetzky, E., Midrio, P., Aminoff, D., Bagolan, P., Iacobelli, B., de Blaauw, I., van der Steeg, H., Broens, P., Çavuşoğlu, H., Deluggi, S., Ludwiczek, J., Divarci, E., Fanjul, M., Fascetti-Leon, F., ... Witvliet, M. (2022). Clinical Differentiation between a Normal Anus, Anterior Anus, Congenital Anal Stenosis, and Perineal Fistula: Definitions and Consequences-The ARM-Net Consortium Consensus. *Children (Basel, Switzerland)*, 9(6). <https://doi.org/10.3390/CHILDREN9060831>
- Bhavsar, R., Ray, S., Verma, M., Agarwal, S. K., & Nundy, S. (2021). Single stage repair of anorectal malformation with rectovestibular fistula in adult. *Annals of Medicine and Surgery*, 72, 103057. <https://doi.org/10.1016/J.AMSU.2021.103057>
- Davidson, J., Zaparackaite, I., Holbrook, C., & Thakkar, H. (2024). Late diagnosis of anorectal malformation: how good is good enough? *Pediatric Surgery International*, 40(1), 175. <https://doi.org/10.1007/S00383-024-05765-2>
- Hageman, I. C., Midrio, P., van der Steeg, H. J. J., Jenetzky, E., Iacobelli, B. D., Morandi, A., Sloots, C. E. J., Schmiedeke, E., Broens, P. M. A., Leon, F. F., Çavuşoğlu, Y. H., Gorter, R. R., Trajanovska, M., King, S. K., Aminoff, D., Schwarzer, N., Haanen, M., de Blaauw, I., & van Rooij, I. A. L. M. (2024). The European Anorectal Malformation Network (ARM-Net) patient registry: 10-year review of clinical and surgical characteristics. *The British Journal of Surgery*, 111(2). <https://doi.org/10.1093/BJS/ZNAE019>
- Hakalmaz, A. E., & Topuzlu Tekant, G. (2023). Anorectal Malformations and Late-Term Problems. *Turkish Archives of Pediatrics*, 58(6), 572-579. <https://doi.org/10.5152/TURKARCHPEDIATR.2023.23090>
- Huijgen, D., Versteegh, H. P., Wijnen, R. M. H., Galjaard, S., Peters, N. C. J., & Sloots, C. E. J. (2025). Prenatal Ultrasound in the Diagnosis of Anorectal Malformations: Correlating Prenatal Signs With Postnatal Outcomes. *Prenatal Diagnosis*, 45(1), 35-43. <https://doi.org/10.1002/PD.6723;PAGE=STRING:ARTICLE/CHAPTER>
- Kancherla, V., Sundar, M., Tandaki, L., Lux, A., Bakker, M. K., Bergman, J. E. H., Bermejo-Sánchez, E., Canfield, M. A., Dastgiri, S., Feldkamp, M. L., Gatt, M., Groisman, B., Hurtado-Villa, P., Kallen, K., Landau, D., Lelong, N., Lopez-Camelo, J., Martinez, L. E., Mastroiacovo, P., ... Rissmann, A. (2023). Prevalence and mortality among children with anorectal malformation: A multi-country analysis. *Birth Defects Research*, 115(3), 390-404. <https://doi.org/10.1002/BDR2.2129>
- King, S. K., Cooksey, R., Atkinson, J., Mcleod, E., & Levitt, M. (2016). Colonic perforation in a neonate with an anorectal malformation. *ANZ Journal of Surgery*, 86(5), 418-419. <https://doi.org/10.1111/ANS.12704>
- Kruger, P., Teague, W. J., Khanal, R., Hutson, J. M., & King, S. K. (2019). Delayed diagnosis of anorectal malformations in neonates. *ANZ Journal of Surgery*, 89(10), 1253-1255. <https://doi.org/10.1111/ANS.15374>
- Long, A. M., Davidson, J. R., Tyraskis, A., Knight, M., & De Coppi, P. (2024). A Population-Based Cohort Study on Diagnosis and Early Management of Anorectal Malformation in the UK and Ireland. *Journal of Pediatric Surgery*, 59(8), 1463-1469. <https://doi.org/10.1016/J.JPESURG.2024.03.009>
- Parelkar, S. V., Kapadnis, S. P., Sanghvi, B. V., Joshi, P. B., Mundada, D. D., & Oak, S. N. (2016). Neonatal Sigmoid Colon Perforation: A Rare Occurrence in Low Anorectal Malformation and Review of the Literature. *Pediatrics and Neonatology*, 57(3), 232-235. <https://doi.org/10.1016/j.pedneo.2013.07.011>
- Riccabona, M., Lobo, M. L., Ording-Muller, L. S., Thomas Augdal, A., Fred Avni, E., Blickman, J., Bruno, C., Damasio, B., Darge, K., Ntoulia, A., Papadopoulou, F., & Vivier, P. H. (2017). European Society of Paediatric Radiology abdominal imaging task force recommendations in paediatric urology, part IX: Imaging in anorectal and cloacal malformation, imaging in childhood ovarian torsion, and efforts in standardising paediatric urology.... *Pediatric Radiology*, 47(10), 1369-1380. <https://doi.org/10.1007/S00247-017-3837-6>
- Singh, R., Suman, B., Sinha, A., & Kumar, B. (2023). Neonatal Sigmoid Colon Perforation in Low Anorectal Malformation. *Journal of Indian Association of Pediatric Surgeons*, 28(4), 342-344. https://doi.org/10.4103/JIAPS.JIAPS_36_23
- Sirichamratsakul, K., Laochareonsuk, W., Surachat, K., & Sangkhathat, S. (2023). Population-based prevalence study of common congenital malformations of the alimentary tract and abdominal wall in Thailand: a study using data from the National Health Security Office. *World Journal of Pediatric Surgery*, 6(3), e000540. <https://doi.org/10.1136/WJPS-2022-000540>
- van der Steeg, H. J. J., van Rooij, I. A. L. M., Iacobelli, B. D., Sloots, C. E. J., Morandi, A., Broens, P. M. A., Makedonsky, I., Leon, F. F., Schmiedeke, E., Vázquez, A. G., Miserez, M., Lisi, G., Midrio, P., Amerstorfer, E. E., Fanjul, M., Ludwiczek, J., Stenström, P., van der Steeg, A. F. W., & de Blaauw, I.

- (2022). Bowel function and associated risk factors at preschool and early childhood age in children with anorectal malformation type rectovestibular fistula: An ARM-Net consortium study. *Journal of Pediatric Surgery*, 57(9), 89–96. <https://doi.org/10.1016/J.JPESURG.2022.02.015>
- Xiao, X., Feng, W., Wang, Z., Chen, P., Hou, J., & Wang, Y. (2026). Colostomy in patients with anorectal malformation with rectoperineal fistula or rectovestibular fistula: a single-center experience. *BMC Gastroenterology*, 26(1). <https://doi.org/10.1186/S12876-026-04734-4>