

<https://jkk-fk.ejournal.unsri.ac.id/index.php/jkk>



HomeCurrentArchivesAnnouncementsAboutdianasamara

About the Journal

Jurnal Kedokteran dan Kesehatan : Publikasi Ilmiah Fakultas Kedokteran Universitas Sriwijaya

Published by Fakultas Kedokteran Universitas Sriwijaya

Online ISSN 2614-0411 Print ISSN 2406-7431

Home / About the Journal

Jurnal Kedokteran dan Kesehatan: Publikasi Ilmiah Fakultas Kedokteran Universitas Sriwijaya (JKK) with registered number ISSN 2614-0411 (Online) and ISSN 2406-7431 (Print) and , is a scientific journal published by Faculty of Medicine Universitas Sriwijaya, Indonesia.

JKK is a peer-reviewed, multidisciplinary, and scientific journal that publishes original research articles relevant to national health issues, including the areas of Biomedical Sciences, Clinical Medicine, and Public Health Science. JKK is an open-access journal published three times yearly (January, April, and October) to promote research in health-related fields.

JKK has been nationally accredited Sinta 4 by the Ministry of Research Technology and Higher Education of the Republic of Indonesia based on SK No. 10/C/C3/DT.05.00/2025, March 21 , 2025)



ISSN 2614-0411 (Online)
ISSN 2406-7431 (Print)

Jurnal Kedokteran dan Kesehatan: Publikasi Ilmiah Fakultas Kedokteran Universitas Sriwijaya is expected to participate and contribute towards better known as a result of scientific studies and research findings that academic circles and researchers can access. JKK has been collaborating with ARIKESI (Asosiasi Riset Ilmu Kesehatan Indonesia) since 2024, intending to help increase the visibility of the results of scientific studies and research findings that academic circles and researchers can access. JKK has been **published in the English language since April 2024** to broaden its reach to all countries globally and to collaborate with editors and reviewers from other countries.

Editorial Team
Reviewer
Contact

Author Guidelines
Publication Ethic
Copyright Notice

Article Processing Charges

Indexed by



[Home](#) [Current](#) [Archives](#) [Announcements](#) [About](#) [dianasamara](#)

Jurnal Kedokteran dan Kesehatan : Publikasi Ilmiah Fakultas Kedokteran Universitas Sriwijaya

Published by Fakultas Kedokteran Universitas Sriwijaya

Online ISSN 2614-0411 Print ISSN 2406-7431

[Current Issue](#) [About the Journal](#)



Vol. 13 No. 2 (2026): Jurnal Kedokteran dan Kesehatan : Publikasi Ilmiah Fakultas Kedokteran Universitas Sriwijaya

2026-04-25

Has been published Jurnal Kedokteran dan Kesehatan : Publikasi Ilmiah Fakultas Kedokteran Universitas Sriwijaya Vol. 13 No. 2 2026. There are 15 articles whose authors come from agencies/universities. Congratulations to all authors and thanks to the team of editors, layouter, proofreader, IT Support and reviewers involved in this issue.



ISSN 2614-0411 (Online)

ISSN 2406-7431 (Print)

Editorial History

Jurnal Kedokteran dan Kesehatan : Publikasi Ilmiah Fakultas Kedokteran Universitas Sriwijaya

Published by Fakultas Kedokteran Universitas Sriwijaya

Online ISSN 2614-0411 Print ISSN 2406-7431

[Home](#) / Editorial History


This section lists past contributors.

Editor in Chief

- [Indri Seta Septadina](#), Scopus ID: 57210232380 Universitas Sriwijaya, Indonesia

Managing Editor

- [Msy. Rulan Adnindya](#), Scopus ID: 57195990858, Universitas Sriwijaya, Indonesia



International Editorial Board Advisory Regional Asia

- [Sharifa Ezat Wan Puteh](#), Scopus ID: 16834115500, Faculty of Medicine. University Kebangsaan Malaysia (UKM)
- [Dito Anurogo](#), Scopus ID: 57224408026, Fakultas Kedokteran dan Ilmu Kesehatan (FKIK) Universitas Muhammadiyah (Unismuh) Makassar and International PhD Program for Cell Therapy and Regeneration Medicine (IPCTRM), Taipei Medical University (TMU), Taiwan, Indonesia
- [Ahmad Ghiffari](#), Scopus ID: 57200047086, Universitas Muhammadiyah Palembang, Goethe-Universität Frankfurt am Main · Institute of Occupational Medicine, Social Medicine and Environmental Medicine, Indonesia
- [Krisna Murti](#), Scopus ID: 55232295600, Universitas Sriwijaya, Indonesia
- [Minarma Siagian](#), Scopus ID : 6505956355, Universitas Indonesia Jakarta, Indonesia
- [Rina agustina](#), Scopus ID : 57214141404, Universitas Indonesia, Indonesia
- [Muhammad Irsan Saleh](#), Scopus ID: 57192919902, Universitas Sriwijaya, Palembang, Indonesia
- [Achadiyani Achadiyani](#), Scopus ID : 57195942091, Universitas Padjadjaran Bandung, Indonesia
- [Andi Muhammad Ihsan](#), Scopus ID : 57215357485, Universitas Hasanudin Makassar, Indonesia

Articles

THE ROLE OF VITAMIN D3 SUPPLEMENTATION ON MYASTHENIA GRAVIS COMPOSITE SCALE: A CASE REPORT

— Meidistya Ayu Mardhani, Wina Sinaga, Diyah Eka Andayani



Page: 137-143

NEUROLYTIC SPLANCHNIC NERVE BLOCK WITH ALCOHOL FOR UPPER ABDOMINAL CANCER PATIENTS: A SYSTEMATIC REVIEW

— Aidyl Fitriyah, Nico Adithia Prawira, Devadiza Friankasari



Page: 144-153

A POSSIBLE MODALITY OF INTERVENTION IN A SURGICALLY IMPOSSIBLE CASE OF GIANT CORONARY ANEURYSM: A SUCCESSFUL CASE

— Sidhi Laksono, Axel Jusuf, Hillary Kusharsamita



Page: 154-165

RESTING ECOLOGY AND SPATIAL HETEROGENEITY OF FILARIASIS MOSQUITO VECTORS IN TROPICAL WETLANDS: IMPLICATIONS FOR INTEGRATED VECTOR CONTROL IN BANYUASIN, INDONESIA

— Rini Pratiwi, Peni Kusumasari, Fitriyanti



Page: 166-182



ISSN 2614-0411 (Online)

ISSN 2406-7431 (Print)

Editorial Team

Reviewer

Contact

Author Guidelines

Publication Ethic

Copyright Notice



Article Processing Charges

VALIDITY, RELIABILITY, AND DIAGNOSTIC PERFORMANCE OF GERIATRIC-8 AS A SCREENING TOOL FOR ABNORMAL COMPREHENSIVE GERIATRIC ASSESSMENT

— Ridzqie Dibyantari, Nina Kemala Sari, Edy Rizal Wahyudi, Muhadi, Aulia Rizka, Arif Mansjoer, Noorwati Sutandyo



Page: 183-191

A CROSS-SECTIONAL STUDY OF NUTRITIONAL STATUS AND ATOPIC DERMATITIS IN ADOLESCENTS

— Ida Effendi, Puti Maharani Diwa



Page: 192-198

THE RELATIONSHIP BETWEEN LENGTH OF SENTENCE AND LEVELS OF ANXIETY AND DEPRESSION AMONG INMATES AT THE RUTAN KELAS IIB TRENGGALEK

— Yohana Christy, Syeikh Alfa Ridho



Page: 199-205

RELATIONSHIP BETWEEN BLOOD TRANSFUSION FREQUENCY AND BLOOD IRON CONCENTRATION IN PATIENT WITH THALASSEMIA MAJOR

— Vanessa Ainum Rizqita, Tjam Diana Samara



Page: 206-213



MOLECULAR DOCKING ANALYSIS OF THE BIOACTIVE COMPOUND EMODIN WITH VEGFR2 IN CERVICAL CANCER

— Innayah Rania Farzana, Rara Inggarsih, Ziske Maritska, Ayesah Augusta Rosdah, Soilia Fertilita



DOI

Page: 214-224

PATERNAL EMOTIONAL SUPPORT, INFANT TEMPERAMENT, AND BREASTFEEDING DURATION AT PUSKESMAS BOGOR TIMUR IN 2026

— Rina Afrina, Tuti Rusmiati



DOI

Page: 225-233

THE IMPACT OF OCCUPATIONAL NOISE EXPOSURE ON AUDITORY FUNCTION AND POSTURAL STABILITY: A CROSS-SECTIONAL STUDY IN AN INDUSTRIAL FACILITY IN SUMATERA

— Yulia Anggraini, Abila Ghanie, Ahmad Hifni



DOI

Page: 234-242

ASSOCIATION BETWEEN MCISAAC SCORE AND ANTIBIOTIC PRESCRIBING IN SUSPECTED BACTERIAL TONSILLITIS

— Yustiana Suwanto, Aryaputri Nabila Haurani



DOI

Page: 243-249

FORENSIC ANALYSIS OF COMPLETE HANGING WITH A TYPICAL KNOT POSITION RESULTING IN MECHANICAL ASPHYXIA IN A SUICIDE CASE (CASE REPORT) FORENSIC ANALYSIS OF COMPLETE HANGING

— Kristina Sihalohe, Widyastuti, Peni Kusumasari

DOI

Page: 258-265

ANTIBIOTIC PROFILE IN THE TREATMENT OF CHRONIC SUPPURATIVE OTITIS MEDIA (CSOM) AT IBNU SINA HOSPITAL IN 2023-2025

— Annisa Shezan Nazafarin, Andi Sitti Fahirah Aarsal, Ahmad Ardhani Pratama, Andi Tenri Sanna, Mila Habibasari



DOI

Page: 266-273

[View All Issues »](#)

**Vol. 13 No. 2 (2026): Jurnal Kedokteran dan Kesehatan : Publikasi Ilmiah
Fakultas Kedokteran Universitas Sriwijaya****RELATIONSHIP BETWEEN BLOOD TRANSFUSION FREQUENCY AND BLOOD IRON
CONCENTRATION IN PATIENT WITH THALASSEMIA MAJOR**

Home / Archives / Vol. 13 No. 2 (2026): Jurnal Kedokteran dan Kesehatan : Publikasi Ilmiah Fakultas Kedokteran Universitas Sriwijaya / Articles

Vanessa Ainum Rizqita

— Anatomi Department, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Tjam Diana Samara

— Anatomi Department, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

 <https://doi.org/10.32539/jkk.v13i2.734> **Keywords:** *frequency, blood transfusion, ferritin level, thalassemia major, children*Abstract

Routine blood transfusions performed on thalassemia major patients can cause excessive iron accumulation

Routine blood transfusions performed on thalassemia major patients can cause excessive iron accumulation in the body, which carries the risk of complications. Serum ferritin levels are used to assess body iron stores. This study aims to determine the correlation between transfusion frequency and iron accumulation in patients with thalassemia major. This study used a cross-sectional design with analytical observation. Data was collected by consecutive, non-random sampling from the medical records of 55 patients with thalassemia major aged 5-18 years who underwent blood transfusions at Hermina Jatinegara Hospital from January to December 2023. The independent variable was the frequency of blood transfusions per year, while the dependent variable was the average serum ferritin level, measured three times a year (every 4 months). The data obtained were analyzed using the Spearman correlation test. The results of this study found that the frequency of blood transfusions in thalassemia major patients over 1 year was 15 per year (based on the median). The average serum ferritin level was 3957.73 ng/mL (based on the median value). Spearman's correlation test showed a weak correlation between blood transfusion frequency and ferritin levels ($r = 0.292$; $p = 0.031$). The study concluded that increased blood transfusion frequency was associated with higher blood iron levels, but the correlation was weak.

Author Biography**Vanessa Ainum Rizqita, Anatomi Department, Faculty of Medicine,
Universitas Trisakti, Jakarta, Indonesia**

Medical Study Program, Medical Faculty, Universitas Trisakti

References

Download this PDF file

 pdf

Published:
2026-04-25

Issue

Download this PDF file pdf**Published:**

2026-04-25

IssueVol. 13 No. 2 (2026): Jurnal Kedokteran dan
Kesehatan : Publikasi Ilmiah Fakultas
Kedokteran Universitas Sriwijaya**Section**

Articles

LicenseThis work is licensed under a Creative
Commons Attribution-ShareAlike 4.0
International License.**How to Cite**Rizqita, V. A., & Samara, T. D. (2026).
RELATIONSHIP BETWEEN BLOOD
TRANSFUSION FREQUENCY AND BLOOD IRON

References

1. Wahidiyat P, Permono B. Hemoglobinopati dan thalassemia anak. In: Permono B, Sutaryo I, Ugrasena ID, Windastuti E, Abdulsalam M, editors. Buku ajar hematologi-onkologi anak. Jakarta: Ikatan Dokter Anak Indonesia; 2018:65-8.
2. Grentina, Wahidayat PA. Mengenal thalassemia [Internet]. Jakarta: Ikatan Dokter Anak Indonesia; 2016 [cited 2024 Apr 7]. Available from: <https://www.idai.or.id/artikel/seputar-kesehatan-anak/mengenal-thalassemia>
3. Rujito L, Siswandari W, Lestari DW, editors. Talasemia: genetik dasar dan pengelolaan terkini. Purwokerto: UNSOED Press; 2019:90.
4. Khikmah FA, Prijoprano HS. Prevalensi thalassemia beta minor dengan menggunakan indeks mentzer pada pasien anemia Suku Sabu di RSUD Sabu Raijua. Journal of Ners Community. 2022;13(2):301-6. doi:10.55129/jnerscommunity.v13i2.1917
5. World Health Organization. Regional desk review of haemoglobinopathies with an emphasis on thalassaemia and accessibility and availability of safe blood and blood products as per these patients' requirement in South-East Asia under universal health coverage [Internet]. 2023 [cited 2024 Apr 30]. Available from: <https://www.who.int/publications/i/item/9789290228516>
6. Fianza PI, Rahmawati A, Widiastha SH, Afifah S, Ghazali M, Indrajaya A. Iron overload in transfusion-dependent Indonesian thalassemic patients. Anemia. 2021;5581831. doi:10.1155/2021/5581831
7. Rochman F, Mulyantari NK, Sutirtayasa IW. Hubungan jumlah transfusi darah dan penggunaan kelasi besi dengan kadar ferritin pada pasien talasemia. Medika Udayana. 2019;8(9):1-6
8. Lal A, Wong T, Keel S, Pagano M, Chung J, Kamdar A. The transfusion management of beta thalassemia in the United States. Transfusion. 2021;61(10):3027-39. doi:10.1111/trf.16640
9. Taher AT, Cappellini MD. How I manage transfusional iron overload in patients with thalassemia. Blood. 2021;137(6):667-75

10. Bhalodiya VR, Valiya LG, Mehta NA, Padhariya BB. Correlation of serum ferritin level in transfusion-dependant thalassemia major patients: a study at a medical college affiliated hospital in Gujarat region. *Int J Contemp Pediatr*. 2023;10(3):330-3. doi:10.18203/2349-3291.ijcp20230429
 11. Pangestika AD, Astuti TD, Aryani T. Literature review: gambaran kadar hemoglobin dan ferritin serum pada anak thalassemia β mayor paska transfusi [Internet]. 2022 [cited 2024 Mar 15]. Available from: http://digilib.unisayogya.ac.id/6629/1/NASKAH%20PUBLIKASI_AYU%20DWI%20PANGESTIKA_1811304020%20%20Ayu%20Dwi%20Pangestika
 12. Yusuf AK, Magdaleni AR, Karunia D. Hubungan kadar feritin serum terhadap fraksi ejeksi ventrikel kiri pada pasien thalassemia β mayor anak di RSUD Kabupaten Kediri. *Jurnal Verdure*. 2022;4(1):123–8.
 13. University of Rochester Medical Center. Ferritin (Blood) [Internet]. URM Health Encyclopedia; 2023 [cited 2026 Feb 24]. Available from: https://www.urmc.rochester.edu/encyclopedia/content?contenttypeid=167&contentid=ferritin_blood
 14. Fianza PI, Rahmawati A, Wijaya I, Oehadian A, Prasetya D, Vidyaniati P. Gender disparities in psychological disturbances and quality of life among adolescent and adult patients with thalassemia: a review. *J Multidiscip Healthc*. 2024;17:1663-9. doi:10.2147/jmdh.s444592
 15. Sadiq IZ, Abubakar FS, Usman HS, Abdullahi AD, Ibrahim B, Kastayal BS. Thalassemia: pathophysiology, diagnosis, and advances in treatment. *Thalass Rep*. 2024;14(4):81-102. doi:10.3390/thalassrep14040010
 16. Almahmoud R, Hussein A, Khaja FA, Soliman AF, Dewedar H, Shareef ZA. Growth and endocrinopathies among children with β -thalassemia major treated at Dubai Thalassemia centre. *BMC Pediatr*. 2024;24:244. doi:10.1186/s12887-024-04670-w
 17. Nassim MS, Mahmoud MA, Shady HA, Mohammed EAER. Assessment of disease knowledge gaps among beta thalassemia major patients and their caregivers. *Egypt Pediatr Assoc Gaz*. 2022;70:49. doi:10.1186/s43054-022-00143-w
 18. Akbar TIS, Riyanti EY, Khairunnisa K, Wahyuni H. No association between frequency of blood transfusions and incompatibility of blood transfusions in thalassemia patients. *J Indones Med Assoc*. 2022;78(4):488-92. doi:10.15405/jaas.2022.78.4.488-492
-

19. Saporizki RS, Sumantri AF, Achmad S. Pengaruh kepatuhan minum obat kelasi besi dengan kadar ferritin pada pasien thalassemia di RSUD Al-Ihsan Bandung tahun 2021/2022. Bandung Conference Series: Medical Science. 2023;3(1):178-9. doi:10.29313/bcsms.v3i1.5824
20. Kurban LA, Almarri BK, Alshamsi MH, Abdelrahman SS, Alwahshi SG, Alhorani Q. Optimized serum ferritin prediction of iron overload in transfusion-dependent thalassemia: likelihood ratio and age-adjustment approach. Ann Saudi Med. 2023;43(2):90-6. doi:10.5144/0256-4947.2023.90
21. Sarkar K, Pramanik N, Goswami RP, Mandal PK, Chowdhury P, Jana K. Correlation of liver function test with different age and sex group and with ferritin level in thalassemia. Indian J Public Health Res Dev. 2020;11(6):452-7. doi:10.37506/ijphrd.v11i6.9819
22. Weiss M, Parisi Jun M, Sheth S. Clinical and economic burden of regularly transfused adult patients with β -thalassemia in the United States: a retrospective cohort study using payer claims. Am J Hematol. 2019;94(5):E129-32. doi:10.1002/ajh.25429
23. Shah F, Huey K, Deshpande S, Turner M, Chitnis M, Schiller E. Relationship between serum ferritin and outcomes in β -thalassemia: a systematic literature review. J Clin Med. 2022;11(15):4448. doi:10.3390/jcm11154448
24. Coates TD. Iron overload in transfusion-dependent patients. Hematology Am Soc Hematol Educ Program. 2019;2019(1):337-44. doi:10.1182/hematology.2019000036
25. Spasiano A, Meloni A, Costantini S, Quaia E, Cademartiti F, Cinque P. Setting for "normal" serum ferritin levels in patients with transfusion-dependent thalassemia: our current strategy. J Clin Med. 2021;10(24):5985. doi:10.3390/jcm10245985
26. Mohamed YA, Morgan DS, Hussein AM, Mohamed AA. Noninvasive assessment and risk factors of liver fibrosis in pediatric patients with beta thalassemia major using transient elastography. Beni Suef Univ J Basic Appl Sci. 2022;11:114. doi:10.1186/s43088-022-00314-z

RELATIONSHIP BETWEEN BLOOD TRANSFUSION FREQUENCY AND BLOOD IRON CONCENTRATION IN PATIENT WITH THALASSEMIA MAJOR

Vanessa Ainum Rizqita¹, Tjam Diana Samara^{2*}

¹ Undergraduate Medical Study Program, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

² Anatomi Department, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

ARTICLE INFO

Corresponding author :

Tjam Diana Samara
Faculty of Medicine,
Universitas Trisakti

Email:

dianasamara@trisakti.ac.id

Kata kunci:

Frekuensi
Transfusi darah
Kadar feritin
Talasemia mayor
Anak

Keywords:

Frequency
Blood transfusion
Ferritin level
Thalassemia major
Children

Original submission:

September 11, 2025

Accepted:

March 26, 2026

Published:

April 25, 2026

ABSTRAK

Transfusi darah rutin yang dilakukan oleh pasien talasemia mayor dapat menyebabkan akumulasi zat besi berlebih dalam tubuh, yang berisiko menimbulkan komplikasi. Kadar feritin dalam serum digunakan untuk menilai akumulasi zat besi dalam tubuh. Penelitian ini bertujuan untuk mengetahui korelasi antara frekuensi transfusi dengan akumulasi zat besi pada pasien talasemia mayor. Penelitian ini menggunakan desain potong lintang dengan observasi analitik. Data dikumpulkan secara consecutive non random sampling dari rekam medis sebanyak 55 pasien talasemia mayor berusia 5-18 tahun yang menjalani transfusi darah di RS Hermina Jatinegara dari Januari hingga Desember 2023. Variabel bebas adalah frekuensi transfusi darah dalam setahun, sedangkan variabel tergantung adalah kadar rata-rata feritin dalam serum yang diukur tiga kali selama setahun (per empat bulan sekali). Data yang didapat dianalisis menggunakan uji korelasi Spearman. Hasil pada penelitian ini ditemukan frekuensi transfusi darah pasien talasemia mayor dalam satu tahun sebanyak 15 kali per tahun (berdasarkan nilai median). Rata-rata kadar feritin dalam serum sebesar 3957,73 ng/mL. (berdasarkan nilai median). Uji korelasi Spearman menunjukkan adanya korelasi lemah antara frekuensi transfusi darah dan kadar feritin ($r = 0,292$; $p = 0,031$). Simpulan penelitian ini adalah kenaikan frekuensi transfusi darah diikuti dengan kenaikan kadar besi dalam darah namun dengan kekuatan korelasi yang lemah.

ABSTRACT

Relationship Between Blood Transfusion Frequency and Blood Iron Concentration in Patient with Thalassemia Major. Routine blood transfusions performed on thalassemia major patients can cause excessive iron accumulation in the body, which carries the risk of complications. Serum ferritin levels are used to assess body iron stores. This study aims to determine the correlation between transfusion frequency and iron accumulation in patients with thalassemia major. This study used a cross-sectional design with analytical observation. Data was collected by consecutive, non-random sampling from the medical records of 55 patients with thalassemia major aged 5-18 years who underwent blood transfusions at Hermina Jatinegara Hospital from January to December 2023. The independent variable was the frequency of blood transfusions per year, while the dependent variable was the average serum ferritin level, measured three times a year (every 4 months). The data obtained were analyzed using the Spearman correlation test. The results of this study found that the frequency of blood transfusions in thalassemia major patients over 1 year was 15 per year (based on the median). The average serum ferritin level was 3957.73 ng/mL (based on the median value). Spearman's correlation test showed a weak correlation between blood transfusion frequency and ferritin levels ($r = 0.292$; $p = 0.031$). The study concluded that increased blood transfusion frequency was associated with higher blood iron levels, but the correlation was weak.

INTRODUCTION

Thalassemia is a chronic hemoglobin (Hb) disorder inherited in an autosomal recessive pattern. It is characterized by a reduction in the synthesis of globin chains, which may involve either the α - or β -globin chains.¹ According to the type of globin chain affected, thalassemia is classified

into α -thalassemia and β -thalassemia.² Thalassemia is classified into three types based on its clinical manifestations. Thalassemia minor (trait) is often asymptomatic or only mildly anemic. Thalassemia intermedia is characterized by symptoms that appear later compared to thalassemia major, with patients experiencing mild to moderate anemia. Thalassemia major represents the most severe form, requiring lifelong blood transfusions, and is typically detected during infancy or early childhood.³

Indonesia is one of the countries located within the global thalassemia belt, characterized by a high frequency of thalassemia genes.⁴ The prevalence of thalassemia carriers in Indonesia is estimated at 3–8% of the total population, with a birth rate of 23 per 1,000 among a population of approximately 240 million. According to 2021 data, there were 10,531 thalassemia patients in Indonesia, with more than 2,500 infants born each year with the condition.⁵ Thalassemia major cases requiring regular blood transfusions occur predominantly among children and adolescents. The majority of patients with thalassemia major who receive blood transfusions fall within the 5–18-year age group, as transfusion requirements during this period tend to remain stable to support growth and daily activities, particularly during the phase of active development.⁶

The primary management of anemia in patients with thalassemia major is blood transfusion. The goal of transfusion therapy is to maintain hemoglobin (Hb) levels above 9–9.5 g/dL.⁷ Blood transfusions are typically performed at intervals of 3–4 weeks, with the volume tailored to the patient's body weight and hemoglobin concentration.⁸ Blood transfusion therapy, monitoring, and management of iron overload are of primary concern. Since the human body lacks an active mechanism for eliminating excess iron, prolonged transfusion may lead to iron accumulation.⁷ Recent evidence consistently demonstrates that cumulative transfusional exposure is a major determinant of iron overload in transfusion-dependent thalassemia. Each unit of transfused red blood cells delivers approximately 200–250 mg of iron, and in the absence of a physiologic mechanism for active iron excretion, iron progressively accumulates over time.⁹ Moreover, clinical cohort data have shown a statistically significant association between total transfusion burden and serum ferritin concentration in patients with transfusion-dependent β -thalassemia major, indicating that greater cumulative exposure to transfusion contributes to higher iron loading as measured by ferritin levels.¹⁰ However, a literature review on post-transfusion ferritin levels in pediatric β -thalassemia major patients reported that mean ferritin concentrations in several analyzed studies were within recommended clinical targets. This observation indicates that ferritin levels may be maintained within acceptable limits when appropriate management strategies are implemented.¹¹ Serum ferritin measurement is the most commonly used method to evaluate iron accumulation (hemosiderosis) in thalassemia patients undergoing regular transfusions.¹²

The novelty of this study lies in its analytical cross-sectional approach in evaluating transfusion frequency as a quantitative variable associated with serum ferritin levels in thalassemia major patients within a specific local population. By providing updated local data, this study aims to clarify inconsistent findings reported in previous studies and to strengthen the evidence base for iron overload monitoring in routinely transfused patients.

METHOD

This study used a cross-sectional design with an analytical observational approach. The study population comprised patients with thalassemia major undergoing blood transfusions at Hermina Jatinegara Hospital, East Jakarta. Although thalassemia major consists of alpha thalassemia major dan beta thalassemia major², in this study they were not differentiated.

Based on the results of the sample-size calculation using the correlation test, 55 samples were obtained from 160 patients between January 1 and December 31, 2023. Sampling used the consecutive non-random sampling method. Samples were selected based on predefined inclusion and exclusion criteria. Inclusion criteria were patients with thalassemia major aged 5–18 years who had received blood transfusions and possessed complete medical records, including age, sex, annual transfusion frequency, and serial ferritin levels recorded between January and December 2023. Exclusion criteria were patients with coexisting hematological disorders, such as leukemia, hemophilia, or sickle cell anemia. Respondents were excluded if these diseases or disorders were listed in their medical records. If they were not listed, the patient was assumed not to have any other blood disorders.

Serum ferritin levels in this study were the average of three blood draws per year, performed at four-month intervals. The blood transfusion frequency in this study was the number of transfusions per year. All data were taken from medical records.

Data were analyzed using the Statistical Package for Social Sciences (SPSS) version 30. The Kolmogorov-Smirnov test was used to assess normality, as the sample size was >50. Since the data were not normally distributed, the Spearman correlation test was employed, with statistical significance set at $p < 0.05$.

RESULT

This study used medical records of patients with thalassemia major from the blood transfusion unit at Hermina Jatinegara Hospital in 2023, involving a total of 55 subjects.

Table 1. Distribution of study subject characteristics

Characteristic	Mean $\pm$ SD	Median (Range)	Frequency (n = 55)	Percentage (%)
Age	10,56 $\pm$ 3,711			
Gender				
Male			32	58,2
Female			23	41,8
Serum ferritin (ng/mL)		3957,73 (383,87 – 28566,16)		
Transfusion frequency (times/year)		15 (9 – 23)		

Note: Data are presented as mean $\pm$ standard deviation for normally distributed variables and median (range) for non-normally distributed variables.


SD: standard deviation

Based on Table 1, of the 55 subjects in the thalassemia blood transfusion unit, 32 (58.2%) were male, and 23 (41.8%) were female. Since the age data were normally distributed, age is presented as mean $\pm$ SD. The subjects' ages ranged from 5 to 18 years, with a mean age of 10.56 years and a standard deviation of 3.711 years.

In this study, because the data were not normally distributed, the distributions of serum ferritin levels and blood transfusion frequency were presented as medians. Based on Table 2, serum ferritin data obtained from serial measurements throughout 2023 showed 3,957.73 ng/mL

(according to a median value), with a minimum of 383.87 ng/mL and a maximum of 28,566.16 ng/mL. The frequency of blood transfusions in patients with thalassemia major in 2023 was 15 times per year (median), ranging from 9 to 23. Normal serum ferritin level for children are 7-140 ng/mL.¹³

Relationship between blood transfusion frequency and iron levels in patients with thalassemia major



*: Spearman's Correlation Test

Figure 1. Correlation between blood transfusion frequency and ferritin levels

The one-sample Kolmogorov–Smirnov test yielded a p-value of <0.001, indicating that the residuals were not normally distributed. Therefore, bivariate analysis was performed using Spearman's correlation test. The results showed a weak correlation between the frequency of blood transfusions and serum ferritin levels ($r = 0.292$, $p = 0.031$).

DISCUSSION

Characteristics of the study subjects

In this study, 55 subjects were included; the majority were male (32, 58.2%). This result is in line with the findings of Bhalodiya et al., who reported that 59% of their study population were male.¹⁰ However, the studies by Fianza et al. and Rochman et al. reported different findings, with a higher proportion of female subjects, accounting for 53.1% and 58.3%, respectively.^{7,14} This variation supports the theory that thalassemia is a genetic disorder with an autosomal recessive inheritance pattern and is not related to the sex chromosomes.¹⁵

The age distribution in this study ranged from 5 to 18 years, with a mean of 10.56 and a standard deviation of 3.711. This finding is consistent with the study by Almahmoud et al., which demonstrated a predominance of children (2–9 years) and adolescents (10–18 years) among patients with thalassemia major.¹⁶ The study by Nassim et al. also supports this finding, reporting an age distribution of subjects ranging from 2 to 14 years, with a mean age of 8.7 years.¹⁷ A considerable number of thalassemia cases are detected during childhood, when children present with worsening clinical symptoms, particularly increasing pallor, and subsequently require routine blood transfusions to maintain growth and daily activities.¹⁸

Distribution of the results of the blood iron level and blood transfusion frequency

In this study, the median ferritin level among subjects was 3957.73 ng/mL, which is consistent with the findings of Saparizki et al., who reported that the majority of thalassemia patients had ferritin levels exceeding 2500 ng/mL. Ferritin levels above 1000 ng/mL generally indicate iron overload (hyperferritinemia).¹⁹ The study by Kurban et al. also reported a median ferritin level of 1548 ng/mL, with a range of 161–3968 ng/mL, exceeding the recommended threshold value of 1000 ng/mL.²⁰ These findings suggest a substantial iron overload as a consequence of repeated blood transfusions among patients with thalassemia.²¹

This study found that patients with thalassemia major had a median annual transfusion frequency of 15 sessions. Aligning with the findings of Weiss et al., who reported a mean frequency of ≥ 12 transfusions per year.²² On the other hand, Akbar et al. reported that most patients received 12 or fewer transfusions annually.¹⁸ The frequency of blood transfusions in thalassemia patients varies depending on hemoglobin (Hb) levels and clinical condition. Transfusions are indicated when Hb levels fall below 7 g/dL in two consecutive measurements taken two weeks apart, or when Hb levels exceed 7 g/dL but are accompanied by clinical manifestations such as growth retardation, suspected bone fractures due to extramedullary hematopoiesis, and Cooley's facies.¹⁸

Relationship between blood transfusion frequency and iron levels in patients with thalassemia major

This study demonstrated a weak correlation between blood transfusion frequency and ferritin levels in patients with thalassemia major ($p = 0.031$; $r = 0.292$). This finding is consistent with the study by Bhalodiya et al., which reported a p-value of 0.00, indicating a significant increase in ferritin levels with rising total annual blood transfusion frequency.¹⁰ Similarly, Rochman et al. demonstrated a significant relationship between the number of transfusions and increased ferritin levels ($p = 0.003$).⁷ Thalassemia patients receiving repeated transfusions for anemia management are prone to elevated ferritin levels due to iron accumulation, as each unit of blood contains approximately 200–250 mg of iron, and the body lacks an inherent mechanism to excrete the excess.²³ The progressive accumulation of iron over time results in higher ferritin concentrations, thereby positioning transfusion frequency as a significant factor influencing iron overload.²⁴

Although no studies have been identified that explicitly state the absence of a correlation between transfusion frequency and ferritin levels, Pangestika et al. reported that hemoglobin and serum ferritin levels in children with beta-thalassemia major remained within the normal range (100 ng/mL) after transfusion.¹¹ This finding is consistent with the study by Spasiano et al., which reported that serum ferritin levels below 500 ng/mL can be achieved safely in some transfusion-dependent thalassemia (TDT) patients, indicating that transfusion frequency does not necessarily correlate directly with ferritin elevation in certain cases.²⁵ Even though a positive correlation between transfusion frequency and ferritin levels is observed, this relationship is significantly influenced by other factors, notably adherence to iron chelation therapy, which can markedly lower ferritin levels.²⁴ In addition, ferritin levels are influenced by other factors such as inflammation, infection, liver function, and genetic variations, which can cause differences in ferritin concentrations even among patients with similar transfusion histories.²⁶

CONCLUSION

In general, serum ferritin levels were higher than normal in study subjects who received more than 12 blood transfusions per year. However, the findings of this study indicate that the frequency of blood transfusions has only a weak effect on the increase in blood iron levels.

Future studies are advised to involve participants across a wider age range to gain a more comprehensive understanding of serum ferritin levels among thalassemia patients. In addition, subsequent research should address other potential determinants of ferritin levels, including variations in chelation therapy, treatment duration and compliance, nutritional status, and coexisting medical conditions.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest in this study.

ACKNOWLEDGMENTS

The authors are deeply grateful to Hermina Jatinegara Hospital for the permission and support provided throughout the data collection phase of this study.

REFERENCES

1. Wahidiyat P, Permono B. Hemoglobinopati dan thalassemia anak. In: Permono B, Sutaryo I, Ugrasena ID, Windastuti E, Abdulsalam M, editors. Buku ajar hematologi-onkologi anak. Jakarta: Ikatan Dokter Anak Indonesia; 2018:65-8.
2. Grentina, Wahidayat PA. Mengenal thalassemia [Internet]. Jakarta: Ikatan Dokter Anak Indonesia; 2016 [cited 2024 Apr 7]. Available from: <https://www.idai.or.id/artikel/seputar-kesehatan-anak/mengenal-thalassemia>
3. Rujito L, Siswandari W, Lestari DW, editors. Talasemia: genetik dasar dan pengelolaan terkini. Purwokerto: UNSOED Press; 2019:90.
4. Khikmah FA, Prijopranto HS. Prevalensi thalassemia beta minor dengan menggunakan indeks mentzer pada pasien anemia Suku Sabu di RSUD Sabu Rajua. Journal of Ners Community. 2022;13(2):301-6. doi:10.55129/jnerscommunity.v13i2.1917
5. World Health Organization. Regional desk review of haemoglobinopathies with an emphasis on thalassaemia and accessibility and availability of safe blood and blood products as per these patients' requirement in South-East Asia under universal health coverage [Internet]. 2023 [cited 2024 Apr 30]. Available from: <https://www.who.int/publications/i/item/9789290228516>
6. Fianza PI, Rahmawati A, Widihashta SH, Afifah S, Ghazali M, Indrajaya A. Iron overload in transfusion-dependent Indonesian thalassemic patients. Anemia. 2021;5581831. doi:10.1155/2021/5581831
7. Rochman F, Mulyantari NK, Sutirtayasa IW. Hubungan jumlah transfusi darah dan penggunaan kelasi besi dengan kadar ferritin pada pasien talasemia. Medika Udayana. 2019;8(9):1-6
8. Lal A, Wong T, Keel S, Pagano M, Chung J, Kamdar A. The transfusion management of beta thalassemia in the United States. Transfusion. 2021;61(10):3027-39. doi:10.1111/trf.16640
9. Taher AT, Cappellini MD. How I manage transfusional iron overload in patients with thalassemia. Blood. 2021;137(6):667-75
10. Bhalodiya VR, Valiya LG, Mehta NA, Padhariya BB. Correlation of serum ferritin level in transfusion-dependant thalassemia major patients: a study at a medical college affiliated hospital in Gujarat region. Int J Contemp Pediatr. 2023;10(3):330-3. doi:10.18203/2349-

3291.ijcp20230429

11. Pangestika AD, Astuti TD, Aryani T. Literature review: gambaran kadar hemoglobin dan ferritin serum pada anak thalassemia β mayor paska transfusi [Internet]. 2022 [cited 2024 Mar 15]. Available from: http://digilib.unisayogya.ac.id/6629/1/NASKAH%20PUBLIKASI_AYU%20DWI%20PANGESTIKA_1811304020%20-%20Ayu%20Dwi%20Pangestika
12. Yusuf AK, Magdalen AR, Karunia D. Hubungan kadar ferritin serum terhadap fraksi ejeksi ventrikel kiri pada pasien thalassemia β mayor anak di RSUD Kabupaten Kediri. *Jurnal Verdure*. 2022;4(1):123–8.
13. University of Rochester Medical Center. Ferritin (Blood) [Internet]. URM Health Encyclopedia; 2023 [cited 2026 Feb 24]. Available from: https://www.urmc.rochester.edu/encyclopedia/content?contenttypeid=167&contentid=ferritin_blood
14. Fianza PI, Rahmawati A, Wijaya I, Oehadian A, Prasetya D, Vidyaniati P. Gender disparities in psychological disturbances and quality of life among adolescent and adult patients with thalassemia: a review. *J Multidiscip Healthc*. 2024;17:1663-9. doi:10.2147/jmdh.s444592
15. Sadiq IZ, Abubakar FS, Usman HS, Abdullahi AD, Ibrahim B, Kastayal BS. Thalassemia: pathophysiology, diagnosis, and advances in treatment. *Thalass Rep*. 2024;14(4):81-102. doi:10.3390/thalassrep14040010
16. Almahmoud R, Hussein A, Khaja FA, Soliman AF, Dewedar H, Shareef ZA. Growth and endocrinopathies among children with β -thalassemia major treated at Dubai Thalassemia centre. *BMC Pediatr*. 2024;24:244. doi:10.1186/s12887-024-04670-w
17. Nassim MS, Mahmoud MA, Shady HA, Mohammed EAER. Assessment of disease knowledge gaps among beta thalassemia major patients and their caregivers. *Egypt Pediatr Assoc Gaz*. 2022;70:49. doi:10.1186/s43054-022-00143-w
18. Akbar TIS, Riyanti EY, Khairunnisa K, Wahyuni H. No association between frequency of blood transfusions and incompatibility of blood transfusions in thalassemia patients. *J Indones Med Assoc*. 2023;73(4):183-90. doi:10.47830/jinma-vol.73.4-2023-1069
19. Saparizki RS, Sumantri AF, Achmad S. Pengaruh kepatuhan minum obat kelasi besi dengan kadar ferritin pada pasien thalassemia di RSUD Al-Ihsan Bandung tahun 2021/2022. *Bandung Conference Series: Medical Science*. 2023;3(1):178-9. doi:10.29313/bcsms.v3i1.5824
20. Kurban LA, Almarri BK, Alshamsi MH, Abdelrahman SS, Alwahshi SG, Alhorani Q. Optimized serum ferritin prediction of iron overload in transfusion-dependent thalassemia: likelihood ratio and age-adjustment approach. *Ann Saudi Med*. 2023;43(2):90-6. doi:10.5144/0256-4947.2023.90
21. Sarkar K, Pramanik N, Goswami RP, Mandal PK, Chowdhury P, Jana K. Correlation of liver function test with different age and sex group and with ferritin level in thalassemia. *Indian J Public Health Res Dev*. 2020;11(6):452-7. doi:10.37506/ijphrd.v11i6.9819
22. Weiss M, Parisi Jun M, Sheth S. Clinical and economic burden of regularly transfused adult patients with β -thalassemia in the United States: a retrospective cohort study using payer claims. *Am J Hematol*. 2019;94(5):E129-32. doi:10.1002/ajh.25429
23. Shah F, Huey K, Deshpande S, Turner M, Chitnis M, Schiller E. Relationship between serum ferritin and outcomes in β -thalassemia: a systematic literature review. *J Clin Med*. 2022;11(15):4448. doi:10.3390/jcm11154448
24. Coates TD. Iron overload in transfusion-dependent patients. *Hematology Am Soc Hematol Educ Program*. 2019;2019(1):337-44. doi:10.1182/hematology.2019000036
25. Spasiano A, Meloni A, Costantini S, Quaia E, Cademartiti F, Cinque P. Setting for "normal" serum ferritin levels in patients with transfusion-dependent thalassemia: our current strategy. *J*

Clin Med. 2021;10(24):5985. doi:10.3390/jcm10245985

26. Mohamed YA, Morgan DS, Hussein AM, Mohamed AA. Noninvasive assessment and risk factors of liver fibrosis in pediatric patients with beta thalassemia major using transient elastography. Beni Suef Univ J Basic Appl Sci. 2022;11:114. doi:10.1186/s43088-022-00314-z

Diana Samara

Thalasemia major

 Pengecekan artikel

Document Details

Submission ID

trn:oid:::3618:138446885

Submission Date

May 11, 2026, 9:17 AM GMT+7

Download Date

May 11, 2026, 12:22 PM GMT+7

File Name

Thalasemia major.pdf

File Size

493.6 KB

8 Pages

3,653 Words

20,919 Characters

13% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

Exclusions

- 71 Excluded Matches

Match Groups

-  **32 Not Cited or Quoted 8%**
Matches with neither in-text citation nor quotation marks
-  **17 Missing Quotations 4%**
Matches that are still very similar to source material
-  **0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 9%  Internet sources
- 10%  Publications
- 8%  Submitted works (Student Papers)

Integrity Flags

0 Integrity Flags for Review

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups

- 32 Not Cited or Quoted 8%**
Matches with neither in-text citation nor quotation marks
- 17 Missing Quotations 4%**
Matches that are still very similar to source material
- 0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
- 0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 9% Internet sources
- 10% Publications
- 8% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

- 1** Student papers
University of Massachusetts-Dartmouth on 2026-04-28 <1%
- 2** Internet
www.ncbi.nlm.nih.gov <1%
- 3** Internet
etd.repository.ugm.ac.id <1%
- 4** Student papers
Chulalongkorn University on 2015-03-19 <1%
- 5** Internet
eglj.springeropen.com <1%
- 6** Internet
file.scirp.org <1%
- 7** Student papers
Brown University on 2025-10-30 <1%
- 8** Student papers
Los Angeles Mission College on 2023-07-12 <1%
- 9** Student papers
Manipal Academy of Higher Education (MAHE) on 2025-05-05 <1%
- 10** Student papers
Universitas Diponegoro on 2023-05-30 <1%

11	Internet	scholar.unand.ac.id	<1%
12	Publication	Wulan Purnama, Masfufah Masfufah, Nurdiana Nurdiana. "The Relationship Betw...	<1%
13	Publication	FUTTERWEIT, WALTER. "Serum Alkaline Phosphatase Activity in Infectious Monon...	<1%
14	Publication	Víctor Juan Vera-Ponce, Joan A. Loayza-Castro, Jhosmer Ballena-Caicedo, Lupita A...	<1%
15	Publication	Adam Wandell, Alexander Papanastassiou, Izabela Tarasiewicz, Mark Miller. "Wha...	<1%
16	Publication	Alicja Kotula, Natalia Kucy, Adrianna Czachor, Paula Kula et al. "Gaucher disease -...	<1%
17	Internet	bmcpediatr.biomedcentral.com	<1%
18	Publication	Khaled Salama, Amina Abdelsalam, Hadeel Seif Eldin, Eman Youness et al. "Iron o...	<1%
19	Internet	jsmc.univsul.edu.iq	<1%
20	Internet	ir.uitm.edu.my	<1%
21	Internet	www.mdpi.com	<1%
22	Student papers	Fakultas Kedokteran Gigi Universitas Trisakti on 2020-11-28	<1%
23	Internet	eprints.uns.ac.id	<1%
24	Publication	Rahmi Amtha, Ferry Sandra, Rosalina Tjandrawinata, Indrayadi Gunardi, Anggrae...	<1%

25	Publication	Safyudin, Dita Tri Ramadianti, Subandrate, Liniyanti D. Oswari, Eka Handayani O...	<1%
26	Student papers	Sriwijaya University on 2024-04-24	<1%
27	Internet	dspace.alquds.edu	<1%
28	Internet	inabj.org	<1%
29	Internet	journals.lww.com	<1%
30	Internet	revista.uepb.edu.br	<1%
31	Internet	www.hematologyresjournal.com	<1%
32	Internet	repository.unsri.ac.id	<1%
33	Publication	Hurriyati Ratih, Tjahjono Benny, GafarAbdullah Ade, Sulastri, Lisnawati. "Advance...	<1%
34	Publication	Qin Ru, Yusheng Li, Wenqing Xie, Yilan Ding, Lin Chen, Guodong Xu, Yuxiang Wu, ...	<1%
35	Student papers	Çankırı Karatekin University on 2022-10-18	<1%
36	Publication	Anita Mehta, Samreen Parveen, Ankur Kumar, Kaleem Ahmad. "Hepatic iron over...	<1%
37	Student papers	Lincoln University College on 2023-01-25	<1%
38	Publication	Antonella Meloni, Laura Pistoia, Paolo Ricchi, Sergio Bagnato et al. "Impact of ge...	<1%

39

Publication

J. Yuan, H.F. Gallardo, T. Rasalan, R. Ranganathan et al. "In vitro expansion of Ag-s... <1%

40

Internet

ejournal.ipinternasional.com <1%

RELATIONSHIP BETWEEN BLOOD TRANSFUSION FREQUENCY AND BLOOD IRON CONCENTRATION IN PATIENT WITH THALASSEMIA MAJOR

Vanessa Ainum Rizqita¹, Tjam Diana Samara^{2*}

¹Undergraduate Medical Study Program, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

²Anatomy Department, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

ARTICLE INFO

Corresponding author :

Tjam Diana Samara
Faculty of Medicine,
Universitas Trisakti

Email:

dianasamara@trisakti.ac.id

Kata kunci:

Frekuensi
Transfusi darah
Kadar feritin
Talasemia mayor
Anak

Keywords:

Frequency
Blood transfusion
Ferritin level
Thalassemia major
Children

Original submission:

September 11, 2025

Accepted:

March 26, 2026

Published:

April 25, 2026

ABSTRAK

Transfusi darah rutin yang dilakukan oleh pasien talasemia mayor dapat menyebabkan akumulasi zat besi berlebih dalam tubuh, yang berisiko menimbulkan komplikasi. Kadar feritin dalam serum digunakan untuk menilai akumulasi zat besi dalam tubuh. Penelitian ini bertujuan untuk mengetahui korelasi antara frekuensi transfusi dengan akumulasi zat besi pada pasien talasemia mayor. Penelitian ini menggunakan desain potong lintang dengan observasi analitik. Data dikumpulkan secara consecutive non random sampling dari rekam medis sebanyak 55 pasien talasemia mayor berusia 5-18 tahun yang menjalani transfusi darah di RS Hermina Jatinegara dari Januari hingga Desember 2023. Variabel bebas adalah frekuensi transfusi darah dalam setahun, sedangkan variabel tergantung adalah kadar rata-rata feritin dalam serum yang diukur tiga kali selama setahun (per empat bulan sekali). Data yang didapat dianalisis menggunakan uji korelasi Spearman. Hasil pada penelitian ini ditemukan frekuensi transfusi darah pasien talasemia mayor dalam satu tahun sebanyak 15 kali per tahun (berdasarkan nilai median). Rata-rata kadar feritin dalam serum sebesar 3957,73 ng/mL. (berdasarkan nilai median). Uji korelasi Spearman menunjukkan adanya korelasi lemah antara frekuensi transfusi darah dan kadar feritin ($r = 0,292$; $p = 0,031$). Simpulan penelitian ini adalah kenaikan frekuensi transfusi darah diikuti dengan kenaikan kadar besi dalam darah namun dengan kekuatan korelasi yang lemah.

ABSTRACT

Relationship Between Blood Transfusion Frequency and Blood Iron Concentration in Patient with Thalassemia Major. Routine blood transfusions performed on thalassemia major patients can cause excessive iron accumulation in the body, which carries the risk of complications. Serum ferritin levels are used to assess body iron stores. This study aims to determine the correlation between transfusion frequency and iron accumulation in patients with thalassemia major. This study used a cross-sectional design with analytical observation. Data was collected by consecutive, non-random sampling from the medical records of 55 patients with thalassemia major aged 5-18 years who underwent blood transfusions at Hermina Jatinegara Hospital from January to December 2023. The independent variable was the frequency of blood transfusions per year, while the dependent variable was the average serum ferritin level, measured three times a year (every 4 months). The data obtained were analyzed using the Spearman correlation test. The results of this study found that the frequency of blood transfusions in thalassemia major patients over 1 year was 15 per year (based on the median). The average serum ferritin level was 3957.73 ng/mL (based on the median value). Spearman's correlation test showed a weak correlation between blood transfusion frequency and ferritin levels ($r = 0.292$; $p = 0.031$). The study concluded that increased blood transfusion frequency was associated with higher blood iron levels, but the correlation was weak.

INTRODUCTION

Thalassemia is a chronic hemoglobin (Hb) disorder inherited in an autosomal recessive pattern. It is characterized by a reduction in the synthesis of globin chains, which may involve either the α - or β -globin chains.¹ According to the type of globin chain affected, thalassemia is classified

into α -thalassemia and β -thalassemia.² Thalassemia is classified into three types based on its clinical manifestations. Thalassemia minor (trait) is often asymptomatic or only mildly anemic. Thalassemia intermedia is characterized by symptoms that appear later compared to thalassemia major, with patients experiencing mild to moderate anemia. Thalassemia major represents the most severe form, requiring lifelong blood transfusions, and is typically detected during infancy or early childhood.³

Indonesia is one of the countries located within the global thalassemia belt, characterized by a high frequency of thalassemia genes.⁴ The prevalence of thalassemia carriers in Indonesia is estimated at 3–8% of the total population, with a birth rate of 23 per 1,000 among a population of approximately 240 million. According to 2021 data, there were 10,531 thalassemia patients in Indonesia, with more than 2,500 infants born each year with the condition.⁵ Thalassemia major cases requiring regular blood transfusions occur predominantly among children and adolescents. The majority of patients with thalassemia major who receive blood transfusions fall within the 5–18-year age group, as transfusion requirements during this period tend to remain stable to support growth and daily activities, particularly during the phase of active development.⁶

The primary management of anemia in patients with thalassemia major is blood transfusion. The goal of transfusion therapy is to maintain hemoglobin (Hb) levels above 9–9.5 g/dL.⁷ Blood transfusions are typically performed at intervals of 3–4 weeks, with the volume tailored to the patient's body weight and hemoglobin concentration.⁸ Blood transfusion therapy, monitoring, and management of iron overload are of primary concern. Since the human body lacks an active mechanism for eliminating excess iron, prolonged transfusion may lead to iron accumulation.⁷ Recent evidence consistently demonstrates that cumulative transfusional exposure is a major determinant of iron overload in transfusion-dependent thalassemia. Each unit of transfused red blood cells delivers approximately 200–250 mg of iron, and in the absence of a physiologic mechanism for active iron excretion, iron progressively accumulates over time.⁹ Moreover, clinical cohort data have shown a statistically significant association between total transfusion burden and serum ferritin concentration in patients with transfusion-dependent β -thalassemia major, indicating that greater cumulative exposure to transfusion contributes to higher iron loading as measured by ferritin levels.¹⁰ However, a literature review on post-transfusion ferritin levels in pediatric β -thalassemia major patients reported that mean ferritin concentrations in several analyzed studies were within recommended clinical targets. This observation indicates that ferritin levels may be maintained within acceptable limits when appropriate management strategies are implemented.¹¹ Serum ferritin measurement is the most commonly used method to evaluate iron accumulation (hemosiderosis) in thalassemia patients undergoing regular transfusions.¹²

The novelty of this study lies in its analytical cross-sectional approach in evaluating transfusion frequency as a quantitative variable associated with serum ferritin levels in thalassemia major patients within a specific local population. By providing updated local data, this study aims to clarify inconsistent findings reported in previous studies and to strengthen the evidence base for iron overload monitoring in routinely transfused patients.

METHOD

This study used a cross-sectional design with an analytical observational approach. The study population comprised patients with thalassemia major undergoing blood transfusions at Hermina Jatinegara Hospital, East Jakarta. Although thalassemia major consists of alpha thalassemia major dan beta thalassemia major², in this study they were not differentiated.

Based on the results of the sample-size calculation using the correlation test, 55 samples were obtained from 160 patients between January 1 and December 31, 2023. Sampling used the consecutive non-random sampling method. Samples were selected based on predefined inclusion and exclusion criteria. Inclusion criteria were patients with thalassemia major aged 5–18 years who had received blood transfusions and possessed complete medical records, including age, sex, annual transfusion frequency, and serial ferritin levels recorded between January and December 2023. Exclusion criteria were patients with coexisting hematological disorders, such as leukemia, hemophilia, or sickle cell anemia. Respondents were excluded if these diseases or disorders were listed in their medical records. If they were not listed, the patient was assumed not to have any other blood disorders.

Serum ferritin levels in this study were the average of three blood draws per year, performed at four-month intervals. The blood transfusion frequency in this study was the number of transfusions per year. All data were taken from medical records.

Data were analyzed using the Statistical Package for Social Sciences (SPSS) version 30. The Kolmogorov-Smirnov test was used to assess normality, as the sample size was >50. Since the data were not normally distributed, the Spearman correlation test was employed, with statistical significance set at $p < 0.05$.

RESULT

This study used medical records of patients with thalassemia major from the blood transfusion unit at Hermina Jatinegara Hospital in 2023, involving a total of 55 subjects.

Table 1. Distribution of study subject characteristics

Characteristic	Mean $\pm$ SD	Median (Range)	Frequency (n = 55)	Percentage (%)
Age	10,56 $\pm$ 3,711			
Gender				
Male			32	58,2
Female			23	41,8
Serum ferritin (ng/mL)		3957,73 (383,87 – 28566,16)		
Transfusion frequency (times/year)		15 (9 – 23)		

Note: Data are presented as mean $\pm$ standard deviation for normally distributed variables and median (range) for non-normally distributed variables.


SD: standard deviation

Based on Table 1, of the 55 subjects in the thalassemia blood transfusion unit, 32 (58.2%) were male, and 23 (41.8%) were female. Since the age data were normally distributed, age is presented as mean $\pm$ SD. The subjects' ages ranged from 5 to 18 years, with a mean age of 10.56 years and a standard deviation of 3.711 years.

In this study, because the data were not normally distributed, the distributions of serum ferritin levels and blood transfusion frequency were presented as medians. Based on Table 2, serum ferritin data obtained from serial measurements throughout 2023 showed 3,957.73 ng/mL

(according to a median value), with a minimum of 383.87 ng/mL and a maximum of 28,566.16 ng/mL. The frequency of blood transfusions in patients with thalassemia major in 2023 was 15 times per year (median), ranging from 9 to 23. Normal serum ferritin level for children are 7-140 ng/mL.¹³

Relationship between blood transfusion frequency and iron levels in patients with thalassemia major



*: Spearman's Correlation Test

Figure 1. Correlation between blood transfusion frequency and ferritin levels

The one-sample Kolmogorov–Smirnov test yielded a p-value of <0.001, indicating that the residuals were not normally distributed. Therefore, bivariate analysis was performed using Spearman's correlation test. The results showed a weak correlation between the frequency of blood transfusions and serum ferritin levels ($r = 0.292$, $p = 0.031$).

DISCUSSION

Characteristics of the study subjects

In this study, 55 subjects were included; the majority were male (32, 58.2%). This result is in line with the findings of Bhalodiya et al., who reported that 59% of their study population were male.¹⁰ However, the studies by Fianza et al. and Rochman et al. reported different findings, with a higher proportion of female subjects, accounting for 53.1% and 58.3%, respectively.^{7,14} This variation supports the theory that thalassemia is a genetic disorder with an autosomal recessive inheritance pattern and is not related to the sex chromosomes.¹⁵

The age distribution in this study ranged from 5 to 18 years, with a mean of 10.56 and a standard deviation of 3.711. This finding is consistent with the study by Almahmoud et al., which demonstrated a predominance of children (2–9 years) and adolescents (10–18 years) among patients with thalassemia major.¹⁶ The study by Nassim et al. also supports this finding, reporting an age distribution of subjects ranging from 2 to 14 years, with a mean age of 8.7 years.¹⁷ A considerable number of thalassemia cases are detected during childhood, when children present with worsening clinical symptoms, particularly increasing pallor, and subsequently require routine blood transfusions to maintain growth and daily activities.¹⁸

Distribution of the results of the blood iron level and blood transfusion frequency

In this study, the median ferritin level among subjects was 3957.73 ng/mL, which is consistent with the findings of Saparizki et al., who reported that the majority of thalassemia patients had ferritin levels exceeding 2500 ng/mL. Ferritin levels above 1000 ng/mL generally indicate iron overload (hyperferritinemia).¹⁹ The study by Kurban et al. also reported a median ferritin level of 1548 ng/mL, with a range of 161–3968 ng/mL, exceeding the recommended threshold value of 1000 ng/mL.²⁰ These findings suggest a substantial iron overload as a consequence of repeated blood transfusions among patients with thalassemia.²¹

This study found that patients with thalassemia major had a median annual transfusion frequency of 15 sessions. Aligning with the findings of Weiss et al., who reported a mean frequency of ≥ 12 transfusions per year.²² On the other hand, Akbar et al. reported that most patients received 12 or fewer transfusions annually.¹⁸ The frequency of blood transfusions in thalassemia patients varies depending on hemoglobin (Hb) levels and clinical condition. Transfusions are indicated when Hb levels fall below 7 g/dL in two consecutive measurements taken two weeks apart, or when Hb levels exceed 7 g/dL but are accompanied by clinical manifestations such as growth retardation, suspected bone fractures due to extramedullary hematopoiesis, and Cooley's facies.¹⁸

Relationship between blood transfusion frequency and iron levels in patients with thalassemia major

This study demonstrated a weak correlation between blood transfusion frequency and ferritin levels in patients with thalassemia major ($p = 0.031$; $r = 0.292$). This finding is consistent with the study by Bhalodiya et al., which reported a p-value of 0.00, indicating a significant increase in ferritin levels with rising total annual blood transfusion frequency.¹⁰ Similarly, Rochman et al. demonstrated a significant relationship between the number of transfusions and increased ferritin levels ($p = 0.003$).⁷ Thalassemia patients receiving repeated transfusions for anemia management are prone to elevated ferritin levels due to iron accumulation, as each unit of blood contains approximately 200–250 mg of iron, and the body lacks an inherent mechanism to excrete the excess.²³ The progressive accumulation of iron over time results in higher ferritin concentrations, thereby positioning transfusion frequency as a significant factor influencing iron overload.²⁴

Although no studies have been identified that explicitly state the absence of a correlation between transfusion frequency and ferritin levels, Pangestika et al. reported that hemoglobin and serum ferritin levels in children with beta-thalassemia major remained within the normal range (100 ng/mL) after transfusion.¹¹ This finding is consistent with the study by Spasiano et al., which reported that serum ferritin levels below 500 ng/mL can be achieved safely in some transfusion-dependent thalassemia (TDT) patients, indicating that transfusion frequency does not necessarily correlate directly with ferritin elevation in certain cases.²⁵ Even though a positive correlation between transfusion frequency and ferritin levels is observed, this relationship is significantly influenced by other factors, notably adherence to iron chelation therapy, which can markedly lower ferritin levels.²⁴ In addition, ferritin levels are influenced by other factors such as inflammation, infection, liver function, and genetic variations, which can cause differences in ferritin concentrations even among patients with similar transfusion histories.²⁶

CONCLUSION

In general, serum ferritin levels were higher than normal in study subjects who received more than 12 blood transfusions per year. However, the findings of this study indicate that the frequency of blood transfusions has only a weak effect on the increase in blood iron levels.

Future studies are advised to involve participants across a wider age range to gain a more comprehensive understanding of serum ferritin levels among thalassemia patients. In addition, subsequent research should address other potential determinants of ferritin levels, including variations in chelation therapy, treatment duration and compliance, nutritional status, and coexisting medical conditions.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest in this study.

ACKNOWLEDGMENTS

The authors are deeply grateful to Hermina Jatinegara Hospital for the permission and support provided throughout the data collection phase of this study.

REFERENCES

1. Wahidiyat P, Permono B. Hemoglobinopati dan thalassemia anak. In: Permono B, Sutaryo I, Ugrasena ID, Windastuti E, Abdulsalam M, editors. Buku ajar hematologi-onkologi anak. Jakarta: Ikatan Dokter Anak Indonesia; 2018:65-8.
2. Grentina, Wahidayat PA. Mengenal thalassemia [Internet]. Jakarta: Ikatan Dokter Anak Indonesia; 2016 [cited 2024 Apr 7]. Available from: <https://www.idai.or.id/artikel/seputar-kesehatan-anak/mengenal-thalassemia>
3. Rujito L, Siswandari W, Lestari DW, editors. Talasemia: genetik dasar dan pengelolaan terkini. Purwokerto: UNSOED Press; 2019:90.
4. Khikmah FA, Prijopranto HS. Prevalensi thalassemia beta minor dengan menggunakan indeks mentzer pada pasien anemia Suku Sabu di RSUD Sabu Raijua. Journal of Ners Community. 2022;13(2):301-6. doi:10.55129/jnerscommunity.v13i2.1917
5. World Health Organization. Regional desk review of haemoglobinopathies with an emphasis on thalassaemia and accessibility and availability of safe blood and blood products as per these patients' requirement in South-East Asia under universal health coverage [Internet]. 2023 [cited 2024 Apr 30]. Available from: <https://www.who.int/publications/i/item/9789290228516>
6. Fianza PI, Rahmawati A, Widiastha SH, Afifah S, Ghazali M, Indrajaya A. Iron overload in transfusion-dependent Indonesian thalassemic patients. Anemia. 2021;5581831. doi:10.1155/2021/5581831
7. Rochman F, Mulyantari NK, Sutirtayasa IW. Hubungan jumlah transfusi darah dan penggunaan kelasi besi dengan kadar ferritin pada pasien talasemia. Medika Udayana. 2019;8(9):1-6
8. Lal A, Wong T, Keel S, Pagano M, Chung J, Kamdar A. The transfusion management of beta thalassemia in the United States. Transfusion. 2021;61(10):3027-39. doi:10.1111/trf.16640
9. Taher AT, Cappellini MD. How I manage transfusional iron overload in patients with thalassemia. Blood. 2021;137(6):667-75
10. Bhalodiya VR, Valiya LG, Mehta NA, Padhariya BB. Correlation of serum ferritin level in transfusion-dependant thalassemia major patients: a study at a medical college affiliated hospital in Gujarat region. Int J Contemp Pediatr. 2023;10(3):330-3. doi:10.18203/2349-

3291.ijcp20230429

11. Pangestika AD, Astuti TD, Aryani T. Literature review: gambaran kadar hemoglobin dan ferritin serum pada anak thalassemia β mayor paska transfusi [Internet]. 2022 [cited 2024 Mar 15]. Available from: http://digilib.unisayogya.ac.id/6629/1/NASKAH%20PUBLIKASI_AYU%20DWI%20PANGESTIKA_1811304020%20-%20Ayu%20Dwi%20Pangestika
12. Yusuf AK, Magdalen AR, Karunia D. Hubungan kadar ferritin serum terhadap fraksi ejeksi ventrikel kiri pada pasien thalassemia β mayor anak di RSUD Kabupaten Kediri. *Jurnal Verdure*. 2022;4(1):123–8.
13. University of Rochester Medical Center. Ferritin (Blood) [Internet]. URM Health Encyclopedia; 2023 [cited 2026 Feb 24]. Available from: https://www.urmc.rochester.edu/encyclopedia/content?contenttypeid=167&contentid=ferritin_blood
14. Fianza PI, Rahmawati A, Wijaya I, Oehadian A, Prasetya D, Vidyaniati P. Gender disparities in psychological disturbances and quality of life among adolescent and adult patients with thalassemia: a review. *J Multidiscip Healthc*. 2024;17:1663-9. doi:10.2147/jmdh.s444592
15. Sadiq IZ, Abubakar FS, Usman HS, Abdullahi AD, Ibrahim B, Kastayal BS. Thalassemia: pathophysiology, diagnosis, and advances in treatment. *Thalass Rep*. 2024;14(4):81-102. doi:10.3390/thalassrep14040010
16. Almahmoud R, Hussein A, Khaja FA, Soliman AF, Dewedar H, Shareef ZA. Growth and endocrinopathies among children with β -thalassemia major treated at Dubai Thalassemia centre. *BMC Pediatr*. 2024;24:244. doi:10.1186/s12887-024-04670-w
17. Nassim MS, Mahmoud MA, Shady HA, Mohammed EAER. Assessment of disease knowledge gaps among beta thalassemia major patients and their caregivers. *Egypt Pediatr Assoc Gaz*. 2022;70:49. doi:10.1186/s43054-022-00143-w
18. Akbar TIS, Riyanti EY, Khairunnisa K, Wahyuni H. No association between frequency of blood transfusions and incompatibility of blood transfusions in thalassemia patients. *J Indones Med Assoc*. 2023;73(4):183-90. doi:10.47830/jinma-vol.73.4-2023-1069
19. Saporizki RS, Sumantri AF, Achmad S. Pengaruh kepatuhan minum obat kelasi besi dengan kadar ferritin pada pasien thalassemia di RSUD Al-Ihsan Bandung tahun 2021/2022. *Bandung Conference Series: Medical Science*. 2023;3(1):178-9. doi:10.29313/bcsms.v3i1.5824
20. Kurban LA, Almarri BK, Alshamsi MH, Abdelrahman SS, Alwahshi SG, Alhorani Q. Optimized serum ferritin prediction of iron overload in transfusion-dependent thalassemia: likelihood ratio and age-adjustment approach. *Ann Saudi Med*. 2023;43(2):90-6. doi:10.5144/0256-4947.2023.90
21. Sarkar K, Pramanik N, Goswami RP, Mandal PK, Chowdhury P, Jana K. Correlation of liver function test with different age and sex group and with ferritin level in thalassemia. *Indian J Public Health Res Dev*. 2020;11(6):452-7. doi:10.37506/ijphrd.v11i6.9819
22. Weiss M, Parisi Jun M, Sheth S. Clinical and economic burden of regularly transfused adult patients with β -thalassemia in the United States: a retrospective cohort study using payer claims. *Am J Hematol*. 2019;94(5):E129-32. doi:10.1002/ajh.25429
23. Shah F, Huey K, Deshpande S, Turner M, Chitnis M, Schiller E. Relationship between serum ferritin and outcomes in β -thalassemia: a systematic literature review. *J Clin Med*. 2022;11(15):4448. doi:10.3390/jcm11154448
24. Coates TD. Iron overload in transfusion-dependent patients. *Hematology Am Soc Hematol Educ Program*. 2019;2019(1):337-44. doi:10.1182/hematology.2019000036
25. Spasiano A, Meloni A, Costantini S, Quaia E, Cademartiti F, Cinque P. Setting for "normal" serum ferritin levels in patients with transfusion-dependent thalassemia: our current strategy. *J*

Clin Med. 2021;10(24):5985. doi:10.3390/jcm10245985

26. Mohamed YA, Morgan DS, Hussein AM, Mohamed AA. Noninvasive assessment and risk factors of liver fibrosis in pediatric patients with beta thalassemia major using transient elastography. Beni Suf Univ J Basic Appl Sci. 2022;11:114. doi:10.1186/s43088-022-00314-z

[Dashboard](#)

[Explore SINTA](#)

[Mutation History](#)

[List Verificator PT](#)

[My SINTA](#)

DASHBOARD

Your Overview

SINTA SCORE OVERALL

792

SINTA SCORE 3YR

117

AFFILIATION SCORE OVERALL

792

AFFILIATION SCORE 3YR

117

SCOPUS H-INDEX

1

WOS H-INDEX

3

GOOGLE SCHOLAR H-INDEX

10

[Request Score Recalculation](#)

[Review affiliation](#)
Help colleagues find you.
[REVIEW](#)

[Add co-authors](#)
We have co-authors suggestions.
[ADD](#)

Tjam Diana Samara

Lecturer, Faculty of Medicine Universitas Trisakti
Verified email at trisakti.ac.id
[Anatomy](#) [Occupational Health](#)

[FOLLOW](#)

☐

TITLE

CITED BY

YEAR

☐ Relationship between blood transfusion frequency and blood iron concentration in patient with thalassemia major
VA Rizqila, TD Samara
Jurnal Kedokteran dan Kesehatan: Publikasi Ilmiah Fakultas Kedokteran ...
2026

☐ 41 The Historical Evolution of Occupational Exposure to Asbestos, Silica, and Inorganic Dusts in Indonesia: Regulation, Industry Influence, and Shifting Recognition of ...
N Hainunisa, A Dwi Lestari, H Nuriyah, A Mohamad Ridwan, L T Merjanti, ...
Journal of Occupational Medicine and Toxicology 21 (Suppl1), 41
2026

☐ Relationship between Serum 25-Hydroxyvitamin D Levels and Cardiovascular Disease Risk among Road Sanitations Workers
M Wartono, TD Samara, A Kosasih, D Mediana
Althea Medical Journal 13 (1), 28-34
2026

☐ Perbedaan jumlah leukosit pada pasien gagal ginjal kronik berdasarkan rutinitas hemodialisis
TA Derajat, TD Samara
Jurnal Kesehatan Tambusai 7 (1), 776-781
2026

☐ Correlation between COHb, HIF-1 α levels, and smoking habits with physical fatigue in online motorcycle taxi drivers
M Wartono, TD Samara, AD Lestari, A Kosasih, N Novendy
Indonesian Journal of Biomedicine and Clinical Sciences 58 (26), 1-9
2026

☐ Perbedaan profil kolesterol darah antara kehamilan pre-eklampsia dan kehamilan normal
SI Kemala, TD Samara
Jurnal Penelitian dan Karya Ilmiah Lembaga Penelitian Universitas Trisakti ...
2026

☐ Hubungan performa akademik dan jenis kelamin dengan Nomophobia pada siswa SMA
TD Samara, ML Simon
Jurnal Lentera Kesehatan Masyarakat 4 (2), 104-111
2025

Cited by

[VIEW ALL](#)

	All	Since 2021
Citations	471	222
h-index	10	7
i10-index	10	5

Co-authors

[EDIT](#)

Maadalea Wartono

Magdalena Wartono
Universitas Trisakti

Purnamawati Tjhin
Universitas Trisakti

Dian Mediana
Universitas Trisakti

Maria Regina Rachmawati
University of Gunadarma

Ani Retno Prijanti
Universitas Indonesia

Isabella Kurnia Liem
Universitas Indonesia

Lie Tanu Merjanti Susanto
Universitas Trisakti