

Website Maintenance 2026-02-04

Due to website maintenance, if you can not log in to our Journal, please clear the cache of your browser on your computer.

The fastest method is pressing Ctrl + Shift + Delete (Windows) or Cmd + Shift + Delete (Mac) to open the clearing menu, select "Cookies and site data, and Temporary cached files and pages" or "Cached images and files" (set time range to "All time"), and click "Clear Data".

[READ MORE >](#)**Old Website:****CURRENT ISSUE****Vol 58 No 1 (2026)****PUBLISHED:** 2026-01-20**ARTICLES****Modulation of Caspase-3 Expression and Spermatogenic Cells by Urtica dioica Extract in Obesity-Induced Male Rats**

Kabir Ardiansyah Tangkari, Jurnalís Gempañing Tyas, Harni Sutiani, Zaenudin, Dicky Mochammad Rizal, Jajar Setyawan

 Abstract views: 50 |  views: 48**Non-contact electro capacitive cancer therapy (ECCT) modulate the mRNA expression of p53, Apaf-1, survivin, NF-κB, TSP-1 and bFGF in DMBA-induced breast cancer rat**

Nurul Hidayah, Agung Putra, Firman Alamsyah, Rarastoeti Pratiwi

 Abstract views: 47 |  views: 40**Evaluation of the current clinical and bacteriological profile in the tubotympanic type and the atticotympanic type chronic suppurative otitis media**

Anton Budhi Darmawan, Bella Jovita Darmayan, Vitasari Indriani

 Abstract views: 54 |  views: 43



Correlation between COHb, HIF-1 α levels, and smoking habits with physical fatigue in online motorcycle taxi drivers

Magdalena Wartono, Diana Samara Tjam, Ade Dwi Lestari, Adrianus Kosasih, Novendy Novendy



Abstract views: 34 | views: 18

Hand-foot-mouth disease in the elderly: A case report

Rizky R. Wijayanti, Amelia Pungky, Cut A.W. Sawitri, Agnes S. Siswati, Flandiana Yogianti, Nabila Arkania



Abstract views: 30 | views: 29

Clinical effects of Spirulina supplementation on hemoglobin levels in anemic pregnant women: a systematic review and meta-analysis

Made Favian Budi Gunawan, Cindy Thiovany Soetomo, Putu Bagus Darmayasa, I Gusti Ayu Cintya Paramyta, I Putu Prabawa Jayakusuma, I Made Indera Gunawan



Abstract views: 30 | views: 29

Immune status in pregnant women against Mycobacterium tuberculosis: A scoping review

Qonita Adzkia, Maria Jeniffer Ann Siburian, Vycke Yunivita



Abstract views: 42 | views: 29

[VIEW ALL ISSUES >](#)

Editor in Chief

- [Hardyanto Soebono](#), Department of Dermatology and Venereology Faculty of Medicine, Public Health, and Nursing Universitas Gadjah Mada, Indonesia, Indonesia, Indonesia

Associate Editors

- [Berry Juliandi](#), Department of Biology, IPB University, Indonesia
- [Dewi Kartikawati Paramita](#), Department of Histology and Cell Biology, Faculty of Medicine, Universitas Gadjah Mada, Indonesia
- [Eti Nurwening Sholikhah](#), Department of Pharmacology and Therapy Faculty of Medicine, Universitas Gadjah Mada, Indonesia
- [Fara Silvia Yuliani](#), Department of Pharmacology and Therapy Faculty of Medicine, Universitas Gadjah Mada, Indonesia
- [Gunadi](#), Universitas Gadjah Mada, Sleman, Indonesia
- [Junaedy Yunus](#), Faculty of Medicine, Public Health, and Nursing, Universitas Gadjah Mada, Indonesia
- [Mae Sri Hartati Wahyuningsih](#), Faculty of Medicine, Public Health, and Nursing Universitas Gadjah Mada, Indonesia
- [Mei Neni Sitaresmi](#), Department of Pediatric Faculty of Medicine, Universitas Gadjah Mada, Indonesia
- [Mustofa](#), Faculty of Medicine, Public Health, and Nursing Universitas Gadjah Mada, Indonesia
- [Loeki Enggar Fitri](#), Department of Parasitology, Faculty of Medicine, University of Brawijaya, Indonesia
- [Pamungkas Bagus Satriyo](#), Department of Pharmacology and Therapy Faculty of Medicine, Universitas Gadjah Mada, Indonesia
- [Retno Danarti](#), Faculty of Medicine, Public Health, and Nursing, Universitas Gadjah Mada, Indonesia
- [Rina Agustina](#), Department of Nutrition, Faculty of Medicine, Universitas Indonesia – Dr. Cipto Mangunkusumo General Hospital, Jakarta, Indonesia
- [Sultana MH Faradz](#), Center for Biomedical Research , Faculty of Medicine, Universitas Diponegoro, Semarang, Indonesia
- [Supangat](#), Department of Pharmacology, Faculty of Medicine, Universitas Jember, Indonesia
- [Yohanes Widodo Wirohadidjojo](#), Faculty of Medicine, Public Health, and Nursing Universitas Gadjah Mada, Indonesia

International Advisory Board

- [Agostino Pierro](#), Department of Surgery, University Toronto, Canada
- [Alexis Valentin](#), Pharmacochimie des Substances Naturelles et Pharmacophores Redox Université Paul Sabatier, Faculté de Pharmacie, France
- [Carina Hanashima](#), RIKEN Center for Developmental Biology, Japan
- [Dorothy E. Oorschot](#), University of Otago, New Zealand, New Zealand
- [Fatima Shad Kaneez](#), University of Technology Sydney, Australia
- [Françoise Benoit-Vical](#), Service de Parasitologie et Mycologie - Hôpital Rangueil-Larrey, France
- [Gan Siew Hua](#), Director, Human Genome Centre, School of Medical Sciences, Universiti Sains Malaysia, Malaysia
- [Hisahide Nishio](#), Division of Public Health, Department of Environmental Health and Safety, Faculty of Medicine, Kobe University Graduate School of Medicine, Japan
- [Hugo A. Heij](#), University Medical Centre of Utrecht/Wilhelmina Children's Hospital and Princess Maxima Centre for Paediatric Oncology, Netherlands
- [I. Bing Tan](#), The Netherlands Cancer Institute – Antoni van Leeuwenhoek Hospital, Department of Head & Neck Oncology & Surgery, Netherlands
- [Isaak Effendy](#), Dermatologische Klinik, Philipps-Universität Marburg, Germany
- [Jaap Middeldorp](#), Department of Pathology, VU University Medical Center, Netherlands
- [Mulyoto Pangestu](#), Monash Institute of Medical Research Monash University, Australia
- [Rik J. Scheper](#), Department of Pathology, VU University Medical Center, Netherlands
- [Teguh Haryo Sasongko](#), Human Genome Center, School of Medical Sciences, Universiti Sains Malaysia, Malaysia
- [William R Faber](#), Academic Medical Center, University of Amsterdam Department of Dermatology, Academic Medical Centre, Netherlands



INDONESIAN JOURNAL OF BIOMEDICINE AND CLINICAL SCIENCES

UNIVERSITAS GADJAH MADA

P-ISSN : 01261312 <> E-ISSN : 23563931 Subject Area : Health

 0.375 Impact	 1496 Google Citations	 Sinta 2 Current Accreditation
--	---	---

[Google Scholar](#) [Garuda](#) [Website](#) [Editor URL](#)

History Accreditation

Correlation between COHb, HIF-1 α levels, and smoking habits with physical fatigue in online motorcycle taxi drivers

Magdalena Wartono^{1,2,3*}, Tjam Diana Samara^{1,2,3}, Ade Dwi Lestari^{2,3}, Adrianus Kosasih⁴, Novendy Novendy⁵

¹Department of Anatomy, Faculty of Medicine, Universitas Trisakti, Jakarta 11440, Indonesia,

²Department of Occupational Medicine, Faculty of Medicine, Universitas Trisakti, Jakarta 11440, Indonesia,

³Occupational Medicine Study Center, Faculty of Medicine, Universitas Trisakti, Jakarta 11440, Indonesia,

⁴Department of Internal Medicine, Faculty of Medicine, Universitas Trisakti, Jakarta 11440, Indonesia,

⁵Department of Public Health, Faculty of Medicine, Universitas Pelita Harapan, Jakarta 11440, Indonesia

<https://doi.org/10.22146/inajbcs.v58i1.23175>

ABSTRACT

Submitted: 2025-07-18

Accepted : 2026-01-02

Online motorcycle taxi drivers have varying working hours and rest periods which can influence the occurrence of fatigue among them. Additionally, working on the road made them exposed to air pollution, including carbon monoxide. Several studies have reported a relationship between carbon monoxide and fatigue levels in specific types of occupations. Therefore, the aim of this study is to determine the correlation between smoking habits, occupational factors, carboxyhemoglobin (COHb) levels, and work fatigue, as well as the correlation between hypoxia inducible factor 1-alpha (HIF-1 α) and COHb levels. Demographic characteristics, smoking habits, and occupational factors were measured using a questionnaire. COHb and HIF-1 α levels were measured through blood biomarker examinations, while fatigue levels were assessed using the IFRC questionnaire. Data analysis was conducted using Spearman correlation tests in the SPSS program. The results showed that majority of respondents were male (87%), with the highest age range being 19-44 yr (85.5%). More than 50% of respondents were classified as overweight or obese. The majority of them were light smokers (78.3%). Most respondents had normal COHb levels, and in terms of fatigue levels, a larger proportion experienced mild fatigue (66.7%). No correlation between COHb levels, smoking habits, occupational factors, and fatigue levels as well as between HIF-1 α and COHb levels ($p>0.05$). In conclusion, no correlation is found between COHb levels, smoking habits, occupational factors, and fatigue levels. Similarly, there is no correlation between HIF-1 α and COHb levels.

ABSTRAK

Pengemudi ojek online memiliki jam kerja dan waktu istirahat yang bervariasi, yang dapat menyebabkan kelelahan. Selain itu, bekerja di jalan raya membuat pengemudi terpapar polusi udara, termasuk karbon monoksida. Beberapa penelitian melaporkan adanya hubungan antara karbon monoksida dan tingkat kelelahan pada jenis pekerjaan tertentu. Tujuan penelitian ini adalah untuk mengetahui hubungan antara kebiasaan merokok, faktor pekerjaan, kadar karboksihemoglobin (COHb), dan kelelahan kerja, serta hubungan antara HIF-1 α dan kadar COHb. Karakteristik demografis, kebiasaan merokok, dan faktor pekerjaan diukur menggunakan kuesioner. Kadar COHb dan HIF-1 α diukur melalui pemeriksaan biomarker darah, sedangkan tingkat kelelahan dinilai menggunakan kuesioner *Industrial Fatigue Research Committee* (IFRC). Analisis data dilakukan dengan uji korelasi Spearman menggunakan program SPSS. Hasil penelitian menunjukkan mayoritas responden laki-laki (87%), dengan rentang usia terbanyak 19–44 tahun (85,5%). Lebih dari 50% responden tergolong kelebihan berat badan atau obesitas. Sebagian besar responden merupakan perokok ringan (78,3%). Mayoritas responden memiliki kadar COHb normal, dan berdasarkan tingkat kelelahan, proporsi terbesar mengalami kelelahan ringan (66,7%). Tidak ditemukan hubungan antara kadar COHb, kebiasaan merokok, faktor pekerjaan, dan tingkat kelelahan, serta antara HIF-1 α dan kadar COHb ($p>0,05$). Simpulan, tidak ada hubungan antara kadar COHb, kebiasaan merokok, faktor pekerjaan, dan tingkat kelelahan. Demikian pula, tidak ada hubungan antara HIF-1 α dan kadar COHb.

Keywords:

COHb;

HIF-1 α ;

smoking habits;

work fatigue;

online motorcycle taxi drivers

*corresponding author: magdalena_w@trisakti.ac.id

INTRODUCTION

Online motorcycle taxis, known as *ojek online* had been one of the transportations in Indonesia since 2010. According to the Joint Action Presidium of Two Wheeled Vehicles (Garda) Indonesia, the number of online motorcycle taxi drivers in 2020 was estimated to be around 4 million people, with a quarter of them located in the Jabodetabek area (Jakarta, Bogor, Tangerang, and Bekasi).¹ Motorcycle taxi drivers often work long hours, exposes to weather changes, air pollution, and traffic conditions.² Working in such hazardous environments makes them vulnerable to work related health problems.³

A study conducted in Vietnam reported that fatigue and lower back pain had been found significantly higher compared to other health issues among motorcycle taxi drivers.² Approximately 22.59% of them were reported to experience fatigue. Fatigue is the result of prolonged mental or physical activities that can affect a person's performance and impair their mental alertness, leading to dangerous errors. Fatigue typically refers to a decline in performance. Psychologically, fatigue can be interpreted as not having enough energy to perform work and experiencing a reluctance to continue tasks. Thus, a fatigued individual receives signals from their body that ongoing activities, both physical and mental, should be terminated. Essentially, the effects include a decrease in the capacity to perform mental and physical tasks.⁴ Long-term fatigue can have impacts on health, such as musculoskeletal disorders, poor mental health status, and increased errors and accidents in the workplace.⁵

A cohort study of Ahanhanzo *et al.*,⁶ in Benin found that fatigue is a significant risk factor for road crashes on two-wheeler especially higher at nighttime. Fatigue has a multidimensional etiology and can originate from both inside and outside the workplace. Fatigue

encompasses physical, emotional, behavioral, and cognitive components. Previous research has often discussed job-related factors and sleep-related factors as contributors to work-related fatigue. Monotonous or repetitive work, long working hours, shift work (day or night), work patterns (permanent or rotating), and the duration of shift work have been found to be associated with fatigue. Inadequate work environment factors, such as lighting, temperature, humidity, noise, and ventilation, are also work-related risk factors for fatigue.⁷

A study showed that taxi motorcycle drivers who complained of fatigue are 10.6% among non-smokers and 14.3% among smokers. Although the onset of fatigue can be caused by various factors, complaints such as headaches, irritability, visual perceptual disturbances, fatigue, and hyporeflexia were more frequently found significantly in motorcycle taxi drivers with high levels of carboxyhemoglobin (COHb).⁸ With the binding of hemoglobin to CO, tissue oxygenation is disrupted, leading to tissue hypoxia, especially in organs sensitive to oxygen deficiency such as the kidneys, brain, and heart, which can have an impact on a person's health. During hypoxia, the main effector response is hypoxia inducible factor 1-alpha (HIF1- α), a transcription factor. This protein is responsive to low oxygen levels in the cellular environment and accumulates rapidly under hypoxic conditions.^{9,10} Due to the significant impact of CO exposure, the study aimed to investigate the correlation between smoking habits, work-related factors (year of service and duration of work), and COHb levels with the fatigue level of online taxi motorcycle drivers. We also intended to examine the correlation between HIF1- α and COHb levels.

MATERIAL AND METHODS

The study was conducted during Covid-19 pandemic period in September 2022 at the Biomedical Laboratory 1, Faculty of Medicine, Universitas Trisakti,

West Jakarta. This research was a cross-sectional approach. The study population consisted of online motorcycle taxi drivers working in Jakarta and its surroundings. The subjects were 69 online motorcycle taxi drivers who had been working for at least 1 year and were willing to sign an informed consent. The exclusion criteria included online motorcycle taxi drivers with chronic respiratory diseases such as chronic obstructive pulmonary disease or other chronic conditions such as heart failure, kidney failure, and malignancy. This study has obtained ethical clearance issued by the Faculty of Medicine, Universitas Trisakti with number 162/KER/FK/VIII/2022.

Questionnaire

The data collected in this study were primary data obtained using a questionnaire that assessed sociodemographic characteristics like age, sex, and respondents' occupations (years and duration of work). The smoking habits were assessed using the Brinkman Index by multiplying the quantity of cigarettes smoked per day by the total duration of the smoking habit in years. The results were classified as light (0-199), moderate (200-599) and heavy smoker (>600). Body mass index (BMI) was obtained by measuring body weight (BW) in kg and height (H) in m and were calculated using a formula ($BMI = BW/H^2$). The classification of BMI was according to World Health Organization (WHO) for Asian.

Fatigue level assessment

Fatigue level assessment was conducted through interviews using the Industrial Fatigue Research Committee (IFRC) questionnaire.¹¹ The classification used was as follows:

Examination of COHb levels

The COHb level were obtained from venous blood sample. The blood was

collected using a 3 mL no anticoagulant red-top vacutainer tube. The sample was then placed in an icebox at a temperature of 11.7°C and stored in a refrigerator overnight. The examination of COHb levels using the Conway diffusion method was conducted at Prodia Laboratory. The reference values for COHb levels are <1.50% for non-smokers and 3-15% for smokers based on the reference provided by the laboratory.

Examination of HIF-1 α

Human hypoxia-inducible factor 1-alpha (HIF-1 α) in serum is measured using The Invitrogen™ HIF-1 Alpha ELISA Kit from ThermoFisher, Austria, conducted at Prodia Laboratory.

Data analysis

Univariate analysis is performed by describing the research results following the principles of descriptive statistics and presenting them in a frequency distribution table. Bivariate analysis is conducted using the Spearman correlation formula due to the non-normal distribution of the data. Data analysis is performed using SPSS 21.0 software for Windows. A p value < 0.05 is considered significance.

RESULTS

This study was conducted on 69 respondents consisting of online motorcycle taxi drivers from various areas in Jakarta and its surroundings. Out of the 69 respondents, 87% were male drivers, with the majority falling within the age range of 19-44 yr (85.5%). More than 50% of the respondents were classified as overweight or obese. The majority of them were light smokers (78.3%). Most of the respondents had normal COHb levels, and in terms of fatigue levels, the majority experienced mild fatigue (66.7%), none of them experienced high level of fatigue (TABLE 2)

TABLE 1. Classification of subjective fatigue level based on total score.¹¹

Fatigue level	Individual total score	Fatigue classification	Improvement measurement
1	30 – 52	Low	No improvement action is required.
2	53 – 75	Medium	Improvement action may be required in the future.
3	76 – 98	High	Immediate action is required.
4	99 – 120	Very high	Comprehensive action is required as soon as possible.

TABLE 2. Frequency distribution of subject characteristics in the study

Variables	Frequency (n)/ median (range)	Percentage (%)
Sex		
Male	60	87
Female	9	13
Ages (yr)		
19 – 44	59	85.5
45 – 59	9	13
≥ 60	1	1.4
Body mass index		
< 18.5	6	8.7
18.5 – 22.9	20	29
23 – 24.9	7	10.1
25 – 29.9	18	26.1
≥ 30	18	26.1
Smoking habit (Brinkman index)		
Light (0 – 199)	54	78.3
Moderate (200 – 599)	14	20.3
Heavy (≥ 600)	1	1.4
Years of service (yr)	5 (1 – 11)	-
Duration of work (hr/d)	12 (1 – 16)	-
HIF1- α (pg/mL)	4.9 (2.58 – 72.09)	-
COHb level (%)	0.39 (0.10 – 6.70)	-
COHb level ategories		
Normal	48	69.6
Not Normal	21	30.4
Fatigue level		
Low (30 – 52)	46	66.7
Medium (53 – 75)	23	33.3
High (76 – 98)	0	0
Very High (99 – 120)	0	0

Regarding the normality test of the data, the Kolmogorov-Smirnov test was used since the sample size was only 69. All variable data had a non-normal distribution ($p > 0.05$). Therefore, the Spearman correlation test was used to analyze the correlation between non-parametric data.

TABLE 3 shows that among all

examined parameters (age, BMI, smoking habit, work duration, length of service, COHb concentration), there were no correlations between these parameters and fatigue level. The analysis results between HIF1- α and COHb levels also did not show any correlation between these two variables (TABLE 4).

TABLE3. Correlation between demographic characteristics, occupational factors, COHb levels, and fatigue levels in online motorcycle taxi drivers.

Variable	Fatigue level	
	r	p*
Ages	0.159	0.192
BMI	0.127	0.299
Smoking habit	0.116	0.345
Working hours	0.052	0.673
Duration of employment	-0.039	0.749
COHb level	-0.42	0.735

*Spearman's correlation test

TABLE 4 Correlation between HIF1- α and COHb levels

Variable	COHb levels	
	r	p*
HIF1-a	-0.017	0.192

*Spearman's correlation test

DISCUSSION

Just like a study in Taiwan, there were more men working as food delivery riders compared to women (74% vs 26%). Similarly, in this study, male respondents were more dominant. In terms of age range, there was not much difference compared to Chen's study where the age range of online motorcycle taxi drivers was 31-50 yo.¹² The median duration of work per day for the study subjects was 12 hr, like online motorcycle taxi drivers in Medan who have an average of 12 hy of work per day.¹³ The CO levels in the blood of the respondents mostly fell into the normal category. During the data collection in September 2022, the concentration of CO in the air in Jakarta ranged from 0.58 to 2.34 mg/m³, which is below the safe limit of 10 mg/m³. Therefore, the amount of CO concentration exposure was not significant.¹⁴ The results showed that 66.7% of the respondents experienced mild fatigue, similar to the study in Bekasi where most of online motorcycle taxi drivers experienced mild fatigue (50.9%).¹⁵ In Indonesia, there are three online motorcycle companies where the drivers do not have fixed working hours; they can work as they please. As a result, the duration of work and rest varies greatly, which certainly affects the fatigue level of online motorcycle taxi drivers. However, a study in Depok found that 43% of the surveyed online motorcycle taxi drivers had experienced near-missed accidents, and one of the causes was fatigue, although the level of fatigue, whether mild or moderate, was not specified.¹⁶

In this study, smoking habits were not correlated with the level of fatigue. This is different from a study in Turkey which found that in smokers, the total fatigue scores were higher than in non-smokers ($p < 0.05$).¹⁷ Similarly, a study by Li *et al.*¹⁸ demonstrated that subjects who still smoked had nearly twice the

risk of experiencing fatigue (OR=1.956). The nicotine content in cigarettes can lead to a state of decreased energy (fatigue) characterized by a decrease in energy intake and an increase in energy expenditure. Energy intake and expenditure depend on regulation in the brain. Nicotine can affect energy expenditure by increasing the release of norepinephrine, serotonin, and other factors. These factors influence the brain to modify appetite and metabolic rate.¹⁹ Another study by Corwin²⁰ reported that research subjects who were heavy smokers tended to experience fatigue, while in this study, there were more subjects who belonged to the group of light smokers.

There was no correlation between work hours and duration of employment with the level of fatigue ($p > 0.05$). Consistent with a study on online motorcycle taxi drivers in Bekasi, work duration and duration of employment were not significantly associated with the occurrence of fatigue in the respondents ($p > 0.05$), but were related to previous medical history.¹⁵ This is different from the findings of a study by Ferusgel *et al.*,¹³ which demonstrated a relationship between work duration and the level of fatigue, as indicated by increased blood lactate levels. Workers with longer employment tend to experience work fatigue more frequently due to declining stamina, especially caused by boredom.^{21,22}

The COHb levels in this study were also not correlated with the level of fatigue. This differs from a study by Carter, which found a significant association between fatigue and the level of CO concentration in the air on the same day ($p = 0.001$). However, this significance was only observed in subjects who experienced physical complaints within a year, as these subjects experienced concurrent immune and inflammatory disturbances that could contribute to physical fatigue.²³ In this study, subjects

with chronic diseases were excluded, and the CO levels in the air during the study were within the permissible limits.

The analysis results between HIF1- α and COHb levels also did not show any correlation between these two variables. The binding of Hb with CO can reduce tissue oxygenation capacity.²⁴ Intermittent hypoxia stimulates the carotid body, inducing the formation of reactive oxygen species (ROS) and increasing intracellular Ca²⁺ levels, both of which enhance the expression of hypoxia-inducible factor HIF-1 α .¹⁰ Significant CO exposure increases COHb concentrations above 10% and CO poisoning is twice as likely at that level.²⁵ However, in this study, the median COHb level was 0.39 (0.1-6.7) %, which showed the subjects were not significantly exposed to CO.

No significant correlation was found between the measured parameters and fatigue levels in online motor taxi drivers, suggesting these factors remain within tolerable limits. Nevertheless, with 33.3% of the population reporting moderate fatigue, preventative measures are essential to avoid more severe conditions. Intervention strategies should focus on health education, specifically regarding respiratory protection, smoking cessation, nutrition, and sufficient recovery time.

Some limitations were observed in this study such as fatigue levels were still subjectively measured using a questionnaire and did not include muscle fatigue biomarkers, nor did it exclude mental fatigue. Blood samples for COHb level examination were not taken after work, and during data collection, the air pollution levels in Jakarta were still categorized as good due to the pandemic period. The strength of this study is the previous research on COHb and HIF-1 α levels among online motorcycle taxi drivers in the Jakarta area were scarce.

CONCLUSION

This study shows no correlation between smoking habits, occupational factors and COHb levels with fatigue levels among online motorcycle taxi drivers. Similarly, there is no correlation between the hypoxia expression factor HIF-1 α and COHb levels. For future research, it is recommended to measure COHb levels and fatigue levels before and after work, accompanied by a comparison with air pollution levels.

ACKNOWLEDGEMENT

No conflict of interest in this study.

REFERENCES

1. Azka RM. Berapa sih jumlah pengemudi ojek online?. Bisnis.com. 2019 Nov. <https://ekonomi.bisnis.com/read/20191112/98/1169620/berapa-sih-jumlah-pengemudi-ojek-online-simak-penelusuran-bisnis.com> Accessed at 3 January 2022
2. Truong L, Tay R, Nguyen H. Investigating health issues of motorcycle taxi drivers: A case study of Vietnam. *J Transp Health* 2021; 20(3):100999. <https://doi.org/10.1016/j.jth.2020.100999>
3. Sanchez HM, Sanchez EGM, Barbosa MA, Porto CC, Approbato MS. Comparison of quality of life and work ability of taxi and motorcycle taxi drivers: Evidence from Brazil. *Int J Environ Res Public Health* 2019; 16(4):666. <https://doi.org/10.3390/ijerph16040666>
4. Sadeghniiat-Haghighi K, Yazdi Z. Fatigue management in the workplace. *Ind Psychiatry J* 2015; 24(1):12-7. <https://doi.org/10.4103/0972-6748.160915>
5. Tian F, Shu Q, Cui Q, Wang L,

- Liu C, Wu H. The mediating role of psychological capital in the relationship between occupational stress and fatigue: A cross-sectional study among 1,104 Chinese physicians. *Front Public Health* 2020; 8:12.
<https://doi.org/10.3389/fpubh.2020.00012>
6. Glèlè-Ahanhanzo Y, Kpozèhouen A, Paraïso N, Makoutodé P, Biaou C, Remacle E, *et al.* Disability and related factors among road traffic accident victims in benin: study from five public and faith-based hospitals in urban and suburban areas. *Open J Epidemiol* 2018; 08(04):226-41.
<https://doi.org/10.4236/ojepi.2018.84018>
7. Tang FC, Li RH, Huang SL. The association between job-related psychosocial factors and prolonged fatigue among industrial employees in Taiwan. *PLoS One* 2016; 11(3):e0150429.
<https://doi.org/10.1371/journal.pone.0150429>
8. Silva LA da, Robazzi ML do CC, Terra F de S. Relation between workplace accidents and the levels of carboxyhemoglobin in motorcycle taxi drivers. *Rev Lat Am Enfermagem* 2013; 21(5):1119-26.
<https://doi.org/10.1590/S0104-11692013000500015>
9. Harrison H, Pegg HJ, Thompson J, Bates C, Shore P. HIF1-alpha expressing cells induce a hypoxic-like response in neighbouring cancer cells. *BMC Cancer* 2018; 18(1):674.
<https://doi.org/10.1186/s12885-018-4577-1>
10. Semenza GL, Prabhakar NR. The role of hypoxia-inducible factors in carotid body (patho) physiology. *J Physiol* 2018; 596(15):2977-83.
<https://doi.org/10.1113/JP275696>
11. Tarwaka, Solikhul H, Sudiajeng L. Ergonomi untuk keselamatan, kesehatan kerja dan produktivitas. [Tesis]. Surakarta: UNIBA; 2004.
12. Chen CF. Investigating the effects of job stress on the distraction and risky driving behaviors of food delivery motorcycle riders. *Saf Health Work* 2023; 14(2):207-14.
<https://doi.org/10.1016/j.shaw.2023.03.004>
13. Ferusgel A, Butar-Butar MH, Widya A, Napitupulu LH, Chaniago AD. Risk factors of an online motorcycle taxi (ojek online) fatigue in Medan. *Adv Health Sci Res* 2020; 24:76-80.
<https://doi.org/10.2991/ahsr.k.200311.014>
14. Simatupang MM, Veronika E, Irfandi A, Garmini R. Environmental health risk assessment of air pollutants in online motorcycle taxi drivers in the special capital region of Jakarta. *J Kes Ling* 2022; 14(2):114-21.
<https://doi.org/10.20473/jkl.v14i2.2022.114-121>
15. Effendi L, Syadiah T. Analysis of factors associated with subjective fatigue among motorcycle drivers in online ojek. In: *Internationalization of Islamic Higher Education Institutions Towards Global Competiveness*. Semarang; 2018. p. 139-45.
16. Nurhafizhah T, Susilowati IH, Maulana A, Habibullah MF. Safety and fatigue risk factors among online motorcycle drivers in Depok City, Indonesia. *KnE Life Sci* 2018; 4(5):702.
<https://doi.org/10.18502/cls.v4i5.2600>
17. Ergin C, Yurdalan SU, Demirbiken I, Zengin O. Fatigue and physical activity levels of smoking and non-smoking healthy sedentary individuals. *Clin Exp Health Sci* 2016; 6(2):51-5.
<https://doi.org/10.5152/clinexphealthsci.2016.036>
18. Li H, Zhao J, Liang J, Song X. Exploring causal effects of smoking and alcohol related lifestyle factors on self-report tiredness: A Mendelian randomization study. *PLoS One* 2023; 18(6):e0287027.
<https://doi.org/10.1371/journal.pone.0287027>
19. Chen H, Saad S, Sandow SL, Bertrand

- PP. Cigarette smoking and brain regulation of energy homeostasis. *Front Pharmacol* 2012; 3:147.
<https://doi.org/10.3389/fphar.2012.00147>
20. Corwin EJ, Klein LC, Rickelman K. Predictors of fatigue in healthy young adults: moderating effects of cigarette smoking and gender. *Biol Res Nurs* 2002; 3(4):222-33.
<https://doi.org/10.1177/109980040200300407>
21. Ferreira AI, Ferreira P, Cooper C, Oliveira D. How daily negative affect and emotional exhaustion correlates with work engagement and presenteeism-constrained productivity. *Int J Stress Manag* 2019; 26(3):261-71.
<https://doi.org/10.1037/str0000114>
22. Härmä M, Karhula K, Puttonen S, Ropponen A, Koskinen A, Ojajärvi A, et al. Shift work with and without night work as a risk factor for fatigue and changes in sleep length: A cohort study with linkage to records on daily working hours. *J Sleep Res* 2019; 28(3):e12658.
<https://doi.org/10.1111/jsr.12658>
23. Armstrong-Carter E, Fuligni AJ, Wu X, Gonzales N, Telzer EH. A 28-day, 2-year study reveals that adolescents are more fatigued and distressed on days with greater NO₂ and CO air pollution. *Sci Rep* 2022; 12(1):17015.
<https://doi.org/s41598-022-20602-z>
24. Rose JJ, Wang L, Xu Q, McTiernan CF, Shiva S, Tejero J, et al. Carbon monoxide poisoning: pathogenesis, management, and future directions of therapy. *Am J Respir Crit Care Med* 2017; 195(5):596-606.
<https://doi.org/10.1164/rccm.201606-1275CI>
25. Acharya SP, Purpura A, Kao L, House DR. Incidence and risk factors for carbon monoxide poisoning in an emergency department in Nepal. *Toxicol Commun* 2022; 6(1):13-9.
<https://doi.org/10.1080/24734306.2021.2010955>

Turnitin.libraryfk

PERBEDAAN PROFIL KOLESTEROL DARAH ANTARA KEHAMILAN PRE-EKLAMPSIA DAN KEHAMILAN NORMAL

Artikel Jurnal 1

Document Details

Submission ID

trn:oid::3618:127829152

Submission Date

Feb 9, 2026, 1:16 PM GMT+7

Download Date

Feb 9, 2026, 1:35 PM GMT+7

File Name

Preeklamsia dan kolesterol_Sri Indah_19012026.pdf

File Size

330.6 KB

12 Pages

3,329 Words

19,233 Characters

13% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

Filtered from the Report

- Bibliography
- Quoted Text
- Small Matches (less than 10 words)

Exclusions

- 18 Excluded Matches

Top Sources

- 13%  Internet sources
- 0%  Publications
- 0%  Submitted works (Student Papers)

Integrity Flags

0 Integrity Flags for Review

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Top Sources

13%  Internet sources
0%  Publications
0%  Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Internet	www.repository.trisakti.ac.id	5%
2	Internet	scholar.unand.ac.id	3%
3	Internet	e-journal.trisakti.ac.id	2%
4	Internet	core.ac.uk	2%
5	Internet	www.trijurnal.lemlit.trisakti.ac.id	<1%
6	Internet	conferences.unusa.ac.id	<1%
7	Internet	www.sicardiologia.it	<1%
8	Internet	medicinaudayana.ejournals.ca	<1%



PERBEDAAN PROFIL KOLESTEROL DARAH ANTARA KEHAMILAN PRE-EKLAMPSIA DAN KEHAMILAN NORMAL

Sri Indah Kemala¹, Tjam Diana Samara^{2*}

¹ Program Studi Sarjana Kedokteran Fakultas Kedokteran, Universitas Trisakti, Jakarta, 11440, Indonesia

² Departemen Anatomi, Fakultas Kedokteran Universitas Trisakti, Jakarta, 11440, Indonesia

*Penulis koresponden: dianasamara@trisakti.ac.id

ABSTRAK

Kolesterol dan lipid lainnya merupakan sesuatu yang dibutuhkan untuk pertumbuhan dan perkembangan janin, kolesterol juga berfungsi sebagai prekursor untuk sintesis progesteron. Metabolisme lipid yang abnormal mempunyai pengaruh yang signifikan terhadap patogenesis kehamilan pada preeklampsia, dimana kadar tinggi remnant kolesterol dan profil lipid aterogenik (tingginya kadar kolesterol total, trigliserida, LDL dan penurunan kadar HDL) merupakan faktor risiko untuk inisiasi dan perkembangan aterosklerosis yang pada akhirnya mengarah pada CVD yang di duga perannya sama pada preeklampsia, karena ditemukan lesi arteri berupa akumulasi sel busa (*lipid-laden macrophages*) di dinding pembuluh arteri spiralis dari plasenta yang diamati, dan biasanya keadaan ini di temukan pada kasus aterosklerosis akut. Tujuan kajian pustaka ini yaitu menganalisis beberapa penelitian mengenai perbedaan profil kolesterol darah antara kehamilan preeklampsia dan kehamilan normal. Dari enam artikel yang dipilih memiliki hasil yang bervariasi, dimana pada dua artikel didapatkan hasil didapatkan peningkatan kadar kolesterol total, trigliserida, VLDL, LDL dan penurunan kadar HDL pada preeklampsia dibanding dengan hamil normal, hasil ini berkaitan dengan risiko terjadinya aterosklerosis. Sedangkan empat penelitian lainnya didapatkan hasil yang sedikit berbeda pada beberapa kolesterol. Untuk itu masih perlu penelitian lebih lanjut mengenai perbedaan profil kolesterol darah antara kehamilan preeklampsia dan kehamilan normal terkait dengan risiko terjadinya preeklampsia.

Kata kunci: Kolesterol, Lipid, Preeklampsia, Kehamilan normal

SEJARAH ARTIKEL

Diterima
Juli 2025
Revisi
Agustus 2025
Disetujui
Desember 2025
Terbit online
Januari 2026

KATA KUNCI

- Kolesterol
- Lipid
- Pre-eklampsia
- Kehamilan normal

ABSTRACT

Cholesterol and other lipids are something needed for fetal growth and development, cholesterol also serves as a precursor for synthetic progesterone. Abnormal lipid metabolism has a significant impact on preeclampsia pregnancies, where a high cholesterol remnant and an atherogenic lipid profile

(high cholesterol, triglycerides, VLDL, LDL and falling HDL) is a risk factor for initial development and development of atherosclerosis, which is thought to be the same role in preeclampsia. Since lesions have been found in the arteries of an accumulation of foam cells (lipid-laden macrophages) on the walls of the spiral artery of the observed placenta, this is usually found in the case of acute atherosclerosis. The purpose of this literature review is to analyze several studies on differences in blood cholesterol profiles between preeclampsia and normal pregnancy. From the six articles selected had mixed results, where in the two articles, the results showed an increase in total cholesterol, triglyceride, VLDL, LDL levels and decreased HDL levels in preeclampsia compared to normal pregnancy, these results are associated with the risk of developing atherosclerosis. Meanwhile, four other studies found slightly different results on some cholesterol. This requires further study on the difference in blood cholesterol between preeclampsia pregnancy and normal pregnancy associated with the risk of preeclampsia.

Keywords: Cholesterol, Lipid, Preeclampsia, Normal Pregnancy

1. PENDAHULUAN

Preeklampsia merupakan penyebab utama dari morbiditas dan mortalitas pada ibu dan bayi diseluruh dunia, dan merupakan kelainan hipertensi yang biasanya terjadi setelah 20 minggu kehamilan.⁽¹⁾ Preeklampsia di definisikan sebagai tekanan darah ≥ 140 mmHg sistolik/diastolik ≥ 90 mmHg setelah 20 minggu kehamilan pada dua kali pengukuran selang 4 jam pada ibu yang sebelumnya normotensi dan proteinuria, atau bila tidak terdapat proteinuria, hipertensi onset baru dengan satu atau lebih keadaan berikut: insufisiensi renal, gangguan fungsi hati, gejala pada cerebral atau visual, trombositopenia, edema paru.⁽²⁾ Menurut Profil Kesehatan Indonesia Tahun 2014, salah satu akibat AKI di Indonesia yaitu hipertensi dalam kehamilan (25%).⁽³⁾ Di Indonesia sendiri frekuensi kejadian preeklampsia yaitu sekitar 3-10%.⁽⁴⁾

Sampai saat ini preeklampsia masih menjadi “Disease of Theory” dimana tidak terdapat kepastian tentang mekanisme terjadinya.⁽⁵⁾ Pada saat kehamilan adaptasi metabolisme lipid dibutuhkan untuk pertumbuhan dan perkembangan janin.⁽⁶⁾ Abnormalitas pada metabolisme lipid mempunyai pengaruh yang signifikan terhadap patogenesis kehamilan pada preeklampsia.⁽⁷⁾ Kadar tinggi dari remnant kolesterol dan profil lipid aterogenik (tingginya kadar kolesterol total, trigliserida, LDL-c dan penurunan kadar HDL-c) merupakan faktor risiko untuk inisiasi dan perkembangan aterosklerosis yang pada akhirnya mengarah pada CVD (Cardiovascular Disease).⁽⁸⁻¹²⁾

Studi prospektif yang dilakukan oleh Anuradha R dan Durga T menunjukkan terdapat peningkatan lipid pada serum pada preeklampsia dibandingkan dengan kehamilan normal, dimana terdapat peningkatan kolesterol, LDL, VLDL, TG pada preeklampsia ringan sampai berat.⁽¹³⁾ Penelitian Siringoringo et al didapatkan adanya perbedaan kadar HDL dan LDL pada preeklampsia dengan

kehamilan normal pada etnik Minangkabau, dengan rasio lebih tingginya perbandingan LDL/HDL pada preeklampsia daripada kehamilan normal, sedangkan pada TG dan juga kolesterol total tidak ada perbedaan pada preeklampsia dengan kehamilan normal. ⁽¹⁴⁾

Berdasarkan penelitian-penelitian tersebut terdapat perbedaan hasil profil kolesterol darah pada kadar kolesterol total dan trigliserida pada preeklampsia, oleh karena masih adanya perbedaan hasil penelitian tersebut, maka penulis tertarik untuk menulis karya tulis lebih lanjut mengenai perbedaan profil kolesterol darah antara kehamilan preeklampsia dan kehamilan normal.

Tinjauan pustaka ini disusun dengan tujuan untuk menurunkan kejadian preeklampsia dengan mengetahui perbedaan profil kolesterol darah antara kehamilan preeklampsia dan kehamilan normal. Diharapkan juga dapat memberikan informasi dan menambah pengetahuan tentang perbedaan profil kolesterol darah antara kehamilan preeklampsia dan kehamilan normal juga menjadi salah satu rujukan untuk penelitian-penelitian selanjutnya.

2. PEMBAHASAN

Adaptasi dari metabolisme lipid pada saat kehamilan merupakan sesuatu yang dibutuhkan untuk pertumbuhan dan perkembangan janin. ⁽⁶⁾ Selama kehamilan terjadi peningkatan yang bermakna konsentrasi lemak, lipoprotein, dan apolipoprotein. ⁽¹⁵⁾ Pada preeklampsia, metabolisme lipid yang abnormal mempunyai pengaruh yang signifikan terhadap patogenesis kehamilan pada preeklampsia. ⁽⁷⁾ Pada wanita dengan preeklampsia konsentrasi TC, LDL, VLDL & Trigliserida ditemukan meningkat secara signifikan dibandingkan dengan wanita dengan kehamilan normal & wanita yang tidak hamil, sedangkan HDL konsentrasinya menurun. ⁽¹⁶⁾

Hal ini didukung dengan penelitian yang dilakukan Yadav S, et al dimana hasilnya terdapat peningkatan yang signifikan dalam serum TC, TG, LDL-C, VLDL-C, dan penurunan HDL-C pada preeklampsia dibandingkan wanita hamil normal. ⁽¹⁷⁾ Sependapat dengan studi cross sectional yang dilakukan Mohammed GK, et al di dapatkan hasil peningkatan yang signifikan di semua parameter profil lipid diantara kedua kelompok dengan tingkat kolesterol, TGS, LDL dan VLDL lebih tinggi pada kelompok preeklampsia, namun tingkat HDL rendah pada kelompok preeklampsia disbanding kelompok kehamilan normotensif sebagai kontrol. ⁽¹⁸⁾

Berbeda dengan penelitian yang dilakukan oleh Anuradha R, Durga T menunjukkan terdapat peningkatan lipid serum kolesterol, LDL, VLDL, TG pada preeklampsia ringan sampai berat pada preeklampsia dibandingkan dengan kehamilan normal, sedangkan pada HDL secara signifikan tidak ada perbedaan antara kehamilan normal dan preeklampsia. ⁽¹³⁾

Perbedaan Profil Kolesterol Darah Antara Kehamilan Pre-Eklampsia dan Kehamilan Normal

Kemala, Samara

p-ISSN 0853-7720; e-ISSN 2541-4275, Volume 11, Nomor 1, halaman 33 – 44, Januari 2026

DOI: <https://doi.org/10.25105/pdk.v11i1.23750>

Studi lain yang dilakukan Siringoringo et al, hasilnya terdapat perbedaan kadar HDL dan LDL pada preeklampsia dengan kehamilan normal pada etnik Minangkabau, dengan rasio perbandingan LDL/HDL lebih tinggi pada preeklampsia daripada kehamilan normal, sedangkan pada kolesterol total dan TG pada preeklampsia dengan kehamilan normal tidak ada perbedaan.⁽¹⁴⁾ Hal ini kontras dengan penelitian yang dilakukan Lima et al didapatkan hasil nilai VLDL dan TG dari wanita dengan preeklampsia secara signifikan lebih tinggi daripada wanita sehat.⁽¹⁹⁾ Didapatkan hasil yang sama juga Kashinakunti et al., pada kelompok preeklampsia TG serum meningkat secara signifikan dibandingkan dengan wanita hamil normal. Parameter lain seperti TC, LDL-C, VLDL-C dan HDL-C tidak meningkat secara signifikan.⁽²⁰⁾

Perbedaan Profil Kolesterol Darah Antara Kehamilan Pre-Eklampsia dan Kehamilan Normal

Kemala, Samara

p-ISSN 0853-7720; e-ISSN 2541-4275, Volume 11, Nomor 1, halaman 33 – 44, Januari 2026

DOI: <https://doi.org/10.25105/pdk.v11i1.23750>

Hasil penelitian mengenai kolesterol darah kehamilan preeklampsia dan kehamilan normal

Tabel. 2 Ringkasan pustaka

Penelitian	Lokasi penelitian	Desain studi	Subjek penelitian	Variabel yang di teliti	Hasil
Yadav, et al. Lama penelitian : 2 tahun (oktober 2015- juli 2017)	Maharashtra, India	<i>prospective case–control comparative study</i>	100 wanita hamil normotensif (kelompok kontrol) dan 100 wanita pra-eklamptik (kelompok studi) dalam trimester ketiga	profil lipid, kehamilan normotensif, preeklampsia	terdapat peningkatan yang signifikan dalam serum TC, TG, LDL-C, VLDL-C, dan penurunan HDL-C pada kelompok studi (wanita preeklampsia) dibandingkan dengan kontrol (wanita hamil normal)
Kehamilan normal (kelompok kontrol)					
TC	TG	VLDL	LDL	HDL	
149.94(mg%)	151.61(mg%)	30.70(mg%)	59.92(mg%)	59.32(mg%)	
Preeklampsia (kelompok studi)					
TC	TG	VLDL	LDL	HDL	
221.92(mg%)	255.68(mg%)	51.19(mg%)	137.10(mg%)	32.93(mg%)	
p-value					
< 0.01	< 0.001	< 0.001	< 0.01	< 0.01	

Perbedaan Profil Kolesterol Darah Antara Kehamilan Pre-Eklampsia dan Kehamilan Normal

Kemala, Samara

p-ISSN 0853-7720; e-ISSN 2541-4275, Volume 11, Nomor 1, halaman 33 – 44, Januari 2026

DOI: <https://doi.org/10.25105/pdk.v11i1.23750>

Penelitian	Lokasi penelitian	Desain studi	Subjek penelitian	Variabel yang di teliti	Hasil
Mohammed GK, et al	Mesir	<i>cross-sectional comparative study</i>	70 wanita hamil dengan 35 wanita hamil normal sebagai kontrol dan 35 preeklampsia	profil lipid, kehamilan normal, preeklampsia	Terdapat perbedaan yang signifikan di semua parameter profil lipid diantara kedua kelompok dengan tingkat kolesterol, TGS, LDL dan VLDL lebih tinggi pada kelompok preeklampsia, namun tingkat HDL rendah pada kelompok preeklampsia dibanding kelompok kehamilan normotensif sebagai kontrol
Kehamilan normal (kelompok kontrol)					
TC	TG	VLDL	LDL	HDL	
135.82 mg/dl	122.59 mg/dl	24.52 mg/dl	68.35 mg/dl	43.14 mg/dl	
Preeklampsia (kelompok studi)					
TC	TG	VLDL	LDL	HDL	
153.73 mg/dl	141.42 mg/dl	28.28 mg/dl	102.22 mg/dl	23.23 mg/dl	
p-value					
<0.001*	0.029*	0.015*	<0.001*	<0.001*	

Perbedaan Profil Kolesterol Darah Antara Kehamilan Pre-Eklampsia dan Kehamilan Normal

Kemala, Samara

p-ISSN 0853-7720; e-ISSN 2541-4275, Volume 11, Nomor 1, halaman 33 – 44, Januari 2026

DOI: <https://doi.org/10.25105/pdk.v11i1.23750>

Penelitian	Lokasi penelitian	Desain studi	Subjek penelitian	Variabel yang di teliti	Hasil
Anuradha R, et al	India	prospective study	Sebanyak 80 kasus dipilih secara acak, 40 wanita dengan hamil normal dan 40 wanita dengan kehamilan preeklampsia	Serum lipid, preeklampsia, kehamilan normal	terdapat peningkatan lipid serum kolesterol, LDL, VLDL, TG pada preeklampsia ringan sampai berat pada preeklampsia dibandingkan dengan kehamilan normal, sedangkan pada HDL secara signifikan tidak ada perbedaan antara kehamilan normal dan preeklampsia.
Kehamilan normal (kelompok kontrol)					
TC	TG	VLDL	LDL	HDL	
136.85 mg/dl	126.85 mg/dl	24.7 mg/dl	72.75 mg/dl	39 mg/dl	
Preeklampsia (kelompok studi)					
TC	TG	VLDL	LDL	HDL	
160.9 mg/dl	210.32 mg/dl	41.95 mg/dl	84.5 mg/dl	38.57 mg/dl	
p-value					
0.0452	0.0001	0.0001	0.0163	0.2835	

Perbedaan Profil Kolesterol Darah Antara Kehamilan Pre-Eklampsia dan Kehamilan Normal

Kemala, Samara

p-ISSN 0853-7720; e-ISSN 2541-4275, Volume 11, Nomor 1, halaman 33 – 44, Januari 2026

DOI: <https://doi.org/10.25105/pdk.v11i1.23750>

Penelitian	Lokasi penelitian	Desain studi	Subjek penelitian	Variabel yang di teliti	Hasil
Siringoringo HE, et al Lama penelitian : Agustus-Desember 2014.	RSUP dr. M. Djamil Padang, RS Reksodiwiryo Padang dan UPTD Laboratorium Kesehatan Daerah Sumatera Barat, Indonesi	observasional dengan rancangan <i>cross sectional</i>	54 orang ibu hamil yang memenuhi kriteria inklusi. 27 orang ibu hamil dengan preeklampsia dan 27 orang ibu hamil dengan kehamilan normal.	serum lipid, kehamilan normal, preeklampsia	terdapat perbedaan kadar HDL dan LDL pada preeklampsia dengan kehamilan normal pada etnik Minangkabau, dengan rasio perbandingan LDL/HDL lebih tinggi pada preeklampsia daripada kehamilan normal, sedangkan pada kolesterol total dan trigliserida pada preeklampsia dengan kehamilan normal tidak ada perbedaan
Kehamilan normal (kelompok kontrol)					
TC	TG	LDL/HDL	LDL	HDL	
247,56 mg/dl	272,89 mg/dl	2,08 mg/dl	129,67 mg/dl	63,33 mg/dl	
Preeklampsia (kelompok studi)					
TC	TG	LDL/HDL	LDL	HDL	
270,19 mg/dl	296,07 mg/dl	3,86 mg/dl	159 mg/dl	51,93 mg/dl	
p-value					
0.158	0.504	0.005	0.023	0.012	
Penelitian	Lokasi penelitian	Desain studi	Subjek penelitian	Variabel yang di teliti	

Perbedaan Profil Kolesterol Darah Antara Kehamilan Pre-Eklampsia dan Kehamilan Normal

Kemala, Samara

p-ISSN 0853-7720; e-ISSN 2541-4275, Volume 11, Nomor 1, halaman 33 – 44, Januari 2026

DOI: <https://doi.org/10.25105/pdk.v11i1.23750>

Lima VJ et al.	Brazil	Cross-sectional analytical study	42 wanita preeklamsia dan 35 wanita hamil sehat pada trimester ketiga kehamilan sebagai kontrol	serum lipid, kehamilan normal, preeklamsia	nilai VLDL dan trigliserida dari wanita dengan preeklamsia secara signifikan lebih tinggi daripada wanita sehat.
Kehamilan normal (kelompok kontrol)					
TC	TG	VLDL	LDL	HDL	
229.0 mg/dl	216 mg/dl	43.3 mg/dl	133.0 mg/dl	53.0 mg/dl	
Preeklamsia (kelompok studi) Proteinuria (mg/dl over 24 hours)					
	0.3 - 0.9	1-2	>2	p-value	
TC	225.0 mg/dl	231.0 mg/dl	273.0 mg/dl	0.06	
TG	241 mg/dl	298 mg/dl	339 mg/dl	< 0.0001	
VLDL	48.3 mg/dl	59.7 mg/dl	67.8 mg/dl	< 0.0001	
LDL	121.8 mg/dl	123.0 mg/dl	150.2 mg/dl	0.30	
HDL	55.0 mg/dl	49.0 mg/dl	56.0 mg/dl	0.60	

Perbedaan Profil Kolesterol Darah Antara Kehamilan Pre-Eklampsia dan Kehamilan Normal

Kemala, Samara

p-ISSN 0853-7720; e-ISSN 2541-4275, Volume 11, Nomor 1, halaman 33 – 44, Januari 2026

DOI: <https://doi.org/10.25105/pdk.v11i1.23750>

Penelitian	Lokasi penelitian	Desain studi	Subjek penelitian	Variabel yang di teliti	Hasil
Kashinakunti SV., et al	India	Case control study	90 wanita preeklampsia dan 90 wanita hamil normal sebagai kontrol	profil lipid, wanita hamil normal, preeklampsia	pada kelompok preeklamsia trigliserida serum meningkat secara signifikan dibandingkan dengan wanita hamil normal. Parameter lain seperti TC, LDL-C, VLDL-C dan HDL-C tidak meningkat secara signifikan
Kehamilan normal (kelompok kontrol)					
TC	TG	VLDL	LDL	HDL	
203.1 mg/dl	187.7 mg/dl	37.47 mg/dl	113.97 mg/dl	41.43 mg/dl	
Preeklampsia (kelompok studi)					
TC	TG	VLDL	LDL	HDL	
193.37 mg/dl	214.86 mg/dl	40.33 mg/dl	111.43 mg/dl	43.2 mg/dl	
p-value					
>0.05	<0.02	>0.05	>0.05	>0.05	

Perbedaan Profil Kolesterol Darah Antara Kehamilan Pre-Eklampsia dan Kehamilan Normal

Kemala, Samara

p-ISSN 0853-7720; e-ISSN 2541-4275, Volume 11, Nomor 1, halaman 33 – 44, Januari 2026

DOI: <https://doi.org/10.25105/pdk.v11i1.23750>

3. KESIMPULAN

Pada beberapa penelitian yang membandingkan kadar kolesterol dan lipid lainnya pada wanita dengan kehamilan normal dan preeklampsia memiliki hasil yang bervariasi, dimana ada beberapa penelitian dengan hasil terjadi peningkatan TC, TG, LDL, VLDL tetapi HDL menurun pada preeklampsia dibanding dengan kelompok kontrol, dimana hasil penelitian ini mendukung peran dislipidemia pada terjadinya preeklampsia, namun ada juga penelitian yang mendapat hasil tidak ada perbedaan dari kadar HDL antara preeklampsia dan kehamilan normal. Penelitian lainnya mendapat hasil hanya terjadi peningkatan trigliserid pada preeklampsia dibandingkan dengan wanita hamil normal sedangkan TC, LDL-C, VLDL-C dan HDL-C tidak meningkat secara signifikan, akan tetapi ada juga yang mendapat hasil yang sedikit berbeda dimana ada TG dan VLDL meningkat pada preeklampsia dibandingkan dengan kehamilan normal. Selanjutnya terdapat juga hasil dimana hanya terdapat perbedaan kadar HDL dan LDL pada preeklampsia dibandingkan dengan wanita hamil normal sedangkan TC dan TG tidak ada perbedaan.

Oleh karena didapatkan hasil yang berbeda pada beberapa penelitian, maka penulis menyarankan perlu untuk dilakukan penelitian lebih lanjut lagi mengenai perbedaan profil kolesterol darah antara kehamilan preeklampsia dan kehamilan normal dengan jumlah responden yang lebih besar menggunakan metode cohort dengan disertakan juga umur dan IMT sebelum dan saat hamil mengingat dua hal ini berpengaruh terhadap serum lipid. Untuk di Indonesia sendiri bisa menyertakan secara ras mengingat di Indonesia terdiri dari banyak suku dan ras.

5. DAFTAR PUSTAKA

1. Pregnancy report of the American College of Obstetricians and Gynecologists' task force on. Hypertens Pregnancy Obs Gynecol. 2013;122:1122–31.09, 544.e541–544.e512, <https://doi.org/10.1016/j.ajog.2013.08.019> (2013).
2. American College of Obstetricians and Gynecologists and Task Force on Hypertension in Pregnancy, "Hypertension in pregnancy. Report of the American college of obstetricians and gynecologists' task force on hypertension in pregnancy," Obstetrics & Gynecology, vol. 122, no. 5, pp. 1122–1131, 2013.
3. Kementerian Kesehatan Republik Indonesia. 2015. Profil Kesehatan Indonesia Tahun 2014. Jakarta: Kemenkes RI.
4. SDKI. (2012). Survei Demografi Kesehatan Indonesia. Jakarta.
5. Perdana RA, Surya IGP, Sanjaya INH. Obesitas dan resistensi insulin merupakan factor risiko terjadinya preeklampsia. MEDICINA. 2019;50(3): 493-7.
6. Herrera E, Ortega-Senovilla H. Lipid metabolism during pregnancy and its implications for fetal growth. Curr Pharm Biotechnol 2014;15:24–31.

Perbedaan Profil Kolesterol Darah Antara Kehamilan Pre-Eklampsia dan Kehamilan Normal

Kemala, Samara

p-ISSN 0853-7720; e-ISSN 2541-4275, Volume 11, Nomor 1, halaman 33 – 44, Januari 2026

DOI: <https://doi.org/10.25105/pdk.v11i1.23750>

doi: 10.2174/1389201015666140330192345

7. Sayed A. Study of Lipid Peroxidation and Antioxidant Status in Preeclampsia. J Krishna Inst Med Scien Univ. 2013; 2:69-76.
8. Higashi Y, Sukhanov S, Anwar A, Shai SY, Delafontaine P. IGF-1, oxidative stress, and atheroprotection. Trends Endocrinol Metab 2010;21:245–54.
doi: 10.1016/j.tem.2009.12.005.
9. Gupta A, Baradaran H, Schweitzer AD, et al. Carotid plaque MRI and stroke risk: a systematic review and meta-analysis. Stroke 2013;44:3071–7.
doi: 10.1161/STROKEAHA.113.002551.
10. Jellinger PS, Handelsman Y, Rosenblit PD, et al. American Association of Clinical Endocrinologists and American College of Endocrinology guidelines for management of dyslipidemia and prevention of cardiovascular disease. Endocr Pract. 2017;23(Suppl 2):1-87.
11. Varbo A, Nordestgaard BG. Remnant cholesterol and risk of ischemic stroke in 112, 512 individuals from the general population. Ann Neurol. 2019;85:550–9.
doi: 10.1002/ana.25432.
12. Varbo A, Benn M, Tybjaerg-Hansen A, Jorgensen AB, Frikke-Schmidt R, Nordestgaard BG. Remnant cholesterol as a causal risk factor for ischemic heart disease. J Am Coll Cardiol. 2013;61:427–36
doi: 10.1016/j.jacc.2012.08.1026.
13. Anuradha R, Durga T. *Estimation of lipid profile among Preeclampsia women by comparing with normal pregnancy.* International Journal of Contemporary Medical Research. 2016;3:1958-61.
14. Siringoringo HE, Yusrawati, Chundrayetti E. Perbedaan Rerata Kadar Profil Lipid pada Preeklampsia dengan Kehamilan Normal pada Etnik Minangkabau. Jurnal Kesehatan Andalas. 2016; 5(1):30-5.
doi: <https://doi.org/10.25077/jka.v5i1.438>
15. Sabrida O, Hariadi, Yantri E. Hubungan Kadar LDL dan HDL Serum Ibu Hamil Aterm dengan Berat Lahir Bayi. Jurnal Kesehatan Andalas. 2014;3(3):289-96.
16. Gohil JT, Patel PK, Priyanka G. Estimation of Lipid Profile in Subjects of Preeclampsia. The Journal of Obstetrics and Gynecology of India. 2011;61(4):399-403.
doi: 10.1007/s13224-011-0057-0
17. Yadav S, Agrawal M, Hariharan C, et al. A Comparative Study of Serum Lipid Profile of Women with Preeclampsia and Normotensive Pregnancy. Journal of Datta Meghe Institute of Medical Sciences University. 2018;13(2):83-6.
18. Mohammed GK, Ayad WA, El-Gazzar UB, et al. Maternal Plasma Lipid Concentration in Third Trimester of Women with Preeclampsia and Normotensive Pregnancy. IJMA. 2020;2(2): 405-11.
doi: 10.21608/ijma.2020.23839.1101
19. Lima VJ, Andrade CR, Ruschi G, Sass N. Serum Lipid Levels in Pregnancies complicated by Preeklampsia. Sao Paulo Med. J 2011;129(2):73-6.
20. Kashinakunti S. V., Sunitha. H. et at. Lipid Profile in Preeclampsia. JCDR. 2010;(4): 2748-51.