



Editorial

The role of artificial intelligence in nutritional assessment in clinical practice

Food & Nutrition: Functional Food

- A scoping review: Potential nutraceutical values of bioactive compounds and antioxidant activity in durian seed – An exotic tropical plant

Clinical Nutrition: Nutrition & Metabolism

- Effect of omega-3 fatty acid supplementation on reduce body weight and body fat mass in obesity: A case study with an evidence-based approach
- Effect of magnesium supplementation on glycemic control in type 2 diabetes mellitus: evidence-based case report
- Relationship between protein intake and nutritional status with response therapy of intensive phase in pulmonary tuberculosis patients in Dili and Oecusse
- Leucine-included liquid diet as medical nutrition therapy improved handgrip strength and mid-upper circumference in clinically malnourished pulmonary tuberculosis patient with high neutrophil lymphocyte ratio: A case report from Universitas Indonesia hospital
- Evaluation of the prevalence, and relationship between nutritional status, malnutrition, and quality of life in cancer patients undergoing chemotherapy in Mashhad, Iran in 2023

Clinical Nutrition: Critical Care Nutrition

- Correlation of energy and protein intake with nitrogen balance changes in late acute phase critically ill patients

Community Nutrition: Nutrition Through Life Cycle

- Correlation between body mass index and 25(OH)D levels in pregnant women
- Socio-demographic, exercise, and anthropometric profiles of adults participating in fitness exercise programs in Vhembe district, Limpopo province, South Africa
- Association between mother's purchase intention of iron-fortified infant cereal and iron intake among children aged 6-23 months in Tangerang, Indonesia
- The implementation and challenges of breastfeeding counselling program in Majene district, West Sulawesi: A qualitative study among health workers
- Gender, education, eating patterns, and nutritional status among high school students in the tourist area of Bukittinggi, Indonesia
- A qualitative study on the experiences of the prospective brides about nutritional preparation during the preconception period
- Association between maternal anemia and low birth weight among stunting children 12–23 months in Dili, Timor Leste

World Nutrition Journal Editorial Office

Wisma Nugraha Building, Suite 510, 5th Floor

Jl. Raden Saleh No. 6 Jakarta Pusat

Website : <http://www.worldnutritionjournal.org/>

Phone: +622131905330 Email: worldnutritionjournal@gmail.com

Cover design by: Harryarts / Freepik

Aim and Scope

World Nutrition Journal (abbreviated: W Nutr J) is an international, English language, peer-reviewed, and open access journal upholding recent evidence related to nutrition sciences. The journal accepts manuscripts in terms of original paper, case report, editorial, and letter to editor.

About

World Nutrition Journal was founded in 2016 as the official journal of Indonesian Nutrition Association. It aims to publish high quality articles in the field of community, clinical, and critical care aspects of nutrition sciences

Peer Review Process

All manuscripts submitted will be screened by an editor whether they have the requirements and accordingly to author instructions. Then, the editor will transfer them to Editor-in-chief. Editor-in-chief will choose peer-reviewer for the manuscripts, according to their field of expertise, with help from associate editors and editorial board members. Peer-review process should be based on Committee on Publication Ethics (COPE) standards. This journal uses double-blind review. After the peer review process and the manuscripts are approved, they will be sent back to Editor-in-chief. Afterwards, Editor-in-chief will send the approved manuscripts back to the editor for final editing, layout editing, and proofreading. Finally, the approved and edited manuscripts will be published.

Publication Frequency

This journal is published bi-annually

Open Access Policy

World Nutrition Journal is an open access journal which allows readers to access the full text of all published articles without charge. Readers are free to read, download, distribute, print, search, or link the full texts of all articles in World Nutrition Journal

Advertising Policy

Editorial materials are not influenced by any advertisement. Advertisement will appear in the online or printed version depending on request. For all inquiries, please contact World Nutrition Journal editorial office at Wisma Nugraha, Suite 510, 5th Floor, Jl Raden Saleh No. 6 Jakarta Pusat; phone: +622131905330; email: worldnutritionjournal@gmail.com.

Copy Right Notice

Indonesian Nutrition Association as publisher reserves the right of first publication of all published materials. All statements in articles are the responsibility of the authors.

Subscription

Submission fee per manuscript are: for outside Indonesia USD 100, for Indonesia IDR 1,000,000.

Editorial Board

Editor-in-chief	Luciana B. Sutanto Indonesia
Associate Editors	Hamid Jan Bin Jan Mohamed Malaysia Joseph Varon USA
Editorial Board Members	Imelda Angeles-Agdeppa Philippines Marek Nalos Australia Yvan Vandenplas Belgium Abdolreza Norouzi Iran Ina. S. Timan Indonesia
Manuscript Editors and Proof Readers	Soemilah Sastroamidjojo Indonesia Pittara Pansawira Indonesia Juwalita Surapsari Indonesia Isa Rosalia Ruslim Indonesia
Website Developer	Dita Nur Oktavia Indonesia Mia Puspita Ratih Indonesia
Project Manager	Dian Novita Chandra Indonesia
Assistant Project Manager	Diana Sunardi Indonesia Mia Puspita Ratih Indonesia
Editorial Office	World Nutrition Journal Wisma Nugraha, Suite 501, 5th Floor Jl Raden Saleh No. 6 Jakarta Pusat Phone: +622131905330 Email: worldnutritionjournal@gmail.com
Publisher	Indonesian Nutrition Association Wisma Nugraha, Suite 510, 5th Floor Jl Raden Saleh No. 6 Jakarta Pusat Phone: +622131905330 Email: ina.nutri@yahoo.co.id

Table of Content**Page**

Volume 08, Issue 02, February 2025 | page 1 - 138, | eISSN: 2580-7013

Editorial

Pansawira P	The role of artificial intelligence in nutritional assessment in clinical practice	i
-------------	--	---

Food and Nutrition: Functional Food

Aisyah S Harioputro D R Nurwati I	A scoping review: Potential nutraceutical values of bioactive compounds and antioxidant activity in durian seed – An exotic tropical plant	1
---	--	---

Clinical Nutrition: Nutrition & Metabolism

Anissa L Lestari W	Effect of omega-3 fatty acid supplementation on reduce body weight and body fat mass in obesity: A case study with an evidence-based approach	11
G Apriyanti A J Sinaga W	Effect of magnesium supplementation on glycemic control in type 2 diabetes mellitus: evidence-based case report	19
Lafo M M Chandra D N Nurwidya F Sequeira E M D C	Relationship between protein intake and nutritional status with response therapy of intensive phase in pulmonary tuberculosis patients in Dili and Oecusse	28
Singal A M Afifa N N Wardhani W I Baskoro H	Leucine-included liquid diet as medical nutrition therapy improved handgrip strength and mid-upper circumference in clinically malnourished pulmonary tuberculosis patient with high neutrophil lymphocyte ratio: A case report from Universitas Indonesia hospital	38
Zeyghami A Moghadam H K Babaei A Bari A R Shahraki S H Shahidsales S Dahri M	Evaluation of the prevalence, and relationship between nutritional status, malnutrition, and quality of life in cancer patients undergoing chemotherapy in Mashhad, Iran in 2023	45

Clinical Nutrition : Critical Care Nutrition

Sari G P P Witjaksono F Sutanto K Singal A M	Correlation of energy and protein intake with nitrogen balance changes in late acute phase critically ill patients	55
---	---	-----------

Clinical Nutrition : Nutrition Through Life Cycle

Sudarma V Lim an P B Istriana E Mohamed H J B J M Mediana A	Correlation between body mass index and 25(OH)D levels in pregnant women	63
Ramashidzha RR Makuse SH Masoga S	Socio-demographic, exercise, and anthropometric profiles of adults participating in fitness exercise programs in Vhembe district, Limpopo province, South Africa	69
Sim atupang YR Chandra DN Khusun H Februhartanty J Iswarawanti D Sari OL	Association between mother's purchase intention of iron-fortified infant cereal and iron intake among children aged 6-23 months in Tangerang, Indonesia	78
Febriyanti D Februhartanty J Wiradnyani L A A	The implementation and challenges of breastfeeding counselling program in Majene district, West Sulawesi: A qualitative study among health workers	89
Taufiq a Z Maise BA Rahman Zam Z S Rizqillah F Alem NK Amril F A	Gender, education, eating patterns, and nutritional status among high school students in the tourist area of Bukittinggi, Indonesia	97
Emirza P Wiradnyani L A A Ermayani E	A qualitative study on the experiences of the prospective brides about nutritional preparation during the preconception period	109
Hornay F G J D C Chandra D N Lestari W	Association between maternal anemia and low birth weight among stunting children 12-23 months in Dili, Timor Leste	121



Correlation between body mass index and 25(OH)D levels in pregnant women

Verawati Sudarma^{1,2}, Patricia Budihartanti Liman^{1,2}, Erita Istriana³, Hamid Jan B Jan Mohamed⁴, Achmad Mediana⁵

1. Department of Nutrition, Faculty of Medicine, Universitas Trisakti
2. Nutrition Study Center, Faculty of Medicine, Universitas Trisakti
3. Department of Psychiatry, Faculty of Medicine, Universitas Trisakti
4. Universiti Sains Malaysia
5. AMS Clinic-Kemang Jakarta

Received 11 August 2024
Accepted 11 November 2024
Published 28 February 2025

Link to DOI:
[10.25220/WNJ.V08.i2.0008](https://doi.org/10.25220/WNJ.V08.i2.0008)

Citation: Sudarma V, Liman P B, Istriana E, Mohamed H J B J M, Mediana A. Correlation between body mass index and 25(OH)D levels in pregnant women. World Nutrition Journal. 2025 February 28;8(12): 63-68.



Copyright: © 2025 by the authors. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution [CC BY] license (<https://creativecommons.org/licenses/by/4.0/>).
<http://www.worldnutrjournal.org>

Abstract

Background: Vitamin D is a vitamin that has a positive effect on maternal and infant health. The mother's nutritional status affects vitamin D levels, where one of nutritional status is expressed in body mass index (BMI). It has been assumed that BMI has the potential to influence the mother's vitamin D levels.

Objective: This study analyzes the correlation between BMI and 25(OH)D levels in pregnant women.

Methods: This observational analytical study examines data from the medical records of pregnant women who underwent antenatal examinations at the AMS Clinic Kemang from January 2022 to December 2023. The data analyzed were maternal age, gestational age at the time of vitamin D examination, gestational age at delivery, 25(OH)D levels, and maternal BMI. The Pearson correlation or Spearman Rank test was used in bivariate analysis, with a significance level of $p < 0.05$.

Results: The mean maternal age was 33.45 ± 6.24 years, with the median gestational age at the time of vitamin D examination being 6.5 mg. The median gestational age at birth was 39 weeks, with vitamin D levels of 20.8 ± 7.8 , and a median BMI of 24.22 kg/m^2 . Based on the results of the Spearman statistical test analysis, there is a strong negative significant correlation between body mass index and levels of 25(OH)D in pregnant women ($r = -0.747$; $p = 0.008$).

Conclusion: Pregnant women's body mass index correlates negatively with levels 25(OH)D. Therefore, it is recommended that pregnant women maintain a BMI within the normal range.

Keywords: body mass index, vitamin D, pregnancy

Corresponding author:

Name : Verawati Sudarma

Affiliation: Department of Nutrition, Faculty of Medicine, Universitas Trisakti

Email : verasudarma@trisakti.ac.id

Introduction

Vitamin D is a vitamin that has recently received attention because of its effects on health. Vitamin D's primary role is to maintain calcium and phosphorus homeostasis. However, nearly all human cells and tissues contain vitamin D receptors, suggesting that this vitamin has extra-

skeletal effects, particularly on the immune and cardiovascular systems.¹ Many studies show a high prevalence of vitamin D deficiency globally, including in Indonesia.²

Several factors influence vitamin D levels, including sun exposure, ethnicity, altitude, season, use of sun protection, and body mass index (BMI).³ Serum concentrations of 25(OH)D can be used for both D2 and D3 to evaluate an individual's vitamin D nutritional status. Research reports show that 25–55% of pregnant women experience vitamin D deficiency.⁴ The fetus is dependent on the mother's 25(OH)D concentration level, which affects the transfer of calcium, phosphorus, hormones, and immune balance, all of which are critical for bone growth processes and fetoplacental integrity.⁵ Observational studies have found a link between low levels of 25(OH)D during pregnancy and more problems for the mother, such as pre-eclampsia, gestational diabetes⁶, bacterial vaginosis, osteomalacia⁷, and needing a primary cesarean section. Similarly, the fetus will also experience complications, such as impaired fetal growth, increased risk of allergies, premature birth, abortion, neonatal hypocalcaemia, and obesity over the long term.⁸

Although the number of studies is limited, observational studies show that obese pregnant women (BMI >30 kg/m²) have significantly lower vitamin D levels than non-obese pregnant women.⁹ Andersen et al.¹⁰ reported that every 5 kg increase in BMI/m² is associated with a decrease in vitamin D levels of 1.48 ng/mL in pregnant women. Karlsson et al.⁹ reported that despite consuming higher vitamin D levels in the diet and using the same vitamin D supplements, obese pregnant women had lower vitamin D levels than normal-weight pregnant women in the first trimester. Among women, the obese group exhibited decreased mean vitamin D levels compared to both the overweight and normal weight groups during winter and summer, frequently attaining vitamin D insufficiency status (<10 ng/mL).¹¹

Other studies revealed different and conflicting results. Study in West Sumatra indicating a higher prevalence of vitamin D deficiency in pregnant women with low pre-pregnancy weight.¹² Similarly, Shen et al.'s study in southeast China at

the first-trimester phase revealed a correlation between a higher pre-pregnancy BMI and higher 25(OH)D levels, a pattern that continues into the second and third trimesters.¹³ Some studies show there is no relationship between pre-pregnancy BMI and vitamin D deficiency, as seen in the research of Woon et al.¹⁴ The differences in results from these studies were caused, among other things, by different subject populations, differences in vitamin D calcification based on 25 (OH)D levels, and differences in sampling times, especially in the study area with four seasons.

Even though maintaining optimal levels of vitamin D in the body is important in theory, research has not shown conclusive results. Only a few publications in Indonesia describe vitamin D levels in pregnant women, specifically in West Sumatra¹¹. Jakarta is the province with the highest prevalence of obesity among women aged over 18, namely 38.8%.¹⁵ This can put women in the Jakarta area at risk of experiencing high pre-pregnancy BMIs and weight gain during pregnancy. This study aims to determine the correlation between BMI and 25(OH)D levels. The data can be used to recommend that pregnant women maintain BMI to improve vitamin D levels.

Methods

The Trisakti University Faculty of Medicine Ethics Committee no. 036/KER/FK/I/2024 granted ethical approval for the research. This research is an observational analytical study. Secondary data was collected from the medical records of pregnant women who had antenatal visits at AMS Kemang Clinic, South Jakarta, from January 2022 to December 2023. Inclusion criteria: medical records of pregnant women who underwent antenatal examinations from January 2022 to December 2023, maternal age between 21- and 35-years during pregnancy, data on body mass index before pregnancy in the medical record, and data on 25(OH)D examination results in medical records during pregnancy. Exclusion criteria: medical records of pregnant women with kidney disorders or hyperparathyroidism, taking atorvastatin and thiazide diuretics. Samples were selected based on consecutive sampling.

All data will be edited, coded, and entered into the computer. SPSS for Windows version 20.0 will be used for data analysis. Univariate analysis will be carried out to determine the distribution of all variables. The normality test will be carried out using the Kolmogorov-Sminov test with a p-value >0.05 as normally distributed data. For continuous data, it will be presented in mean ± standard deviation, and for categorical data, data will be presented in n (%) for those that are normally distributed and geometric data median (min, max) for those that are not normally distributed. For bivariate analysis, the Pearson correlation test will

be used if the data is normally distributed or the Spearman Rank test if the data is not normally distributed with a significance if p-value <0.05.

Results

This study obtained data from 155 respondents who met the inclusion criteria in accordance with the minimum sample size required. The characteristics of the subject can be seen in **Table 1**.

Table 1. Subject characteristics

Variables	Mean±SD /median (min, max)	95% CI
Mother's age (age)	33.45±6.24	29.27 – 37.95
Gestational age at Vitamin D examination (weeks)	6.5 (5, 22)	6.26 – 13.03
Gestational age at labor (weeks)	39 (37, 40)	38.09 – 39.18
25(OH)D level (mg/dL)	20.8±7.8	15.56 – 26.04
Body mass index (kg/m ²)	24.22 (20.22, 41.62)	21.9 – 30.4

The mean maternal age is 33.45±6.24 years, with the median gestational age at the time of vitamin D examination being 6.5 mg. Gestational age during pregnancy with a median of 39 weeks, vitamin D levels 20.8±7.8 with a median BMI of 24.22 kg/m².

Table 2 shows the correlation of variables with 25(OH)D levels. Based on the Spearman statistical test analysis results, a strong negative significant correlation exists between body mass index and 25(OH)D levels in pregnant women (r -0.747; p = 0.008).

Table 2. Correlation of independent variables with 25(OH)D levels in pregnant women

Variables	25(OH)D level (mg/dL)	
	r	p
Mother's age	0.132	0.652 ^p
Gestational age at Vitamin D examination	-0.163	0.579 ^s
Gestational age at labor	0.038	0.897 ^s
Body mass index	-0.747	0.008 ^{*s}

p, Pearson correlation test; s, Spearman correlation test; *p<0,05

Discussion

The mean age of mothers in this study was 33.45±6.24; this age was in the range of 20-35 years, which corresponds to the ideal pregnancy age. WHO recommends that women be between 20 and 35 years old when experiencing pregnancy and childbirth. Pregnant women who are less than 20 years old or older than 35 years old are particularly vulnerable because their uterus and pelvis have not

yet reached adult size, making pregnancy and childbirth more prone to complications, including miscarriage. As the uterus gets older (more than 35 years old), pregnancy complications will easily occur, and the incidence of chromosomal abnormalities will also increase.¹⁶

The gestational age at the time of vitamin D examination in this study was a median of 6.5 weeks, with a minimum gestational age of 5 weeks and a maximum of 22 weeks. According to the

literature, increasing gestational age will affect vitamin D levels, and early pregnancy is an essential window for preventing vitamin D deficiency in pregnancy, which affects perinatal outcomes, including preeclampsia and spontaneous premature birth.¹⁷ There are no recommendations for when is the best time to check vitamin D levels in pregnant women. However, maternal 25(OH)D levels progressively decrease during pregnancy due to the fetus' increased physiological needs.¹⁸ However, there is insufficient evidence to support the recommendation to screen all pregnant women for vitamin D deficiency.

The median gestational age is 39 mg; this shows that the gestational age at parturition is within normal limits, namely 37-40 weeks. These new gestational age categories were suggested by the American College of Obstetricians and Gynecologists and the Society for Maternal-Fetal Medicine in 2012. They are early term (37 0/7 weeks of gestation to 38 6/7 weeks of gestation), full term (39 0/7 weeks of gestation to 40 6/7 weeks of gestation), late-term (41 0/7 weeks of gestation to gestational age 41 6/7 weeks), and post-term (gestational age 42 0/7 weeks and beyond) to more accurately describe births that happen at or after 37 0/7 weeks of gestation.¹⁹

The 25(OH)D level is in an insufficiency condition, namely 20.8 ± 7.8 mg/dL. Several studies show that pregnant women worldwide have a high prevalence of vitamin D deficiency in each trimester of pregnancy.²⁰ Factors determining 25(OH)D concentrations in pregnant women include skin pigmentation, UV radiation, extensive use of skin coverings due to religious or cultural reasons, and, more significantly, social deprivation. Several studies link low 25(OH)D levels with an increased risk of preeclampsia (PE), gestational diabetes mellitus (GDM), indications for cesarean section, premature birth, low birth weight (LBW), and small for gestational age (SGA). Vitamin D status is also critical for the fetus in pregnant women. Early in pregnancy, 25(OH)D crosses the placenta from mother to fetus, and levels measured in cord blood at birth depend on maternal status, averaging 80% of maternal values.²¹ If the mother is deficient, the same happens to the fetus.²² The placenta and fetal

tissues express 1 α -hydroxylase, which produces bioactive vitamin D in the fetal circulation.

Late hypocalcemia and nutritional rickets are the classic effects of vitamin D deficiency during pregnancy and in the neonate. Some studies suggest that prenatal vitamin D status affects offspring's susceptibility to developing asthma later in life.²³ It may also contribute to destroying pancreatic beta cells due to its action on lymphocytes and type 1 helper cytokines.²⁴ Vitamin D deficit during maternal pregnancy may also be a risk factor. Early brain development, which plays a relevant role in neuronal differentiation and synaptic function, causes multiple sclerosis in adulthood. The Body Mass Index (BMI) during vitamin D examination showed a median of 24.22 kg/m², which indicates that the mother was already overweight at the start of pregnancy. Bivariate analysis showed that of the various variables, only BMI showed a significant, strong negative correlation with 25(OH)D levels. This is in accordance with the theory, which states that an individual's body fat can influence vitamin D levels. Research from Giacoia et al. shows that obesity (BMI >30 kg/m²) in pregnant women indicates a severe vitamin D deficit risk.²⁵ 25(OH)D levels <20 ng/mL are considered an indication of vitamin D deficiency, although the Endocrine Society and another expert group consider vitamin D insufficiency if the 25(OH)D level is 20–29 ng/mL and sufficient if the level is ≥ 30 ng/mL as sufficient; for non-classical measures, some authors suggest achieving levels >40 ng/mL.

Savard et al.'s research on 79 pregnant women showed that in the first and third trimesters, pregnant women with a pre-pregnancy BMI ≥ 30 kg/m² had lower total serum 25(OH)D concentrations than women with a pre-pregnancy BMI <25 kg/m².²⁶ Serum total 25(OH)D concentration was inversely related to pre-pregnancy BMI in all trimesters but more strongly in the first trimester. Vitamin D supplement intake is adequate in improving vitamin D status among pregnant women with a higher pre-pregnancy BMI.²⁶ In assessing vitamin D status, BMI is important because obesity and adiposity are associated with higher total serum 25(OH)D

concentrations, and because most women now enter pregnancy with a long-term BMI in the overweight or obese range. Obese adults also often require larger doses of vitamin D to achieve a serum total of 25(OH)D concentrations, similar to adults with a BMI in the recommended range.²⁷

From the results of this study, it can be concluded that the need to maintain normal 25(OH)D levels is useful for preventing pregnancy complications. There was also a significant, strong negative correlation between BMI and 25(OH)D levels. This demonstrates a strong correlation between pregnant women's higher BMI and lower 25(OH)D levels. This study has several limitations. This study utilized secondary data, which may have biases and inaccuracies on pre-pregnancy BMI. Secondly, the sample size of this study is limited, requiring bigger sample sizes in further studies. The evaluation of maternal vitamin D levels occurred primarily during the first trimester of pregnancy, rendering the fluctuations in vitamin D status during early gestation uncertain. These constraints may influence the applicability of the findings to different populations. Additional potential factors influencing vitamin D levels, including skin type, physical activity, season, and genetic background, necessitate more investigation.

Conclusion

The results of this study suggest that women entering pregnancy should maintain their nutritional status in normal conditions, characterized by a normal BMI, in order to maintain 25(OH)D levels. Optimal 25(OH)D levels lead to optimal pregnancy outcomes for both the mother and the baby. Future studies using randomized controlled trials with more subjects and more factors influencing vitamin D are strongly recommended.

Conflict of interest

The authors declared no conflict of interest regarding this article.

Acknowledgement

The authors thank the AMS Clinic Team for assisting and supporting this study throughout all aspects of our study.

Open Access

This article is distributed under the terms of the Creative Commons Attribution 4.0 International License

(<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

1. De La Guía-Galipienso F, Martínez-Ferran M, Vallecillo N, Lavie CJ, Sanchis-Gomar F, Pareja-Galeano H. Vitamin D and cardiovascular health. *Clin Nutr*. 2021 May;40(5):2946–57.
2. Sudama V, Halim L. High skeletal muscle mass is associated with increased serum 25(OH)D levels in elderly. *Universa Med*. 2017 Dec 14;36(3):236–42.
3. Van Schoor N, Lips P. Global Overview of Vitamin D Status. *Endocrinol Metab Clin North Am*. 2017 Dec;46(4):845–70.
4. Davies-Tuck M, Yim C, Knight M, Hodges R, Doery JCG, Wallace E. Vitamin D testing in pregnancy: Does one size fit all? *Aust N Z J Obstet Gynaecol*. 2015 Apr;55(2):149–55.
5. Olmos-Ortiz A, Avila E, Durand-Carbajal M, Díaz L. Regulation of Calcitriol Biosynthesis and Activity: Focus on Gestational Vitamin D Deficiency and Adverse Pregnancy Outcomes. *Nutrients*. 2015 Jan 9;7(1):443–80.
6. Zhang MX, Pan GT, Guo JF, Li BY, Qin LQ, Zhang ZL. Vitamin D Deficiency Increases the Risk of Gestational Diabetes Mellitus: A Meta-Analysis of Observational Studies. *Nutrients*. 2015 Oct 1;7(10):8366–75.
7. Aghajafari F, Nagulesapillai T, Ronksley PE, Tough SC, O'Beirne M, Rabi DM. Association between maternal serum 25-hydroxyvitamin D level and pregnancy and neonatal outcomes: systematic review and meta-analysis of observational studies. *BMJ*. 2013 Mar 26;346(mar26 4):f1169–f1169.
8. Feng H, Xun P, Pike K, Wills AK, Chawes BL, Bisgaard H, et al. In utero exposure to 25-hydroxyvitamin D and risk of childhood asthma, wheeze, and respiratory tract infections: A meta-analysis of birth cohort studies. *J Allergy Clin Immunol*. 2017 May;139(5):1508–17.

9. Karlsson T, Andersson L, Hussain A, Bosaeus M, Jansson N, Osmaneovic A, et al. Lower vitamin D status in obese compared with normal-weight women despite higher vitamin D intake in early pregnancy. *Clin Nutr*. 2015 Oct;34(5):892–8.
10. Andersen LB, Abrahamsen B, Dalgård C, Kyhl HB, Beck-Nielsen SS, Frost-Nielsen M, et al. Parity and tanned white skin as novel predictors of vitamin D status in early pregnancy: a population-based cohort study. *Clin Endocrinol (Oxf)*. 2013 Sep;79(3):333–41.
11. Delle Monache S, Di Fulvio P, Iannetti E, Valeni L, Capone L, Nespoli MG, et al. Body mass index represents a good predictor of vitamin D status in women independently from age. *Clin Nutr*. 2019 Apr;38(2):829–34.
12. Aji AS, Yenizel E, Desmawati null, Lipoeto NI. The association between lifestyle and maternal vitamin D during pregnancy in West Sumatra, Indonesia. *Asia Pac J Clin Nutr*. 2018;27(6):1286–93.
13. Shen Y, Pu L, Si S, Xin X, Mo M, Shao B, et al. Vitamin D nutrient status during pregnancy and its influencing factors. *Clin Nutr Edinb Scotl*. 2020 May;39(5):1432–9.
14. Woon FC, Chin YS, Ismail IH, Batterham M, Abdul Latiff AH, Gan WY, et al. Vitamin D deficiency during pregnancy and its associated factors among third trimester Malaysian pregnant women. Gebreselassie SG, editor. *PLOS ONE*. 2019 Jun 24;14(6):e0216439.
15. Kemenkes RI. *Survei Kesehatan Indonesia (SKI) 2023* [Internet]. Available from: <https://www.badankebijakan.kemkes.go.id/ski-2023-dalam-angka/>
16. Dereje T, Muluneh S, Kebebe B. Practice of child spacing and its associated factors among women of child bearing age (15 to 49 years) in Illubabor zone, South West Ethiopia. *Int J Nurs Midwifery*. 2017 Jul 31;9(7):102–8.
17. Kiely ME, Wagner CL, Roth DE. Vitamin D in pregnancy: Where we are and where we should go. *J Steroid Biochem Mol Biol*. 2020 Jul;201:105669.
18. Pérez-López FR, Pilz S, Chedraui P. Vitamin D supplementation during pregnancy: an overview. *Curr Opin Obstet Gynecol*. 2020 Oct;32(5):316–21.
19. The American College of Obstetricians and Gynecologists. Definition of Term Pregnancy [Internet]. Available from: <https://www.acog.org/clinical/clinical-guidance/committee-opinion/articles/2013/11/definition-of-term-pregnancy>
20. Karras S, Paschou SA, Kandaraki E, Anagnostis P, Annweiler C, Tarlatzis BC, et al. Hypovitaminosis D in pregnancy in the Mediterranean region: a systematic review. *Eur J Clin Nutr*. 2016 Sep;70(9):979–86.
21. Ortigosa Gómez S, García-Algar O, Mur Sierra A, Ferrer Costa R, Carrascosa Lezcano A, Yeste Fernández D. Concentraciones plasmáticas de 25-OH vitamina D y parathormona en sangre de cordón umbilical. *Rev Esp Salud Pública*. 2015 Feb;89(1):75–83.
22. Dawodu A, Tsang RC. Maternal Vitamin D Status: Effect on Milk Vitamin D Content and Vitamin D Status of Breastfeeding Infants. *Adv Nutr*. 2012 May 1;3(3):353–61.
23. Wolsk HM, Chawes BL, Litorjua AA, Hollis BW, Waage J, Stokholm J, et al. Prenatal vitamin D supplementation reduces risk of asthma/recurrent wheeze in early childhood: A combined analysis of two randomized controlled trials. Cameron DW, editor. *PLOS ONE*. 2017 Oct 27;12(10):e0186657.
24. Infante M, Ricordi C, Sanchez J, Clare-Salzler MJ, Padilla N, Fuenmayor V, et al. Influence of Vitamin D on Islet Autoimmunity and Beta-Cell Function in Type 1 Diabetes. *Nutrients*. 2019 Sep 11;11(9):2185.
25. Giacoia E, Costanzo P, Mansur J. Variación estacional de los niveles de vitamina D y su relación con la obesidad en una población de embarazadas de alto riesgo en Buenos Aires. *REV ARGENT ENDOCRINOL METAB*. 2019;56(4):27–32.
26. Savard C, Bielecki A, Plante AS, Lemieux S, Gagnon C, Weiler HA, et al. Longitudinal Assessment of Vitamin D Status across Trimesters of Pregnancy. *J Nutr*. 2021 Jul;151(7):1937–46.
27. Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, et al. Evaluation, Treatment, and Prevention of Vitamin D Deficiency: an Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab*. 2011 Jul;96(7):1911–30.



ORIGINAL PAPER

7
Correlation between body mass index and 25(OH)D levels in pregnant women

Verawati Sudarma^{1,2}, Patricia Budihartanti Liman^{1,2}, Erita Istriana³, Hamid Jan B Jan Mohamed⁴, Achmad Mediana⁵

1. Department of Nutrition, Faculty of Medicine, Universitas Trisakti
2. Nutrition Study Center, Faculty of Medicine, Universitas Trisakti
3. Department of Psychiatry, Faculty of Medicine, Universitas Trisakti
4. Universiti Sains Malaysia
5. AMS Clinic-Kemang Jakarta

Abstract

Background: Vitamin D is a vitamin that has a positive effect on maternal and infant health. The mother's nutritional status affects vitamin D levels, where one of nutritional status is expressed in body mass index (BMI). It has been assumed that BMI has the potential to influence the mother's vitamin D levels.

Objective: This study analyzes the correlation between BMI and 25(OH)D levels in pregnant women.

Methods: This observational analytical study examines data from the medical records of pregnant women who underwent antenatal examinations at the AMS Clinic Kemang from January 2022 to December 2023. The data analyzed were maternal age, gestational age at the time of vitamin D examination, gestational age at delivery, 25(OH)D levels, and maternal BMI. The Pearson correlation or Spearman Rank test was used in bivariate analysis, with a significance level of $p < 0.05$.

Results: The mean maternal age was 33.45 ± 6.24 years, with the median gestational age at the time of vitamin D examination being 6.5 mg. The median gestational age at birth was 39 weeks, with vitamin D levels of 20.8 ± 7.8 , and a median BMI of 24.22 kg/m^2 . Based on the results of the Spearman statistical test analysis, there is a strong negative significant correlation between body mass index and levels of 25(OH)D in pregnant women ($r = -0.747$; $p = 0.008$).

Conclusion: Pregnant women's body mass index correlates negatively with levels 25(OH)D. Therefore, it is recommended that pregnant women maintain a BMI within the normal range.

Keywords: body mass index, vitamin D, pregnancy

Received 11 August 2024
Accepted 11 November 2024
Published 28 February 2025

Link to DOI:
[10.25220/WNJ.V08.I2.0008](https://doi.org/10.25220/WNJ.V08.I2.0008)

Citation: Sudarma V, Liman P B, Istriana E, Mohamed H J B J M, Mediana A. Correlation between body mass index and 25(OH)D levels in pregnant women. World Nutrition Journal. 2025 February 28;8(12): 63-68.



Copyright: © 2025 by the authors. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).
<http://www.worldnutrjournal.org>

Corresponding author:
Name : Verawati Sudarma
Affiliation: Department of Nutrition, Faculty of Medicine, Universitas Trisakti
Email : verasudarma@trisakti.ac.id

Introduction

Vitamin D is a vitamin that has recently received attention because of its effects on health. Vitamin D's primary role is to maintain calcium and phosphorus homeostasis. However, nearly all human cells and tissues contain vitamin D receptors, suggesting that this vitamin has extra-

skeletal effects, particularly on the immune and cardiovascular systems.¹ Many studies show a high prevalence of vitamin D deficiency globally, including in Indonesia.²

Several factors influence vitamin D levels, including sun exposure, ethnicity, altitude, season, use of sun protection, and body mass index (BMI).³ Serum concentrations of 25(OH)D can be used for both D2 and D3 to evaluate an individual's vitamin D nutritional status. Research reports show that 25–55% of pregnant women experience vitamin D deficiency.⁴ The fetus is dependent on the mother's 25(OH)D concentration level, which affects the transfer of calcium, phosphorus, hormones, and immune balance, all of which are critical for bone growth processes and fetoplacental integrity.⁵ Observational studies have found a link between low levels of 25(OH)D during pregnancy and more problems for the mother, such as pre-eclampsia, gestational diabetes,⁶ bacterial vaginosis, osteomalacia⁷, and needing a primary cesarean section. Similarly, the fetus will also experience complications, such as impaired fetal growth, increased risk of allergies, premature birth, abortion, neonatal hypocalcaemia, and obesity over the long term.⁸

Although the number of studies is limited, observational studies show that obese pregnant women (BMI >30 kg/m²) have significantly lower vitamin D levels than non-obese pregnant women.⁹ Andersen et al.¹⁰ reported that every 5 kg increase in BMI/m² is associated with a decrease in vitamin D levels of 1.48 ng/mL in pregnant women. Karlsson et al.⁹ reported that despite consuming higher vitamin D levels in the diet and taking the same vitamin D supplements, obese pregnant women had lower vitamin D levels than normal-weight pregnant women in the first trimester. Among women, the obese group exhibited decreased mean vitamin D levels compared to both the overweight and normal weight groups during winter and summer, frequently attaining vitamin D insufficiency status (<10 ng/mL).¹¹

Other studies revealed different and conflicting results. Study in West Sumatra indicating a higher prevalence of vitamin D deficiency in pregnant women with low pre-pregnancy weight.¹² Similarly, Shen et al.'s study in southeast China at

the first-trimester phase revealed a correlation between a higher pre-pregnancy BMI and higher 25(OH)D levels, a pattern that continues into the second and third trimesters.¹³ Some studies show there is no relationship between pre-pregnancy BMI and vitamin D deficiency, as seen in the research of Woon et al.¹⁴ The differences in results from these studies were caused, among other things, by different subject populations, differences in vitamin D calcification based on 25 (OH)D levels, and differences in sampling times, especially in the study area with four seasons.¹⁵

Even though maintaining optimal levels of vitamin D in the body is important in theory, research has not shown conclusive results. Only a few publications in Indonesia describe vitamin D levels in pregnant women, specifically in West Sumatra¹¹. Jakarta is the province with the highest prevalence of obesity among women aged over 18, namely 38.8%.¹⁵ This can put women in the Jakarta area at risk of experiencing high pre-pregnancy BMIs and weight gain during pregnancy. This study aims to determine the correlation between BMI and 25(OH)D levels. The data can be used to recommend that pregnant women maintain BMI to improve vitamin D levels.

Methods

The Trisakti University Faculty of Medicine Ethics Committee no. 036/KER/FK/1/2024 granted ethical approval for the research. This research is an observational analytical study. Secondary data was collected from the medical records of pregnant women who had antenatal visits at AMS Kemang Clinic, South Jakarta, from January 2022 to December 2023. Inclusion criteria: medical records of pregnant women who underwent antenatal examinations from January 2022 to December 2023, maternal age between 21- and 35-years during pregnancy, data on body mass index before pregnancy in the medical record, and data on 25(OH)D examination results in medical records during pregnancy. Exclusion criteria: medical records of pregnant women with kidney disorders or hyperparathyroidism, taking atorvastatin and thiazide diuretics. Samples were selected based on consecutive sampling.

All data will be edited, coded, and entered into the computer. SPSS for Windows version 20.0 will be used for data analysis. Univariate analysis will be carried out to determine the distribution of all variables. The normality test will be carried out using the Kolmogorov-Sminov test with a p-value >0.05 as normally distributed data. For continuous data, it will be presented in mean ± standard deviation, and for categorical data, data will be presented in n (%) for those that are normally distributed and geometric data median (min, max) for those that are not normally distributed. For bivariate analysis, the Pearson correlation test will

be used if the data is normally distributed or the Spearman Rank test if the data is not normally distributed with a significance if p-value <0.05.

Results

This study obtained data from 155 respondents who met the inclusion criteria in accordance with the minimum sample size required. The characteristics of the subject can be seen in **Table 1**.

Table 1. Subject characteristics

Variables	Mean±SD /median (min, max)	95% CI
Mother's age (age)	33.45±6.24	29.27 – 37.95
Gestational age at Vitamin D examination (weeks)	6.5 (5, 22)	6.26 – 13.03
Gestational age at labor (weeks)	39 (37, 40)	38.09 – 39.18
25(OH)D level (mg/dL)	20.8±7.8	15.56 – 26.04
Body mass index (kg/m ²)	24.22 (20.22, 41.62)	21.9 – 30.4

The mean maternal age is 33.45±6.24 years, with the median gestational age at the time of vitamin D examination being 6.5 mg. Gestational age during pregnancy with a median of 39 weeks, vitamin D levels 20.8±7.8 with a median BMI of 24.22 kg/m².

Table 2 shows the correlation of variables with 25(OH)D levels. Based on the Spearman statistical test analysis results, a strong negative significant correlation exists between body mass index and 25(OH)D levels in pregnant women (r -0.747; p = 0.008).

Table 2. Correlation of independent variables with 25(OH)D levels in pregnant women

Variables	25(OH)D level (mg/dL)	
	r	p
Mother's age	0.132	0.652 ^p
Gestational age at Vitamin D examination	-0.163	0.579 ^s
Gestational age at labor	0.038	0.897 ^s
Body mass index	-0.747	0.008 ^{qs}

p, Pearson correlation test; s, Spearman correlation test; *p<0.05

Discussion

The mean age of mothers in this study was 33.45±6.24; this age was in the range of 20-35 years, which corresponds to the ideal pregnancy age. WHO recommends that women be between 20 and 35 years old when experiencing pregnancy and childbirth. Pregnant women who are less than 20 years old or older than 35 years old are particularly vulnerable because their uterus and pelvis have not

yet reached adult size, making pregnancy and childbirth more prone to complications, including miscarriage. As the uterus gets older (more than 35 years old), pregnancy complications will easily occur, and the incidence of chromosomal abnormalities will also increase.¹⁶

The gestational age at the time of vitamin D examination in this study was a median of 6.5 weeks, with a minimum gestational age of 5 weeks and a maximum of 22 weeks. According to the

literature, increasing gestational age will affect vitamin D levels, and early pregnancy is an essential window for preventing vitamin D deficiency in pregnancy, which affects perinatal outcomes, including preeclampsia and spontaneous premature birth.¹⁷ There are no recommendations for when is the best time to check vitamin D levels in pregnant women. However, maternal 25(OH)D levels progressively decrease during pregnancy due to the fetus' increased physiological needs.¹⁸ However, there is insufficient evidence to support the recommendation to screen all pregnant women for vitamin D deficiency.

The median gestational age is 39 mg; this shows that the gestational age at parturition is within normal limits, namely 37-40 weeks. The new gestational age categories were suggested by the American College of Obstetricians and Gynecologists and the Society for Maternal-Fetal Medicine in 2012. They are early term (37 0/7 weeks of gestation to 38 6/7 weeks of gestation), full term (39 0/7 weeks of gestation to 40 6/7 weeks of gestation), late-term (41 0/7 weeks of gestation to gestational age 41 6/7 weeks), and post-term (gestational age 42 0/7 weeks and beyond) to more accurately describe births that happen at or after 37 0/7 weeks of gestation.¹⁹

The 25(OH)D level is in an insufficiency condition, namely 20.8 ± 7.8 mg/dL. Several studies show that pregnant women worldwide have a high prevalence of vitamin D deficiency in each trimester of pregnancy.²⁰ Factors determining 25(OH)D concentrations in pregnant women include skin pigmentation, UV radiation, extensive use of skin coverings due to religious or cultural reasons, and, more significantly, social deprivation. Several studies link low 25(OH)D levels with an increased risk of preeclampsia (PE), gestational diabetes mellitus (GDM), indications for cesarean section, premature birth, low birth weight (LBW), and small for gestational age (SGA). Vitamin D status is also critical for the fetus in pregnant women. Early in pregnancy, 25(OH)D crosses the placenta from mother to fetus, and levels measured in cord blood at birth depend on maternal status, averaging 80% of maternal values.²¹ If the mother is deficient, the same happens to the fetus.²² The placenta and fetal

tissues express 1 α -hydroxylase, which produces bioactive vitamin D in the fetal circulation.¹ Late hypocalcemia and nutritional rickets are the classic effects of vitamin D deficiency during pregnancy and in the neonate. Some studies suggest that prenatal vitamin D status affects offspring's susceptibility to developing asthma later in life.²³ It may also contribute to destroying pancreatic beta cells due to its action on lymphocytes and type 1 helper cytokines.²⁴ Vitamin D deficit during maternal pregnancy may also be a risk factor. Early brain development, which plays a relevant role in neuronal differentiation and synaptic function, causes multiple sclerosis in adulthood. The Body Mass Index (BMI) during vitamin D examination showed a median of 24.22 kg/m², which indicates that the mother was already overweight at the start of pregnancy. Bivariate analysis showed that of the various variables, only BMI showed a significant, strong negative correlation with 25(OH)D levels. This is in accordance with the theory, which states that an individual's body fat can influence vitamin D levels. Research from Giacoia et al. shows that obesity (BMI >30 kg/m²) in pregnant women indicates a severe vitamin D deficit risk.²⁵ 25(OH)D levels <20 ng/mL are considered an indication of vitamin D deficiency, although the Endocrine Society and another expert group consider vitamin D insufficiency if the 25(OH)D level is 20–29 ng/mL and sufficient if the level is ≥ 30 ng/mL as sufficient; for non-classical measures, some authors suggest achieving levels >40 ng/mL.

Savard et al.'s research on 79 pregnant women showed that in the first and third trimesters, pregnant women with a pre-pregnancy BMI ≥ 30 kg/m² had lower total serum 25(OH)D concentrations than women with a pre-pregnancy BMI <25 kg/m².²⁶ Serum total 25(OH)D concentration was inversely related to pre-pregnancy BMI in all trimesters but more strongly in the first trimester. Vitamin D supplement intake is adequate in improving vitamin D status among pregnant women with a higher pre-pregnancy BMI.²⁶ In assessing vitamin D status, BMI is important because obesity and adiposity are associated with higher total serum 25(OH)D

concentrations, and because most women now enter pregnancy with a long-term BMI in the overweight or obese range. Obese adults also often require larger doses of vitamin D to achieve a serum total of 25(OH)D concentrations, similar to adults with a BMI in the recommended range.²⁷

From the results of this study, it can be concluded that the need to maintain normal 25(OH)D levels is useful for preventing pregnancy complications. There was also a significant, strong negative correlation between BMI and 25(OH)D levels. This demonstrates a strong correlation between pregnant women's higher BMI and lower 25(OH)D levels. This study has several limitations. This study utilized secondary data, which may have biases and inaccuracies on pre-pregnancy BMI. Secondly, the sample size of this study is limited, requiring bigger sample sizes in further studies. The evaluation of maternal vitamin D levels occurred primarily during the first trimester of pregnancy, rendering the fluctuations in vitamin D status during early gestation uncertain. These constraints may influence the applicability of the findings to different populations. Additional potential factors influencing vitamin D levels, including skin type, physical activity, season, and genetic background, necessitate more investigation.

Conclusion

The results of this study suggest that women entering pregnancy should maintain their nutritional status in normal conditions, characterized by a normal BMI, in order to maintain 25(OH)D levels. Optimal 25(OH)D levels lead to optimal pregnancy outcomes for both the mother and the baby. Future studies using randomized controlled trials with more subjects and more factors influencing vitamin D are strongly recommended.

Conflict of interest

The authors declared no conflict of interest regarding this article.

Acknowledgement

The authors thank the AMS Clinic Team for assisting and supporting this study throughout all aspects of our study.

Open Access

This article is distributed under the terms of the Creative Commons Attribution 4.0 International Licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

1. De La Guía-Galipienso F, Martínez-Ferran M, Vallecillo N, Lavie CJ, Sanchis-Gomar F, Pareja-Galeano H. Vitamin D and cardiovascular health. *Clin Nutr*. 2021 May;40(5):2946–57.
2. Sudarma V, Halim L. High skeletal muscle mass is associated with increased serum 25(OH)D levels in elderly. *Universa Med*. 2017 Dec 14;36(3):236–42.
3. Van Schoor N, Lips P. Global Overview of Vitamin D Status. *Endocrinol Metab Clin North Am*. 2017 Dec;46(4):845–70.
4. Davies-Tuck M, Yim C, Knight M, Hodges R, Doery JCG, Wallace E. Vitamin D testing in pregnancy: Does one size fit all? *Aust N Z J Obstet Gynaecol*. 2015 Apr;55(2):149–55.
5. Olmos-Ortiz A, Avila E, Durand-Carbajal M, Díaz L. Regulation of Calcitriol Biosynthesis and Activity: Focus on Gestational Vitamin D Deficiency and Adverse Pregnancy Outcomes. *Nutrients*. 2015 Jan 9;7(1):443–80.
6. Zhang MX, Pan GT, Guo JF, Li BY, Qin LQ, Zhang ZL. Vitamin D Deficiency Increases the Risk of Gestational Diabetes Mellitus: A Meta-Analysis of Observational Studies. *Nutrients*. 2015 Oct 1;7(10):8366–75.
7. Aghajafari F, Nagulesapillai T, Ronksley PE, Tough SC, O'Beirne M, Rabi DM. Association between maternal serum 25-hydroxyvitamin D level and pregnancy and neonatal outcomes: systematic review and meta-analysis of observational studies. *BMJ*. 2013 Mar 26;346(mar26 4):f1169–f1169.
8. Feng H, Xun P, Pike K, Wills AK, Chawes BL, Bisgaard H, et al. In utero exposure to 25-hydroxyvitamin D and risk of childhood asthma, wheeze, and respiratory tract infections: A meta-analysis of birth cohort studies. *J Allergy Clin Immunol*. 2017 May;139(5):1508–17.

9. Karlsson T, Andersson L, Hussain A, Bosaeus M, Jansson N, Osancevic A, et al. Lower vitamin D status in obese compared with normal-weight women despite higher vitamin D intake in early pregnancy. *Clin Nutr*. 2015 Oct;34(5):892–8.
10. Andersen LB, Abrahamsen B, Dalgård C, Kyhl HB, Beck-Nielsen SS, Frost-Nielsen M, et al. Parity and tanned white skin as novel predictors of vitamin D status in early pregnancy: a population-based cohort study. *Clin Endocrinol (Oxf)*. 2013 Sep;79(3):333–41.
11. Delle Monache S, Di Fulvio P, Iannetti E, Valerii L, Capone L, Nespoli MG, et al. Body mass index represents a good predictor of vitamin D status in women independently from age. *Clin Nutr*. 2019 Apr;38(2):829–34.
12. Aji AS, Yerizel E, Desmawati null, Lipoeto NI. The association between lifestyle and maternal vitamin D during pregnancy in West Sumatra, Indonesia. *Asia Pac J Clin Nutr*. 2018;27(6):1286–93.
13. Shen Y, Pu L, Si S, Xin X, Mo M, Shao B, et al. Vitamin D nutrient status during pregnancy and its influencing factors. *Clin Nutr Edinb Scotl*. 2020 May;39(5):1432–9.
14. Woon FC, Chin YS, Ismail IH, Batterham M, Abdul Latiff AH, Gan WY, et al. Vitamin D deficiency during pregnancy and its associated factors among third trimester Malaysian pregnant women. *Gebreselassie SG, editor. PLOS ONE*. 2019 Jun 24;14(6):e0216439.
15. Kemenkes RI. *Survei Kesehatan Indonesia (SKI) 2023* [Internet]. Available from: <https://www.badankebijakan.kemkes.go.id/ski-2023-dalam-angka/>
16. Dereje T, Muluneh S, Kebebe B. Practice of child spacing and its associated factors among women of child bearing age (15 to 49 years) in Illubabor zone, South West Ethiopia. *Int J Nurs Midwifery*. 2017 Jul 31;9(7):102–8.
17. Kiely ME, Wagner CL, Roth DE. Vitamin D in pregnancy: Where we are and where we should go. *J Steroid Biochem Mol Biol*. 2020 Jul;201:105669.
18. Pérez-López FR, Pilz S, Chedraui P. Vitamin D supplementation during pregnancy: an overview. *Curr Opin Obstet Gynecol*. 2020 Oct;32(5):316–21.
19. The American College of Obstetricians and Gynecologists. Definition of Term Pregnancy [Internet]. Available from: <https://www.acog.org/clinical/clinical-guidance/committee-opinion/articles/2013/11/definition-of-term-pregnancy>
20. Karras S, Paschou SA, Kandaraki E, Anagnostis P, Annweiler C, Tarlatzis BC, et al. Hypovitaminosis D in pregnancy in the Mediterranean region: a systematic review. *Eur J Clin Nutr*. 2016 Sep;70(9):979–86.
21. Ortigosa Gómez S, García-Algar O, Mur Sierra A, Ferrer Costa R, Carrascosa Lezcano A, Yeste Fernández D. Concentraciones plasmáticas de 25-OH vitamina D y parathormona en sangre de cordón umbilical. *Rev Esp Salud Pública*. 2015 Feb;89(1):75–83.
22. Dawodu A, Tsang RC. Maternal Vitamin D Status: Effect on Milk Vitamin D Content and Vitamin D Status of Breastfeeding Infants. *Adv Nutr*. 2012 May 1;3(3):353–61.
23. Wolsk HM, Chawes BL, Litonjua AA, Hollis BW, Waage J, Stokholm J, et al. Prenatal vitamin D supplementation reduces risk of asthma/recurrent wheeze in early childhood: A combined analysis of two randomized controlled trials. *Cameron DW, editor. PLOS ONE*. 2017 Oct 27;12(10):e0186657.
24. Infante M, Ricordi C, Sanchez J, Clare-Salzler MJ, Padilla N, Fuenmayor V, et al. Influence of Vitamin D on Islet Autoimmunity and Beta-Cell Function in Type 1 Diabetes. *Nutrients*. 2019 Sep 11;11(9):2185.
25. Giacoia E, Costanzo P, Mansur J. Variación estacional de los niveles de vitamina D y su relación con la obesidad en una población de embarazadas de alto riesgo en Buenos Aires. *REV ARGENT ENDOCRINOL METAB*. 2019;56(4):27–32.
26. Savard C, Bielecki A, Plante AS, Lemieux S, Gagnon C, Weiler HA, et al. Longitudinal Assessment of Vitamin D Status across Trimesters of Pregnancy. *J Nutr*. 2021 Jul;151(7):1937–46.
27. Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, et al. Evaluation, Treatment, and Prevention of Vitamin D Deficiency: an Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab*. 2011 Jul;96(7):1911–30.

ORIGINALITY REPORT

28% SIMILARITY INDEX	27% INTERNET SOURCES	30% PUBLICATIONS	9% STUDENT PAPERS
--------------------------------	--------------------------------	----------------------------	-----------------------------

PRIMARY SOURCES

1	pmc.ncbi.nlm.nih.gov Internet Source	5%
2	worldnutrijournal.org Internet Source	5%
3	hdl.handle.net Internet Source	3%
4	pure.ulster.ac.uk Internet Source	2%
5	www.lsmuni.it Internet Source	2%
6	bmcpregnancychildbirth.biomedcentral.com Internet Source	2%
7	Choi, Rihwa, Seonwoo Kim, Heejin Yoo, Yoon Cho, Sun Kim, Jae Chung, Soo-young Oh, and Soo-Youn Lee. "High Prevalence of Vitamin D Deficiency in Pregnant Korean Women: The First Trimester and the Winter Season as Risk Factors for Vitamin D Deficiency", <i>Nutrients</i> , 2015. Publication	1%
8	Ulfi Rahma Yunita, Helda Khusun, Fiastuti Witjaksono. "The association between knowledge and attitude towards nutrition fact panels (NFP) with sugar intake of the Indonesian adolescents", <i>World Nutrition Journal</i> , 2024 Publication	1%
9	ijn.zotarellifilhoscientificworks.com Internet Source	1%

10	ro.uow.edu.au Internet Source	1%
11	Submitted to Manchester Metropolitan University Student Paper	1%
12	journals.lww.com Internet Source	1%
13	maljnutr.org.my Internet Source	1%
14	oaji.net Internet Source	1%
15	www.ncbi.nlm.nih.gov Internet Source	1%
16	Hung N. Winn, Frank A. Chervenak, Roberto Romero. "Clinical Maternal-Fetal Medicine ", CRC Press, 2021 Publication	1%
17	Isaac Blickstein, Louis G. Keith. "Multiple Pregnancy - Epidemiology, Gestation, and Perinatal Outcome", CRC Press, 2019 Publication	1%
18	health.gov.au Internet Source	1%

Exclude quotes On

Exclude matches < 15 words

Exclude bibliography On