







MULTIDISCIPLINARY SCIENCE JOURNAL



 **Malque**
Publishing

August



Editorial Team

Editors-in-Chief



Dr. Maiko Dantas

*Federal Rural University of
Semi-Arid, Brazil*
maiko.dantas@malque.pub



Dr. Fatemeh Mollaamin

Kastamonu University, Turkey
smollaamin@gmail.com



Associate Editors



Dr. Thibério Castelo

*Federal Rural University of
Semi-Arid, Brazil*
thiberio.castelo@malque.pub



Dr. Hitesh Panchal

*Government Engineering
College Patan, India*
panchal_hitesh1@gtu.edu.in



Editorial Board Members

Dr. Ala Ali Nemer Amourah

Sohar University, Oman
Email: aamourah@su.edu.om



Dr. Ahmed Obaid

University of Kufa, Iraq
Email: ahmedj.aljanaby@uokufa.edu.iq



Dr. Alejandro Regalado

Universidad del Mar, Mexico
Email: alejandro.regalado33@gmail.com



Dr. Aman B. Upaganlawar

*SNJB's Shriman Suresh Dada
Jain College of Pharmacy,
India*
Email: amanrxy@gmail.com



Dr. Anam Azam

*Fraunhofer Institute for Systems
and Innovation Research,
Germany*
Email: anamrafiq95@gmail.com



Dr. Anand Joshi

Parul University, India
Email: anandyjoshi@gmail.com



Dr. Andee Dzulkarnaen Zakaria

*Universiti Sains Malaysia,
Malaysia*
Email: andee778@gmail.com



Dr. Anis Ahmad Chaudhary

*Al-Imam Muhammad Ibn Saud
Islamic University, Riyadh, Saudi
Arabia*
Email: anis.chaudhary@gmail.com



Dr. Anjanna Matta

*ICFAI Foundation for Higher
Education, India*
Email: anjireddyitm@gmail.com



Dr. Atif Amin Baig

*Management & Science
University, Malaysia*
Email: atif_amin@msu.edu.my



Dr. Ayo Olasupo

*KU School of Engineering, United
States*
Email: ayoolasupo@ku.edu



Dr. Bahtiar Mohamad

*Universiti Utara Malaysia,
Malaysia*
Email: mabahtiar@uum.edu.my



**Dr. Carlos Alberto Acosta-
Olivo**

*Universidad Autónoma de
Nuevo León, Mexico*
Email:
dr.carlosacosta@gmail.com



Dr. Ágnes Csiszárík-Kocsir

*Obuda University, Budapest,
Hungary*
Email: kocsir.agnes@kgk.uni-obuda.hu



Dr. Denys Miroshnichenko

*National Technical University
Kharkiv Polytechnic Institute,
Ukraine*
Email: dvmir79@gmail.com



Dr. Farooz Ahmad Lone

*Sher-e-Kashmir University of
Agricultural Sciences and
Technology of Kashmir, India*
Email:
dr.farooz462@gmail.com



Dr. Jafar Alasad Alshraideh

The University of Jordan, Jordan
Email: jalasad@ju.edu.jo



Dr. João Queiroz

*Federal University of Roraima,
Brazil*
Email:
jp.fernandes@yahoo.com.br



Dr. K. Karthick

GMR Institute of Technology,

Dr. Kehinde Obayomi

Victoria University Melbourne,

Dr. Khongdet Phasinam

Pibulsongkram Rajabhat

India

Email: karthick.k@gmrit.edu.in



Australia

Email:

obayomikehindeshola@gmail.com



University, Thailand

Email: phasinam@psru.ac.th



**Dr. Leonel Avendaño
Reyes**

*Universidad Autónoma de
Baja California, Mexico*

Email: lar62@uabc.edu.mx



Dr. Lukman Raimi

*Universiti Brunei Darussalam,
Brunei*

Email: luq_man2001@yahoo.com



Dr. Manish Kumar Dubey

Chandigarh University, India

Email: mkmkdubey@gmail.com



Dr. Md. Fahad Jubayer

*Sylhet Agricultural University,
Bangladesh*

Email:

fahadbau21@hotmail.com



**Dr. Mohamed Nor Azhari
Azman**

*Universiti Pendidikan Sultan Idris,
Malaysia*

Email: mnazhari@ftv.upsi.edu.my



Dr. Musa Danladi Ibrahim

Kogi State University, Nigeria

Email: dimusa55@gmail.com



Dr. Nítalo Machado

*Federal University of
Maranhao, Brazil*

Email: nitalo.farias@ufma.br



Dr. Palanivel Velmurugan

*Bharath Institute of Higher
Education and Research, India*

Email:

palanivelmurugan2008gmail.com



**Dr. Potula Sree
Brahmanandam**

*Shri Vishnu Engineering College
for Women, India*

Email:

dranandpotula@svecw.edu.in



Dr. Pranav Mehta

*Dharmsinh Desai University,
India*

Email:

pranavmehta.mech@ddu.ac.in



Dr. Rakesh Kumar Ameta

*Gandhinagar University,
Gandhinagar, India*

Email:

ametarakesh40@gmail.com



**Dr. Ram Chandra
Chaurasia**

*Indian Institute of Technology,
India*

Email: rcc.0761@gmail.com



Dr. Ramona Birau

Craiova University, Romania

Email:

ramona.f.birau@gmail.com



Dr. Rohail Hassan

*Universiti Utara Malaysia,
Malaysia*

Email:

rohail.hassan@uum.edu.my



Dr. Rohit Sahu

*GL Bajaj Institute of Technology
and Management, India*

Email: rohitsahu87@gmail.com



Dr. Roman Mylostyvyi

*Dnipro State Agrarian and
Economic University, Ukraine*

Email:

mylostyvyi.r.v@dsau.dp.ua



Dr. Romanus Osabohien

*Universiti Tenaga Nasional,
Malaysia*

Email: romik247@gmail.com



Dr. Ruchi Juyal

*Himalayan Institute of Medical
Sciences, India*

Email: ruchijuyal@srhu.edu.in



Dr. Stanly Tebug

*International Committee of
the Red Cross, South Sudan*

Dr. Vinod Joshi

Sharda University, India
Email: vinodjoshidmrc@gmail.com

Dr. Wunhong Su

Hangzhou Dianzi University,

Email:

stanlytebug@yahoo.com



China

Email: whsu@hdu.edu.cn



Most Read Articles

Acceptability of sweet potato (<i>Ipomoea batatas</i>) flour and toasted rice (<i>Oryza sativa</i>) powder as base ingredients in making polvoron	
Refardo et al.	663
Peculiarities of the impact of stress on physical and psychological health	
Ovsiannikova et al.	583
Emotional attachment to artificial intelligence and perceived social isolation among university students: An application of Sternberg's triangular theory of love	
Tran et al.	533
The influence of co-branding, service quality, lifestyle, and brand image to customer's loyalty: A survey of global coffee brand in Indonesia	
Sahabu et al.	422
Learning skills for the future – implementing the 21st-century learning	
Avdiu et al.	421

Publish with Malque Publishing and Advance your Academic Career

[Publisher home](#)

[About us](#)

[Register](#)

[Login](#)

Our Journals

[Applied Veterinary Research](#)

[Humanities Journal](#)

[Journal of Animal Behaviour and Biometeorology](#)

[Multidisciplinary Science Journal](#)

[Multidisciplinary Reviews](#)

[Malque Health](#)

[NexusSDGs](#)

[NexusTech](#)

Editorial Office

Malque Publishing, Edifício T&D - R. Juvenal Lamartine, 894 - Sala 402 - Centro, Mossoró - RN, CEP: 59618-740, Brazil.

multiscience@malque.pub



 [_malquepub](#)

 [_malquepub](#)

 [_malquepub](#)

 [_Whatsapp](#)

Exclusive OJS theme plugin developed with ♥ by



Vol. 7 Issue 8 (2025)



Artificial intelligence and data privacy in international trade law

Alazzam F A F, Aldrou K. K. A. R.

Research Article | The rapid advancement of artificial intelligence (AI) technologies has highlighted the critical importance of data privacy within the framework of international trade law. This study aims to...



Published in 06 February 2025

In Progress

Editor-in-chief
Dr. Maiko Dantas



Determining factors of lung tuberculosis among children in community health centers: A cross-sectional study

Machrumnizar M, Yuliana Y, et al.



Research Article | Tuberculosis is a serious global health issue and is particularly prevalent among children. Early and correct diagnosis is very important for effective management, but tuberculosis diagnosis in...



Published in 06 February 2025

Teachers' performance and experiences: an explanatory mixed method design in individual performance commitment and review form compliance

Parcon B S, Alum J G, et al.



Research Article | This explanatory mixed-method study examines the extent of teachers' performance, the challenges they face, and their strategies for compliance with the Individual

Performance Commitment and...



Published in 05 February 2025

Communication of women's empowerment through MSMEs as a CSR program in the cement subsector industry.




Jannah A, Mani L.

Research Article | This study aims to explain the CSR (Corporate Social Responsibility) strategy of cement subsector companies in Central Java, Indonesia, in contributing to the Sustainable Development Goal (SGD)...



Published in 05 February 2025

Students' critical thinking in numeracy problem-solving through moderate self-Efficacy: A mixed-methods study.



Yustitia V, Kusmaharti D, et al.

Research Article | The urgency of improving c in solving numeracy problems in elementary school students underlines the importance of this study. This study aims to determine the effect of self-efficacy on...




Published in 04 February 2025

Marital quality of disaster survivor farming families in cianjur district, Indonesia

Kumalasari B, Sunarti E, et al.

Research Article | Life at the micro level, particularly within families, is disrupted by earthquakes. Farming families face additional losses due to the disaster's impact, requiring substantial time and resources...



Published in 03 February 2025

Social-emotional development in adolescents: The role of peer interaction, social media engagement, and self-esteem

Rachman Y A, Sumarwan U, et al.

Research Article | Adolescence is closely associated with social-emotional development, which is characterized by an increase in emotions and the complexity of social relationships. This study is aimed at...



Published in 03 February 2025

iDDCS: Cervical intraepithelial neoplasia severity detection during vagina birth using artificial intelligence approach

Kapruwan A, Sharma S, et al.

Research Article | To diagnose cervical cancer, this paper uses patient Pap tests to look for abnormal tissue growth in the cervix region of pregnant women. By undergoing routine tests to identify any precancers...



Published in 02 February 2025

Maternal knowledge of Cushuro (*Nostoc* sp.) and prevalence of iron deficiency anemia: A study at the “Liberación Social” Health Center, Trujillo - 2024

Solansh N D C, Yosalin M V Y, et al.

Research Article | The study aimed to assess the level of knowledge about Cushuro and the prevalence of iron deficiency anemia at the Social Liberation Health Center in Trujillo in 2024. Utilizing a...



Published in 02 February 2025

Advancing innovation through university-industry collaboration: Empirical evidence from a developed economy

Hajrizi B, Shaqiri A B, et al.

Research Article | This study investigates the impact of university-industry collaboration (UIC) and job mobility on researchers' performance in Germany. It examines the temporal effects of collaboration,...



Published in 02 February 2025

Disinformation's Influence on Public Perception and Defense Policy Implementation in Indonesia

Sarjito A, Lelyana N.

Research Article | Disinformation poses a growing and complex challenge to public perception and policy implementation, especially in sensitive sectors like national defense. In Indonesia, where defense policies...




Published in 01 February 2025

Effectiveness of ethnoscience oriented project to improve students performance

Khery Y, Hakim A, et al.

Research Article | This study endeavors to harness this invaluable traditional wisdom and leverage it for pedagogical purposes by integrating the ethnoscience of the sasak tribe into science education,...




Published in 08 February 2025

Factors influencing the intention to use artificial intelligence for online advertising on social networks

Giang N T P, Duy N B P, et al.

Research Article | The rapid advancement of artificial intelligence (AI) has significantly transformed various industries, including online advertising on social networks. Businesses are increasingly investing in...



The effect of CG on tax avoidance: The moderating role of company size

Chen C Y, Nor-Ahmad S N H J N, et al.

Research Article | This paper focuses on the impact of different corporate governance characteristics, such as board size, board independence, board gender diversity, audit committee effectiveness and...



Published in 07 February 2025

Design and implementation of outdoor home smart farming based on raspberry.pi for home assistant management


Maslan M N, Ramli I R, et al.

Research Article | Outdoor home smart farming, also known as urban farming, is the practice of cultivating fruits, herbs, or vegetables for personal consumption on a small scale within a residential area. It is...



Published in 08 February 2025

Harmonizing heritage: A two-decade scoping bibliometric analysis of music in cultural tourism



Mazlan C A N, Abdullah M H, et al.

Research Article | This study provides a scoping bibliometric analysis of research on the integration of music within cultural tourism from 2004 to 2024. Analyzing 12 key publications from the Scopus database, the...



PDF

Published in 08 February 2025

A postmenopausal woman with generalized osteoporosis and risk for fracture: A case report

Stena R, Tahiraj D, et al.

Case Report/Clinical Report | Osteoporosis is one of the most common metabolic bone diseases. Patient X, a 67-year-old female, was presented for the first time in 2017 to a specialist doctor at the Polyclinic of Specialties...



PDF

Published in 04 February 2025

Novel ayurveda management of central serous retinopathy. (c.s.r.): an explorative case report

Mali N, Nibrad C, et al.

Case Report/Clinical Report | CSR is a retinal disorder that presents with the formation of fluid at the subretinal level, indeed bringing about vision impairment. It has some relation to hormonal and fluid imbalances due to...



Published in 07 February 2025

**Publish with
Malque Publishing
and Advance your
Academic Career**

[Publisher home](#)

Our Journals

[Applied Veterinary Research](#)

[Humanities Journal](#)

[Journal of Animal Behaviour and
Biometeorology](#)

[Multidisciplinary Science Journal](#)

Editorial Office

Malque Publishing, Edifício T&D -
R. Juvenal Lamartine, 894 - Sala
402 - Centro, Mossoró - RN, CEP:
59618-740, Brazil.

multiscience@malque.pub



 [malquepub](#)

 [malquepub](#)

[About us](#)

[Register](#)

[Login](#)

[Multidisciplinary Reviews](#)

[Research in Alternative Animal](#)

[Feeds](#)

[Wild Animals](#)

 [_malquepub](#)

 [Whatsapp](#)

Exclusive OJS theme plugin developed with ♥ by 



Determining factors of lung tuberculosis among children in community health centers: A cross-sectional study

Machrumnizar Machrumnizar^{ab}   | Yuliana Yuliana^b  | Adang Bachtiar^c 

^aDoctoral Program, Faculty of Public Health, Universitas Indonesia, Indonesia.

^bDepartment of Parasitology, Faculty of Medicine, Universitas Trisakti, Indonesia.

^cDepartment of Health Policy and Administration, Faculty of Public Health, Universitas Indonesia, Indonesia.

Abstract Tuberculosis is a serious global health issue and is particularly prevalent among children. Early and correct diagnosis is very important for effective management, but tuberculosis diagnosis in children is often difficult because the clinical symptoms are nonspecific and difficult to collect specimens. This study aims to determine risk factors based on sociodemographic, socioeconomic, nutrition and clinical status associated with tuberculosis. This study is a cross-sectional study. Children aged 0 to 14 years who came to the community health center with one or more clinical symptoms related to TB from January to December 2023 were studied. Research participants underwent two diagnostic methods, clinical and bacteriological. Positive Clinical diagnosis was found to be 72.4%. Meanwhile, bacteriologically confirmed participants were 10.3%, the remainder were unable to produce sputum. Most respondents were men (62.8%), with 66% of respondents aged ≥ 5 years, and those sociodemographic factors have a significant relationship with tuberculosis in children ($p < 0.05$). This study found that 64.7% of respondents had a BMI < 18.5 and significantly related to tuberculosis in children ($p < 0.05$). Furthermore, tuberculosis incidence and malnutrition also have a significant relationship ($p < 0.05$). Contact with active tuberculosis cases shows a significant difference in the proportion of tuberculosis and nontuberculosis cases ($p = 0.009$), while 57.1% of children were exposed to cigarette smoke ($p < 0.05$). Government health insurance (BPJS) was used by 70.5% of respondents who were tested positive for tuberculosis, and it is statistically significant ($p < 0.05$). Other socioeconomic factors were also significant ($p < 0.05$) for tuberculosis cases in children whose parents had low education at 85.9%; have 96.8% of jobs with a household income equal to or more than the provincial minimum wage (UMP) of 60.3%. In the future, these factors can be utilized to construct prediction models for clinical diagnosis in children suspected of tuberculosis, especially machine learning-based models, in addition to tuberculosis-related clinical symptoms.

Keywords: pediatrics, exposure to tuberculosis, tuberculosis, risk factor

1. Introduction

Among the leading sources of morbidity and death in children aged 0–14 years are tuberculosis (TB), which accounts for approximately 12% of the total 10.6 million TB cases in the world in 2023 and contributes to 3–25% of the total TB cases (World Health Organization, 2023). Children might represent up to 20% of the global burden of TB at endemic locations; hence, the real situation can vary from forecasts given adequate reporting systems and appropriate diagnoses (Richens et al., 2020). Nevertheless, there is a fairly modest rate of TB-identified cases and notifications, particularly in developing nations with limited resources, such as Indonesia; the estimated TB incidence of children aged 0–14 in 2022 is 110,881 children, or approximately 15.3% of all TB cases in Indonesia (Ministry of Health of the Republic of Indonesia, 2023). According to a survey by the United Nations International Children's Emergency Fund (UNICEF), 35% of instances of TB among children between the ages of 0 and 14 are unreported, meaning that these 110,881 children constitute only 65% of the total child TB cases in 2022 (UNICEF Indonesia, 2022). These data indicate that the level of community transmission is relatively high.

Bacteriological testing is the standard diagnostic method for adult TB but has low sensitivity and specificity when used in children (Basile et al., 2022). Since the prevalence of TB in children serves as an immediate indicator and surrogate for TB spread in the community, there is a dearth of novel, easily used, and inexpensive diagnostic methods for TB, especially in remote areas with limited laboratory and X-ray facilities, which adds to existing problems (UNICEF Indonesia, 2022). The diagnosis of tuberculosis in children aged 0–14 years without specific symptoms is a major challenge for doctors, academics, and program managers because it can hinder diagnosis and lead to worse treatment outcomes (Siddalingaiah et al., 2023). Current diagnostic methods have limited sensitivity, especially in children aged 0–14 years (Namuganga et al., 2021). The

scoring system, an alternative for diagnosing TB in children, has not yet been fully implemented according to guidelines (Ministry of Health of the Republic of Indonesia, 2023; World Health Organization, 2022).

In addition to clinical and biological variables, several socioeconomic and environmental variables increase the likelihood of TB in children between the ages of 0 and 14. These risk factors are very useful for predicting the development of TB disease. Therefore, identifying the risk factors for TB in children is crucial for making a diagnosis and initiating treatment as soon as possible for the best outcomes. Community health centers in Indonesia are the main place where TB patients seek treatment, so they are the starting point for TB case care, which includes early diagnosis, treatment, and reporting. The purpose of this study was to assess the sociodemographic, socioeconomic, and clinical risk factors for TB in children aged 0 to 14 years who exhibited one or more TB-related clinical symptoms.

2. Materials and methods

2.1. Data collection

This research was conducted at a community health center in the administrative area of South Jakarta, Indonesia. The respondents were children aged 0–14 years who came with a cough to receive treatment at community health centers between January and December 2023.

2.2. Research design

We used a cross-sectional design and studied children with normal nutritional status or malnutrition, including all genders, aged 0–14 years, who presented with a cough to receive treatment at community health centers between January and December 2023. The number of respondents was determined via consecutive sampling, and the sample size was determined via the total sampling method. All respondents who met the inclusion criteria were included in the study. TB diagnosis is categorized on the basis of clinical diagnosis (accompanied by X-rays and/or tuberculin tests) and bacteriology. The clinical diagnosis was determined by the occurrence or absence of symptoms of a persistent cough lasting more than 2 weeks, weight depletion or no weight gain in the previous two months, fever ≥ 2 weeks, and malaise ≥ 2 weeks. Sociodemographic factors (age, sex, age group, weight, height, nutritional status) were recorded during the medical history. Contact with active TB cases, history of BCG immunization, and exposure to cigarette smoke/air pollution were categorized dichotomously. Socioeconomic factors (education, employment, parental income, and health insurance) were also categorized dichotomously. This study also recorded the presence or absence of comorbid factors (HIV and DM). A child's nutritional status is determined by body mass index (BMI), which is calculated as the ratio of body weight in kilograms to height in meters squared. Furthermore, the BMI value of children aged 0–60 months must be compared with the WHO 2005 standard BMI value (WHO, 2006) and BMI values for children and adolescents (5–19 years old) in comparison to the WHO/NCHS 2007 reference (WHO, 2007).

2.3. Data analysis

All the data that had been collected and entered into standard Microsoft Excel were then analyzed via SPSS for Windows (SPSS Inc. version 25.0, Chicago, IL). The variables were initially examined univariately, and then a chi-square statistic was used to evaluate categorical variables to determine risk factors in children with tuberculosis. The McNemar statistic was used to estimate the proportions of risk variables between TB patients and non-TB patients. A statistically significant probability is smaller than 0.05. The odds ratio (OR) and 95% confidence interval (CI) were computed to ascertain the strength of the link.

3. Results and Discussion

The purpose of this cross-sectional study was to identify the predictive variables for both TB and non-TB diagnoses among children aged 0–14 years in Community Health Center facilities in Indonesia. TB risk factors such as HIV and diabetes were not found in the child respondents, so no analysis was carried out.

In this study, 113 out of 156 respondents were diagnosed with pulmonary TB from January to December 2023. Among the TB patients who received treatment, 72.4% of the respondents were clinically diagnosed with pulmonary TB. Bacteriologically confirmed TB cases were found in 10.3% of the respondents, and 89.7% of the respondents were unable to produce phlegm. Most of the respondents (62.8%) were boys, with 34% being less than five years old and 66% being older than five years (Table 1). Negative smear test results reportedly contribute to 13–20% of transmission and are, on average, 20–25% more infectious than positive test results (Asadi et al., 2022). Previous studies reported that most TB cases were diagnosed on the basis of clinical criteria (50–58%) (Chilyabanyama et al., 2024; Zürcher et al., 2019). Similarly, a previous cohort study reported that 16% (58/372), 95% CI, and 12–19% of the cases were smear-negative (Chilyabanyama et al., 2024), indicating that 84% of the TB cases were smear positive. However, another study reported that 65.2% of TB patients had negative BTA results and that 34.8% had positive BTA results (Campos et al., 2016).

Table 1 Characteristics related to sociodemographic factors, TB contact, BCG immunization, nutritional status, and exposure to smoking/air pollution between TB and non-TB respondents.

Characteristics	Frequency		TB	Non-TB	p Value	Odd Ratio (OR)	95% CI
	N, mean	%, SD					
Bacteriological diagnosis							
Yes	16	10.3	16	0	0.009*	-	-
No	140	89.7	97	43			
Age (year)							
<5 years	53	34	38	15	0.000¶	0.964	0.594 – 1.564
≥5 years	103	66	75	28		1.019	0.790 – 1.315
Gender							
Boy	98	62.8	72	26	0.000¶	1.054	0.797 – 1.393
Girl	58	37.2	41	17		0.918	0.589 – 1.429
Weight (kg)	19.6218	8.94918			0.000*	-	18.21 – 21.04
Height (cm)	104.3910	26.23235			0.000*	-	100.24 – 108.54
Body Mass Index (BMI)							
<18.5	101	64.7	91	10	0.000*	3.463	1.997 – 6.005
>18.4	55	35.3	22	33		0.254	0.168 – 0.382
Contacts with active TB patients							
Yes	85	54.5	62	23	0.002¶	1.026	0.741 – 1.420
None	71	45.5	51	20		0.970	0.664 – 1.418
BCG immunization							
Yes	114	73.1	85	29	0.328*	1.115	0.883 – 1.408
No	42	26.9	28	14		0.761	0.445 – 1.302
nutritional status							
Malnutrition	100	64.1	90	10	0.000*	3.425	1.974 – 5.941
Normal	56	35.9	23	33		0.265	0.178 – 0.396
Smoke exposure/air pollution							
Yes	89	57.1	65	24	0.006¶	1.031	0.756 – 1.405
None	67	42.9	48	19		0.961	0.645 – 1.432

*Chi-square test, p value <0.05; ¶McNemar test, p value <0.05.

TB disease is generally determined on the basis of clinical diagnosis and is accepted when bacteriological evidence is unavailable (Abdullahi et al., 2021). As is often reported, most respondents were boys. Various studies have reported similar findings, with TB cases occurring more often in boys (Alao et al., 2020; Campos et al., 2016; Chidambaram et al., 2021) and variations in the age range of children who contract TB. Younger children are believed to be more affected than older children are (Hamid et al., 2019; Ogbudebe et al., 2018; Tilahun & Gebre-Selassie, 2016; Vukugah et al., 2022) due to household contact with active TB patients and immature immunity. However, exposure to active TB can also occur in the community. Boys tend toward physical activity and high socialization, so they enjoy playing outside the house. This causes boys to potentially encounter adult active TB and be exposed to cigarette smoke or air pollution (Martínez-Andrés et al., 2017; Miller et al., 2021).

Malnutrition and tuberculosis (TB) are reciprocally associated: malnutrition increases the likelihood of having active TB by six to ten times, and TB contributes to malnutrition in patients (Feleke et al., 2019; Musuenge et al., 2020). Patients with active TB tend to be underweight and have a BMI less than 18.5 kg/m², which is considered an index of malnutrition. Thus, TB incidence is strongly correlated with poor nutrition and low BMI (Musuenge et al., 2020). The possibility of TB increases due to malnutrition, both micro- and macrodeficits, and compromised immunity. Infection with TB can intensify metabolism and require more energy, which can lead to eased appetite and nutritional malabsorption, ultimately increasing the possibility of being underweight (Feleke et al., 2019; Vonasek et al., 2022). A previous study revealed that, compared with the control group, TB patients had a greater likelihood of having a BMI of less than 18.5 kg/m² (Kitonsa et al., 2020). The nutritional status in this study was assessed via BMI; 64.7% of the respondents had a BMI <18.5, which was significantly associated with the incidence of childhood TB (p<0.05).

This study revealed that 54.5% of the respondents had a history of interaction with active TB patients. The proportions of TB and non-TB cases in the child group differed significantly when the children were exposed to cases of active TB (p value=0.002). The strong connection between TB disease in children and a history of interaction with the source of TB cases is crucial since it is well known that the risk of developing TB is directly proportional to the number of germs to which a child is exposed (Koura et al., 2023). Multivariate analysis in previous research revealed a significant relationship between household contacts of children with active TB, with OR = 15 288 and 95% CI: 5,378–43 457; household contacts with smokers, with OR = 7,094 and 95% CI: 2,128–23 648; and contacts > 18 hours with TB sufferers, with OR = 4.681 and 95% CI: 1.198–18.294 (Laghari et al., 2019). Children under the age of five, especially those under 1 year, who meet a source of TB cases usually have a 50% chance of contracting TB, and ninety-five percent of TB cases occur within a year after contact. Home contacts, caregivers, and

medical professionals are considered close contacts of active tuberculosis patients (Patra et al., 2015) because they have a greater chance of contracting *Mycobacterium tuberculosis* and developing primary active tuberculosis (Rea et al., 2022; Reichler et al., 2018).

This study revealed that 73.1% of the respondents had been vaccinated against BCG. The sole licensed and amply used vaccine to prevent TB prompted by *M. tuberculosis* is Bacillus Calmette-Guérin (BCG) (Casey et al., 2016; Katelaris et al., 2019). BCG immunization is highly effective in preventing meningitis and miliary TB in children, with a 70 to 80% success rate (Katelaris et al., 2019; Roy et al., 2019). An 18-year cohort study in Tunisia reported that 36.5% of TB cases occurred in individuals who received BCG vaccination, whereas 63.3% of unvaccinated individuals did (Bennasrallah et al., 2019).

The respondents in this study more often had a history of exposure to cigarette smoke (57.1%), but none of the respondents had a history of HIV or diabetes (Table 1). In numerous countries where the prevalence of tuberculosis is high, most children are exposed to cigarette smoke or secondhand smoke. A comprehensive analysis encompassing eighteen observational studies revealed that exposure to tobacco-related smoke resulted in children being 3.41 times more likely to contract tuberculosis and 1.64 times more likely to develop latent tuberculosis infection (Patra et al., 2015). Research in Spain reported that exposure to cigarette smoke in children (95% CI: 9.63–18.17) increased the prevalence of TB (13.48%) (Altet et al., 2022). Greater vulnerability to pulmonary tuberculosis has been attributed to several biological factors, including poor mucosal secretion clearance, impaired alveolar macrophage phagocytic capacity, diminished immunological reactions, and/or CD4+ lymphopenia caused by cigarette nicotine (Burusie, Enquesilassie, Addissie, Dessalegn, & Lamaro, 2020; Wang et al., 2020). Another study that examined the relationship between TB disease and exposure to cigarette smoke reported that the detrimental impact of smoking on tuberculosis transmission was significantly reduced immediately after smoking cessation, highlighting the value of smoking cessation programs in the fight against TB (Chu et al., 2021). Case-control studies carried out in Brazil and India have shown that firewood or biomass smoke, apart from cigarette smoke, is a separate threat to TB. However, there is currently still a dearth of information about the pathway through which biomass smoke results in chronic pulmonary illness (Báez-Saldaña et al., 2021; Sarkar et al., 2017).

The World Health Organization (WHO) lists fever, inexplicable weight decrease, lethargy or exhaustion, sweating at night, chest pain, and cough lasting longer than two weeks as manifestations and indicators of TB (World Health Organization, 2024). Given the risk of overdue diagnosis in those infected and subsequent spread via close contacts, over time, cough is the most noticed manifestation of tuberculosis, thus summoning inquiry, particularly in resource-constrained states where tuberculosis incidence rises (defined as ≥ 100 in 100 000 inhabitants) (Field et al., 2018). Most respondents in the current study, 79.5%, came to the health center with cough symptoms, and the remaining 20.5% did not experience cough complaints. The change in body weight was not very different; only 49.4% of the respondents experienced weight loss. A total of 55.8% of the respondents had a fever, and 44.2% did not have a fever; malaise was present in 59.6% of the respondents, and 40.4% did not have malaise (Table 1). In the study population conducted in the Philippines, which provided positive TB test results, 84.9% of patients experienced cough ≥ 2 weeks, fever ≥ 2 weeks of unexplained cause (6.9%), and considerable and unintended weight loss (9.5%), and 6.3% of patients experienced exhaustion, sluggishness, and malaise (Lee et al., 2019). The findings of these studies revealed that all participants with cough symptoms, regardless of the length of time for TB examination, presented increased sensitivity from forty-two percent to fifty-one percent (Cheng et al., 2015).

This research revealed that 70.5% of respondents who had national health insurance facilities (BPJS) and 75.5% of respondents tested positive for pulmonary TB. Meanwhile, research in the Philippines reported that 16% of individuals who tested positive for TB were enrolled in a health insurance program (51%) (Lee et al., 2019). Since TB is known to be a disease of poverty, the national health insurance program in Indonesia both improves the capacity of TB patients to seek treatment and contributes significantly to the TB reporting rate. Another study revealed that 60.3% of patients' household income was either per or exceeded the provincial minimum wage (UMP). Almost all the respondents' parents had jobs, 96.8% of

Whom were grouped as nongovernment workers (85.1%) or government workers (10.9%), but only 14.1% had higher education (Table 2). A systematic review revealed a correlation between tuberculosis and poor educational attainment, with a pooled odds ratio of 2.34 (Ricvan Dana et al., 2024). In a South African study including adolescents aged 12 to 18 years, the characteristics that were predictive of TB were older age, ethnic background, low household earnings, and basic schooling or less for both parents (Tchakounte et al., 2022).

With respect to socioeconomic factors (Table 2), TB in childhood occurs more often in families with lower parental education ($p = 0.005$), parents who have jobs ($p < 0.001$), parents with incomes greater than or equal to the UMP ($p < 0.001$), and those with health insurance ($p < 0.001$). The associations between clinical features and the incidence of bacteriological TB in pediatric patients are displayed in Table 3. Symptoms of cough ($p < 0.001$), subfebrile fever ($p < 0.001$), malaise ($p < 0.001$), and weight loss or no weight gain ($p < 0.001$) were significantly different between TB and non-TB events. Owing to the correlation with the increasing prevalence of TB, socioeconomic determinants are linked to the availability of healthcare services, the capacity to fulfill nutritional demands, and the availability of decent and standardized living spaces (Hameed et al., 2019). Working parents are linked to the external environment and to tuberculosis infection, which can spread to children. The exposure that children obtain from a household member who has TB is nearly fourfold inclined to contract the disease (Long et al., 2022; Martinez et al., 2017).

Table 2 Characteristics related to socioeconomic factors, clinical symptoms, and smoke exposure between TB and non-TB respondents.

Characteristics	Frequency		TB	Non-TB	p Value*	Odd Ratio (OR)	95% CI
	N, mean	%, SD					
Parents education							
Low	134	85.9	97	37	0.005¶	0.998	0.66 – 1.150
High	22	14.1	16	6		1.015	0.425 – 2.422
Parents occupation							
Not working	5	3.2	4	1	0.000¶	1.522	0.175 – 13.238
Working	151	96.8	109	42		0.988	0.932 – 1.047
Household income							
< UMP	62	39.7	42	20	0.000¶	0.799	0.536 – 1.192
>= UMP	94	60.3	71	23		1.175	0.859 – 1.606
Government health insurance							
None	46	29.5	30	16	0.000¶	0.713	0.435 – 1.170
Yes	110	70.5	83	27		1.170	0.906 – 1.510
Chronic coughing ≥2 weeks							
Yes	124	79.5	90	34	0.937*	1.007	0.842 – 1.206
None	32	20.5	23	9		0.972	0.490 – 1.931
Weight loss or does not increase in the last 2 months							
Yes	77	49.4	60	17	0.000¶	1.343	0.893 – 2.020
None	79	50.6	53	26		0.776	0.568 – 1.059
Subfebris febris ≥2 weeks							
Yes	87	55.8	63	24	0.003¶	0.999	0.731 – 1.365
None	69	44.2	50	19		1.001	0.675 – 1.486
Malaise ≥ 2 weeks							
Yes	93	59.6	67	26	0.024¶	0.981	0.737 – 1.305
None	63	40.4	46	17		1.030	0.669 – 1.585

*Chi-square test, p value <0.05; ¶McNemar test, p value <0.05. UMP = provincial minimum wage.

Table 3 Characteristics related to clinical factors among TB and non-TB respondents on the basis of bacteriological diagnosis.

Characteristics	Frequency		TB	Non-TB	p Value*	Odd Ratio (OR)	95% CI
	N, mean	%, SD					
Chronic cough ≥2 weeks							
Yes	124	79.5	14	110	0.000¶	1.114	0.908 – 1.366
2. None	32	20.5	2	30		0.583	0.154 – 2.216
Weight drop or did not rise in the previous 2 months							
Yes	77	49.4	7	70	0.000¶	0.875	0.490 – 1.562
2. None	79	50.6	9	70		1.125	0.708 – 1.787
Subfebris fever ≥2 weeks							
Yes	87	55.8	11	76	0.000¶	1.266	0.880 – 1.822
2. None	69	44.2	5	64		0.684	0.323 – 1.446
Malaise ≥ 2 weeks							
Yes	93	59.6	10	83	0.000¶	1.054	0.704 – 1.578
2. None	63	40.4	6	57		0.921	0.474 – 1.788

¶McNemar test, p value <0.05.

4. Conclusions

Despite this modest study, this study underscores the importance of sociodemographic factors, socioeconomic factors, active TB contacts, malnutrition conditions, and exposure to cigarette smoke or air pollution in the diagnosis of TB in children, in addition to TB-related clinical symptoms (cough, fever, malaise, and weight loss).

Ethical considerations

All procedures involving patients in this study were approved by the Ethical Review Committee of the Faculty of Medicine, Trisakti University No.055/KER/FK/I/2024. Informed consent was signed by all patients' legal guardians.

Conflict of interest

The authors have disclosed no conflicts of interest.

Funding

This research received no external financial support.

References

- Abdullahi, O., Moses, N., Sanga, D., & Annie, W. (2021). The effect of empirical and laboratory-confirmed tuberculosis on treatment outcomes. *Scientific Reports*, 11(1), 14854. doi:10.1038/s41598-021-94153-0
- Alao, M. A., Maroushek, S. R., Chan, Y. H., Asinobi, A. O., Slusher, T. M., & Gbadero, D. A. (2020). Treatment outcomes of Nigerian patients with tuberculosis: A retrospective 25-year review in a regional medical center. *PLoS one*, 15(10), e0239225. doi:10.1371/journal.pone.0239225
- Altet, N., Latorre, I., Jiménez-Fuentes, M. Á., Soriano-Arandes, A., Villar-Hernández, R., Millà, C., . . . Group, o. b. o. P. S. S. W. (2022). Tobacco Smoking and Second-Hand Smoke Exposure Impact on Tuberculosis in Children. *Journal of Clinical Medicine*, 11(7), 2000. Retrieved from <https://www.mdpi.com/2077-0383/11/7/2000>
- Asadi, L., Croxen, M., Heffernan, C., Dhillon, M., Paulsen, C., Egedahl, M. L., . . . Long, R. (2022). How much do smear-negative patients really contribute to tuberculosis transmissions? Re-examining an old question with new tools. *eClinicalMedicine*, 43. doi:10.1016/j.eclinm.2021.101250
- Báez-Saldaña, R., Canseco-Raymundo, A., Ixcot-Mejía, B., Juárez-Verdugo, I., Escobar-Rojas, A., Rumbo-Nava, U., . . . Arrieta, O. (2021). Case-control study about magnitude of exposure to wood smoke and risk of developing lung cancer. *European Journal of Cancer Prevention*, 30(6), 462-468. doi:10.1097/cej.0000000000000644
- Basile, F. W., Nabeta, P., Ruhwald, M., & Song, R. (2022). Pediatric Tuberculosis Diagnostics: Present and Future. *Journal of the Pediatric Infectious Diseases Society*, 11(Supplement_3), S85-S93. doi:10.1093/jpids/piac082
- Bennasrallah, C., Kacem, M., Dhoub, W., Zemni, I., Ben Fredj, M., Abroug, H., . . . Sriha Belguith, A. (2019). BCG vaccination and tuberculosis prevention: A forty years cohort study, Monastir, Tunisia. *PLoS one*, 14(8), e0219991. doi:10.1371/journal.pone.0219991
- Burusie, A., Enquesilassie, F., Addissie, A., Dessalegn, B., & Lamaro, T. (2020). Effect of smoking on tuberculosis treatment outcomes: A systematic review and meta-analysis. *PLoS one*, 15(9), e0239333. doi:10.1371/journal.pone.0239333
- Campos, L. C., Rocha, M. V. V., Willers, D. M. C., & Silva, D. R. (2016). Characteristics of Patients with Smear-Negative Pulmonary Tuberculosis (TB) in a Region with High TB and HIV Prevalence. *PLoS one*, 11(1), e0147933. doi:10.1371/journal.pone.0147933
- Casey, R. M., Dumolard, L., Danovaro-Holliday, M. C., Gacic-Dobo, M., Diallo, M. S., Hampton, L. M., & Wallace, A. S. (2016). Global Routine Vaccination Coverage, 2015. *MMWR Morb Mortal Wkly Rep*, 65(45), 1270-1273. doi:10.15585/mmwr.mm6545a5
- Cheng, J., Wang, L., Zhang, H., & Xia, Y. (2015). Diagnostic Value of Symptom Screening for Pulmonary Tuberculosis in China. *PLoS one*, 10(5), e0127725. doi:10.1371/journal.pone.0127725
- Chidambaram, V., Tun, N. L., Majella, M. G., Ruelas Castillo, J., Ayeh, S. K., Kumar, A., . . . Karakousis, P. C. (2021). Male Sex Is Associated With Worse Microbiological and Clinical Outcomes Following Tuberculosis Treatment: A Retrospective Cohort Study, a Systematic Review of the Literature, and Meta-analysis. *Clinical Infectious Diseases*, 73(9), 1580-1588. doi:10.1093/cid/ciab527
- Chilyabanyama, R., Kamanga, N., & Mwandia, J. N. (2024). Factors associated with tuberculosis treatment outcomes among TB patients aged 15 years and older at chawama level one hospital in Lusaka, Zambia. *Global Public Health*, 19(1), 2307979. doi:10.1080/17441692.2024.2307979
- Chu, A. L., Lecca, L. W., Calderón, R. I., Contreras, C. C., Yataco, R. M., Zhang, Z., . . . Huang, C.-C. (2021). Smoking Cessation in Tuberculosis Patients and the Risk of Tuberculosis Infection in Child Household Contacts. *Clinical Infectious Diseases*, 73(8), 1500-1506. doi:10.1093/cid/ciab504
- Feleke, B. E., Feleke, T. E., & Biadlegne, F. (2019). Nutritional status of tuberculosis patients, a comparative cross-sectional study. *BMC Pulmonary Medicine*, 19(1), 182. doi:10.1186/s12890-019-0953-0
- Field, S. K., Escalante, P., Fisher, D. A., Ireland, B., Irwin, R. S., Adams, T. M., . . . Weir, K. (2018). Cough Due to TB and Other Chronic Infections: CHEST Guideline and Expert Panel Report. *CHEST*, 153(2), 467-497. doi:10.1016/j.chest.2017.11.018
- Hameed, S., Zuberi, F. F., Hussain, S., & Ali, S. K. (2019). Risk factors for mortality among inpatients with smear positive pulmonary tuberculosis: Smear positive pulmonary tuberculosis. *Pakistan Journal of Medical Sciences*, 35(5). doi:10.12669/pjms.35.5.919
- Hamid, M., Brooks, M. B., Madhani, F., Ali, H., Naseer, M. J., The Childhood Tuberculosis Karachi, G., . . . Amanullah, F. (2019). Risk factors for unsuccessful tuberculosis treatment outcomes in children. *PLoS one*, 14(9), e0222776. doi:10.1371/journal.pone.0222776
- Katellaris, A. L., Jackson, C., Southern, J., Gupta, R. K., Drobniowski, F., Lalvani, A., . . . Abubakar, I. (2019). Effectiveness of BCG Vaccination Against Mycobacterium tuberculosis Infection in Adults: A Cross-sectional Analysis of a UK-Based Cohort. *The Journal of Infectious Diseases*, 221(1), 146-155. doi:10.1093/infdis/jiz430
- Kitonsa, P. J., Nalutaaya, A., Mukibi, J., Nakasolya, O., Isooba, D., Kamoga, C., . . . Kendall, E. A. (2020). Evaluation of underweight status may improve identification of the highest-risk patients during outpatient evaluation for pulmonary tuberculosis. *PLoS one*, 15(12), e0243542. doi:10.1371/journal.pone.0243542
- Koura, K. G., Mbitikon, O. B., Fogbé, A. A., Ouédraogo, A. R., Kuate Kuate, A., Magassouba, A. S., . . . Badoum, G. (2023). Programmatic Implementation of Contact Investigation in Eight African Countries. *Tropical Medicine and Infectious Disease*, 8(1), 29. Retrieved from <https://www.mdpi.com/2414-6366/8/1/29>
- Laghari, M., Sulaiman, S. A. S., Khan, A. H., Talpur, B. A., Bhatti, Z., & Memon, N. (2019). Contact screening and risk factors for TB among the household contact of children with active TB: a way to find source case and new TB cases. *BMC Public Health*, 19(1), 1274. doi:10.1186/s12889-019-7597-0
- Lee, S., Lau, L., Lim, K., Ferma, J., Dodd, W., & Cole, D. (2019). The Presence of Cough and Tuberculosis: Active Case Finding Outcomes in the Philippines. *Tuberculosis Research and Treatment*, 2019(1), 4578329. doi:https://doi.org/10.1155/2019/4578329
- Long, R., Divangahi, M., & Schwartzman, K. (2022). Chapter 2: Transmission and pathogenesis of tuberculosis. *Canadian Journal of Respiratory, Critical Care, and Sleep Medicine*, 6(sup1), 22-32. doi:10.1080/24745332.2022.2035540
- Martínez-Andrés, M., Bartolomé-Gutiérrez, R., Rodríguez-Martín, B., Pardo-Guijarro, M. J., & Martínez-Vizcaino, V. (2017). "Football is a boys' game": children's perceptions about barriers for physical activity during recess time. *International Journal of Qualitative Studies on Health and Well-being*, 12(1), 1379338. doi:10.1080/17482631.2017.1379338
- Martinez, L., Shen, Y., Mupere, E., Kizza, A., Hill, P. C., & Whalen, C. C. (2017). Transmission of Mycobacterium Tuberculosis in Households and the Community: A Systematic Review and Meta-Analysis. *American Journal of Epidemiology*, 185(12), 1327-1339. doi:10.1093/aje/kwx025
- Miller, P. B., Zalwango, S., Galiwango, R., Kakaire, R., Sekandi, J., Steinbaum, L., . . . Kiwanuka, N. (2021). Association between tuberculosis in men and social network structure in Kampala, Uganda. *BMC Infectious Diseases*, 21(1), 1023. doi:10.1186/s12879-021-06475-z

- Ministry of Health of the Republic of Indonesia. Tata Laksana Tuberkulosis Anak dan Remaja. Ministry of Health of the Republic of Indonesia Tahun 2023. Accessed on May 2, 2024. <https://tbindonesia.or.id/wp-content/uploads/2024/02/Final-Petunjuk-Teknis-Tata-Laksana-TBC-Anak-Remaja-2023.pdf>.
- Musunge, B. B., Poda, G. G., & Chen, P.-C. (2020). Nutritional Status of Patients with Tuberculosis and Associated Factors in the Health Centre Region of Burkina Faso. *Nutrients*, 12(9), 2540. Retrieved from <https://www.mdpi.com/2072-6643/12/9/2540>
- Namuganga, A. R., Chegou, N. N., & Mayanja-Kizza, H. (2021). Past and Present Approaches to Diagnosis of Active Pulmonary Tuberculosis. *Frontiers in Medicine*, 8. doi:10.3389/fmed.2021.709793
- Ogbudebe, C. L., Adepoju, V., Ekerete-Udofia, C., Abu, E., Egesemba, G., Chukwueme, N., & Gidado, M. (2018). Childhood Tuberculosis in Nigeria: Disease Presentation and Treatment Outcomes. *Health Services Insights*, 11, 1178632918757490. doi:10.1177/1178632918757490
- Patra, J., Bhatia, M., Suraweera, W., Morris, S. K., Patra, C., Gupta, P. C., & Jha, P. (2015). Exposure to Second-Hand Smoke and the Risk of Tuberculosis in Children and Adults: A Systematic Review and Meta-Analysis of 18 Observational Studies. *PLOS Medicine*, 12(6), e1001835. doi:10.1371/journal.pmed.1001835
- Rea, E., Huard, J., & Lee, R. (2022). Chapter 11: Tuberculosis contact investigation and outbreak management. *Canadian Journal of Respiratory, Critical Care, and Sleep Medicine*, 6(sup1), 167-183. doi:10.1080/24745332.2022.2037909
- Reichler, M. R., Khan, A., Sterling, T. R., Zhao, H., Moran, J., McAuley, J., . . . Team, T. E. S. C. T. O. (2018). Risk and Timing of Tuberculosis Among Close Contacts of Persons with Infectious Tuberculosis. *The Journal of Infectious Diseases*, 218(6), 1000-1008. doi:10.1093/infdis/jiy265
- Richens, J. G., Lee, C. M., & Johri, S. (2020). Improving the accuracy of medical diagnosis with causal machine learning. *Nature Communications*, 11(1), 3923. doi:10.1038/s41467-020-17419-7
- Ricvan Dana, N., Rika, S., Pudia M, I., Benny Alexander, M., Muthia, S., Linda, R., . . . Mochammad Fariz, A. (2024). MODIFIABLE AND NON-MODIFIABLE RISK FACTORS FOR TUBERCULOSIS AMONG ADULTS IN INDONESIA: A SYSTEMATIC REVIEW AND META-ANALYSIS. *African Journal of Infectious Diseases (AJID)*, 18(2), 19-28. doi:10.21010/Ajidv18i2.3
- Roy, P., Vekemans, J., Clark, A., Sanderson, C., Harris, R. C., & White, R. G. (2019). Potential effect of age of BCG vaccination on global paediatric tuberculosis mortality: a modelling study. *The Lancet Global Health*, 7(12), e1655-e1663. doi:10.1016/S2214-109X(19)30444-9
- Sarkar, M., Srinivasa, Madabhavi, I., & Kumar, K. (2017). Tuberculosis associated chronic obstructive pulmonary disease. *The Clinical Respiratory Journal*, 11(3), 285-295. doi:<https://doi.org/10.1111/crj.12621>
- Siddalingaiah, N., Chawla, K., Nagaraja, S. B., & Hazra, D. (2023). Risk factors for the development of tuberculosis among the pediatric population: a systematic review and meta-analysis. *European Journal of Pediatrics*, 182(7), 3007-3019. doi:10.1007/s00431-023-04988-0
- Tchakounte Youngui, B., Tchounga, B. K., Graham, S. M., & Bonnet, M. (2022). Tuberculosis Infection in Children and Adolescents. *Pathogens*, 11(12), 1512. Retrieved from <https://www.mdpi.com/2076-0817/11/12/1512>
- Tilahun, G., & Gebre-Selassie, S. (2016). Treatment outcomes of childhood tuberculosis in Addis Ababa: a five-year retrospective analysis. *BMC Public Health*, 16(1), 612. doi:10.1186/s12889-016-3193-8
- UNICEF Indonesia. Desk Review: Pediatric Tuberculosis with a Focus on Indonesia 2022. *UNICEF Indonesia*. Accessed on May 2, 2024. <https://www.unicef.org/indonesia/reports/desk-review-pediatric-tuberculosis-focus-indonesia>.
- Vonasek, B. J., Radtke, K. K., Vaz, P., Buck, W. C., Chabala, C., McCollum, E. D., . . . Garcia-Prats, A. J. (2022). Tuberculosis in children with severe acute malnutrition. *Expert Review of Respiratory Medicine*, 16(3), 273-284. doi:10.1080/17476348.2022.2043747
- Vukugah, T. A., Akoku, D. A., Tchoupa, M. M., & Lambert, E. (2022). Epidemiology of Pediatric Tuberculosis and Factors Associated with Unsuccessful Treatment Outcomes in the Centre Region of Cameroon: A Three-Year Retrospective Cohort Study. *Interdisciplinary Perspectives on Infectious Diseases*, 2022(1), 2236110. doi:<https://doi.org/10.1155/2022/2236110>
- Wang, E. Y., Arrazola, R. A., Mathema, B., Ahluwalia, I. B., & Mase, S. R. (2020). The impact of smoking on tuberculosis treatment outcomes: a meta-analysis. *Int J Tuberc Lung Dis*, 24(2), 170-175. doi:10.5588/ijtld.19.0002
- World Health Organization. World health statistics 2024: monitoring health for the SDGs, Sustainable Development Goals. Geneva: WHO. Accessed on August 21, 2024. <https://www.who.int/data/gho/publications/world-health-statistics>
- World Health Organization. WHO Global Tuberculosis Report 2023. Geneva: WHO. Accessed on May 2, 2024. <https://www.who.int/teams/global-tuberculosis-programme/data>.
- World Health Organization. WHO consolidated guidelines on tuberculosis: Module 5: Management of tuberculosis in children and adolescents 2022. Geneva: WHO. Accessed on May 2, 2024. https://www.ncbi.nlm.nih.gov/books/NBK579387/pdf/Bookshelf_NBK579387.pdf.
- Zürcher, K., Ballif, M., Kiertiburanakul, S., Chenal, H., Yotebieng, M., Grinsztejn, B., . . . consortium, t. I. E. D. t. E. A. (2019). Diagnosis and clinical outcomes of extrapulmonary tuberculosis in antiretroviral therapy programmes in low- and middle-income countries: a multicohort study. *Journal of the International AIDS Society*, 22(9), e25392. doi:<https://doi.org/10.1002/jia2.25392>

dr.Machrumnizar

Determining Factors of Lung Tuberculosis among Indonesian Children

 Artikel Jurnal 1

Document Details

Submission ID

trn:oid::3618:120908310

Submission Date

Nov 11, 2025, 2:48 PM GMT+7

Download Date

Nov 11, 2025, 3:43 PM GMT+7

File Name

Determining Factors of Lung Tuberculosis among Indonesian Children.docx

File Size

57.9 KB

16 Pages

5,469 Words

31,958 Characters

6% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.

Filtered from the Report

- Bibliography
- Quoted Text
- Small Matches (less than 10 words)

Exclusions

- 63 Excluded Matches

Match Groups

-  **6 Not Cited or Quoted 4%**
Matches with neither in-text citation nor quotation marks
-  **4 Missing Quotations 2%**
Matches that are still very similar to source material
-  **0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 1%  Internet sources
- 6%  Publications
- 0%  Submitted works (Student Papers)

Integrity Flags

0 Integrity Flags for Review

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups

- 6 Not Cited or Quoted 4%**
Matches with neither in-text citation nor quotation marks
- 4 Missing Quotations 2%**
Matches that are still very similar to source material
- 0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
- 0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 1% Internet sources
- 6% Publications
- 0% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

- 1 Publication**
Machrumnizar Machrumnizar, Yuliana Yuliana, Adang Bachtiar. "Determining fac..." 6%
- 2 Internet**
discovery.researcher.life <1%

Determining Factors of Lung Tuberculosis among Indonesian Children in Community Health Centers: A Cross-sectional Study

Machrumnizar Machrumnizar^{1*}, Yuliana Yuliana², Adang Bachtiar^{3*}

¹Doctoral Program, Faculty of Public Health, University of Indonesia, Indonesia

²Departement of Parasitology, Faculty of Medicine, Universitas Trisakti

³Department of Health Analysis and Policy, Faculty of Public Health, University of Indonesia, Indonesia

Corresponding author: adang@post.harvard.edu*

Abstract

Background: Tuberculosis is a serious global health issue and is particularly prevalent among children. Early and correct diagnosis is very important for effective management, but tuberculosis diagnosis in children is often difficult because the clinical symptoms are non-specific and difficult to collect specimens. This study aims to determine risk factors based on sociodemographic, socioeconomic, nutrition and clinical status associated with tuberculosis. **Methods:** This study is a cross-sectional study. Children aged 0 to 14 years who came to the community health center with one or more clinical symptoms related to TB from January to December 2023 were studied. Research participants underwent two diagnostic methods, clinical and bacteriological. **Results:** Positif Clinical diagnosis was found to be 72.4%. Meanwhile, bacteriologically confirmed participants were 10.3%, the remainder were unable to produce sputum. Most respondents were men (62.8%), with 66% of respondents aged ≥ 5 years, and those sociodemographic factors have a significant relationship with tuberculosis in children ($p < 0.05$). This study found that 64.7% of respondents had a BMI < 18.5 and significantly related to tuberculosis in children ($p < 0.05$). Furthermore, tuberculosis incidence and malnutrition also have a significant relationship ($p < 0.05$). Contact with active tuberculosis cases shows a significant difference in the proportion of tuberculosis and non-tuberculosis cases ($p = 0.009$), while 57.1% of children were exposed to cigarette smoke ($p < 0.05$). Government health insurance (BPJS) was used by 70.5% of respondents who were tested positive for tuberculosis, and it is statistically significant ($p < 0.05$). Other socio-economic factors were also significant ($p < 0.05$) for tuberculosis cases in children whose parents had low education at 85.9%; have 96.8% of jobs with a household income equal to or more than the provincial minimum wage (UMP) of 60.3%. **Conclusion:** In the future, these factors can be utilized to construct prediction models for clinical diagnosis in children suspected of tuberculosis, especially machine learning-based models, in addition to tuberculosis-related clinical symptoms.

Keyword: pediatrics, contact with active tuberculosis cases, tuberculosis, risk factor, comorbidity

Introduction

Among the leading source for morbidity and death in children aged 0 – 14 years is tuberculosis (TB), which is around 12% of the total 10.6 million TB sufferers in the world in 2023 and contributes to 3 – 25% of the total TB caseload [1]. Children might represent up to 20% of the global burden of TB in endemic spots, hence the real situation can vary from forecasts given

adequate reporting systems and appropriate diagnosis [2]. Still, there is a fairly modest rate of TB identified cases and notification, particularly in developing nations with limited resources, such as Indonesia, the estimated TB incidence of children aged 0 – 14 in 2022 is 110,881 children or around 15.3% of all TB cases in Indonesia [3]. According to a survey by The United Nations International Children's Emergency Fund (UNICEF), 35 percent of instances of TB among children between the ages of 0 and 14 go unreported, meaning these 110,881 children are only 65% of the total child TB cases in 2022 [4]. Mentioned data indicate the level of community transmission is relatively high.

Bacteriological testing is the standard diagnosis in adult TB but has low sensitivity and specificity when used in children [5]. Since the prevalence of TB in children serves as an immediate clue and surrogate for TB spreads in the community, there is a dearth of novel, easily used, and inexpensive diagnostic methods for TB, especially in remote areas with limited laboratory and x-ray facilities, adds to existing problems [4]. Diagnosis of tuberculosis in children aged 0 – 14 years without specific symptoms is a big challenge for doctors, academics, and program managers because it can hinder diagnosis and lead to worse treatment outcomes [6]. Current diagnostic methods have limited sensitivity, especially in children aged 0 – 14 years [7]. The scoring system, which is an alternative for diagnosing TB in children, is not yet fully implemented according to guidelines [3,8].

Despite clinical and biological variables, there are several socioeconomic and environmental variables that increase the likelihood of TB in children between the ages of 0 and 14. These risk factors are very useful for predicting the development of TB disease. Therefore, knowing the risk factors for TB in children is crucial for making the diagnosis and initiating treatment as soon as possible for the best outcomes. Community health centers in Indonesia are the main place where TB patients seek treatment, so they are the starting point for TB case care which includes early diagnosis, treatment, and reporting. The purpose of this study is to assess the sociodemographic, socioeconomic, and clinical risk factors for TB in children ages 0 to 14 who exhibit one or more TB-related clinical symptoms.

Material and Methods

Data collection

This research was conducted at a community health center in the administrative area of South Jakarta, Indonesia. The respondents are children, aged 0 to 14 years, who came with a cough to receive treatment at the community health centers, between January and December 2023.

Research design

We used a cross-sectional design and studied children with normal nutritional status or malnutrition, including all genders, aged 0 to 14 years, who came with a cough to receive treatment at the community health centers, between January and December 2023. Respondents were determined using consecutive sampling and the sample size was determined using the total sampling method, all respondents who met the inclusion criteria would be included in the study. TB diagnosis is categorized based on clinical diagnosis (accompanied by x-rays and/or tuberculin tests) and bacteriology. The clinical diagnosis was determined by the occurrence or absence of symptoms of a persistent cough lasting more than 2 weeks, weight depletion or no weight gain in the previous two months, fever ≥ 2 weeks, and malaise ≥ 2 weeks. Socio-demographic factors (age, gender, age group, weight, height, nutritional status) were recorded during the history taking.

Contact with active TB cases, history of BCG immunization and exposure to cigarette smoke/air pollution were categorized dichotomously. Socio-economic factors (education, employment and parental income, and health insurance) were also categorized dichotomously. This study also recorded the presence or absence of comorbid factors (HIV and DM). A child's nutritional status is determined by the body mass index, or BMI, is calculated using the ratio between body weight in kilograms and height in meters squared. Furthermore, the BMI value of children aged 0-60 months must be compared with the WHO 2005 standard BMI value (WHO, 2006); and BMI values for children and adolescents (5–19 years old) in comparison to the WHO/NCHS 2007 reference (WHO, 2007).

Data analysis

All data that had been collected and entered standard Microsoft Excel was then analyzed using SPSS for Windows (SPSS Inc version 25.0, Chicago, IL). Variables were initially examined univariately, and then a Chi-square statistic was used to evaluate categorical variables in order to determine risk factors in children with tuberculosis. The McNemar statistic was used to estimate the proportion of risk variables between TB and non-TB patients. A statistically significant probability if smaller than 0.05. The odds ratio (OR) and 95% confidence interval (CI) were computed to ascertain the strength of the link.

Results

In this study, 113 out of 156 respondents were diagnosed with pulmonary TB from January to December 2023. Of the TB cases who received treatment, 72.4% of respondents were clinically diagnosed with pulmonary TB. Meanwhile, bacteriologically confirmed TB cases were found in 10.3% of respondents and 89.7% of respondents were unable to produce phlegm. Most of responders (62.8%) were boys, with 34% being less than five years old and 66% being older than five (**Table 1**).

The nutritional status was assessed using the BMI, 64.7% of respondents had a BMI <18.5 and it was significantly associated with the incidence of childhood TB ($p < 0.05$). A previous record of interaction with active tuberculosis patients was reported by 54.5% of the respondents. The proportion of TB and non-TB occurrences in the child group differs significantly when exposed to cases of active TB ($p\text{-value} = 0.002$). Most respondents received BCG immunization, namely 73.1%. Respondents more often had a history of exposure to cigarette smoke, namely 57.1%, but all respondents did not have a history of HIV and diabetes (**Table 1**).

Most respondents came to the health center with cough symptoms, namely 79.5% and the remaining 20.5% did not experience cough complaints. The change in body weight was not much different, only 49.4% of respondents experienced weight loss. 55.8% of respondents had fever and 44.2% did not have fever, malaise was present in 59.6% of respondents and 40.4% did not have malaise (**Table 1**). All respondents did not use private insurance, but most respondents used BPJS insurance facilities, 70.5%, while 29.5% did not have insurance. Most of the patient's household income corresponds to or exceeds the provincial minimum wage (UMP) of 60.3%. Almost all the respondents' parents had jobs, 96.8% of whom were grouped as non-PNS 85.9% and PNS 10.9%, but only 14.1% had higher education (**Table 2**).

Most respondents were diagnosed with TB based on clinical diagnosis, because it was difficult to obtain children's sputum specimens for bacteriological examination. A bacteriological diagnosis could only be made in 10.3% of respondents and the majority (81.25%) were >10 years old. According to sociodemographic data (**Table 1**) indicates that there are statistically significant

2 variations in proportions of boys and girls ($p<0.001$), body weight ($p<0.001$), height ($p<0.001$). This study showed a notable association between the incidence of TB and nutritional status ($p=0.035$), exposure to cigarette smoke or air pollution ($p=0.006$), but not significantly with BCG immunization ($p=0.328$).

Regarding socioeconomic factors (**Table 2**), TB in childhood occurs more often in families with lower parental education ($p=0.005$), parents who have jobs ($p<0.001$), parental income greater than or equal to the UMP ($p<0.001$) and have health insurance ($p<0.001$). The association among clinical features and the incidence of pediatric TB patients with bacteriological diagnosis is displayed in **Table 3**. Symptoms of cough ($p<0.001$), subfebrile fever ($p<0.001$), malaise ($p<0.001$), and weight loss or no weight gain ($p<0.001$) showed significant differences in proportion between TB and non-TB events.

Table 1. Characteristics related to sociodemographic factors, TB contact, BCG immunization, nutritional status, and Exposure to smoking/air pollution between TB and non-TB respondents.

Characteristics	Frequency		TB	Non-TB	p-Value	Odd Ratio (OR)	95% CI
	N, mean	%, SD					
Bacteriological diagnosis							
1.Yes	16	10.3	16	0	0.009*	-	-
2.No	140	89.7	97	43			
Age (year)							
1.<5 years	53	34	38	15	0.000¶	0.964	0.594 – 1.564
2.≥5 years	103	66	75	28		1.019	0.790 – 1.315
Gender							
1.Boy	98	62.8	72	26	0.000¶	1.054	0.797 – 1.393
2.Girl	58	37.2	41	17		0.918	0.589 – 1.429
Weight (kg)	19.6218	8.94918			0.000*	-	18.21 – 21.04
Height (cm)	104.3910	26.23235			0.000*	-	100.24 – 108.54
Body Mass Index (BMI)							
1.<18.5	101	64.7	91	10	0.000*	3.463	1.997 – 6.005
2.>18.4	55	35.3	22	33		0.254	0.168 – 0.382
Contacts with active TB patients							
1.Yes	85	54.5	62	23	0.002¶	1.026	0.741 – 1.420
2.None	71	45.5	51	20		0.970	0.664 – 1.418
BCG immunization							
1.Yes	114	73.1	85	29	0.328*	1.115	0.883 – 1.408
2.No	42	26.9	28	14		0.761	0.445 – 1.302
nutritional status							
1.Malnutrition	100	64.1	90	10	0.000*	3.425	1.974 – 5.941
2.Normal	56	35.9	23	33		0.265	0.178 – 0.396

Smoke exposure/air pollution							
1. Yes	89	57.1	65	24	0.006¶	1.031	0.756 – 1.405
2. None	67	42.9	48	19		0.961	0.645 – 1.432

*Chi-square test, $p\text{-value} < 0.05$; ¶McNemar test, $p\text{-value} < 0.05$

Table 2. Characteristics related to socioeconomic factors, clinical symptoms, and smoke exposure between TB and non-TB respondents.

Characteristics	Frequency		TB	Non-TB	p-Value*	Odd Ratio (OR)	95% CI
	N, mean	%, SD					
Parents education							
1.Low	134	85.9	97	37	0.005¶	0.998	0.66 – 1.150
2.High	22	14.1	16	6		1.015	0.425 – 2.422
Parents occupation							
• Not working	5	3.2	4	1	0.000¶	1.522	0.175 –
• Working	151	96.8	109	42		0.988	13.238 0.932 – 1.047
Household income							
• < UMP	62	39.7	42	20	0.000¶	0.799	0.536 – 1.192
• ≥ UMP	94	60.3	71	23		1.175	0.859 – 1.606
Government health insurance							
1. None	46	29.5	30	16	0.000¶	0.713	0.435 – 1.170
2. Yes	110	70.5	83	27		1.170	0.906 – 1.510
Chronic coughing ≥2 weeks							
1. Yes	124	79.5	90	34	0.937*	1.007	0.842 – 1.206
2. None	32	20.5	23	9		0.972	0.490 – 1.931
Weight loss or does not increase in the last 2 months							
1. Yes	77	49.4	60	17	0.000¶	1.343	0.893 – 2.020
2. None	79	50.6	53	26		0.776	0.568 – 1.059
Subfebris febris ≥2 weeks							
1. Yes	87	55.8	63	24	0.003¶	0.999	0.731 – 1.365
2. None	69	44.2	50	19		1.001	0.675 – 1.486

Malaise $\geq$ 2 weeks							
1. Yes	93	59.6	67	26	0.024¶	0.981	0.737 – 1.305
2. None	63	40.4	46	17		1.030	0.669 – 1.585

*Chi-square test, p -value < 0.05 ; ¶McNemar test, p -value < 0.05 .

Table 3. Characteristics related to clinical factors among TB and non-TB respondents based on bacteriological diagnosis

Characteristics	Frequency		TB	Non-TB	p-Value*	Odd Ratio (OR)	95% CI
	N, mean	%, SD					
Chronic cough ≥ 2 weeks							
1. Yes	124	79.5	14	110	0.000¶	1.114	0.908 – 1.366
2. None	32	20.5	2	30		0.583	0.154 – 2.216
weight drop or did not rise in the previous 2 months							
1. Yes	77	49.4	7	70	0.000¶	0.875	0.490 – 1.562
2. None	79	50.6	9	70		1.125	0.708 – 1.787
Subfebris fever ≥ 2 weeks							
1. Yes	87	55.8	11	76	0.000¶	1.266	0.880 – 1.822
2. None	69	44.2	5	64		0.684	0.323 – 1.446
Malaise ≥ 2 weeks							
1. Yes	93	59.6	10	83	0.000¶	1.054	0.704 – 1.578
2. None	63	40.4	6	57		0.921	0.474 – 1.788

¶McNemar test, *p*-value <0.05

Discussion

The purpose of this cross-sectional study was to identify the predictive variables for both TB and non-TB diagnoses among children aged 0 – 14 years in Community Health Center facilities in Indonesia. TB risk factors such as HIV and diabetes were not found in child respondents, so no analysis was carried out.

Participants in this study underwent two diagnostic methods, namely clinical and bacteriological. Negative smear test results reportedly contribute to 13–20% of transmission and are, on average, 20–25% more infectious than those with positive test results [9]. Previous studies reported that most TB cases were diagnosed based on clinical criteria (50 – 58%) [10,11]. Likewise, a previous cohort study found 16% (58/372), 95% CI, 12-19% of cases came from smear negative sources [9], meaning 84% of TB cases were smear positive. However, another study reported that 65.2% of TB patients had negative BTA results and 34.8% had positive BTA results [12]. In children, TB disease is generally determined based on clinical diagnosis, which is the accepted method when bacteriological evidence is not available [13]. As is often reported, most respondents were boys. Various studies have similar findings where TB cases occur more often in boys [12,14,15] and variations in the age range of children who contract TB. Younger children are believed to be more affected than older children [16-19] due to household contacts of active TB patients and immature immunity. However, exposure to active TB can also occur in the community. Boys have a tendency for physical activity and high socialization, so they enjoy playing outside the house. This causes boys to potentially encounter adult active TB and be exposed to cigarette smoke or air pollution [20,21].

Malnutrition and tuberculosis (TB) have a reciprocal association in which malnutrition raises the likelihood of getting active TB by six to ten times and TB contributes patients to malnutrition [22,23]. Patients with active TB tend to be underweight, having a BMI below 18.5 kg/m², which is considered an index of malnutrition. Thus, TB incidence is strongly correlated with poor nutrition and lowly BMI [23]. The possibility of TB increases due to malnutrition, both micro and macro deficits, compromised immunity. Being infected with TB can intensify metabolism and require more energy, which can lead to ease appetites and nutrition malabsorption, ultimately raising the possibility of underweight [22,24]. Previous study found that, compared to the control group, TB patients had a higher likelihood of having a BMI of less than 18.5 kg/m² [25]. Numerous countries where tuberculosis prevalence is elevated, most children are exposed to cigarette smoke or secondhand smoke. A comprehensive analysis encompassing eighteen observational studies revealed that exposure to tobacco-related smoke resulted in children being 3.41 times more likely to contract tuberculosis and a 1.64-times more likely to develop latent tuberculosis infection [26]. Research in Spain reported that exposure to cigarette smoke in children (95% CI: 9.63–18.17) and had an increased prevalence of TB (13.48%) [27]. Higher vulnerability to pulmonary tuberculosis illness has been attributed to several biological factors, including poor mucosal secretion clearance, impaired alveolar macrophage phagocytic capacity, diminished immunological reaction, and/or CD4⁺ lymphopenia brought on by cigarette nicotine [28,29]. Another study that looked at the relationship between TB disease and exposure to cigarette smoke also reported that the detrimental impact of smoking on tuberculosis transmission was significantly reduced immediately after smoking cessation, hence highlighting the value of smoking cessation programs in the fight against TB [30]. Case-control studies carried out in Brazil and India have shown that firewood or biomass smoke, apart from cigarette smoke, is a separate threat to TB. But there is currently still a dearth of information about the pathway via which biomass smoke results in chronic pulmonary illness [31,32].

Noticing a high connection between TB disease in children and a history of interaction with the source of TB cases is crucial since it is well-known that the risk of developing TB is directly

proportionate to the quantity of germs one receives exposure to [33]. Multivariate analysis in previous research showed a significant relationship between household contacts of children with active TB, obtained OR = 15,288 and 95% CI: 5,378–43,457, household contacts with smokers obtained OR = 7,094 and 95% CI: 2,128–23,648, contacts > 18 hours with TB sufferers obtained OR = 4.681 and 95% CI: 1.198–18.294 [34]. Children beneath the age of five, especially under 1 year, who meet a source of TB cases usually have a 50% chance of contracting TB, and ninety-five percent of TB occurs within a year after contact. Home contacts, caregivers, and medical professionals are considered close contacts of active tuberculosis cases [26] because they have a greater chance of contracting *Mycobacterium tuberculosis* and developing primary active tuberculosis [35,36].

The sole licensed and amply used vaccine to prevent TB prompted by *M. tuberculosis* is Bacillus Calmette-Guérin (BCG) [37,38]. The BCG immunization has been shown to be highly effective in preventing meningitis and miliary TB in children, with a 70 to 80 percent success rate [37, 39]. An 18-year cohort study in Tunisia found 36.5% of TB cases in individuals who received BCG vaccination compared to 63.3% of unvaccinated individuals [40]. The World Health Organization (WHO) lists fever, inexplicable weight decrease, lethargy or exhaustion, sweating at night, chest pain, and cough lasting longer than two weeks as manifestation and indicators of TB. Given the risk of overdue diagnosis in those infected and subsequent spread via close contacts, overtime cough is the most noticed manifestation in tuberculosis thus summons inquiry, particularly in resource-constrained states where tuberculosis incidence rises (defined as ≥ 100 in 100,000 inhabitants) [41]. In the study population conducted in the Philippines, providing positive TB test results, 84.9% experienced cough ≥ 2 weeks, fever ≥ 2 weeks of unexplained cause (6.9%), considerable and unintended weight loss (9.5%), and 6.3% of patients experienced exhaustion, sluggishness, and malaise [42]. Study finding concluded that all participants with cough symptoms of whichever length for TB examining versed rose sensitivity from forty-two percent to fifty-one percent [43].

This research shows that 70.5% of respondents who have national health insurance facilities (BPJS), 75.5% of respondents tested positive for pulmonary TB. Meanwhile, research in the Philippines reported that 16% of individuals who tested positive for TB were enrolled in a health insurance program (51%) [42]. Since TB is known to be a disease of poverty, national health insurance program in Indonesia's both improves the capacity of TB patients to seek treatment and contributes significantly to TB reporting rate. The study's finding revealed the respondents' parents who were positive for TB had non-PNS jobs with a household income equal to or more than the provincial minimum wage but had low education. A systematic review of studies found a correlation between tuberculosis and a poor educational attainment with a pooled odds ratio = 2.34 [44]. In a South African study including adolescents aged 12 to 18, characteristics that were predictive of TB were older age, ethnic background, low household earning, and basic schooling or lesser for both parents [45]. Due to the correlation with the increasing prevalence of TB, socioeconomic determinants are linked to the availability of healthcare services, the capacity to fulfill nutrition demands, also the availability of a decent and standardized living space [46]. Working parents are linked to external environment and tuberculosis infection, which can spread to children. Exposure that children obtain from a household member who has TB is nearly fourfold inclined to contract the disease [47,48].

Conclusion

Despite the modest study, this study underscores the importance of sociodemographic, socioeconomic, active TB contacts, malnutrition conditions, and exposure to cigarette smoke or air pollution in the diagnosis of tuberculosis in children, in addition to TB-related clinical symptoms (cough, fever, malaise, and weight loss).

Study limitations

This research was conducted at a community health center and there were no patient home visits so it did not analyze environmental factors which are one of the determinants of TB. However, other factors managed to show meaningful results.

Ethical considerations

All procedures involving patients in this study were approved by the Ethical Review Committee of the Faculty of Medicine, Trisakti University No.055/KER/FK/I/2024. Informed consent was signed by all patients' legal guardians.

Acknowledgments

Not Applicable.

Conflict of Interest

The authors have disclosed no conflicts of interest.

Availability of data and materials

The corresponding author will make the datasets used and/or analyzed in this work accessible upon an acceptable request.

Authors' contributions

M.M: study concept, writing up, data interpretation, methodology writing, statistical analysis, Finalize the manuscript revisions and provide approval.

A.B: study concept, revise manuscript, and provide final approval.

Y.Y: data collection, revise manuscript, and provide final approval.

References

1. WHO Global Tuberculosis Report [Internet]. Geneva: World Health Organization; 2023. Available at: <https://www.who.int/teams/global-tuberculosis-programme/data>. Accessed on Mei 2nd, 2024.
2. Richens, J.G., Lee, C.M. & Johri, S. Improving the accuracy of medical diagnosis with causal machine learning. *Nat Commun* 11, 3923 (2020). Doi: 10.1038/s41467-020-17419-7.
3. Keboosterian Kesehatan RI. Direktorat Jenderal Pencegahan dan Pengendalian Penyakit Petunjuk Teknis Tata Laksana Tuberkulosis Anak dan Remaja [Internet]. Jakarta: Keboosterian Kesehatan RI. 2023. Available at <https://tbindonesia.or.id/wp-content/uploads/2024/02/Final-Petunjuk-Teknis-Tata-Laksana-TBC-Anak-Remaja-2023.pdf>
4. UNICEF Indonesia. Desk Review: Pediatric Tuberculosis with a Focus on Indonesia [Internet]. 2022. Accessed on Mei 2nd, 2024. Available at: <https://www.unicef.org/indonesia/reports/desk-review-pediatric-tuberculosis-focus-indonesia>
5. Basile, F.W., Nabeta, P., Ruhwald, M., Song, R. (2022). Pediatric Tuberculosis Diagnostics: Present and Future. *J Pediatric Infect Dis Soc*, 11(Suppleboyst_3), S85-S93. doi: 10.1093/jpids/piac082
6. Siddalingaiah N, Chawla K, Nagaraja SB, Hazra D. Risk factors for the development of tuberculosis among the pediatric population: a systematic review and meta-analysis. *Eur J Pediatr*. 2023 Jul;182(7):3007-3019. doi: 10.1007/s00431-023-04988-0.
7. Namuganga AR, Chegou NN, Mayanja-Kizza H. Past and Present Approaches to Diagnosis of Active Pulmonary Tuberculosis. *Front Med (Lausanne)*. 2021 Sep 23; 8: 709793. doi: 10.3389/fmed.2021.709793. PMID: 34631731; PMCID: PMC8495065.

8. WHO consolidated guidelines on tuberculosis: Module 5: Management of tuberculosis in children and adolescents [Internet]. Geneva: World Health Organization; 2022. Available at https://www.ncbi.nlm.nih.gov/books/NBK579387/pdf/Bookshelf_NBK579387.pdf. Accessed on Mei 2nd, 2024.
9. Asadi, L., Croxen, M., Heffernan, C., Dhillon, M., Paulsen, C., Egedahl, M. L., Tyrrell, G., Doroshenko, A., & Long, R. (2022). How much do smear-negative patients really contribute to tuberculosis transmissions? Re-examining an old question with new tools. *EClinicalMedicine*, 43, 101250. <https://doi.org/10.1016/j.eclinm.2021.101250>
10. Chilyabanyama, R., Kamanga, N., & Mwandia, J. N. (2024). Factors associated with tuberculosis treatment outcomes among TB patients aged 15 years and older at chawama level one hospital in Lusaka, Zambia. *Global Public Health*, 19(1). <https://doi.org/10.1080/17441692.2024.2307979>
11. Zürcher K, Ballif M, Kiertiburanakul S, Chenal H, Yotebieng M, Grinsztejn B, Michael D, Sterling TR, Ngonyani KM, Mandalakas AM, Egger M, Pettit AC, Fenner L; International Epidemiology Databases to Evaluate AIDS (IeDEA) consortium. Diagnosis and clinical outcomes of extrapulmonary tuberculosis in antiretroviral therapy programmes in low- and middle-income countries: a multicohort study. *J Int AIDS Soc*. 2019 Sep;22(9): e25392. doi: 10.1002/jia2.25392
12. Campos LC, Rocha MVV, Willers DMC, Silva DR (2016) Characteristics of Patients with Smear-Negative Pulmonary Tuberculosis (TB) in a Region with High TB and HIV Prevalence. *PLoS ONE* 11(1): e0147933. <https://doi.org/10.1371/journal.pone.0147933>
13. Abdullahi, O., Moses, N., Sanga, D. *et al.* The effect of empirical and laboratory-confirmed tuberculosis on treatment outcomes. *Sci Rep* 11, 14854 (2021). <https://doi.org/10.1038/s41598-021-94153-0>
14. Chidambaram, V., Tun, N. L., Majella, M. G., Castillo, J. R., Ayeh, S. K., Kumar, A., Neupane, P., Sivakumar, R. K., Win, E. P., Abbey, E. J., Wang, S., Zimmerman, A., Blanck, J., Gupte, A., Wang, J.-Y., & Karakousis, P. C. (2021). Sex differences in TB treatment outcomes: Retrospective cohort study and meta-analysis. *medRxiv*, 2021.04.26.21256155. <https://doi.org/10.1101/2021.04.26.21256155>
15. Alao MA, Maroushek SR, Chan YH, Asinobi AO, Slusher TM, Gbadero DA. Treatment outcomes of Nigerian patients with tuberculosis: A retrospective 25-year review in a regional medical center. *PLoS One*. 2020 Oct 29;15(10): e0239225. doi: 10.1371/journal.pone.0239225
16. Vukugah, T. A., Akoku, D. A., Tchoupa, M. M., & Lambert, E. (2022). Epidemiology of Pediatric Tuberculosis and Factors Associated with Unsuccessful Treatment Outcomes in the Centre Region of Cameroon: A Three-Year Retrospective Cohort Study. *Interdisciplinary perspectives on infectious diseases*, 2022, 2236110. <https://doi.org/10.1155/2022/2236110>
17. Ogbudebe CL, Adepoju V, Ekerete-Udofia C, Abu E, Egesemba G, Chukwueme N, Gidado M. Childhood Tuberculosis in Nigeria: Disease Presentation and Treatment Outcomes. *Health Serv Insights*. 2018; 11: 1178632918757490. doi: 10.1177/1178632918757490
18. Tilahun, G., & Gebre-Selassie, S. (2016). Treatment outcomes of childhood tuberculosis in Addis Ababa: a five-year retrospective analysis. *BMC public health*, 16, 612. <https://doi.org/10.1186/s12889-016-3193-8>
19. Hamid M, Brooks MB, Madhani F, Ali H, Naseer MJ; Childhood Tuberculosis Karachi Group; Becerra M, Amanullah F. Risk factors for unsuccessful tuberculosis treatment outcomes in children. *PLoS One*. 2019 Sep 25;14(9): e0222776. doi: 10.1371/journal.pone.0222776
20. Martínez-Andrés M, Bartolomé-Gutiérrez R, Rodríguez-Martín B, Pardo-Guijarro MJ, Martínez-Vizcaino V. (2017). "Football is a boys' game": children's perceptions about barriers for physical

- activity during recess time. *Int J Qual Stud Health Well-being*, 12(sup2): 1379338. doi: 10.1080/17482631.2017.1379338.
21. Miller, P.B., Zalwango, S., Galiwango, R. et al. (2021). Association between tuberculosis in men and social network structure in Kampala, Uganda. *BMC Infect Dis*, 21, 1023. <https://doi.org/10.1186/s12879-021-06475-z>
 22. Feleke, B. E., Feleke, T. E., & Biadlegne, F. (2019). Nutritional status of tuberculosis patients, a comparative cross-sectional study. *BMC pulmonary medicine*, 19(1), 182. <https://doi.org/10.1186/s12890-019-0953-0>
 23. Musuenge, B. B., Poda, G. G., & Chen, P. C. (2020). Nutritional Status of Patients with Tuberculosis and Associated Factors in the Health Centre Region of Burkina Faso. *Nutrients*, 12(9), 2540. <https://doi.org/10.3390/nu12092540>
 24. Vonasek, B. J., Radtke, K. K., Vaz, P., Buck, W. C., Chabala, C., McCollum, E. D., Marcy, O., Fitzgerald, E., Kondwani, A., & Garcia-Prats, A. J. (2022). Tuberculosis in children with severe acute malnutrition. *Expert review of respiratory medicine*, 16(3), 273–284. <https://doi.org/10.1080/17476348.2022.2043747>
 25. Kitonsa, P. J., Nalutaaya, A., Mukiibi, J., Nakasolya, O., Isooba, D., Kamoga, C., Baik, Y., Robsky, K., Dowdy, D. W., Katamba, A., & Kendall, E. A. (2020). Evaluation of underweight status may improve identification of the highest-risk patients during outpatient evaluation for pulmonary tuberculosis. *PloS one*, 15(12), e0243542. <https://doi.org/10.1371/journal.pone.0243542>
 26. Patra J, Bhatia M, Suraweera W, Morris SK, Patra C, Gupta PC, et al. (2015) Exposure to Second-Hand Smoke and the Risk of Tuberculosis in Children and Adults: A Systematic Review and Meta-Analysis of 18 Observational Studies. *PLoS Med* 12(6): e1001835. <https://doi.org/10.1371/journal.pmed.1001835>
 27. Altet, N., Latorre, I., Jiménez-Fuentes, M. Á., Soriano-Arandes, A., Villar-Hernández, R., Milà, C., Rodríguez-Fernández, P., Muriel-Moreno, B., Comella-Del-Barrio, P., Godoy, P., Millet, J. P., de Souza-Galvão, M. L., Jiménez-Ruiz, C. A., Domínguez, J., & On Behalf Of Pii Smoking Separ Working Group (2022). Tobacco Smoking and Second-Hand Smoke Exposure Impact on Tuberculosis in Children. *Journal of clinical medicine*, 11(7), 2000. <https://doi.org/10.3390/jcm11072000>
 28. Burusie, A., Enquesilassie, F., Addissie, A., Dessalegn, B., & Lamaro, T. (2020). Effect of smoking on tuberculosis treatment outcomes: A systematic review and meta-analysis. *PloS one*, 15(9), e0239333. <https://doi.org/10.1371/journal.pone.0239333>
 29. Wang, E. Y., Arrazola, R. A., Mathema, B., Ahluwalia, I. B., & Mase, S. R. (2020). The impact of smoking on tuberculosis treatment outcomes: a meta-analysis. *The international journal of tuberculosis and lung disease: the official journal of the International Union against Tuberculosis and Lung Disease*, 24(2), 170–175. <https://doi.org/10.5588/ijtld.19.0002>
 30. Chu, A. L., Lecca, L. W., Calderón, R. I., Contreras, C. C., Yataco, R. M., Zhang, Z., Becerra, M. C., Murray, M. B., & Huang, C. C. (2021). Smoking Cessation in Tuberculosis Patients and the Risk of Tuberculosis Infection in Child Household Contacts. *Clinical infectious diseases: an official publication of the Infectious Diseases Society of America*, 73(8), 1500–1506. <https://doi.org/10.1093/cid/ciab504>
 31. Báez-Saldaña R, Canseco-Raymundo A, Ixcot-Mejía B, Juárez-Verdugo I, Escobar-Rojas A, Rumbo-Nava U, Castillo-González P, León-Dueñas S, Arrieta O. Case-control study about magnitude of exposure to wood smoke and risk of developing lung cancer. *Eur J Cancer Prev*. 2021 Nov 1;30(6):462-468. doi: 10.1097/CEJ.0000000000000644

32. Sarkar M, Srinivasa, Madabhavi I, Kumar K. Tuberculosis associated chronic obstructive pulmonary disease. *Clin Respir J*. 2017 May;11(3):285-295. doi: 10.1111/crj.12621
33. Shaw J.B., Wynn-Williams N. Infectivity of pulmonary tuberculosis in relation to sputum status. *Am Rev Tuberc*. 1954 May;69(5):724-32. doi: 10.1164/art.1954.69.5.724
34. Laghari, M., Sulaiman, S.A.S., Khan, A.H. *et al*. Contact screening and risk factors for TB among the household contact of children with active TB: a way to find source case and new TB cases. *BMC Public Health* 19, 1274 (2019). <https://doi.org/10.1186/s12889-019-7597-0>
35. Rea, E., Huard, J., & Lee, R. (2022). Chapter 11: Tuberculosis contact investigation and outbreak management. *Canadian Journal of Respiratory, Critical Care, and Sleep Medicine*, 6(sup1), 167–183. <https://doi.org/10.1080/24745332.2022.2037909>
36. Reichler, M. R., Khan, A., Sterling, T. R., Zhao, H., Moran, J., McAuley, J., Bessler, P., Mangura, B., & Tuberculosis Epidemiologic Studies Consortium Task Order 2 Team (2018). Risk and Timing of Tuberculosis Among Close Contacts of Persons with Infectious Tuberculosis. *The Journal of infectious diseases*, 218(6), 1000–1008. <https://doi.org/10.1093/infdis/jiy265>
37. Katelaris AL, Jackson C, Southern J, Gupta RK, Drobniewski F, Lalvani A, Lipman M, Mangtani P, Abubakar I. Effectiveness of BCG Vaccination Against Mycobacterium Tuberculosis Infection in Adults: A Cross-sectional Analysis of a UK-Based Cohort. *J Infect Dis*. 2020 Jan 1;221(1):146-155. doi: 10.1093/infdis/jiz430
38. Casey RM, Dumolard L, Danovaro-Holliday MC, Gacic-Dobo M, Diallo MS, Hampton LM, Wallace AS. Global Routine Vaccination Coverage, 2015. *MMWR Morb Mortal Wkly Rep*. 2016 Nov 18;65(45):1270-1273. doi: 10.15585/mmwr.mm6545a5
39. Trunz BB, Fine P, Dye C. Effect of BCG vaccination on childhood tuberculous meningitis and miliary tuberculosis worldwide: a meta-analysis and assessment of cost-effectiveness. *Lancet*. 2006 Apr 8;367(9517):1173-80. doi: 10.1016/S0140-6736(06)68507-3
40. Bennasrallah C, Kacem M, Dhouib W, Zemni I, Ben Fredj M, Abroug H, et al. (2019) BCG vaccination and tuberculosis prevention: A forty-year cohort study, Monastir, Tunisia. *PLoS ONE* 14(8): e0219991. <https://doi.org/10.1371/journal.pone.0219991>
41. Field, S. K., Escalante, P., Fisher, D. A., Ireland, B., Irwin, R. S., & CHEST Expert Cough Panel (2018). Cough Due to TB and Other Chronic Infections: CHEST Guideline and Expert Panel Report. *Chest*, 153(2), 467–497. <https://doi.org/10.1016/j.chest.2017.11.018>
42. Lee, S., Lau, L., Lim, K., Ferma, J., Dodd, W., & Cole, D. (2019). The Presence of Cough and Tuberculosis: Active Case Finding Outcomes in the Philippines. *Tuberculosis research and treatment*, 2019, 4578329. <https://doi.org/10.1155/2019/4578329>
43. Cheng J, Wang L, Zhang H, Xia Y. Diagnostic value of symptom screening for pulmonary tuberculosis in China. *PLoS One*. 2015 May 22;10(5): e0127725. doi: 10.1371/journal.pone.0127725.
44. Dana, N. R., Rika, S., M, I. P., Alexander, M. B., Muthia, S., Linda, R., Astri, W., Zuhrah, T., Rahman, A. D., Rahmi, F., Nomira, P., Setia, N. D. A. W., Arif, L. B. L., Ainil, M., Octarini, E. M., Nova, L., Tamia, M. Y., Sri, R. A., Permata, S. A., Mimin, O., ... Fariz, A. M. (2024). Modifiable and non-modifiable risk factors for tuberculosis among adults in Indonesia: A Systematic review and Meta-analysis. *African journal of infectious diseases*, 18(2), 19–28. <https://doi.org/10.21010/Ajidv18i2.3>
45. Tchakounte Youngui B, Tchounga BK, Graham SM, Bonnet M. Tuberculosis Infection in Children and Adolescents. *Pathogens*. 2022 Dec 9;11(12):1512. doi: 10.3390/pathogens11121512.

46. Khaliq, A., Khan, I.H., Akhtar, M.W., & Chaudhry, M.N. (2015). Environmental Risk Factors and Social Determinants of Pulmonary Tuberculosis in Pakistan. *Epidemiology* (Sunnyvale) 2015;2015(5):3. doi: <https://doi.org/10.4172/2161-1165.1000201>
47. 45. Long R., Diwangahi M., Schwartzman K. Chapter 2: Transmission and Pathogenesis of Tuberculosis. *Can. J. Breathing. Critics. Sleep Care Med.* 2022; 6 :22–32. doi: 10.1080/24745332.2022.2035540.
48. 46. Martinez L., Shen Y., Mupere E., Kizza A., Hill PC, Whalen CC Transmission of Mycobacterium tuberculosis in Households and Communities: A Systematic Review and Meta-Analysis. *I. J. Epidemiol.* 2017; 185 :1327–1339. doi: 10.1093/aje/kwx025.