



## Complexity Journal Manuscript

Join Leading Researchers in the Field of Complexity. Publish Your Next Research With Us

Hindawi Open

# Journal of Oral Microbiology

### COUNTRY

United Kingdom



Universities and research institutions in United Kingdom

### SUBJECT AREA AND CATEGORY

- Dentistry
- Dentistry (miscellaneous)
- Medicine
- Infectious Diseases
- Microbiology (medical)

### PUBLISHER

Taylor and Francis Ltd.

### H-INDEX

# 45

### PUBLICATION TYPE

Journals

### ISSN

20002297

### COVERAGE

2009-2021

### INFORMATION

- [Homepage](#)
- [How to publish in this journal](#)
- [Contact](#)

### SCOPE

As the first Open Access journal in its field, the Journal of Oral Microbiology aims to be an influential source of knowledge on the aetiological agents behind oral infectious diseases. The journal is an international forum for original research on all aspects of 'oral health'. Articles which seek to understand 'oral health' through exploration of the pathogenesis, virulence, host-parasite interactions, and immunology of oral infections are of particular interest. However, the journal also welcomes work that



## Complexity Journal Manuscript

Join Leading Researchers in the Field of Complexity. Publish Your Next Research With Us Hindawi

Open

FIND SIMILAR JOURNALS ?

<p>1</p> <p><b>Molecular Oral Microbiology</b></p> <p>USA</p> <p><b>71%</b> similarity</p>	<p>2</p> <p><b>npj Biofilms and Microbiomes</b></p> <p>GBR</p> <p><b>57%</b> similarity</p>	<p>3</p> <p><b>BMC Microbiology</b></p> <p>GBR</p> <p><b>56%</b> similarity</p>	<p>4</p> <p><b>Critical Reviews Microbiology</b></p> <p>GBR</p> <p><b>52%</b> similarity</p>
--	---	---	--


### Complexity Journal Manuscript

Join Leading Researchers in the Field of Complexity. Publish Your Next Research With Us


Hindawi

Open


SJR



Total Documents




Total Cites    Self-Cites



Citations per document

External Cites per Doc    Cites per Doc



### Complexity Journal Manuscript


Join Leading Researchers in the Field of Complexity. Publish Your Next Research With Us Hindawi

% International Collaboration




Citable documents

Non-citable documents



Cited documents

Uncited documents



**Journal of Oral Microbiology**

Q1 Dentistry (miscellaneous) best quartile

SJR 2021 1.15

powered by scimagojr.com


Show this widget in your own website

Just copy the code below and paste within your html code:

`<a href="https://www.scimaç`

### SCImago Graphica

Explore, visually communicate and make sense of data with our [new data visualization tool](#).



**Sigma-Aldrich** **BENCHTOP BREAKTHROUGHS** **Roche** **MERCK**

Evolve your PCR with KAPA Biosystems

Learn More

Metrics based on Scopus® data as of April 2022

A **Afaf El-Ansary** 4 months ago

I want to check the status of my submitted manuscript entitled "Gum-gut axis: the potential role of salivary biomarkers in the diagnosis and prognosis of inflammatory bowel diseases

reply

### Complexity Journal Manuscript

Join Leading Researchers in the Field of Complexity. Publish Your Next Research With Us Hindawi



# Journal of Oral Microbiology



Taylor & Francis  
Taylor & Francis Group

[www.journals.tandf.co.uk/journals](http://www.journals.tandf.co.uk/journals)

## Journal of Oral Microbiology

[Submit](#) [About](#) [Browse](#) [Subscribe](#) 

# Editorial board

## Announcing renewed Editorial Board

It gives me great pleasure to announce that Bruce Paster (The Forsyth Institute and Harvard School of Dental Medicine) has joined Journal of Oral Microbiology as Deputy Editor-in-Chief. In addition, we have added numerous new members to the Editorial Board. Please join me in welcoming the new members and thanking those members who have renewed their commitment to the journal. I also acknowledge outgoing board members for their years of service who helped make Journal of Oral Microbiology the success it is today.

Ingar Olsen, Editor-in-Chief

### Editor-In-Chief

Ingar Olsen - University of Oslo, Norway

### Deputy Editor-in-Chief

Bruce Paster - The Forsyth Institute and Harvard School of Dental Medicine, USA

### Associate Editors

Alpdogan Kantarci - The Forsyth Institute, USA

Ann Progulsk-Fox - College of Dentistry, Florida, USA

Kazuhisa Yamazaki - RIKEN Center for Integrative Medical Sciences (IMS), Japan

## **Editorial Board**

Nezar Al-Hebshi - Temple University, Philadelphia, USA

Atsuo Amano - Osaka University, Japan

Daniel Belstrøm - University of Copenhagen, Denmark

Paulo H Braz-Silva - University of Sao Paulo, Brazil

Casey Chen - University of Southern California, USA

Georg Conrads - RWTH Aachen University Hospital, Germany

Patricia Corby - University of Pennsylvania, USA

Mike Curtis - King's College London, UK

Gunnar Dahlén - University of Gothenburg, Sweden

Stuart Dashper - University of Melbourne, Australia

Mary Ellen Davey - University of Florida, USA

Patricia Diaz-Moreno - University of Buffalo, USA

Anna Dongari-Bagtzoglou - University of Connecticut Health Center, USA

Magda Feres - University of Guarulhos, Sao Paulo, Brazil

Jorge Frias-Lopez - University of Florida, USA

Frank Gibson - University of Florida, USA

Octavio Gonzalez - University of Kentucky, USA

Dana Graves - University of Pennsylvania, USA

Daniel Grenier - Laval University, Canada

Yoshiaki Hasegawa - Aichi-Gakuin University, Nagoya, Japan

Xuesong He - The Forsyth Institute, USA

Hatice Hasturk - The Forsyth Institute, USA

Olivier Huck - University of Strasbourg, France

Eija Könönen - University of Turku, Finland

Tomoko Kurita-Ochiai - Nihon University, Japan

Richard L Lamont - University of Louisville, USA

Yuqing Li - Sichuan University, China

Suzanne M Michalek - University of Alabama at Birmingham, USA

Pallavi Murugkar, Stanford University, California, USA

Keiji Nagano - Health Sciences University of Hokkaido, Japan

Jan S Potempa - University of Louisville, USA

Pirkko J Pussinen - University of Helsinki, Finland



Eric C Reynolds - University of Melbourne, Australia  
Aurea Simon-Soro - University of Pennsylvania, USA  
José F. Siqueira - Estácio de Sá University, Brazil  
Graham Stafford - University of Sheffield, UK  
Flavia Teles - University of Pennsylvania, USA  
Thomas E Van Dyke - The Forsyth Institute, USA  
William Wade - Kings College London, UK  
Marvin Whiteley - Georgia Institute of Technology, USA  
David Williams - Cardiff University, UK  
Özlem Yilmaz - Medical University of South Carolina, USA  
Qing Yu - The Forsyth Institute, USA  
Egija Zuara - University of Amsterdam, The Netherlands  
Xuedong Zhou - Sichuan University, China

## Information for

[Authors](#)

[R&D professionals](#)

[Editors](#)

[Librarians](#)

[Societies](#)

## Opportunities

[Reprints and e-prints](#)

[Advertising solutions](#)

[Accelerated publication](#)

[Corporate access solutions](#)

## Open access

[Overview](#)

[Open journals](#)

[Open Select](#)

[Dove Medical Press](#)

[F1000Research](#)

## Help and information

[Help and contact](#)

[Newsroom](#)

[All journals](#)

[Books](#)

## Keep up to date

Register to receive personalised research and resources by email

Sign me up

[Copyright © 2022 Informa UK Limited](#) [Privacy policy](#) [Cookies](#) [Terms & conditions](#)

[Accessibility](#)



Taylor & Francis Group  
an **informa** business

Registered in England & Wales No. 3099067  
5 Howick Place | London | SW1P 1WG



## Journal of Oral Microbiology

[Submit](#)

[About](#)

[Browse](#)

[Subscribe](#)



## Journal of Oral Microbiology, Volume 14, Issue 1 (2022)



Volume 14, 2022

Vol 13, 2021

Vol 12, 20 



Issue

1



Download citations

Download PDFs



Browse by section (All)



Display order (Default)



## Issue In Progress

### Review Article



Review Article

**VBNC, previously unrecognized in the life cycle of *Porphyromonas gingivalis*?** 

A Progulske-Fox, SS Chukkapalli, H Getachew, WA Dunn & JD Oliver

Article: 1952838

**Published online:** 13 Jan 2022

1021

Views

0

CrossRef citations

0


Altmetric



### Original Article



Article

**Nanoparticle carrier co-delivery of complementary antibiofilm drugs abrogates dual species cariogenic biofilm formation *in vitro*** 

Guilherme Roncari Rocha, Kenneth R. Sims Jr., Baixue Xiao, Marlise I. Klein & Danielle S.W. Benoit

Article: 1997230

**Published online:** 25 Nov 2021

1340

Views

0

CrossRef citations

4

Altmetric



Article  
○ [Global proteomics of fibroblast cells treated with bacterial cyclic dinucleotides, c-di-GMP and c-di-AMP](#) □

Kenneth I. Onyedibe, Samira Elmanfi, Uma K. Aryal, Eija Könönen, Ulvi Kahraman Gürsoy & Herman O. Sintim

Article: 2003617

**Published online:** 29 Dec 2021

---

1370	0	1
Views	CrossRef citations	Altmetric

---



Article  
○ [The oral microbiome in alcohol use disorder: a longitudinal analysis during inpatient treatment](#) □

JJ Barb, KA Maki, N Kazmi, BK Meeks, M Krumlauf, RT Tuason, AT Brooks, NJ Ames, D Goldman & GR Wallen

Article: 2004790

**Published online:** 01 Dec 2021

---

971	0	4
Views	CrossRef citations	Altmetric

---



Article  
○ [Quantification of torque teno virus \(TTV\) DNA in saliva and plasma samples in patients at short time before and after kidney transplantation](#) □

Alexandre Mendes Batista, Matheus W. Caetano, Maria A. Stincarelli, Ana C. Mamana, Rodrigo Melim Zerbinati, Dmitry J. S. Sarmiento, Marina Gallottini, Rafael A. V. Caixeta, José Medina-Pestana, Bengt Hasséus, Louise Zanella, Tania R. Tozetto-Mendoza, Simone Gianecchini & Paulo H. Braz-Silva

Article: 2008140

**Published online:** 12 Dec 2021

---

734	0	1
Views	CrossRef citations	Altmetric

---



- Article  
**A genomic sequence of the type II-A clustered regularly interspaced short palindromic repeats (CRISPR)/CRISPR-associated system in *Mycoplasma salivarium* strain ATCC 29803** □

Harumi Mizuki, Yu Shimoyama, Taichi Ishikawa & Minoru Sasaki

Article: 2008153

**Published online:** 02 Jan 2022

---

936	0	1
Views	CrossRef citations	Altmetric

---



- Article  
**Development of antibacterial composite resin containing chitosan/fluoride microparticles as pit and fissure sealant to prevent caries** □

Chun-Cheng Lai, Chun-Pin Lin & Yin-Lin Wang

Article: 2008615

**Published online:** 27 Dec 2021

---

1162	0	0
Views	CrossRef citations	Altmetric

---



- Article  
**Thermoresponsive sol-gel containing probiotic's cell free supernatant for dental caries prophylaxis** □

Panithi Raknam, Neelam Balekar, Rawee Teanpaisan & Thanaporn Amnuaiakit

Article: 2012390

**Published online:** 19 Dec 2021

---

884	0	0
Views	CrossRef citations	Altmetric

---



- Article  
**Microtubule affinity regulating kinase 4 promoted activation of the NLRP3 inflammasome-mediated pyroptosis in periodontitis** □

Lulu Wang, Wenchen Pu, Chun Wang, Lang Lei & Houxuan Li

Article: 2015130

**Published online:** 27 Dec 2021

---

1358	0	2
Views	CrossRef citations	Altmetric

---



Article

○ [Arginine-induced metabolomic perturbation in \*Streptococcus mutans\*](#) □

Yudong Liu, Shanshan Liu, Qinghui Zhi, Peilin Zhuang, Rongxiu Zhang, Zhenzhen Zhang, Kai Zhang & Yu Sun

Article: 2015166

**Published online:** 07 Jan 2022

---

940	0	1
Views	CrossRef citations	Altmetric

---



Article

○ [Assessing the drug resistance profiles of oral probiotic lozenges](#) □

Yi Wang, Jingya Dong, Junyi Wang, Wei Chi, Wei Zhou, Qiwen Tian, Yue Hong, Xuan Zhou, Hailv Ye, Xuechen Tian, Rongdang Hu & Aloysius Wong

Article: 2019992

**Published online:** 07 Jan 2022

---

1828	0	
Views	CrossRef citations	Altmetric

---



Article

○ [Cetylpyridinium chloride promotes disaggregation of SARS-CoV-2 virus-like particles](#) □

Manuel Bañó-Polo, Luis Martínez-Gil, Manuel M. Sánchez del Pino, Alberto Massoli, Ismael Mingarro, Rubén León & Maria Jesus Garcia-Murria

Article: 2030094

**Published online:** 24 Jan 2022

---

7150 | 0  
Views | CrossRef citations | Altmetric

---



Article  
○ [Locational effects on oral microbiota among long-term care patients](#) □

Fa-Tzu Tsai, Ding-Han Wang, Cheng-Chieh Yang, Yu-Cheng Lin, Lin-Jack Huang, Wei-Yu Tsai, Chang-Wei Li, Wun-Eng Hsu, Hsi-Feng Tu & Ming-Lun Hsu

Article: 2033003

**Published online:** 14 Feb 2022

646 | 0  
Views | CrossRef citations | Altmetric

---



Article  
○ [Diversity of site-specific microbes of occlusal and proximal lesions in severe- early childhood caries \(S-ECC\)](#) □

Kausar Sadia Fakhruddin, Lakshman Perera Samaranayake, Rifat Akram Hamoudi, Hien Chi Ngo & Hiroshi Egusa

Article: 2037832

**Published online:** 13 Feb 2022

604 | 0  
Views | CrossRef citations | Altmetric

---



Article  
○ [Unraveling the binding microprocess of individual \*Streptococcus mutans\* cells via sucrose-dependent adhesion based on surface plasmon resonance imaging](#) □

Yuhao Guo, Bo Li, Tengfei Ma, Emily R. Moore, Huixu Xie, Chenzhou Wu & Longjiang Li

Article: 2038906

**Published online:** 17 Feb 2022

565 | 0  
Views | CrossRef citations | Altmetric

---



Article  
○ *Porphyromonas gingivalis* lipopolysaccharide induced RIPK3/MLKL-mediated necroptosis of oral epithelial cells and the further regulation in macrophage activation □

Fengxue Geng, Junchao Liu, Chengcheng Yin, Shuwei Zhang, Yaping Pan & Hongchen Sun

Article: 2041790

**Published online:** 27 Feb 2022

---

614	0	
Views	CrossRef citations	Altmetric

---



Article  
○ Microbiome profiles of non-responding and responding paired periodontitis sites within the same participants following non-surgical treatment □

SJ Byrne, D Chang, GG Adams, CA Butler, EC Reynolds, IB Darby & SG Dashper

Article: 2043595

**Published online:** 13 Mar 2022

---

311	0	
Views	CrossRef citations	Altmetric

---



Article  
○ MALDI-TOF mass spectrometry of saliva samples as a prognostic tool for COVID-19 □

Lucas C. Lazari, Rodrigo M. Zerbinati, Livia Rosa-Fernandes, Veronica Feijoli Santiago, Klaise F. Rosa, Claudia B. Angeli, Gabriela Schwab, Michelle Palmieri, Dmitry J. S. Sarmento, Claudio R. F. Marinho, Janete Dias Almeida, Kelvin To, Simone Gianecchini, Carsten Wrenger, Ester C. Sabino, Herculano Martinho, José A. L. Lindoso, Edison L. Durigon, Paulo H. Braz-Silva & Giuseppe Palmisano

Article: 2043651

**Published online:** 27 Feb 2022

---

683	0	
Views	CrossRef citations	Altmetric

---



Article  
○ [High iron-mediated increased oral fungal burden, oral-to-gut transmission, and changes to pathogenicity of \*Candida albicans\* in oropharyngeal candidiasis](#) □

Aparna Tripathi, Anubhav Nahar, Rishabh Sharma, Trevor Kanaskie, Nezar Al-Hebshi & Sumant Puri

Article: 2044110

**Published online:** 01 Mar 2022

---

373	0	
Views	CrossRef citations	Altmetric

---



Article  
○ [In vitro and in vivo anti-\*Candida\* activity of citral in combination with fluconazole](#) □

Katherine Miranda-Cadena, Cristina Marcos-Arias, Aitzol Perez-Rodriguez, Iván Cabello-Beitia, Estibaliz Mateo, Elena Sevillano, Lucila Madariaga, Guillermo Quindós & Elena Eraso

Article: 2045813

**Published online:** 02 Mar 2022

---

415	0	
Views	CrossRef citations	Altmetric

---



Article  
○ [Streptococcus mutans-associated bacteria in dental plaque of severe early childhood caries](#) □

Yixin Zhang, Jiakun Fang, Jingyi Yang, Xiaolei Gao, Liying Dong, Xuan Zheng, Liangjie Sun, Bin Xia, Na Zhao, Zeyun Ma & Yixiang Wang

Article: 2046309

**Published online:** 02 Mar 2022

---

627	0	
Views	CrossRef citations	Altmetric

---



Article  
○ [Lack of direct association between oral mucosal lesions and SARS-CoV- 2 in a cohort of patients hospitalised with COVID-19](#) □



Gabriela Schwab, Michelle Palmieri, Rodrigo M. Zerbinati, Dmitry J. S. Sarmento, Thais Reis, Karem L. Ortega, Italo T. Kano, Rafael A. V. Caixeta, Bengt Hasséus, Dipak Sapkota, Roger Junges, Simone Gianecchini, André L. F. Costa, Sumatra M. C. P. Jales, José A. L. Lindoso, Camila Barros Gallo & Paulo H. Braz-Silva

Article: 2047491

**Published online:** 10 Mar 2022

---

507	0	
Views	CrossRef citations	Altmetric

---



○ Article  
[Mapping bacterial diversity and metabolic functionality of the human respiratory tract microbiome](#) □

Leonardo Mancabelli, Christian Milani, Federico Fontana, Gabriele Andrea Lugli, Chiara Tarracchini, Francesca Turrone, Douwe van Sinderen & Marco Ventura

Article: 2051336

**Published online:** 16 Mar 2022

---

375	0	
Views	CrossRef citations	Altmetric

---



○ Article  
[Maturation of the oral microbiota during primary teeth eruption: a longitudinal, preliminary study](#) □

He Xu, Bijun Tian, Weihua Shi, Jing Tian, Wenjun Wang & Man Qin

Article: 2051352

**Published online:** 16 Mar 2022

---

346	0	
Views	CrossRef citations	Altmetric

---



○ Article  
[Comparative analysis of the oral microbiome of burning mouth syndrome patients](#) □

Byeong-Min Lee, Ji Woon Park, Jung Hwan Jo, Bumjo Oh & Gehoon Chung

Article: 2052632

Published online: 20 Mar 2022

---

332	0	
Views	CrossRef citations	Altmetric

---



○ Article  
[Profile of the oral microbiota from preconception to the third trimester of pregnancy and its association with oral hygiene practices](#) □

Xuena La, Hong Jiang, An Chen, Huajun Zheng, Liandi Shen, Weiyi Chen, Fengyun Yang, Lifeng Zhang, Xushan Cai, Hongfang Mao & Lu Cheng

Article: 2053389

Published online: 20 Mar 2022

---

306	0	
Views	CrossRef citations	Altmetric

---



○ Article  
[The reciprocal interaction between fluoride release of glass ionomers and acid production of \*Streptococcus mutans\* biofilm](#) □

Jian-Na Cai, Hyeon-Mi Choi, Kwang-Yeob Song & Jae-Gyu Jeon

Article: 2055267

Published online: 23 Mar 2022

---

233	0	
Views	CrossRef citations	Altmetric

---



○ Article  
[Anti-biofilm activities of coumarin as quorum sensing inhibitor for \*Porphyromonas gingivalis\*](#) □

Zhiyan He, Wei Jiang, Yiting Jiang, Jiachen Dong, Zhongchen Song, Jianrong Xu & Wei Zhou

Article: 2055523

Published online: 29 Mar 2022

---

392	0	
Views	CrossRef citations	Altmetric

---



Article  
○ **Nicotinamide could reduce growth and cariogenic virulence of *Streptococcus mutans*** □

Yongwang Lin, Tao Gong, Qizhao Ma, Meiling Jing, Ting Zheng, Jiangchuan Yan, Jiamin Chen, Yangyang Pan, Qun Sun, Xuedong Zhou & Yuqing Li

Article: 2056291

**Published online:** 23 Mar 2022

---

285	0	
Views	CrossRef citations	Altmetric

---



Article  
○ **Maintaining oral health for a hundred years and more? - An analysis of microbial and salivary factors in a cohort of centenarians** □

Caroline Sekundo, Eva Langowski, Diana Wolff, Sébastien Boutin & Cornelia Frese

Article: 2059891

**Published online:** 04 Apr 2022

---

198	0	
Views	CrossRef citations	Altmetric

---



Article  
○ **The suppression effect of SCH-79797 on *Streptococcus mutans* biofilm formation** □

Lingjun Zhang, Yan Shen, Lili Qiu, Fangzheng Yu, Xiangyu Hu, Min Wang, Yan Sun, Yihuai Pan & Keke Zhang

Article: 2061113

**Published online:** 18 Apr 2022

---

54	0	
Views	CrossRef citations	Altmetric

---



○ Article  
**Short- term effect of probiotic *Lactobacillus reuteri* consumption on the salivary microbiome profile of subjects undergoing orthodontic treatment with fixed appliances** □

Armelia Sari Widyarman, Nadeeka S. Udawatte, Moehamad Orliando Roeslan, Muhammad Ihsan Rizal, Mario Richi, Joko Kusnoto & Chaminda Jayampath Seneviratne

Article: 2067103

**Published online:** 25 Apr 2022

---

0	0	
Views	CrossRef citations	Altmetric

---



## Latest articles

Most read articles

Most cited articles

Information for

Authors

R&D professionals

Editors

Librarians

Societies

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Open access

Overview

Open journals

Open Select

Dove Medical Press

F1000Research

Help and information

Help and contact

Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources by

email

Sign me up



Copyright © 2022 Informa UK Limited [Privacy policy](#) [Cookies](#) [Terms & conditions](#)

[Accessibility](#)



Taylor & Francis Group  
an **informa** business

Registered in England & Wales No. 3099067  
5 Howick Place | London | SW1P 1WG



## Short- term effect of probiotic *Lactobacillus reuteri* consumption on the salivary microbiome profile of subjects undergoing orthodontic treatment with fixed appliances

Armelia Sari Widyarman, Nadeeka S. Udawatte, Moehamad Orliando Roeslan, Muhammad Ihsan Rizal, Mario Richi, Joko Kusnoto & Chaminda Jayampath Seneviratne

To cite this article: Armelia Sari Widyarman, Nadeeka S. Udawatte, Moehamad Orliando Roeslan, Muhammad Ihsan Rizal, Mario Richi, Joko Kusnoto & Chaminda Jayampath Seneviratne (2022) Short- term effect of probiotic *Lactobacillus reuteri* consumption on the salivary microbiome profile of subjects undergoing orthodontic treatment with fixed appliances, Journal of Oral Microbiology, 14:1, 2067103, DOI: [10.1080/20002297.2022.2067103](https://doi.org/10.1080/20002297.2022.2067103)

To link to this article: <https://doi.org/10.1080/20002297.2022.2067103>



© 2022 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



[View supplementary material](#)



Published online: 25 Apr 2022.



[Submit your article to this journal](#)



[View related articles](#)



[View Crossmark data](#)

## Short- term effect of probiotic *Lactobacillus reuteri* consumption on the salivary microbiome profile of subjects undergoing orthodontic treatment with fixed appliances

Armelia Sari Widyarman <sup>a</sup>, Nadeeka S. Udawatte <sup>b</sup>, Moehamad Orliando Roeslan <sup>c</sup>,  
Muhammad Ihsan Rizal <sup>c</sup>, Mario Richi <sup>d</sup>, Joko Kusnoto<sup>e</sup> and Chaminda Jayampath Seneviratne<sup>b,f</sup>

<sup>a</sup>Department of Microbiology, Faculty of Dentistry, Trisakti University, Jakarta, Indonesia; <sup>b</sup>Singapore Oral Microbiomics Initiative, National Dental Research Institute Singapore (NDRIS), National Dental Centre Singapore, Oral Health ACP, Duke NUS Medical School, Singapore; <sup>c</sup>Department of Oral Biology, Faculty of Dentistry, Trisakti University, Jakarta, Indonesia; <sup>d</sup>MICORE Laboratory, Faculty of Dentistry, Trisakti University, Jakarta, Indonesia; <sup>e</sup>Department of Orthodontics, Faculty of Dentistry, Trisakti University, Jakarta, Indonesia; <sup>f</sup>School of Dentistry, The University of Queensland, Brisbane, Australia

### ABSTRACT

**Objective:** This prospective clinical study aim was to analyze the effect of the probiotic *Lactobacillus reuteri* Prodentis lozenges on salivary microbiome of subjects wearing fixed orthodontic appliances.

**Methods:** Saliva samples were collected prior to consumption and 14th-day post probiotic lozenges consumption (n=40, age 18–23). Oral hygiene index-score (OHI-S) and papilla bleeding index (PBI) were recorded. The salivary microbiome was profiled by next-generation sequencing using the V3-V4 region of 16S-rRNA. Microbial composition, diversity and taxonomic biomarkers were analysed in comparison to probiotic intervention and the clinical characteristics of the cohort using standard bioinformatics tools.

**Results:** The diversity and bacterial community structures did not change significantly in salivary microbiome of periodontally healthy subjects during short-term probiotic intervention. Probiotic consumption correlated with reduction of OHI and PBI scores (50% reduction of scores,  $P < 0.001$ ). The reduction of clinical indices was evident in conjunction with significantly reduced abundance of oral pathogens, such as *Porphyromonas pasteri*, *Treponema sp.*, *Fretibacterium fastidiosum*, *Kingella oralis* and *Propionibacterium acnes*.

**Conclusion:** Short-term probiotic intervention helped maintaining good oral health in patients undergoing fixed orthodontic therapy. Although overall oral microbiome structure remained largely unchanged, a significant alteration in the abundance of health and disease-associated species highlighted the beneficial effect of probiotic.

### ARTICLE HISTORY

Received 10 February 2022  
Revised 7 April 2022  
Accepted 14 April 2022

### KEYWORDS





Probiotic; *Lactobacillus reuteri*; oral microbiome; fixed orthodontic appliance


## Background

Fixed orthodontic treatment is a risk factor for the development of various oral diseases, such as dental caries, periodontal disease, and candidiasis. The presence of brackets, bands, and arch wire complicates oral hygiene practices and increases the accumulation of dental plaque. Hence, patients wearing a fixed orthodontic treatment are at higher risk for developing oral diseases [1]. Therefore, orthodontic patients need supplementary oral hygiene measures in addition to personal mechanical oral hygiene practices. Probiotics, which are defined as ‘living microorganisms which, when administered in adequate amounts, confer a health benefit on the host’ by the Food and Agriculture Organization of the United Nations/World Health Organization (FAO/WHO) have been shown to confer various health benefits to humans, including improvement of oral health [2].

*Lactobacillus reuteri* is a probiotic bacterium known to confer various health benefits to humans [3]. It produces the antimicrobial substances reuterin and reutericyclin, which are active against a wide range of pathogenic organisms [4]. Previously, we performed a pilot study on the beneficial health effects of lozenges containing the probiotic *L. reuteri* on a few orthodontic patients [5]. We found that consumption of probiotics for 2 weeks significantly reduced the number of pathogenic bacteria in the patients’ saliva. Moreover, other studies using appropriate *in vitro* and *in vivo* models have demonstrated that *L. reuteri* is able to modulate host inflammatory responses [6]. However, clinical studies that examine the effect of the probiotic *L. reuteri* on the oral microbiome are sparse in the literature.

To our knowledge, there is no information available regarding how *L. reuteri* probiotics regulate the oral microbiome in periodontally healthy subjects

**CONTACT** Armelia Sari Widyarman  [armeliasari@trisakti.ac.id](mailto:armeliasari@trisakti.ac.id)  Department of Microbiology, Faculty of Dentistry, Trisakti University, Indonesia; Chaminda Jayampath Seneviratne  [jaya.seneviratne@uq.edu.au](mailto:jaya.seneviratne@uq.edu.au)  School of Dentistry, The University of Queensland, Australia

 Supplemental data for this article can be accessed here.

© 2022 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

using fixed orthodontic appliances. Considering this research gap, we aimed to analyze the short-term effect of the consumption of the probiotic *L. reuteri* Prodentis lozenges on the salivary microbiome profile of subjects wearing fixed orthodontic appliances.

## Materials and methods

### Cohort characteristics

The subjects undergoing fixed orthodontic therapy were included in this open-labelled prospective clinical trial. Ethical approval for the study was obtained from the Institutional Review Board, Faculty of Dentistry, Universitas Indonesia (approval number 100,701,020). The inclusion criteria included an age above 18 years, undergoing orthodontic treatment with a fixed appliance of a minimum of 1 year and no consumption of probiotics or antibiotics within the past 3 months. Exclusion criteria included subjects who had systemic diseases, such as hypertension or diabetes, or were on systemic drugs *i.e.* anti-hypertensives, analgetics, hormonal drugs, sedatives, anti-seizure medication and subjects with features of severe periodontal diseases and who were allergic to probiotics. The lozenges containing *L. reuteri* Prodentis were obtained from BioGaia, Stockholm, Sweden. BioGaia Prodentis lozenges (800 mg of one lozenge contained a minimum of  $2 \times 10^8$  live *L. reuteri* Prodentis) were a food supplement for oral health containing the patented lactic acid bacterium *Limosilactobacillus reuteri* (formerly known as *Lactobacillus reuteri*) Prodentis. They also contained a strain combination of *L. reuteri* DSM 17938 and *L. reuteri* ATCC PTA 5289 that was expected to help maintain a good oral health.

To standardize the oral hygiene practice, each participant was given a toothpaste and a toothbrush during the study period. The participants were given oral hygiene instructions and were requested to brush their teeth twice a day. Each subject was given one lozenge per day, containing the probiotic *L. reuteri* for 14 days to be taken once a day after brushing teeth, followed by breakfast. Saliva samples were collected from the subjects on the day of the recruitment prior to the consumption of probiotic lozenges on the 14th-day post-consumption of the probiotics.

In addition, clinical data pertaining to the oral hygiene index score (OHI-S) and papilla-bleeding index (PBI) were recorded at each visit. Previously, it had been found that consumption of probiotic *L. reuteri* Prodentis over 14 day-duration is able to exert considerable impact on reducing clinical parameters (Periodontal Index-PI, OHI-S and Gingival index-GI) in healthy test subjects compared to a placebo control [7,8].

### Sample collection and DNA extraction

Prior to sample collection, subjects had to take a rapid antigen test for SARS-CoV-2 detection due to the COVID-19 pandemic (as of 30 September 2020) and the result of the test had to be negative for taking part in the study. Following this confirmation, sample specimens from the subjects were collected with a cover of a complete set of personal protective equipment. A standard method was used for the saliva sample collection [9]. In brief, the subject was made to sit quietly, bending the head forward and opening the mouth to allow the saliva to drip passively from the lower lip. Saliva was collected using sterile, pre-weighted tubes. Subjects were requested to refrain from food for 2 h prior to the sample collection. Following the collection, the saliva samples were stored at  $-80^{\circ}\text{C}$  prior to DNA extraction.

Microbiome profiling of the saliva samples was performed using a previously established protocol with some modification [10]. In brief, DNA was extracted from the samples using the QIAamp® DNA Mini Kit (Qiagen, Hilden, Germany) per the manufacturer's protocol. Extracted DNA was subjected to a quality check using Nanodrop™ 2000 (Thermo Fisher, Waltham, MA) for its purity and its concentration of gDNA using a Qubit 3.0 fluorometer (Thermo Fisher, Waltham, MA).

### 16S rRNA amplicon library preparation and sequencing

Polymerase chain reaction (PCR) amplifications had been finished for library preparation prior to sequencing as previously described [11]. Primers targeted for the V3-V4 region of 16S rRNA and modified with Illumina overhang adapters had been amplified by using PCR and the primers: F (5' - TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCTACGGGNGGC WGCAG-3') and R (5' GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGG ACTACHVGGGTATCTAATCC -3'). The PCR amplification was done as described previously [10]. The amplified library was subjected to purification using AMPure XP beads (Beckman Coulter Genomics, MA), visualized via agarose gel electrophoresis and quantified by using the Qubit 3.0 Fluorometer (Thermo Fisher Scientific, Waltham, MA). The amplified library after the second purification using AMPure XP Beads was eluted, normalized, and pooled to the recommended concentration as per the manufacturer's protocol (80 pM). An internal 5% spike-in control using the Phi-X manage v3 library (Illumina, San Diego, CA) was pooled together with the normalized amplicon library. Sequencing was performed using the iSeq 100 instrument with the i1 v2 reagent (Illumina, San Diego, CA) with the



parameters of  $2 \times 150$  cycles paired-end reads, with 10 cycles for every index, in line with the manufacturer's protocols.

### Data analysis

Sequencing runs were first analysed in the Illumina's Sequencing evaluation Viewer version 2.4.7 for analytical run quality controls assessments [12–14]. Pre-processing of the raw sequencing data and the Quality Control (QC) was assessed as previously described [15]. The 16S metagenomics workflow based on the Local Run Manager v2.0 used the Illumina-curated version of the Greengenes database ([greengenes.secondgenome.com/downloads/data-base/13\\_5](http://greengenes.secondgenome.com/downloads/data-base/13_5)) for taxonomic classification of 16S rRNA centred amplicon reads. It generated a total of 1,282 operational taxonomic units (OTUs) clustered at 97% identification with an open-reference OTU picking approach. The resulting accumulation curves showed reasonable sequence saturation for subsequent analysis (Supplementary Figure 1).

Alpha diversity (Chao1, Shannon, ACE), along with rarefaction curves, was evaluated and visualized using the MicrobiomeAnalystR Platform and the 'phyloseq' and 'ggplot' and 'microbiomeseq' [16,17] codes obtained from GitHub (<https://github.com/xia-lab/MicrobiomeAnalystR>). The microbial community structure was assessed by beta diversity with the respective algorithms implemented in the MicrobiomeAnalystR package using Principal Coordinate Analysis (PCoA) and the Bray–Curtis's dissimilarity indexes retrieved from sample pairwise comparisons based totally on weighted Unifrac distances. Significance of the community structure analysis was performed using the permutational multivariate analysis of variance (PERMANOVA) test [18].

DESeq2 [19] analysis was run under default settings and *q* values were calculated using the Benjamini–Hochberg procedure to control for false discovery rates applying the MicrobiomeAnalystR. Based on both significance and biological relevance, differentially abundant microbial biomarkers were characterized using the linear discriminant analysis effect size (LEfSe) method with default parameters and an LDA log score threshold set to 2 with a *P* value less than 0.05 considered as statistically significant [20,21]. Species with a *P* value <0.1 on the univariate analysis with probiotic intervention were selected for further analysis. The relative abundance differences in the top 27 bacterial species between pre- and post-probiotic cohorts were compared by Welch's *t*-test using STAMP (v2.1.3) [22]. Further, the random forest (RF) algorithm was applied to determine the proper number of microbial predictors in random forest classifiers by maximizing the area under the curve of the receiver operator characteristic (AUC) with the default parameters of the

'RandomForest' (RF) package in R (v. 4.6.1). To avoid over-fitting of the data in the model, 5-time and 10-fold cross-validations were made. The resulting model was subsequently used for the validation cohort [23,24].

To identify key members of the microbiome (based on relative abundance) associated with 50% reduction of the oral hygiene index score from the baseline value after probiotic intake, the Boruta algorithm [Boruta package for R (v. 5.1)] was used to perceive species with the highest predictive power [25]. The relative abundance >0.1% and presence in >50% of the samples in at least one group were used for the above analyses unless otherwise specified.

### Statistical analyses

Statistical analyses were carried out using the GraphPad Prism software (version 8) and the R package (<http://www.r-project.org/>). The Mann–Whitney rank sum test or the Kruskal–Wallis test was used to evaluate variations among groups in the diversity indices of the oral microbiota. False discovery rate (FDR) values were used for the evaluation of the Benjamini–Hochberg method to control for multiple testing – adjusted *P* values lower than 0.05 were considered as statistically significant [19]. The analysis was confined to species with a prevalence greater than 10% and a maximum proportion (relative abundance) greater than 0.002. An FDR-adjusted *P* value (or *Q*-value) <5% was considered as significant.

The Boruta algorithm was used to identify the most predictive taxonomic biomarker based on the importance values determined by RF, indicated as 'confirmed' in the Boruta output [25]. The importance value pertaining to a taxon was calculated by RF based on the loss of accuracy by random permutation of the abundance profile of the taxon. To assess whether the importance was significant, the Boruta algorithm compared the observed importance to those produced by the spiked-in 'shadow' taxa, which were randomized versions of original taxa; hence, this algorithm typically gave more strength to identify taxa that jointly predicted a phenotype [26].

## Results


### Effect of probiotic intake on the clinical parameters

The mean age of the participants (*n* = 20; male – 4, female – 16) was 21 years (SD  $\pm$  1.59) across all samples (Figure 1A). Consumption of probiotic *L. reuteri* lozenges for 14 days improved the oral health parameters of the subjects as measured by the reduction of the OHI-S (*P* < 0.0001) and PBI scores (*P* < 0.0056) (Figure 1B).


### Effect of probiotics on taxonomic composition and complexity of the salivary microbiome

We then taxonomically profiled each group based on the relative abundances of microbial clades at distinctive taxonomic tiers. Five phyla made up >1% of the community in at least one sample with two phyla naming, *Firmicutes* and *Proteobacteria*, dominated the groups with relative abundance (33–30% and 33–26% in the pre and post-probiotic groups, respectively) among the top 10 predominant phylum groups.

*Actinobacteria* and *Bacteroidetes* were present in all samples, however to a lesser extent (16–11% and 22–10% in the pre- and post-probiotic groups, respectively) (Figure 2A). Lower-abundance phyla such as *Spirochaetes* were increased in post-probiotic organisms, and the so-called ‘environmental’ phyla such as SR1 were detected in similar level of consistency. In the top 15 genus level, *Streptococcus* and *Haemophilus* made up for almost 28% of the total reads, with 27% and 28% in the pre- and post-probiotic groups, respectively. *Rothia*, *Actinomyces*, *Fusobacterium*, *Neisseria*,




**Figure 1.** Demographic and clinical evaluation of individuals with good oral hygiene using orthodontic-fixed appliances prior and post-probiotic *L. reuteri* lozenges supplementation. **(A)** Baseline characteristics of the study participants in both intervention groups. **(B)** Two clinical outcomes were evaluated before and after 14 days of probiotic intervention. The Simplified Oral Hygiene index (OHIS) for determining the soft sediments classified with microbial signatures designed no plaque (0), mild [1- (1/3) covered with soft plaque], moderate [2- (1/3-2/3) covered with soft plaque], bad [3- (> 2/3) covered with soft plaque] according to the Green – Vermillion – Hirschman index. The Papillary Bleeding Index (PBI) used to determine the presence or absence of inter- dental plaque; designed no bleeding (0), only one bleeding point appearing [1], several isolated bleeding points or a small blood area appearing [2], inter dental triangle filled with blood soon after probing [3], and profuse bleeding when probing led to blood spreads towards the marginal gingiva [4]. All subjects showed significant oral health improvement, as measured by the reduction of the OHIS and PBI scores ( $P < 0.001$ ), by the end of the probiotic supplementation.



**Figure 2.** Taxonomic composition of the salivary microbiome before and after probiotic supplementation. **(A)** The top 10 relative abundances of the microbial phyla highlights the major role played by *Firmicutes* and *Proteobacteria* (green and brown) in the microbiome, and to a less extent by *Actinobacteria* and *Bacteroidetes* (light brown and purple). **(B)** At the top 15 genus level, *Haemophilus*, *Streptococcus*, *Veillonella*, *Rothia* and *Neisseria* (maroon, blue, dark green, light green and light purple) are shown to be the most abundant clades. **(C)** Out of the genera accounting for less than 1% of the samples which are grouped in the ‘Other’ category, *Schlegella*, *Bacillus*, *Pseudoxanthomonas*, *Lactobacillus*, *Microbacterium* and *Anoxybacillus* were significantly increased after probiotic intake. Letters indicate statistical differences between samples are \* $P < 0.05$ , \*\* $P < 0.01$ , and \*\*\* $P < 0.001$ .

*Veillonella*, *Prevotella*, *Selenomonas* and *Leptotrichia* were also detected in all samples with uneven quantitative distributions (Figure 2B). At the species level, *Haemophilus parainfluenzae*, *Rothia dentocariosa*, *Leptotrichia hongkongensis*, *Haemophilus haemolyticus* and *Neisseria elongata* were the main species components in both the pre- and post-probiotic groups (relative abundance >3%). Overall, taxonomic phyla and genus abundance distributions of these dominant oral microbes (top 10 taxa) were minimally influenced by short-term probiotic intervention. Out of lesser abundant genera (relative abundance <1%), *Schlegelella* ( $P = 0.00043$ ), *Bacillus* ( $P = 0.0011$ ), *Pseudoxanthomonas* ( $P = 0.003$ ), *Lactobacillus* ( $P = 0.037$ ), *Microbacterium* ( $P = 0.043$ ), and *Anoxybacillus* ( $P = 0.037$ ) were significantly increased following the probiotic intervention (Figure 2C).

The microbial community composition of the saliva samples at baseline and 2-week post-consumption was analysed. There was no significant difference in the alpha diversity (Shannon –  $P$  value 0.698; [Mann–Whitney]) and evenness (Chao1 –  $P$  value 0.881; ACE –  $P$  value 0.989; Simpson –  $P$  value 0.62 [Mann–Whitney]) of the post-consumption samples compared to the baseline (Figure 3A). Post-consumption saliva samples had a slightly higher alpha diversity than pre-probiotic intake samples, although the difference was not statistically significant. To understand whether the quantitative structure of the microbiome changed, PCoA on the taxonomic profiles using the weighted UniFrac measure was performed. Ordination analysis showed no obvious clustering by pre-defined groups, suggesting that the overall community structure of the saliva microbiome was not stricken by short-term



**Figure 3.** Structural composition and diversity comparison of the salivary microbiome before and after probiotic intervention. The overall structure of the saliva microbiome is not affected by short-term probiotic intake. (A)  $\alpha$ -diversity boxplot (Shannon, Chao1 and Simpson-reciprocal indices) of the salivary microbiome between pre- and post-probiotic cohorts. Boxes represented the 25th to 75th percentile of the distribution; the median was shown as a thick dot in the middle of the box; whiskers extend between the 25th and 75th percentiles. (B) No apparent clustering trend can be noticed ( $P > 0.05$ ) in the PCoA space when the non-quantitative unweighted UniFrac measure is used as beta-diversity, suggesting that a change in the relative abundances of the main components of the salivary microbiome is not the driving feature characterizing between probiotic and respective baseline samples. Species with a  $P$  value < 0.1 on the univariate analysis with probiotic intervention were selected to define the top most abundantly differentiated across the study cohort. (C) Heatmap of the top 27 abundant species differentiates between probiotic and baseline samples. The color intensity in each box indicates the relative percentage of species in each sample. Relative abundance data were z-scored normalized by row. To show the distribution of abundant OTUs, the relative abundances of OTU data were normalized to have a mean of 0 and a standard deviation of 1 (z-score normalization). (D) Random Forest algorithm with 5 times 10-fold-cross validation used to identify a minimal set of bacterial species that maximally differentiated between pre- and post-probiotic intervention. *Schlegelella aquatica*, *Microbacterium mitrae*, *Microbacterium ginsengisoli*, and *Streptococcus parasuis* ranked the predominant difference between two cohorts under probiotic intervention. (E)  $P < 0.01$  to detect species with the most statistically significant differences filtered using a q-value of 0.05 and effective size of 0.05 threshold in STAMP illustrating significant bacterial species in variations of different proportions using Welch's t-test. See also Supplementary Figure 2 for LefSe results.



probiotic intake (PERMANOVA  $P$  value  $<0.688$ ) (Supplementary Table 1) (Figure 3B).

### Increased complexity of the salivary microbiome following short-term probiotic consumption

We performed a linear discriminant analysis effect size (LEfSe) to determine which microbial clades, if any, could explain the slight increase in the diversity of the salivary microbiome following short-term probiotic intervention. The LEfSe analyses identified 12 species in the salivary microbiome whose over-abundance discriminated between pre- and post-probiotic samples (Figure 3 B and C) (full details are given in Supplementary Table 2 and Supplementary Figures 2 and 3). Pre-probiotic samples had a significant enrichment of species (LDA log score at a threshold of 3.0): *Schlegelella aquatica* (Proteobacteria), *Streptococcus parasuis* (Firmicutes), *Anoxybacillus pushchinoensis* (Firmicutes), *Anoxybacillus bogrovensis* (Firmicutes), *Pseudoxanthomonas taiwanensis* (Proteobacteria). On the other hand, *Sediminibacterium goheungense* (Bacteroidetes) and *Microbacterium ginsengisoli* (Actinobacteria) OTUs gave significance at the same effect size threshold for discriminating samples with probiotic consumption. Further analysis using 10-cross validation RF modeling (Figure 3D) and STAMP methods (Figure 3E) showed that few of these identified species, *S. aquatica*, *Microbacterium mitrae*, *M. ginsengisoli* and *S. parasuis*, reflected the predominant difference between pre- and post-probiotic cohorts under the short-time probiotic intervention.

We used the Boruta feature selection algorithm coupled with the RF classifier to examine which differential microbiome markers (predictors) at the species level accounted for 50% reduction of observed clinical indexes *i.e.* OHIS and PBI scores (response variables) with probiotic intervention. As evidenced by a decreased predictive mean square error compared to a random guess, a predictive model utilized 18 species with significant predictive power substantiated by LEfSe analysis (Figure 4 C and 4D) (Supplementary Table 3). Among the identified microbiome markers, there was a reduction in disease-associated pathogens, such as *P. pasteri*, *Treponema* sp. and *Prevotella* sp., which correlated with the 50% reduction of the OHIS index score group (OHIS50-Y/Yes) (Figure 4G). Based on feature selection by Boruta, over-abundance of *M. ginsengisoli*, *S. canadensis* and *A. suimastitidis*, and reduced relative abundance of *F. fastidiosum* were highly discriminative for the 50% reduction of the OHIS index score with probiotic intervention (Supplementary Figure and Table 3) (Figure 4C). On the other hand, by both LEfSe and Boruta feature selection; overabundance of *C. leadbetteri* and decreased abundance of *K. oralis*, *P. acnes*,

*S. aquatica*, and *A. bogrovensis* were identified in the 50% reduction of the PBI index score group by probiotic intake (PBI50-Y/Yes) (Supplementary Figure and Table 3) (Figure 4D). These species may independently be useful for predicting the 50% reduction of the PBI index score with probiotic intervention. The key species affected by probiotic intervention according to the clinical parameters are presented in heatmaps (Figure 4F and 4G). Moreover, the chord diagram was used to delineate species influenced by different clinical states before and after probiotic intake (Figure 4E). The chord diagram was divided into five sections representing the two clinical states of PBI and OHI-S before and after probiotic intake. Key species identified in two or more clinical states are indicated by arcs (chords) connecting the different clinical states.

Re-analysis of the overall salivary microbiome profiles using only the foregoing species with significant predictive power yielded some interesting results. PERMANOVA analysis significantly differentiated the salivary microbiome composition following probiotic intervention [between OHIS50-N/No and OHIS50-Y/Yes groups ( $P = 0.008$ ) and PBI50-N/No and PBI50-Y/Yes ( $P = 0.001$ )] (Figure 4 A and B).


### Persistence of the *Lactobacillus* genus in the salivary microbiome after probiotic intake

The *Lactobacillus* strain present in the probiotic product (*L. reuteri*) was not detected in the probiotic-intake samples. However, we observed the presence of a significantly higher-abundance OTU attributable to the *Lactobacillus* genus in the post-probiotic samples (Figure 2C).

The foregoing results suggested that the short course of the probiotic *L. reuteri* lozenges could potentially modulate the salivary microbiome, albeit the effect was limited in both the breadth of the change and in its temporal persistence.

### Discussion

There has been some evidence from clinical and fundamental studies suggesting that fixed orthodontic appliances affect the oral health of the patients [27,28]. The effect of probiotics on various microbiological and clinical parameters in fixed orthodontic patients has produced conflicting evidence [1,29–35]. The present study evaluated the effect of oral probiotics on the salivary microbiome of periodontally healthy subjects under orthodontic treatment with fixed appliances using a prospective interventional study. The oral health of the subjects was significantly improved following 14 days' intake of the probiotic *L. reuteri* Prodentis lozenges. Although the overall diversity or evenness of the salivary microbiome did



**Figure 4.** Increased complexity of the salivary microbiome in conjunction with 50% reduction of the oral hygiene index scores in subjects with fixed orthodontic appliances. **(A)** and **(B)** Apparent clustering in structural salivary composition observed when coupled with 50% reduction of the oral hygiene index scores (OHIS and PBI) after probiotic intervention. **(C)** and **(D)** Significant differentially abundant taxonomic biomarkers contribute to the change in the microbiome composition identified by LEfSe analysis. **(E)** The chord diagrams show the key taxa identified by the five different strategies (Baseline, post-probiotic; OHIS50-Y, OHIS50-N, PBI50-Y and PBI50-N). Key species identified in two or more clinical states are indicated by arcs (chords) connecting the different clinical states. **(F)** and **(G)** Heatmap of most abundant species among the OHIS50 groups differentiates between probiotic and baseline samples. Species highlighted in dotted blue squares are distinguished to have a strong predictive power for 50% reduction of both the OHIS and PBI index scores with probiotic intervention identified by the Machine learning Boruta algorithm coupled with the Random Forest classifier. See also Supplementary Figure 4. Abbreviations: PRPT- Pre-Probiotic group, POPT- Post-Probiotic group, OHIS50-N/No and PBI50-N/No – baseline and no change of OHIS and PBI index score groups, OHIS50-Y/Yes and PBI50-Y/Yes – 50% reduction of OHIS and PBI index score groups in probiotic intake.

not change significantly, there were minor changes in the abundance of disease-associated species following probiotic intervention.

Fixed orthodontic treatment may pose a significant challenge to maintain oral health of the patients as the appliance traps dental plaque biofilm leading to increased level of potential oral pathogens [36,37]. The use of probiotics as an adjunct therapy has been suggested for various diseases including periodontitis. Overall, the administration of probiotics, either as an adjunct to mechanical plaque control or as a single intervention, has yielded conflicting results [38]. Some studies have proven the advantages of probiotic consumption in reducing the extent of gingival

inflammation and the amount of plaque [39,40]. In contrast, other studies have failed to exhibit additional clinical benefits [41–43]. Moreover, several investigations documented the varying impact of long-term probiotic intervention (maximum up to 1 month) on maintaining gingival health in orthodontic patients, some with beneficial improvements [44,45] while some without improvement [30,31]. There are ample evidence in the literature supporting the beneficial effects of oral *L. reuteri* such as its capability: (i) to inhibit plaque accumulation [46], (ii) to reduce cariogenic *mutans* streptococci [47,48], (iii) to improve clinical parameters when used as an adjunct therapy for periodontitis [7,8,46] and (iv) to exert beneficial

immunomodulatory and anti-inflammation effects [46,49]. *L. reuteri* has been considered as a potential probiotic adjunct together with non-surgical periodontal therapy for periodontitis patients [7,8,50]. In our study, we observed reduced gingival bleeding with improved PBI scores and overall oral hygiene improvement by the OHIS index in periodontally healthy individuals undergoing fixed appliances for orthodontic treatment after intake of the probiotic *L. reuteri* Prodentis lozenges. The observed difference in the improved oral health may be attributed to the short-term probiotic intervention. However, according to a recently published review, no consensus can be reached on the significant effect of oral probiotics on gingival inflammation and enamel decalcification in patients undergoing treatment with fixed orthodontic appliances [29]. Therefore, in the presented study, we investigated whether the impact of probiotic would be more pronounced by defining our study cohort with respect to 50% reduction of the oral clinical state from their respective baseline score.

In this context, there were some interesting findings with regard to the change in key species associated with the salivary microbiome. Two weeks' probiotic intake reduced the disease-associated pathogens, such as *Porphyromonas* sp., *Treponema* sp., *Prevotella* sp. and *F. fastidiosum*, which correlated with the 50% reduction of the OHIS index score group (OHIS50-Y), suggesting improvement of oral health. Moreover, probiotic interventions decreased the abundance of the periodontal pathogens *K. oralis* and *P. acnes*, which correlated with the 50% reduction of the PBI index score group (PBI50-Y). Hence, probiotic intervention considerably eliminated the dysbiotic oral microbiome by reducing disease-associated pathogens and enriching health-associated species. *K. oralis* is normally found in small numbers in the oral cavity in dental plaque biofilm [51]. It has been reported that the species increased significantly in periodontitis *i.e.* more than 5% of the total microbiota in periodontitis sites compared to the meagre number (0.4%) in periodontally healthy individuals. Additionally, *P. acnes* and *F. fastidiosum* have been shown to be associated with gingivitis [52] and periodontitis [53], respectively. Significant reduction of these species indicated the potential modulatory activity of probiotic intake. Moreover, *Capnocytophaga* species, known to be associated with healthy periodontium, increased following probiotic intervention [54].

Fixed orthodontic appliance could be considered as a condition that may significantly change the local micro-environment in dental plaque biofilm, consequently resulting in an altered salivary microbiome [55]. Previous studies have shown significant dysbiosis in both the saliva microbiome and supragingival plaque microbiome with substantial replacement of anaerobic

bacteria in patients wearing fixed orthodontic appliances [36,37]. In the present study, we observed the presence of certain facultative anaerobes like *S. parasuis*, *Anoxybacillus* sp., *Bacillus thermoamylovorans* and mesophilic anaerobes like *M. mitrae* in the salivary microbiome of patients under fixed orthodontic treatment. Probiotic intervention reduced the proportion of these taxa. However, the absence of a placebo control group in the present study is a major limitation for a comprehensive comparison. Moreover, we did not detect the presence of the administered probiotic strain *L. reuteri* in the post-intervention salivary microbiome. However, this does not exclude the probability that this probiotic strain colonizes in dental plaque biofilms. Therefore, future studies should aim at analyzing both saliva and dental plaque samples from the subjects. On the other hand, according to the key-stone plaque hypothesis, species with low-abundance may be able to modulate the niche environment [56,57]. Therefore, further research at a molecular level is warranted to investigate the molecular mechanism behind the modulation of the oral microbiome by the probiotic *L. reuteri* strain. It is also noteworthy that our results do not imply an explicit causative relationship between the microbiome changes with the reduction of clinical parameters, considering the potential confounders and a smaller sample size. Nevertheless, the changes of certain key microbes associated with health and disease are restructured during the transition from the baseline to the 50% reduction of the clinical state. This change could be attributed to the probiotic intervention.

Future studies should be designed to examine the effect of long-term probiotic intervention

with an appropriate placebo control group in a large clinical cohort for more convincing evidence. Changes of the oral microbiota by the end of probiotic consumption after removal of fixed appliance as well as comparative evaluation of these changes prior to the bonding of orthodontic brackets should also be considered. Despite the foregoing limitation, the present study has provided valuable clinical and microbiological evidence on the health benefits of short-time administration of the probiotic *L. reuteri* lozenges on oral health. Hence, probiotic consumption may provide better oral health benefits by preventing accumulation of pathogenic microbiota while enriching symbiotic organisms during orthodontic treatment. These changes in the oral microbiome may have played a role in the observed improvement of the clinical parameters in the present study.

## Conclusion

Within the limitations of the present study, it can be concluded that a short-term probiotic intervention



helped maintaining good oral health in patients undergoing fixed orthodontic therapy. Although there was no overall change to the microbiome structure, significant modulation in the abundance of the health and disease-associated species highlighted the health benefits of probiotic intervention on the oral microbiome and oral health of patients undergoing orthodontic treatment.

## Acknowledgments

The authors thank the Faculty of Dentistry, Trisakti University for funding this study and the Microbiology Center of Research of Education (MiCORE) Laboratory, University of Trisakti for invaluable support in this study.

## Disclosure statement

No potential conflict of interest was reported by the author(s).

## Funding

This work was supported by the Universitas Trisakti


## Authorship contribution statement

ASW, CJS and JK designed the study. ASW, MIH and MOR collected the samples. MR performed the laboratory experiments. NS and CJS analyzed the data. ASW, NS and CJS wrote the paper. All authors discussed the results and commented on the manuscript.

## ORCID

Armelia Sari Widyarman  <http://orcid.org/0000-0003-1435-2392>

Nadeeka S. Udawatte  <http://orcid.org/0000-0001-6676-7704>

Moehamad Orliando Roeslan  <http://orcid.org/0000-0001-9260-7755>

Muhammad Ihsan Rizal  <http://orcid.org/0000-0002-2370-038X>

Mario Richi  <http://orcid.org/0000-0002-4729-8436>


## References

- [1] Julien KC, Buschang PH, Campbell PM. Prevalence of white spot lesion formation during orthodontic treatment. *Angle Orthod.* 2013;83(4):641–647.
- [2] Joint FAO/WHO expert consultation on evaluation of health and nutritional properties of probiotics in food including powder milk with live lactic acid bacteria. 2001;34.
- [3] Meurman JH. Probiotics: do they have a role in oral medicine and dentistry? *Eur J Oral Sci.* 2005 Jun;113(3):188–196.
- [4] Urbańska M, Szajewska H. The efficacy of *Lactobacillus reuteri* DSM 17938 in infants and children: a review of the current evidence. *Eur J Pediatr.* 2014;173(10):1327–1337.
- [5] Widyarman AS, Hartono V, Marjani LI, et al. *Lactobacillus reuteri* containing probiotic lozenges consumption reduces *streptococcus mutans*, *streptococcus sobrinus*, *porphyromonas gingivalis*, and *aggregatibacter actinomycetemcomitans* in orthodontic patients. *J Int Dental Med Res.* 2018 Jan 1;11(2):628–633.
- [6] Widyarman AS, Drestia AM, Bachtiar EW, et al. The anti-inflammatory effects of glycerol-supplemented probiotic *Lactobacillus reuteri* on infected epithelial cells in vitro. *Contemp Clin Dent.* 2018;9(2):298–303.
- [7] Twetman S, Derawi B, Keller M, et al. Short-term effect of chewing gums containing probiotic *Lactobacillus reuteri* on the levels of inflammatory mediators in gingival crevicular fluid. *Acta Odontol Scand.* 2009;67(1):19–24.
- [8] Krasse P, Carlsson B, Dahl C, et al. Decreased gum bleeding and reduced gingivitis by the probiotic *Lactobacillus reuteri*. *Swed Dent J.* 2006;30(2):55–60.
- [9] Ali Alasadi Z, Ahmed Qasim A. Impact of fixed orthodontic therapy on salivary characteristics in relation to weight status. *Biomed Pharmacol J.* 2018 Sep 28;11(3):1463–1470.
- [10] Widyarman AS, Udawatte NS, Theodorea CF, et al. Casein phosphopeptide–amorphous calcium phosphate fluoride treatment enriches the symbiotic dental plaque microbiome in children. *J Dent.* 2021 Mar 1;106:103582.
- [11] 16S rRNA microbiome protocol for Illumina iSeq 100 – ezBioCloud Help center. [Internet]. cited 2022 Apr 4. Available from: <https://help.ezbiocloud.net/16s-mtp-protocol-for-illumina-iseq-100/>
- [12] Afgan E, Baker D, Batut B, et al. The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. *Nucleic Acids Res.* 2018 Jul 2;46(W1):537–544.
- [13] Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods.* 2012 Apr;9(4):357–359.
- [14] DePristo MA, Banks E, Poplin R, et al. A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat Genet.* 2011 May;43(5):491–498.
- [15] Widyarman AS, Theodorea CF, Udawatte NS, et al. Diversity of oral microbiome of women from urban and rural areas of Indonesia: a pilot study. *Front Oral Health.* [Internet]. 2021 [cited 2022 Apr 4];2. Available from. [10.3389/froh.2021.738306](https://doi.org/10.3389/froh.2021.738306)
- [16] Chong J, Liu P, Zhou G, et al. Using MicrobiomeAnalyst for comprehensive statistical, functional, and meta-analysis of microbiome data. *Nat Protoc.* 2020 Mar;15(3):799–821.
- [17] McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLOS ONE.* 2013 Apr 22;8(4):e61217.
- [18] Anderson MJ. A new method for non-parametric multivariate analysis of variance. *Austral Ecol.* 2001;26(1):32–46.
- [19] Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Bioinformatics* [Internet].; 2014 Feb [cited 2022 Apr 4]. Available from. [10.1101/002832](https://doi.org/10.1101/002832)
- [20] Segata N, Izard J, Waldron L, et al. Metagenomic biomarker discovery and explanation. *Genome Biol.* 2011 Jun 24;12(6):R60.
- [21] Blankenberg D, Von Kuster G, Coraor N, et al. Galaxy, a web-based genome analysis tool for experimentalists. *Curr Protoc Mol Biol.* 2010 Jan;89(1). doi:[10.1002/0471142727.mb1910s89](https://doi.org/10.1002/0471142727.mb1910s89).
- [22] Parks DH, Tyson GW, Hugenholtz P, et al. STAMP: statistical analysis of taxonomic and functional profiles. *Bioinformatics.* 2014 Nov 1;30(21):3123–3124.

- [23] Liaw A, Wiener M. Classification and regression by randomForest. 2002;2:5. <https://cogns.northwestern.edu/cbm/LiawAndWiener2002.pdf>
- [24] Hastie T. The elements of statistical learning: data mining, inference and prediction. [Internet]. Universitas Indonesia Library. Springer; 2017 cited 2022 Apr 4. Available from: <https://lib.ui.ac.id>
- [25] Kursa MB, Rudnicki WR. Feature selection with the boruta package. J Stat Softw. 2010 Sep 16;36:1–13.
- [26] Breiman L. Random Forests. Mach Learn. 2001 Oct 1;45(1):5–32.
- [27] van Gastel J, Quirynen M, Teughels W, et al. Longitudinal changes in microbiology and clinical periodontal variables after placement of fixed orthodontic appliances. J Periodontol. 2008 Nov;79(11):2078–2086.
- [28] Uzuner FD, Kaygisiz E, Cankaya ZT. Effect of the bracket types on microbial colonization and periodontal status. Angle Orthod. 2014 Nov;84(6):1062–1067.
- [29] Hadj-Hamou R, Senok AC, Athanasiou AE, et al. Do probiotics promote oral health during orthodontic treatment with fixed appliances? A systematic review. BMC Oral Health. 2020 Apr 25;20(1):126.
- [30] Benic GZ, Farella M, Morgan XC, et al. Oral probiotics reduce halitosis in patients wearing orthodontic braces: a randomized, triple-blind, placebo-controlled trial. J Breath Res. 2019 May 31;13(3):036010.
- [31] Habib S. Assessment of the therapeutic potential of a dental probiotic in orthodontic patients affected by gingivitis: a randomized control trial [Internet] [Thesis]. 2016 cited 2022 Apr 4. Available from: <https://tspace.library.utoronto.ca/handle/1807/79737>
- [32] Jivraj FE. Effectiveness of the lorodent probiotic lozenge in reducing plaque and *S. mutans* levels in orthodontic patients: a randomized, double-blind, placebo-controlled trial [Internet] [Thesis]. 2015 cited 2022 Apr 4. Available from: <https://tspace.library.utoronto.ca/handle/1807/70287>
- [33] Jose JE, Padmanabhan S, Chitharanjan AB. Systemic consumption of probiotic curd and use of probiotic toothpaste to reduce *Streptococcus mutans* in plaque around orthodontic brackets. Am J Orthod Dentofacial Orthop. 2013 Jul;144(1):67–72.
- [34] Alp S, Baka ZM. Effects of probiotics on salivary *Streptococcus mutans* and *Lactobacillus* levels in orthodontic patients. Am J Orthod Dentofacial Orthop. 2018 Oct;154(4):517–523.
- [35] Pinto GS, Cenci MS, Azevedo MS, et al. Effect of yogurt containing *Bifidobacterium animalis* subsp. *lactis* DN-173010 probiotic on dental plaque and saliva in orthodontic patients. Caries Res. 2014;48(1):63–68.
- [36] Kado I, Hisatsune J, Tsuruda K, et al. The impact of fixed orthodontic appliances on oral microbiome dynamics in Japanese patients. Sci Rep. 2020 Dec 15;10(1):21989.
- [37] Shokeen B, Vilorio E, Duong E, et al. The impact of fixed orthodontic appliances and clear aligners on the oral microbiome and the association with clinical parameters: a longitudinal comparative study. Am J Orthod Dentofacial Orthop. 2022 Mar 2;(22):S0889–5406. DOI:10.1016/j.ajodo.2021.10.015
- [38] Agossa K, Dubar M, Lemaire G, et al. Effect of *Lactobacillus reuteri* on gingival inflammation and composition of the oral microbiota in patients undergoing treatment with fixed orthodontic appliances: study protocol of a randomized control trial. Pathogens. 2022 Feb;11(2):112.
- [39] Kuru BE, Laleman I, Yalnızoğlu T, et al. The Influence of a *Bifidobacterium animalis* probiotic on gingival health: a randomized controlled clinical trial. J Periodontol. 2017;88(11):1115–1123.
- [40] Toiviainen A, Jalasvuori H, Lahti E, et al. Impact of orally administered lozenges with *Lactobacillus rhamnosus* GG and *Bifidobacterium animalis* subsp. *lactis* BB-12 on the number of salivary mutans streptococci, amount of plaque, gingival inflammation and the oral microbiome in healthy adults. Clin Oral Investig. 2015;19(1):77–83.
- [41] Alkaya B, Laleman I, Keceli S, et al. Clinical effects of probiotics containing *Bacillus* species on gingivitis: a pilot randomized controlled trial. J Periodontol Res. 2017 Jun;52(3):497–504.
- [42] Iniesta M, Herrera D, Montero E, et al. Probiotic effects of orally administered *Lactobacillus reuteri*-containing tablets on the subgingival and salivary microbiota in patients with gingivitis. A randomized clinical trial. J Clin Periodontol. 2012 Aug;39(8):736–744.
- [43] Hallström H, Lindgren S, Yucel-Lindberg T, et al. Effect of probiotic lozenges on inflammatory reactions and oral biofilm during experimental gingivitis. Acta Odontol Scand. 2013 Jul;71(3–4):828–833.
- [44] Kohar N, Emmanuel V, Astuti L. Comparison between probiotic lozenges and drinks towards periodontal status improvement of orthodontic patients. Dental J (Majalah Kedokteran Gigi). 2015 Sep 1; 48(3):126.
- [45] Shah SS, Nambiar S, Kamath D, et al. Comparative evaluation of plaque inhibitory and antimicrobial efficacy of probiotic and chlorhexidine oral rinses in orthodontic patients: a randomized clinical trial. Int J Dent. 2019 Feb 20;e1964158. <https://doi.org/10.1155/2019/1964158>
- [46] Vivekananda MR, Vandana KL, Bhat KG. Effect of the probiotic *Lactobacilli reuteri* (Prodentis) in the management of periodontal disease: a preliminary randomized clinical trial. J Oral Microbiol. 2010 Nov 2;2:10.3402/jom.v2i0.5344.
- [47] Caglar E, Kavaloglu SC, Kuscu OO, et al. Effect of chewing gums containing xylitol or probiotic bacteria on salivary mutans streptococci and lactobacilli. Clin Oral Investig. 2007 Dec;11(4):425–429.
- [48] Nikawa H, Makihira S, Fukushima H, et al. *Lactobacillus reuteri* in bovine milk fermented decreases the oral carriage of mutans streptococci. Int J Food Microbiol. 2004 Sep 1;95(2):219–223.
- [49] Azad MAK, Sarker M, Wan D. Immunomodulatory effects of probiotics on cytokine profiles. Biomed Res Int. 2018 Oct 23:e8063647. <https://www.hindawi.com/journals/bmri/2018/8063647/>
- [50] Morita H, Toh H, Fukuda S, et al. Comparative genome analysis of *Lactobacillus reuteri* and *Lactobacillus fermentum* reveal a genomic Island for reuterin and cobalamin production. DNA Res. 2008 Jun 30;15(3):151–161.
- [51] Chen C. Distribution of a newly described species, *Kingella oralis*, in the human oral cavity. Oral Microbiol Immunol. 1996 Dec;11(6):425–427.
- [52] McDowell A, Patrick S, Eishi Y, et al. *Propionibacterium acnes* in human health and disease. Biomed Res Int. 2013;2013:493564.
- [53] Oliveira RRDS, Fermiano D, Feres M, et al. Levels of candidate periodontal pathogens in subgingival biofilm. J Dent Res. 2016 Jun;95(6):711–718.
- [54] Ehrmann E, Handal T, Tamanai-Shacoori Z, et al. High prevalence of  $\beta$ -lactam and macrolide resistance genes in human oral *Capnocytophaga* species. J Antimicrob Chemother. 2014 Feb;69(2):381–384.



- [55] Jing D, Hao J, Shen Y, et al. Effect of fixed orthodontic treatment on oral microbiota and salivary proteins. *Exp Ther Med.* 2019 May;17(5):4237–4243.
- [56] Dassi E, Ferretti P, Covello G, et al. The short-term impact of probiotic consumption on the oral cavity microbiome. *Sci Rep.* 2018 Jul 11;8(1):10476.
- [57] Dassi E, Ballarini A, Covello G. HTM-CMB2013. Quattrone A, Jousson O, et al. Enhanced microbial diversity in the saliva microbiome induced by short-term probiotic intake revealed by 16S rRNA sequencing on the IonTorrent PGM platform. *J Biotechnol.* 2014 Nov 20;190:30–39.

 Join the conversation about this journal

## Complexity Journal Manuscript

Join Leading Researchers in the Field of Complexity. Publish Your Next Research With Us

Hindawi


Open

## Complexity Journal Manuscript

Join Leading Researchers in the Field of Complexity. Publish Your Next Research With Us

Hindawi

Open

 Quartiles



## Complexity Journal Manuscript

Join Leading Researchers in the Field of Complexity. Publish Your Next Research With Us

Hindawi

Open

## Complexity Journal Manuscript

Join Leading Researchers in the Field of Complexity. Publish Your Next Research With Us Hindawi

Short- term effect of probiotic  
Lactobacillus reuteri  
consumption on the salivary  
microbiome profile of subjects  
undergoing orthodontic  
treatment with fixed appliances

*by* Joko Kusnoto

---

**Submission date:** 22-Sep-2022 09:16AM (UTC+0700)






**Submission ID:** 1905862282

**File name:** r\_Joko\_Kusnoto\_-\_07\_SCOPUS\_1\_-\_TAMBAHAN\_8\_SEPT\_2022\_1\_-20-30.pdf (5.05M)

**Word count:** 7358

**Character count:** 42615

## Short-term effect of probiotic *Lactobacillus reuteri* consumption on the salivary microbiome profile of subjects undergoing orthodontic treatment with fixed appliances

Armelia Sari Widyarman <sup>a</sup>, Nadeeka S. Udawatte <sup>b</sup>, Moehamad Orliando Roeslan <sup>c</sup>,  
Muhammad Ihsan Rizal <sup>c</sup>, Mario Richi <sup>d</sup>, Joko Kusnoto<sup>e</sup> and Chaminda Jayampath Seneviratne<sup>b,f</sup>

<sup>a</sup>Department of Microbiology, Faculty of Dentistry, Trisakti University, Jakarta, Indonesia; <sup>b</sup>Singapore Oral Microbiomics Initiative, National Dental Research Institute Singapore (NDRIS), National Dental Centre Singapore, Oral Health ACP, Duke NUS Medical School, Singapore; <sup>c</sup>Department of Oral Biology, Faculty of Dentistry, Trisakti University, Jakarta, Indonesia; <sup>d</sup>MiCORE Laboratory, Faculty of Dentistry, Trisakti University, Jakarta, Indonesia; <sup>e</sup>Department of Orthodontics, Faculty of Dentistry, Trisakti University, Jakarta, Indonesia; <sup>f</sup>School of Dentistry, The University of Queensland, Brisbane, Australia

### ABSTRACT

**Objective:** This prospective clinical study aim was to analyze the effect of the probiotic *Lactobacillus reuteri* Prodentis lozenges on salivary microbiome of subjects wearing fixed orthodontic appliances.

**Methods:** Saliva samples were collected prior to consumption and 14th-day post probiotic lozenges consumption (n=40, age 18–23). Oral hygiene index-score (OHI-S) and papilla bleeding index (PBI) were recorded. The salivary microbiome was profiled by next-generation sequencing using the V3–V4 region of 16S-rRNA. Microbial composition, diversity and taxonomic biomarkers were analysed in comparison to probiotic intervention and the clinical characteristics of the cohort using standard bioinformatics tools.

**Results:** The diversity and bacterial community structures did not change significantly in salivary microbiome of periodontally healthy subjects during short-term probiotic intervention. Probiotic consumption correlated with reduction of OHI and PBI scores (50% reduction of scores, P<0.001). The reduction of clinical indices was evident in conjunction with significantly reduced abundance of oral pathogens, such as *Porphyromonas pasteri*, *Treponema sp.*, *Fretibacterium fastidiosum*, *Kingella oralis* and *Propionibacterium acnes*.

**Conclusion:** Short-term probiotic intervention helped maintaining good oral health in patients undergoing fixed orthodontic therapy. Although overall oral microbiome structure remained largely unchanged, a significant alteration in the abundance of health and disease-associated species highlighted the beneficial effect of probiotic.

### ARTICLE HISTORY

Received 10 February 2022  
Revised 7 April 2022  
Accepted 14 April 2022

### KEYWORDS

Probiotic; *Lactobacillus reuteri*; oral microbiome; fixed orthodontic appliance

## Background

Fixed orthodontic treatment is a risk factor for the development of various oral diseases, such as dental caries, periodontal disease, and candidiasis. The presence of brackets, bands, and arch wire complicates oral hygiene practices and increases the accumulation of dental plaque. Hence, patients wearing a fixed orthodontic treatment are at higher risk for developing oral diseases [1]. Therefore, orthodontic patients need supplementary oral hygiene measures in addition to personal mechanical oral hygiene practices. Probiotics, which are defined as ‘living microorganisms which, when administered in adequate amounts, confer a health benefit on the host’ by the Food and Agriculture Organization of the United Nations/World Health Organization (FAO/WHO) have been shown to confer various health benefits to humans, including improvement of oral health [2].

*Lactobacillus reuteri* is a probiotic bacterium known to confer various health benefits to humans [3]. It produces the antimicrobial substances reuterin and reutericyclin, which are active against a wide range of pathogenic organisms [4]. Previously, we performed a pilot study on the beneficial health effects of lozenges containing the probiotic *L. reuteri* on a few orthodontic patients [5]. We found that consumption of probiotics for 2 weeks significantly reduced the number of pathogenic bacteria in the patients’ saliva. Moreover, other studies using appropriate *in vitro* and *in vivo* models have demonstrated that *L. reuteri* is able to modulate host inflammatory responses [6]. However, clinical studies that examine the effect of the probiotic *L. reuteri* on the oral microbiome are sparse in the literature.

To our knowledge, there is no information available regarding how *L. reuteri* probiotics regulate the oral microbiome in periodontally healthy subjects

**CONTACT** Armelia Sari Widyarman  [armeliasari@trisakti.ac.id](mailto:armeliasari@trisakti.ac.id)  Department of Microbiology, Faculty of Dentistry, Trisakti University, Indonesia; Chaminda Jayampath Seneviratne  [jaya.seneviratne@uq.edu.au](mailto:jaya.seneviratne@uq.edu.au)  School of Dentistry, The University of Queensland, Australia

 Supplemental data for this article can be accessed here.

© 2022 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.



using fixed orthodontic appliances. Considering this research gap, we aimed to analyze the short-term effect of the consumption of the probiotic *L. reuteri* Prodentis lozenges on the salivary microbiome profile of subjects wearing fixed orthodontic appliances.

## Materials and methods

### Cohort characteristics

The subjects undergoing fixed orthodontic therapy were included in this open-labelled prospective clinical trial. Ethical approval for the study was obtained from the Institutional Review Board, Faculty of Dentistry, Universitas Indonesia (approval number 100,701,020). The inclusion criteria included an age above 18 years, undergoing orthodontic treatment with a fixed appliance of a minimum of 1 year and no consumption of probiotics or antibiotics within the past 3 months. Exclusion criteria included subjects who had systemic diseases, such as hypertension or diabetes, or were on systemic drugs *i.e.* anti-hypertensives, analgetics, hormonal drugs, sedatives, anti-seizure medication and subjects with features of severe periodontal diseases and who were allergic to probiotics. The lozenges containing *L. reuteri* Prodentis were obtained from BioGaia, Stockholm, Sweden. BioGaia Prodentis lozenges (800 mg of one lozenge contained a minimum of  $2 \times 10^8$  live *L. reuteri* Prodentis) were a food supplement for oral health containing the patented lactic acid bacterium *Limosilactobacillus reuteri* (formerly known as *Lactobacillus reuteri*) Prodentis. They also contained a strain combination of *L. reuteri* DSM 17938 and *L. reuteri* ATCC PTA 5289 that was expected to help maintain a good oral health.

To standardize the oral hygiene practice, each participant was given a toothpaste and a toothbrush during the study period. The participants were given oral hygiene instructions and were requested to brush their teeth twice a day. Each subject was given one lozenge per day, containing the probiotic *L. reuteri* for 14 days to be taken once a day after brushing teeth, followed by breakfast. Saliva samples were collected from the subjects on the day of the recruitment prior to the consumption of probiotic lozenges on the 14th-day post-consumption of the probiotics.

In addition, clinical data pertaining to the oral hygiene index score (OHI-S) and papilla-bleeding index (PBI) were recorded at each visit. Previously, it had been found that consumption of probiotic *L. reuteri* Prodentis over 14 day-duration is able to exert considerable impact on reducing clinical parameters (Periodontal Index-PI, OHI-S and Gingival index-GI) in healthy test subjects compared to a placebo control [7,8].

### Sample collection and DNA extraction

Prior to sample collection, subjects had to take a rapid antigen test for SARS-CoV-2 detection due to the COVID-19 pandemic (as of 30 September 2020) and the result of the test had to be negative for taking part in the study. Following this confirmation, sample specimens from the subjects were collected with a cover of a complete set of personal protective equipment. A standard method was used for the saliva sample collection [9]. In brief, the subject was made to sit quietly, bending the head forward and opening the mouth to allow the saliva to drip passively from the lower lip. Saliva was collected using sterile, pre-weighted tubes. Subjects were requested to refrain from food for 2 h prior to the sample collection. Following the collection, the saliva samples were stored at  $-80^{\circ}\text{C}$  prior to DNA extraction.

Microbiome profiling of the saliva samples was performed using a previously established protocol with some modification [10]. In brief, DNA was extracted from the samples using the QIAamp<sup>®</sup> DNA Mini Kit (Qiagen, Hilden, Germany) per the manufacturer's protocol. Extracted DNA was subjected to a quality check using Nanodrop<sup>™</sup> 2000 (Thermo Fisher, Waltham, MA) for its purity and its concentration of gDNA using a Qubit 3.0 fluorometer (Thermo Fisher, Waltham, MA).

### 16S rRNA amplicon library preparation and sequencing

Polymerase chain reaction (PCR) amplifications had been finished for library preparation prior to sequencing as previously described [11]. Primers targeted for the V3-V4 region of 16S rRNA and modified with Illumina overhang adapters had been amplified by using PCR and the primers: F (5' - TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCCTACGGGNGGC WGCAG-3') and R (5' GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGG ACTACHVGGGTATCTAATCC -3'). The PCR amplification was done as described previously [10]. The amplified library was subjected to purification using AMPure XP beads (Beckman Coulter Genomics, MA), visualized via agarose gel electrophoresis and quantified by using the Qubit 3.0 Fluorometer (Thermo Fisher Scientific, Waltham, MA). The amplified library after the second purification using AMPure XP Beads was cluted, normalized, and pooled to the recommended concentration as per the manufacturer's protocol (80 pM). An internal 5% spike-in control using the Phi-X manage v3 library (Illumina, San Diego, CA) was pooled together with the normalized amplicon library. Sequencing was performed using the iSeq 100 instrument with the i1 v2 reagent (Illumina, San Diego, CA) with the



parameters of  $2 \times 150$  cycles paired-end reads, with 10 cycles for every index, in line with the manufacturer's protocols.

### Data analysis

Sequencing runs were first analysed in the Illumina's Sequencing evaluation Viewer version 2.4.7 for analytical run quality controls assessments [12–14]. Pre-processing of the raw sequencing data and the Quality Control (QC) was assessed as previously described [15]. The 16S metagenomics workflow based on the Local Run Manager v2.0 used the Illumina-curated version of the Greengenes database ([greengenes.secondgenome.com/downloads/database/13\\_5](http://greengenes.secondgenome.com/downloads/database/13_5)) for taxonomic classification of 16S rRNA centred amplicon reads. It generated a total of 1,282 operational taxonomic units (OTUs) clustered at 97% identification with an open-reference OTU picking approach. The resulting accumulation curves showed reasonable sequence saturation for subsequent analysis (Supplementary Figure 1).

Alpha diversity (Chao1, Shannon, ACE), along with rarefaction curves, was evaluated and visualized using the MicrobiomeAnalystR Platform and the 'phyloseq' and 'ggplot' and 'microbiomeseq' [16,17] codes obtained from GitHub (<https://github.com/xia-lab/MicrobiomeAnalystR>). The microbial community structure was assessed by beta diversity with the respective algorithms implemented in the MicrobiomeAnalystR package using Principal Coordinate Analysis (PCoA) and the Bray–Curtis's dissimilarity indexes retrieved from sample pairwise comparisons based totally on weighted Unifrac distances. Significance of the community structure analysis was performed using the permutational multivariate analysis of variance (PERMANOVA) test [18].

DESeq2 [19] analysis was run under default settings and *q* values were calculated using the Benjamini–Hochberg procedure to control for false discovery rates applying the MicrobiomeAnalystR. Based on both significance and biological relevance, differentially abundant microbial biomarkers were characterized using the linear discriminant analysis effect size (LEfSe) method with default parameters and an LDA log score threshold set to 2 with a *P* value less than 0.05 considered as statistically significant [20,21]. Species with a *P* value <0.1 on the univariate analysis with probiotic intervention were selected for further analysis. The relative abundance differences in the top 27 bacterial species between pre- and post-probiotic cohorts were compared by Welch's *t*-test using STAMP (v2.1.3) [22]. Further, the random forest (RF) algorithm was applied to determine the proper number of microbial predictors in random forest classifiers by maximizing the area under the curve of the receiver operator characteristic (AUC) with the default parameters of the

'RandomForest' (RF) package in R (v. 4.6.1). To avoid over-fitting of the data in the model, 5-time and 10-fold cross-validations were made. The resulting model was subsequently used for the validation cohort [23,24].

To identify key members of the microbiome (based on relative abundance) associated with 50% reduction of the oral hygiene index score from the baseline value after probiotic intake, the Boruta algorithm [Boruta package for R (v. 5.1)] was used to perceive species with the highest predictive power [25]. The relative abundance >0.1% and presence in >50% of the samples in at least one group were used for the above analyses unless otherwise specified.

### Statistical analyses

Statistical analyses were carried out using the GraphPad Prism software (version 8) and the R package (<http://www.r-project.org/>). The Mann–Whitney rank sum test or the Kruskal–Wallis test was used to evaluate variations among groups in the diversity indices of the oral microbiota. False discovery rate (FDR) values were used for the evaluation of the Benjamini–Hochberg method to control for multiple testing – adjusted *P* values lower than 0.05 were considered as statistically significant [19]. The analysis was confined to species with a prevalence greater than 10% and a maximum proportion (relative abundance) greater than 0.002. An FDR-adjusted *P* value (or *Q*-value) <5% was considered as significant.

The Boruta algorithm was used to identify the most predictive taxonomic biomarker based on the importance values determined by RF, indicated as 'confirmed' in the Boruta output [25]. The importance value pertaining to a taxon was calculated by RF based on the loss of accuracy by random permutation of the abundance profile of the taxon. To assess whether the importance was significant, the Boruta algorithm compared the observed importance to those produced by the spiked-in 'shadow' taxa, which were randomized versions of original taxa; hence, this algorithm typically gave more strength to identify taxa that jointly predicted a phenotype [26].

## Results


### Effect of probiotic intake on the clinical parameters

The mean age of the participants (*n* = 20; male – 4, female – 16) was 21 years (SD ± 1.59) across all samples (Figure 1A). Consumption of probiotic *L. reuteri* lozenges for 14 days improved the oral health parameters of the subjects as measured by the reduction of the OHI-S (*P* < 0.0001) and PBI scores (*P* < 0.0056) (Figure 1B).


### Effect of probiotics on taxonomic composition and complexity of the salivary microbiome

We then taxonomically profiled each group based on the relative abundances of microbial clades at distinctive taxonomic tiers. Five phyla made up >1% of the community in at least one sample with two phyla naming, *Firmicutes* and *Proteobacteria*, dominated the groups with relative abundance (33–30% and 33–26% in the pre and post-probiotic groups, respectively) among the top 10 predominant phylum groups.

*Actinobacteria* and *Bacteroidetes* were present in all samples, however to a lesser extent (16–11% and 22–10% in the pre- and post-probiotic groups, respectively) (Figure 2A). Lower-abundance phyla such as *Spirochaetes* were increased in post-probiotic organisms, and the so-called 'environmental' phyla such as SR1 were detected in similar level of consistency. In the top 15 genus level, *Streptococcus* and *Haemophilus* made up for almost 28% of the total reads, with 27% and 28% in the pre- and post-probiotic groups, respectively. *Rothia*, *Actinomyces*, *Fusobacterium*, *Neisseria*,



**Figure 1.** Demographic and clinical evaluation of individuals with good oral hygiene using orthodontic-fixed appliances prior and post-probiotic *L. reuteri* lozenges supplementation. **(A)** Baseline characteristics of the study participants in both intervention groups. **(B)** Two clinical outcomes were evaluated before and after 14 days of probiotic intervention. The Simplified Oral Hygiene index (OHIS) for determining the soft sediments classified with microbial signatures designed no plaque (0), mild [1- (1/3) covered with soft plaque], moderate [2-(1/3-2/3) covered with soft plaque], bad [3- (> 2/3) covered with soft plaque] according to the Green – Vermillion – Hirschman index. The Papillary Bleeding Index (PBI) used to determine the presence or absence of inter-dental plaque; designed no bleeding (0), only one bleeding point appearing [1], several isolated bleeding points or a small dental area appearing [2], inter dental triangle filled with blood soon after probing [3], and profuse bleeding when probing led to blood spreads towards the marginal gingiva [4]. All subjects showed significant oral health improvement, as measured by the reduction of the OHIS and PBI scores ( $P < 0.001$ ), by the end of the probiotic supplementation.




**Figure 2.** Taxonomic composition of the salivary microbiome before and after probiotic supplementation. **(A)** The top 10 relative abundances of the microbial phyla highlights the major role played by *Firmicutes* and *Proteobacteria* (green and brown) in the microbiome, and to a less extent by *Actinobacteria* and *Bacteroidetes* (light brown and purple). **(B)** At the top 15 genus level, *Haemophilus*, *Streptococcus*, *Veillonella*, *Rothia* and *Neisseria* (maroon, blue, dark green, light green and light purple) are shown to be the most abundant clades. **(C)** Out of the genera accounting for less than 1% of the samples which are grouped in the 'Other' category, *Schlegelella*, *Bacillus*, *Pseudoxanthomonas*, *Lactobacillus*, *Microbacterium* and *Anoxybacillus* were significantly increased after probiotic intake. Letters indicate statistical differences between samples are \* $P < 0.05$ , \*\* $P < 0.01$ , and \*\*\* $P < 0.001$ .



*Veillonella*, *Prevotella*, *Selenomonas* and *Leptotrichia* were also detected in all samples with uneven quantitative distributions (Figure 2B). At the species level, *Haemophilus parainfluenzae*, *Rothia dentocariosa*, *Leptotrichia hongkongensis*, *Haemophilus haemolyticus* and *Neisseria elongata* were the main species components in both the pre- and post-probiotic groups (relative abundance >3%). Overall, taxonomic phyla and genus abundance distributions of these dominant oral microbes (top 10 taxa) were minimally influenced by short-term probiotic intervention. Out of lesser abundant genera (relative abundance <1%), *Schlegelella* ( $P = 0.00043$ ), *Bacillus* ( $P = 0.0011$ ), *Pseudoxanthomonas* ( $P = 0.003$ ), *Lactobacillus* ( $P = 0.037$ ), *Microbacterium* ( $P = 0.043$ ), and *Anoxybacillus* ( $P = 0.037$ ) were significantly increased following the probiotic intervention (Figure 2C).

The microbial community composition of the saliva samples at baseline and 2-week post-consumption was analysed. There was no significant difference in the alpha diversity (Shannon –  $P$  value 0.698; [Mann-Whitney]) and evenness (Chao1 –  $P$  value 0.881; ACE –  $P$  value 0.989; Simpson –  $P$  value 0.62 [Mann-Whitney]) of the post-consumption samples compared to the baseline (Figure 3A). Post-consumption saliva samples had a slightly higher alpha diversity than pre-probiotic intake samples, although the difference was not statistically significant. To understand whether the quantitative structure of the microbiome changed, PCoA on the taxonomic profiles using the weighted UniFrac measure was performed. Ordination analysis showed no obvious clustering by pre-defined groups, suggesting that the overall community structure of the saliva microbiome was not stricken by short-term



**Figure 3.** Structural composition and diversity comparison of the salivary microbiome before and after probiotic intervention. The overall structure of the saliva microbiome is not affected by short-term probiotic intake. **(A)**  $\alpha$ -diversity boxplot (Shannon, Chao1 and Simpson-reciprocal indices) of the salivary microbiome between pre- and post-probiotic cohorts. Boxes represented the 25th to 75th percentile of the distribution; the median was shown as a thick dot in the middle of the box; whiskers extend between the 25th and 75th percentiles. **(B)** No apparent clustering trend can be noticed ( $P > 0.05$ ) in the PCoA space when the non-quantitative unweighted UniFrac measure is used as beta-diversity, suggesting that a change in the relative abundances of the main components of the salivary microbiome is not the driving feature characterizing between probiotic and respective baseline samples. Species with a  $P$  value < 0.1 on the univariate analysis with probiotic intervention were selected to define the top most abundantly differentiated across the study cohort. **(C)** Heatmap of the top 27 abundant species differentiates between probiotic and baseline samples. The color intensity in each box indicates the relative percentage of species in each sample. Relative abundance data were z-scored normalized by row. To show the distribution of abundant OTUs, the relative abundances of OTU data were normalized to have a mean of 0 and a standard deviation of 1 (z-score normalization). **(D)** Random Forest algorithm with 5 times 10-fold-cross validation used to identify a minimal set of bacterial species that maximally differentiated between pre- and post-probiotic intervention. *Schlegelella aquatica*, *Microbacterium mitrae*, *Microbacterium ginsengisoli*, and *Streptococcus parasuis* ranked the predominant difference between two cohorts under probiotic intervention. **(E)**  $P < 0.01$  to detect species with the most statistically significant differences filtered using a q-value of 0.05 and effective size of 0.05 threshold in STAMP illustrating significant bacterial species in variations of different proportions using Welch's t-test. See also Supplementary Figure 2 for LEfSe results.



probiotic intake (PERMANOVA  $P$  value  $<0.688$ ) (Supplementary Table 1) (Figure 3B).

### Increased complexity of the salivary microbiome following short-term probiotic consumption

We performed a linear discriminant analysis effect size (LEfSe) to determine which microbial clades, if any, could explain the slight increase in the diversity of the salivary microbiome following short-term probiotic intervention. The LEfSe analyses identified 12 species in the salivary microbiome whose over-abundance discriminated between pre- and post-probiotic samples (Figure 3 B and C) (full details are given in Supplementary Table 2 and Supplementary Figures 2 and 3). Pre-probiotic samples had a significant enrichment of species (LDA log score at a threshold of 3.0): *Schlegelella aquatica* (Proteobacteria), *Streptococcus parasuis* (Firmicutes), *Anoxybacillus pushchinoensis* (Firmicutes), *Anoxybacillus bogrovensis* (Firmicutes), *Pseudoxanthomonas taiwanensis* (Proteobacteria). On the other hand, *Sediminibacterium goheungense* (Bacteroidetes) and *Microbacterium ginsengisoli* (Actinobacteria) OTUs gave significance at the same effect size threshold for discriminating samples with probiotic consumption. Further analysis using 10-cross validation RF modeling (Figure 3D) and STAMP methods (Figure 3E) showed that few of these identified species, *S. aquatica*, *Microbacterium mitrae*, *M. ginsengisoli* and *S. parasuis*, reflected the predominant difference between pre- and post-probiotic cohorts under the short-time probiotic intervention.

We used the Boruta feature selection algorithm coupled with the RF classifier to examine which differential microbiome markers (predictors) at the species level accounted for 50% reduction of observed clinical indexes *i.e.* OHIS and PBI scores (response variables) with probiotic intervention. As evidenced by a decreased predictive mean square error compared to a random guess, a predictive model utilized 18 species with significant predictive power substantiated by LEfSe analysis (Figure 4 C and 4D) (Supplementary Table 3). Among the identified microbiome markers, there was a reduction in disease-associated pathogens, such as *P. pasteri*, *Treponema* sp. and *Prevotella* sp., which correlated with the 50% reduction of the OHIS index score group (OHIS50-Y/Yes) (Figure 4G). Based on feature selection by Boruta, over-abundance of *M. ginsengisoli*, *S. canadensis* and *A. suimastitidis*, and reduced relative abundance of *F. fastidiosum* were highly discriminative for the 50% reduction of the OHIS index score with probiotic intervention (Supplementary Figure and Table 3) (Figure 4C). On the other hand, by both LEfSe and Boruta feature selection; overabundance of *C. leadbetteri* and decreased abundance of *K. oralis*, *P. acnes*,

*S. aquatica*, and *A. bogrovensis* were identified in the 50% reduction of the PBI index score group by probiotic intake (PBI50-Y/Yes) (Supplementary Figure and Table 3) (Figure 4D). These species may independently be useful for predicting the 50% reduction of the PBI index score with probiotic intervention. The key species affected by probiotic intervention according to the clinical parameters are presented in heatmaps (Figure 4F and 4G). Moreover, the chord diagram was used to delineate species influenced by different clinical states before and after probiotic intake (Figure 4E). The chord diagram was divided into five sections representing the two clinical states of PBI and OHI-S before and after probiotic intake. Key species identified in two or more clinical states are indicated by arcs (chords) connecting the different clinical states.

Re-analysis of the overall salivary microbiome profiles using only the foregoing species with significant predictive power yielded some interesting results. PERMANOVA analysis significantly differentiated the salivary microbiome composition following probiotic intervention [between OHIS50-N/No and OHIS50-Y/Yes groups ( $P = 0.008$ ) and PBI50-N/No and PBI50-Y/Yes ( $P = 0.001$ )] (Figure 4 A and B).


### Persistence of the *Lactobacillus* genus in the salivary microbiome after probiotic intake

The *Lactobacillus* strain present in the probiotic product (*L. reuteri*) was not detected in the probiotic-intake samples. However, we observed the presence of a significantly higher-abundance OTU attributable to the *Lactobacillus* genus in the post-probiotic samples (Figure 2C).

The foregoing results suggested that the short course of the probiotic *L. reuteri* lozenges could potentially modulate the salivary microbiome, albeit the effect was limited in both the breadth of the change and in its temporal persistence.

### Discussion

There has been some evidence from clinical and fundamental studies suggesting that fixed orthodontic appliances affect the oral health of the patients [27,28]. The effect of probiotics on various microbiological and clinical parameters in fixed orthodontic patients has produced conflicting evidence [1,29–35]. The present study evaluated the effect of oral probiotics on the salivary microbiome of periodontally healthy subjects under orthodontic treatment with fixed appliances using a prospective interventional study. The oral health of the subjects was significantly improved following 14 days' intake of the probiotic *L. reuteri* Prodentis lozenges. Although the overall diversity or evenness of the salivary microbiome did



**Figure 4.** Increased complexity of the salivary microbiome in conjunction with 50% reduction of the oral hygiene index scores in subjects with fixed orthodontic appliances. **(A)** and **(B)** Apparent clustering in structural salivary composition observed when coupled with 50% reduction of the oral hygiene index scores (OHIS and PBI) after probiotic intervention. **(C)** and **(D)** Significant differentially abundant taxonomic biomarkers contribute to the change in the microbiome composition identified by LEfSe analysis. **(E)** The chord diagrams show the key taxa identified by the five different strategies (Baseline, post-probiotic; OHIS50-Y, OHIS50-N, PBI50-Y and PBI50-N). Key species identified in two or more clinical states are indicated by arcs (chords) connecting the different clinical states. **(F)** and **(G)** Heatmap of most abundant species among the OHIS50 groups differentiates between probiotic and baseline samples. Species highlighted in dotted blue squares are distinguished to have a strong predictive power for 50% reduction of both the OHIS and PBI index scores with probiotic intervention identified by the Machine learning Boruta algorithm coupled with the Random Forest classifier. See also Supplementary Figure 4. Abbreviations: PRPT- Pre-Probiotic group, POPT- Post-Probiotic group, OHIS50-N/No and PBI50-N/No – baseline and no change of OHIS and PBI index score groups, OHIS50-Y/Yes and PBI50-Y/Yes – 50% reduction of OHIS and PBI index score groups in probiotic intake.

not change significantly, there were minor changes in the abundance of disease-associated species following probiotic intervention.

Fixed orthodontic treatment may pose a significant challenge to maintain oral health of the patients as the appliance traps dental plaque biofilm leading to increased level of potential oral pathogens [36,37]. The use of probiotics as an adjunct therapy has been suggested for various diseases including periodontitis. Overall, the administration of probiotics, either as an adjunct to mechanical plaque control or as a single intervention, has yielded conflicting results [38]. Some studies have proven the advantages of probiotic consumption in reducing the extent of gingival

inflammation and the amount of plaque [39,40]. In contrast, other studies have failed to exhibit additional clinical benefits [41–43]. Moreover, several investigations documented the varying impact of long-term probiotic intervention (maximum up to 1 month) on maintaining gingival health in orthodontic patients, some with beneficial improvements [44,45] while some without improvement [30,31]. There are ample evidence in the literature supporting the beneficial effects of oral *L. reuteri* such as its capability: (i) to inhibit plaque accumulation [46], (ii) to reduce cariogenic *mutans* streptococci [47,48], (iii) to improve clinical parameters when used as an adjunct therapy for periodontitis [7,8,46] and (iv) to exert beneficial



immunomodulatory and anti-inflammation effects [46,49]. *L. reuteri* has been considered as a potential probiotic adjunct together with non-surgical periodontal therapy for periodontitis patients [7,8,50]. In our study, we observed reduced gingival bleeding with improved PBI scores and overall oral hygiene improvement by the OHIS index in periodontally healthy individuals undergoing fixed appliances for orthodontic treatment after intake of the probiotic *L. reuteri* Prodentis lozenges. The observed difference in the improved oral health may be attributed to the short-term probiotic intervention. However, according to a recently published review, no consensus can be reached on the significant effect of oral probiotics on gingival inflammation and enamel decalcification in patients undergoing treatment with fixed orthodontic appliances [29]. Therefore, in the presented study, we investigated whether the impact of probiotic would be more pronounced by defining our study cohort with respect to 50% reduction of the oral clinical state from their respective baseline score.

In this context, there were some interesting findings with regard to the change in key species associated with the salivary microbiome. Two weeks' probiotic intake reduced the disease-associated pathogens, such as *Porphyromonas* sp., *Treponema* sp., *Prevotella* sp. and *F. fastidiosum*, which correlated with the 50% reduction of the OHIS index score group (OHIS50-Y), suggesting improvement of oral health. Moreover, probiotic interventions decreased the abundance of the periodontal pathogens *K. oralis* and *P. acnes*, which correlated with the 50% reduction of the PBI index score group (PBI50-Y). Hence, probiotic intervention considerably eliminated the dysbiotic oral microbiome by reducing disease-associated pathogens and enriching health-associated species. *K. oralis* is normally found in small numbers in the oral cavity in dental plaque biofilm [51]. It has been reported that the species increased significantly in periodontitis *i.e.* more than 5% of the total microbiota in periodontitis sites compared to the meagre number (0.4%) in periodontally healthy individuals. Additionally, *P. acnes* and *F. fastidiosum* have been shown to be associated with gingivitis [52] and periodontitis [53], respectively. Significant reduction of these species indicated the potential modulatory activity of probiotic intake. Moreover, *Capnocytophaga* species, known to be associated with healthy periodontium, increased following probiotic intervention [54].

Fixed orthodontic appliance could be considered as a condition that may significantly change the local micro-environment in dental plaque biofilm, consequently resulting in an altered salivary microbiome [55]. Previous studies have shown significant dysbiosis in both the saliva microbiome and supragingival plaque microbiome with substantial replacement of anaerobic

bacteria in patients wearing fixed orthodontic appliances [36,37]. In the present study, we observed the presence of certain facultative anaerobes like *S. parasuis*, *Anoxybacillus* sp., *Bacillus thermoamylovorans* and mesophilic anaerobes like *M. mitrae* in the salivary microbiome of patients under fixed orthodontic treatment. Probiotic intervention reduced the proportion of these taxa. However, the absence of a placebo control group in the present study is a major limitation for a comprehensive comparison. Moreover, we did not detect the presence of the administered probiotic strain *L. reuteri* in the post-intervention salivary microbiome. However, this does not exclude the probability that this probiotic strain colonizes in dental plaque biofilms. Therefore, future studies should aim at analyzing both saliva and dental plaque samples from the subjects. On the other hand, according to the key-stone plaque hypothesis, species with low-abundance may be able to modulate the niche environment [56,57]. Therefore, further research at a molecular level is warranted to investigate the molecular mechanism behind the modulation of the oral microbiome by the probiotic *L. reuteri* strain. It is also noteworthy that our results do not imply an explicit causative relationship between the microbiome changes with the reduction of clinical parameters, considering the potential confounders and a smaller sample size. Nevertheless, the changes of certain key microbes associated with health and disease are restructured during the transition from the baseline to the 50% reduction of the clinical state. This change could be attributed to the probiotic intervention.

Future studies should be designed to examine the effect of long-term probiotic intervention

with an appropriate placebo control group in a large clinical cohort for more convincing evidence. Changes of the oral microbiota by the end of probiotic consumption after removal of fixed appliance as well as comparative evaluation of these changes prior to the bonding of orthodontic brackets should also be considered. Despite the foregoing limitation, the present study has provided valuable clinical and microbiological evidence on the health benefits of short-time administration of the probiotic *L. reuteri* lozenges on oral health. Hence, probiotic consumption may provide better oral health benefits by preventing accumulation of pathogenic microbiota while enriching symbiotic organisms during orthodontic treatment. These changes in the oral microbiome may have played a role in the observed improvement of the clinical parameters in the present study.

## Conclusion

Within the limitations of the present study, it can be concluded that a short-term probiotic intervention



helped maintaining good oral health in patients undergoing fixed orthodontic therapy. Although there was no overall change to the microbiome structure, significant modulation in the abundance of the health and disease-associated species highlighted the health benefits of probiotic intervention on the oral microbiome and oral health of patients undergoing orthodontic treatment.

### Acknowledgments

The authors thank the Faculty of Dentistry, Trisakti University for funding this study and the Microbiology Center of Research of Education (MiCORE) Laboratory, University of Trisakti for invaluable support in this study.

### Disclosure statement

No potential conflict of interest was reported by the author(s).

### Funding

This work was supported by the Universitas Trisakti


### Authorship contribution statement

ASW, CJS and JK designed the study. ASW, MIH and MOR collected the samples. MR performed the laboratory experiments. NS and CJS analyzed the data. ASW, NS and CJS wrote the paper. All authors discussed the results and commented on the manuscript.

### ORCID

Armelia Sari Widyarman  <http://orcid.org/0000-0003-1435-2392>

Nadeeka S. Udawatte  <http://orcid.org/0000-0001-6676-7704>

Moehamad Orliando Roeslan  <http://orcid.org/0000-0001-9260-7755>

Muhammad Ihsan Rizal  <http://orcid.org/0000-0002-2370-038X>

Mario Richi  <http://orcid.org/0000-0002-4729-8436>

### References

- [1] Julien KC, Buschang PH, Campbell PM. Prevalence of white spot lesion formation during orthodontic treatment. *Angle Orthod.* 2013;83(4):641–647.
- [2] Joint FAO/WHO expert consultation on evaluation of health and nutritional properties of probiotics in food including powder milk with live lactic acid bacteria. 2001;34.
- [3] Meurman JH. Probiotics: do they have a role in oral medicine and dentistry? *Eur J Oral Sci.* 2005 Jun;113(3):188–196.
- [4] Urbańska M, Szajewska H. The efficacy of *Lactobacillus reuteri* DSM 17938 in infants and children: a review of the current evidence. *Eur J Pediatr.* 2014;173(10):1327–1337.
- [5] Widyarman AS, Hartono V, Marjani LI, et al. *Lactobacillus reuteri* containing probiotic lozenges consumption reduces *streptococcus mutans*, *streptococcus sobrinus*, *porphyromonas gingivalis*, and *aggregatibacter actinomycetemcomitans* in orthodontic patients. *J Int Dental Med Res.* 2018 Jan 1;11(2):628–633.
- [6] Widyarman AS, Drestia AM, Bachtiar EW, et al. The anti-inflammatory effects of glycerol-supplemented probiotic *Lactobacillus reuteri* on infected epithelial cells in vitro. *Contemp Clin Dent.* 2018;9(2):298–303.
- [7] Twetman S, Derawi B, Keller M, et al. Short-term effect of chewing gums containing probiotic *Lactobacillus reuteri* on the levels of inflammatory mediators in gingival crevicular fluid. *Acta Odontol Scand.* 2009;67(1):19–24.
- [8] Krasse P, Carlsson B, Dahl C, et al. Decreased gum bleeding and reduced gingivitis by the probiotic *Lactobacillus reuteri*. *Swed Dent J.* 2006;30(2):55–60.
- [9] Ali Alasadi Z, Ahmed Qasim A. Impact of fixed orthodontic therapy on salivary characteristics in relation to weight status. *Biomed Pharmacol J.* 2018 Sep 28;11(3):1463–1470.
- [10] Widyarman AS, Udawatte NS, Theodorea CF, et al. Casein phosphopeptide–amorphous calcium phosphate fluoride treatment enriches the symbiotic dental plaque microbiome in children. *J Dent.* 2021 Mar 1;106:103582.
- [11] 16S rRNA microbiome protocol for Illumina iSeq 100 – ezBioCloud Help center. [Internet]. cited 2022 Apr 4. Available from: <https://help.ezbiocloud.net/16s-mtp-protocol-for-illumina-iseq-100/>
- [12] Afgan E, Baker D, Batut B, et al. The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. *Nucleic Acids Res.* 2018 Jul 2;46(W1):537–544.
- [13] Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods.* 2012 Apr;9(4):357–359.
- [14] DePristo MA, Banks E, Poplin R, et al. A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat Genet.* 2011 May;43(5):491–498.
- [15] Widyarman AS, Theodorea CF, Udawatte NS, et al. Diversity of oral microbiome of women from urban and rural areas of Indonesia: a pilot study. *Front Oral Health.* [Internet]. 2021 [cited 2022 Apr 4];2. Available from: 10.3389/froh.2021.738306
- [16] Chong J, Liu P, Zhou G, et al. Using MicrobiomeAnalyst for comprehensive statistical, functional, and meta-analysis of microbiome data. *Nat Protoc.* 2020 Mar;15(3):799–821.
- [17] McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLOS ONE.* 2013 Apr 22;8(4):e61217.
- [18] Anderson MJ. A new method for non-parametric multivariate analysis of variance. *Austral Ecol.* 2001;26(1):32–46.
- [19] Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Bioinformatics* [Internet].; 2014 Feb [cited 2022 Apr 4]. Available from: 10.1101/002832
- [20] Segata N, Izard J, Waldron L, et al. Metagenomic biomarker discovery and explanation. *Genome Biol.* 2011 Jun 24;12(6):R60.
- [21] Blankenberg D, Von Kuster G, Coraor N, et al. Galaxy, a web-based genome analysis tool for experimentalists. *Curr Protoc Mol Biol.* 2010 Jan;89(1). doi:10.1002/0471142727.mb1910s89.
- [22] Parks DH, Tyson GW, Hugenholtz P, et al. STAMP: statistical analysis of taxonomic and functional profiles. *Bioinformatics.* 2014 Nov 1;30(21):3123–3124.



- [23] Liaw A, Wiener M. Classification and regression by randomForest. 2002;2:5. <https://cogns.northwestern.edu/cbmg/LiawAndWiener2002.pdf>
- [24] Hastie T. The elements of statistical learning: data mining, inference and prediction. [Internet]. Universitas Indonesia Library. Springer; 2017 cited 2022 Apr 4. Available from: <https://lib.ui.ac.id>
- [25] Kursa MB, Rudnicki WR. Feature selection with the boruta package. J Stat Softw. 2010 Sep 16;36:1-13.
- [26] Breiman L. Random Forests. Mach Learn. 2001 Oct 1;45(1):5-32.
- [27] van Gastel J, Quirynen M, Teughels W, et al. Longitudinal changes in microbiology and clinical periodontal variables after placement of fixed orthodontic appliances. J Periodontol. 2008 Nov;79(11):2078-2086.
- [28] Uzuner FD, Kaygisiz E, Cankaya ZT. Effect of the bracket types on microbial colonization and periodontal status. Angle Orthod. 2014 Nov;84(6):1062-1067.
- [29] Hadj-Hamou R, Senok AC, Athanasios AE, et al. Do probiotics promote oral health during orthodontic treatment with fixed appliances? A systematic review. BMC Oral Health. 2020 Apr 25;20(1):126.
- [30] Benic GZ, Farella M, Morgan XC, et al. Oral probiotics reduce halitosis in patients wearing orthodontic braces: a randomized, triple-blind, placebo-controlled trial. J Breath Res. 2019 May 31;13(3):036010.
- [31] Habib S. Assessment of the therapeutic potential of a dental probiotic in orthodontic patients affected by gingivitis: a randomized control trial [Internet] [Thesis]. 2016 cited 2022 Apr 4. Available from: <https://tspace.library.utoronto.ca/handle/1807/79737>
- [32] Jivraj FE. Effectiveness of the lorodent probiotic lozenge in reducing plaque and *S. mutans* levels in orthodontic patients: a randomized, double-blind, placebo-controlled trial [Internet] [Thesis]. 2015 cited 2022 Apr 4. Available from: <https://tspace.library.utoronto.ca/handle/1807/70287>
- [33] Jose JE, Padmanabhan S, Chitharanjan AB. Systemic consumption of probiotic curd and use of probiotic toothpaste to reduce *Streptococcus mutans* in plaque around orthodontic brackets. Am J Orthod Dentofacial Orthop. 2013 Jul;144(1):67-72.
- [34] Alp S, Baka ZM. Effects of probiotics on salivary *Streptococcus mutans* and *Lactobacillus* levels in orthodontic patients. Am J Orthod Dentofacial Orthop. 2018 Oct;154(4):517-523.
- [35] Pinto GS, Cenci MS, Azevedo MS, et al. Effect of yogurt containing *Bifidobacterium animalis* subsp. *lactis* DN-173010 probiotic on dental plaque and saliva in orthodontic patients. Caries Res. 2014;48(1):63-68.
- [36] Kado I, Hisatsune J, Tsuruda K, et al. The impact of fixed orthodontic appliances on oral microbiome dynamics in Japanese patients. Sci Rep. 2020 Dec 15;10(1):21989.
- [37] Shokeen B, Vilorio E, Duong E, et al. The impact of fixed orthodontic appliances and clear aligners on the oral microbiome and the association with clinical parameters: a longitudinal comparative study. Am J Orthod Dentofacial Orthop. 2022 Mar 2;(22):S0889-5406. DOI:10.1016/j.ajodo.2021.10.015
- [38] Agossa K, Dubar M, Lemaire G, et al. Effect of *Lactobacillus reuteri* on gingival inflammation and composition of the oral microbiota in patients undergoing treatment with fixed orthodontic appliances: study protocol of a randomized control trial. Pathogens. 2022 Feb;11(2):112.
- [39] Kuru BE, Laleman I, Yalnızoğlu T, et al. The Influence of a *Bifidobacterium animalis* probiotic on gingival health: a randomized controlled clinical trial. J Periodontol. 2017;88(11):1115-1123.
- [40] Toiviainen A, Jalasvuori H, Lahti E, et al. Impact of orally administered lozenges with *Lactobacillus rhamnosus* GG and *Bifidobacterium animalis* subsp. *lactis* BB-12 on the number of salivary mutans streptococci, amount of plaque, gingival inflammation and the oral microbiome in healthy adults. Clin Oral Investig. 2015;19(1):77-83.
- [41] Alkaya B, Laleman I, Keceli S, et al. Clinical effects of probiotics containing *Bacillus* species on gingivitis: a pilot randomized controlled trial. J Periodontol Res. 2017 Jun;52(3):497-504.
- [42] Iniesta M, Herrera D, Montero E, et al. Probiotic effects of orally administered *Lactobacillus reuteri*-containing tablets on the subgingival and salivary microbiota in patients with gingivitis. A randomized clinical trial. J Clin Periodontol. 2012 Aug;39(8):736-744.
- [43] Hallström H, Lindgren S, Yucel-Lindberg T, et al. Effect of probiotic lozenges on inflammatory reactions and oral biofilm during experimental gingivitis. Acta Odontol Scand. 2013 Jul;71(3-4):828-833.
- [44] Kohar N, Emmanuel V, Astuti L. Comparison between probiotic lozenges and drinks towards periodontal status improvement of orthodontic patients. Dental J (Majalah Kedokteran Gigi). 2015 Sep 1; 48(3):126.
- [45] Shah SS, Nambiar S, Kamath D, et al. Comparative evaluation of plaque inhibitory and antimicrobial efficacy of probiotic and chlorhexidine oral rinses in orthodontic patients: a randomized clinical trial. Int J Dent. 2019 Feb 20;e1964158. <https://doi.org/10.1155/2019/1964158>
- [46] Vivekananda MR, Vandana KL, Bhat KG. Effect of the probiotic *Lactobacilli reuteri* (Prodentis) in the management of periodontal disease: a preliminary randomized clinical trial. J Oral Microbiol. 2010 Nov 2;2:10.3402/jom.v2i0.5344.
- [47] Caglar E, Kavaloglu SC, Kuscü OO, et al. Effect of chewing gums containing xylitol or probiotic bacteria on salivary mutans streptococci and lactobacilli. Clin Oral Investig. 2007 Dec;11(4):425-429.
- [48] Nikawa H, Makihira S, Fukushima H, et al. *Lactobacillus reuteri* in bovine milk fermented decreases the oral carriage of mutans streptococci. Int J Food Microbiol. 2004 Sep 1;95(2):219-223.
- [49] Azad MAK, Sarker M, Wan D. Immunomodulatory effects of probiotics on cytokine profiles. Biomed Res Int. 2018 Oct 23:e8063647. <https://www.hindawi.com/journals/bmri/2018/8063647/>
- [50] Morita H, Toh H, Fukuda S, et al. Comparative genome analysis of *Lactobacillus reuteri* and *Lactobacillus fermentum* reveal a genomic Island for reuterin and cobalamin production. DNA Res. 2008 Jun 30;15(3):151-161.
- [51] Chen C. Distribution of a newly described species, *Kingella oralis*, in the human oral cavity. Oral Microbiol Immunol. 1996 Dec;11(6):425-427.
- [52] McDowell A, Patrick S, Eishi Y, et al. *Propionibacterium acnes* in human health and disease. Biomed Res Int. 2013;2013:493564.
- [53] Oliveira RRDS, Fermiano D, Feres M, et al. Levels of candidate periodontal pathogens in subgingival biofilm. J Dent Res. 2016 Jun;95(6):711-718.
- [54] Ehrmann E, Handal T, Tamanai-Shacoori Z, et al. High prevalence of  $\beta$ -lactam and macrolide resistance genes in human oral *Capnocytophaga* species. J Antimicrob Chemother. 2014 Feb;69(2):381-384.

- [55] Jing D, Hao J, Shen Y, et al. Effect of fixed orthodontic treatment on oral microbiota and salivary proteins. *Exp Ther Med.* 2019 May;17(5):4237-4243.
- [56] Dassi E, Ferretti P, Covello G, et al. The short-term impact of probiotic consumption on the oral cavity microbiome. *Sci Rep.* 2018 Jul 11;8(1):10476.
- [57] Dassi E, Ballarini A, Covello G. HTM-CMB2013. Quattrone A, Jousson O, et al. Enhanced microbial diversity in the saliva microbiome induced by short-term probiotic intake revealed by 16S rRNA sequencing on the IonTorrent PGM platform. *J Biotechnol.* 2014 Nov 20;190:30-39.

# Short- term effect of probiotic Lactobacillus reuteri consumption on the salivary microbiome profile of subjects undergoing orthodontic treatment with fixed appliances

---

ORIGINALITY REPORT

---

18%

SIMILARITY INDEX

11%

INTERNET SOURCES

15%

PUBLICATIONS

4%

STUDENT PAPERS

---

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

---

3%

★ [www.x-mol.com](http://www.x-mol.com)

Internet Source

---

Exclude quotes  On

Exclude matches  < 15 words

Exclude bibliography  On



# Short- term effect of probiotic Lactobacillus reuteri consumption on the salivary microbiome profile of subjects undergoing orthodontic treatment with fixed appliances

GRADEMARK REPORT

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11