

Cover Jurnal



- Users Online: 59
-  [Print this page](#)
-  [Email this page](#)

-  [Home](#)
- [About](#)
 - [About Journal](#)
 - [Editorial Board](#)
- [Articles](#)
 - [Ahead of Print](#)
 - [Current Issue](#)
 - [Archives](#)
- [Authors](#)
 - [Submit Article](#)
 - [Instructions](#)
- [Search](#)
 - [Simple Search](#)
 - [Advanced Search](#)
 - [Image Search](#)
 - [Medline Search](#)
- [Subscribe](#)
- [Contact Us](#)
- [Reader Login](#)
 - [Sign Up](#)

Editorial Board

Editor-in-Chief:
Dr. Armelia Sari Widyarman
Department of Microbiology, Faculty of Dentistry, Trisakti University, Jakarta, Indonesia
[\[email protected\]](#)
[\[email protected\]](#)

Executive Editor:
Dr. Rosalina Tjandrawinata,
Department of Dental Material, Faculty of Dentistry, Trisakti University, Jakarta, Indonesia
[\[email protected\]](#)

Editorial Board:
Dr. Carolina Marpaung
Department of Prosthodontics, Faculty of Dentistry, Trisakti University, Jakarta, Indonesia
[\[email protected\]](#)

Dr. Chaminda Jayampath Seneviratne,
Faculty of Dentistry, National University of Singapore, Singapore
[\[email protected\]](#)

Prof. SG Damlee
Pediatric and Preventive Dentistry, Maharishi Markandeshwar University, India
[\[email protected\]](#)

Dr. Risa Chaisuparat
Associate Professor, Department of Oral Pathology,
Faculty of Dentistry, Chulalongkorn University, Thailand
[\[email protected\]](#)

Prof. Sittichai Koontongkaew
Department of Oral Biology, Faculty of Dentistry, Thammasat University, Thailand
[\[email protected\]](#)

Prof. Rahmi Amtha
Department of Oral Medicine, Faculty of Dentistry, Trisakti University, Indonesia
[\[email protected\]](#)

Dr. M. Orliando Roeslan
Department of Histology, Faculty of Dentistry, Trisakti University, Indonesia
[\[email protected\]](#)

Dr. Muhammad Ihsan Rizal
Deparment of Biochemistry and Molecular Biology, Division of Oral Biology, Faculty of Dentistry, Trisakti University, Indonesia
[\[email protected\]](#)

Administration:
Dr. Enrita Dian Rahmadini
Department of Pedodontics, Faculty of Dentistry, Trisakti University, Indonesia
[\[email protected\]](#)

Mega Cahyati
Faculty of Dentistry, Trisakti University, Indonesia
[\[email protected\]](#)

Mario Richi
MiCORE Laboratory, Faculty of Dentistry, Trisakti University, Indonesia
[\[email protected\]](#)

Moh Shidqon
Faculty of Economics and Business, Trisakti University, Indonesia
[\[email protected\]](#)

About the journal

- [POPULAR ARTICLES](#)
- [JOIN AS REVIEWER](#)
- [GET EMAIL ALERTS](#)
- [RECOMMEND](#)

- [Sitemap](#)
- [What's New](#)
- [Feedback](#)
- [Disclaimer](#)

- Users Online: 59
- Print this page
- Email this page

- Home
- About
 - About Journal
 - Editorial Board
- Articles
 - Ahead of Print
 - Current Issue
 - Archives
- Authors
 - Submit Article
 - Instructions
- Search
 - Simple Search
 - Advanced Search
 - Image Search
 - Medline Search
- Subscribe
- Contact Us
- Login
 - Sign Up

Export selected to

☐ Endnote

☐ Reference Manager

☐ Procite

☐ Medlars Format

☐ RefWorks Format

☐ BibTex Format

Apply

[Next Issue](#)
[Previous Issue](#)

- [POPULAR ARTICLES](#)
- [JOIN AS REVIEWER](#)
- [GET EMAIL ALERTS](#)
- [RECOMMEND](#)

Table of Contents

Coverpage

September-December 2021

Volume 5 | Issue 3

Page Nos. 101-154

View issue as eBook

Issue statistics

RSS

Online since Monday, October 18, 2021

Accessed 8,958 times.

PDF access policy

Journal allows immediate open access to content in HTML + PDF




Show all abstracts

Show selected abstracts

Export selected to ▼


Add to my list

ORIGINAL ARTICLES

	Effect of <i>Clinacanthus nutans</i> leaf extract on oral mucosal burns and tongue wounds: An <i>in-vivo</i> study Moehamad Orliando Roeslan, Salsabila Hany Reskia, Fathin Raihanah Firman, Hastania Sakti Dewa, Komariah Komariah, Del Afriadi Bustami DOI:10.4103/SDJ.SDJ_92_21 [ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}	p. 101
	Anterior tooth trauma among schoolchildren aged 10—19 years in Mewat district, India Amita Sharma, Sarika Sharma, Sonam Sharma, Rachit Sharma DOI:10.4103/SDJ.SDJ_90_21 [ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}	p. 110
	A comparison between panoramic radiography and dental study model for space maintainer Muhammad Chair Effendi, Berty Pramartika, Dini Rachmawati DOI:10.4103/SDJ.SDJ_82_21 [ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}	p. 115
	Root canal cleaning effectiveness with passive ultrasonic irrigation: A comparative <i>in vitro</i> study between Irrisafe and EndoUltra tip Erlando Sartiono, Bernard Iskandar, Aryadi, Widijanto Sudhana DOI:10.4103/SDJ.SDJ_65_21 [ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}	p. 122
	Antimicrobial efficacy of copper nanoparticles against <i>Aggregatibacter actinomycetemcomitans</i> and <i>Porphyromonas gingivalis</i>: An <i>in-vitro</i> study Swapna Arunkumar Mahale, Shraddha Rajendra Shimpi, Kunal S Sethi, Dipali Dhanraj Chaudhari, Pooja S Kadam, Ankita A Katkurwar DOI:10.4103/SDJ.SDJ_84_21 [ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}	p. 128
	Peppermint flavor oil in fluoride varnishes enhances fluoride release Yosi Kusuma Eriwati, Dhea Putriani, Karen Geraldine, Heri Hermansyah DOI:10.4103/SDJ.SDJ_78_21 [ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}	p. 133
	The effect of garlic extract (<i>Allium sativum</i> L.) (Amaryllidaceae) to eradicate the <i>Porphyromonas endodontalis</i> biofilm: An <i>in-vitro</i> research Adi Dharma Widjaya, Meiny Faudah Amin, Aryadi, Boedi Oetomo Roeslan DOI:10.4103/SDJ.SDJ_79_21 [ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}	p. 138
	Relationship between nutritional status and mandibular length in subjects aged 10–16 years Hilda Fitria Lubis, Rachel Tiong DOI:10.4103/SDJ.SDJ_32_20 [ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}	p. 144

CASE REPORT



	Challenges in the management of oral manifestations in a patient with limited systemic sclerosis Yannie Febby Martina Lefaan, Riani Setiadhi DOI:10.4103/SDJ.SDJ_93_21 [ABSTRACT] [HTML Full text] [PDF] [Mobile Full text] [EPub] [Sword Plugin for Repository]^{Beta}	p. 148
--	---	-------------------

- [Sitemap](#)
- [What's New](#)
- [Feedback](#)
- [Disclaimer](#)

Effect of *Clinacanthus nutans* Leaf Chloroform Extract on Reducing Nitric Oxide in Fibroblasts

Erdita Hapsari, Rezky Anggraeni¹, Komariah Komariah¹, Moehamad Orliando Roeslan¹

Student of Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia, ¹Department of Oral Biology, Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia

ABSTRACT

Background: *Clinacanthus nutans* or dandang gendis is a plant that can be an effective herbal treatment for inflammation, in addition the Indonesian people are still looking for alternative treatments. Research shows that the antiinflammatory properties of *C. nutans* leaves are effective in reducing inflammation against fibroblasts. Nitric oxide (NO) is one of the inflammatory mediators, that are released by fibroblast when inflamed. **Objective:** To observe the effect of *C. nutans* leaf chloroform extract on reducing NO in inflamed fibroblasts. **Methods:** Fibroblasts stimulated with *Escherichia coli* lipopolysaccharide were treated with chloroform extracts of *C. nutans* leaves at concentrations of 25, 50, 100, and 200 µg/mL for 24 h, then a NO inhibition test was performed. **Result:** The *C. nutans* leaf extract at a concentration of 200 µg/mL resulted in significantly ($P < 0.05$) lower NO production than the negative control. **Conclusion:** *C. nutans* leaf chloroform extract has the effect of reducing NO in fibroblasts. This extract has the potential as an antiinflammatory agent.

KEYWORDS: Dandang gendis (*Clinacanthus nutans*), fibroblasts, nitric oxide

Received: 07-Feb-2023
Revised: 03-Aug-2023
Accepted: 04-Dec-2023
Published: 31-Jan-2024

BACKGROUND

Inflammation, a biological reaction of the immune system, may be caused by pathogens, injured cells, and poisonous substances.^[1] Inflammation is the result of prolonged exposure to stimulation or an inappropriate reaction to the cell's molecules and may result in an acute or chronic phase, where tissue damage and fibrosis may occur.^[2] The acute phase is when inflammation occurs for a short time and functions as the body's defense against infection—where immune cells migrate to the injured area, facilitated by soluble mediators, such as cytokines, chemokines, and proteins. If the inflammation occurs repeatedly, it progresses to the second stage—chronic inflammation, which lasts for a long period and can trigger various chronic diseases, such as arthritis, cancer, cardiovascular disease, diabetes, neurological disease, and respiratory disease.^[3] Inflammatory conditions depend on the severity,

chronicity, and mechanisms involved in the process, as well as the species and capacity of the individual's immune system to respond and adapt. Inflammation has a role in the development of several diseases.^[4] Most human diseases have an inflammatory component and inflammatory mediators, such as inducible nitric oxide synthase (iNOS), which produces nitric oxide (NO) from l-arginine.^[5] NO is a reactive gas molecule that has several functions in biological systems, depending on its concentration. At low concentrations, NO functions as a signaling molecule, whereas at high concentrations, it becomes highly toxic due to its ability to react with several cellular targets.^[6] NO is a potent vasodilatory and antiinflammatory signaling molecule.^[7] NO is also

Address for correspondence: Dr. Moehamad Orliando Roeslan, Department of Oral Biology, Faculty of Dentistry, Universitas Trisakti, Jakarta 11440, Indonesia.
E-mail: orliando.roeslan@trisakti.ac.id

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Hapsari E, Anggraeni R, Komariah K, Roeslan MO. Effect of *Clinacanthus nutans* leaf chloroform extract on reducing nitric oxide in fibroblasts. *Sci Dent J* 2023;7:120-4.

Access this article online

Quick Response Code:



Website:
www.scidentj.com

DOI:
10.4103/SDJ.SDJ_9_23

an inflammatory mediator^[8] formed by iNOS, which has many homeostatic functions and plays an important role in inflammation. Astrocytes and microglia in an inflamed brain produce high NO levels during inflammation. When inflammation occurs, NO can cause vasodilation and increased leukocyte adhesion.^[9] There are several nitrovasodilator compounds capable of releasing NO, such as glyceryl trinitrate, isosorbide dinitrate, sodium nitroprusside, and s-nitroso glutathione.^[10]

Fibroblasts are cells in connective tissue that play a major role in wound healing. When fibroblasts are injured, NO is secreted and provides an antiinflammatory effect.^[11] When a tissue experiences an inflammatory state, fibroblasts move toward the wound to proliferate and synthesize collagen and an extracellular matrix, and then a new epithelium forms on the mucosa to cover the wound. Fibroblast proliferation indicates rapid healing.^[12] Fibroblasts also assume the role of inflammatory cells and recruit leukocytes, promote angiogenesis, and enable chronic inflammation in tissues.^[13]

Clinacanthus nutans is a plant that belongs to the family Acanthaceae. This plant has antiinflammatory properties,^[14] and the leaf extract can be an effective alternative herbal treatment.^[15] *C. nutans* has attracted worldwide attention for its hidden therapeutic potential.^[16] The biological activities of *C. nutans* include antioxidant, antiinflammatory, anticancer, antidiabetic, antiviral, antimicrobial, and antiproliferative effects. As such, it is used to treat several antiinflammatory conditions.^[17]

Research on the effect of *C. nutans* extracts on NO in fibroblasts has not been conducted. Therefore, this study aimed to investigate the effect of the chloroform extract of *C. nutans* leaves on NO production in fibroblasts.

MATERIALS AND METHODS

This study was an *in vitro* laboratory experimental research to investigate the effects of *C. nutans* chloroform extract on NO in fibroblasts. The study location was the BioCore Laboratory, Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia.

The leaves of *C. nutans*, obtained from the Indonesian Spice and Medicinal Crops Research Institute, Bogor, Indonesia, were washed and dried. Then, the extraction stage was performed by the maceration method using 85 g of *C. nutans* leaf powder and 425 mL of chloroform solvent with a ratio of 1:5 at room temperature. Periodic agitation was performed during maceration for 3 days

(every 15 min for 8 h/day). Then the suspension was filtered using Whatman filter paper no. 1, and the filtrate was evaporated using a rotary evaporator to separate the chloroform from the extract at a set temperature of 61°C. This procedure was repeated three times.

Fibroblasts were cultured in Dulbecco's Modified Eagle Medium (Gibco, New York, NY, USA) supplemented with 20% fetal bovine serum (Gibco), and 1% amphotericin B/1% penicillin–streptomycin (Invitrogen, Waltham, MA, USA) at 37°C in a 5% CO₂ incubator. The fibroblasts were subcultured when the cells reached 80%–90% confluency in the flask.

Fibroblasts were incubated for 24 h at 37°C and 5% CO₂. The medium was then aspirated, and cells were stimulated with 2 µg/mL of *Escherichia coli* lipopolysaccharide (LPS, 0111:B4, Sigma–Aldrich, Burlington, MA, United States) for 24 h. Then the medium was replaced with various concentrations of the extract (25, 50, 100, and 200 µg/mL) and genistein 1000 µM as a positive control. The concentration of nitrite, a stable metabolite of NO, in the supernatant, was measured as an indicator of NO production using an NO detection kit (Griess Reagent System, Promega, Madison, WI, USA), according to the manufacturer's instructions. The 96-well plate containing 50 µL of supernatant and Griess reagent was incubated at room temperature for 10 min. The absorbance was then measured in a microplate reader at a wavelength of 490 nm. The nitric concentration is determined from the standard sodium nitrite curve.

The data obtained will be presented as mean ± standard deviation. The results of the detection of NO secretion in fibroblasts, which were analyzed using the NO detection kit, were then tested for normality using the Shapiro–Wilk method. If the results of the normality test showed that the distribution of the data was not normal, a nonparametric test was conducted. If the results of the test showed normal data ($P > 0.05$), then the analysis proceeded to a one-way Analysis of variance test. If the results showed a significant difference ($P < 0.05$), then Tukey's *post hoc* test was indicated.

RESULTS

In this study, the variables used were *C. nutans* leaf chloroform extracts with respective concentrations of 25, 50, 100, and 200 µg/mL. The negative control was fibroblasts with 2 µg/mL LPS *E. coli* added without treatment, whereas the positive control used Genistein 1000 µM. Each variable was tested for its ability to determine the effect of *C. nutans* leaf chloroform extract on inflamed fibroblasts.




Figure 1: Bar graph of nitric oxide inhibition test results with the chloroform extract of *C. nutans* leaves against inflamed fibroblasts. *Indicates a significant difference from the negative control ($P < 0.05$)

The results revealed that *C. nutans* at a concentration of 200 µg/mL, and the positive control group had a significant difference in NO production compared with the negative control group [Figure 1]. Other concentrations (25, 50, and 100 µg/mL) have no significant difference compared with the negative control group.

DISCUSSION

The biological activities of *C. nutans* leaves are antioxidant, antiinflammatory, anticancer, antidiabetic and antiviral, antimicrobial, and antiproliferative effects.^[18] In previous studies, *C. nutans* leaves indicated the presence of flavonoids.^[19] In addition, the results of phytochemical tests performed in previous studies indicated that the chloroform extract of *C. nutans* leaves has the highest total flavonoid and phenolic content compared to extracts with other solvents.^[20]

In this study, *C. nutans* leaf chloroform extract at a concentration of 200 µg/mL produced lower levels of NO compared with the negative control. The positive control group (Genistein 1000 µM) also produced lower NO compared with the negative control group. At concentrations of 25, 50, and 100 µg/mL, the extracts did not produce lower NO levels compared with the negative control. This implies that *C. nutans* leaf chloroform extract requires a minimum concentration of 200 µg/mL to reduce NO production in fibroblasts stimulated by *E. coli*.

A previous study indicated that *C. nutans* leaf chloroform extract with a concentration of 100 µg/

mL can reduce NO production in macrophages.^[19] However, in this study, NO production decreased at a concentration of 200 µg/mL. This difference may be due to variations in the cell types used; this study utilized fibroblasts, whereas the previous research used macrophages. Differences in extraction methods could also account for variations in results. The previous study employed the Soxhlet method for extraction, whereas this study used the maceration method. Variances in extraction methods can influence the compounds present in the extract.^[19] In this study, fibroblasts were stimulated with 2 µg/mL LPS, whereas in the previous study, macrophages were stimulated with 100 ng/mL of LPS.^[19] The difference in concentration is due to differences in the cells used. This difference in LPS concentration caused a difference in the amount of NO produced by cells.

The anti-inflammatory activity of *C. nutans* can be attributed to its flavonoid content. Flavonoids can inhibit the tyrosine kinase enzyme by binding to adenosine triphosphate in the active enzyme.^[20] Inhibition of this enzyme can result in a downregulation of various proteins, such as NO synthase, cyclooxygenase, and lipoxygenase, which can stop the proliferation of inflammatory cells and inhibit inflammation.^[21] The phenolic compounds contained in the chloroform extract of *C. nutans* leaves also play an active role as an antiinflammatory. The antiinflammatory activity of its natural compounds exerts its biological properties by blocking two main signaling pathways: the nuclear factor kappa B (NF-κB) and mitogen-activated protein kinase pathways, which play a major role in the production of various proinflammatory mediators.^[22]

In addition, the phenolic compounds in the chloroform extract of *C. nutans* leaves also have the highest levels of antioxidant activity.^[23] In a normal and healthy body, there is a balance between the formation of reactive oxygen/free radicals and endogenous antioxidant defense mechanisms. However, if this balance is disturbed, it can cause oxidative stress, which in turn can result in injury to all cellular components (such as DNA, proteins, and membrane lipids) and cell death. Antioxidants based on natural compounds play a preventive role in protecting against the formation of free radicals. Antioxidant compounds can inhibit and prevent oxidation processes that can produce free radicals and break chains that can damage cells and tissues; hence, these compounds can be used as a therapeutic option for inflammation.^[24]

In general, antiinflammatory properties can suppress the NF-κB pathway and the production of iNOS and proinflammatory cytokines, such as interleukin-1β

and tumor necrosis factor alpha.^[25] Inflammation is prevented by the binding of LPS to toll-like receptor 4 (TLR4). Simultaneously, LPS and TLR4 are catalyzed by lipopolysaccharide-binding protein (LBP) and cluster of differentiation 14 (CD14). After LPS is released, it LPS binds to LBP to form LPS-LBP and targets CD14 on macrophages. LBP catalyzes LPS to CD14 so that LPS combines with CD14, which transfers LPS to TLR4 to then activate the TLR pathway and be directed at inflammatory factors. LPS that binds to TLRs can be neutralized by adenosine monophosphate and can inhibit the release of inflammatory factors.^[24] Another antiinflammatory mechanism is the inhibition and regulation of inflammatory-related signaling pathways. Signal transduction is dependent on a toll-interleukin-1 receptor (TIR) domain-containing adapter protein (TRIF/TIR domain-containing adaptor-inducing interferon- β) associated with endocytosis of activated TLR4. LPS binding to TLRs activated by TLR4 can also activate the NF- κ B and mitogen-activated protein kinase pathways, where inhibition of TLR4 endocytosis can be performed to suppress inflammatory pathways.^[26]

In a previous study, *C. nutans* extracts showed positive results on a fibroblast migration assay compared to *Strobilanthes crispus* extracts. Chloroform extract of *C. nutans* has also been demonstrated to accelerate healing, including in inflammatory processes, compared with acetone and ethanol extracts. However, *C. nutans* extracts showed less antimicrobial activity against *Staphylococcus aureus*.^[27] In addition, previous studies have shown that chloroform extract of *C. nutans* is the most active at inhibiting NO production compared with hexane and ethanol extracts.^[19] In addition, chloroform extracts of *C. nutans* and aloe vera can upregulate the synthesis of basic fibroblast growth factor (bFGF) in wound healing.^[28] Another previous study showed that *C. nutans* leaf could increase fibroblast count in mucosal burn and tongue wounds *in vivo*.^[29] This proves that the chloroform extract of *C. nutans* leaves is more effective in reducing or inhibiting inflammation.

During inflammation, macrophages and fibroblasts are the main source of NO synthesis.^[30] NO has a small size and lacks charge, so the penetration of NO molecules through cell membranes and subcellular organelles is high. Moreover, NO can be transferred by S-nitrosothiols through cells with protein binding. Therefore, when inflammation occurs, NO is not only in the cells where it is produced but also in the surrounding cells. Among the immune cells that are involved in inflammation, macrophages synthesize the most NO.^[31] Several other factors influence the effect of the chloroform extract

of *C. nutans* leaves on inflamed fibroblasts, such as differences in the metabolite compounds contained in the extract, the amount of extract concentration used, and the extraction method.^[32,33]

The limitation of this study was the use of a limited concentration range. If concentrations above 200 μ g/mL and up to 1000 μ g/mL were used, we could have determined definitively whether 200 μ g/mL is indeed the minimum concentration needed to reduce NO production in fibroblasts.

CONCLUSION

Based on the study results, the chloroform extract of *C. nutans* leaves at a concentration of 200 μ g/mL has the effect of reducing NO synthesis in inflamed fibroblasts. This extract has the potential to be an antiinflammatory agent.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Varela ML, Mogildea M, Moreno I, Lopes A. Acute inflammation and metabolism. *Inflammation* 2018;41:1115-27.
- Germolec DR, Shipkowski KA, Frawley RP, Evans E. Markers of inflammation. *Methods Mol Biol* 2018;1803:57-79.
- Kunnumakkara AB, Sailo BL, Banik K, Harsha C, Prasad S, Gupta SC, et al. Chronic diseases, inflammation, and spices: How are they linked? *J Transl Med* 2018;16:1-25.
- Strandberg TE, Nieminen T. Future perspectives on the role of frailty in cardiovascular diseases. *Adv Exp Med Biol* 2020;1216:149-52.
- Wang R, Ghahary A, Shen YJ, Scott PG, Tredget EE. Human dermal fibroblasts produce nitric oxide and express both constitutive and inducible nitric oxide synthase isoforms. *J Invest Dermatol* 1996;106:419-27.
- Salas A, Cabrera JJ, Jiménez-Leiva A, Mesa S, Bedmar EJ, Richardson DJ, et al. Bacterial nitric oxide metabolism: Recent insights in rhizobia. *Adv Microb Physiol* 2021;78:259-315.
- Cyr AR, Huckaby LV, Shiva SS, Zuckerbraun BS. Nitric oxide and endothelial dysfunction. *Crit Care Clin* 2020;36:307-21.
- Mo X, Chen J, Wang X, Pan Z, Ke Y, Zhou Z, et al. Krüppel-like factor 4 regulates the expression of inducible nitric oxide synthase induced by TNF- α in human fibroblast-like synoviocyte MH7A cells. *Mol Cell Biochem* 2018;438:77-84.
- Liy PM, Puzi NNA, Jose S, Vidyadaran S. Nitric oxide modulation in neuroinflammation and the role of mesenchymal stem cells. *Exp Biol Med* 2021;246:2399-406.
- da Silva GM, da Silva MC, Nascimento DVG, Lima Silva EM, Gouvêa FFF, de França Lopes LG, et al. Nitric oxide as a central molecule in hypertension: Focus on the vasorelaxant activity of new nitric oxide donors. *Biology (Basel)* 2021;10:1041.
- Kwesiga MP, Cook E, Hannon J, Wayward S, Gwaltney C, Rao S, et al. Investigative study on nitric oxide production in human dermal fibroblast cells under normal and high glucose conditions. *Med Sci* 2018;6:99.

12. Ardiana T, Rizkia A, Kusuma P, Firdausy MD, Lqwr G, Urxs J, et al. Efektivitas daun jambu biji terhadap jumlah sel fibroblas pasca luka pencabutan gigi pada marmut (*Cavia cobaya*). ODONTO Dent J 2015;2:64-70.
13. Marsh LJ, Kemble S, Reis Nisa P, Singh R, Croft AP. Fibroblast pathology in inflammatory joint disease. Immunol Rev 2021;302:163-83.
14. Kao MH, Wu JS, Cheung WM, Chen JJ, Sun GY, Ong WY, et al. *Clinacanthus nutans* mitigates neuronal death and reduces ischemic brain injury: Role of NF- κ B-driven IL-1 β transcription. Neuromolecular Med 2021;23:199-210.
15. Widjaja S, Rusdiana, Ichwan M. Enhanced cytotoxic effects of *Clinacanthus nutans* and doxorubicin in combination toward breast cancer cell lines. J Adv Pharm Technol Res 2021;12:152-6.
16. Zulkipli IN, Rajabalaya R, Idris A, Sulaiman NA, David SR. *Clinacanthus nutans*: A review on ethnomedicinal uses, chemical constituents and pharmacological properties. Pharm Biol 2017;55:1093-113.
17. Roeslan MO, Reskia SH, Firman FR, Dewa HS, Komariah K, Bustami DA. Effect of *Clinacanthus nutans* leaf extract on oral mucosal burns and tongue wounds: An in-vivo study. Sci Dent J 2022;5:101-9.
18. Goshi E, Zhou G, He Q. Nitric oxide detection methods in vitro and in vivo. Med Gas Res 2019;9:192-207.
19. Roeslan MO, Ayudhya TDN, Yingyongnarongkul B, Koontongkaew S. Anti-biofilm, nitric oxide inhibition and wound healing potential of purpurin-18 phytyl ester isolated from *Clinacanthus nutans* leaves. Biomed Pharmacother = Biomed Pharmacotherap 2019;113:108724.
20. Al-Khayri JM, Sahana GR, Nagella P, Joseph BV, Alessa FM, Al-Mssallem MQ. Flavonoids as potential anti-inflammatory molecules: A review. Molecules 2022;27:2901.
21. Islam MI, Shanta MA, Mondal M, Hoque N, Majumder S, Ahmed T, et al. Protective effect of chloroform extract of *Stereospermum chelonoides* bark against amyloid beta42 induced cell death in SH-SY5Y cells and against inflammation in Swiss albino mice. J Basic Clin Physiol Pharmacol 2018;29:621-30.
22. Arulselvan P, Fard MT, Tan WS, Gothai S, Fakurazi S, Norhaizan ME, et al. Role of antioxidants and natural products in inflammation. Oxid Med Cell Longevity 2016;2016:5276130.
23. Alam A, Ferdosh S, Ghafoor K, Hakim A, Juraimi AS, Khatib A, et al. *Clinacanthus nutans*: A review of the medicinal uses, pharmacology and phytochemistry. Asian Paci J Trop Med 2016;9:402-9.
24. Ikrima K, Amalia R, Levita J. Peran spesies oksigen reaktif pada inflamasi serta antioksidan alami sebagai fitoterapi. Farmaka 2020; 17:198-211.
25. Ong WY, Herr DR, Sun GY, Lin TN. Anti-inflammatory effects of phytochemical components of *Clinacanthus nutans*. Molecules 2022;27:3607.
26. Luo Y, Song Y. Mechanism of antimicrobial peptides: Antimicrobial, anti-inflammatory and antibiofilm activities. Int J Mol Sci 2021;22:11401.
27. Ban WK, Fong IL, Khong HY, Phung JHY. Wound healing, antimicrobial and antioxidant properties of *Clinacanthus nutans* (burm.f.) Lindau and *Strobilanthes crispus* (l) Blume extracts. Molecules 2022;27:1722.
28. Anggraeni R, Roeslan MO. Effects of *Clinacanthus nutans* and *Aloe vera* extracts on bFGF synthesis in fibroblasts. J Indones Dent Assoc 2022;5:21-6.
29. Roeslan MO, Reskia SH, Firman FR, Dewa HS, Komariah K, Bustami DA. Effect of *Clinacanthus nutans* leaf extract on oral mucosal burns and tongue wounds: An In-vivo study. Sci Dent J 2021;5:101-9.
30. Schäffer MR, Tantry U, Barbul A. Wound fluid inhibits wound fibroblast nitric oxide synthesis. J Surg Res 2004;122:43-8.
31. Holotiuk V, Kryzhanivska A, Churpiy I, Tataryn B, Ivasiutyn D. Role of nitric oxide in pathogenesis of tumor growth and its possible application in cancer treatment. Exp Oncol 2019;41:210-15.
32. Pn P, Ma H, Studi P, Kedokteran S, Dokter DP. Efek ekstrak metanol daun sirsak (*Annona muricata*) dalam menghambat pertumbuhan bakteri *Salmonella typhi* secara in vitro. E-J Medika Udayana 2021;10:2021.
33. Mangkulion S, Roeslan MO, Monthanpisut P. The potential of cogon grass (*Imperata cylindrica*) ethanol extract in inhibiting nitric oxide secretion in fibroblast Sci Dent J 2023;7:11-4.

Effect of Clinacanthus nutans Leaf Chloroform Extract on Reducing Nitric Oxide in Fibroblasts

by Moehamad Orliando Roeslan

Submission date: 04-Jul-2025 03:21PM (UTC+0700)

Submission ID: 2708664051

File name: Artikel_Effect_of_clinacanthus_nutans_leaf_chloroform.pdf (420.94K)

Word count: 3601

Character count: 20123

Original Article

Effect of *Clinacanthus nutans* Leaf Chloroform Extract on Reducing Nitric Oxide in Fibroblasts

Erdita Hapsari, Rezky Anggraeni¹, Komariah Komariah¹, Moehamad Orliando Roeslan¹

¹⁵
Student of Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia, ¹Department of Oral Biology, Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia

ABSTRACT

Background: *Clinacanthus nutans* or dandang gendis is a plant that can be an effective herbal treatment for inflammation, in addition the Indonesian people are still looking for alternative treatments. Research shows that the antiinflammatory properties of *C. nutans* leaves are effective in reducing inflammation against fibroblasts. Nitric oxide (NO) is one of the inflammatory mediators, that are released by fibroblast when inflamed. **Objective:** To observe the effect of *C. nutans* leaf chloroform extract on reducing NO in inflamed fibroblasts. **Methods:** Fibroblasts stimulated with *Escherichia coli* lipopolysaccharide were treated with chloroform extracts of *C. nutans* leaves at concentrations of 25, 50, 100, and 200 µg/mL for 24h, then a NO inhibition test was performed. **Result:** The *C. nutans* leaf extract at a concentration of 200 µg/mL resulted in significantly ($P < 0.05$) lower NO production than the negative control. **Conclusion:** *C. nutans* leaf chloroform extract has the effect of reducing NO in fibroblasts. This extract has the potential as an antiinflammatory agent.

KEYWORDS: Dandang gendis (*Clinacanthus nutans*), fibroblasts, nitric oxide

Received: 07-Feb-2023
Revised: 03-Aug-2023
Accepted: 04-Dec-2023
Published: 31-Jan-2024

BACKGROUND

Inflammation, a biological reaction of the immune system, may be caused by pathogens, injured cells, and poisonous substances.^[1] Inflammation is the result of prolonged exposure to stimulation or an inappropriate reaction to the cell's molecules and may result in an acute or chronic phase, where tissue damage and fibrosis may occur.^[2] The acute phase is when inflammation occurs for a short time and functions as the body's defense against infection—where immune cells migrate to the injured area, facilitated by soluble mediators, such as cytokines, chemokines, and proteins. If the inflammation occurs repeatedly, it progresses to the second stage—chronic inflammation, which lasts for a long period and can trigger various chronic diseases, such as arthritis, cancer, cardiovascular disease, diabetes, neurological disease, and respiratory disease.^[3] Inflammatory conditions depend on the severity,

⁶
chronicity, and mechanisms involved in the process, as well as the species and capacity of the individual's immune system to respond and adapt. Inflammation has a role in the development of several diseases.^[4] Most human diseases have an inflammatory component and inflammatory mediators, such as inducible nitric oxide synthase (iNOS), which produces nitric oxide (NO) from L-arginine.^[5] NO is a reactive gas molecule that has several functions in biological systems, depending on its concentration. At low concentrations, NO functions as a signaling molecule, whereas at high concentrations, it becomes highly toxic due to its ability to react with several cellular targets.^[6] NO is a potent vasodilatory and antiinflammatory signaling molecule.^[7] NO is also

¹⁶
Address for correspondence: Dr. Moehamad Orliando Roeslan, Department of Oral Biology, Faculty of Dentistry, Universitas Trisakti, Jakarta 11440, Indonesia. E-mail: orliando.roeslan@trisakti.ac.id

³
This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Hapsari E, Anggraeni R, Komariah K, Roeslan MO. Effect of *Clinacanthus nutans* leaf chloroform extract on reducing nitric oxide in fibroblasts. Sci Dent J 2023;7:120-4.

Access this article online	
Quick Response Code:	Website: www.scidentj.com
	DOI: 10.4103/SDJ.SDJ_9_23

an inflammatory mediator^[8] formed by iNOS, which has many homeostatic functions and plays an important role in inflammation. Astrocytes and microglia in an inflamed brain produce high NO levels during inflammation. When inflammation occurs, NO can cause vasodilation and increased leukocyte adhesion.^[9] There are several nitrovasodilator compounds capable of releasing NO, such as glyceryl trinitrate, isosorbide dinitrate, sodium nitroprusside, and s-nitroso glutathione.^[10]

Fibroblasts are cells in connective tissue that play a major role in wound healing. When fibroblasts are injured, NO is secreted and provides an antiinflammatory effect.^[11] When a tissue experiences an inflammatory state, fibroblasts move toward the wound to proliferate and synthesize collagen and an extracellular matrix, and then a new epithelium forms on the mucosa to cover the wound. Fibroblast proliferation indicates rapid healing.^[12] Fibroblasts also assume the role of inflammatory cells and recruit leukocytes, promote angiogenesis, and enable chronic inflammation in tissues.^[13]

Clinacanthus nutans is a plant that belongs to the family Acanthaceae. This plant has antiinflammatory properties,^[14] and the leaf extract can be an effective alternative herbal treatment.^[15] *C. nutans* has attracted worldwide attention for its hidden therapeutic potential.^[16] The biological activities of *C. nutans* include antioxidant, antiinflammatory, anticancer, antidiabetic, antiviral, antimicrobial, and antiproliferative effects. As such, it is used to treat several antiinflammatory conditions.^[17]

Research on the effect of *C. nutans* extracts on NO in fibroblasts has not been conducted. Therefore, this study aimed to investigate the effect of the chloroform extract of *C. nutans* leaves on NO production in fibroblasts.

MATERIALS AND METHODS

This study was an *in vitro* laboratory experimental research to investigate the effects of *C. nutans* chloroform extract on NO in fibroblasts. The study location was the BioCore Laboratory, Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia.

The leaves of *C. nutans*, obtained from the Indonesian Spice and Medicinal Crops Research Institute, Bogor, Indonesia, were washed and dried. Then, the extraction stage was performed by the maceration method using 85 g of *C. nutans* leaf powder and 425 mL of chloroform solvent with a ratio of 1:5 at room temperature. Periodic agitation was performed during maceration for 3 days

(every 15 min for 8 h/day). Then the suspension was filtered using Whatman filter paper no. 1, and the filtrate was evaporated using a rotary evaporator to separate the chloroform from the extract at a set temperature of 61°C. This procedure was repeated three times.

Fibroblasts were cultured in Dulbecco's Modified Eagle Medium (Gibco, New York, NY, USA) supplemented with 20% fetal bovine serum (Gibco), and 1% amphotericin B/1% penicillin-streptomycin (Invitrogen, Waltham, MA, USA) at 37°C in a 5% CO₂ incubator. The fibroblasts were subcultured when the cells reached 80%-90% confluency in the flask.

Fibroblasts were incubated for 24 h at 37°C and 5% CO₂. The medium was then aspirated, and cells were stimulated with 2 µg/mL of *Escherichia coli* lipopolysaccharide (LPS, 0111:B4, Sigma-Aldrich, Burlington, MA, United States) for 24 h. Then the medium was replaced with various concentrations of the extract (25, 50, 100, and 200 µg/mL) and genistein 1000 µM as a positive control. The concentration of nitrite, a stable metabolite of NO, in the supernatant, was measured as an indicator of NO production using an NO detection kit (Griess Reagent System, Promega, Madison, WI, USA), according to the manufacturer's instructions. The 96-well plate containing 50 µL of supernatant and Griess reagent was incubated at room temperature for 10 min. The absorbance was then measured in a microplate reader at a wavelength of 490 nm. The nitric concentration is determined from the standard sodium nitrite curve.

The data obtained will be presented as mean ± standard deviation. The results of the detection of NO secretion in fibroblasts, which were analyzed using the NO detection kit, were then tested for normality using the Shapiro-Wilk method. If the results of the normality test showed that the distribution of the data was not normal, a nonparametric test was conducted. If the results of the test showed normal data ($P > 0.05$), then the analysis proceeded to a one-way Analysis of variance test. If the results showed a significant difference ($P < 0.05$), then Tukey's *post hoc* test was indicated.

RESULTS

In this study, the variables used were *C. nutans* leaf chloroform extracts with respective concentrations of 25, 50, 100, and 200 µg/mL. The negative control was fibroblasts with 2 µg/mL LPS *E. coli* added without treatment, whereas the positive control used Genistein 1000 µM. Each variable was tested for its ability to determine the effect of *C. nutans* leaf chloroform extract on inflamed fibroblasts.




Figure 1: Bar graph of nitric oxide inhibition test results with the chloroform extract of *C. nutans* leaves against inflamed fibroblasts. *Indicates a significant difference from the negative control ($P < 0.05$)

The results revealed that *C. nutans* at a concentration of 200 µg/mL, and the positive control group had a significant difference in NO production compared with the negative control group [Figure 1]. Other concentrations (25, 50, and 100 µg/mL) have no significant difference compared with the negative control group.

DISCUSSION

The biological activities of *C. nutans* leaves are antioxidant, antiinflammatory, anticancer, antidiabetic and antiviral, antimicrobial, and antiproliferative effects.^[18] In previous studies, *C. nutans* leaves indicated the presence of flavonoids.^[19] In addition, the results of phytochemical tests performed in previous studies indicated that the chloroform extract of *C. nutans* leaves has the highest total flavonoid and phenolic content compared to extracts with other solvents.^[20]

In this study, *C. nutans* leaf chloroform extract at a concentration of 200 µg/mL produced lower levels of NO compared with the negative control. The positive control group (Genistein 1000 µM) also produced lower NO compared with the negative control group. At concentrations of 25, 50, and 100 µg/mL, the extracts did not produce lower NO levels compared with the negative control. This implies that *C. nutans* leaf chloroform extract requires a minimum concentration of 200 µg/mL to reduce NO production in fibroblasts stimulated by *E. coli*.

A previous study indicated that *C. nutans* leaf chloroform extract with a concentration of 100 µg/

mL can reduce NO production in macrophages.^[19] However, in this study, NO production decreased at a concentration of 200 µg/mL. This difference may be due to variations in the cell types used; this study utilized fibroblasts, whereas the previous research used macrophages. Differences in extraction methods could also account for variations in results. The previous study employed the Soxhlet method for extraction, whereas this study used the maceration method. Variances in extraction methods can influence the compounds present in the extract.^[19] In this study, fibroblasts were stimulated with 2 µg/mL LPS, whereas in the previous study, macrophages were stimulated with 100 ng/mL of LPS.^[19] The difference in concentration is due to differences in the cells used. This difference in LPS concentration caused a difference in the amount of NO produced by cells.

The anti-inflammatory activity of *C. nutans* can be attributed to its flavonoid content. Flavonoids can inhibit the tyrosine kinase enzyme by binding to adenosine triphosphate in the active enzyme.^[20] Inhibition of this enzyme can result in a downregulation of various proteins, such as NO synthase, cyclooxygenase, and lipoxygenase, which can stop the proliferation of inflammatory cells and inhibit inflammation.^[21] The phenolic compounds contained in the chloroform extract of *C. nutans* leaves also play an active role as an antiinflammatory. The antiinflammatory activity of its natural compounds exerts its biological properties by blocking two main signaling pathways: the nuclear factor kappa B (NF-κB) and mitogen-activated protein kinase pathways, which play a major role in the production of various proinflammatory mediators.^[22]

In addition, the phenolic compounds in the chloroform extract of *C. nutans* leaves also have the highest levels of antioxidant activity.^[23] In a normal and healthy body, there is a balance between the formation of reactive oxygen/free radicals and endogenous antioxidant defense mechanisms. However, if this balance is disturbed, it can cause oxidative stress, which in turn can result in injury to all cellular components (such as DNA, proteins, and membrane lipids) and cell death. Antioxidants based on natural compounds play a preventive role in protecting against the formation of free radicals. Antioxidant compounds can inhibit and prevent oxidation processes that can produce free radicals and break chains that can damage cells and tissues; hence, these compounds can be used as a therapeutic option for inflammation.^[24]

In general, antiinflammatory properties can suppress the NF-κB pathway and the production of iNOS and proinflammatory cytokines, such as interleukin-1β

and tumor necrosis factor alpha.^[25] Inflammation is prevented by the binding of LPS to toll-like receptor 4 (TLR4). Simultaneously, LPS and TLR4 are catalyzed by lipopolysaccharide-binding protein (LBP) and cluster of differentiation 14 (CD14). After LPS is released, it LPS binds to LBP to form LPS-LBP and targets CD14 on macrophages. LBP catalyzes LPS to CD14 so that LPS combines with CD14, which transfers LPS to TLR4 to then activate the TLR pathway and be directed at inflammatory factors. LPS that binds to TLRs can be neutralized by adenosine monophosphate and can inhibit the release of inflammatory factors.^[24] Another antiinflammatory mechanism is the inhibition and regulation of inflammatory-related signaling pathways. Signal transduction is dependent on a toll-interleukin-1 receptor (TIR) domain-containing adaptor protein (TRIF/TIR domain-containing adaptor-inducing interferon- β) associated with endocytosis of activated TLR4. LPS binding to TLRs activated by TLR4 can also activate the NF- κ B and mitogen-activated protein kinase pathways, where inhibition of TLR4 endocytosis can be performed to suppress inflammatory pathways.^[26]

In a previous study, *C. nutans* extracts showed positive results on a fibroblast migration assay compared to *Strobilanthes crispus* extracts. Chloroform extract of *C. nutans* has also been demonstrated to accelerate healing, including in inflammatory processes, compared with acetone and ethanol extracts. However, *C. nutans* extracts showed less antimicrobial activity against *Staphylococcus aureus*.^[27] In addition, previous studies have shown that chloroform extract of *C. nutans* is the most active at inhibiting NO production compared with hexane and ethanol extracts.^[19] In addition, chloroform extracts of *C. nutans* and aloe vera can upregulate the synthesis of basic fibroblast growth factor (bFGF) in wound healing.^[28] Another previous study showed that *C. nutans* leaf could increase fibroblast count in mucosal burn and tongue wounds *in vivo*.^[29] This proves that the chloroform extract of *C. nutans* leaves is more effective in reducing or inhibiting inflammation.

During inflammation, macrophages and fibroblasts are the main source of NO synthesis.^[30] NO has a small size and lacks charge, so the penetration of NO molecules through cell membranes and subcellular organelles is high. Moreover, NO can be transferred by S-nitrosothiols through cells with protein binding. Therefore, when inflammation occurs, NO is not only in the cells where it is produced but also in the surrounding cells. Among the immune cells that are involved in inflammation, macrophages synthesize the most NO.^[31] Several other factors influence the effect of the chloroform extract

of *C. nutans* leaves on inflamed fibroblasts, such as differences in the metabolite compounds contained in the extract, the amount of extract concentration used, and the extraction method.^[32,33]

The limitation of this study was the use of a limited concentration range. If concentrations above 200 μ g/mL and up to 1000 μ g/mL were used, we could have determined definitively whether 200 μ g/mL is indeed the minimum concentration needed to reduce NO production in fibroblasts.

CONCLUSION

Based on the study results, the chloroform extract of *C. nutans* leaves at a concentration of 200 μ g/mL has the effect of reducing NO synthesis in inflamed fibroblasts. This extract has the potential to be an antiinflammatory agent.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Varela ML, Mogildea M, Moreno I, Lopes A. Acute inflammation and metabolism. *Inflammation* 2018;41:1115-27.
- Germolec DR, Shipkowski KA, Frawley RP, Evans E. Markers of inflammation. *Methods Mol Biol* 2018;1803:57-79.
- Kunnumakkara AB, Sailo BL, Banik K, Harsha C, Prasad S, Gupta SC, et al. Chronic diseases, inflammation, and spices: How are they linked? *J Transl Med* 2018;16:1-25.
- Strandberg TE, Nieminen T. Future perspectives on the role of frailty in cardiovascular diseases. *Adv Exp Med Biol* 2020;1216:149-52.
- Wang R, Ghahary A, Shen YJ, Scott PG, Tredget EE. Human dermal fibroblasts produce nitric oxide and express both constitutive and inducible nitric oxide synthase isoforms. *J Invest Dermatol* 1996;106:419-27.
- Salas A, Cabrera JJ, Jiménez-Leiva A, Mesa S, Bedmar EJ, Richardson DJ, et al. Bacterial nitric oxide metabolism: Recent insights in rhizobia. *Adv Microb Physiol* 2021;78:259-315.
- Cyr AR, Huckaby LV, Shiva SS, Zuckerbraun BS. Nitric oxide and endothelial dysfunction. *Crit Care Clin* 2020;36:307-21.
- Mo X, Chen J, Wang X, Pan Z, Ke Y, Zhou Z, et al. Krüppel-like factor 4 regulates the expression of inducible nitric oxide synthase induced by TNF- α in human fibroblast-like synovial cells. *Mol Cell Biochem* 2018;438:77-84.
- Liy PM, Puzi NNA, Jose S, Vidyadaran S. Nitric oxide modulation in neuroinflammation and the role of mesenchymal stem cells. *Exp Biol Med* 2021;246:2399-406.
- da Silva GM, da Silva MC, Nascimento DVG, Lima Silva EM, Gouvêa FFF, de França Lopes LG, et al. Nitric oxide as a central molecule in hypertension: Focus on the vasorelaxant activity of new nitric oxide donors. *Biology (Basel)* 2021;10:1041.
- Kwesiga MP, Cook E, Hannon J, Wayward S, Gwaltney C, Rao S, et al. Investigative study on nitric oxide production in human dermal fibroblast cells under normal and high glucose conditions. *Med Sci* 2018;6:99.

12. Ardiana T, Rizkia A, Kusuma P, Firdausy MD, Lqwr G, Urxs J, et al. Efektivitas daun jambu biji terhadap jumlah sel fibroblas pasca luka pencabutan gigi pada marmut (*Cavia cobaya*). ODONTO Dent J 2015;2:64-70.
13. Marsh LJ, Kemble S, Reis Nisa P, Singh R, Croft AP. Fibroblast pathology in inflammatory joint disease. Immunol Rev 2021;302:163-83.
14. Kao MH, Wu JS, Cheung WM, Chen JJ, Sun GY, Ong WY, et al. *Clinacanthus nutans* mitigates neuronal death and reduces ischemic brain injury: Role of NF- κ B-driven IL-1 β transcription. Neuromolecular Med 2021;23:199-210.
15. Widjaja S, Rusdiana, Ichwan M. Enhanced cytotoxic effects of *Clinacanthus nutans* and doxorubicin in combination toward breast cancer cell lines. J Adv Pharm Technol Res 2021;12:152-6.
16. Zulkipli IN, Rajabalaya R, Idris A, Sulaiman NA, David SR. *Clinacanthus nutans*: A review on ethnomedicinal uses, chemical constituents and pharmacological properties. Pharm Biol 2017;55:1093-113.
17. Roeslan MO, Reskia SH, Firman FR, Dewa HS, Komariah K, Bustami DA. Effect of *Clinacanthus nutans* leaf extract on oral mucosal burns and tongue wounds: An in-vivo study. Sci Dent J 2022;5:101-9.
18. Goshi E, Zhou G, He Q. Nitric oxide detection methods in vitro and in vivo. Med Gas Res 2019;9:192-207.
19. Roeslan MO, Ayudhya TDN, Yingyongnarongkul B, Koontongkaew S. Anti-biofilm, nitric oxide inhibition and wound healing potential of purpurin-18 phytol ester isolated from *Clinacanthus nutans* leaves. Biomed Pharmacother = Biomed Pharmacotherap 2019;113:108724.
20. Al-Khayri JM, Sahana GR, Nagella P, Joseph BV, Alessa FM, Al-Mssallem MQ. Flavonoids as potential anti-inflammatory molecules: A review. Molecules 2022;27:2901.
21. Islam MI, Shanta MA, Mondal M, Hoque N, Majumder S, Ahmed T, et al. Protective effect of chloroform extract of *Stereospermum chelonoides* bark against amyloid beta42 induced cell death in SH-SY5Y cells and against inflammation in Swiss albino mice. J Basic Clin Physiol Pharmacol 2018;29:621-30.
22. Arulselvan P, Fard MT, Tan WS, Gothai S, Fakurazi S, Norhaizan ME, et al. Role of antioxidants and natural products in inflammation. Oxid Med Cell Longevity 2016;2016:5276130.
23. Alam A, Ferdosh S, Ghafoor K, Hakim A, Juraimi AS, Khatib A, et al. *Clinacanthus nutans*: A review of the medicinal uses, pharmacology and phytochemistry. Asian Paci J Trop Med 2016;9:402-9.
24. Ikrima K, Amalia R, Levita J. Peran spesies oksigen reaktif pada inflamasi serta antioksidan alami sebagai fitoterapi. Farmaka 2020;17:198-211.
25. Ong WY, Herr DR, Sun GY, Lin TN. Anti-inflammatory effects of phytochemical components of *Clinacanthus nutans*. Molecules 2022;27:3607.
26. Luo Y, Song Y. Mechanism of antimicrobial peptides: Antimicrobial, anti-inflammatory and antibiofilm activities. Int J Mol Sci 2021;22:11401.
27. Ban WK, Fong IL, Khong HY, Phung JHY. Wound healing, antimicrobial and antioxidant properties of *Clinacanthus nutans* (burm.f) Lindau and *Strobilanthes crispus* (l) Blume extracts. Molecules 2022;27:1722.
28. Anggraeni R, Roeslan MO. Effects of *Clinacanthus nutans* and *Aloe vera* extracts on bFGF synthesis in fibroblasts. J Indones Dent Assoc 2022;5:21-6.
29. Roeslan MO, Reskia SH, Firman FR, Dewa HS, Komariah K, Bustami DA. Effect of *Clinacanthus nutans* leaf extract on oral mucosal burns and tongue wounds: An In-vivo study. Sci Dent J 2021;5:101-9.
30. Schäffer MR, Tantry U, Barbul A. Wound fluid inhibits wound fibroblast nitric oxide synthesis. J Surg Res 2004;122:43-8.
31. Holotiuik V, Kryzhanivska A, Churpiy I, Tataryn B, Ivasiutyn D. Role of nitric oxide in pathogenesis of tumor growth and its possible application in cancer treatment. Exp Oncol 2019;41:210-15.
32. Pn P, Ma H, Studi P, Kedokteran S, Dokter DP. Efek ekstrak metanol daun sirsak (*Ammon muricata*) dalam menghambat pertumbuhan bakteri *Salmonella typhi* secara in vitro. E-J Medika Udayana 2021;10:2021.
33. Mangkulion S, Roeslan MO, Monthanpisut P. The potential of cogon grass (*Imperata cylindrica*) ethanol extract in inhibiting nitric oxide secretion in fibroblast Sci Dent J 2023;7:11-4.

Effect of Clinacanthus nutans Leaf Chloroform Extract on Reducing Nitric Oxide in Fibroblasts

ORIGINALITY REPORT

21%

SIMILARITY INDEX

15%

INTERNET SOURCES

22%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1

keralasurgj.com

Internet Source

2%

2

A La Teng Zhu La, K.M. Pierce, W.H. Liu, S.T. Gao, D.P. Bu, L. Ma. "Supplementation with Schizochytrium sp. enhances growth performance and antioxidant capability of dairy calves before weaning", Animal Feed Science and Technology, 2020

Publication

2%

3

v3r.esp.org

Internet Source

1%

4

Ying Luo, Yuzhu Song. "Mechanism of Antimicrobial Peptides: Antimicrobial, Anti-Inflammatory and Antibiofilm Activities", International Journal of Molecular Sciences, 2021

Publication

1%

5

academic.oup.com

Internet Source

1%

6

pubmed.ncbi.nlm.nih.gov

Internet Source

1%

7

Submitted to Kaohsiung Medical University

Student Paper

1%

8

www.degruyter.com

Internet Source

1%

9

Submitted to Indian Institute of Technology, Kharagpure

Student Paper

1%

10 Md. Imamul Islam, Meena Afroze Shanta, Milon Mondal, Nazia Hoque, Senjuti Majumder, Taksim Ahmed, Md. Sohel Rana. "Protective effect of chloroform extract of Stereospermum chelonoides bark against amyloid beta42 induced cell death in SH-SY5Y cells and against inflammation in Swiss albino mice", Journal of Basic and Clinical Physiology and Pharmacology, 2018
Publication

11 assets.researchsquare.com 1 %
Internet Source

12 Saruda Thongyim, Thomas A. Wright, Pachara Sattayawat, Thida Kaewkod et al. "Clinacanthus nutans extract lowers periodontal inflammation under high-glucose conditions via inhibiting NF-κB signaling pathway", Frontiers in Pharmacology, 2024
Publication

13 www.jidmr.com 1 %
Internet Source

14 Yessy Ariesanti, Putri Graesya Melani Octavianus, Annisa Tri Handayani, Basril Abbas. "Characterization of Polyvinyl Alcohol–Collagen–Hydroxyapatite Composite Membrane from Lates calcarifer Scales for Guided Tissue and Bone Regeneration", European Journal of Dentistry, 2023
Publication

15 Areta Vania Bhanuwati, Alfred Pakpahan. "Cytotoxic test of different solvents of soursop (Annona muricata) leaf extract against HSC-3 cell line", Dental Journal (Majalah Kedokteran Gigi), 2022
Publication

16 Yoshua Sander, Yessy Ariesanti, M. Orliando Roeslan. "Effect of Moringa oleifera Leaves on

Human Blood Coagulation Process", 2021
IEEE International Conference on Health,
Instrumentation & Measurement, and Natural
Sciences (InHeNce), 2021

Publication

17 Mozaniel Santana de Oliveira, Leo M.L. Nollet.
"Bioactive Compounds - Identification and
Characterization of their Food and
Pharmacological Potential", CRC Press, 2025

Publication

18 encyclopedia.pub 1 %

Internet Source

19 repozitorij.mef.unizg.hr 1 %

Internet Source

20 Sureeporn Suriyaprom, Pornpimon
Ngamsaard, Varachaya Intachaisri, Nitsanat
Cheepchirasuk et al. "Inhibition of Oral
Pathogenic Bacteria, Suppression of Bacterial
Adhesion and Invasion on Human Squamous
Carcinoma Cell Line (HSC-4 Cells), and
Antioxidant Activity of Plant Extracts from
Acanthaceae Family", Plants, 2024

Publication

21 Submitted to Universiti Sains Malaysia 1 %

Student Paper

22 Yugal Kishore Mohanta, Bishwambhar Mishra,
Tamanna Bhuyan. "Microbial Nanotechnology
for Sustainable Future - Industrial and
Environmental Perspectives", CRC Press, 2025

Publication

23 jcd.org.in 1 %

Internet Source

Exclude quotes On

Exclude matches < 15 words

Exclude bibliography On



orliando roeslan <orliandoichol@gmail.com>

[SDJ] Manuscript for revision: SDJ_9_23 - OVERDUE

Scientific Dental Journal <sdj@trisakti.ac.id>
Kepada: orliando roeslan <orliandoichol@gmail.com>
Cc: ihsan.rizal@trisakti.ac.id

1 Agustus 2023 pukul 09.01

Dear Dr. M. Orliando Roeslan,

With reference to your manuscript SDJ_9_23 entitled "Effect of Clinacanthus nutans leaf chloroform extract on reducing Nitric oxide on Fibroblast". The manuscript would be reconsidered after requisite modifications as per the comments and instructions provided by the journal.

If you wish to continue with the publication process, kindly make the changes according to the comments and upload the revised manuscript along with clarifications for all the comments clearly indicating the areas where the changes have been made.

The journal allows not more than 24 hours for the revision of the manuscript.

We thank you for submitting your valuable research work to Scientific Dental Journal.

Sincerely yours,
Muhammad Ihsan Rizal
Editor in-Chief
Scientific Dental Journal
ihsan.rizal@trisakti.ac.id



UNIVERSITAS TRISAKTI
"Is a one stop learning for sustainable development"
Kampus A, [Jl. Kyai Tapa No.1, Grogol](#)
[Jakarta Barat 11440 - INDONESIA](#)
www.trisakti.ac.id
(t) +62-21.566 3232, (f) +62-21.567 3001



SDJ_9_23.docx
64K



orliando roeslan <orliandoichol@gmail.com>

[SDJ] Manuscript for revision: SDJ_9_23

Scientific Dental Journal <sdj@trisakti.ac.id>
Kepada: orliando roeslan <orliandoichol@gmail.com>
Cc: ihsan.rizal@trisakti.ac.id

1 Desember 2023 pukul 14.21

Dear Dr. M. Orliando Roeslan,

With reference to your manuscript SDJ_9_23 entitled "Effect of Clinacanthus nutans leaf chloroform extract on reducing Nitric oxide on Fibroblast". The manuscript would be reconsidered after requisite modifications as per the comments and instructions provided by the journal.

If you wish to continue with the publication process, kindly make the changes according to the comments and upload the revised manuscript along with clarifications for all the comments clearly indicating the areas where the changes have been made.

The journal allows not more than 24 hours for the revision of the manuscript.

We thank you for submitting your valuable research work to Scientific Dental Journal.

Sincerely yours,
Muhammad Ihsan Rizal
Editor in-Chief
Scientific Dental Journal
ihsan.rizal@trisakti.ac.id



UNIVERSITAS TRISAKTI
"Is a one stop learning for sustainable development"
Kampus A, [Jl. Kyai Tapa No.1, Grogol](#)
[Jakarta Barat 11440 - INDONESIA](#)
www.trisakti.ac.id
(t) +62-21.566 3232, (f) +62-21.567 3001



SDJ_9_23.docx
71K