



PADJADJARAN JOURNAL OF DENTISTRY

UNIVERSITAS PADJADJARAN

P-ISSN : 19790201 <> E-ISSN : 25496212 Subject Area : Health, Science

0.75
Impact Factor

1226
Google Citations

Sinta 2
Current Accreditation

[Google Scholar](#) [Garuda](#) [Website](#) [Editor URL](#)

History Accreditation

2018 2019 2020 2021 2022 2023 2024 2025 2026

[Garuda](#) [Google Scholar](#)

[Antibacterial potential of celery leaf extract toothpaste on the growth of Streptococcus mutans ATCC 25175](#)

Faculty of Dentistry Universitas Padjadjaran [Padjadjaran Journal of Dentistry Vol 35, No 1 \(2023\): March 2023 1-4](#)

2023 [DOI: 10.24198/pjd.vol35no1.30624](#) [Accred : Sinta 2](#)

[Effect of varnish coating agent application on surface hardness of glass ionomer cement after immersion in carbonated drinks](#)

Faculty of Dentistry Universitas Padjadjaran [Padjadjaran Journal of Dentistry Vol 35, No 1 \(2023\): March 2023 47-52](#)

2023 [DOI: 10.24198/pjd.vol35no1.33139](#) [Accred : Sinta 2](#)

[The effectiveness of using local exhaust ventilation, HEPA filter, and dental aerosol suction on indoor air quality](#)

Faculty of Dentistry Universitas Padjadjaran [Padjadjaran Journal of Dentistry Vol 35, No 1 \(2023\): March 2023 23-28](#)

2023 [DOI: 10.24198/pjd.vol35no1.33566](#) [Accred : Sinta 2](#)

[Relationship between mother's oral health literacy level with oral hygiene behavior and self-reported oral health status in sociodemographic scope](#)

Faculty of Dentistry Universitas Padjadjaran [Padjadjaran Journal of Dentistry Vol 35, No 1 \(2023\): March 2023 40-46](#)


2023 [DOI: 10.24198/pjd.vol35no1.35411](#) [Accred : Sinta 2](#)

[Effectiveness of mixed tea plant extract gel with chitosan on fibroblast cells after tooth extraction of Wistar rats](#)

Faculty of Dentistry Universitas Padjadjaran [Padjadjaran Journal of Dentistry Vol 35, No 1 \(2023\): March 2023 10-15](#)

2023 [DOI: 10.24198/pjd.vol35no1.36563](#) [Accred : Sinta 2](#)

Citation Per Year By Google Scholar



Journal By Google Scholar

	All	Since 2018
Citation	1226	1135
h-index	13	12
i10-index	24	22

[Prevention and control of COVID-19 infection at the community health center during the new normal era](#)

Faculty of Dentistry Universitas Padjadjaran [Padjadjaran Journal of Dentistry Vol 35, No 1 \(2023\): March 2023 70-79](#)

[2023](#) [DOI: 10.24198/pjd.vol35no1.39563](#) [Accred : Sinta 2](#)

[Relationship between oral health service quality and patient satisfaction at community health centers in Kupang city during COVID-19 pandemic](#)

Faculty of Dentistry Universitas Padjadjaran [Padjadjaran Journal of Dentistry Vol 35, No 1 \(2023\): March 2023 86-90](#)

[2023](#) [DOI: 10.24198/pjd.vol35no1.40111](#) [Accred : Sinta 2](#)

[The effect of xerostomia on the quality of life of controlled and uncontrolled type 2 diabetes mellitus patients](#)

Faculty of Dentistry Universitas Padjadjaran [Padjadjaran Journal of Dentistry Vol 35, No 1 \(2023\): March 2023 5-9](#)

[2023](#) [DOI: 10.24198/pjd.vol35no1.41996](#) [Accred : Sinta 2](#)

[Aloe vera extract wound healing sheet and free-range chicken egg albumin \(gallus domesticus\) accelerate angiogenesis on gingival incision wound in rats](#)

Faculty of Dentistry Universitas Padjadjaran [Padjadjaran Journal of Dentistry Vol 35, No 1 \(2023\): March 2023 16-22](#)

[2023](#) [DOI: 10.24198/pjd.vol35no1.42775](#) [Accred : Sinta 2](#)

[The association of cariogenic foods consumption frequency on the occurrence of dental caries in children with autism spectrum disorders](#)

Faculty of Dentistry Universitas Padjadjaran [Padjadjaran Journal of Dentistry Vol 35, No 1 \(2023\): March 2023 34-39](#)

[2023](#) [DOI: 10.24198/pjd.vol35no1.42892](#) [Accred : Sinta 2](#)

[View more ...](#)

Get More with
SINTA Insight

[Go to Insight](#)

Citation Per Year By Google Scholar

Journal By Google Scholar

	All	Since 2018
Citation	1226	1135
h-index	13	12
i10-index	24	22

PADJADJARAN JOURNAL OF DENTISTRY

p-ISSN : 1979-0201 e-ISSN : 2549-6212

In affiliation with
Indonesia Dental Association



Published by
Universitas Padjadjaran
Faculty of Dentistry

e ISSN 2549-6212
p ISSN 1979-0201

PADJADJARAN JOURNAL OF DENTISTRY

<http://jurnal.unpad.ac.id/pjd>

ISSUE	TITLE	
Vol 35, No 1 (2023): March 2023	Prevention and control of COVID-19 infection at the community health center during the new normal era	ABSTRACT PDF
	<i>Gilang Ferara Daktari, Wahyu Hidayat, Netty Suryanti</i>	
Vol 34, No 2 (2022): July	Indonesian dentists knowledge, attitudes and practices in COVID-19 pandemic	ABSTRACT PDF
	<i>Anne Agustina Suwargiani, Erry Mochamad Arief, Asty Samiaty Setiawan, Fidya Meditia Putri, Ardena Maulidia Hamdani, Azlina Nuur Sanjaya, Gilang Yubiliani, Sri Susilawati, Dudi Aripin, Nina Djuastina, Sunardhi Widyaputra</i>	
Vol 35, No 1 (2023): March 2023	Relationship between mother's oral health literacy level with oral hygiene behavior and self-reported oral health status in sociodemographic scope	ABSTRACT PDF
	<i>Tiarani Maudi, Netty Suryanti, Asty Samiaty Setiawan</i>	
Vol 35, No 1 (2023): March 2023	Differences in the visual perception of the upper dental midline deviation between orthodontically completely treated and untreated patients	ABSTRACT PDF
	<i>Avi Laviana, Imamanda Dwipura Yakin Hasbullah, Elih Sayuti, Endah Mardiaty, Indah Yuri Noviaranny</i>	
Vol 35, No 1 (2023): March 2023	The effectiveness of using local exhaust ventilation, HEPA filter, and dental aerosol suction on indoor air quality	ABSTRACT PDF
	<i>Retno Hayati Alchusnah, Dewi Sarastuti, Lia Fetti Hidayati, Fatimah Septiantari, Belladina Yusi Lasara</i>	
Vol 35, No 1 (2023): March 2023	Aloe vera extract wound healing sheet and free-range chicken egg albumin (<i>gallus domesticus</i>) accelerate angiogenesis on gingival incision wound in rats	ABSTRACT PDF
	<i>Myrna Evana Amanda Putri, Christiana Cahyani Prihastuti, Mutia Rochmawati, Alya Ghina Rosyada, Wizni A'dila A'ziza</i>	
Vol 35, No 1 (2023): March 2023	The effect of ZrCl ₄ , Al(NO ₃) ₃ and Na ₂ SiO ₃ precursors concentration on particle size as dental composites filler through spray pyrolysis method	ABSTRACT PDF
	<i>Elin Karlina, Khairina Idznihaq, Renny Febrida, Niekla Survia Andiesta</i>	

Vol 35, No 1 (2023): March 2023 The resistance of the intracanal retention thickness with flowable short fiber reinforced composite (SFRC) materials towards fracture toughness [ABSTRACT](#) [PDF](#)

Isabella Audy Tania, Wiena Widyastuti

Vol 35, No 1 (2023): March 2023 Effect of varnish coating agent application on surface hardness of glass ionomer cement after immersion in carbonated drinks [ABSTRACT](#) [PDF](#)

Natasya Arsanti, Aryo Megantoro, Ratna Meidyawati

Vol 35, No 1 (2023): March 2023 The association of cariogenic foods consumption frequency on the occurrence of dental caries in children with autism spectrum disorders [ABSTRACT](#) [PDF](#)

Chita Sintya Dewi, Risti Saptarini Primarti, Inne Suherma Sasmita

Vol 35, No 1 (2023): March 2023 Preparation of micro-calcite powder using fine-bubble assisted carbonation as toothpaste material [ABSTRACT](#) [PDF](#)

Renny Febrida, Dinda Jihan Nabillah Salma, Fauziah Azzahra, Elin Karlina, Niekla Survia Andiesta, Yanwar Faza

Vol 35, No 1 (2023): March 2023 Effectiveness of mixed tea plant extract gel with chitosan on fibroblast cells after tooth extraction of Wistar rats [ABSTRACT](#) [PDF](#)

Nyoman Ayu Anggayanti, Putu Lestari Sudirman, Ni Nyoman Rian Permata Sari, I Gusti Ayu Putu Diah Suryani

Vol 35, No 1 (2023): March 2023 Mixture of Indonesian White Portland Cements, Bi₂O₃, and Light Cured Methacrylate-based resin as a potential candidate for pulp capping material [ABSTRACT](#) [PDF](#)

Denny Nurdin, Beatrix Lamria Simanjuntak, Rahmi Alma Farah Adang, Arief Cahyanto

Vol 35, No 1 (2023): March 2023 The effect of xerostomia on the quality of life of controlled and uncontrolled type 2 diabetes mellitus patients [ABSTRACT](#) [PDF](#)

Sectio Aprista, Surya Nelis, Avina Anin Nasia, Yoghi Bagus Prabowo

Vol 35, No 1 (2023): March 2023 Relationship between oral health service quality and patient satisfaction at community health centers in Kupang city during COVID-19 pandemic [ABSTRACT](#) [PDF](#)

Friska Deli Simamora, Jacob Matheos Ratu, Anderias Uumbu Roga, Pius Weraman, Hotlif Arkilaus Nope

Chief Editor

- [Prof. Sunardhi Widyaputra, drg, MS, PhD](#), Scopus ID= 6602995626; Department of Oral Biology, Faculty of Dentistry Universitas Padjadjaran, Indonesia

Handling Editor

- [Prof. Dr. Nina Djustiana, drg, MKes](#), Scopus ID= 57189578833; Department of Dental Materials, Science, and Technology, Faculty of Dentistry Universitas Padjadjaran, Indonesia
- [Dr Anne Agustina Suwargiani, drg, MKM](#), Scopus ID= 57203020093; Department Ilmu Kesehatan Gigi Masyarakat, Fakultas Kedokteran Gigi Universitas Padjadjaran, Indonesia
- [Zulia Hasratiningsih, drg, MDSc](#), Scopus ID= 37045476800; Departemen Ilmu Teknologi dan Material Kedokteran Gigi, Fakultas Kedokteran Gigi Universitas Padjadjaran, Indonesia, Indonesia
- [Dr. Netty Suryanti, drg, MARS.](#), Scopus ID= 57210117266; Department of Community Dental Health, Faculty of Dentistry, Padjadjaran University, Indonesia

Editorial Board

- [Dr. Ali Mohammed](#), Scopus ID= 57652411300; Faculty of Medicine, Dentistry and Health Sciences, The University of Melbourne, Australia, Australia
- [Cortino Sukotjo, DDS, PhD, MMsC, FACP](#), Scopus ID= 6508194317; Department of Restorative Dentistry and Advanced Prosthodontics, College of Dentistry, University of Illinois, United States
- [Prof. Mariko Naito](#), Scopus ID= 57204325415; Department of Oral Epidemiology, Graduate School of Biomedical and Health Sciences Hiroshima University, Japan
- [Drg Rizky Indrameikha Sugianto, MPH, PhD](#), Scopus ID= 57201006215, Postdoctoral Researcher, Hannover Medical School,, Germany
- [drg. Niekla Survia Andiesta, BDS, MDS](#), Scopus ID= 57202599268; Division of Children and Community Oral Health, School of Dentistry, International Medical University, Malaysia
- [Prof. Mohamed Ebrahim Parker](#), Scopus ID= 7403672513; Department of Diagnostic Sciences Radiology, Maxillofacial and Forensic Sciences, University of The Western Cape, South Africa
- [Prof. Kotaro Tanimoto](#), Scopus ID= 57191990083; Graduate School of Biomedical and Health Sciences Dentistry & Oral Health Sciences, Hiroshima University, Japan
- [Prof. Dr. Mohammad Tariqur Rahman](#), Scopus ID = 55457946600, Dean Office Faculty of Dentistry, universiti malaya, Malaysia

- [Prof. Kotaro Tanimoto](#), Scopus ID= 57191990083; Graduate School of Biomedical and Health Sciences Dentistry & Oral Health Sciences, Hiroshima University, Japan
- [Prof. Dr. Mohammad Tariqur Rahman](#), Scopus ID = 55457946600, Dean Office Faculty of Dentistry, universiti malaya, Malaysia
- [Prof. Yoshizo Matsuka](#), Scopus ID= 7003862097; Department of Stomatognathic Function and Occlusal Reconstruction, Graduate School of Biomedical Sciences Tokushima University, Japan
- [Prof. Dr. Zamros Yuzadi Yusof](#), Scopus ID = 22939737100, Department of Community Oral Health & Clinical Prevention Faculty of Dentistry, Universiti Malaya, Malaysia, Malaysia
- [Associate Professor Dr Akram Hassan](#), Scopus ID= 55832848700; Department of Periodontics, School of Dental Sciences Universiti Sains Malaysia, Malaysia
- [Dr. Solachuddin Jauhari Arief, DDS., PhD. Ichwan](#), Scopus ID= 6504103591; PAPRSB Institute of Health Sciences, International Islamic University, Brunei Darussalam
- [Prof. Dr. Arlette Suzy Puspa Pertiwi Setiawan, drg., Sp KGA., M.Si.](#), Scopus ID= 56044838600; Department of Pediatric, Faculty of Dentistry Padjadjaran University,, Indonesia
- [Dr Arief Cahyanto, MT., Ph.D](#), Scopus ID= 55532851800; Department of Dentistry Material Science and Technology, Faculty of Dentistry, Padjadjaran University, Indonesia, Indonesia
- [Dr Elizabeth Fitriana Sari](#), Scopus ID= 57219228212 Departemen Ilmu Penyakit Mukut, Fakultas Kedokteran Gigi Universitas Padjadjaran,, Indonesia
- [Dr. Rasmi Rikmasari, drg., Sp.Pros., Subsp. OGST \(K\)](#), Scopus ID= 57191990083; Department of Prosthodontics, Faculty of Dentistry, Padjadjaran University, Indonesia, Indonesia
- [Dr. Sri Tjahajawati, drg., M.Kes.AIFM.](#), Scopus ID= 57197722254; Department of Oral Biology, Faculty of Dentistry Universitas Padjadjaran, Indonesia
- [Dr. Hendra Dian Adhita Dharsono, drg, Sp.KG., Subsp.KE\[K\]](#), Scopus ID= 57204917449; Department of Dental Conservation, Faculty of Dentistry, Padjadjaran University, Indonesia
- [Fahmi Oscandar, drg., M.Kes., SpOF., SubSp OFK \(K\), Ph.D. \(Cr.Img\).](#), Scopus ID= 57199734614; Departemen Radiologi Kedokteran Gigi, Fakultas Kedokteran Gigi, Universitas Padjadjaran, Indonesia
- [Dr. Endang Sjamsudin, drg, Sp.BMMF, Subsp.TMF-TMJ\[K\]](#), Scopus ID= 57192257503; Department of Oral Surgery, Faculty of Dentistry, Padjadjaran University, Indonesia
- [R. Tantry Maulina, drg., M.Kes., Ph.D.](#), Scopus ID= 57191972242; Department of Oral Surgery, Faculty of Dentistry, Padjadjaran University, Indonesia, Indonesia
- [Dr. Avi Laviana, drg., Sp.Ort., Subsp.DDTK\(K\).](#), Scopus ID= 57211331865; Doctor of Orthodontics Department of Orthodontics Faculty of Dentistry, Padjadjaran University, Indonesia
- [Prof. Dr. Inna Sufiawati, drg., Sp.PM., Subsp.Inff\[K\]](#), Scopus ID= 56081844700; Departemen Ilmu Penyakit Mukut, Fakultas Kedokteran Gigi Universitas Padjadjaran, Indonesia, Indonesia
- [Amaliya, drg., M.Sc., Ph.D.](#), Scopus ID= 56584444300; Department of Periodontics, Faculty of Dentistry, Padjadjaran University, Indonesia
- [Lusi Epsilawati, drg., Sp.RKG., Subsp.Rad.P\(K\), M.Kes.](#), Scopus ID= 55523245700; Department of Radiology, Faculty of Dentistry, Padjadjaran University, Indonesia

Reviewer

- [Prof. Jie Yang](#), Scopus ID= 57051770100; Department of Oral and Maxillofacial Pathology, Medicine, and Surgery, Kornberg School of Dentistry Temple University, United States
- [Prof. Erry Mochamad Arief, BDS, MS](#), Scopus ID= 6508283549; Department of Periodontics, School of Dental Sciences, Universiti Sains Malaysia, Malaysia
- [DR. Christopher John Nile](#), Scopus ID= 6507923698; Institute of Infection Immunity & Inflammation, School of Medicine, Dentistry and Nursing University of Glasgow, United Kingdom
- [Epita Sarah Pane, DDS, GradDipClinDent, MDSc, PhD](#), Scopus ID= 6603159739; Department of Endodontics, Melbourne Dental School, University of Melbourne, Australia
- [Associate Prof. Dr. Noor Azlin Yahya](#), Scopus ID=3 6611095600 Department of Restorative Dentistry, Faculty of Dentistry, Universiti Malaya, malaysia, Malaysia
- [Mr. Rozaini Abdul Rahim](#), Scopus ID= 57194208647 Fakulti Pergigian, Universiti Malaya, Malaysia
- [Prof. Dr Jennifer Geraldine Doss](#), Scopus ID= 14631803900 Department of Community Oral Health & Clinical Prevention Faculty of Dentistry,Universiti Malaya, Malaysia, Malaysia
- [Sazan Manaf Azeez](#), Conservative department assistant lecturer at college of dentistry, Hawler Medical University, Kurdistan Region,Iraq, Iraq
- [Dr. Madhuri Pattamatta,Phd](#), Quality and Safety of Oral Health Care, Radboud University, Nijmegen, Netherlands, Netherlands
- [Profesor Dr Fathilah Abdul Razak](#), Scopus ID = 16642630700 Department of Oral and Craniofacial Sciences Faculty of Dentistry,Universiti Malaya, Malaysia, Malaysia
- [Prof. Dr. Mieke Hemiawati Satari Hemiawati Satari, drg., M.Kes.](#), Scopus ID= 56884598800; Department of Oral Biology, Faculty of Dentistry, Padjadjaran University, Indonesia
- [Prof. Dr. Unang Supratman, M.Si](#), Scopus ID= 6508000748; Professor of Natural Products Chemistry Department of Chemistry Faculty of Mathematics and Natural Sciences, Padjadjaran University, Indonesia
- [Prof. Dr. Williyanti Suwondo, drg., Sp.Ped., Subsp. AIBK\(K\).](#), Scopus ID= 56045377200; Department of Pediatric Dentistry, Faculty of Dentistry, Padjadjaran University, Indonesia
- [Prof. Dr. Endah Mardiaty, drg., Sp.Ort., Subsp.DDTK\(K\), MS.](#), Scopus ID= 57217914378; Department of Orthodontics, Faculty of Dentistry, Padjadjaran University, Indonesia
- [Prof. Dr. Eriska Riyanti, drg., Sp.KGA., Subsp.AIBK\(K\).](#), Scopus ID= 57195358674; Department of Pediatric Dentistry, Faculty of Dentistry, Padjadjaran University, Indonesia
- [Prof. Dr. Keri Lestari, M.Si., Apt.](#), Scopus ID= 55070933000; Department of Pharmacology and Clinical Pharmacy, Faculty of Pharmacy, Padjadjaran University, Indonesia
- [Prof. Dr. Risdiana, S.Si, M. Eng Silaka](#), Scopus ID= 25633171800; Department of Physics, Faculty of Mathematics and Natural Sciences, Padjadjaran University,, Indonesia
- [Prof. Dr. Harmas Yazid Yusuf, drg., SpBM](#), Scopus ID= 57197229289; Department of Oral and Maxillofacial Surgery Faculty of Dentistry Universitas Padjadjaran, Indonesia

- [Dr. Azhari, drg., Sp.RKG, Subsp.Rad.D\(K\), M.Kes.](#), Scopus ID= 57193963704; Department of Dental Radiology, Faculty of Dentistry, Padjadjaran University, Indonesia
- [Dr. Agus Susanto, drg., Sp.Perio., Subsp. RPID\(K\), M.Kes.](#), Scopus ID=57220399229 ; Department of Periodontics, Faculty of Dentistry, Padjadjaran University, Indonesia
- [Dr. Sri Susilawati, drg., M.Kes.](#), Scopus ID= 57201867946; Department of Public Dental Health, Faculty of Dentistry, Padjadjaran University, Indonesia
- [Dr. Nanan Nur'aeny, drg., Sp.PM., Subsp.Noninf\(K\),](#) Scopus ID= 57222990248; Department Of Oral Medicine , Faculty of Dentistry, Padjadjaran University,, Indonesia
- [Prof. Dr. Dewi Marhaeni Diah Herawati, drg, MSi,](#) Scopus ID= 57063296400, Department of Public Health Sciences, Faculty of Medicine, Padjadjaran University, Indonesia
- [Dr. Setiawan Soetopo, Sp.Rad, Sp.Onk,](#) Scopus ID= 57196436060; Department of Radiation Oncology, Radiotherapy Installation, Hasan Sadikin Hospital
- [Prof. Dr. Inna Sufiawati, drg., Sp.PM., Subsp.Inf\[K\],](#) Scopus ID= 56081844700; Departemen Ilmu Penyakit Mukut, Fakultas Kedokteran Gigi Universitas Padjadjaran, Indonesia, Indonesia
- [Dr. Irmaleny , drg., Sp.KG., Subsp.KR\(K\),](#) Scopus ID= 57291776000; Departemen Konservasi, Fakultas Kedokteran Gigi Universitas Padjadjaran, Indonesia
- [Dr. Achadiyani, dr.,MKes,](#) Scopus ID= 57195942091; Department of Anatomy, Physiology, and Cellular Biology, Faculty of Medicine, Padjadjaran University, Indonesia
- [Dr. Hening Tjaturina Pramesti, dra, MS,](#) Scopus ID= 55258696800; Department of Oral Biology, Faculty of Dentistry, Padjadjaran University, Indonesia
- [Amaliya, drg., M.Sc., Ph.D.,](#) Scopus ID= 56584444300; Department of Periodontics, Faculty of Dentistry, Padjadjaran University, Indonesia
- [Deby Fajar Mardhian, S.Si., MT., Ph.D.,](#) Scopus ID= 57196040281, Department of Dental Materials Science and Technology, Faculty of Dentistry, Padjadjaran University, Indonesia
- [Lusi Epsilawati, drg., Sp.RKG., Subsp.Rad.P\(K\), M.Kes.,](#) Scopus ID= 55523245700; Department of Radiology, Faculty of Dentistry, Padjadjaran University, Indonesia
- [Yuti Malinda, drg., MM., M.Kes.,](#) Scopus ID= 57200731581; Department of Oral Biologi, Fakultas Kedokteran Gigi, Universitas Padjadjaran, Indonesia, Indonesia
- [Diani Prisinda, drg., Sp.KG., Subsp.KE\(K\), MARS.,](#) Scopus ID= 57197812277; Department of Dental Conservation, Faculty of Dentistry, Padjadjaran University, Indonesia
- [Dewi Zakiawati, drg., Sp.PM., MSc.,](#) Scopus ID= 57204158372; Department of Oral Medicine, Faculty of Dentistry, Padjadjaran University, Indonesia
- [Veni Takarini, drg., M.Kes,](#) Scopus ID= 57189580491; Department of Dental Materials Science and Technology, Faculty of Dentistry, Padjadjaran University, Indonesia
- [Felisha Febriane Balafif, drg., M.Kes.,](#) Scopus ID= 57220592980, Department of Oral Biology, Faculty of Dentistry, Padjadjaran University, Indonesia

ORIGINAL ARTICLE

The resistance of the intracanal retention thickness with flowable short fiber reinforced composite (SFRC) materials towards fracture toughness

Isabella Audy Tania^{1*}
Wienna Widyastuti¹

¹Department of Conservative Dentistry,
Faculty of Dentistry, Universitas Trisakti,
Indonesia

*Correspondence:
wiena@trisakti.ac.id

Received: 07 February 2023
Revised: 17 March 2023
Accepted: 31 March 2023
Published: 31 March 2023
DOI: [10.24198/pjd.vol35no1.44892](https://doi.org/10.24198/pjd.vol35no1.44892)

Citation:

Tania IA, Widyastuti W. The resistance of the intracanal retention thickness with flowable short fiber reinforced composite (SFRC) materials towards fracture toughness. *Padj J Dent*, March. 2023;35(1):80-85

ABSTRACT

Introduction: Dental hard tissue problems were relatively high in Indonesia. If these problems were ignored, they would cause pulpal necrosis. The main treatment for this case was Root Canal Treatment (RCT). The selection of post-RCT treatment and suitable materials was important. One of the examples was short fiber reinforced composite (SFRC) which can be an option for intracanal retention treatment. This study aims to analyze the resistance of the intracanal retention thickness with flowable fiber reinforced composite materials toward fracture toughness. **Methods:** This research was using a correlative laboratory experimental method with a post-test group-only design. This study's samples were SFRC material inserted in 27 extracted mandibular first premolars with a length of 21 mm ± 2 mm and root canal walls width of 2 mm. All samples were given intracanal retention at a depth of 2 mm below the orifice and divided into 3 groups: intracanal retention with thickness of (A) 6 mm, (B) 5 mm, and (C) 4 mm. RCT was performed using the crown-down technique. The bonding agent was applied, and then light cured for 20 seconds. SFRC was applied according to the sample group and light cured for 20 seconds per 2 mm, followed by filling with composite resin. The sample was implanted in the dental stone 2 mm below the CEJ. The fracture toughness test was performed using UTM on the occlusal surface with a 0° tilt and crosshead of 1 mm/min. **Results:** The fracture toughness results of group A, B, and C are 735.44, 756.78, and 829 respectively. Statistical tests with one-way ANOVA showed no significant difference with p value = 0.412 (p < 0.05). **Conclusion:** There is no difference in resistance of the intracanal retention thickness with flowable fiber reinforced composite materials towards fracture toughness.

KEYWORDS

post-RCT treatments, intracanal retentions, SFRC, fracture toughness

INTRODUCTION

Oral health is an important thing that everyone should be aware of. However, according to the Basic Health Research (Risikesdas) in 2018, awareness level of oral health in Indonesia still needs to be improved. Nowadays, the main problems are hard and soft tissue diseases. Risikesdas stated that the level of dental hard tissue disease in Indonesian people is still high, namely 45.3% who experience dental caries.¹ Caries can become severe problems, such as pain and discomfort, and also teeth becoming easily broken due to loss of tooth tissue structure that can lead to acute and chronic infections.² Pulp necrosis is a condition of pulp tissue death that is usually asymptomatic. The etiology of pulp tissue death can be mechanical, chemical irritants, or bacterial microorganisms.³ Tooth trauma is also one of the main causes of tooth fracture, which can involve damage to the tooth pulp.⁴

According to the 2020 American Association of Endodontists (AAE), root canal treatment (RCT) is the procedure for removing necrotic tissue from the root canal, followed by shaping and obturating the root canal with a filling material.⁵ Post-RCT teeth will become more brittle and prone to fracture. This is because the water content in the dentine tissue decreases, causing it to become dehydrated, which causes the dentine collagen to shrink. Also, the post-RCT teeth can no longer form the secondary dentine tissue, which makes dentin support for enamel reduced.⁶ As a result of these factors, post-RCT teeth become prone to fracture when they receive functional pressure or mastication.⁷

Fracture toughness is a mechanical property that describes the ability of a material to resist crack propagation and defines the level of tolerance of a material against damage.⁸ The fracture toughness value depends on the physical properties and chemical composition of the components of the restorative material. In previous studies, it was shown that there is a strong correlation between the fracture toughness of the material and the fracture rate of dental restoration materials.⁹ Therefore, materials with high fracture toughness values tend to be able to withstand the risk of micro fractures that may occur during the mastication process. In other words, materials that have higher fracture toughness values tend to have a better ability to resist the formation and spread of fractures.

It is known that fiber reinforced composite (FRC) materials have high fracture toughness values.^{10,11} The properties of FRC are highly dependent on fiber diameter, fiber length, fiber orientation or direction, fiber loading, and also the adhesion or attachment between the fibers and the polymer resin matrix.¹² The development of FRC materials is urgently needed to reduce the failure rate of RCT in dentistry, especially fractures. This study aims to analyze the resistance of the

intracanal retention thickness with flowable fiber reinforced composite materials toward fracture toughness.

METHODS

This research was conducted using a correlative laboratory experimental method with a post-test group-only design carried out at the RSGM-P Faculty of Dentistry, Universitas Trisakti and Industrial Metrology Laboratory, Universitas Trisakti, Jakarta, Indonesia. The research sample used a flowable short fiber-reinforced resin composite (SFRC) (GC EverX Flow) inserted into the mandibular first premolar as intracanal retention.

A sample of 27 extracted mandibular first premolars with single and straight root, intact crown with no caries, without root caries, and completely closed apex teeth were cleaned and then soaked in saline water. Each sample was selected by measuring the working length using a caliper, with a length of $21 \text{ mm} \pm 2 \text{ mm}$ (Figure 1).



Figure 1. Measurement with millimeter blocks.

Next, a digital radiographic photo was taken to measure the root canal walls and to ensure the length of the sample using a digital millimeter block. The length of the teeth was calculated using 2 points, namely the cusp of the premolar and the apex. The thickness of the root canal walls should be 2 mm thick from the CEJ on the outside to the root canal walls on the inside. It was then grouped into 3 groups: samples that had undergone root canal treatment and were given 2 mm intracanal retention below the orifice using SFRC flowable material with a thickness of (A) 6 mm; (B) 5 mm; (C) 4 mm (Figure 2).




Figure 2. The treatment scheme of the sample: Group A. Sample with 6 mm thickness of flowable SFRC intracanal retention; Group B: Sample with 5 mm thickness of flowable SFRC intracanal retention. Group C: Sample with 4 mm thickness of flowable SFRC intracanal retention

RCT was performed using the crown down technique using ProTaper (Dentsply Maillefer) up to size F3. At each file change, 2 ml of 5.25% NaOCl solution was irrigated in the root canals and then dried using three-way syringe and paper points. Then obturation was carried out using the single cone technique using gutta-percha and a sealer was applied using a root canal sealer (Dentsply AH 26). Finally, the obturation material was removed using a large reamer #3 (Dentsply Maillefer) 2 mm below the orifice for all samples.

Bonding material (GC Solare Bond) was applied to the tooth sample and dried, then light cured for 20 seconds. Next, intracanal retention material was applied 2 mm deep under the orifice for all samples and light cured for 20 seconds per 2 mm thickness. Furthermore, the filling was carried out using a composite resin filling material (GC Solare Sculpt Composite). These thicknesses from group A, B, and C were chosen based on previous studies¹³, which found that the fiber length differences between 1–2 mm could change the pattern of the fractures and greatly enhance the fracture resistance of the composite (Figure 3).

The tooth sample was implanted in type II dental stone 2 mm deep below the CEJ so that it resembled the position of the teeth on the alveolar bone. Furthermore, the fracture toughness test was performed using the Universal Testing Machine (Hounsfield: 50kN, England). The pressure was applied using a fracture toughness jig on the occlusal surface with a slope of 0° along the tooth axis apically. Constant pressure was applied at a crosshead speed of 1 mm/min (American Society for Testing Materials Standard) until the tooth failed in the form of a macroscopic fracture in the

intracanal retention of the SFRC material. Fractures of fillings and teeth that do not involve intracanal retention were ignored.



Figure 3. Fracture on intracanal retention with SFRC materials.

Statistical analysis was performed on the fracture toughness data. Saphiro-Wilk normality test was performed, and the p-value > 0.05 was considered as normal data distribution. Then followed by one-way ANOVA test data analysis. if the One Way ANOVA test obtained p-value < 0.05, the analysis was continued with the post hoc test with Tukey's HSD. If the p-value shows > 0.05, the conclusion was that there is no significant difference in the fracture toughness values of the sample groups A, B, and C, and the post hoc test was not carried out.

RESULTS

The statistical results showed in the Shapiro-Wilk normality test (Table 1), data is considered normally distributed if the p-value>0,05. They are then followed by One Way ANOVA data analysis (Table 3) with a p-value = 0.412. This means that there is no significant difference in the fracture toughness values of the sample groups A, B, and C.

Table 1. Shapiro-Wilk test of Normality on fracture toughness data.

Flowable SFRC	Shapiro-Wilk		
	Statistic	df	p-value
6 mm	0.200	9	0.692*
5 mm	0.200	9	0.949*
4 mm	0.200	9	0.356*

The results of the One Way ANOVA analysis (Table 2) showed that the average fracture toughness value in group A with an intracanal retention thickness of 6 mm had the lowest fracture toughness value at 735.44 N (±146.407). Group B, with an intracanal retention of 5 mm, had a fracture toughness value of 756.78 N (±153.791). While the sample in group C with intracanal retention thickness of 4 mm had the highest fracture toughness value at 829.00 N (±159.290). As can be seen in bar diagram (Figure 4). The results of the homogeneity test showed that there was no significant relationship between the sample groups (p > 0.05).

Table 2. One Way ANOVA analysis on fracture toughness data.

	n	Mean	Std. Deviation	Std. Error	Lower Bound	Upper Bound	Min	Max
6 mm	9	735.44	146.407	48.802	622.91	847.98	548	1002
5 mm	9	756.78	153.791	51.264	638.56	874.98	502	972
4 mm	9	829.00	159.290	53.097	706.56	951.44	589	1013
Total	27	773.74	152.788	29.404	713.90	834.18	502	1013




Figure 4. Fracture toughness data of group A, B, and C.

Table 3. One Way ANOVA analysis.

	Sum of Squares	df	Mean square	F	p-value
Between groups	43271.407	2	21635.704	0.921	0.412
Within groups	563679.778	24	23486.657		
Total	606951.185	26			

DISCUSSION

The test data using UTM found that the average fracture toughness value was group C > B > A. Based on the Shapiro-Wilk normality test, group A has a significance value of 0.692, group B has 0.949, and group C has 0.356. So that the significance value (p/Sig.) in this normality test can be concluded that the data is normally distributed. This conclusion is obtained from the significance value of all groups of more than 0.05. Therefore, the assumption of normality of the data is met and data analysis can be continued using the one-way ANOVA test. On the One Way ANOVA test data, it is known that the average fracture toughness value in group A is 735.44 N (± 146.407), in group B is 756.78 N (± 153.791), and group C is 829.00 N (± 159.290). Thus, it can be known that the strongest fracture toughness value is group C with shortest intracanal retention thickness.

These results could be caused by the nature of the SFRC material, which is a resin matrix combined with e-glass fiber and inorganic particulate fillers.¹⁴ The fiber functions as a reinforcement embedded in a polymer resin matrix to transmit the pressure received by the tooth to the surrounding tissue and protects the tooth from mechanical and environmental damage in the form of fracture.¹⁵ Basically, the matrix of the resin composites has much lower modulus than the fiber which makes the matrix easily strains more. Fibers that are incorporated into the matrix form a stress transfer and have the ability to stop the crack propagation through the material. This is the reason why fibers could enhance the physical properties of a material.¹³ In previous studies stated that flowable SFRC as intracanal retention can improve the integrity of the restoration and enable efficient stress distribution that is able to withstand fractures in composite restorations.¹⁰ In this study, flowable SFRC as intracanal retention was tested with 3 groups of thickness and does not show significant results. These results are consistent with the results of a previous study regarding the effect of fiber post length on fracture resistance conducted by Adanir and Belli¹⁶, where they also found no significant relationship between fiber post length and fracture resistance.

Resin materials have several drawbacks, one of which is experiencing shrinkage during the polymerization process (polymerization shrinkage). Polymerization is a chemical reaction process when the resin monomer molecules bind and form chain bonds which are called polymers.¹⁷ Polymerization shrinkage can cause microleakage, which causes teeth to fracture easily.³ During the polymerization process, the chains of triethylene glycol dimethacrylate (TEGDMA) can experience higher shrinkage which caused by the higher conversion of double bonds in highly flowable restorative materials such as SFRC.¹⁸ In the previous research conducted by Agwu and Ozoegwu¹⁹, cited from Tezvergil et al. that SFRC is anisotropic, which makes SFRC material with different mechanical and thermal properties in all directions. This causes shrinkage in SFRC materials to also occur in all directions with varying shrinkage levels. In addition, shrinkage will occur greater in fibers perpendicular to the direction of polymerization because the bond between the fiber and the matrix is broader in the direction of the fiber orientation that is lead to a lower fracture toughness value with more intracanal retention length. According to a study conducted by Gupta et al.²⁰ that cited from Giachetti, the degree of polymerization shrinkage stress becomes smaller when the thickness of the composite resin material is thinner. When the thickness of the material is thinner, the volume of SFRC material involved is less, which causes the stress that is formed to be smaller.^{20,21} This explains the relation between intracanal retention length and fracture toughness value, and the reason why group C with shorter intracanal retention thickness have a higher fracture toughness value. The more length of intracanal retention of SFRC, the higher shrinkage polymerization will occur, causing fracture toughness value to decrease.

According to research conducted by Sakaguchi et al., Ping et al.²² stated the stress given to the polymerization process would cause vacancies in the interfacial bonds or voids in the bond volume between fibers that would lead to material failures. Polymerization on a larger surface area or volume will result in a larger particle interfacial bond vacancy.¹⁹ The void at the fiber particle interface is caused by a decrease in wetting in the matrix, which ends in a reduction of mechanical properties and failure of the composite system. Besides that, other factors can affect the results of the study, one of which is the cavity configuration factor or c-factor. The c-factor is the surface ratio of the bonded and unbound restoration areas to the cavity walls. The more surface area of the bonded resin, the value of the c-factor will increase, resulting in an increase in polymerization shrinkage.⁶ The discussion could suggest exploring other mechanical properties of SFRC material such as diametral tensile strength, or developing the studies with another bonding technique.

CONCLUSION

This study showed no resistance of the intracanal retention thickness with flowable fiber reinforced composite materials toward fracture toughness. However, in the research that has been done, SFRC material as intracanal retention with a thickness of 4 mm shows the highest fracture toughness value.

Author Contributions: Conceptualization, T.I.A. and W.W.; methodology, T.I.A. and W.W.; software, T.I.A.; validation, T.I.A. and W.W.; formal analysis, T.I.A.; investigation, W.W.; resources, T.I.A.; data curation, T.I.A.; writing original draft preparation, T.I.A.; writing review and editing, T.I.A.; visualization, T.I.A.; supervision, W.W.; project administration, T.I.A.; funding acquisition, T.I.A. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: The study was approved by the Ethics Committee of Faculty of Dentistry, Universitas Trisakti (546/S1/KEPK/FKG/7/2022).

Informed Consent Statement: Not applicable.

Data Availability Statement: http://repository.trisakti.ac.id/usaktiana/index.php/home/detail/detail_koleksi/0/SKR/pembimbing/000000000000011341/widyastuti#menu

Conflicts of Interest: The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

REFERENCES

1. Riset Kesehatan Dasar (Riskesdas). Badan Penelitian dan Pengembangan Kesehatan Kementerian RI tahun 2018. Jakarta; 2018;7:179-217.
2. Colak H, Dülgergil CT, Dalli M, Hamidi MM. Early childhood caries update - a review of causes, diagnoses, and treatments. *J Nat Sci Biol Med.* 2013;4(1):29-38. DOI: [10.4103/0976-9668.107257](https://doi.org/10.4103/0976-9668.107257).
3. Garg N, Garg A. *Textbook of Endodontic.* 4th ed. New Delhi, India: Jaypee Brothers Medical (P). 2019;3(11):34-35.
4. Andreasen JO, Andreasen FM, Andersson L. *Textbook and color atlas of traumatic injuries to the teeth.* 5th ed. Wiley-Blackwell (P) Ltd., Oxford:UK. 2018;413-42. ISBN: 9781119167051.
5. American Association of Endodontist. *Glossary of Endodontic Terms.* 10th ed. Chicago; 2020. [cited May 2022] Available from: <https://www.aae.org/specialty/clinical-resources/glossary-endodontic-terms/>
6. Hilton TJ, Ferracane JL, Broome JC. *Summitt's fundamentals of operative dentistry: a contemporary approach.* 4th ed. Brazil, Quintessence (P) co. Inc. 2013;21:523-30. ISBN: 978-0-86715-528-0
7. Alvanforoush N, Palamara J, Wong RH, Burrow MF. Comparison between published clinical success of direct resin composite restorations in vital posterior teeth in 1995-2005 and 2006-2016 periods. *Aust. Dent. Journal.* 2016;62(2):132-145. DOI: [10.1111/adj.12487](https://doi.org/10.1111/adj.12487).
8. Heintze SD, Ilie N, Hickel R, Reis A, Loguercio A, Rousson V. Laboratory mechanical parameters of composite resins and their relation to fractures and wear in clinical trials: a systematic review. *Dent. Mater Journal.* 2016;33(3):101-114. DOI: [10.1016/j.dental.2016.11.013](https://doi.org/10.1016/j.dental.2016.11.013).
9. Lassila L, Keulemans F, Vallittu PK, Garoushi S. Characterization of restorative short fiber reinforced dental composites. *Dental Material Journal.* 2020;39(6):1-8. DOI: [10.4012/dmj.2019-088](https://doi.org/10.4012/dmj.2019-088).
10. Lassila L, Keulemans F, Säilynoja E, Vallittu PK, Garoushi S. Mechanical properties and fracture behavior of flowable fiber reinforced composite restorations. *Dental Material Journal: Elsevier Ltd.* 2018;34:598-606. DOI: [10.1016/j.dental.2018.01.002](https://doi.org/10.1016/j.dental.2018.01.002).
11. Garoushi S, Vallittu P, Lassila L. Mechanical properties and radiopacity of flowable fiber-reinforced composite. *Dental Material Journal: Elsevier Ltd.* 2019;38(2):196-202. DOI: [10.4012/dmj.2018-102](https://doi.org/10.4012/dmj.2018-102).
12. Vallittu PK. High-aspect ratio fillers: fiber-reinforced composites and their anisotropic properties. *Dental Material Journal: Elsevier Ltd.* 2015;31(1):1-7. DOI: [10.1016/j.dental.2014.07.009](https://doi.org/10.1016/j.dental.2014.07.009).
13. Abouelleil H, Pradelle N, Villat C, Attik N, Colon P, Grosogoeat B. Comparison of mechanical properties of a new fiber reinforced composite and bulk filling composites. *Restor Dent Endod.* 2015;40(4):262-70. DOI : [10.5395/rde.2015.40.4.262](https://doi.org/10.5395/rde.2015.40.4.262).
14. Garoushi S, Tanner J, Vallittu PK, Lassila L. Preliminary clinical evaluation of short fiber-reinforced composite resin in posterior teeth: 12-months report. *The Open Dentistry Journal.* 2012;6(1):41-45. DOI: [10.2174/1874210601206010041](https://doi.org/10.2174/1874210601206010041).
15. Mergulhão VA, de Mendonça LS, de Albuquerque MS, Braz R. Fracture resistance of endodontically treated maxillary premolars restored with different methods. *Operative Dentistry Journal.* 2019;44(1):1-11. DOI: [10.2341/17-262-L](https://doi.org/10.2341/17-262-L).
16. Adanir N, Belli S. Evaluation of different post lengths' effects on fracture resistance of a glass fiber post system. *European J Denstistry.* 2018;23-28.
17. Eakle WS, Bastin KG. *Dental material clinical application for dental assistants and dental hygienist* 4th ed. St. Louis: Elsevier Inc. 2021;6(4):83-116. ISBN: 978-0-323-59658-9.
18. Abbasi M, Moradi Z, Mirzaei M, Kharazifard MJ, Rezaei S. Polymerization shrinkage of five bulk-fill composite resins in comparison with a conventional composite resin. *J Dent (Tehran).* 2018;15(6):365-374.
19. Agwu N, Ozoegwu C. Critical investigation on the effect of fiber geometry and orientation on the effective mechanical properties of fiber-reinforced polymer composites. *Mechanics of Advanced Materials and Structures.* 2022. DOI: [10.1080/15376494.2022.2068206](https://doi.org/10.1080/15376494.2022.2068206).
20. Gupta S, Gupta T, Mehra M, Kaur B. Comparative evaluation of fracture resistance of endodontically treated teeth restored with different build-up materials: an in vitro study. *International Journal of Pediatric Dentistry.* 2021;14(1):51-58. DOI: [10.5005/ip-journals-10005-1901](https://doi.org/10.5005/ip-journals-10005-1901).
21. Son SA, Roh HM, Bock H, Kwon YH, Park JK. The effect of resin thickness of polymerization characteristic of siloxane-based composite resin: research article. *J Restorative Dentistry and Endodontics.* 2014;39(4):310-318. DOI: [10.5395/rde.2014.39.4.310](https://doi.org/10.5395/rde.2014.39.4.310).
22. Ping T, Zhou Y, He Y, Tang Y, Yang JL, Akram MY, Nie J. Preparation and characterization of yellowing resistance and low volume shrinkage of fluorinated polysiloxane urethane acrylate. *J Progress in Organic Coatings.* 2016;97:74-81. DOI: [10.1016/j.porgcoat.2016.03.023](https://doi.org/10.1016/j.porgcoat.2016.03.023)

The resistance of the intracanal retention thickness with flowable short fiber reinforced composite (SFRC) materials towards fracture toughness

by Wiena Widyastuti FKG

Submission date: 31-Oct-2023 08:35AM (UTC+0700)

Submission ID: 2212611289

File name: 45151-173867-1-PB.pdf (211.68K)

Word count: 3882

Character count: 20500

ORIGINAL ARTICLE

The resistance of the intracanal retention thickness with flowable short fiber reinforced composite (SFRC) materials towards fracture toughness

Isabella Audy Tania^{1*}
Wienna Widyastuti¹

¹Department of Conservative Dentistry,
Faculty of Dentistry, Universitas Trisakti,
Indonesia

*Correspondence:
wiena@trisakti.ac.id

Received: 07 February 2023
Revised: 17 March 2023
Accepted: 31 March 2023
Published: 31 March 2023
DOI: [10.24198/pjd.vol35no1.44892](https://doi.org/10.24198/pjd.vol35no1.44892)

Citation:

Tania IA, Widyastuti W. The resistance of the intracanal retention thickness with flowable short fiber reinforced composite (SFRC) materials towards fracture toughness. *Padj J Dent*. 2023;35(1):80-85

ABSTRACT

Introduction: Dental hard tissue problems were relatively high in Indonesia. If these problems were ignored, they would cause pulpal necrosis. The main treatment for this case was Root Canal Treatment (RCT). The selection of post-RCT treatment and suitable materials was important. One of the examples was short fiber reinforced composite (SFRC) which can be an option for intracanal retention treatment. This study aims to analyze the resistance of the intracanal retention thickness with flowable fiber reinforced composite materials toward fracture toughness. **Methods:** This research was using a correlative laboratory experimental method with a post-test group-only design. This study's samples were SFRC material inserted in 27 extracted mandibular first premolars with a length of 21 mm \pm 2 mm and root canal walls width of 2 mm. All samples were given intracanal retention at a depth of 2 mm below the orifice and divided into 3 groups: intracanal retention with thickness of (A) 6 mm, (B) 5 mm, and (C) 4 mm. RCT was performed using the crown-down technique. The bonding agent was applied, and then light cured for 20 seconds. SFRC was applied according to the sample group and light cured for 20 seconds per 2 mm, followed by filling with composite resin. The sample was implanted in the dental stone 2 mm below the CEJ. The fracture toughness test was performed using UTM on the occlusal surface with a 0° tilt and crosshead of 1 mm/min. **Results:** The fracture toughness results of group A, B, and C are 735.44, 756.78, and 829 respectively. Statistical tests with one-way ANOVA showed no significant difference with p value = 0.412 ($p < 0.05$). **Conclusion:** There is no difference in resistance of the intracanal retention thickness with flowable fiber reinforced composite materials towards fracture toughness.

KEYWORDS

post-RCT treatments, intracanal retentions, SFRC, fracture toughness

INTRODUCTION

Oral health is an important thing that everyone should be aware of. However, according to the Basic Health Research (Risikedas) in 2018, awareness level of oral health in Indonesia still needs to be improved. Nowadays, the main problems are hard and soft tissue diseases. Risikedas stated that the level of dental hard tissue disease in Indonesian people is still high, namely 45.3% who experience dental caries.¹ Caries can become severe problems, such as pain and discomfort, and also teeth becoming easily broken due to loss of tooth tissue structure that can lead to acute and chronic infections.² Pulp necrosis is a condition of pulp tissue death that is usually asymptomatic. The etiology of pulp tissue death can be mechanical, chemical irritants, or bacterial microorganisms.³ Tooth trauma is also one of the main causes of tooth fracture, which can involve damage to the tooth pulp.⁴

According to the 2020 American Association of Endodontists (AAE), root canal treatment (RCT) is the procedure for removing necrotic tissue from the root canal, followed by shaping and obturating the root canal with a filling material.⁵ Post-RCT teeth will become more brittle and prone to fracture. This is because the water content in the dentine tissue decreases, causing it to become dehydrated, which causes the dentine collagen to shrink. Also, the post-RCT teeth can no longer form the secondary dentine tissue, which makes dentin support for enamel reduced.⁶ As a result of these factors, post-RCT teeth become prone to fracture when they receive functional pressure or mastication.⁷

Fracture toughness is a mechanical property that describes the ability of a material to resist crack propagation and defines the level of tolerance of a material against damage.⁸ The fracture toughness value depends on the physical properties and chemical composition of the components of the restorative material. In previous studies, it was shown that there is a strong correlation between the fracture toughness of the material and the fracture rate of dental restoration materials.⁹ Therefore, materials with high fracture toughness values tend to be able to withstand the risk of micro fractures that may occur during the mastication process. In other words, materials that have higher fracture toughness values tend to have a better ability to resist the formation and spread of fractures.

It is known that fiber reinforced composite (FRC) materials have high fracture toughness values.^{10,11} The properties of FRC are highly dependent on fiber diameter, fiber length, fiber orientation or direction, fiber loading, and also the adhesion or attachment between the fibers and the polymer resin matrix.¹² The development of FRC materials is urgently needed to reduce the failure rate of RCT in dentistry, especially fractures. This study aims to analyze the resistance of the

intracanal retention thickness with flowable fiber reinforced composite materials toward fracture toughness.

METHODS

This research was conducted using a correlative laboratory experimental method with a post-test group-only design carried out at the RSGM-P Faculty of Dentistry, Universitas Trisakti and Industrial Metrology Laboratory, Universitas Trisakti, Jakarta, Indonesia. The research sample used a flowable short fiber-reinforced resin composite (SFRC) (GC EverX Flow) inserted into the mandibular first premolar as intracanal retention.

A sample of 27 extracted mandibular first premolars with single and straight root, intact crown with no caries, without root caries, and completely closed apex teeth were cleaned and then soaked in saline water. Each sample was selected by measuring the working length using a caliper, with a length of $21 \text{ mm} \pm 2 \text{ mm}$ (Figure 1).



Figure 1. Measurement with millimeter blocks.

Next, a digital radiographic photo was taken to measure the root canal walls and to ensure the length of the sample using a digital millimeter block. The length of the teeth was calculated using 2 points, namely the cusp of the premolar and the apex. The thickness of the root canal walls should be 2 mm thick from the CEJ on the outside to the root canal walls on the inside. It was then grouped into 3 groups: samples that had undergone root canal treatment and were given 2 mm intracanal retention below the orifice using SFRC flowable material with a thickness of (A) 6 mm; (B) 5 mm; (C) 4 mm (Figure 2).




Figure 2. The treatment scheme of the sample: Group A. Sample with 6 mm thickness of flowable SFRC intracanal retention; Group B: Sample with 5 mm thickness of flowable SFRC intracanal retention. Group C: Sample with 4 mm thickness of flowable SFRC intracanal retention

RCT was performed using the crown down technique using ProTaper (Dentsply Maillefer) up to size F3. At each file change, 2 ml of 5.25% NaOCl solution was irrigated in the root canals and then dried using three-way syringe and paper points. Then obturation was carried out using the single cone technique using gutta-percha and a sealer was applied using a root canal sealer (Dentsply AH 26). Finally, the obturation material was removed using a large reamer #3 (Dentsply Maillefer) 2 mm below the orifice for all samples.

Bonding material (GC Solare Bond) was applied to the tooth sample and dried, then light cured for 20 seconds. Next, intracanal retention material was applied 2 mm deep under the orifice for all samples and light cured for 20 seconds per 2 mm thickness. Furthermore, the filling was carried out using a composite resin filling material (GC Solare Sculpt Composite). These thicknesses from group A, B, and C were chosen based on previous studies¹³, which found that the fiber length differences between 1–2 mm could change the pattern of the fractures and greatly enhance the fracture resistance of the composite (Figure 3).

The tooth sample was implanted in type II dental stone 2 mm deep below the CEJ so that it resembled the position of the teeth on the alveolar bone. Furthermore, the fracture toughness test was performed using the Universal Testing Machine (Hounsfield: 50kN, England). The pressure was applied using a fracture toughness jig on the occlusal surface with a slope of 0° along the tooth axis apically. Constant pressure was applied at a crosshead speed of 1 mm/min (American Society for Testing Materials Standard) until the tooth failed in the form of a macroscopic fracture in the

intracanal retention of the SFRC material. Fractures of fillings and teeth that do not involve intracanal retention were ignored.



Figure 3. Fracture on intracanal retention with SFRC materials.

Statistical analysis was performed on the fracture toughness data. Shapiro-Wilk normality test was performed, and the p -value > 0.05 was considered as normal data distribution. Then followed by one-way ANOVA test data analysis. If the One Way ANOVA test obtained p -value < 0.05 , the analysis was continued with the post hoc test with Tukey's HSD. If the p -value shows > 0.05 , the conclusion was that there is no significant difference in the fracture toughness values of the sample groups A, B, and C, and the post hoc test was not carried out.

RESULTS

The statistical results showed in the Shapiro-Wilk normality test (Table 1), data is considered normally distributed if the p -value > 0.05 . They are then followed by One Way ANOVA data analysis (Table 3) with a p -value = 0.412. This means that there is no significant difference in the fracture toughness values of the sample groups A, B, and C.

Table 1. Shapiro-Wilk test of Normality on fracture toughness data.

Flowable SFRC	Shapiro-Wilk		
	Statistic	df	p-value
6 mm	0.200	9	0.692*
5 mm	0.200	9	0.949*
4 mm	0.200	9	0.356*

The results of the One Way ANOVA analysis (Table 2) showed that the average fracture toughness value in group A with an intracanal retention thickness of 6 mm had the lowest fracture toughness value at 735.44 N (± 146.407). Group B, with an intracanal retention of 5 mm, had a fracture toughness value of 756.78 N (± 153.791). While the sample in group C with intracanal retention thickness of 4 mm had the highest fracture toughness value at 829.00 N (± 159.290). As can be seen in bar diagram (Figure 4). The results of the homogeneity test showed that there was no significant relationship between the sample groups ($p > 0.05$).

Table 2. One Way ANOVA analysis on fracture toughness data.

	n	Mean	Std. Deviation	Std. Error	Lower Bound	Upper Bound	Min	Max
6 mm	9	735.44	146.407	48.802	622.91	847.98	548	1002
5 mm	9	756.78	153.791	51.264	638.56	874.98	502	972
4 mm	9	829.00	159.290	53.097	706.56	951.44	589	1013
Total	27	773.74	152.788	29.404	713.90	834.18	502	1013




Figure 4. Fracture toughness data of group A, B, and C.

7
Table 3. One Way ANOVA analysis.

	Sum of Squares	df	Mean square	F	p-value
Between groups	43271.407	2	21635.704	0.921	0.412
Within groups	563679.778	24	23486.657		
Total	606951.185	26			

DISCUSSION

The test data using UTM found that the average fracture toughness value was group C > B > A. Based on the Shapiro-Wilk normality test, group A has a significance value of 0.692, group B has 0.949, and group C has 0.356. So that the significance value (p/Sig.) in this normality test can be concluded that the data is normally distributed. This conclusion is obtained from the significance value of all groups of more than 0.05. Therefore, the assumption of normality of the data is met and data analysis can be continued using the one-way ANOVA test. On the One Way ANOVA test data, it is known that the average fracture toughness value in group A is 735.44 N (± 146.407), in group B is 756.78 N (± 153.791), and group C is 829.00 N (± 159.290). Thus, it can be known that the strongest fracture toughness value is group C with shortest intracanal retention thickness.

These results could be caused by the nature of the SFRC material, which is a resin matrix combined with e-glass fiber and inorganic particulate fillers.¹⁴ The fiber functions as a reinforcement embedded in a polymer resin matrix to transmit the pressure received by the tooth to the surrounding tissue and protects the tooth from mechanical and environmental damage in the form of fracture.¹⁵ Basically, the matrix of the resin composites has much lower modulus than the fiber which makes the matrix easily strains more. Fibers that are incorporated into the matrix form a stress transfer and have the ability to stop the crack propagation through the material. This is the reason why fibers could enhance the physical properties of a material.¹³ In previous studies stated that flowable SFRC as intracanal retention can improve the integrity of the restoration and enable efficient stress distribution that is able to withstand fractures in composite restorations.¹⁰ In this study, flowable SFRC as intracanal retention was tested with 3 groups of thickness and does not show significant results. These results are consistent with the results of a previous study regarding the effect of fiber post length on fracture resistance conducted by Adanir and Belli¹⁶, where they also found no significant relationship between fiber post length and fracture resistance.

Resin materials have several drawbacks, one of which is experiencing shrinkage during the polymerization process (polymerization shrinkage). Polymerization is a chemical reaction process when the resin monomer molecules bind and form chain bonds which are called polymers.¹⁷ Polymerization shrinkage can cause microleakage, which causes teeth to fracture easily.³ During the polymerization process, the chains of triethylene glycol dimethacrylate (TEGDMA) can experience higher shrinkage which caused by the higher conversion of double bonds in highly flowable restorative materials such as SFRC.¹⁸ In the previous research conducted by Agwu and Ozoegwu¹⁹, cited from Tezvergil et al. that SFRC is anisotropic, which makes SFRC material with different mechanical and thermal properties in all directions. This causes shrinkage in SFRC materials to also occur in all directions with varying shrinkage levels. In addition, shrinkage will occur greater in fibers perpendicular to the direction of polymerization because the bond between the fiber and the matrix is broader in the direction of the fiber orientation that is lead to a lower fracture toughness value with more intracanal retention length. According to a study conducted by Gupta et al.²⁰ that cited from Giachetti, the degree of polymerization shrinkage stress becomes smaller when the thickness of the composite resin material is thinner. When the thickness of the material is thinner, the volume of SFRC material involved is less, which causes the stress that is formed to be smaller.^{20,21} This explains the relation between intracanal retention length and fracture toughness value, and the reason why group C with shorter intracanal retention thickness have a higher fracture toughness value. The more length of intracanal retention of SFRC, the higher shrinkage polymerization will occur, causing fracture toughness value to decrease.

According to research conducted by Sakaguchi et al., Ping et al.²² stated the stress given to the polymerization process would cause vacancies in the interfacial bonds or voids in the bond volume between fibers that would lead to material failures. Polymerization on a larger surface area or volume will result in a larger particle interfacial bond vacancy.¹⁹ The void at the fiber particle interface is caused by a decrease in wetting in the matrix, which ends in a reduction of mechanical properties and failure of the composite system. Besides that, other factors can affect the results of the study, one of which is the cavity configuration factor or c-factor. The c-factor is the surface ratio of the bonded and unbound restoration areas to the cavity walls. The more surface area of the bonded resin, the value of the c-factor will increase, resulting in an increase in polymerization shrinkage.⁵ The discussion could suggest exploring other mechanical properties of SFRC material such as diametral tensile strength, or developing the studies with another bonding technique.

CONCLUSION

This study showed no resistance of the intracanal retention thickness with flowable fiber reinforced composite materials toward fracture toughness. However, in the research that has been done, SFRC material as intracanal retention with a thickness of 4 mm shows the highest fracture toughness value.

Author Contributions: Conceptualization, T.I.A. and W.W.; methodology, T.I.A. and W.W.; software, T.I.A.; validation, T.I.A. and W.W.; formal analysis, T.I.A.; investigation, W.W.; resources, T.I.A.; data curation, T.I.A.; writing original draft preparation, T.I.A.; writing review and editing, T.I.A.; visualization, T.I.A.; supervision, W.W.; project administration, T.I.A.; funding acquisition, T.I.A. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: The study was approved by the Ethics Committee of Faculty of Dentistry, Universitas Trisakti (546/S1/KEPK/FKG/7/2022).

Informed Consent Statement: Not applicable.

Data Availability Statement: http://repository.trisakti.ac.id/usaktiana/index.php/home/detail/detail_koleksi/0/SKR/pembimbing/00000000000011341/widyastuti#menu

Conflicts of Interest: The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

REFERENCES

- Riset Kesehatan Dasar (Riskesdas). Badan Penelitian dan Pengembangan Kesehatan Kementerian RI tahun 2018. Jakarta; 2018;7:179-217.
- Colak H, Dülgeril CT, Dalli M, Hamidi MM. Early childhood caries update - a review of causes, diagnoses, and treatments. *J Nat Sci Biol Med.* 2013;4(1):29-38. DOI: [10.4103/0976-9668.107257](https://doi.org/10.4103/0976-9668.107257).
- Garg N, Garg A. *Textbook of Endodontic.* 4th ed. New Delhi, India: Jaypee Brothers Medical (P). 2019;3(11):34-35.
- Andreasen JO, Andreasen FM, Andersson L. *Textbook and color atlas of traumatic injuries to the teeth.* 5th ed. Wiley-Blackwell (P) Ltd., Oxford:UK. 2018;413-42. ISBN: 9781119167051.
- American Association of Endodontist. *Glossary of Endodontic Terms.* 10th ed. Chicago; 2020. [cited May 2022] Available from: <https://www.aae.org/specialty/clinical-resources/glossary-endodontic-terms/>
- Hilton TJ, Ferracane JL, Broome JC. *Summitt's fundamentals of operative dentistry: a contemporary approach.* 4th ed. Brazil, Quintessence (P) co. Inc. 2013;21:523-30. ISBN: 978-0-86715-528-0
- Alvanforoush N, Palamara J, Wong RH, Burrow MF. Comparison between published clinical success of direct resin composite restorations in vital posterior teeth in 1995-2005 and 2006-2016 periods. *Aust. Dent. Journal.* 2016;62(2):132-145. DOI: [10.1111/adj.12487](https://doi.org/10.1111/adj.12487).
- Heintze SD, Ilie N, Hickel R, Reis A, Loguercio A, Rousson V. Laboratory mechanical parameters of composite resins and their relation to fractures and wear in clinical trials: a systematic review. *Dent. Mater Journal.* 2016;33(3):101-114. DOI: [10.1016/j.dental.2016.11.013](https://doi.org/10.1016/j.dental.2016.11.013).
- Lassila L, Keulemans F, Vallittu PK, Garoushi S. Characterization of restorative short fiber reinforced dental composites. *Dental Material Journal.* 2020;39(6):1-8. DOI: [10.4012/dmj.2019-088](https://doi.org/10.4012/dmj.2019-088).
- Lassila L, Keulemans F, Säilynoja E, Vallittu PK, Garoushi S. Mechanical properties and fracture behavior of flowable fiber reinforced composite restorations. *Dental Material Journal: Elsevier Ltd.* 2018;34:598-606. DOI: [10.1016/j.dental.2018.01.002](https://doi.org/10.1016/j.dental.2018.01.002).
- Garoushi S, Vallittu P, Lassila L. Mechanical properties and radiopacity of flowable fiber-reinforced composite. *Dental Material Journal: Elsevier Ltd.* 2019;38(2):196-202. DOI: [10.4012/dmj.2018-102](https://doi.org/10.4012/dmj.2018-102).
- Vallittu PK. High-aspect ratio fillers: fiber-reinforced composites and their anisotropic properties. *Dental Material Journal: Elsevier Ltd.* 2015;31(1):1-7. DOI: [10.1016/j.dental.2014.07.009](https://doi.org/10.1016/j.dental.2014.07.009).
- Abouelleil H, Pradelle N, Villat C, Attik N, Colon P, Grosogeat B. Comparison of mechanical properties of a new fiber reinforced composite and bulk filling composites. *Restor Dent Endod.* 2015;40(4):262-70. DOI: [10.5395/rde.2015.40.4.262](https://doi.org/10.5395/rde.2015.40.4.262).
- Garoushi S, Tanner J, Vallittu PK, Lassila L. Preliminary clinical evaluation of short fiber-reinforced composite resin in posterior teeth: 12-months report. *The Open Dentistry Journal.* 2012;6(1):41-45. DOI: [10.2174/1874210601206010041](https://doi.org/10.2174/1874210601206010041).
- Mergulhão VA, de Mendonça LS, de Albuquerque MS, Braz R. Fracture resistance of endodontically treated maxillary premolars restored with different methods. *Operative Dentistry Journal.* 2019;44(1):1-11. DOI: [10.2341/17-262-L](https://doi.org/10.2341/17-262-L).
- Adanir N, Belli S. Evaluation of different post lengths' effects on fracture resistance of a glass fiber post system. *European J Dentistry.* 2018;23-28.
- Eakle WS, Bastin KG. *Dental material clinical application for dental assistants and dental hygienist* 4th ed. St. Louis: Elsevier Inc. 2021;6(4):83-116. ISBN: 978-0-323-59658-9.
- Abbasi M, Moradi Z, Mirzaei M, Kharazifard MJ, Rezaei S. Polymerization shrinkage of five bulk-fill composite resins in comparison with a conventional composite resin. *J Dent (Tehran).* 2018;15(6):365-374.
- Agwu N, Ozoegwu C. Critical investigation on the effect of fiber geometry and orientation on the effective mechanical properties of fiber-reinforced polymer composites. *Mechanics of Advanced Materials and Structures.* 2022. DOI: [10.1080/15376494.2022.2068206](https://doi.org/10.1080/15376494.2022.2068206).
- Gupta S, Gupta T, Mehra M, Kaur B. Comparative evaluation of fracture resistance of endodontically treated teeth restored with different build-up materials: an in vitro study. *International Journal of Pediatric Dentistry.* 2021;14(1):51-58. DOI: [10.5005/ip-journals-10005-1901](https://doi.org/10.5005/ip-journals-10005-1901).
- Son SA, Roh HM, Bock H, Kwon YH, Park JK. The effect of resin thickness of polymerization characteristic of siloxane-based composite resin: research article. *J Restorative Dentistry and Endodontics.* 2014;39(4):310-318. DOI: [10.5395/rde.2014.39.4.310](https://doi.org/10.5395/rde.2014.39.4.310).
- Ping T, Zhou Y, He Y, Tang Y, Yang JL, Akram MY, Nie J. Preparation and characterization of yellowing resistance and low volume shrinkage of fluorinated polysiloxane urethane acrylate. *J Progress in Organic Coatings.* 2016;97:74-81. DOI: [10.1016/j.porgcoat.2016.03.023](https://doi.org/10.1016/j.porgcoat.2016.03.023)

<https://doi.org/10.24198/pjd.vol35no1.44892> Copyright: © 2022 by the authors. Submitted to Padjadjaran journal of dentistry for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

The resistance of the intracanal retention thickness with flowable short fiber reinforced composite (SFRC) materials towards fracture toughness

ORIGINALITY REPORT

6%

SIMILARITY INDEX

4%

INTERNET SOURCES

5%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Universitas Diponegoro Student Paper	2%
2	www.jstage.jst.go.jp Internet Source	1%
3	www.thejcdp.com Internet Source	1%
4	wjarr.com Internet Source	1%
5	"Full Issue PDF", Operative Dentistry, 2020 Publication	1%
6	Submitted to Udayana University Student Paper	1%
7	neuropsychiatricinvestigation.org Internet Source	1%
8	link.springer.com Internet Source	<1%

Exclude quotes On

Exclude matches < 15 words

Exclude bibliography On

The resistance of the intracanal retention thickness with flowable short fiber reinforced composite (SFRC) materials towards fracture toughness

GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/0

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
