

JURNAL KESMAS

COVER JURNAL



EDITORIAL BOARD

Jurnal Kesehatan Masyarakat
[Home](#) [Current](#) [Archives](#) [Archive 2009 - 2023](#) [About](#) ▼

[Home](#) / [Editorial Team](#)

Editorial Team

Editor-in-Chief

- Prof. Dr. dr. Oktia Woro Kasmini Handayani, M.Kes, {SCOPUS ID : 57192428885} {Public Health Nutrition} Universitas Negeri Semarang, Indonesia

Editorial Advisory Regional America

- Dr. Gina Samaan, {SCOPUS ID : 6602382950} {Epidemiology} US. Centers for Disease Control and Prevention, United States

Editorial Advisory Regional Asia

- Prof. Kathirvelu Baskar, {SCOPUS ID : 55092286200} {Entomology} Loyola College India, Entomology Research

Editorial Advisory Regional Asia

- Prof. Kathirvelu Baskar, {SCOPUS ID : 55092286200} {Entomology} Loyola College India, Entomology Research Institute, India
- Dr. Khalid M. Al Aboud, {SCOPUS ID : 7003345190} {Dermatology} King Faisal Specialist Hospital and Research Centre, Saudi Arabia
- Prof. Dato'. Syed Mohamed Aljunid, {SCOPUS ID : 6504304159} {Health Economics, Policy and Management} Universiti Kebangsaan Malaysia, Malaysia
- Dr. Dina Nur Anggraini Ningrum, {SCOPUS ID : 57195329470} {Health Information System} Taipei Medical University, Taiwan, Province of China
- Dr. Mahalul Azam, {SCOPUS ID: 57194196255} {Medical} Universitas Negeri Semarang, Indonesia
- Dr. Songpol Tornee, {SCOPUS ID : 6506180249} {Health Education} Shrinakharinwirot University, Thailand

Editorial Advisory Regional Australia

- Prof. Doune Macdonald, PhD, {SCOPUS ID : 7401463393} {Health Education} University of Queensland, Australia

Editorial Advisory Regional Africa

- Assoc. Prof. Dr. Henry Odhianosen Imhonde, {SCOPUS ID : 36069265600} {Psychology} Ambrose Alli University, Nigeria

Editorial Board

- Muhammad Azinar, S.K.M, M.Kes, {SCOPUS ID : 57194193079} {Health Promotion} Universitas Negeri Semarang, Indonesia
- Nur Siyam, S.K.M, M.PH, {SCOPUS ID : 57222668801} {Maternal and Child Health} Universitas Negeri Semarang, Indonesia
- Efa Nugroho, S.K.M, M.Kes, {SCOPUS ID : 57192436111} {Reproduction Health} Universitas Negeri Semarang, Indonesia

Administration

- Satria Adi Rachim, Universitas Negeri Semarang, Indonesia
- Widiyanto Widiyanto, Universitas Terbuka, Indonesia

PARAMETER INDEXING



KEMAS: Jurnal Kesehatan Masyarakat [P-ISSN 1858-1196 | E-ISSN 2355-3596] published by Universitas Negeri Semarang in collaboration with [Ikatan Ahli Kesehatan Masyarakat Indonesia](#) (IAKMI Tingkat Pusat) and [Jejaring Nasional Pendidikan Kesehatan](#) (JNPK). KEMAS publishes the article based on research or equivalent to research results in public health or other disciplines related to public health that has not been loaded/published by other media. The journal contains articles about epidemiology and biostatistics, health policy and administration, public health nutrition, environmental health, occupational health and safety, health promotion, reproductive health, maternal and child health, and other related articles in public health. The journal can be used by health practitioners, health caregivers, teachers, medical students, and people who are interested in public health issues. The journal was first published in July 2005. Since 2022, the journal were published quarterly a year in July, October, January, and April. Starting October 2022, articles in KEMAS Journal have been accepted for indexing to [Scopus](#).

Abstracting & Indexing

[GARUDA](#) | [SINTA](#) | [DOAJ](#) | [DIMENSION](#) | [SCOPUS](#)

PARAMETER INDEXING



DAFTAR ISI

Vol. 21 No. 2 (2025)

DOI: <https://doi.org/10.15294/kemas.v21i2>

Published: 2025-10-03

Articles

Environmental Factors and Efficacy of Castor Seed Influencing *Aedes aegypti* Larval Presence

DOI: <https://doi.org/10.15294/kemas.v21i2.16069>

Indra Chahaya, Winni R. E. Tumanggor, Lanova Dwi Arde, Khairunnisa, Najwa Shadrina, Lyra Caroline D Purba (Author) 261-271
Article ID 16069



hpmA Gene as a Detection Method of *Proteus mirabilis* Bacteria using real-time Polymerase Chain Reaction

DOI: <https://doi.org/10.15294/kemas.v21i2.16214>

Muktiningsih Nurjayadi, Gladys Indira Putri, Anisa Fitriyanti, Jefferson Lynford Declan, Novitasari, Bassam Abomoelak (Author) 272-280
Article ID 16214



Breeding Site Preferences and Resistance Status of *Aedes aegypti* in Malang City

DOI: <https://doi.org/10.15294/kemas.v21i2.21666>

Moh Mirza Nuryady, Ely Purwanti, Tutut Indria Permana, Kiky Martha Ariesaka (Author) 281-289
Article ID 21666



Vitamin D and Lifestyle Factors in Active Smoker in Indonesia

DOI: <https://doi.org/10.15294/kemas.v21i2.22618>

Amelia Lorensia, Rivan Virlando Suryadinata, Marthy Meliana Ariyanti Jalmar, Pande Made Ayu Aprianti, Angela Nofvianti Cahyo Wati, Zahwa Dhiba (Author) 290-300
Article ID 22618



PARAMETER INDEXING

Kemas

Q4

Health Policy

best quartile

SJR 2024

0.14

powered by scimagojr.com

Jurnal Kesehatan Masyarakat

S1

H-INDEX 54

I5-INDEX 47



Work-Related Factors, Exercise Habits, and Individual Characteristics on Musculoskeletal Disorders among Indonesian Young Dentists

DOI: <https://doi.org/10.15294/kemas.v21i2.17220>

Dwi Windu Kinanti Arti, Tiwakron Prachalboon, Zita Aprillia, Diki Bima Prasetyo (Author)

301-308

Article ID 17220

 PDF

Health Beliefs and Husband's Preparedness in Supporting Postpartum Contraceptive Use

DOI: <https://doi.org/10.15294/kemas.v21i2.24085>

Pipit Feriani, Komang Menik Sri Krisnawati, Rini Ernawati, Yuliani Winarti (Author)

309-320

Article ID 24085

 PDF

Moringa Leaves-Added Food, Baby Massage, and Tui Na Massage to Reduce Stunting Rate on Young Children in Indonesia

DOI: <https://doi.org/10.15294/kemas.v21i2.24094>

Eka Sutrisna, Husna Maulida, Fauzan Saputra, Yunitasari, Arista Ardila, Suriani, Ristiani (Author)

321-332

Article ID 24094

 PDF

Quality of Life Among Women at Menopause

DOI: <https://doi.org/10.15294/kemas.v21i2.24246>

Marni Marni, Siti Farida, Putri Halimu Husna , Wahyuningsih, Domingos Soares, Kresna Agung Yidhianto (Author)

333-340

Article ID 24246

 PDF

Telemonitoring-Based Cardiac Care to Reduce Readmissions in Coronary Heart Disease Patients

DOI: <https://doi.org/10.15294/kemas.v21i2.24553>

Hariyono Hariyono, Sri Pantja Madyawati, Leo Yosdimyati Romli, Chin Xuan Tan, Shifa Fauziyah, Teguh Hari Sucipto (Author)

341-347

Article ID 24553

 PDF

Nutritional Status of Children Aged 6-59 Months Based on Composite Index of Anthropometric Failure

DOI: <https://doi.org/10.15294/kemas.v21i2.25264>

Rizka Pratiwi, Trini Sudiarti (Author)

348-359

Article ID 25264

 PDF

Handgrip Strength of Public Works Personnel in West Jakarta

DOI: <https://doi.org/10.15294/kemas.v21i2.25917>

Alvina Alvina, Pusparini Pusparini, Mario Mario, Yasmine Mashabi (Author)

360-368

Article ID 25917



Perinatal Mental Health Disorders in Indonesia: A Systematic Review of Quantitative Studies (2015–2024)

DOI: <https://doi.org/10.15294/kemas.v21i2.26267>

Noviyati Rahardjo Putri, Ulinuha Aufa Rahmah, Grhasta Dian Perestroika (Author)

369-377

Article ID 26267



Understanding Cadre Coping Mechanisms in Utilizing Digital Tools for Stunting Intervention Programs

DOI: <https://doi.org/10.15294/kemas.v21i2.24587>

Astriaana Marta Batubara, Judhiastuty Februhartanty, Dwi Nastiti Iswarawanti (Author)

378-385

Article ID 24587



Hookah Smoking in Incidence with Asymptomatic Gout

DOI: <https://doi.org/10.15294/kemas.v21i2.25033>

Israa A. Mohammed Jumaah Jumaah, Abdulrahman Ahmed Mahmood Mahmood, Shatha A. Demerchi (Author)

386-391

Article ID 25033



Brand Image as a Determinant of Patient Decision-Making in Inpatient Healthcare Utilization

DOI: <https://doi.org/10.15294/kemas.v21i2.25500>

Martianawati, Tuffaillah Muawina Nur, Ahmad Lathifi (Author)

392-401

Article ID 25500



Preoperative Anxiety Levels and the Incidence of Postoperative Nausea and Vomiting in Patients Undergoing General Anesthesia

DOI: <https://doi.org/10.15294/kemas.v21i2.27115>

Widigdo Rekso Negoro, Sindu Sintara, Annes Rindy Permana, Muhammad Rodli, Reko Priyonggo, Suryanto (Author)

402-410

Article ID 27115



Pentahelix Collaboration in Achieving Disaster Preparedness through Resilient Villages: A Systematic Literature Review

DOI: <https://doi.org/10.15294/kemas.v21i2.29453>

Dwi Yunanto Hermawan, Evi Widowati, Sofwan Indarjo, Efa Nugroho, Annisa Novanda Maharani Utami (Author)

411-422

Article ID 29453



The Role of Decentralized Health Systems in Shaping Service Quality: A Systematic Review in Low- and Middle-Income Countries

DOI: <https://doi.org/10.15294/kemas.v21i2.29250>

Eri Witcahyo Witcahyo, Asobat Gani, I Made Suwandi (Author)

423-433

Article ID 29250



Giving Moringa Soy Milk on Weight Gain and Upper Arm Circumference of Pregnant Women

DOI: <https://doi.org/10.15294/kemas.v21i2.29036>

Syafaatun Nahdyah, Mardiana Ahmad, Veni Hadju, Suryani As'ad, Jumrah (Author)

434-442

Article ID 29036



Giving Moringa Soy Milk on Weight Gain and Upper Arm Circumference of Pregnant Women

DOI: <https://doi.org/10.15294/kemas.v21i2.29036>

Syafaatun Nahdyah, Mardiana Ahmad, Veni Hadju, Suryani As'ad, Jumrah (Author)

434-442

Article ID 29036



Screen Time, Anthropometric Parameter, Insulin and HOMA IR in Adolescents

DOI: <https://doi.org/10.15294/kemas.v21i2.6265>

Nur Aisiyah Widjaja, Rizka Arifani, Tausiyah Rohmah Noviyanti, Yoppi Yeremia Alexander, Edi Hermanto, Iitdrie Iitdrie, Eva Ardianah, Rino Tryanto Keya (Author)

443-453

Article ID 6265





Handgrip Strength of Public Works Personnel in West Jakarta

Alvina^{1✉}, Pusparini¹, Mario¹, Yasmine Mashabi¹

¹Department of Clinical Pathology, Faculty of Medicine, Universitas Trisakti

Article Info

Article History:

Submitted: May 2025

Accepted: September 2025

Published: October 2025

Keywords:

Handgrip strength; Body mass index; Hip circumference; Waist circumference; Fasting blood glucose

DOI

<https://doi.org/10.15294/kemas.v21i2.25917>

Abstract

Public Works Personnel (PWP) in Jakarta perform their routine duties manually instead of using equipment. Handgrip strength (HGS) measures maximum hand strength as a quality indicator for muscle strength and mass. Factors affecting muscle strength are age, sex, body mass index (BMI), waist circumference (WC), and hip circumference (HC). Increased fasting blood glucose (FBG) is also associated with muscle quality, muscle strength, and physical performance. This study aimed to determine the factors associated with handgrip strength in PWP. A total of 192 male PWP from the Cengkareng district were recruited by simple random sampling. The collected data were demographics, BMI, FBG, WC, HC, and HGS. Statistical analysis was done using Pearson's correlation test and multiple regression analysis. There was a significant positive correlation of BMI, WC, and HC with HGS ($p=0.006$; $r=0.20$, $p=0.009$; $r=0.19$, and $p=0.005$; $r=0.20$) and a significant negative correlation between age and HGS ($p=0.008$; $r=-0.19$) but not between FBG and HGS ($p=0.847$). Multiple regression analysis showed height, BMI, and HC positively predicting HGS ($R^2=0.19$). Handgrip strength of public works personnel is associated with BMI, age, waist circumference, and hip circumference, but not with fasting blood glucose. Height, BMI, and hip circumference predict handgrip strength

Introduction

Personnel for the management of public infrastructure and facilities at the village level are workers who manage village public works for a given time period on the basis of a work order. The tasks of village-level public works personnel (PWP) consist of managing the infrastructure and facilities, such as streets, drains, and parks, including installed lights (Gubernur Provinsi Daerah Khusus Ibukota Jakarta, 2017). The work performed by village PWP consists mainly of intense physical activity involving the hands, such that handgrip strength is essential in their tasks. Lowering of handgrip strength may indicate reduced muscle strength, which may especially interfere with the daily activities of village PWP that consist mainly of manual labor.

Handgrip strength measures maximal hand strength and indicates muscle quality that reflects muscle strength and mass. Muscle

strength refers to the ability of muscle groups to raise a given load. Low or poor grip strength is frequently associated with damaging health impacts, such as chronic disease, and is the initial sign of reduced muscle mass, limitations in physical functions, and mortality risk. Poor handgrip strength may also reduce a person's ability to perform activities of daily living and, therefore, impact quality of life (Soraya *et al.*, 2023; Nathania *et al.*, 2023). Muscle mass accounts for approximately 40% of total weight, such that knowledge of muscle mass conditions is essential in maintaining health and activities (Handayani *et al.*, 2018).

Handgrip strength may represent the strength of all muscles in the body. Several factors possibly influencing an individual's muscle strength are age, sex, and body mass index (BMI) (Soraya *et al.*, 2023). Obesity may also affect BMI. Obesity in adults is not only associated with genetic factors but also with

✉ Correspondence Address:
Department of Clinical Pathology, Faculty of Medicine, Universitas Trisakti
Email: dr.alvina@trisakti.ac.id

environmental factors during development (Fauzi *et al.*, 2022). Handgrip strength is also associated with chronic diseases such as diabetes, metabolic syndrome, and cardiovascular disease. Increased glucose concentration is also associated with disorders of muscle quality and strength, as well as physical performance (Chen *et al.*, 2023). A reduction in muscle mass also affects an individual's strength and physical endurance. A combination of proteins can maintain muscle function, thereby building body strength and physical endurance (Soenyoto *et al.*, 2017). In the present study, we included fasting blood glucose concentrations in the data collection to obtain more accurate measures of glucose concentrations that have not yet been influenced by food intake.

Handgrip strength may be a health indicator that provides information on a person's health (Vaishya *et al.*, 2024). However, its clinical significance in determining the prevention and treatment of poor handgrip remains limited because of the still unclear etiology of muscle weakness, especially because the clinical outcomes associated with handgrip strength are extremely diverse (McGrath *et al.*, 2020). The relationship between BMI and handgrip strength is still controversial, with several investigators finding a significant positive association and others reporting a non-significant relationship (Al Asadi, 2018). There are few studies on handgrip strength, especially on workers in the sanitation sector, such as public works personnel, which is why the present investigators were interested in studying this topic. The present study aimed to determine the factors associated with handgrip strength in public works personnel.

Method

This cross-sectional study involving male public works personnel in Cengkareng District, West Jakarta, was conducted from September to October 2024. The study sample was selected by simple random sampling. Calculation of the sample size for a correlation with $r=0.2$, $Z\alpha=0.05$, and $Z\beta=0.1$ resulted in a sample of 192 subjects who met the inclusion criterion of being male and 25-60 years old, and the exclusion criteria of having an abnormality of the hands, heart disease, pulmonary disease,

and malignancies.

Measurements were obtained on height in cm, weight in kg, as well as waist circumference and hip circumference in cm. Height and weight were determined using microtoise and digital scales, respectively. BMI was calculated using the formula of weight divided by height squared. Measurement of waist and hip circumferences was done using a measuring tape. Waist circumference was determined with the subject in the upright position, at the end of a normal expiration, at the smallest abdominal circumference between the iliac crest and twelfth rib. The cut-off values for waist circumference in males and females are ≥ 90 cm and ≥ 80 cm, respectively. Hip circumference was measured using a measuring tape that was applied around the hips at the level of the symphysis pubis and the maximal gluteal circumference (Rokhmah *et al.*, 2015).

Fasting blood glucose (FBG) concentration was determined on venous blood by the hexokinase method after the respondents had fasted for approximately 10 hours prior. Handgrip strength was measured with a handgrip strength dynamometer to the nearest kilogram as follows: The respondent was asked to sit down comfortably without reclining and with the feet completely relaxed. The hips and knees were positioned at a 90° angle, the gripping hand with the shoulder in adduction, the elbow flexed 90° , and the forearm and wrist in the neutral position. The gripping duration was 3 seconds. The respondent gripped the instrument with the dominant hand as hard as possible, and the value indicated by the needle was recorded. In accordance with the protocol recommended by the American Society of Hand Therapists (ASHT), the recorded handgrip strength was the average of three handgrip strength tests (Nathania *et al.*, 2023).

Before data analysis by means of the correlation test, the normality of the data distribution was determined using the Kolmogorov-Smirnov test. The Pearson correlation test was used for a normal data distribution, whereas for a non-normal data distribution, the Spearman correlation test was done, followed by a multiple regression analysis. Ethics approval was issued by the Ethics Committee of the Faculty of Medicine,

Universitas Trisakti, under No. 033/KER/FK/08/2024.

Result and Discussion

In this study, the respondents comprised 192 male public works personnel of Cengkareng District who consisted of the villages (subdistricts) of Kedaung, Duri Kosambi, West Cengkareng, Kapuk, and Rawa Buaya. A total of

26% of the respondents were residents of Duri Kosambi, where Muslims comprised 98.4% of the population. The majority of the respondents, at 77.6% had an educational level of senior high school. Nearly all of the respondents, or 99%, were right-handed, using the right hand as the dominant one at work, and most were of the Betawi (Jakarta) ethnic group at 51.6% (Table 1).

Table 1. Demographic Data of the Respondents

Characteristic	N (%)
Village	
Kedaung	40 (20.8)
Duri Kosambi	50 (26)
Cengkareng Barat	33 (17.2)
Kapuk	43 (22.4)
Rawa Buaya	26 (13.5)
Religion	
Islamic	189 (98.4)
Christian	2 (1)
Catholic	1 (0.5)
Level of education	
Elementary school	7 (3.6)
Junior high school	29 (15.1)
Senior high school	149 (77.6)
Academy-University	7 (3.6)
Dominant hand	
Right hand	190 (99)
Left hand	2 (1)
Ethnic group	
Javanese	53 (27.6)
Betawi	99 (51.6)
Sundanese	25 (13)
Other	14 (7.3)

Legend: Other ethnicities comprised groups from Palembang, Lampung, Bima, and Kupang
Source: primary data 2024 (questionnaires and interviews with respondents)

Table 2. Respondents' Characteristics

Variable	Mean ± SD
Age (years)	40.9 ± 8.9
Height (cm)	164.6 ± 6.7
Weight (kg)	64.7 ± 13.1
BMI (kg/m²)	23.8 ± 4.3
Waist circumference (cm)	85.4 ± 10.4
Hip circumference (cm)	96.9 ± 8.4

HGS (kg)	33.1 ± 6.9
FBG (mg/dL)	87 (64-280)*

Legend: BMI: body mass index. *FBG: fasting blood glucose, expressed as Median (Min-Max).

HGS: handgrip strength

Source: primary data (respondents' measurements)

Table 3. Analysis of the Relationship of Age, Height, Weight, BMI, Waist Circumference, Hip Circumference, and Fasting Blood Glucose Concentration with Handgrip Strength

Variable	R	P value
Age	- 0.19	0.008*
Height	0.36	0.000*
Weight	0.32	0.000*
BMI	0.20	0.006*
Waist circumference	0.19	0.009*
Hip circumference	0.20	0.005*
FBG	0.02	0.847 [#]

*P value with Pearson's test, [#] Spearman's test

Source: primary data

Table 4. Multiple Regression Analysis on Handgrip Strength

Determinant	b	SE	p
Height	0.48	0.08	0.000
BMI	0.79	0.22	0.000
Hip circumference	- 0.29	0.12	0.015
Constant	- 36.98		

R²=0,19

Source: primary data

The bivariate analysis found a significant positive correlation of BMI, height, weight, waist circumference, and hip circumference with handgrip strength ($p=0.006$, $r=0.20$; $p<0.001$, $r=0.36$; $p<0.001$, $r=0.32$; $p=0.009$, $r=0.19$; $p=0.005$; $r=0.20$, respectively). This showed that the larger the BMI, height, weight, waist circumference, and hip circumference, the stronger the handgrip. However, there was no significant relationship between fasting glucose concentration and handgrip strength ($p=0.847$). This study also found that age had a significant negative correlation with handgrip strength ($p=0.011$; $r=-0.182$), signifying that the older a person, the lower the handgrip strength (Table 3). Results of multiple regression analysis on handgrip strength are listed in Table 4.

This study found that most of the public works personnel in Duri Kosambi village were Muslims, had a senior high school level of education, and were of Betawi ethnicity. This was in line with the demographic data in the Cengkareng district, in that Duri Kosambi was

the village with the greatest area, most of the residents were male, and the Islamic religion had the most believers (Badan Pusat Statistik Jakarta Barat, 2020). Among our respondents, more than ten percent were in the diabetes category. Socio-demographic factors capable of affecting dietary habits in patients with diabetes are the level of education and the type of family (Rondhianto *et al.*, 2024). Among the respondents of this study, the dominant right hand was used for working. In the adult population, a total of 90% of the residents preferred to use the right hand for manual tasks, whereas around 10% used the left hand (Zaccagni *et al.*, 2020). According to Agtuahene *et al.* (Agtuahene *et al.*, 2023), the dominant hand is generally around 10% stronger than the non-dominant hand. Many activities in daily life need a high level of exertion of the muscles of the hand and forearm (Hammed *et al.*, 2017). The frequent use of the dominant hand in daily living results in the muscles of the hand and forearm providing strength in gripping (Afaq *et*

al., 2019).

The mean handgrip strength in this study was 33.1 ± 6.9 kg, which is relatively low if compared with the study results of Kurniawan *et al.* (Kurniawan *et al.*, 2018). In healthy young Indonesian adults, the mean handgrip strength in males was 37.37 ± 8.29 kg, and the lower limit of muscle strength in Indonesian males was 20.79 kg. The handgrip strength results in both studies are still below those of males in other Southeast Asian countries, according to the study results of Leong *et al.* (Leong *et al.*, 2016). These investigators found that the mean handgrip strength in Southeast Asian males aged 35-40 years was 40 kg (34-44 kg); for the age range of 41-50 years, the handgrip strength was 38 kg (33-44 kg), for the age range of 51-60 years it was 34 kg (30-40 kg), and for ages 61-70 years 30 kg (24-34 kg). This may have been caused by ethnic differences mediated by genetic differences that also affect individual muscle strength. The highest handgrip strength is found in the European and North American populations, whereas the lowest handgrip strength is found in African, South Asian, and Southeast Asian populations. These variations in muscle strength in various countries may also be partly caused by differences in socioeconomic status (Leong *et al.*, 2016).

According to Agtuahene *et al.* (Agtuahene *et al.*, 2023), handgrip strength is a physiological variable influenced by several factors, such as age, sex, and body measurements. In contrast, according to Zaccagni *et al.* (Zaccagni *et al.*, 2020), handgrip strength, apart from being affected by age, may also be influenced by physical activity, grip measurement, height, and body mass. The present study found a significant positive correlation between BMI and handgrip strength. According to Hammed *et al.*, this may be caused by the higher percentage of skeletal muscle mass compared to the percentage of fat mass, thereby resulting in a higher handgrip strength. Our study's results are identical to those of Agtuahene *et al.* (Agtuahene *et al.*, 2023), who found a significant positive correlation ($r=0.290$) of handgrip strength with BMI and height. This shows that taller individuals with higher BMI apparently have greater handgrip strength. Our study is also in line with the study of Krakauer in 2020, which

showed a positive correlation between BMI and handgrip strength (Krakauer *et al.*, 2020).

The study of Cooper *et al.* (Cooper *et al.*, 2022) concluded that a high childhood BMI is associated with higher grip strength at age 46 years. This is because fat mass acts as a mechanical load that elicits an anabolic response that stimulates muscle growth and function. This anabolic response is usually greater in males than in females because of the higher testosterone concentration. However, it should also be noted that BMI does not differentiate between fat mass and lean mass as adipocyte indicators that may vary with sex. It should be noted that BMI is a measure of body fat that does not differentiate between weight changes because of an increase or decrease in muscle mass or body fat (Zaccagni *et al.*, 2020). According to Jafar *et al.* (Jaafar *et al.*, 2023), heavier individuals have a greater muscle mass, resulting in greater handgrip strength, than lighter individuals. In the study of Heidy *et al.* (Heidy *et al.*, 2019), the investigators found a positive correlation between height and forearm length. The study by Sirajudeen *et al.* (Sirajudeen *et al.*, 2012) found a positive correlation between hand length and handgrip strength. This shows that height affects the longer extremity or arm, thus approaching optimal muscle length that may result in greater grip strength. According to a prospective cohort study, handgrip strength in middle age appears to be associated with birth weight and the increase in height at puberty. Pubertal growth and handgrip strength possibly represent the development of muscle size and growth (Schrager *et al.*, 2007).

Our study found no relationship between blood glucose concentration and handgrip strength. Our study agrees with the study of Giglio *et al.* (Giglio *et al.*, 2018), which showed no significant relationship between muscle mass strength and blood glucose concentration. According to Giglio, this is because most respondents are individuals who routinely perform structured physical activity, even though they are in the light category. This is similar to our study respondents, who were public works personnel and who had performed physical activity such as street sweeping and cleaning storm drains. Glucose is the main fuel

for muscle contraction, entering the muscle cells by diffusion through the GLUT4 glucose transporter, which, after muscle contraction, moves from its intracellular storage site to the plasma membrane and T-tubules (Richter *et al.*, 2013). Glucose metabolism has an influence on handgrip strength through various mechanisms, such as disorders of glucose metabolism, particularly glycogenolysis, that may result in the loss of muscle strength. Insulin resistance in hyperglycemia may cause muscle degradation because hyperglycemia causes low muscle strength through effects on skeletal muscle mitochondria. The lowering of muscle strength by hyperglycemia occurs through Krüppel-like factor 15, which regulates skeletal muscle lipid flux. Hyperglycemia also increases pro-inflammatory cytokines, which may increase the catabolism that plays a role in the further decrease in muscle mass and quality (Gamayanti *et al.*, 2023). The amount of carbohydrates also influences the glycemic response and post-prandial insulin level, impacting the supply of glycogen to the muscles (Safitri *et al.*, 2020).

This study has a significant negative correlation between age and handgrip strength, which decreases with age (Manoharan *et al.*, 2015). Handgrip strength initially experiences an increase with increasing age, reaches its peak between the ages of 30 and 40 years, and decreases afterward (Zaccagni *et al.*, 2020). The age-related changes in skeletal muscle composition, particularly the accumulation of lipids in skeletal muscle fibers, play a role in the poor quality of the muscles and may result in metabolic abnormalities such as insulin resistance. Apart from changes in muscle quality, there is also a reduction in muscle mass that consistently occurs during aging. The increase in oxidative stress and chronic inflammation that commonly occurs in the aging process apparently also plays a role in the development of decreased muscle mass and strength (Mainous *et al.*, 2015). The decrease in muscle mass in the elderly corresponds to a 10% reduction in grip strength per decade, starting from the age of 40 years (Jaafar *et al.*, 2023). Skeletal muscle plays an important role in glucose metabolism and has a significant influence on insulin sensitivity. Increased

insulin resistance in older adults may be closely associated with skeletal muscle aging. A reduction in GLUT4 expression as a result of a decrease in muscle volume causes a reduction in insulin sensitivity in aging skeletal muscle (Joo *et al.*, 2022). The results of this study showed a correlation of waist circumference and hip circumference with handgrip strength. This is in line with the study of Widjaja *et al.* (Widjaja *et al.*, 2018), who stated that there was a significant correlation of waist circumference and hip circumference with handgrip strength ($p < 0.001$, $r = 0.302$; and $p < 0.001$, $r = 0.296$). A greater waist circumference is associated with a greater handgrip strength in males (Hardy *et al.*, 2013).

In the multiple regression analysis of HGS, we removed the data on body weight because of the occurrence of multicollinearity between weight and BMI, with a correlation coefficient of more than 0.9, indicating an extremely strong relationship between these two variables. Conceptually, only one of the variables is used in multivariate analysis. The multiple regression analysis included age, height, BMI, waist circumference, and hip circumference, with HGS as the dependent variable in a stepwise manner. After three iterations, only height, BMI, and hip circumference played a role in predicting HGS by means of the regression formula, which resulted in $R^2 = 0.19$. This signifies that for known values of height, BMI, and hip circumference, we can predict the HGS value 19% more accurately than without knowledge of these data. For a clinical diagnosis, a minimum increase of 10% is needed. This study found that height, BMI, and hip circumference play a role in predicting handgrip strength through the regression formula, i.e., $\text{handgrip strength} = -36.98 + 0.48 \times \text{height} + 0.79 \times \text{BMI} - 0.29 \times \text{hip circumference}$. Handgrip strength and waist circumference are surrogate muscle strength and visceral adiposity markers, respectively (Nakanishi *et al.*, 2024). Waist circumference is also more sensitive in determining body fat distribution, particularly in the abdomen. Waist circumference correlates better with abdominal fat distribution than BMI (Rokhmah *et al.*, 2015). Waist circumference and hip circumference are associated with obesity, in which the fat

content is higher, and the skeletal muscle and fascicular content is lower, thereby resulting in a lower contractile capacity of skeletal muscle (Afaq *et al.*, 2019). The limitation of this study is that it did not evaluate other factors that may affect handgrip strength, such as nutrient intake. From the anthropometric measures of the hands and feet in this study, the causes and effects of the decrease in handgrip strength could not be determined because this study used a cross-sectional design.

Conclusion

Three factors may be used to estimate handgrip strength in PWP, namely height, BMI, and hip circumference, whereas blood glucose concentration is not correlated with handgrip strength in PWP. It is recommended to pay attention to height, BMI, and hip circumference, particularly in the acceptance criteria of PWP candidates, and to conduct further studies with consideration of other factors that affect handgrip strength, such as food intake and anthropometric features of the hands and arms.

Acknowledgments

Financial support was provided by Universitas Trisakti for this research.

References

- Afaq, E., Hassan, S.H., Khan, M.R., Zaidi, D.A., & Baktashi, H., 2019. Correlation of Hand Dynamometry With Body Mass Index and Waist-Hip Ratio in Medical Students. *Pak J Physiol.*, 15, pp.19-21.
- Agtuahene, M.A., Quartey, J., & Kwakye, S., 2023. Influence of Hand Dominance, Gender, and Body Mass Index on Hand Grip Strength. *S Afr J Physiother.*, 79, pp.1-6.
- Al-Asadi, J.N., 2018. Handgrip Strength in Medical Students: Correlation with Body Mass Index and Hand Dimensions. *Asian Journal of Medical Sciences*, 9, pp.21-26.
- Badan Pusat Statistik Kota Administrasi Jakarta Barat., 2020. *Kecamatan Cengkareng Dalam Angka*, pp.1-114.
- Chen, D., Zhu, Y., Ni, W., Li, Y., Yin, G., Shao, Z., & Zhu, J., 2023. Hand Grip Strength is Inversely Associated with Total Daily Insulin Dose Requirement in Patients with Type 2 Diabetes Mellitus: A Cross-Sectional Study. *PeerJ.*, pp.1-13.
- Cooper, R., Tomlinson, D., Hamer, M., & Pereira, S.M.P., 2022. Lifetime Body Mass Index and Grip Strength at Age 46 Years: The 1970 British Cohort Study. *J Cachexia Sarcopenia Muscle*, 13, pp.1-10.
- Fauzi, L., Handayani, O.W.K., Susilo, M.T., Kurnia, A.R., Rahayu, S.R., Irawan, F.A., Horng Lu, F.J., Lin, C., Lai, M.F., & Yu, Y.C., 2022. Obesity in Indonesian and Taiwanese Adolescents Related to Self Perception, Diet, Exercise, and Body Image. *KEMAS*, 17(3), pp.453-461.
- Gamayanti, K.A.V., Aryana, I.G.P.S., & Gotera, W., 2023. Hubungan Antara Kekuatan Genggaman Tangan Dengan Kadar Gula Darah Sewaktu Pada Lansia Di Desa Melinggih, Kecamatan Payangan, Kabupaten Gianyar. *Intisari Sains Medis*, 14(2), pp.483-488.
- Giglio, B.M., Mota, J.F., Wall, B.T., & Pimentel, G.D., 2018. Low Handgrip Strength is Not Associated With Type 2 Diabetes Mellitus and Hyperglycemia: A Population-Based Study. *Clin Nutr Res.*, 7(2), pp.112-116.
- Hammed, A.I., & Obaseki, C.O., 2017. Interdependence of Body Mass Index with Handgrip Strength and Endurance Among Apparently Healthy Teenagers. *Turk J Kin.*, 4, pp.1-7.
- Handayani, M.D.N., Sadewa, A.H., Farmawati, A., & Rochmah, W., 2018. Anthropometric Prediction Equations for Estimating Muscle Mass of Elderly Women. *KEMAS*, 14(2), pp.195-204.
- Hardy, R., Cooper, R., Sayer, A.A., Shlomo, Y.B., Cooper, C., Deary, I.J., Demakakos, P., Gallacher, J., Martin, R.M., McNeill, G., Starr, J.M., Steptoe, A., Syddall, H., & Kuh, D., 2013. Body Mass Index, Muscle Strength, and Physical Performance in Older Adults From Eight Cohort Studies: The Halcyon Programme. *PLoS One*, 8(2), pp.e56483.
- Heidy., Djuartina, T., & Irawan, R., 2019. Korelasi Kekuatan Genggaman Tangan Dengan Karakter Antropometri Lengan Bawah dan Tangan Serta Indeks Massa Tubuh. *Damianus Journal of Medicine*, 18, pp.1-7.
- Ilmi, A.F., & Utari, D.M., 2020. Hubungan Lingkar Pinggang dan Rasio Lingkar Pinggang-Panggul (RLPP) Terhadap Kadar Glukosa Darah Puasa Pada Mahasiswa. *Journal of Nutrition College*, 9, pp.222-227.
- Jaafar, M.H., Ismail, R., Ismail, N.H., Isa, Z.M., Tamil, A.M., Nasir, N.M., Keat, N.K., Razak, N.H., Abidin, N.Z., & Yusof, K.H., 2023. Normative Reference Values and Predicting Factors of Handgrip Strength For Dominant

- and Non-Dominant Hands Among Healthy Malay Adults in Malaysia. *BMC Musculoskeletal Disord.*, 24, pp.1-9.
- Joo, K.C., Son, D.H., & Park, J.M., 2022. Association Between Relative Handgrip Strength and Insulin Resistance in Korean Elderly Men Without Diabetes: Findings Of The 2015 Korea National Health Nutrition Examination Survey. *Korean J Fam Med.*, 43, pp.199-205.
- Krakauer, N.Y., & Krakauer, J.C., 2020. Association of Body Shape Index (ABSI) with Hand Grip Strength. *Int J Environ Res Public Health*, 17, pp.1-12.
- Kurniawan, A., Widjaja, D., & Lugito, N.P.H., 2018. *Mean and Cutoff Value of Hand Grip Strength For Healthy Indonesian Young Adults*. World Congress on Osteoporosis, Osteoarthritis and Musculoskeletal Diseases; Krakow, Poland.
- Leong, D.P., Teo, K.K., Rangarajan, S., Kutty, R., Lanas, F., Hui, C., Quanyong, X., Zhenzhen, Q., Jinhua, T., Noorhassim, I., AlHabib, K.F., Moss, S.J., Rosengren, A., Akalin, A.A., Rahman, O., Chifamba, J., Orlandini, A., Kumar, R., Yeates, K., Gupta, R., Yusufali, A., Dans, A., Avezum, A., Lopez-Jaramillo, P., Poirier, P., Heidari, H., Zatonska, K., Iqbal, R., Khatib, R., & Yusuf, S., 2016. Reference Ranges of Handgrip Strength From 125,462 Healthy Adults in 21 Countries: A Prospective Urban Rural Epidemiologic (PURE) Study. *J Cachexia Sarcopenia Muscle*, 7, pp.535-546.
- Mainous, A.G., Tanner, R.J., Anton, S.D., & Jo, A., 2015. Grip Strength as A Marker of Hypertension and Diabetes in Healthy Weight Adults. *Am J Prev Med*, 49(6), pp.850-858.
- Manoharan, V.S., Sundaram, S.G., & Jason, J.I., 2015. Factors Affecting Hand Grip Strength and Its Evaluation: A Systemic Review. *Int J Physiother Res.*, 3(6), pp.1288-1293.
- McGrath, R., Johnson, N., Klawitter, L., Mahoney, S., Trautman, K., Carlson, C., Rockstad, E., & Hackney, K.J., 2020. What Are The Association Patterns Between Handgrip Strength and Adverse Health Conditions? A Topical Review. *SAGE Open Medicine*, 8, pp.1-12.
- Nakanishi, S., Shimoda, M., Kimura, T., Sanada, J., Fushimi, Y., Iwamoto, Y., Iwamoto, H., Dan, K., Mune, T., Kaku, K., & Kaneto, H., 2024. The Impact of Handgrip Strength and Waist Circumference on Glycemic Control: Prospective, Observational Study Using Outpatient Clinical Data in Japanese Patients With Type 2 Diabetes Mellitus. *J Diabetes Investig.*, 15(7), pp.892-898.
- Nathania, N.P.S., Nugraha, M.H.S., Primayanti, I.D.A., & Saraswati, P.A.S., 2023. Handgrip Strength is Associated With Balance in Elderly Women. *Majalah Ilmiah Fisioterapi Indonesia*, 11, pp.70-75.
- Gubernur Provinsi Daerah Khusus Ibukota Jakarta., 2017. *Peraturan Gubernur Provinsi Daerah Khusus Ibukota Jakarta Nomor 7 tahun 2017 Tentang Penanganan Prasarana dan Sarana Umum Tingkat Kelurahan*. Jakarta 2017.
- Richter, E.A., & Hargreaves, M., 2013. Exercise, GLUT4, and Skeletal Muscle Glucose Uptake. *Physiol Rev.*, 93(3), pp.993-1017.
- Rokhmah, F.D., Handayani, D., & Al-Rasyid, H., 2015. Korelasi Lingkar Pinggang dan Rasio Lingkar Pinggang-Panggul Terhadap Kadar Glukosa Plasma Menggunakan Tes Toleransi Glukosa Oral. *Jurnal Gizi Klinik Indonesia*, 12, pp.28-35.
- Rondhianto., Ridla, A.Z., & Hasan, H., 2024. Sociodemographic Factors Affecting Diabetic Dietary Behavior in People with Type 2 Diabetes Mellitus. *KEMAS*, 19(3), pp.429-437.
- Safitri, I., Setyarsih, L., Susanto, H., Suhartono., & Fitranti, D.Y., 2020. The Effect of Low and High Glycemic Load Diet on Muscle Fatigue of Young Soccer Athletes. *KEMAS*, 16(1), pp.111-120.
- Schrager, M.A., Metter, E.J., Simonsick, E., Ble, A., Bandinelli, S., Lauretani, F., & Ferrucci, L., 2007. Sarcopenic Obesity and Inflammation in The InChianti Study. *J Appl Physiol.*, 102(3), pp.919-925.
- Sirajudeen, M.S., Shah, U.N., Pillai, P.S., Mohasin, N., & Shantaram, M., 2012. Correlation Between Grip Strength and Physical Factors in Men. *International Journal of Health and Rehabilitation Sciences*, 1, pp.58-63.
- Soenyoto, T., Mollap, K., & Mungpong, J., 2017. Correlation of Energy and Protein Consumption Levels with Physical Endurance of Rhythmic Gymnast Athletes. *KEMAS*, 13(1), pp.131-136.
- Soraya, N., & Parwanto, E., 2023. The Controversial Relationship Between Body Mass Index and Handgrip Strength in The Elderly: An Overview. *Malays J Med Sci.*, 30, pp.73-83.
- Vaishya, R., Misra, A., Vaish, A., Ursino, N., & D'Ambrosi, R., 2024. Hand Grip Strength as A Proposed New Vital Sign Of Health: A Narrative Review of Evidences. *Journal of Health, Population and Nutrition*, 43, pp.1-

14.

- Widjaja, D., Tjhai, K., Dermawan, K., Kalim, J., Tanzil, S., Pramusinta, A., Phenca, H.K., Kurniawan, A., 2018. *Correlations Between Hand Grip Strength and Anthropometric Status in Young Indonesian Adults*. World Congress on Osteoporosis, Osteoarthritis and Musculoskeletal Diseases; Krakow, Poland.
- Zaccagni, L., Toselli, S., Bramanti, B., Gualdi-Russo, E., Mongillo, J., & Rinaldo, N., 2020. Handgrip Strength in Young Adults: Association With Anthropometric Variables and Laterality. *Int J Environ Res Public Health*, 17, pp.1-18.

Handgrip strength of public works personnel in West Jakarta

by Alvina

Submission date: 06-Oct-2025 10:54AM (UTC+0700)

Submission ID: 2400879299

File name: as-Handgripstrength_of_public_work_personel_in_West_Jakarta.pdf (211.87K)

Word count: 4941

Character count: 25964

Handgrip Strength of Public Works Personnel in West Jakarta

Alvina^{1✉}, Pusparini¹, Mario¹, Yasmine Mashabi¹

Article Info

Keywords:
Handgrip strength; Body mass index; Hip circumference; Waist circumference; Fasting blood glucose

Abstract

Public Works Personnel (PWP) in Jakarta perform their routine duties manually instead of using equipment. Handgrip strength (HGS) measures maximum hand strength as a quality indicator for muscle strength and mass. Factors affecting muscle strength are age, sex, body mass index (BMI), waist circumference (WC), and hip circumference (HC). Increased fasting blood glucose (FBG) is also associated with muscle quality, muscle strength, and physical performance. This study aimed to determine the factors associated with handgrip strength in PWP. A total of 192 male PWP from the Cengkareng district were recruited by simple random sampling. The collected data demographics, BMI, FBG, WC, HC, and HGS. Statistical analysis was done using Pearson's correlation test and multiple regression analysis. There was a significant positive correlation of BMI, WC, and HC with HGS ($p=0.006$; $r=0.20$, $p=0.009$; $r=0.19$, and $p=0.005$; $r=0.20$) and a significant negative correlation between age and HGS ($p=0.008$; $r=-0.19$) but not between FBG and HGS ($p=0.847$). Multiple regression analysis showed height, BMI, and HC positively predicting HGS ($R^2=0.19$). Handgrip strength of public works personnel is associated with BMI, age, waist circumference, and hip circumference, but not with fasting blood glucose. Height, BMI, and hip circumference predict handgrip strength

Introduction

Personnel for the management of public infrastructure and facilities at the village level are workers who manage village public works for a given time period on the basis of a work order. The tasks of village-level public works personnel (PWP) consist of managing the infrastructure and facilities, such as streets, drains, and parks, including installed lights (Gubernur Provinsi Daerah Khusus Ibukota Jakarta, 2017). The work performed by village PWP consists mainly of intense physical activity involving the hands, such that handgrip strength is essential in their tasks. Lowering of handgrip strength may indicate reduced muscle strength, which may especially interfere with the daily activities of village PWP that consist mainly of manual labor.

Handgrip strength measures maximal hand strength and indicates muscle quality that reflects muscle strength and mass. Muscle

strength refers to the ability of muscle groups to raise a given load. Low or poor grip strength is frequently associated with damaging health impacts, such as chronic disease, and is the initial sign of reduced muscle mass, limitations in physical functions, and mortality risk. Poor handgrip strength may also reduce a person's ability to perform activities of daily living and, therefore, impact quality of life (Soraya *et al.*, 2023; Nathania *et al.*, 2023). Muscle mass accounts for approximately 40% of total weight, such that knowledge of muscle mass conditions is essential in maintaining health and activities (Handayani *et al.*, 2018).

Handgrip strength may represent the strength of all muscles in the body. Several factors possibly influencing an individual's muscle strength are age, sex, and body mass index (BMI) (Soraya *et al.*, 2023). Obesity may also affect BMI. Obesity in adults is not only associated with genetic factors but also with

environmental factors during development (Izumi *et al.*, 2022). Handgrip strength is also associated with chronic diseases such as diabetes, metabolic syndrome, and cardiovascular disease. Increased glucose concentration is also associated with disorders of muscle quality and strength, as well as physical performance (Chen *et al.*, 2023). A reduction in muscle mass also affects an individual's strength and physical endurance. A combination of proteins can maintain muscle function, thereby building body strength and physical endurance (Soenyoto *et al.*, 2017). In the present study, we included fasting blood glucose concentrations in the data collection to obtain more accurate measures of glucose concentrations that have not yet been influenced by food intake.

Handgrip strength may be a health indicator that provides information on a person's health (Vaishya *et al.*, 2024). However, its clinical significance in determining the prevention and treatment of poor handgrip remains limited because of the still unclear etiology of muscle weakness, especially because the clinical outcomes associated with handgrip strength are extremely diverse (McGrath *et al.*, 2020). The relationship between BMI and handgrip strength is still controversial, with several investigators finding a significant positive association and others reporting a non-significant relationship (Al Asadi, 2018). There are few studies on handgrip strength, especially on workers in the sanitation sector, such as public works personnel, which is why the present investigators were interested in studying this topic. The present study aimed to determine the factors associated with handgrip strength in public works personnel.

Method

This cross-sectional study involving male public works personnel in Cengkareng District, West Jakarta, was conducted from September to October 2024. The study sample was selected by simple random sampling. Calculation of the sample size for a correlation with $r=0.2$, $Z_{\alpha}=0.05$, and $Z_{\beta}=0.1$ resulted in a sample of 192 subjects who met the inclusion criterion of being male and 25-60 years old, and the exclusion criteria of having an abnormality of the hands, heart disease, pulmonary disease,

and malignancies.

Measurements were obtained on height in cm, weight in kg, as well as waist circumference and hip circumference in cm. Height and weight were determined using microtoise and digital scales, respectively. BMI was calculated using the formula of weight divided by height squared. Measurement of waist and hip circumferences was done using a measuring tape. Waist circumference was determined with the subject in the upright position, at the end of a normal expiration, at the smallest abdominal circumference between the xiphoid crest and twelfth rib. The cut-off values for waist circumference in males and females are ≥ 90 cm and ≥ 80 cm, respectively. Hip circumference was measured using a measuring tape that was applied around the hips at the level of the symphysis pubis and the maximal gluteal circumference (Rokhmah *et al.*, 2015).

Fasting blood glucose (FBG) concentration was determined on venous blood by the hexokinase method after the respondents had fasted for approximately 10 hours prior. Handgrip strength was measured with a handgrip strength dynamometer to the nearest kilogram as follows: The respondent was asked to sit down comfortably without reclining and with the feet completely relaxed. The hips and knees were positioned at a 90° angle, the gripping hand with the shoulder in adduction, the elbow flexed 90° , and the forearm and wrist in the neutral position. The gripping duration was 3 seconds. The respondent gripped the instrument with the dominant hand as hard as possible, and the value indicated by the needle was recorded. In accordance with the protocol recommended by the American Society of Hand Therapists (ASHT), the recorded handgrip strength was the average of three handgrip strength tests (Nathania *et al.*, 2023).

Before data analysis by means of the correlation test, the normality of the data distribution was determined using the Kolmogorov-Smirnov test. The Pearson correlation test was used for a normal data distribution, whereas for a non-normal data distribution, the Spearman correlation test was done, followed by a multiple regression analysis. Ethics approval was issued by the Ethics Committee of the Faculty of Medicine,

Universitas Trisakti, under No. 033/KER/ FK/08/2024.

Result and Discussion

In this study, the respondents comprised 192 male public works personnel of Cengkareng District who consisted of the villages (subdistricts) of Kedaung, Duri Kosambi, West Cengkareng, Kapuk, and Rawa Buaya. A total of

26% of the respondents were residents of Duri Kosambi, where Muslims comprised 98.4% of the population. The majority of the respondents, at 77.6% had an educational level of senior high school. Nearly all of the respondents, or 99%, were right-handed, using the right hand as the dominant one at work, and most were of the Betawi (Jakarta) ethnic group at 51.6% (Table 1).

Table 1. Demographic Data of the Respondents

Characteristic	N (%)
Village	
Kedaung	40 (20.8)
Duri Kosambi	50 (26)
Cengkareng Barat	33 (17.2)
Kapuk	43 (22.4)
Rawa Buaya	26 (13.5)
Religion	
Islamic	189 (98.4)
Christian	2 (1)
Catholic	1 (0.5)
Level of education	
Elementary school	7 (3.6)
Junior high school	29 (15.1)
Senior high school	149 (77.6)
Academy-University	7 (3.6)
Dominant hand	
Right hand	190 (99)
Left hand	2 (1)
Ethnic group	
Javanese	53 (27.6)
Betawi	99 (51.6)
Sundanese	25 (13)
Other	14 (7.3)

Legend: Other ethnicities comprised groups from Palembang, Lampung, Bima, and Kupang
Source: primary data 2024 (questionnaires and interviews with respondents)

Table 2. Respondents' Characteristics

Variable	Mean ± SD
Age (years)	40.9 ± 8.9
Height (cm)	164.6 ± 6.7
Weight (kg)	64.7 ± 13.1
BMI (kg/m²)	23.8 ± 4.3
Waist circumference (cm)	85.4 ± 10.4
Hip circumference (cm)	96.9 ± 8.4

HGS (kg)	33.1 ± 6.9
FBG (mg/dL)	87 (64-280)*

Legend: BMI: body mass index. *FBG: fasting blood glucose, expressed as Median (Min-Max).
HGS: handgrip strength
Source: primary data (respondents' measurements)

Table 3. Analysis of the Relationship of Age, Height, Weight, BMI, Waist Circumference, Hip Circumference, and Fasting Blood Glucose Concentration with Handgrip Strength

Variable	R	P value
Age	- 0.19	0.008*
Height	0.36	0.000*
Weight	0.32	0.000*
BMI	0.20	0.006*
Waist circumference	0.19	0.009*
Hip circumference	0.20	0.005*
FBG	0.02	0.847'

*P value with Pearson's test, 'Spearman's test
Source: primary data

Table 4. Multiple Regression Analysis on Handgrip Strength

Determinant	b	SE	p
Height	0.48	0.08	0.000
BMI	0.79	0.22	0.000
Hip circumference	- 0.29	0.12	0.015
Constant	- 36.98		

R²=0,19
Source: primary data

The bivariate analysis found a significant positive correlation of BMI, height, weight, waist circumference, and hip circumference with handgrip strength (p=0.006, r=0.20; p<0.001, r=0.36; p<0.001, r=0.32; p=0.009, r=0.19; p=0.005; r=0.20, respectively). This showed that the larger the BMI, height, weight, waist circumference, and hip circumference, the stronger the handgrip. However, there was no significant relationship between fasting glucose concentration and handgrip strength (p=0.847). This study also found that age had a significant negative correlation with handgrip strength (p=0.011; r=-0.182), signifying that the older a person, the lower the handgrip strength (Table 3). Results of multiple regression analysis on handgrip strength are listed in Table 4.

This study found that most of the public works personnel in Duri Kosambi village were Muslims, had a senior high school level of education, and were of Betawi ethnicity. This was in line with the demographic data in the Cengkareng district, in that Duri Kosambi was

the village with the greatest area, most of the residents were male, and the Islamic religion had the most believers (Badan Pusat Statistik Jakarta Barat, 2020). Among our respondents, more than ten percent were in the diabetes category. Socio-demographic factors capable of affecting dietary habits in patients with diabetes are the level of education and the type of family (Rondhianto *et al.*, 2024). Among the respondents of this study, the dominant right hand was used for working. In the adult population, a total of 90% of the residents preferred to use the right hand for manual tasks, whereas around 10% used the left hand (Zaccagni *et al.*, 2020). According to Agtuahene *et al.* (Agtuahene *et al.*, 2023), the dominant hand is generally around 10% stronger than the non-dominant hand. Many activities in daily life need a high level of exertion of the muscles of the hand and forearm (Hammed *et al.*, 2017). The frequent use of the dominant hand in daily living results in the muscles of the hand and forearm providing strength in gripping (Afaq *et*

et al., 2019).

The mean handgrip strength in this study was 33.1 ± 6.9 kg, which is relatively low if compared with the study results of Kurniawan *et al.* (Kurniawan *et al.*, 2018). In healthy young Indonesian adults, the mean handgrip strength in males was 37.37 ± 8.29 kg, and the lower limit of muscle strength in Indonesian males was 20.79 kg. The handgrip strength results in both studies are still below those of males in other Southeast Asian countries, according to the study results of Leong *et al.* (Leong *et al.*, 2016). These investigators found that the mean handgrip strength in Southeast Asian males aged 35-40 years was 40 kg (34-44 kg); for the age range of 41-50 years, the handgrip strength was 38 kg (33-44 kg), for the age range of 51-60 years it was 34 kg (30-40 kg), and for ages 61-70 years 30 kg (24-34 kg). This may have been caused by ethnic differences mediated by genetic differences that also affect individual muscle strength. The highest handgrip strength is found in the European and North American populations, whereas the lowest handgrip strength is found in African, South Asian, and Southeast Asian populations. These variations in muscle strength in various countries may also be partly caused by differences in socio-economic status (Leong *et al.*, 2016).

According to Agtuahene *et al.* (Agtuahene *et al.*, 2023), handgrip strength is a physiological variable influenced by several factors, such as age, sex, and body measurements. In contrast, according to Zaccagni *et al.* (Zaccagni *et al.*, 2020), handgrip strength, apart from being affected by age, may also be influenced by physical activity, grip measurement, height, and body mass. The present study found a significant positive correlation between BMI and handgrip strength. According to Hammed *et al.*, this may be caused by the higher percentage of skeletal muscle mass compared to the percentage of fat mass, thereby resulting in a higher handgrip strength. Our study's results are identical to those of Agtuahene *et al.* (Agtuahene *et al.*, 2023), who found a significant positive correlation ($r=0.290$) of handgrip strength with BMI and height. This shows that taller individuals with higher BMI apparently have greater handgrip strength. Our study is also in line with the study of Krakauer in 2020, which

showed a positive correlation between BMI and handgrip strength (Krakauer *et al.*, 2020).

The study of Cooper *et al.* (Cooper *et al.*, 2022) concluded that a high childhood BMI is associated with higher grip strength at age 46 years. This is because fat mass acts as a mechanical load that elicits an anabolic response that stimulates muscle growth and function. This anabolic response is usually greater in males than in females because of the higher testosterone concentration. However, it should also be noted that BMI does not differentiate between fat mass and lean mass as adipocyte indicators that may vary with sex. It should be noted that BMI is a measure of body fat that does not differentiate between weight changes because of an increase or decrease in muscle mass or body fat (Zaccagni *et al.*, 2020). According to Jafar *et al.* (Jaafar *et al.*, 2023), heavier individuals have a greater muscle mass, resulting in greater handgrip strength, than lighter individuals. In the study of Heidt *et al.* (Heidt *et al.*, 2019), the investigators found a positive correlation between height and forearm length. The study by Sirajudeen *et al.* (Sirajudeen *et al.*, 2012) found a positive correlation between hand length and handgrip strength. This shows that height affects the longer extremity or arm, thus approaching optimal muscle length that may result in greater grip strength. According to a prospective cohort study, handgrip strength in middle age appears to be associated with birth weight and the increase in height at puberty. Pubertal growth and handgrip strength possibly represent the development of muscle size and growth (Schrager *et al.*, 2007).

Our study found no relationship between blood glucose concentration and handgrip strength. Our study agrees with the study of Giglio *et al.* (Giglio *et al.*, 2018), which showed no significant relationship between muscle mass strength and blood glucose concentration. According to Giglio, this is because most respondents are individuals who routinely perform structured physical activity, even though they are in the light category. This is similar to our study respondents, who were public works personnel and who had performed physical activity such as street sweeping and cleaning storm drains. Glucose is the main fuel

for muscle contraction, entering the muscle cells by diffusion through the GLUT4 glucose transporter, which, after muscle contraction, moves from its intracellular storage site to the plasma membrane and T-tubules (Richter *et al.*, 2013). Glucose metabolism has an influence on handgrip strength through various mechanisms, such as disorders of glucose metabolism, particularly glycogenolysis, that may result in the loss of muscle strength. Insulin resistance in hyperglycemia may cause muscle degradation because hyperglycemia causes low muscle strength through effects on skeletal muscle mitochondria. The lowering of muscle strength by hyperglycemia occurs through Krüppel-like factor 15, which regulates skeletal muscle lipid flux. Hyperglycemia also increases pro-inflammatory cytokines, which may increase the catabolism that plays a role in the further decrease in muscle mass and quality (Gamayanti *et al.*, 2023). The amount of carbohydrates also influences the glycemic response and post-prandial insulin level, impacting the supply of glycogen to the muscles (Safitri *et al.*, 2020).

This study has a significant negative correlation between age and handgrip strength, which decreases with age (Manoharan *et al.*, 2015). Handgrip strength initially experiences an increase with increasing age, reaches its peak between the ages of 30 and 40 years, and decreases afterward (Zaccagni *et al.*, 2020). The age-related changes in skeletal muscle composition, particularly the accumulation of lipids in skeletal muscle fibers, play a role in the poor quality of the muscles and may result in metabolic abnormalities such as insulin resistance. Apart from changes in muscle quality, there is also a reduction in muscle mass that consistently occurs during aging. The increase in oxidative stress and chronic inflammation that commonly occurs in the aging process apparently also plays a role in the development of decreased muscle mass and strength (Mainous *et al.*, 2015). The decrease in muscle mass in the elderly corresponds to a 10% reduction in grip strength per decade, starting from the age of 40 years (Jaafar *et al.*, 2023). Skeletal muscle plays an important role in glucose metabolism and has a significant influence on insulin sensitivity. Increased

insulin resistance in older adults may be closely associated with skeletal muscle aging. A reduction in GLUT4 expression as a result of a decrease in muscle volume causes a reduction in insulin sensitivity in aging skeletal muscle (Joo *et al.*, 2022). The results of this study showed a correlation of waist circumference and hip circumference with handgrip strength. This is in line with the study of Widjaja *et al.* (Widjaja *et al.*, 2018), who stated that there was a significant correlation of waist circumference and hip circumference with handgrip strength ($p < 0.001$, $r = 0.302$; and $p < 0.001$, $r = 0.296$). A greater waist circumference is associated with a greater handgrip strength in males (Hardy *et al.*, 2013).

In the multiple regression analysis of HGS, we removed the data on body weight because of the occurrence of multicollinearity between weight and BMI, with a correlation coefficient of more than 0.9, indicating an extremely strong relationship between these two variables. Conceptually, only one of the variables is used in multivariate analysis. The multiple regression analysis included age, height, BMI, waist circumference, and hip circumference, with HGS as the dependent variable in a stepwise manner. After three iterations, only height, BMI, and hip circumference played a role in predicting HGS by means of the regression formula, which resulted in $R^2 = 0.19$. This signifies that for known values of height, BMI, and hip circumference, we can predict the HGS value 19% more accurately than without knowledge of these data. For a clinical diagnosis, a minimum increase of 10% is needed. This study found that height, BMI, and hip circumference play a role in predicting handgrip strength through the regression formula, i.e., $\text{handgrip strength} = -36.98 + 0.48 \times \text{height} + 0.79 \times \text{BMI} - 0.29 \times \text{hip circumference}$. Handgrip strength and waist circumference are surrogate muscle strength and visceral adiposity markers, respectively (Nakanishi *et al.*, 2024). Waist circumference is also more sensitive in determining body fat distribution, particularly in the abdomen. Waist circumference correlates better with abdominal fat distribution than BMI (Rokhmah *et al.*, 2015). Waist circumference and hip circumference are associated with obesity, in which the fat

content is higher, and the skeletal muscle and fascicular content is lower, therefore resulting in a lower contractile capacity of skeletal muscle (Afaq *et al.*, 2019). The limitation of this study is that it did not evaluate other factors that may affect handgrip strength, such as nutrient intake. From the anthropometric measures of the hands and feet in this study, the causes and effects of the decrease in handgrip strength could not be determined because this study used a cross-sectional design.

Conclusion

Three factors may be used to estimate handgrip strength in PWP, namely height, BMI, and hip circumference, whereas blood glucose concentration is not correlated with handgrip strength in PWP. It is recommended to pay attention to height, BMI, and hip circumference, particularly in the acceptance criteria of PWP candidates, and to conduct further studies with consideration of other factors that affect handgrip strength, such as food intake and anthropometric features of the hands and arms.

Acknowledgments

Financial support was provided by Universitas Trisakti for this research.

References

- Afaq, E., Hassan, S.H., Khan, M.R., Zaidi, D.A., & Baktashi, H., 2019. Correlation of Hand Dynamometry With Body Mass Index and Waist-Hip Ratio in Medical Students. *Pak J Physiol.*,15, pp.19-21.
- Agtuahene, M.A., Quartey, J., & Kwakye, S., 2023. Influence of Hand Dominance, Gender, and Body Mass Index on Hand Grip Strength. *S Afr J Physiother.*, 79, pp.1-6.
- Al-Asadi, J.N., 2018. Handgrip Strength in Medical Students: Correlation with Body Mass Index and Hand Dimensions. *Asian Journal of Medical Sciences*, 9, pp.21-26.
- Badan Pusat Statistik Kota Administrasi Jakarta Barat., 2020. *Kecamatan Cengkareng Dalam Angka*, pp.1-114.
- Chen, D., Zhu, Y., Ni, W., Li, Y., Yin, G., Shao, Z., & Zhu, J., 2023. Hand Grip Strength is Inversely Associated with Total Daily Insulin Dose Requirement in Patients with Type 2 Diabetes Mellitus: A Cross-Sectional Study. *PeerJ.*, pp.1-13.
- Cooper, R., Tomlinson, D., Hamer, M., & Pereira, S.M.P., 2022. Lifetime Body Mass Index and Grip Strength at Age 46 Years: The 1970 British Cohort Study. *J Cachexia Sarcopenia Muscle*, 13, pp.1-10.
- Fauzi, L., Handayani, O.W.K., Susilo, M.T., Kurnia, A.R., Rahayu, S.R., Irawan, F.A., Horng Lu, F.J., Lin, C., Lai, M.F., & Yu, Y.C., 2022. Obesity in Indonesian and Taiwanese Adolescents Related to Self Perception, Diet, Exercise, and Body Image. *KEMAS*, 17(3), pp.453-461.
- Gamayanti, K.A.V., Aryana, I.G.P.S., & Gotera, W., 2023. Hubungan Antara Kekuatan Genggaman Tangan Dengan Kadar Gula Darah Sewaktu Pada Lansia Di Desa Melinggih, Kecamatan Payangan, Kabupaten Gianyar. *Intisari Sains Medis*, 14(2), pp.483-488.
- Giglio, B.M., Mota, J.F., Wall, B.T., & Pimentel, G.D., 2018. Low Handgrip Strength is Not Associated With Type 2 Diabetes Mellitus and Hyperglycemia: A Population-Based Study. *Clin Nutr Res.*, 7(2), pp.112-116.
- Hammed, A.I., & Obaseki, C.O., 2017. Interdependence of Body Mass Index with Handgrip Strength and Endurance Among Apparently Healthy Teenagers. *Turk J Kin.*, 4, pp.1-7.
- Handayani, M.D.N., Sadewa, A.H., Farmawati, A., & Rochmah, W., 2018. Anthropometric Prediction Equations for Estimating Muscle Mass of Elderly Women. *KEMAS*, 14(2), pp.195-204.
- Hardy, R., Cooper, R., Sayer, A.A., Shlomo, Y.B., Cooper, C., Deary, I.J., Demakakos, P., Gallacher, J., Martin, R.M., McNeill, G., Starr, J.M., Steptoe, A., Syddall, H., & Kuh, D., 2013. Body Mass Index, Muscle Strength, and Physical Performance in Older Adults From Eight Cohort Studies: The Halcyon Programme. *PLoS One*, 8(2), pp.e56483.
- Heidy., Djuartina, T., & Irawan, R., 2019. Korelasi Kekuatan Genggaman Tangan Dengan Karakter Antropometri Lengan Bawah dan Tangan Serta Indeks Massa Tubuh. *Damianus Journal of Medicine*, 18, pp.1-7.
- Ilmi, A.F., & Utari, D.M., 2020. Hubungan Lingkar Pinggang dan Rasio Lingkar Pinggang-Panggul (RLPP) Terhadap Kadar Glukosa Darah Puasa Pada Mahasiswa. *Journal of Nutrition College*, 9, pp.222-227.
- Jaafar, M.H., Ismail, R., Ismail, N.H., Isa, Z.M., Tamil, A.M., Nasir, N.M., Keat, N.K., Razak, N.H., Abidin, N.Z., & Yusof, K.H., 2023. Normative Reference Values and Predicting Factors of Handgrip Strength For Dominant

- and Non-Dominant Hands Among Healthy Malay Adults in Malaysia. *BMC Musculoskelet Disord.*, 24, pp.1-9.
- Joo, K.C., Son, D.H., & Park, J.M., 2022. Association Between Relative Handgrip Strength and Insulin Resistance in Korean Elderly Men Without Diabetes: Findings Of The 2015 Korea National Health Nutrition Examination Survey. *Korean J Fam Med.*, 43, pp.199-205.
- Krakauer, N.Y., & Krakauer, J.C., 2020. Association of Body Shape Index (ABSI) with Hand Grip Strength. *Int J Environ Res Public Health*, 17, pp.1-12.
- Kurniawan, A., Widjaja, D., & Lugito, N.P.H., 2018. *Mean and Cutoff Value of Hand Grip Strength For Healthy Indonesian Young Adults*. World Congress on Osteoporosis, Osteoarthritis and Musculoskeletal Diseases; Krakow, Poland.
- Leong, D.P., Teo, K.K., Rangarajan, S., Kutty, R., Lanas, F., Hui, C., Quanyong, X., Zhenzhen, Q., Jinhua, T., Noorhassim, I., AlHabib, K.F., Moss, S.J., Rosengren, A., Akalin, A.A., Rahman, O., Chifamba, J., Orlandini, A., Kumar, R., Yeates, K., Gupta, R., Yusuf, A., Dans, A., Avezum, A., Lopez-Jaramillo, P., Poirier, P., Heidari, H., Zatonska, K., Iqbal, R., Khatib, R., & Yusuf, S., 2016. Reference Ranges of Handgrip Strength From 125,462 Healthy Adults in 21 Countries: A Prospective Urban Rural Epidemiologic (PURE) Study. *J Cachexia Sarcopenia Muscle*, 7, pp.535-546.
- Mainous, A.G., Tanner, R.J., Anton, S.D., & Jo, A., 2015. Grip Strength as A Marker of Hypertension and Diabetes in Healthy Weight Adults. *Am J Prev Med*, 49(6), pp.850-858.
- Manoharan, V.S., Sundaram, S.G., & Jason, J.I., 2015. Factors Affecting Hand Grip Strength and Its Evaluation: A Systemic Review. *Int J Physiother Res.*, 3(6), pp.1288-1293.
- McGrath, R., Johnson, N., Klawitter, L., Mahoney, S., Trautman, K., Carlson, C., Rockstad, E., & Hackney, K.J., 2020. What Are The Association Patterns Between Handgrip Strength and Adverse Health Conditions? A Topical Review. *SAGE Open Medicine*, 8, pp.1-12.
- Nakanishi, S., Shimoda, M., Kimura, T., Sanada, J., Fushimi, Y., Iwamoto, Y., Iwamoto, H., Dan, K., Mune, T., Kaku, K., & Kaneto, H., 2024. The Impact of Handgrip Strength and Waist Circumference on Glycemic Control: Prospective, Observational Study Using Outpatient Clinical Data in Japanese Patients With Type 2 Diabetes Mellitus. *J Diabetes Investig.*, 15(7), pp.892-898.
- Nathania, N.P.S., Nugraha, M.H.S., Primayanti, I.D.A., & Saraswati, P.A.S., 2023. Handgrip Strength is Associated With Balance in Elderly Women. *Majalah Ilmiah Fisioterapi Indonesia*, 11, pp.70-75.
- Gubernur Provinsi Daerah Khusus Ibukota Jakarta., 2017. *Peraturan Gubernur Provinsi Daerah Khusus Ibukota Jakarta Nomor 7 tahun 2017 Tentang Penanganan Prasarana dan Sarana Umum Tingkat Kelurahan*. Jakarta 2017.
- Richter, E.A., & Hargreaves, M., 2013. Exercise, GLUT4, and Skeletal Muscle Glucose Uptake. *Physiol Rev.*, 93(3), pp.993-1017.
- Rokhmah, F.D., Handayani, D., & Al-Rasyid, H., 2015. Korelasi Lingkar Pinggang dan Rasio Lingkar Pinggang-Panggul Terhadap Kadar Glukosa Plasma Menggunakan Tes Toleransi Glukosa Oral. *Jurnal Gizi Klinik Indonesia*, 12, pp.28-35.
- Rondhianto., Ridla, A.Z., & Hasan, H., 2024. Sociodemographic Factors Affecting Diabetic Dietary Behavior in People with Type 2 Diabetes Mellitus. *KEMAS*, 19(3), pp.429-437.
- Safitri, I., Setyarsih, L., Susanto, H., Suhartono., & Fitranti, D.Y., 2020. The Effect of Low and High Glycemic Load Diet on Muscle Fatigue of Young Soccer Athletes. *KEMAS*, 16(1), pp.111-120.
- Schrager, M.A., Metter, E.J., Simonsick, E., Ble, A., Bandinelli, S., Lauretani, F., & Ferrucci, L., 2007. Sarcopenic Obesity and Inflammation in The InChianti Study. *J Appl Physiol.*, 102(3), pp.919-925.
- Sirajudeen, M.S., Shah, U.N., Pillai, P.S., Mohasin, N., & Shantaram, M., 2012. Correlation Between Grip Strength and Physical Factors in Men. *International Journal of Health and Rehabilitation Sciences*, 1, pp.58-63.
- Soenyoto, T., Mollap, K., & Mungpong, J., 2017. Correlation of Energy and Protein Consumption Levels with Physical Endurance of Rhythmic Gymnast Athletes. *KEMAS*, 13(1), pp.131-136.
- Soraya, N., & Parwanto, E., 2023. The Controversial Relationship Between Body Mass Index and Handgrip Strength in The Elderly: An Overview. *Malays J Med Sci.*, 30, pp.73-83.
- Vaishya, R., Misra, A., Vaish, A., Ursino, N., & D'Ambrosi, R., 2024. Hand Grip Strength as A Proposed New Vital Sign Of Health: A Narrative Review of Evidences. *Journal of Health, Population and Nutrition*, 43, pp.1-

14.

- Widjaja, D., Tjhai, K., Dermawan, K., Kalim, J., Tanzil, S., Pramusinta, A., Phenca, H.K., Kurniawan, A., 2018. *Correlations Between Hand Grip Strength and Anthropometric Status in Young Indonesian Adults*. World Congress on Osteoporosis, Osteoarthritis and Musculoskeletal Diseases; Krakow, Poland.
- Zaccagni, L., Toselli, S., Bramanti, B., Gualdi-Russo, E., Mongillo, J., & Rinaldo, N., 2020. Handgrip Strength in Young Adults: Association With Anthropometric Variables and Laterality. *Int J Environ Res Public Health*, 17, pp.1-18.

Handgrip strength of public works personnel in West Jakarta

ORIGINALITY REPORT

18%

SIMILARITY INDEX

15%

INTERNET SOURCES

12%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1	pmc.ncbi.nlm.nih.gov Internet Source	1%
2	researchonline.jcu.edu.au Internet Source	1%
3	Sarfaraz Alam, Omna Chawla, Anant Sinha, Ankita Juyal, Supriya Singh. " Fasting blood sugar, handgrip strength and central obesity: Exploring the relationship ", National Journal of Physiology, Pharmacy and Pharmacology, 2024 Publication	1%
4	turk-jem.dergisi.org Internet Source	1%
5	www.kjfm.or.kr Internet Source	1%
6	www.researchgate.net Internet Source	1%
7	www.ncbi.nlm.nih.gov Internet Source	1%
8	Ni Putu Laksmi Martini, Dian Mediana. "The Risk Factors Associated with Handgrip Strength and Endurance in Adolescence", Jurnal Biomedika dan Kesehatan, 2023 Publication	1%

9	David Roseveare. "World Congress on Osteoporosis, Osteoarthritis and Musculoskeletal Diseases (WCO-IOF-ESCEO 2018): Poster Abstracts", Osteoporosis International, 2018 Publication	1 %
10	Submitted to University of Stellenbosch, South Africa Student Paper	1 %
11	wco-iof-esceo.org Internet Source	1 %
12	www.wjgnet.com Internet Source	1 %
13	Da-shuang Chen, Yun-qing Zhu, Wen-ji Ni, Yu-jiao Li, Guo-ping Yin, Zi-yue Shao, Jian Zhu. "Hand grip strength is inversely associated with total daily insulin dose requirement in patients with type 2 diabetes mellitus: a cross-sectional study", PeerJ, 2023 Publication	<1 %
14	Zhihao Wei, Tengfei Yu, Xufeng Jin, Guanyi Ma, Xianfeng Meng. "The association between body roundness index and handgrip strength and muscle quality index: A cross-sectional study", PLOS One, 2025 Publication	<1 %
15	theses.bham.ac.uk Internet Source	<1 %
16	www.mjms.usm.my Internet Source	<1 %
17	www.scielo.br Internet Source	<1 %

18	efsupit.ro Internet Source	<1 %
19	www.ptantropologiczne.pl Internet Source	<1 %
20	ejournal.medistra.ac.id Internet Source	<1 %
21	perpustakaan.poltekkes-malang.ac.id Internet Source	<1 %
22	ejournal3.undip.ac.id Internet Source	<1 %
23	www.riped-online.com Internet Source	<1 %
24	Mainous, Arch G., Rebecca J. Tanner, Stephen D. Anton, and Ara Jo. "Grip Strength as a Marker of Hypertension and Diabetes in Healthy Weight Adults", American Journal of Preventive Medicine, 2015. Publication	<1 %
25	huggingface.co Internet Source	<1 %
26	www.researchsquare.com Internet Source	<1 %
27	www.ssph-journal.org Internet Source	<1 %
28	Submitted to Coventry University Student Paper	<1 %
29	publisher.uthm.edu.my Internet Source	<1 %
30	Adriani Thahara, Ika Krisnawati. "Hubungan Rasio Lingkar Pinggang Panggul dan Asupan	<1 %

Lemak dengan Risiko Kejadian Kardiovaskular pada Usia Produktif", Jurnal Biomedika dan Kesehatan, 2023

Publication

31

Mumtaz Gowhar, Farhana Ahad, Javed Hussain Bhat. "Correlation of Hand Dominance and Body Mass Index on Maximal Isometric Handgrip Strength among Students of SKIMS Medical College, Srinagar, India", National Journal of Medical Research, 2025

Publication

<1 %

32

www.mdpi.com

Internet Source

<1 %

33

1library.net

Internet Source

<1 %

34

Chiao-Nan Chen, Ting-Chung Chen, Shiow-Chwen Tsai, Chii-Min Hwu. "Factors associated with relative muscle strength in patients with type 2 diabetes mellitus", Archives of Gerontology and Geriatrics, 2021

Publication

<1 %

Exclude quotes On

Exclude matches < 10 words

Exclude bibliography On



**KEMENTERIAN PENDIDIKAN TINGGI,
SAINS, DAN TEKNOLOGI
DIREKTORAT JENDERAL SAINS DAN TEKNOLOGI**
Jalan Jenderal Sudirman, Senayan, Jakarta 10270
Telepon (021) 57946104, Pusat Panggilan ULT DIKTISAINTEK 126
Laman www.kemdiktisaintek.go.id

Nomor : 244/DST/D.D1/HM.01.01/2026
Lampiran : Satu lampiran
Hal : Apresiasi Capaian Tata Kelola Jurnal Ilmiah

12 Juni 2026

Yth. Pengelola Jurnal Nasional Terakreditasi SINTA 1
(Daftar Terlampir)

Direktorat Jenderal Sains dan Teknologi menyampaikan apresiasi dan penghargaan yang setinggi-tingginya kepada seluruh pengelola jurnal atas capaian sebagai Jurnal Nasional Terakreditasi SINTA 1. Prestasi tersebut merupakan hasil kerja keras, komitmen, dan konsistensi dalam menjaga mutu publikasi ilmiah serta mendukung pengembangan ekosistem riset dan inovasi di Indonesia.

Kami berharap pengelolaan jurnal dapat terus ditingkatkan dan dikembangkan sehingga mampu mencapai dan mempertahankan reputasi sebagai Jurnal Internasional Bereputasi. Upaya tersebut dapat dilakukan melalui penguatan kualitas substansi artikel, peningkatan profesionalisme dalam tata kelola editorial, perluasan jejaring kerja sama internasional, serta pemenuhan standar publikasi ilmiah yang diakui secara global. Di samping itu, kami mengharapkan pengelolaan jurnal senantiasa mengedepankan integritas akademik, transparansi, akuntabilitas, etika publikasi, dan praktik-praktik terbaik dalam penerbitan ilmiah, sehingga jurnal dapat menjadi media diseminasi ilmu pengetahuan yang berkualitas, terpercaya, dan berdaya saing internasional.

Selain itu, kami informasikan bahwa Direktorat Jenderal Sains dan Teknologi akan menyelenggarakan Konvensi Nasional Tata Kelola dan Integritas Publikasi Ilmiah di Surabaya dalam rentang tanggal 7-10 Agustus 2026. Kegiatan ini diselenggarakan bersamaan dengan Konvensi Sains, Teknologi, dan Industri Indonesia (KSTI) serta Konferensi Puncak Pendidikan Tinggi Indonesia (KPPTI) sebagai wadah penguatan tata kelola, integritas, kualitas jurnal serta publikasi ilmiah di Indonesia. Kami berharap kegiatan tersebut dapat semakin mendorong peningkatan mutu dan reputasi jurnal nasional menuju standar internasional.

Demikian apresiasi dan harapan ini kami sampaikan. Semoga capaian yang telah diraih dapat menjadi motivasi untuk terus meningkatkan kualitas dan reputasi jurnal, baik di tingkat nasional maupun internasional.

Atas perhatian dan kerja sama yang baik, kami ucapkan terima kasih.

Direktur Jenderal,



Ahmad Najib Burhani
NIP 197604272005021001

Tembusan:
Sekretaris Direktorat Jenderal Sains, dan Teknologi

Lampiran Surat

Nomor : 244/DST/D.D1/HM.01.01/2026

Tanggal : 12 Juni 2026

DAFTAR JURNAL NASIONAL TERAKREDITASI SINTA 1

NO	NAMA JURNAL	INSTITUSI	TAUTAN JURNAL
1	Sinergi	Universitas Mercu Buana	https://publikasi.mercubuana.ac.id/index.php/sinergi
2	JEBIS (Jurnal Ekonomi dan Bisnis Islam)	Universitas Airlangga	https://e-journal.unair.ac.id/JEBIS
3	Forum Geografi	Universitas Muhammadiyah Surakarta	http://journals.ums.ac.id/index.php/fg/issue/archive
4	Al-Albab	IAIN Pontianak	http://jurnal.iainpontianak.or.id/index.php/alalbab/index
5	Otoritas : Jurnal Ilmu Pemerintahan	Universitas Muhammadiyah Makassar	http://journal.unismuh.ac.id/index.php/Otoritas/
6	Millah: Journal of Religious Studies	Universitas Islam Indonesia	http://journal.uii.ac.id/Millah
7	al-Balagh : Jurnal Dakwah dan Komunikasi	Institut Agama Islam Negeri Surakarta	https://ejournal.uinsaid.ac.id/index.php/al-balagh/
8	Jurnal Ilmiah Psikologi Terapan	Universitas Muhammadiyah Malang	http://ejournal.umm.ac.id/index.php/jipt/
9	Yuridika	Universitas Airlangga	https://e-journal.unair.ac.id/YDK
10	De Jure: Jurnal Hukum dan Syariah	Universitas Islam Negeri Maulana Malik Ibrahim	http://ejournal.uin-malang.ac.id/index.php/syariah
11	Shirkah: Journal of Economics and Business	Institut Agama Islam Negeri Surakarta	http://shirkah.or.id/new-ojs/index.php/home
12	Industria: Jurnal Teknologi dan Manajemen Agroindustri	Universitas Brawijaya	http://industria.ub.ac.id
13	AGRARIS: Journal of Agribusiness and Rural Development Research	Universitas Muhammadiyah Yogyakarta	https://agraris.umy.ac.id/index.php/agraris
14	Script Journal: Journal of Linguistic and English Teaching	Universitas Widya Gama Mahakam Samarinda	https://jurnal.fkip-uwgm.ac.id/index.php/Script/issue/archive
15	Jurnal Keperawatan Soedirman	Universitas Jenderal Soedirman	http://jos.unsoed.ac.id/index.php/jks/index
16	Psikohumaniora: Jurnal Penelitian Psikologi	Universitas Islam Negeri Walisongo	https://journal.walisongo.ac.id/index.php/Psikohumaniora/index

17	Journal of Biomedicine and Translational Research	Universitas Diponegoro	https://ejournal2.undip.ac.id/index.php/jbtr/index
18	Jurnal Hukum Novelty	Universitas Ahmad Dahlan	http://journal.uad.ac.id/index.php/Novelty/index
19	Jurnal Arbitrer	Universitas Andalas	http://arbitrer.fib.unand.ac.id
20	LLT Journal: A Journal on Language and Language Teaching	Universitas Sanata Dharma	http://e-journal.usd.ac.id/index.php/LLT/index
21	Al-Ahkam: Jurnal Ilmu Syariah dan Hukum	Universitas Islam Negeri Raden Mas Said Surakarta	https://ejournal.uinsaid.ac.id/index.php/al-ahkam/
22	Journal of Governance and Public Policy	Universitas Muhammadiyah Yogyakarta	https://journal.umy.ac.id/index.php/GPP/index
23	Jurnal Optimasi Sistem Industri	Universitas Andalas	http://josi.ft.unand.ac.id/index.php/josi/issue/view/21
24	Journal of Indonesian Islam	Universitas Islam Negeri Sunan Ampel	https://jiis.uinsa.ac.id/index.php/JIIs/index
25	Indonesian Journal of Chemistry	Universitas Gadjah Mada	https://journal.ugm.ac.id/index.php/ijc
26	Medical Journal of Indonesia	Universitas Indonesia	http://mji.ui.ac.id/journal/index.php/mji
27	TELKOMNIKA (Telecommunication Computing Electronics and Control)	Universitas Ahmad Dahlan	http://journal.uad.ac.id/index.php/telkomnika
28	Gadjah Mada International Journal of Business	Universitas Gadjah Mada	https://jurnal.ugm.ac.id/gamaijb/index
29	Microbiology Indonesia	Perhimpunan Mikrobiologi Indonesia	https://jurnal.permi.or.id/index.php/mionline
30	Journal of Mathematical and Fundamental Sciences	Institut Teknologi Bandung	https://journals.itb.ac.id/index.php/jmfs
31	Indonesian Journal of Islam and Muslim Societies	IAIN Salatiga	http://ijims.iainsalatiga.ac.id
32	Indonesian Journal of Geography	Universitas Gadjah Mada	http://jurnal.ugm.ac.id/ijg/
33	Jurnal Pendidikan IPA Indonesia (Indonesian Journal of Science Education)	Universitas Negeri Semarang	https://journal.unnes.ac.id/journals/jpii
34	Jurnal Manajemen Hutan Tropika	Institut Pertanian Bogor	http://journal.ipb.ac.id/index.php/jmht
35	TEFLIN Journal	Universitas Negeri Malang	http://www.journal.teflin.org

36	Paramita: Historical Studies Journal	Universitas Negeri Semarang	https://journal.unnes.ac.id/journals/paramita
37	Atom Indonesia	Badan Tenaga Nuklir Nasional	https://atomindonesia.brin.go.id/index.php/aij/index
38	BIOTROPIA	SEAMEO BIOTROP	http://journal.biotrop.org/index.php/biotropia
39	Tropical Animal Science Journal	Institut Pertanian Bogor	https://journal.ipb.ac.id/index.php/tasj
40	Journal of the Indonesian Tropical Animal Agriculture	Universitas Diponegoro	https://ejournal.undip.ac.id/index.php/jitaa/index
41	ACTA MEDICA INDONESIANA	Perhimpunan Dokter Spesialis Penyakit Dalam Indonesia	http://www.actamedindones.org/index.php/ijim
42	Journal of Engineering and Technological Sciences	Institut Teknologi Bandung	https://journals.itb.ac.id/index.php/jets/index
43	Bulletin of Electrical Engineering and Informatics	Universitas Ahmad Dahlan	http://journal.portalgaruda.org/index.php/EEI
44	AGRIVITA, Journal of Agricultural Science	Universitas Brawijaya	https://agrivita.ub.ac.id/index.php/agrivita
45	HAYATI Journal of Biosciences	Institut Pertanian Bogor	http://journal.ipb.ac.id/index.php/hayati/
46	Studia Islamika	Universitas Islam Negeri Syarif Hidayatullah	http://journal.uinjkt.ac.id/index.php/studia-islamika
47	Al-Jami'ah: Journal of Islamic Studies	Universitas Islam Negeri Sunan Kalijaga	http://www.aljamiah.or.id
48	Journal of ICT Research and Applications	Institut Teknologi Bandung	https://journals.itb.ac.id/index.php/jictra
49	Critical Care and Shock	Indonesian Society of Critical Care Medicine	http://criticalcareshock.org/
50	International Journal on Electrical Engineering and Informatics	Institut Teknologi Bandung	http://www.ijeei.org/
51	International Journal of Technology	Universitas Indonesia	http://ijtech.eng.ui.ac.id/archives

52	International Journal of Power Electronics and Drive Systems	Institute of Advanced Engineering and Science (IAES)	http://ijpeds.iaescore.com/index.php/IJPEDS
53	Bulletin of Chemical Reaction Engineering & Catalysis	Universitas Diponegoro	https://journal.bcrec.id/index.php/bcrec
54	Kukila	Indonesian Ornithology	http://kukila.org/index.php/KKL
55	International Journal of Electrical and Computer Engineering	Institute of Advanced Engineering and Science (IAES)	http://ijece.iaescore.com/index.php/IJECE
56	Biodiversitas : Journal of Biological Diversity	Universitas Sebelas Maret	http://biodiversitas.mipa.uns.ac.id/
57	International Journal on Advanced Science, Engineering and Information Technology (IJASEIT)	Indonesian Society for Knowledge and Human Development (Insight)	http://ijaseit.insightsociety.org/
58	Indonesian Journal of Electrical Engineering and Computer Science	Institute of Advanced Engineering and Science (IAES)	http://ijeecs.iaescore.com
59	Cakrawala Pendidikan	Universitas Negeri Yogyakarta	https://journal.uny.ac.id/index.php/cp
60	Civil Engineering Dimension	Universitas Kristen Petra	http://ced.petra.ac.id
61	Jurnal Hama dan Penyakit Tumbuhan Tropika	Universitas Lampung	http://jhpttropika.fp.unila.ac.id/index.php/jhpttropika
62	Ijtihad : Jurnal Wacana Hukum Islam dan Kemanusiaan	IAIN Salatiga	http://ijtihad.iainsalatiga.ac.id
63	Universa Medicina	Universitas Trisakti	https://univmed.org/ejurnal/index.php/medicina
64	Jurnal Ners	Universitas Airlangga	http://e-journal.unair.ac.id/index.php/JNERS/
65	Indonesian Journal of Biotechnology	Universitas Gadjah Mada	https://jurnal.ugm.ac.id/ijbiotech
66	Journal of Dentistry Indonesia	Universitas Indonesia	https://scholarhub.ui.ac.id/jdi/
67	Indonesia Law Review	Universitas Indonesia	https://scholarhub.ui.ac.id/ilrev

68	Jurnal Penelitian dan Evaluasi Pendidikan	Universitas Negeri Yogyakarta	https://journal.uny.ac.id/index.php/jpep/
69	MIQOT: Jurnal Ilmu-ilmu Keislaman	Universitas Islam Negeri Sumatera Utara	http://jurnalmiqotojs.uinsu.ac.id/index.php/jurnalmiqot
70	Jurnal Konstitusi	Mahkamah Konstitusi RI	https://jurnalkonstitusi.mkri.id/index.php/jk
71	Al-Manahij: Jurnal Kajian Hukum Islam	Universitas Islam Negeri Profesor Kiai Haji Saifuddin Zuhri Purwokerto	https://ejournal.uinsaizu.ac.id/index.php/almanahij/index
72	Jurnal Pendidikan Islam	Universitas Islam Negeri Sunan Gunung Djati	http://journal.uinsgd.ac.id/index.php/jpi
73	Al-'Adalah	Universitas Islam Negeri Raden Intan	http://ejournal.radenintan.ac.id/index.php/adalah/index
74	Teosofi: Jurnal Tasawuf dan Pemikiran Islam	Universitas Islam Negeri Sunan Ampel	http://jurnalfuf.uinsby.ac.id/index.php/teosofi
75	Jurnal Kesehatan Masyarakat	Universitas Negeri Semarang	https://journal.unnes.ac.id/journals/kemas
76	Media Kesehatan Masyarakat Indonesia	Universitas Hasanuddin	http://journal.unhas.ac.id/index.php/mkmi
77	ILMU KELAUTAN: Indonesian Journal of Marine Sciences	Universitas Diponegoro	http://ejournal.undip.ac.id/index.php/ijms/index
78	Jurnal Pengolahan Hasil Perikanan Indonesia	Institut Pertanian Bogor	http://journal.ipb.ac.id/index.php/jphpi
79	Jurnal Kajian Bali	Universitas Udayana	https://ejournal1.unud.ac.id/index.php/kajianbali
80	Al-Risalah : Forum Kajian Hukum dan Sosial Kemasyarakatan	Universitas Islam Negeri Sulthan Thaha Saifuddin Jambi	http://e-journal.lp2m.uinjambi.ac.id/ojs/index.php/AL-RISALAH
81	Jurnal Keperawatan Padjadjaran	Universitas Padjadjaran	http://jkp.fkep.unpad.ac.id/index.php/jkp
82	Journal of Mechatronics, Electrical Power, and Vehicular Technology	Lembaga Ilmu Pengetahuan Indonesia	https://mev.brin.go.id/mev
83	Harmonia: Journal of Arts Research and Education	Universitas Negeri Semarang	https://journal.unnes.ac.id/journals/harmonia

84	ADDIN	STAIN Kudus	http://journal.stainkudus.ac.id/index.php/Addin
85	Folia Medica Indonesiana	Universitas Airlangga	https://scholarly.unair.ac.id/fk-fmi/
86	Biosaintifika: Journal of Biology & Biology Education	Universitas Negeri Semarang	https://journal.unnes.ac.id/journals/biosaintifika
87	Molekul: Jurnal Ilmiah Kimia	Universitas Jenderal Soedirman	https://jos.unsoed.ac.id/index.php/jm/index
88	Journal of Regional and City Planning	Institut Teknologi Bandung	http://journals.itb.ac.id/index.php/jpww
89	Buletin Peternakan	Universitas Gadjah Mada	http://buletinpeternakan.fapet.ugm.ac.id
90	Jurnal Pendidikan Vokasi	Universitas Negeri Yogyakarta	http://journal.uny.ac.id/index.php/jpv
91	Economic Journal of Emerging Markets	Universitas Islam Indonesia	http://journal.uui.ac.id/index.php/JEP
92	Paediatrica Indonesiana	Ikatan Dokter Anak Indonesia	https://paediatricaindonesiana.org/
93	Jurnal Ilmu Sosial dan Ilmu Politik	Universitas Gadjah Mada	http://jurnal.ugm.ac.id/jsp
94	Journal of Indonesian Economy and Business	Universitas Gadjah Mada	https://journal.ugm.ac.id/v3/jieb/index
95	ESENSIA: Jurnal Ilmu-Ilmu Ushuluddin	Universitas Islam Negeri Sunan Kalijaga	https://ejournal.uin-suka.ac.id/ushuluddin/esensia/index
96	Ahkam: Jurnal Ilmu Syariah	Universitas Islam Negeri Syarif Hidayatullah	http://journal.uinjkt.ac.id/index.php/ahkam
97	Media Hukum	Universitas Muhammadiyah Yogyakarta	http://journal.umy.ac.id/index.php/jmh/index
98	k@ta	Universitas Kristen Petra	https://kata.petra.ac.id/
98	Jurnal Akuntansi dan Keuangan	Universitas Kristen Petra	https://jurnalakuntansi.petra.ac.id

99	Electronic Journal of Graph Theory and Applications	Indonesian Combinatorial Society (InaCombS), Graph Theory and Applications (GTA) Research	http://www.ejgta.org/index.php/ejgta/index
100	IJOG : Indonesian Journal on Geoscience	Kementerian Energi dan Sumber Daya Mineral	http://ijog.bgl.esdm.go.id
101	Journal of Islamic Architecture	Universitas Islam Negeri Maulana Malik Ibrahim	http://ejournal.uin-malang.ac.id/index.php/JIA
102	Jurnal Penelitian Kehutanan Wallacea	Kementerian Lingkungan Hidup dan Kehutanan	https://journal.unhas.ac.id/index.php/wallacea
103	Jurnal Pengelolaan Sumberdaya Alam dan Lingkungan (Journal of Natural Resources and Environmental Management)	Institut Pertanian Bogor	https://journal.ipb.ac.id/index.php/jpsl
104	Journal of Tropical Life Science : International Journal of Theoretical, Experimental, and Applied Life Sciences	Universitas Brawijaya	http://jtrolis.ub.ac.id/index.php/jtrolis/index
105	Nurse Media Journal of Nursing	Universitas Diponegoro	https://ejournal.undip.ac.id/index.php/medianers/index
106	CommIT (Communication and Information Technology) Journal	Universitas Bina Nusantara	https://journal.binus.ac.id/index.php/commit/index
107	Wacana: Journal of the Humanities of Indonesia	Universitas Indonesia	https://scholarhub.ui.ac.id/wacana/
108	International Journal of Renewable Energy Development : JRED	Universitas Diponegoro	https://ijred.cbiorc.id/index.php/ijred/index
109	Journal of Education and Learning (EduLearn)	Universitas Ahmad Dahlan	http://edulearn.intelektual.org
110	Journal of Degraded and Mining Lands Management	Universitas Brawijaya	https://jdmlm.ub.ac.id/
111	Jurnal Elektronika dan Telekomunikasi	Lembaga Ilmu Pengetahuan Indonesia	http://www.jurnalet.com
112	Brawijaya Law Journal : Journal of Legal Studies	Universitas Brawijaya	http://lawjournal.ub.ac.id/index.php/law/index

113	Geoplanning: Journal of Geomatics and Planning	Universitas Diponegoro	http://ejournal.undip.ac.id/index.php/geoplanning/
114	Hasanuddin Law Review	Universitas Hasanuddin	http://pasca.unhas.ac.id/ojs/index.php/halrev
115	The Indonesian Biomedical Journal	Prodia Education and Research Institute	http://inabj.org/
116	Kesmas: Jurnal Kesehatan Masyarakat Nasional (National Public Health Journal)	Universitas Indonesia	https://journal.fkm.ui.ac.id/kesmas/issue/archive
117	Jurnal Ilmiah Islam Futura	Universitas Islam Negeri Ar-Raniry	https://jurnal.ar-raniry.ac.id/index.php/islamfutura
118	Etikonomi	Universitas Islam Negeri Syarif Hidayatullah	http://journal.uinjkt.ac.id/index.php/etikonomi
119	Mazahib Jurnal Pemikiran Hukum Islam	UIN Sultan Aji Muhammad Idris Samarinda	https://journal.uinsi.ac.id/index.php/mazahib
120	ULUL ALBAB Jurnal Studi Islam	Universitas Islam Negeri Maulana Malik Ibrahim	https://ejournal.uin-malang.ac.id/index.php/ululalbab/index
121	Indonesian Journal of International Law	Universitas Indonesia	https://scholarhub.ui.ac.id/ijil
122	Yustisia	Universitas Sebelas Maret	https://jurnal.uns.ac.id/yustisia/index
123	Indonesian Journal of Pharmacy	Universitas Gadjah Mada	http://indonesianjpharm.farmasi.ugm.ac.id/index.php/3
124	Jurnal Keperawatan Indonesia	Universitas Indonesia	http://jki.ui.ac.id/index.php/jki
125	VALENSI	Universitas Islam Negeri Syarif Hidayatullah	http://journal.uinjkt.ac.id/index.php/valensi
126	International Journal of Advances in Intelligent Informatics	Universitas Ahmad Dahlan	http://ijain.org/index.php/IJAIN/index
127	Jurnal Ilmiah Peuradeun	SCAD Independent	https://journal.scadindependent.org/index.php/jipeuradeun
128	SAINS TANAH - Journal of Soil Science and Agroclimatology	Universitas Sebelas Maret	https://jurnal.uns.ac.id/tanah

129	Indonesian Journal of Electrical Engineering and Informatics (IJEI)	Institute of Advanced Engineering and Science (IAES)	http://section.iaesonline.com/index.php/IJEI/index
130	Indonesian Journal of Applied Linguistics (IJAL)	Universitas Pendidikan Indonesia	http://ejournal.upi.edu/index.php/IJAL
131	Buletin Ekonomi Moneter dan Perbankan	Bank Indonesia	https://www.bmebi.org/index.php/BEMP/
132	International Journal on Emerging Mathematics Education	Universitas Ahmad Dahlan	http://journal.uad.ac.id/index.php/IJEME
133	JURNAL INFOTEL	Institut Teknologi Telkom Purwokerto	https://ejournal.itelkom-pwt.ac.id/index.php/infotel
134	Jurnal Sylva Lestari	Universitas Lampung	http://jurnal.fp.unila.ac.id/index.php/JHT
135	Al-Jabar: Jurnal Pendidikan Matematika	Universitas Islam Negeri Raden Intan	http://ejournal.radenintan.ac.id/index.php/al-jabar/issue/archive
136	Forest and Society	Universitas Hasanuddin	http://journal.unhas.ac.id/index.php/fs/index
137	Journal of the Civil Engineering Forum	Universitas Gadjah Mada	https://jurnal.ugm.ac.id/v3/jcef/
138	JURNAL IUS KAJIAN HUKUM DAN Keadilan	Universitas Mataram	http://jurnalius.ac.id/ojs/index.php/jurnalIUS
139	Public Health of Indonesia	Ikatan Ahli Kesehatan Masyarakat (IAKMI) Pengurus Daerah Sulawesi Tenggara	http://stikbar.org/ycabpublisher/index.php/PHI/index
140	JOIN (Jurnal Online Informatika)	Universitas Islam Negeri Sunan Gunung Djati	https://join.if.uinsgd.ac.id/index.php/join
141	QIJIS (Qudus International Journal Of Islamic Studies)	STAIN Kudus	http://journal.stainkudus.ac.id/index.php/QIJIS
142	Habitat	Universitas Brawijaya	https://habitat.ub.ac.id/index.php/habitat
143	Jurnal Civics: Media Kajian Kewarganegaraan	Universitas Negeri Yogyakarta	https://journal.uny.ac.id/index.php/civics/index
144	Al-Ihkam: Jurnal Hukum dan Pranata Sosial	Institut Agama Islam Negeri Madura	http://ejournal.iainmadura.ac.id/index.php/alihkam/index

145	ASEAN Journal on Science and Tecnology for Development	Universitas Gadjah Mada	http://ajstd.org/index.php/ajstd/index
146	Bali Journal of Anesthesiology	Universitas Udayana	http://www.bjoaonline.com/
147	Belitung Nursing Journal	Belitung Raya Foundation	http://belitungraya.org/BRP/index.php/bnj
148	INDONESIAN JOURNAL OF APPLIED PHYSICS	Universitas Sebelas Maret	https://jurnal.uns.ac.id/ijap
149	Jurnal Infinity	Institut Keguruan dan Ilmu Pendidikan Siliwangi	http://e-journal.stkipsiliwangi.ac.id/index.php/infinity
150	Jurnal Pendidikan Matematika	Universitas Sriwijaya	https://jpm.ejournal.unsri.ac.id/index.php/jpm
151	Studies in English Language and Education	Universitas Syiah Kuala	https://jurnal.usk.ac.id/SiELE
152	Jurnal Gramatika	Universitas PGRI Sumatera Barat	https://ejournal.upgrisba.ac.id/index.php/jurnal-gramatika/index
153	Sriwijaya Law Review	Universitas Sriwijaya	http://journal.fh.unsri.ac.id/index.php/sriwijayalawreview/issue/archive
154	Register: Jurnal Ilmiah Teknologi Sistem Informasi	Universitas Pesantren Tinggi Darul Ulum	http://www.journal.unipdu.ac.id/index.php/register
155	LAW REFORM	Universitas Diponegoro	https://ejournal.undip.ac.id/index.php/lawreform
156	Journal on Mathematics Education	Universitas Sriwijaya	https://jme.ejournal.unsri.ac.id/index.php/jme
157	JAS (Journal of ASEAN Studies)	Universitas Bina Nusantara	https://journal.binus.ac.id/index.php/jas/index
158	JURNAL ILMIAH PENDIDIKAN FISIKA AL BIRUNI	Universitas Islam Negeri Raden Intan	http://ejournal.radenintan.ac.id/index.php/al-biruni/index
159	Jurnal Kedokteran dan Kesehatan Indonesia	Universitas Islam Indonesia	https://journal.uii.ac.id/JKKI
160	Bali Medical Journal	Universitas Udayana	http://www.balimedicaljournal.id

161	Udayana Journal of Law and Culture	Universitas Udayana	https://ejournal4.unud.ac.id/index.php/ujlc/en
162	Indonesian Aquaculture Journal	Kementerian Kelautan dan Perikanan	http://ejournal-balitbang.kkp.go.id/index.php/iaj/index
163	Indonesian Journal of Agricultural Science	Kementerian Pertanian	http://ejurnal.litbang.pertanian.go.id/index.php/ijas
164	Jurnal Ilmu Ternak dan Veteriner	Kementerian Pertanian	https://medpub.appertani.org/index.php/jitv
165	Scientific Contributions Oil and Gas	Kementerian Energi dan Sumber Daya Mineral	https://journal.lemigas.esdm.go.id/index.php/SCOG
166	Operations and Supply Chain Management: An International Journal	Institut Teknologi Sepuluh Nopember	http://journal.oscm-forum.org/
167	Record and Library Journal	Universitas Airlangga	https://e-journal.unair.ac.id/RLJ/index
168	Squalen Bulletin Of Marine And Fisheries Postharvest And Biotechnology	Kementerian Kelautan dan Perikanan	https://www.bbp4b.litbang.kkp.go.id/squalen-bulletin/index.php/squalen/index
169	International Journal of Evaluation and Research in Education (IJERE)	Institute of Advanced Engineering and Science (IAES)	http://ijere.iaescore.com/index.php/IJERE
170	Indonesian Journal of Cancer Chemoprevention	Indonesian Society for Cancer Chemoprevention	https://ijcc.chemoprev.org/index.php/ijcc/
171	Formatif: Jurnal Ilmiah Pendidikan MIPA	Universitas Indraprasta PGRI	http://journal.lppmunindra.ac.id/index.php/Formatif/user
172	Communications in Science and Technology	Komunitas Ilmuwan dan Profesional Muslim Indonesia (KIPMI) Yogyakarta	https://cst.kipmi.or.id
173	Caraka Tani: Journal of Sustainable Agriculture	Universitas Sebelas Maret	http://jurnal.uns.ac.id/carakata ni/
174	BISMA (Bisnis dan Manajemen)	Universitas Negeri Surabaya	https://journal.unesa.ac.id/index.php/bisma
175	Jurnal Administrasi Kesehatan Indonesia	Universitas Airlangga	https://e-journal.unair.ac.id/JAKI/
176	Jurnal Pendidikan Agama Islam	Universitas Islam Negeri Sunan Kalijaga	http://ejournal.uin-suka.ac.id/tarbiyah/index.php/jpai/index

177	Journal of the Indonesian Mathematical Society	Indonesian Mathematical Society (IndoMS)	https://jims-a.org
178	Jurnal Geografi	Universitas Negeri Medan	http://jurnal.unimed.ac.id/2012/index.php/geo/editor
179	Indonesian Journal of Science and Technology	Universitas Pendidikan Indonesia	http://ejournal.upi.edu/index.php/ijost/
180	ARTEKS : Jurnal Teknik Arsitektur	Universitas Katolik Widya Mandira Kupang	https://journal.unwira.ac.id/index.php/ARTEKS/index
181	IJoLE: International Journal of Language Education	Universitas Negeri Makassar	http://ojs.unm.ac.id/ijole
182	The Indonesian Journal Of Occupational Safety and Health	Universitas Airlangga	https://e-journal.unair.ac.id/IJOSH/issue/archive
183	Amerta Nutrition	Universitas Airlangga	https://e-journal.unair.ac.id/AMNT
184	Journal of Southeast Asian Human Rights	Universitas Jember	https://jurnal.unej.ac.id/index.php/JSEHR
185	Jurnal Ilmiah Perikanan dan Kelautan (JIPK)	Universitas Airlangga	https://e-journal.unair.ac.id/JIPK/issue/archive
186	Jurnal Biometrika dan Kependudukan	Universitas Airlangga	https://e-journal.unair.ac.id/GBK/issue/archive
187	Jurnal Ilmiah Al-Syir'ah	IAIN Manado	http://journal.iain-manado.ac.id/index.php/JIS/index
188	JURNAL KESEHATAN LINGKUNGAN (Journal of Environmental Health)	Universitas Airlangga	https://e-journal.unair.ac.id/JKL/index
189	Jurnal Nasional Teknik Elektro	Universitas Andalas	https://jnte.ft.unand.ac.id/index.php/jnte
190	Lentera Hukum	Universitas Jember	https://jurnal.unej.ac.id/index.php/eJLH/issue/archive
98	The Indonesian Journal of Public Health	Universitas Airlangga	https://e-journal.unair.ac.id/IJPH/index
99	Journal of Indonesian Legal Studies	Universitas Negeri Semarang	https://journal.unnes.ac.id/journals/jils

100	JURIS (Jurnal Ilmiah Syariah)	Institut Agama Islam Negeri Batusangkar	https://ejournal.uinmybatusangkar.ac.id/ojs/index.php/Juris
101	pedagogik : jurnal pendidikan	Universitas Nurul Jadid Paiton Probolinggo Jawa Timur	https://ejournal.unuja.ac.id/index.php/pedagogik/index
102	Syariah Jurnal Hukum dan Pemikiran	UIN Antasari Banjarmasin	http://jurnal.uin-antasari.ac.id/index.php/syariah
103	Jurnal Ilmiah Ilmu Terapan Universitas Jambi	Universitas Jambi	https://online-journal.unja.ac.id/index.php/JIITUJ/issue/archive
104	Asy-Syari'ah	Universitas Islam Negeri Sunan Gunung Djati	http://journal.uinsgd.ac.id/index.php/asy-syariah/index
105	JOIV : International Journal on Informatics Visualization	Politeknik Negeri Padang	http://joiv.org/index.php/joiv
106	Indonesian Journal of Urban and Environmental Technology	Universitas Trisakti	https://e-journal.trisakti.ac.id/index.php/urbanenvirotech
107	Islamic Guidance and Counseling Journal	Universitas Ma'arif Lampung	https://journal.iaimnumetrolampung.ac.id/index.php/igcj
108	Legality : Jurnal Ilmiah Hukum	Universitas Muhammadiyah Malang	http://ejournal.umm.ac.id/index.php/legality
109	Tasfiah: Jurnal Pemikiran Islam	Universitas Darussalam Gontor	https://ejournal.unida.gontor.ac.id/index.php/tasfiah
110	Jurnal Medik Veteriner	Universitas Airlangga	https://e-journal.unair.ac.id/JMV/issue/archive
111	SAMARAH: Jurnal Hukum Keluarga dan Hukum Islam	Universitas Islam Negeri Ar-Raniry Banda Aceh	https://jurnal.ar-raniry.ac.id/index.php/samarah/index
112	Nazhruna: Jurnal Pendidikan Islam	Universitas KH. Abdul Chalim	http://e-journal.ikhac.ac.id/index.php/NAZHRUNA
113	Lex Scientia Law Review	Universitas Negeri Semarang	https://journal.unnes.ac.id/journals/lsr
114	El-Mashlahah	Universitas Islam Negeri Palangka Raya	http://e-journal.iain-palangkaraya.ac.id/index.php/maslahah/index
115	Al-Istinbath : Jurnal Hukum Islam	Institut Agama Islam Negeri Curup	http://journal.iaincurup.ac.id/index.php/alistinbath

116	EduReligia : Jurnal Pendidikan Agama Islam	Universitas Nurul Jadid	https://ejournal.unuja.ac.id/index.php/edureligia/index
117	Interiority	Universitas Indonesia	https://interiority.eng.ui.ac.id/index.php/journal
118	Journal of Local Government Issues (Logos)	Universitas Muhammadiyah Malang	http://ejournal.umm.ac.id/index.php/LOGOS/index
119	JP3I (Jurnal Pengukuran Psikologi dan Pendidikan Indonesia)	Universitas Islam Negeri Syarif Hidayatullah Jakarta	http://journal.uinjkt.ac.id/index.php/jp3i
120	Ulul Albab: Jurnal Studi dan Penelitian Hukum Islam	Universitas Islam Sultan Agung	https://jurnal.unissula.ac.id/index.php/ua/index
121	Automotive Experiences	Universitas Muhammadiyah Magelang	http://journal.unimma.ac.id/index.php/AutomotiveExperiences
122	Jurnal Studi Ilmu-ilmu Al-Qur'an dan Hadis	Universitas Islam Negeri Sunan Kalijaga	http://ejournal.uin-suka.ac.id/ushuluddin/alquran
123	Media Publikasi Promosi Kesehatan Indonesia (MPPKI)	Universitas Muhammadiyah Palu	http://jurnal.unismuhpalu.ac.id/index.php/MPPKI
124	Izdihar : Journal of Arabic Language Teaching, Linguistics, and Literature	Universitas Muhammadiyah Malang	http://ejournal.umm.ac.id/index.php/izdihar/index
125	Borneo Journal of Pharmacy	Universitas Muhammadiyah Palangkaraya	https://journal.umpr.ac.id/index.php/bjop
126	Canrea Journal: Food Technology, Nutritions, and Culinary Journal	Universitas Hasanuddin	http://agritech.unhas.ac.id/ojs/index.php/canrea/index
127	Science and Technology Indonesia	Universitas Sriwijaya	http://sciencetechindonesia.com/index.php/jsti
128	Journal of Islamic Monetary Economics and Finance	Bank Indonesia	http://www.jimf-bi.org/index.php/JIMF
129	Communication in Biomathematical Sciences	Institut Teknologi Bandung	http://journals.itb.ac.id/index.php/cbms
130	Volksggeist: Jurnal Ilmu Hukum dan Konstitusi	Institut Agama Islam Negeri Purwokerto	http://ejournal.uinsaizu.ac.id/index.php/volksggeist/issue/archive
131	CHEESA: Chemical Engineering Research Articles	Universitas PGRI Madiun	http://e-journal.unipma.ac.id/index.php/cheesa

132	IAES International Journal of Artificial Intelligence	Institute of Advanced Engineering and Science (IAES)	http://ijai.iaescore.com/index.php/IJAI
133	Jambe Law Journal	Universitas Jambi	http://jlj.unja.ac.id/index.php/home/index
134	Jambura Law Review	Universitas Negeri Gorontalo	http://ejurnal.ung.ac.id/index.php/jalrev
135	Bestuur	Universitas Sebelas Maret	https://jurnal.uns.ac.id/bestuur/index
136	Indonesian Journal of Halal Research	Universitas Islam Negeri Sunan Gunung Djati Bandung	https://journal.uinsgd.ac.id/index.php/ijhar/index
137	International Journal of Education and Learning	Association for Scientific Computing Electronics and Engineering (ASCEE)	http://pubs2.ascee.org/index.php/ijele
138	Indonesian Journal Of Medical Laboratory Science And Technology	Universitas Nahdlatul Ulama Surabaya	https://journal2.unusa.ac.id/index.php/IJMLST
139	Indonesian Food Science and Technology Journal	Universitas Jambi	https://online-journal.unja.ac.id/iftj
140	International Journal of Islamic Khazanah	Universitas Islam Negeri Sunan Gunung Djati Bandung	https://khazanah.uinsgd.ac.id/index.php/ijik
141	El-Usrah : Jurnal Hukum Keluarga	Universitas Islam Negeri Ar-Raniry Banda Aceh	https://jurnal.ar-raniry.ac.id/index.php/usrah
142	Teknomekanik	Universitas Negeri Padang	http://teknomekanik.ppj.unp.ac.id/index.php/teknomekanik
143	Jurnal Hukum	Universitas Islam Sultan Agung	http://jurnal.unissula.ac.id/index.php/jurnalhukum
144	Journal of Innovation in Educational and Cultural Research	Yayasan Keluarga Guru Mandiri	http://jiecr.org/index.php/jiecr
145	Journal of Tropical Biodiversity and Biotechnology	Universitas Gadjah Mada	https://journal.ugm.ac.id/jtbb
146	APTISI Transactions on Technopreneurship	APTISI	https://att.aptisi.or.id/index.php/att
147	International Journal of Reconfigurable and Embedded Systems (IJRES)	Institute of Advanced Engineering and Science	http://ijres.iaescore.com/index.php/IJRES

148	Journal of Law and Legal Reform	Universitas Negeri Semarang	https://journal.unnes.ac.id/journals/jlrr
149	Khazanah Hukum	Universitas Islam Negeri Sunan Gunung Djati Bandung	https://khazanah.uinsgd.ac.id/index.php/kh
150	International Journal of Advances in Applied Sciences	Institute of Advanced Engineering and Science	http://ijaas.iaescore.com/index.php/IJAAS
151	International Journal of Applied Power Engineering (IJAPE)	Institute of Advanced Engineering and Science	http://ijape.iaescore.com/index.php/IJAPE
152	Journal of Applied Engineering and Technological Science (JAETS)	Yayasan Pendidikan Riset dan Pengembangan Intelektual	https://journal.yrpiiku.com/index.php/jaets
153	Journal of Robotics and Control (JRC)	Universitas Muhammadiyah Yogyakarta	https://journal.umy.ac.id/index.php/jrc/index
154	TARBIYA: Journal of Education in Muslim Society	Universitas Islam Negeri Syarif Hidayatullah	http://journal.uinjkt.ac.id/index.php/tarbiya
155	Ulumuna	Universitas Islam Negeri Mataram	https://ulumuna.or.id/index.php/ujis
156	Asean Journal of Science and Engineering (AJSE)	Universitas Pendidikan Indonesia	https://ejournal.upi.edu/index.php/AJSE
157	Jambura Journal of Biomathematics (JJBm)	Universitas Negeri Gorontalo	https://jjbm.fmipa.ung.ac.id/index.php/ejournal/index
158	Advance Sustainable Science, Engineering and Technology (ASSET)	Universitas PGRI Semarang	http://journal.upgris.ac.id/index.php/asset
159	Indonesian Journal of Geography	Universitas Gadjah Mada	https://jurnal.ugm.ac.id/ijg
160	Journal of Islamic Law (JIL)	Institut Agama Islam Negeri Pontianak	http://e-journal.iainptk.ac.id/index.php/jil/index
161	Jurisdictie : Jurnal Hukum dan Syariah	Universitas Islam Negeri Maulana Malik Ibrahim Malang	http://ejournal.uin-malang.ac.id/index.php/jurisdictie
162	Kanz Philosophia: A Journal for Islamic Philosophy and Mysticism	Sekolah Tinggi Agama Islam Sadra	https://journal.sadra.ac.id/ojs/index.php/kanz
163	PETITA: Jurnal Kajian Ilmu Hukum dan Syariah	Universitas Islam Negeri Ar-Raniry Banda Aceh	http://petita.ar-raniry.ac.id/index.php/petita

164	Mechanical Engineering for Society and Industry	Universitas Muhammadiyah Magelang	https://journal.unimma.ac.id/index.php/mesi
165	JHCLS	Lembaga Contrarius Institute	https://www.jhcls.org/index.php/JHCLS
166	Ascarya: Journal of Islamic Science, Culture and Social Studies	<u>Perkumpulan Alumni Dan Santri Mahyajatul Qurro</u>	http://journal.ascarya.or.id/index.php/iscs
167	Journal of Public Health and Pharmacy	Universitas Muhammadiyah Palu	https://jurnal.unismuhpalu.ac.id/index.php/jphp/index
168	Heart Science Journal	Universitas Brawijaya	https://heartscience.ub.ac.id/index.php/heartscience
169	The Indonesian Journal of Socio-Legal Studies	Universitas Indonesia	https://scholarhub.ui.ac.id/ijsls
170	Online Learning in Educational Research	Foundation Of Advanced Education	https://journal.foundae.com/index.php/oler
171	MILRev: Metro Islamic Law Review	Universitas Islam Negeri Jurai Siwo Lampung	https://e-journal.metrouniv.ac.id/index.php/milrev
172	Journal of Coaching and Sports Science	Cv. Foundae	https://journal.foundae.com/index.php/jcss
173	International Journal of Electronics and Communications Systems	Universitas Islam Negeri Raden Intan Lampung	http://ejournal.radenintan.ac.id/index.php/IJECS/index
174	Narra J	Universitas Syiah Kuala	https://narraj.org/main

BUKTI KORESPONDENSI
ARTIKEL JURNAL INTERNASIONAL TERINDEKS SCOPUS Q4

Judul artikel : " **Handgrip Strength of Public Works Personnel in West Jakarta**"

Jurnal : Jurnal Kesehatan Masyarakat, volume 21, no. 2, (2025), Oktober: 360– 368.

Penulis : Alvina,Pusparini, Mario, Yasmine Mashabi.

No	Perihal	Tanggal
1	Bukti konfirmasi submit artikel dan artikel yang disubmit	29 Mei 2025
2	Bukti konfirmasi review dan hasil review pertama	22 Juni 2025
3	Bukti konfirmasi submit revisi pertama, respon kepada reviewer, dan artikel yang diresubmit	22 Juni 2025
4	Bukti konfirmasi review dan hasil review kedua	21 Juli 2025
5	Bukti konfirmasi submit revisi kedua, respon kepada reviewer, dan artikel yang diresubmit	21 Juli 2025
6	Bukti Accepted artikel	1 Agustus 2025
7	Bukti konfirmasi proofreading dan hasil proofreading	23 September 2025
8	Bukti konfirmasi perbaikan proofreading dan hasil perbaikan proofreading	25 September 2025
9	Bukti konfirmasi publikasi artikel dan artikel yang dipublikasi	6 Oktober 2025

1. Bukti konfirmasi submit artikel dan
artikel yang disubmit
(29 Mei 2025)



Alvina fk <dr.alvina@trisakti.ac.id>

Thank you for your submission to Jurnal Kesehatan Masyarakat

1 message

Prof. Dr. dr. Oktia Woro Kasmini Handayani, M.Kes <kemas@mail.unnes.ac.id>
To: Alvina Alvina <dr.alvina@trisakti.ac.id>

Thu, May 29, 2025 at 5:05 PM

Dear Alvina Alvina,

Thank you for your submission to Jurnal Kesehatan Masyarakat. We have received your submission, Factors Associated With Handgrip Strength of Public Works Personnel in West Jakarta, and a member of our editorial team will see it soon. You will be sent an email when an initial decision is made, and we may contact you for further information.

You can view your submission and track its progress through the editorial process at the following location:

Submission URL: <https://journal.unnes.ac.id/journals/kemas/authorDashboard/submission/25917>

If you have been logged out, you can login again with the username alvina.

If you have any questions, please contact me from your [submission dashboard](#).

Thank you for considering Jurnal Kesehatan Masyarakat as a venue for your work.

—

This is an automated message from [Jurnal Kesehatan Masyarakat](#).

Factors Associated With Handgrip Strength of Public Works Personnel in West Jakarta

Alvina^{1,a)}, Pusparini^{2,b)}, Mario^{3,c)}, Yasmine Mashabi^{4,d)}

^{1,2,3,4} Department of Clinical Pathology, Faculty of Medicine, Universitas Trisakti

^{a)}Corresponding author: dr.alvina@trisakti.ac.id

^{b)}pusparini@trisakti.ac.id

^{c)}mario@trisakti.ac.id

^{d)} yasminedrp@gmail.com

Abstract

Public Works Personnel (PWP) in Jakarta perform their routine duties manually instead of using equipment. Handgrip strength (HGS) measures maximum hand strength as a quality indicator for muscle strength and mass. Factors affecting muscle strength are age, sex, body mass index (BMI), waist circumference (WC), and hip circumference (HC). Increased fasting blood glucose (FBG) is also associated with muscle quality, muscle strength, and physical performance. This study aimed to determine the factors associated with handgrip strength in PWP. A total of 192 male PWP from the Cengkareng district were recruited by simple random sampling. Collected data were demographics, BMI, FBG, WC, HC, and HGS. Statistical analysis was done using Pearson's correlation test and multiple regression analysis. There was a significant positive correlation of BMI, WC, and HC with HGS ($p=0.006$; $r=0.20$, $p=0.009$; $r=0.19$ and $p=0.005$; $r=0.20$) and a significant negative correlation between age and HGS ($p=0.008$; $r=-0.19$) but not between FBG and HGS ($p=0.847$). Multiple regression analysis showed height, BMI, and HC positively predicting HGS ($R^2=0.19$). Handgrip strength of public works personnel is associated with BMI, age, waist circumference, and hip circumference, but not with fasting blood glucose. Height, BMI, and hip circumference predict handgrip strength.

Keywords: Handgrip strength, Body mass index, Hip circumference, Waist circumference, Fasting blood glucose

INTRODUCTION

Personnel for the management of public infrastructure and facilities at the village level are workers who manage village public works for a given time period on the basis of a work order. The tasks of village-level public works personnel (PWP) consist of managing the infrastructure and facilities, such as streets, drains, and parks, including installed lights (Peraturan Gubernur DKI Jakarta, 2017). The work performed by village PWP consists mainly of intense physical activity involving the hands, such that handgrip strength is essential in their tasks. Lowering of handgrip strength may indicate reduced muscle strength, which may especially interfere with the daily activities of village PWP that consist mainly of manual labor.

Handgrip strength measures maximal hand strength and indicates muscle quality that reflects muscle strength and mass. Muscle strength refers to the ability of muscle groups to raise a given load. Low or poor grip strength is frequently associated with damaging health impacts, such as chronic disease, and is the initial sign of reduced muscle mass, limitations in physical functions, and mortality risk. Poor handgrip strength may also reduce a person's ability to perform activities of daily living and, therefore, impact quality of life (Soraya et al., 2023; Nathania et al., 2023).

Handgrip strength may represent the strength of all muscles in the body. Several factors possibly influencing an individual's muscle strength are age, sex, and body mass index (BMI) (Soraya et al., 2023). Handgrip strength is also associated with chronic diseases such as diabetes, metabolic syndrome, and cardiovascular disease. Increased glucose concentration is also associated with disorders of muscle quality and strength, as well as physical performance (Chen et al., 2023). In the present study, we included fasting blood glucose concentrations in the data collection in order to obtain more accurate measures of glucose concentrations that have not yet been influenced by food intake.

Handgrip strength may be a health indicator that provides information on a person's health (Vaishya et al., 2024). However, its clinical significance in determining the prevention and treatment of poor handgrip remains limited because of the still unclear etiology of muscle weakness, especially because the clinical outcomes associated with handgrip strength are extremely diverse (Mc Grath et al., 2020). The relationship between BMI and handgrip strength is still controversial, with several investigators finding a significant positive association and others reporting a non-significant relationship (Al Asadi, 2018).

There are few studies on handgrip strength, especially on workers in the sanitation sector such as public works personnel, such that the present investigators were interested in studying this topic. The aim of the present study was to determine the factors associated with handgrip strength in public works personnel.

METHOD

This cross-sectional study involving male public works personnel in Cengkareng District, West Jakarta, was conducted from September to October 2024. The study sample was selected by simple random sampling. Calculation of the sample size for a correlation with $r=0.2$, $Z\alpha=0.05$, and $Z\beta=0.1$, resulted in a sample of 192 subjects who met the inclusion criterion of being male and 25-

60 years old, and the exclusion criteria of having an abnormality of the hands, heart disease, pulmonary disease, and malignancies.

Measurements were obtained on height in cm, weight in kg, as well as waist circumference and hip circumference in cm. Height and weight were determined using microtoise and digital scales, respectively. BMI was calculated using the formula of weight divided by height squared. Measurement of waist and hip circumferences was done using measuring tape. Waist circumference was determined with the subject in the upright position, at the end of a normal expiration, at the smallest abdominal circumference between the iliac crest and twelfth rib. The cut-off values for waist circumference in males and females are ≥ 90 cm and ≥ 80 cm, respectively. Hip circumference was measured using measuring tape that was applied around the hips at the level of the symphysis pubis and the maximal gluteal circumference (Rokhmah et al., 2015; Ilmi et al., 2020).

Fasting blood glucose (FBG) concentration was determined on venous blood by the hexokinase method after the respondents had fasted approximately 10 hours prior. Handgrip strength was measured with a handgrip strength dynamometer to the nearest kilogram as follows: The respondent was asked to sit down comfortably without reclining and with the feet completely relaxed. The hips and knees were positioned at a 90^0 angle, the gripping hand with the shoulder in adduction, the elbow flexed 90^0 , and the forearm and wrist in the neutral position. The gripping duration was 3 seconds. The respondent gripped the instrument with the dominant hand as hard as possible and the value indicated by the needle was recorded. In accordance with the protocol recommended by the American Society of Hand Therapists (ASHT), the recorded handgrip strength was the average of three handgrip strength tests (Nathania et al., 2023).

Prior to data analysis by means of the correlation test, the normality of the data distribution was determined using the Kolmogorov-Smirnov test. The Pearson correlation test was used for a normal data distribution, whereas for a non-normal data distribution, the Spearman correlation test was done, followed by a multiple regression analysis.

Ethics approval was issued by the Ethics Committee of the Faculty of Medicine, Universitas Trisakti, under No. 033/KER/FK/08/2024.

RESULT AND DISCUSSION

In this study the respondents comprised 192 male public works personnel of Cengkareng District that consisted of the villages (subdistricts) of Kedaung, Duri Kosambi, West Cengkareng, Kapuk, and Rawa Buaya. A total of 26% of the respondents were residents of Duri Kosambi, where Muslims comprised 98.4% of the population. The majority of the respondents at 77.6% had an educational level of senior high school, nearly all of the respondents, or 99%, were right-handed, using the right hand as the dominant one at work, and most were of the Betawi (Jakarta) ethnic group at 51.6% (Table 1).

Table 1. Demographic data of the respondents

Characteristic	N (%)
Village	
Kedaung	40 (20.8)
Duri Kosambi	50 (26)
Cengkareng Barat	33 (17.2)
Kapuk	43 (22.4)
Rawa Buaya	26 (13.5)
Religion	
Islamic	189 (98.4)
Christian	2 (1)
Catholic	1 (0.5)
Level of education	
Elementary school	7 (3.6)
Junior high school	29 (15.1)
Senior high school	149 (77.6)
Academy-University	7 (3.6)
Dominant hand	
Right hand	190 (99)
Left hand	2 (1)
Ethnic group	
Javanese	53 (27.6)
Betawi	99 (51.6)
Sundanese	25 (13)
Other	14 (7.3)

Legend: Other ethnicities comprised groups from Palembang, Lampung, Bima, and Kupang

Table 2. Respondents' characteristics

Variable	Mean $\pm$ SD
Age (years)	40.9 $\pm$ 8.9
Height (cm)	164.6 $\pm$ 6.7
Weight (kg)	64.7 $\pm$ 13.1
BMI (kg/m ²)	23.8 $\pm$ 4.3
Waist circumference (cm)	85.4 $\pm$ 10.4
Hip circumference (cm)	96.9 $\pm$ 8.4
HGS (kg)	33.1 $\pm$ 6.9
FBG (mg/dL)	87 (64-280)*

Legend: BMI: body mass index. *FBG: fasting blood glucose, expressed as Median (Min-Max).

HGS: handgrip strength

The bivariate analysis found a significant positive correlation of BMI, height, weight, waist circumference, and hip circumference with handgrip strength ($p=0.006$, $r=0.20$; $p<0.001$, $r=0.36$; $p<0.001$, $r=0.32$; $p=0.009$, $r=0.19$; $p=0.005$; $r=0.20$, respectively). This showed that the larger the BMI, height, weight, waist circumference, and hip circumference, the stronger the handgrip. However, there was no significant relationship between fasting glucose concentration and handgrip strength ($p=0.847$). This study also found that age had a significant negative correlation with handgrip strength ($p=0.011$; $r=-0.182$), signifying that the older a person, the lower the handgrip strength (Table 3). Results of multiple regression analysis on handgrip strength are listed in Table 4.

Table 3. Analysis of the relationship of age, height, weight, BMI, waist circumference, hip circumference, and fasting blood glucose concentration with handgrip strength

Variable	R	P value
Age	- 0.19	0.008*
Height	0.36	0.000*
Weight	0.32	0.000*
BMI	0.20	0.006*
Waist circumference	0.19	0.009*
Hip circumference	0.20	0.005*
FBG	0.02	0.847 [#]

*P value with Pearson's test, [#] Spearman's test

Table 4. Multiple regression analysis on handgrip strength

Determinant	b	SE	p
Height	0.48	0.08	0.000
BMI	0.79	0.22	0.000
Hip circumference	- 0.29	0.12	0.015
Constant	- 36.98		

$R^2=0.19$

This study found that most of the public works personnel in Duri Kosambi village were Muslims, had a senior high school level of education, and were of Betawi ethnicity. This was in line with the demographic data in Cengkareng district, in that Duri Kosambi was the village with the greatest area, the majority of the residents were male, and the Islamic religion had the most believers (Badan Pusat Statistik Jakarta Barat, 2020).

Among the respondents of this study, the dominant right hand was used for working. In the adult population, a total of 90% of the residents preferred to use the right hand for manual tasks, whereas around 10% used the left hand (Zaccagni et al., 2020). According to Agtuahene et al. (Agtuahene et al., 2023), the dominant hand is generally around 10% stronger than the non-dominant hand. Many activities in daily life need a high level of exertion of the muscles of the hand and forearm (Hammed et al., 2017). The frequent use of the dominant hand in daily living results in the muscles of the hand and forearm providing strength in gripping (Afaq et al., 2019).

The mean handgrip strength in this study was 33.1 ± 6.9 kg, which is relatively low if compared with the study results of Kurniawan et al. (Kurniawan et al., 2018). in healthy young

Indonesian adults, where mean handgrip strength in males was 37.37 ± 8.29 kg and the lower limit of muscle strength in Indonesian males was 20.79 kg. The handgrip strength results in both studies are still below those of males in other Southeast Asian countries, according to the study results of Leong et al. (Leong et al., 2016). These investigators found that the mean handgrip strength in Southeast Asian males aged 35-40 years was 40 kg (34-44 kg); for the age range of 41-50 years, the handgrip strength was 38 kg (33-44 kg), for the age range of 51-60 years it was 34 kg (30-40 kg), and for ages 61-70 years 30 kg (24-34 kg). This may have been caused by ethnic differences mediated by genetic differences that also affect individual muscle strength. The highest handgrip strength is found in the European and North American populations, whereas the lowest handgrip strength is found in African, South Asian, and Southeast Asian populations. These variations in muscle strength in various countries may also be partly caused by differences in socio-economic status (Leong et al., 2016).

According to Agtuahene et al. (Agtuahene et al., 2023), handgrip strength is a physiological variable influenced by several factors, such as age, sex, and body measurements. In contrast, according to Zaccagni et al. (Zaccagni et al., 2020), handgrip strength, apart from being affected by age, may also be influenced by physical activity, grip measurement, height, and body mass.

The present study found a significant positive correlation between BMI and handgrip strength. According to Hammed et al., this may be caused by the higher percentage of skeletal muscle mass compared to the percentage of fat mass, thereby resulting in a higher handgrip strength [13]. Our study's results are identical to those of Agtuahene et al. (Agtuahene et al., 2023), who found a significant positive correlation ($r = 0.290$) of handgrip strength with BMI and height. This shows that taller individuals with higher BMI apparently have greater handgrip strength. Our study is also in line with the study of Krakauer in 2020, which showed a positive correlation between BMI and handgrip strength (Krakauer et al., 2020).

The study of Cooper et al. (Cooper et al., 2022) concluded that a high childhood BMI is associated with higher grip strength at age 46 years. This is because fat mass acts as a mechanical load that elicits an anabolic response that stimulates muscle growth and function. This anabolic response is usually greater in males than in females because of the higher testosterone concentration in males. However, it should also be noted that BMI does not differentiate between fat mass and lean mass as adipocyte indicators that may vary with sex. It should be noted that BMI is a measure of body fat that does not differentiate between weight changes as a result of an increase or decrease in muscle mass or body fat (Zaccagni et al., 2020). According to Jafar et al. (Jaafar et al., 2023), heavier individuals have a greater muscle mass, resulting in greater handgrip strength, than lighter individuals.

In the study of Heidy et al. (Heidy et al., 2019), the investigators found a positive correlation between height and forearm length. The study by Sirajudeen et al. (Sirajudeen et al., 2012) found a positive correlation between hand length and handgrip strength. This shows that height affects the longer extremity or arm, thus approaching optimal muscle length that may result in greater grip strength. According to a prospective cohort study, handgrip strength in middle age appears to be associated with birth weight and the increase in height at puberty. Pubertal growth and handgrip strength possibly represent the development of muscle size and growth (Schrager et al., 2007).

Our study found no relationship between blood glucose concentration and handgrip strength. Our study agrees with the study of Giglio et al. (Giglio et al., 2018), which showed no significant relationship between muscle mass strength and blood glucose concentration. According to Giglio, this is because most respondents are individuals who routinely perform structured

physical activity, even though they are in the light category. This is similar to our study respondents, who were public works personnel and who had performed physical activity such as street sweeping and cleaning storm drains.

Glucose is the main fuel for muscle contraction, entering the muscle cells by diffusion through the GLUT4 glucose transporter, which, after muscle contraction, moves from its intracellular storage site to the plasma membrane and T-tubules (Richter et al., 2013). Glucose metabolism has an influence on handgrip strength through various mechanisms, such as disorders of glucose metabolism, particularly glycogenolysis, that may result in the loss of muscle strength. Insulin resistance in hyperglycemia may cause muscle degradation because hyperglycemia causes low muscle strength through effects on skeletal muscle mitochondria. The lowering of muscle strength by hyperglycemia occurs through Krüppel-like factor 15, which regulates skeletal muscle lipid flux. Hyperglycemia also increases pro-inflammatory cytokines, which may increase the catabolism that plays a role in the further decrease in muscle mass and quality (Gamayanti et al., 2023).

This study has a significant negative correlation between age and handgrip strength, which decreases with age (Manoharan et al., 2015). Handgrip strength initially experiences an increase with increasing age, reaches its peak between the age of 30 and 40 years, and decreases afterward (Zaccagni et al., 2020). The age-related changes in skeletal muscle composition, particularly the accumulation of lipids in skeletal muscle fibers, play a role in the poor quality of the muscles and may result in metabolic abnormalities such as insulin resistance. Apart from changes in muscle quality, there is also a reduction in muscle mass that is consistently occurring during aging. The increase in oxidative stress and chronic inflammation that commonly occurs in the aging process apparently also plays a role in the development of decreased muscle mass and strength (Mainous et al., 2015).

The decrease in muscle mass in the elderly corresponds to a 10% reduction in grip strength per decade, starting from the age of 40 years (Jaafar et al., 2023). Skeletal muscle plays an important role in glucose metabolism and has a significant influence on insulin sensitivity. Increased insulin resistance in older adults may be closely associated with skeletal muscle aging. A reduction in GLUT4 expression as a result of a decrease in muscle volume causes a reduction in insulin sensitivity in aging skeletal muscle (Joo et al., 2022).

The results of this study showed a correlation of waist circumference and hip circumference with handgrip strength. This is in line with the study of Widjaja et al. (Widjaja et al., 2018), who stated that there was a significant correlation of waist circumference and hip circumference with handgrip strength ($p < 0.001$, $r = 0.302$; and $p < 0.001$, $r = 0.296$). A greater waist circumference is associated with a greater handgrip strength in males (Hardy et al., 2013).

In the multiple regression analysis of HGS, we removed the data on body weight because of the occurrence of multicollinearity between weight and BMI, with a correlation coefficient of more than 0.9, indicating an extremely strong relationship between these two variables. Conceptually, only one of the variables is used in multivariate analysis. The multiple regression analysis included age, height, BMI, waist circumference, and hip circumference, with HGS as dependent variable in a stepwise dependent manner. After three iterations, only height, BMI, and hip circumference played a role in predicting HGS by means of the regression formula, which resulted in $R^2 = 0.19$. This signifies that for known values of height, BMI, and hip circumference, we can predict the HGS value 19% more accurately than without a knowledge of these data. For a clinical diagnosis, a minimum increase of 10% is needed. This study found that height, BMI, and

hip circumference play a role in predicting handgrip strength through the regression formula, i.e., $\text{handgrip strength} = -36.98 + 0.48 \times \text{height} + 0.79 \times \text{BMI} - 0.29 \times \text{hip circumference}$.

Handgrip strength and waist circumference are surrogate muscle strength and visceral adiposity markers, respectively (Nakanishi et al., 2024). Waist circumference is also more sensitive in determining body fat distribution, particularly in the abdomen. Waist circumference correlates better with abdominal fat distribution than BMI (Rokhmah et al., 2015). Waist circumference and hip circumference are associated with obesity, in which the fat content is higher, and the skeletal muscle and fascicular content is lower, thereby resulting in a lower contractile capacity of skeletal muscle (Afaq et al., 2019).

The limitation of this study is that it did not evaluate other factors that may affect handgrip strength, such as nutrient intake. From the anthropometric measures of the hands and feet in this study, the causes and effects of the decrease in handgrip strength could not be determined because this study used a cross-sectional design.

CONCLUSION

There are three factors that may be used to estimate handgrip strength in PWP, namely height, BMI, and hip circumference, whereas blood glucose concentration is not correlated with handgrip strength in PWP. It is recommended to pay attention to height, BMI, and hip circumference, particularly in the acceptance criteria of PWP candidates, and to conduct further studies with consideration of other factors that affect handgrip strength, such as food intake and anthropometric features of the hands and arms.

ACKNOWLEDGMENTS

Financial support has been provided from Universitas Trisakti for this research.

REFERENCES

- Afaq, E., Hassan, S.H., Khan, M.R., Zaidi, D.A., & Baktashi, H., 2019. Correlation of Hand Dynamometry With Body Mass Index and Waist Hip Ratio in Medical Students. *Pak J Physiol*, 15, pp.19-21.
- Agtuahene, M.A., Quartey, J., & Kwakye, S., 2023. Influence of Hand Dominance, Gender, and Body Mass Index on Hand Grip Strength. *S Afr J Physiother*, 79, pp.1-6. doi: [org/10.4102/sajp.v79i1.1923](https://doi.org/10.4102/sajp.v79i1.1923).
- Al-Asadi, J.N., 2018. Handgrip Strength in Medical Students: Correlation with Body Mass Index and Hand Dimensions. *Asian Journal of Medical Sciences*, 9, pp.21-26. doi: [10.3126/ajms.v9i1.18577](https://doi.org/10.3126/ajms.v9i1.18577).

Badan Pusat Statistik Kota Administrasi Jakarta Barat., 2020. Kecamatan Cengkareng Dalam Angka, pp. 1-114.

Chen, D., Zhu, Y., Ni, W., Li, Y., Yin, G., Shao, Z., & Zhu, J., 2023. Hand Grip Strength is Inversely Associated with Total Daily Insulin Dose Requirement in Patients with Type 2 Diabetes Mellitus: A Cross-Sectional Study. *PeerJ*, pp.1-13. doi: 10.7717/peerj.15761.

Cooper, R., Tomlinson, D., Hamer, M., & Pereira, S.M.P., 2022. Lifetime Body Mass Index and Grip Strength at Age 46 Years: The 1970 British Cohort Study. *J Cachexia Sarcopenia Muscle*, 13, pp.1-10. doi: 10.1002/jcsm.12992.

Gamayanti, K.A.V., Aryana, I.G.P.S., & Gotera, W., 2023. Hubungan Antara Kekuatan Genggaman Tangan Dengan Kadar Gula Darah Sewaktu Pada Lansia Di Desa Melinggih, Kecamatan Payangan, Kabupaten Gianyar. *Intisari Sains Medis*, 14(2): pp.483-488.

Giglio, B.M., Mota, J.F., Wall, B.T., & Pimentel, G.D., 2018. Low Handgrip Strength is Not Associated With Type 2 Diabetes Mellitus and Hyperglycemia: A Population-Based Study. *Clin Nutr Res*, 7(2): pp.112-116. doi: 10.7762/cnr.2018.7.2.112.

Hammed, A.I., & Obaseki, C.O., 2017. Interdependence of Body Mass Index with Handgrip Strength and Endurance Among Apparently Healthy Teenagers. *Turk J Kin*, 4, pp.1-7.

Hardy, R., Cooper, R., Sayer, A.A., Shlomo, Y.B., Cooper, C., Deary, I.J., Demakakos, P., Gallacher, J., Martin, R.M., McNeill, G., et al., 2013. Body Mass Index, Muscle Strength and Physical Performance in Older Adults From Eight Cohort Studies: The Halcyon Programme. *PLoS One*, 8(2): pp.e56483. doi: 10.1371/journal.pone.0056483.

Heidy., Djuartina, T., & Irawan, R., 2019. Korelasi Kekuatan Genggaman Tangan Dengan Karakter Antropometri Lengan Bawah dan Tangan Serta Indeks Massa Tubuh. *Damianus Journal of Medicine*, 18, pp.1-7.

Ilmi, A.F., & Utari, D.M., 2020. Hubungan Lingkar Pinggang dan Rasio Lingkar Pinggang-Panggul (RLPP) Terhadap Kadar Glukosa Darah Puasa Pada Mahasiswa. *Journal of Nutrition College*, 9, pp.222-227.

Jaafar, M.H., Ismail, R., Ismail, N.H., Isa, Z.M., Tamil, A.M., Nasir, N.M., Keat, N.K., Razak, N.H., Abidin, N.Z., & Yusof, K.H., 2023. Normative Reference Values and Predicting Factors of Handgrip Strength For Dominant and Non-Dominant Hands Among Healthy Malay Adults in Malaysia. *BMC Musculoskelet Disord*, 24, pp. 1-9. doi: 10.1186/s12891-023-06181-8.

Joo, K.C., Son, D.H., & Park, J.M., 2022. Association Between Relative Handgrip Strength and Insulin Resistance in Korean Elderly Men Without Diabetes: Findings Of The 2015 Korea National Health Nutrition Examination Survey. *Korean J Fam Med*, 43, pp.199-205. doi:10.4082/kjfm.21.0138.

Krakauer, N.Y., & Krakauer, J.C., 2020. Association of Body Shape Index (ABSI) with Hand Grip Strength. *Int J Environ Res Public Health*, 17, pp.1-12. doi: doi:10.3390/ijerph17186797.

Kurniawan, A., Widjaja, D., & Lugito, N.P.H., 2018. Mean and Cutoff Value of Hand Grip Strength For Healthy Indonesian Young Adults. World Congress on Osteoporosis, Osteoarthritis and Musculoskeletal Diseases; Krakow, Poland.

Leong, D.P., Teo, K.K., Rangarajan, S., Kutty, R., Lanas, F., Hui, C., Quanyong, X., Zhenzhen, Q., Jinhua, T., Noorhassim, I., et al., 2016. Reference Ranges of Handgrip Strength From 125,462 Healthy Adults in 21 Countries: A Prospective Urban Rural Epidemiologic (PURE) Study. *J Cachexia Sarcopenia Muscle*, 7, pp.535-546. doi: 10.1002/jcsm.12112.

Mainous, A.G., Tanner, R.J., Anton, S.D., & Jo, A., 2015. Grip Strength as A Marker of Hypertension and Diabetes in Healthy Weight Adults. *Am J Prev Med*, 49(6): pp.850-858. doi: 10.1016/j.amepre.2015.05.025.

Manoharan, V.S., Sundaram, S.G., & Jason, J.I., 2015. Factors Affecting Hand Grip Strength and Its Evaluation: A Systemic Review. *Int J Physiother Res*, 3(6): pp.1288-1293. doi:10.16965/ijpr.2015.193.

McGrath, R., Johnson, N., Klawitter, L., Mahoney, S., Trautman, K., Carlson, C., Rockstad, E., & Hackney, K.J., 2020. What Are The Association Patterns Between Handgrip Strength and Adverse Health Conditions? A Topical Review. *SAGE Open Medicine*, 8, pp.1-12. doi:10.1177/2050312120910358.

Nakanishi, S., Shimoda, M., Kimura, T., Sanada, J., Fushimi, Y., Iwamoto, Y., Iwamoto, H., Dan, K., Mune, T., Kaku, K., et al., 2024. The Impact of Handgrip Strength and Waist Circumference on Glycemic Control: Prospective, Observational Study Using Outpatient Clinical Data in Japanese Patients With Type 2 Diabetes Mellitus. *J Diabetes Investig*, 15(7): pp.892-898. doi: 10.1111/jdi.14200.

Nathania, N.P.S., Nugraha, M.H.S., Primayanti, I.D.A., & Saraswati, P.A.S., 2023. Handgrip Strength is Associated With Balance in Elderly Women. *Majalah Ilmiah Fisioterapi Indonesia*, 11, pp.70-75.

Peraturan Gubernur Provinsi Daerah Khusus Ibukota Jakarta Nomor 7 tahun 2017 Tentang Penanganan Prasarana dan Sarana Umum Tingkat Kelurahan. Jakarta 2017.

Richter, E.A., & Hargreaves, M., 2013. Exercise, GLUT4, and Skeletal Muscle Glucose Uptake. *Physiol Rev*, 93(3): pp.993-1017. doi: 10.1152/physrev.00038.2012.

Rokhmah, F.D., Handayani, D., & Al-Rasyid, H., 2015. Korelasi Lingkar Pinggang dan Rasio Lingkar Pinggang-Panggul Terhadap Kadar Glukosa Plasma Menggunakan Tes Toleransi Glukosa Oral. *Jurnal Gizi Klinik Indonesia*, 12, pp.28-35.

Schrager, M.A., Metter, E.J., Simonsick, E., Ble, A., Bandinelli, S., Lauretani, F., & Ferrucci, L., 2007. Sarcopenic Obesity and Inflammation in The Inchiati Study. *J Appl Physiol*, 102(3): pp.919-925. doi: 10.1152/jappphysiol.00627.2006.

Sirajudeen, M.S., Shah, U.N., Pillai, P.S., Mohasin, N., & Shantaram, M., 2012. Correlation Between Grip Strength and Physical Factors in Men. *International Journal of Health and Rehabilitation Sciences*, 1, pp.58-63.

Soraya, N., & Parwanto, E., 2023. The Controversial Relationship Between Body Mass Index and Handgrip Strength in The Elderly: An Overview. *Malays J Med Sci*, 30, pp.73-83.

Vaishya, R., Misra, A., Vaish, A., Ursino, N., & D'Ambrosi, R., 2024. Hand Grip Strength as A Proposed New Vital Sign Of Health: A Narrative Review of Evidences. *Journal of Health, Population and Nutrition*, 43, pp.1-14. doi:10.1186/s41043-024-00500-y.

Widjaja, D., Tjhai, K., Dermawan, K., Kalim, J., Tanzil, S., Pramusinta, A., Phenca, H.K., Kurniawan, A., 2018. Correlations Between Hand Grip Strength and Anthropometric Status in Young Indonesian Adults. *World Congress on Osteoporosis, Osteoarthritis and Musculoskeletal Diseases*; Krakow, Poland.

Zaccagni, L., Toselli, S., Bramanti, B., Gualdi-Russo, E., Mongillo, J., & Rinaldo, N., 2020. Handgrip Strength in Young Adults: Association With Anthropometric Variables and Laterality. *Int J Environ Res Public Health*, 17, pp.1-18. doi:10.3390/ijerph17124273.

2. Bukti konfirmasi review dan
hasil review pertama
(22 Juni 2025)



Alvina fk <dr.alvina@trisakti.ac.id>

Discussion (Submission)

2 messages

Oktia woro kasmini handayani <oktia2016@mail.unnes.ac.id>
To: Alvina Alvina <dr.alvina@trisakti.ac.id>

Sun, Jun 22, 2025 at 9:15 AM

Please enter your message.

Reply to this comment at [#25917 Alvina et al.](#) or [unsubscribe](#) from emails sent by [Jurnal Kesehatan Masyarakat](#).



(1)+Alvina-template-kemas+(Revisi)-Factor+Associated+with+Handgripstrength+of+Public+Works+Personeel+in+West+Jakarta.docx
44K

Alvina fk <dr.alvina@trisakti.ac.id>
To: Pusparini Pusparini <pusparini@trisakti.ac.id>

Mon, Jun 23, 2025 at 12:33 PM

Dear Prof Puspa,
Prof, I forward email from Jurnal Kemas Unnes, about input my article from chief editor Jurnal Kemas.
Prof, can you give me advice about that.
Thank you Prof
Regards,
Vina

[Quoted text hidden]



(1)+Alvina-template-kemas+(Revisi)-Factor+Associated+with+Handgripstrength+of+Public+Works+Personeel+in+West+Jakarta.docx
44K

Factors Associated With Handgrip Strength of Public Works Personnel in West Jakarta

Alvina^{1,a)}, Pusparini^{2,b)}, Mario^{3,c)}, Yasmine Mashabi^{4,d)}

^{1,2,3,4} Department of Clinical Pathology, Faculty of Medicine, Universitas Trisakti

^{a)}Corresponding author: dr.alvina@trisakti.ac.id

^{b)}pusparini@trisakti.ac.id

^{c)}mario@trisakti.ac.id

^{d)} yasminedrp@gmail.com

Abstract

Public Works Personnel (PWP) in Jakarta perform their routine duties manually instead of using equipment. Handgrip strength (HGS) measures maximum hand strength as a quality indicator for muscle strength and mass. Factors affecting muscle strength are age, sex, body mass index (BMI), waist circumference (WC), and hip circumference (HC). Increased fasting blood glucose (FBG) is also associated with muscle quality, muscle strength, and physical performance. This study aimed to determine the factors associated with handgrip strength in PWP. A total of 192 male PWP from the Cengkareng district were recruited by simple random sampling. Collected data were demographics, BMI, FBG, WC, HC, and HGS. Statistical analysis was done using Pearson's correlation test and multiple regression analysis. There was a significant positive correlation of BMI, WC, and HC with HGS ($p=0.006$; $r=0.20$, $p=0.009$; $r=0.19$ and $p=0.005$; $r=0.20$) and a significant negative correlation between age and HGS ($p=0.008$; $r=-0.19$) but not between FBG and HGS ($p=0.847$). Multiple regression analysis showed height, BMI, and HC positively predicting HGS ($R^2=0.19$). Handgrip strength of public works personnel is associated with BMI, age, waist circumference, and hip circumference, but not with fasting blood glucose. Height, BMI, and hip circumference predict handgrip strength.

Keywords: Handgrip strength, Body mass index, Hip circumference, Waist circumference, Fasting blood glucose

INTRODUCTION

Personnel for the management of public infrastructure and facilities at the village level are workers who manage village public works for a given time period on the basis of a work order. The tasks of village-level public works personnel (PWP) consist of managing the infrastructure and facilities, such as streets, drains, and parks, including installed lights (Peraturan Gubernur DKI Jakarta, 2017). The work performed by village PWP consists mainly of intense physical activity involving the hands, such that handgrip strength is essential in their tasks. Lowering of handgrip strength may indicate reduced muscle strength, which may especially interfere with the daily activities of village PWP that consist mainly of manual labor.

Handgrip strength measures maximal hand strength and indicates muscle quality that reflects muscle strength and mass. Muscle strength refers to the ability of muscle groups to raise a given load. Low or poor grip strength is frequently associated with damaging health impacts, such as chronic disease, and is the initial sign of reduced muscle mass, limitations in physical functions, and mortality risk. Poor handgrip strength may also reduce a person's ability to perform activities of daily living and, therefore, impact quality of life (Soraya et al., 2023; Nathania et al., 2023). Muscle mass accounts for approximately 40% of total weight, such that knowledge of muscle mass conditions is essential in maintaining health and activities (Handayani et al., 2018).

Handgrip strength may represent the strength of all muscles in the body. Several factors possibly influencing an individual's muscle strength are age, sex, and body mass index (BMI) (Soraya et al., 2023). Obesity may also affect BMI. Obesity in adults is not only associated with genetic factors but also with environmental factors during development (Fauzi et al., 2022). Handgrip strength is also associated with chronic diseases such as diabetes, metabolic syndrome, and cardiovascular disease. Increased glucose concentration is also associated with disorders of muscle quality and strength, as well as physical performance (Chen et al., 2023). A reduction in muscle mass also affects an individual's strength and physical endurance. A combination of proteins is considered to be able to maintain muscle function, thereby building body strength and physical endurance (Soenyoto et al., 2017). In the present study, we included fasting blood glucose concentrations in the data collection in order to obtain more accurate measures of glucose concentrations that have not yet been influenced by food intake.

Handgrip strength may be a health indicator that provides information on a person's health (Vaishya et al., 2024). However, its clinical significance in determining the prevention and treatment of poor handgrip remains limited because of the still unclear etiology of muscle weakness, especially because the clinical outcomes associated with handgrip strength are extremely diverse (Mc Grath et al., 2020). The relationship between BMI and handgrip strength is still controversial, with several investigators finding a significant positive association and others reporting a non-significant relationship (Al Asadi, 2018).

There are few studies on handgrip strength, especially on workers in the sanitation sector such as public works personnel, such that the present investigators were interested in studying this topic. The aim of the present study was to determine the factors associated with handgrip strength in public works personnel.

METHOD

This cross-sectional study involving male public works personnel in Cengkareng District, West Jakarta, was conducted from September to October 2024. The study sample was selected by simple random sampling. Calculation of the sample size for a correlation with $r=0.2$, $Z\alpha=0.05$, and $Z\beta=0.1$, resulted in a sample of 192 subjects who met the inclusion criterion of being male and 25-60 years old, and the exclusion criteria of having an abnormality of the hands, heart disease, pulmonary disease, and malignancies.

Measurements were obtained on height in cm, weight in kg, as well as waist circumference and hip circumference in cm. Height and weight were determined using microtoise and digital scales, respectively. BMI was calculated using the formula of weight divided by height squared. Measurement of waist and hip circumferences was done using measuring tape. Waist circumference was determined with the subject in the upright position, at the end of a normal expiration, at the smallest abdominal circumference between the iliac crest and twelfth rib. The cut-off values for waist circumference in males and females are ≥ 90 cm and ≥ 80 cm, respectively. Hip circumference was measured using measuring tape that was applied around the hips at the level of the symphysis pubis and the maximal gluteal circumference (Rokhmah et al., 2015; Ilmi et al., 2020).

Fasting blood glucose (FBG) concentration was determined on venous blood by the hexokinase method after the respondents had fasted approximately 10 hours prior. Handgrip strength was measured with a handgrip strength dynamometer to the nearest kilogram as follows: The respondent was asked to sit down comfortably without reclining and with the feet completely relaxed. The hips and knees were positioned at a 90° angle, the gripping hand with the shoulder in adduction, the elbow flexed 90° , and the forearm and wrist in the neutral position. The gripping duration was 3 seconds. The respondent gripped the instrument with the dominant hand as hard as possible and the value indicated by the needle was recorded. In accordance with the protocol recommended by the American Society of Hand Therapists (ASHT), the recorded handgrip strength was the average of three handgrip strength tests (Nathania et al., 2023).

Prior to data analysis by means of the correlation test, the normality of the data distribution was determined using the Kolmogorov-Smirnov test. The Pearson correlation test was used for a normal data distribution, whereas for a non-normal data distribution, the Spearman correlation test was done, followed by a multiple regression analysis.

Ethics approval was issued by the Ethics Committee of the Faculty of Medicine, Universitas Trisakti, under No. 033/KER/FK/08/2024.

RESULT AND DISCUSSION

In this study the respondents comprised 192 male public works personnel of Cengkareng District that consisted of the villages (subdistricts) of Kedaung, Duri Kosambi, West Cengkareng, Kapuk, and Rawa Buaya. A total of 26% of the respondents were residents of Duri Kosambi, where Muslims comprised 98.4% of the population. The majority of the respondents at 77.6% had an educational level of senior high school, nearly all of the respondents, or 99%, were right-handed, using the right hand as the dominant one at work, and most were of the Betawi (Jakarta) ethnic group at 51.6% (Table 1).

Table 1. Demographic data of the respondents

Characteristic	N (%)
Village	
Kedaung	40 (20.8)
Duri Kosambi	50 (26)
Cengkareng Barat	33 (17.2)
Kapuk	43 (22.4)
Rawa Buaya	26 (13.5)
Religion	
Islamic	189 (98.4)
Christian	2 (1)
Catholic	1 (0.5)
Level of education	
Elementary school	7 (3.6)
Junior high school	29 (15.1)
Senior high school	149 (77.6)
Academy-University	7 (3.6)
Dominant hand	
Right hand	190 (99)
Left hand	2 (1)
Ethnic group	
Javanese	53 (27.6)
Betawi	99 (51.6)
Sundanese	25 (13)
Other	14 (7.3)

Legend: Other ethnicities comprised groups from Palembang, Lampung, Bima, and Kupang

Source: primary data (questionnaires and interviews with respondents)

Commented [MA1]: Dibawah setiap tabel sebutkan sumbernya. Silahkan mencermati aartikel yang sudah terbit di Jurnal KEMAS

Alvina: sudah ditambahkan sumber data dibawah setiap tabel

Table 2. Respondents' characteristics

Variable	Mean $\pm$ SD
Age (years)	40.9 $\pm$ 8.9
Height (cm)	164.6 $\pm$ 6.7
Weight (kg)	64.7 $\pm$ 13.1
BMI (kg/m ²)	23.8 $\pm$ 4.3
Waist circumference (cm)	85.4 $\pm$ 10.4
Hip circumference (cm)	96.9 $\pm$ 8.4
HGS (kg)	33.1 $\pm$ 6.9
FBG (mg/dL)	87 (64-280)*

Legend: BMI: body mass index. *FBG: fasting blood glucose, expressed as Median (Min-Max).

HGS: handgrip strength

Source: primary data (respondents' measurements)

The bivariate analysis found a significant positive correlation of BMI, height, weight, waist circumference, and hip circumference with handgrip strength ($p=0.006$, $r=0.20$; $p<0.001$, $r=0.36$; $p<0.001$, $r=0.32$; $p=0.009$, $r=0.19$; $p=0.005$, $r=0.20$, respectively). This showed that the larger the BMI, height, weight, waist circumference, and hip circumference, the stronger the handgrip. However, there was no significant relationship between fasting glucose concentration and handgrip strength ($p=0.847$). This study also found that age had a significant negative correlation with handgrip strength ($p=0.011$; $r=-0.182$), signifying that the older a person, the lower the handgrip strength (Table 3). Results of multiple regression analysis on handgrip strength are listed in Table 4.

Table 3. Analysis of the relationship of age, height, weight, BMI, waist circumference, hip circumference, and fasting blood glucose concentration with handgrip strength

Variable	R	P value
Age	- 0.19	0.008*
Height	0.36	0.000*
Weight	0.32	0.000*
BMI	0.20	0.006*
Waist circumference	0.19	0.009*
Hip circumference	0.20	0.005*
FBG	0.02	0.847 [#]

*P value with Pearson's test, [#] Spearman's test

Source: primary data

Table 4. Multiple regression analysis on handgrip strength

Determinant	b	SE	p
Height	0.48	0.08	0.000
BMI	0.79	0.22	0.000
Hip circumference	- 0.29	0.12	0.015
Constant	- 36.98		

R²=0,19

Source: primary data

This study found that most of the public works personnel in Duri Kosambi village were Muslims, had a senior high school level of education, and were of Betawi ethnicity. This was in line with the demographic data in Cengkareng district, in that Duri Kosambi was the village with the greatest area, the majority of the residents were male, and the Islamic religion had the most believers (Badan Pusat Statistik Jakarta Barat, 2020). Among our respondents, more than ten percent were in the diabetes category. Socio-demographic factors capable of affecting dietary habits in patients with diabetes are level of education and type of family (Rondhianto et al., 2024). Among the respondents of this study, the dominant right hand was used for working. In the adult population, a total of 90% of the residents preferred to use the right hand for manual tasks, whereas around 10% used the left hand (Zaccagni et al., 2020). According to Agtuahene et al. (Agtuahene et al., 2023), the dominant hand is generally around 10% stronger than the non-dominant hand. Many activities in daily life need a high level of exertion of the muscles of the hand and forearm (Hammed et al., 2017). The frequent use of the dominant hand in daily living results in the muscles of the hand and forearm providing strength in gripping (Afaq et al., 2019).

The mean handgrip strength in this study was 33.1 ± 6.9 kg, which is relatively low if compared with the study results of Kurniawan et al. (Kurniawan et al., 2018). in healthy young Indonesian adults, where mean handgrip strength in males was 37.37 ± 8.29 kg and the lower limit of muscle strength in Indonesian males was 20.79 kg. The handgrip strength results in both studies are still below those of males in other Southeast Asian countries, according to the study results of Leong et al. (Leong et al., 2016). These investigators found that the mean handgrip strength in Southeast Asian males aged 35-40 years was 40 kg (34-44 kg); for the age range of 41-50 years, the handgrip strength was 38 kg (33-44 kg), for the age range of 51-60 years it was 34 kg (30-40 kg), and for ages 61-70 years 30 kg (24-34 kg). This may have been caused by ethnic differences mediated by genetic differences that also affect individual muscle strength. The highest handgrip strength is found in the European and North American populations, whereas the lowest handgrip strength is found in African, South Asian, and Southeast Asian populations. These variations in muscle strength in various countries may also be partly caused by differences in socio-economic status (Leong et al., 2016).

According to Agtuahene et al. (Agtuahene et al., 2023), handgrip strength is a physiological variable influenced by several factors, such as age, sex, and body measurements. In contrast, according to Zaccagni et al. (Zaccagni et al., 2020), handgrip strength, apart from being affected by age, may also be influenced by physical activity, grip measurement, height, and body mass. The present study found a significant positive correlation between BMI and handgrip strength. According to Hammed et al., this may be caused by the higher percentage of skeletal muscle mass compared to the percentage of fat mass, thereby resulting in a higher handgrip strength [13]. Our study's results are identical to those of Agtuahene et al. (Agtuahene et al., 2023), who found a

Commented [MA2]: Terlalu banyak paragraf ,

Alvina: Paragraf sudah diringkas menjadi 9 paragraf

significant positive correlation ($r = 0.290$) of handgrip strength with BMI and height. This shows that taller individuals with higher BMI apparently have greater handgrip strength. Our study is also in line with the study of Krakauer in 2020, which showed a positive correlation between BMI and handgrip strength (Krakauer et al., 2020).

The study of Cooper et al. (Cooper et al., 2022) concluded that a high childhood BMI is associated with higher grip strength at age 46 years. This is because fat mass acts as a mechanical load that elicits an anabolic response that stimulates muscle growth and function. This anabolic response is usually greater in males than in females because of the higher testosterone concentration in males. However, it should also be noted that BMI does not differentiate between fat mass and lean mass as adipocyte indicators that may vary with sex. It should be noted that BMI is a measure of body fat that does not differentiate between weight changes as a result of an increase or decrease in muscle mass or body fat (Zaccagni et al., 2020). According to Jafar et al. (Jaafar et al., 2023), heavier individuals have a greater muscle mass, resulting in greater handgrip strength, than lighter individuals. In the study of Heidy et al. (Heidy et al., 2019), the investigators found a positive correlation between height and forearm length. The study by Sirajudeen et al. (Sirajudeen et al., 2012) found a positive correlation between hand length and handgrip strength. This shows that height affects the longer extremity or arm, thus approaching optimal muscle length that may result in greater grip strength. According to a prospective cohort study, handgrip strength in middle age appears to be associated with birth weight and the increase in height at puberty. Pubertal growth and handgrip strength possibly represent the development of muscle size and growth (Schrager et al., 2007).

Our study found no relationship between blood glucose concentration and handgrip strength. Our study agrees with the study of Giglio et al. (Giglio et al., 2018), which showed no significant relationship between muscle mass strength and blood glucose concentration. According to Giglio, this is because most respondents are individuals who routinely perform structured physical activity, even though they are in the light category. This is similar to our study respondents, who were public works personnel and who had performed physical activity such as street sweeping and cleaning storm drains. Glucose is the main fuel for muscle contraction, entering the muscle cells by diffusion through the GLUT4 glucose transporter, which, after muscle contraction, moves from its intracellular storage site to the plasma membrane and T-tubules (Richter et al., 2013). Glucose metabolism has an influence on handgrip strength through various mechanisms, such as disorders of glucose metabolism, particularly glycogenolysis, that may result in the loss of muscle strength. Insulin resistance in hyperglycemia may cause muscle degradation because hyperglycemia causes low muscle strength through effects on skeletal muscle mitochondria. The lowering of muscle strength by hyperglycemia occurs through Krüppel-like factor 15, which regulates skeletal muscle lipid flux. Hyperglycemia also increases pro-inflammatory cytokines, which may increase the catabolism that plays a role in the further decrease in muscle mass and quality (Gamayanti et al., 2023). The amount of carbohydrates also influences the glycemic response and post-prandial insulin level, impacting the supply of glycogen to the muscles (Safitri et al., 2020).

This study has a significant negative correlation between age and handgrip strength, which decreases with age (Manoharan et al., 2015). Handgrip strength initially experiences an increase with increasing age, reaches its peak between the age of 30 and 40 years, and decreases afterward (Zaccagni et al., 2020). The age-related changes in skeletal muscle composition, particularly the accumulation of lipids in skeletal muscle fibers, play a role in the poor quality of the muscles and may result in metabolic abnormalities such as insulin resistance. Apart from changes in muscle

quality, there is also a reduction in muscle mass that is consistently occurring during aging. The increase in oxidative stress and chronic inflammation that commonly occurs in the aging process apparently also plays a role in the development of decreased muscle mass and strength (Mainous et al., 2015). The decrease in muscle mass in the elderly corresponds to a 10% reduction in grip strength per decade, starting from the age of 40 years (Jaafar et al., 2023). Skeletal muscle plays an important role in glucose metabolism and has a significant influence on insulin sensitivity. Increased insulin resistance in older adults may be closely associated with skeletal muscle aging. A reduction in GLUT4 expression as a result of a decrease in muscle volume causes a reduction in insulin sensitivity in aging skeletal muscle (Joo et al., 2022).

The results of this study showed a correlation of waist circumference and hip circumference with handgrip strength. This is in line with the study of Widjaja et al. (Widjaja et al., 2018), who stated that there was a significant correlation of waist circumference and hip circumference with handgrip strength ($p < 0.001$, $r = 0.302$; and $p < 0.001$, $r = 0.296$). A greater waist circumference is associated with a greater handgrip strength in males (Hardy et al., 2013).

In the multiple regression analysis of HGS, we removed the data on body weight because of the occurrence of multicollinearity between weight and BMI, with a correlation coefficient of more than 0.9, indicating an extremely strong relationship between these two variables. Conceptually, only one of the variables is used in multivariate analysis. The multiple regression analysis included age, height, BMI, waist circumference, and hip circumference, with HGS as dependent variable in a stepwise dependent manner. After three iterations, only height, BMI, and hip circumference played a role in predicting HGS by means of the regression formula, which resulted in $R^2 = 0.19$. This signifies that for known values of height, BMI, and hip circumference, we can predict the HGS value 19% more accurately than without a knowledge of these data. For a clinical diagnosis, a minimum increase of 10% is needed. This study found that height, BMI, and hip circumference play a role in predicting handgrip strength through the regression formula, i.e., $\text{handgrip strength} = -36.98 + 0.48 \times \text{height} + 0.79 \times \text{BMI} - 0.29 \times \text{hip circumference}$. Handgrip strength and waist circumference are surrogate muscle strength and visceral adiposity markers, respectively (Nakanishi et al., 2024). Waist circumference is also more sensitive in determining body fat distribution, particularly in the abdomen. Waist circumference correlates better with abdominal fat distribution than BMI (Rokhmah et al., 2015). Waist circumference and hip circumference are associated with obesity, in which the fat content is higher, and the skeletal muscle and fascicular content is lower, thereby resulting in a lower contractile capacity of skeletal muscle (Afaq et al., 2019).

The limitation of this study is that it did not evaluate other factors that may affect handgrip strength, such as nutrient intake. From the anthropometric measures of the hands and feet in this study, the causes and effects of the decrease in handgrip strength could not be determined because this study used a cross-sectional design.

CONCLUSION

There are three factors that may be used to estimate handgrip strength in PWP, namely height, BMI, and hip circumference, whereas blood glucose concentration is not correlated with handgrip strength in PWP. It is recommended to pay attention to height, BMI, and hip circumference,

particularly in the acceptance criteria of PWP candidates, and to conduct further studies with consideration of other factors that affect handgrip strength, such as food intake and anthropometric features of the hands and arms.

ACKNOWLEDGMENTS

Financial support was provided by Universitas Trisakti for this research.

REFERENCES

- Afaq, E., Hassan, S.H., Khan, M.R., Zaidi, D.A., & Baktashi, H., 2019. Correlation of Hand Dynamometry With Body Mass Index and Waist Hip Ratio in Medical Students. *Pak J Physiol*, 15, pp.19-21.
- Agtuahene, M.A., Quartey, J., & Kwakye, S., 2023. Influence of Hand Dominance, Gender, and Body Mass Index on Hand Grip Strength. *S Afr J Physiother*, 79, pp.1-6. doi: org/10.4102/sajp.v79i1.1923.
- Al-Asadi, J.N., 2018. Handgrip Strength in Medical Students: Correlation with Body Mass Index and Hand Dimensions. *Asian Journal of Medical Sciences*, 9, pp.21-26. doi: 10.3126/ajms.v9i1.18577.
- Badan Pusat Statistik Kota Administrasi Jakarta Barat., 2020. Kecamatan Cengkareng Dalam Angka, pp. 1-114.
- Chen, D., Zhu, Y., Ni, W., Li, Y., Yin, G., Shao, Z., & Zhu, J., 2023. Hand Grip Strength is Inversely Associated with Total Daily Insulin Dose Requirement in Patients with Type 2 Diabetes Mellitus: A Cross-Sectional Study. *PeerJ*, pp.1-13. doi: 10.7717/peerj.15761.
- Cooper, R., Tomlinson, D., Hamer, M., & Pereira, S.M.P., 2022. Lifetime Body Mass Index and Grip Strength at Age 46 Years: The 1970 British Cohort Study. *J Cachexia Sarcopenia Muscle*, 13, pp.1-10. doi: 10.1002/jcsm.12992.
- Fauzi, L., Handayani, O.W.K., Susilo, M.T., Kurnia, A.R., Rahayu, S.R., Irawan, F.A., Horng Lu, F.J., Lin, C., Lai, M.F., & Yu, Y.C., 2022. Obesity in Indonesian and Taiwanese Adolescents Related to Self Perception, Diet, Exercise and Body Image. *KEMAS*, 17(3):pp.453-461. doi: 10.15294/kemas.v17i3.34396
- Gamayanti, K.A.V., Aryana, I.G.P.S., & Gotera, W., 2023. Hubungan Antara Kekuatan Genggaman Tangan Dengan Kadar Gula Darah Sewaktu Pada Lansia Di Desa Melinggih, Kecamatan Payangan, Kabupaten Gianyar. *Intisari Sains Medis*, 14(2): pp.483-488.
- Giglio, B.M., Mota, J.F., Wall, B.T., & Pimentel, G.D., 2018. Low Handgrip Strength is Not Associated With Type 2 Diabetes Mellitus and Hyperglycemia: A Population-Based Study. *Clin Nutr Res*, 7(2): pp.112-116. doi: 10.7762/cnr.2018.7.2.112.

Commented [MA3]: - Daftar pustaka minimal 20.
- Minimal 80% daftar pustaka adalah ARTIKEL JURNAL INTERNASIONAL
- Pastikan daftar pustaka artikel ada: nama penulis, tahun, judul artikel, nama jurnal, no/volume
- tambahkan 5 sitasi dari jurnal kemas UNNES

Alvina:
-Sudah ditambahkan 5 sitasi dari jurnal KEMAS UNNES

Hammed, A.I., & Obaseki, C.O., 2017. Interdependence of Body Mass Index with Handgrip Strength and Endurance Among Apparently Healthy Teenagers. *Turk J Kin*, 4, pp.1-7.

Handayani, M.D.N., Sadewa, A.H., Farmawati, A., & Rochmah, W., 2018. Anthropometric Prediction Equations for Estimating Muscle Mass of Elderly Women. *KEMAS*, 14(2):pp.195-204. doi: 10.15294/kemas.v14i2.14073

Hardy, R., Cooper, R., Sayer, A.A., Shlomo, Y.B., Cooper, C., Deary, I.J., Demakakos, P., Gallacher, J., Martin, R.M., McNeill, G., et al., 2013. Body Mass Index, Muscle Strength and Physical Performance in Older Adults From Eight Cohort Studies: The Halcyon Programme. *PLoS One*, 8(2): pp.e56483. doi: 10.1371/journal.pone.0056483.

Heidy., Djuartina, T., & Irawan, R., 2019. Korelasi Kekuatan Genggaman Tangan Dengan Karakter Antropometri Lengan Bawah dan Tangan Serta Indeks Massa Tubuh. *Damianus Journal of Medicine*, 18, pp.1-7.

Ilmi, A.F., & Utari, D.M., 2020. Hubungan Lingkar Pinggang dan Rasio Lingkar Pinggang-Panggul (RLPP) Terhadap Kadar Glukosa Darah Puasa Pada Mahasiswa. *Journal of Nutrition College*, 9, pp.222-227.

Jaafar, M.H., Ismail, R., Ismail, N.H., Isa, Z.M., Tamil, A.M., Nasir, N.M., Keat, N.K., Razak, N.H., Abidin, N.Z., & Yusof, K.H., 2023. Normative Reference Values and Predicting Factors of Handgrip Strength For Dominant and Non-Dominant Hands Among Healthy Malay Adults in Malaysia. *BMC Musculoskelet Disord*, 24, pp. 1-9. doi: 10.1186/s12891-023-06181-8.

Joo, K.C., Son, D.H., & Park, J.M., 2022. Association Between Relative Handgrip Strength and Insulin Resistance in Korean Elderly Men Without Diabetes: Findings Of The 2015 Korea National Health Nutrition Examination Survey. *Korean J Fam Med*, 43, pp.199-205. doi:10.4082/kjfm.21.0138.

Krakauer, N.Y., & Krakauer, J.C., 2020. Association of Body Shape Index (ABSI) with Hand Grip Strength. *Int J Environ Res Public Health*, 17, pp.1-12. doi: doi:10.3390/ijerph17186797.

Kurniawan, A., Widjaja, D., & Lugito, N.P.H., 2018. Mean and Cutoff Value of Hand Grip Strength For Healthy Indonesian Young Adults. *World Congress on Osteoporosis, Osteoarthritis and Musculoskeletal Diseases*; Krakow, Poland.

Leong, D.P., Teo, K.K., Rangarajan, S., Kutty, R., Lanas, F., Hui, C., Quanyong, X., Zhenzhen, Q., Jinhua, T., Noorhassim, I., et al., 2016. Reference Ranges of Handgrip Strength From 125,462 Healthy Adults in 21 Countries: A Prospective Urban Rural Epidemiologic (PURE) Study. *J Cachexia Sarcopenia Muscle*, 7, pp.535-546. doi: 10.1002/jcsm.12112.

Mainous, A.G., Tanner, R.J., Anton, S.D., & Jo, A., 2015. Grip Strength as A Marker of Hypertension and Diabetes in Healthy Weight Adults. *Am J Prev Med*, 49(6): pp.850-858. doi: 10.1016/j.amepre.2015.05.025.

Manoharan, V.S., Sundaram, S.G., & Jason, J.I., 2015. Factors Affecting Hand Grip Strength and Its Evaluation: A Systemic Review. *Int J Physiother Res*, 3(6): pp.1288-1293. doi:10.16965/ijpr.2015.193.

McGrath, R., Johnson, N., Klawitter, L., Mahoney, S., Trautman, K., Carlson, C., Rockstad, E., & Hackney, K.J., 2020. What Are The Association Patterns Between Handgrip Strength and Adverse Health Conditions? A Topical Review. *SAGE Open Medicine*, 8, pp.1-12. doi:10.1177/2050312120910358.

Nakanishi, S., Shimoda, M., Kimura, T., Sanada, J., Fushimi, Y., Iwamoto, Y., Iwamoto, H., Dan, K., Mune, T., Kaku, K., et al., 2024. The Impact of Handgrip Strength and Waist Circumference on Glycemic Control: Prospective, Observational Study Using Outpatient Clinical Data in Japanese Patients With Type 2 Diabetes Mellitus. *J Diabetes Investig*, 15(7): pp.892-898. doi: 10.1111/jdi.14200.

Nathania, N.P.S., Nugraha, M.H.S., Primayanti, I.D.A., & Saraswati, P.A.S., 2023. Handgrip Strength is Associated With Balance in Elderly Women. *Majalah Ilmiah Fisioterapi Indonesia*, 11, pp.70-75.

Peraturan Gubernur Provinsi Daerah Khusus Ibukota Jakarta Nomor 7 tahun 2017 Tentang Penanganan Prasarana dan Sarana Umum Tingkat Kelurahan. Jakarta 2017.

Richter, E.A., & Hargreaves, M., 2013. Exercise, GLUT4, and Skeletal Muscle Glucose Uptake. *Physiol Rev*, 93(3): pp.993-1017. doi: 10.1152/physrev.00038.2012.

Rokhmah, F.D., Handayani, D., & Al-Rasyid, H., 2015. Korelasi Lingkar Pinggang dan Rasio Lingkar Pinggang-Panggul Terhadap Kadar Glukosa Plasma Menggunakan Tes Toleransi Glukosa Oral. *Jurnal Gizi Klinik Indonesia*, 12, pp.28-35.

Rondhianto., Ridla, A.Z., & Hasan, H., 2024. Sociodemographic Factors Affecting Diabetic Dietary Behavior in People with Type 2 Diabetes Mellitus. *KEMAS*, 19(3):pp.429-437. doi: 10.15294/kemas.v19i3.37856

Safitri, I., Setyarsih, L., Susanto, H., Suhartono., & Fitranti, D.Y., 2020. The Effect of Low and High Glycemic Load Diet on Muscle Fatigue of Young Soccer. *KEMAS*, 16(1):pp.111-120. doi: 10.15294/kemas.v16i1.23508

Schrager, M.A., Metter, E.J., Simonsick, E., Ble, A., Bandinelli, S., Lauretani, F., & Ferrucci, L., 2007. Sarcopenic Obesity and Inflammation in The Inchiati Study. *J Appl Physiol*, 102(3): pp.919-925. doi: 10.1152/japplphysiol.00627.2006.

Sirajudeen, M.S., Shah, U.N., Pillai, P.S., Mohasin, N., & Shantaram, M., 2012. Correlation Between Grip Strength and Physical Factors in Men. *International Journal of Health and Rehabilitation Sciences*, 1, pp.58-63.

Soenyoto, T., Mollap, K., & Mungpong, J., 2017. Correlation of Energy and Protein Consumption Levels with Physical Endurance of Rhythmic Gymnast Athletes. KEMAS, 13(1):pp.131-136. doi:10.15294/kemas.v13i1.9888

Soraya, N., & Parwanto, E., 2023. The Controversial Relationship Between Body Mass Index and Handgrip Strength in The Elderly: An Overview. Malays J Med Sci, 30, pp.73-83.

Vaishya, R., Misra, A., Vaish, A., Ursino, N., & D'Ambrosi, R., 2024. Hand Grip Strength as A Proposed New Vital Sign Of Health: A Narrative Review of Evidences. Journal of Health, Population and Nutrition, 43, pp.1-14. doi:10.1186/s41043-024-00500-y.

Widjaja, D., Tjhai, K., Dermawan, K., Kalim, J., Tanzil, S., Pramusinta, A., Phenca, H.K., Kurniawan, A., 2018. Correlations Between Hand Grip Strength and Anthropometric Status in Young Indonesian Adults. World Congress on Osteoporosis, Osteoarthritis and Musculoskeletal Diseases; Krakow, Poland.

Zaccagni, L., Toselli, S., Bramanti, B., Gualdi-Russo, E., Mongillo, J., & Rinaldo, N., 2020. Handgrip Strength in Young Adults: Association With Anthropometric Variables and Laterality. Int J Environ Res Public Health, 17, pp.1-18. doi:10.3390/ijerph17124273.

3. Bukti konfirmasi submit revisi pertama, respon
kepada reviewer, dan artikel yang diresubmit
(22 Juni 2025)



Alvina fk <dr.alvina@trisakti.ac.id>

Discussion (Submission)

2 messages

Oktia woro kasmini handayani <oktia2016@mail.unnes.ac.id>
To: Alvina Alvina <dr.alvina@trisakti.ac.id>

Sun, Jun 22, 2025 at 9:15 AM

Please enter your message.

Reply to this comment at [#25917 Alvina et al.](#) or [unsubscribe](#) from emails sent by [Jurnal Kesehatan Masyarakat](#).



(1)+Alvina-template-kemas+(Revisi)-Factor+Associated+with+Handgripstrength+of+Public+Works+Personeel+in+West+Jakarta.docx
44K

Alvina fk <dr.alvina@trisakti.ac.id>
To: Pusparini Pusparini <pusparini@trisakti.ac.id>

Mon, Jun 23, 2025 at 12:33 PM

Dear Prof Puspa,
Prof, I forward email from Jurnal Kemas Unnes, about input my article from chief editor Jurnal Kemas.
Prof, can you give me advice about that.
Thank you Prof
Regards,
Vina

[Quoted text hidden]



(1)+Alvina-template-kemas+(Revisi)-Factor+Associated+with+Handgripstrength+of+Public+Works+Personeel+in+West+Jakarta.docx
44K

Factors Associated With Handgrip Strength of Public Works Personnel in West Jakarta

Alvina^{1,a)}, Pusparini^{2,b)}, Mario^{3,c)}, Yasmine Mashabi^{4,d)}

^{1,2,3,4} Department of Clinical Pathology, Faculty of Medicine, Universitas Trisakti

^{a)}Corresponding author: dr.alvina@trisakti.ac.id

^{b)}pusparini@trisakti.ac.id

^{c)}mario@trisakti.ac.id

^{d)} yasminedrp@gmail.com

Abstract

Public Works Personnel (PWP) in Jakarta perform their routine duties manually instead of using equipment. Handgrip strength (HGS) measures maximum hand strength as a quality indicator for muscle strength and mass. Factors affecting muscle strength are age, sex, body mass index (BMI), waist circumference (WC), and hip circumference (HC). Increased fasting blood glucose (FBG) is also associated with muscle quality, muscle strength, and physical performance. This study aimed to determine the factors associated with handgrip strength in PWP. A total of 192 male PWP from the Cengkareng district were recruited by simple random sampling. Collected data were demographics, BMI, FBG, WC, HC, and HGS. Statistical analysis was done using Pearson's correlation test and multiple regression analysis. There was a significant positive correlation of BMI, WC, and HC with HGS ($p=0.006$; $r=0.20$, $p=0.009$; $r=0.19$ and $p=0.005$; $r=0.20$) and a significant negative correlation between age and HGS ($p=0.008$; $r=-0.19$) but not between FBG and HGS ($p=0.847$). Multiple regression analysis showed height, BMI, and HC positively predicting HGS ($R^2=0.19$). Handgrip strength of public works personnel is associated with BMI, age, waist circumference, and hip circumference, but not with fasting blood glucose. Height, BMI, and hip circumference predict handgrip strength.

Keywords: Handgrip strength, Body mass index, Hip circumference, Waist circumference, Fasting blood glucose

INTRODUCTION

Personnel for the management of public infrastructure and facilities at the village level are workers who manage village public works for a given time period on the basis of a work order. The tasks of village-level public works personnel (PWP) consist of managing the infrastructure and facilities, such as streets, drains, and parks, including installed lights (Peraturan Gubernur DKI Jakarta, 2017). The work performed by village PWP consists mainly of intense physical activity involving the hands, such that handgrip strength is essential in their tasks. Lowering of handgrip strength may indicate reduced muscle strength, which may especially interfere with the daily activities of village PWP that consist mainly of manual labor.

Handgrip strength measures maximal hand strength and indicates muscle quality that reflects muscle strength and mass. Muscle strength refers to the ability of muscle groups to raise a given load. Low or poor grip strength is frequently associated with damaging health impacts, such as chronic disease, and is the initial sign of reduced muscle mass, limitations in physical functions, and mortality risk. Poor handgrip strength may also reduce a person's ability to perform activities of daily living and, therefore, impact quality of life (Soraya et al., 2023; Nathania et al., 2023). Muscle mass accounts for approximately 40% of total weight, such that knowledge of muscle mass conditions is essential in maintaining health and activities (Handayani et al., 2018).

Handgrip strength may represent the strength of all muscles in the body. Several factors possibly influencing an individual's muscle strength are age, sex, and body mass index (BMI) (Soraya et al., 2023). Obesity may also affect BMI. Obesity in adults is not only associated with genetic factors but also with environmental factors during development (Fauzi et al., 2022). Handgrip strength is also associated with chronic diseases such as diabetes, metabolic syndrome, and cardiovascular disease. Increased glucose concentration is also associated with disorders of muscle quality and strength, as well as physical performance (Chen et al., 2023). A reduction in muscle mass also affects an individual's strength and physical endurance. A combination of proteins is considered to be able to maintain muscle function, thereby building body strength and physical endurance (Soenyoto et al., 2017). In the present study, we included fasting blood glucose concentrations in the data collection in order to obtain more accurate measures of glucose concentrations that have not yet been influenced by food intake.

Handgrip strength may be a health indicator that provides information on a person's health (Vaishya et al., 2024). However, its clinical significance in determining the prevention and treatment of poor handgrip remains limited because of the still unclear etiology of muscle weakness, especially because the clinical outcomes associated with handgrip strength are extremely diverse (Mc Grath et al., 2020). The relationship between BMI and handgrip strength is still controversial, with several investigators finding a significant positive association and others reporting a non-significant relationship (Al Asadi, 2018).

There are few studies on handgrip strength, especially on workers in the sanitation sector such as public works personnel, such that the present investigators were interested in studying this topic. The aim of the present study was to determine the factors associated with handgrip strength in public works personnel.

METHOD

This cross-sectional study involving male public works personnel in Cengkareng District, West Jakarta, was conducted from September to October 2024. The study sample was selected by simple random sampling. Calculation of the sample size for a correlation with $r=0.2$, $Z\alpha=0.05$, and $Z\beta=0.1$, resulted in a sample of 192 subjects who met the inclusion criterion of being male and 25-60 years old, and the exclusion criteria of having an abnormality of the hands, heart disease, pulmonary disease, and malignancies.

Measurements were obtained on height in cm, weight in kg, as well as waist circumference and hip circumference in cm. Height and weight were determined using microtoise and digital scales, respectively. BMI was calculated using the formula of weight divided by height squared. Measurement of waist and hip circumferences was done using measuring tape. Waist circumference was determined with the subject in the upright position, at the end of a normal expiration, at the smallest abdominal circumference between the iliac crest and twelfth rib. The cut-off values for waist circumference in males and females are ≥ 90 cm and ≥ 80 cm, respectively. Hip circumference was measured using measuring tape that was applied around the hips at the level of the symphysis pubis and the maximal gluteal circumference (Rokhmah et al., 2015; Ilmi et al., 2020).

Fasting blood glucose (FBG) concentration was determined on venous blood by the hexokinase method after the respondents had fasted approximately 10 hours prior. Handgrip strength was measured with a handgrip strength dynamometer to the nearest kilogram as follows: The respondent was asked to sit down comfortably without reclining and with the feet completely relaxed. The hips and knees were positioned at a 90° angle, the gripping hand with the shoulder in adduction, the elbow flexed 90° , and the forearm and wrist in the neutral position. The gripping duration was 3 seconds. The respondent gripped the instrument with the dominant hand as hard as possible and the value indicated by the needle was recorded. In accordance with the protocol recommended by the American Society of Hand Therapists (ASHT), the recorded handgrip strength was the average of three handgrip strength tests (Nathania et al., 2023).

Prior to data analysis by means of the correlation test, the normality of the data distribution was determined using the Kolmogorov-Smirnov test. The Pearson correlation test was used for a normal data distribution, whereas for a non-normal data distribution, the Spearman correlation test was done, followed by a multiple regression analysis.

Ethics approval was issued by the Ethics Committee of the Faculty of Medicine, Universitas Trisakti, under No. 033/KER/FK/08/2024.

RESULT AND DISCUSSION

In this study the respondents comprised 192 male public works personnel of Cengkareng District that consisted of the villages (subdistricts) of Kedaung, Duri Kosambi, West Cengkareng, Kapuk, and Rawa Buaya. A total of 26% of the respondents were residents of Duri Kosambi, where Muslims comprised 98.4% of the population. The majority of the respondents at 77.6% had an educational level of senior high school, nearly all of the respondents, or 99%, were right-handed, using the right hand as the dominant one at work, and most were of the Betawi (Jakarta) ethnic group at 51.6% (Table 1).

Table 1. Demographic data of the respondents

Characteristic	N (%)
Village	
Kedaung	40 (20.8)
Duri Kosambi	50 (26)
Cengkareng Barat	33 (17.2)
Kapuk	43 (22.4)
Rawa Buaya	26 (13.5)
Religion	
Islamic	189 (98.4)
Christian	2 (1)
Catholic	1 (0.5)
Level of education	
Elementary school	7 (3.6)
Junior high school	29 (15.1)
Senior high school	149 (77.6)
Academy-University	7 (3.6)
Dominant hand	
Right hand	190 (99)
Left hand	2 (1)
Ethnic group	
Javanese	53 (27.6)
Betawi	99 (51.6)
Sundanese	25 (13)
Other	14 (7.3)

Legend: Other ethnicities comprised groups from Palembang, Lampung, Bima, and Kupang

Source: primary data (questionnaires and interviews with respondents)

Commented [MA1]: Dibawah setiap tabel sebutkan sumbernya. Silahkan mencermati aartikel yang sudah terbit di Jurnal KEMAS

Alvina: sudah ditambahkan sumber data dibawah setiap tabel

Table 2. Respondents' characteristics

Variable	Mean $\pm$ SD
Age (years)	40.9 $\pm$ 8.9
Height (cm)	164.6 $\pm$ 6.7
Weight (kg)	64.7 $\pm$ 13.1
BMI (kg/m ²)	23.8 $\pm$ 4.3
Waist circumference (cm)	85.4 $\pm$ 10.4
Hip circumference (cm)	96.9 $\pm$ 8.4
HGS (kg)	33.1 $\pm$ 6.9
FBG (mg/dL)	87 (64-280)*

Legend: BMI: body mass index. *FBG: fasting blood glucose, expressed as Median (Min-Max).

HGS: handgrip strength

Source: primary data (respondents' measurements)

The bivariate analysis found a significant positive correlation of BMI, height, weight, waist circumference, and hip circumference with handgrip strength ($p=0.006$, $r=0.20$; $p<0.001$, $r=0.36$; $p<0.001$, $r=0.32$; $p=0.009$, $r=0.19$; $p=0.005$, $r=0.20$, respectively). This showed that the larger the BMI, height, weight, waist circumference, and hip circumference, the stronger the handgrip. However, there was no significant relationship between fasting glucose concentration and handgrip strength ($p=0.847$). This study also found that age had a significant negative correlation with handgrip strength ($p=0.011$; $r=-0.182$), signifying that the older a person, the lower the handgrip strength (Table 3). Results of multiple regression analysis on handgrip strength are listed in Table 4.

Table 3. Analysis of the relationship of age, height, weight, BMI, waist circumference, hip circumference, and fasting blood glucose concentration with handgrip strength

Variable	R	P value
Age	- 0.19	0.008*
Height	0.36	0.000*
Weight	0.32	0.000*
BMI	0.20	0.006*
Waist circumference	0.19	0.009*
Hip circumference	0.20	0.005*
FBG	0.02	0.847 [#]

*P value with Pearson's test, [#] Spearman's test

Source: primary data

Table 4. Multiple regression analysis on handgrip strength

Determinant	b	SE	p
Height	0.48	0.08	0.000
BMI	0.79	0.22	0.000
Hip circumference	- 0.29	0.12	0.015
Constant	- 36.98		

R²=0,19

Source: primary data

This study found that most of the public works personnel in Duri Kosambi village were Muslims, had a senior high school level of education, and were of Betawi ethnicity. This was in line with the demographic data in Cengkareng district, in that Duri Kosambi was the village with the greatest area, the majority of the residents were male, and the Islamic religion had the most believers (Badan Pusat Statistik Jakarta Barat, 2020). Among our respondents, more than ten percent were in the diabetes category. Socio-demographic factors capable of affecting dietary habits in patients with diabetes are level of education and type of family (Rondhianto et al., 2024). Among the respondents of this study, the dominant right hand was used for working. In the adult population, a total of 90% of the residents preferred to use the right hand for manual tasks, whereas around 10% used the left hand (Zaccagni et al., 2020). According to Agtuahene et al. (Agtuahene et al., 2023), the dominant hand is generally around 10% stronger than the non-dominant hand. Many activities in daily life need a high level of exertion of the muscles of the hand and forearm (Hammed et al., 2017). The frequent use of the dominant hand in daily living results in the muscles of the hand and forearm providing strength in gripping (Afaq et al., 2019).

The mean handgrip strength in this study was 33.1 ± 6.9 kg, which is relatively low if compared with the study results of Kurniawan et al. (Kurniawan et al., 2018). In healthy young Indonesian adults, where mean handgrip strength in males was 37.37 ± 8.29 kg and the lower limit of muscle strength in Indonesian males was 20.79 kg. The handgrip strength results in both studies are still below those of males in other Southeast Asian countries, according to the study results of Leong et al. (Leong et al., 2016). These investigators found that the mean handgrip strength in Southeast Asian males aged 35-40 years was 40 kg (34-44 kg); for the age range of 41-50 years, the handgrip strength was 38 kg (33-44 kg), for the age range of 51-60 years it was 34 kg (30-40 kg), and for ages 61-70 years 30 kg (24-34 kg). This may have been caused by ethnic differences mediated by genetic differences that also affect individual muscle strength. The highest handgrip strength is found in the European and North American populations, whereas the lowest handgrip strength is found in African, South Asian, and Southeast Asian populations. These variations in muscle strength in various countries may also be partly caused by differences in socio-economic status (Leong et al., 2016).

According to Agtuahene et al. (Agtuahene et al., 2023), handgrip strength is a physiological variable influenced by several factors, such as age, sex, and body measurements. In contrast, according to Zaccagni et al. (Zaccagni et al., 2020), handgrip strength, apart from being affected by age, may also be influenced by physical activity, grip measurement, height, and body mass. The present study found a significant positive correlation between BMI and handgrip strength. According to Hammed et al., this may be caused by the higher percentage of skeletal muscle mass compared to the percentage of fat mass, thereby resulting in a higher handgrip strength [13]. Our study's results are identical to those of Agtuahene et al. (Agtuahene et al., 2023), who found a

Commented [MA2]: Terlalu banyak paragraf ,
Alvina: Paragraf sudah diringkas menjadi 9 paragraf

significant positive correlation ($r = 0.290$) of handgrip strength with BMI and height. This shows that taller individuals with higher BMI apparently have greater handgrip strength. Our study is also in line with the study of Krakauer in 2020, which showed a positive correlation between BMI and handgrip strength (Krakauer et al., 2020).

The study of Cooper et al. (Cooper et al., 2022) concluded that a high childhood BMI is associated with higher grip strength at age 46 years. This is because fat mass acts as a mechanical load that elicits an anabolic response that stimulates muscle growth and function. This anabolic response is usually greater in males than in females because of the higher testosterone concentration in males. However, it should also be noted that BMI does not differentiate between fat mass and lean mass as adipocyte indicators that may vary with sex. It should be noted that BMI is a measure of body fat that does not differentiate between weight changes as a result of an increase or decrease in muscle mass or body fat (Zaccagni et al., 2020). According to Jafar et al. (Jaafar et al., 2023), heavier individuals have a greater muscle mass, resulting in greater handgrip strength, than lighter individuals. In the study of Heidy et al. (Heidy et al., 2019), the investigators found a positive correlation between height and forearm length. The study by Sirajudeen et al. (Sirajudeen et al., 2012) found a positive correlation between hand length and handgrip strength. This shows that height affects the longer extremity or arm, thus approaching optimal muscle length that may result in greater grip strength. According to a prospective cohort study, handgrip strength in middle age appears to be associated with birth weight and the increase in height at puberty. Pubertal growth and handgrip strength possibly represent the development of muscle size and growth (Schrager et al., 2007).

Our study found no relationship between blood glucose concentration and handgrip strength. Our study agrees with the study of Giglio et al. (Giglio et al., 2018), which showed no significant relationship between muscle mass strength and blood glucose concentration. According to Giglio, this is because most respondents are individuals who routinely perform structured physical activity, even though they are in the light category. This is similar to our study respondents, who were public works personnel and who had performed physical activity such as street sweeping and cleaning storm drains. Glucose is the main fuel for muscle contraction, entering the muscle cells by diffusion through the GLUT4 glucose transporter, which, after muscle contraction, moves from its intracellular storage site to the plasma membrane and T-tubules (Richter et al., 2013). Glucose metabolism has an influence on handgrip strength through various mechanisms, such as disorders of glucose metabolism, particularly glycogenolysis, that may result in the loss of muscle strength. Insulin resistance in hyperglycemia may cause muscle degradation because hyperglycemia causes low muscle strength through effects on skeletal muscle mitochondria. The lowering of muscle strength by hyperglycemia occurs through Krüppel-like factor 15, which regulates skeletal muscle lipid flux. Hyperglycemia also increases pro-inflammatory cytokines, which may increase the catabolism that plays a role in the further decrease in muscle mass and quality (Gamayanti et al., 2023). The amount of carbohydrates also influences the glycemic response and post-prandial insulin level, impacting the supply of glycogen to the muscles (Safitri et al., 2020).

This study has a significant negative correlation between age and handgrip strength, which decreases with age (Manoharan et al., 2015). Handgrip strength initially experiences an increase with increasing age, reaches its peak between the age of 30 and 40 years, and decreases afterward (Zaccagni et al., 2020). The age-related changes in skeletal muscle composition, particularly the accumulation of lipids in skeletal muscle fibers, play a role in the poor quality of the muscles and may result in metabolic abnormalities such as insulin resistance. Apart from changes in muscle

quality, there is also a reduction in muscle mass that is consistently occurring during aging. The increase in oxidative stress and chronic inflammation that commonly occurs in the aging process apparently also plays a role in the development of decreased muscle mass and strength (Mainous et al., 2015). The decrease in muscle mass in the elderly corresponds to a 10% reduction in grip strength per decade, starting from the age of 40 years (Jaafar et al., 2023). Skeletal muscle plays an important role in glucose metabolism and has a significant influence on insulin sensitivity. Increased insulin resistance in older adults may be closely associated with skeletal muscle aging. A reduction in GLUT4 expression as a result of a decrease in muscle volume causes a reduction in insulin sensitivity in aging skeletal muscle (Joo et al., 2022).

The results of this study showed a correlation of waist circumference and hip circumference with handgrip strength. This is in line with the study of Widjaja et al. (Widjaja et al., 2018), who stated that there was a significant correlation of waist circumference and hip circumference with handgrip strength ($p < 0.001$, $r = 0.302$; and $p < 0.001$, $r = 0.296$). A greater waist circumference is associated with a greater handgrip strength in males (Hardy et al., 2013).

In the multiple regression analysis of HGS, we removed the data on body weight because of the occurrence of multicollinearity between weight and BMI, with a correlation coefficient of more than 0.9, indicating an extremely strong relationship between these two variables. Conceptually, only one of the variables is used in multivariate analysis. The multiple regression analysis included age, height, BMI, waist circumference, and hip circumference, with HGS as dependent variable in a stepwise dependent manner. After three iterations, only height, BMI, and hip circumference played a role in predicting HGS by means of the regression formula, which resulted in $R^2 = 0.19$. This signifies that for known values of height, BMI, and hip circumference, we can predict the HGS value 19% more accurately than without a knowledge of these data. For a clinical diagnosis, a minimum increase of 10% is needed. This study found that height, BMI, and hip circumference play a role in predicting handgrip strength through the regression formula, i.e., $\text{handgrip strength} = -36.98 + 0.48 \times \text{height} + 0.79 \times \text{BMI} - 0.29 \times \text{hip circumference}$. Handgrip strength and waist circumference are surrogate muscle strength and visceral adiposity markers, respectively (Nakanishi et al., 2024). Waist circumference is also more sensitive in determining body fat distribution, particularly in the abdomen. Waist circumference correlates better with abdominal fat distribution than BMI (Rokhmah et al., 2015). Waist circumference and hip circumference are associated with obesity, in which the fat content is higher, and the skeletal muscle and fascicular content is lower, thereby resulting in a lower contractile capacity of skeletal muscle (Afaq et al., 2019).

The limitation of this study is that it did not evaluate other factors that may affect handgrip strength, such as nutrient intake. From the anthropometric measures of the hands and feet in this study, the causes and effects of the decrease in handgrip strength could not be determined because this study used a cross-sectional design.

CONCLUSION

There are three factors that may be used to estimate handgrip strength in PWP, namely height, BMI, and hip circumference, whereas blood glucose concentration is not correlated with handgrip strength in PWP. It is recommended to pay attention to height, BMI, and hip circumference,

particularly in the acceptance criteria of PWP candidates, and to conduct further studies with consideration of other factors that affect handgrip strength, such as food intake and anthropometric features of the hands and arms.

ACKNOWLEDGMENTS

Financial support was provided by Universitas Trisakti for this research.

REFERENCES

- Afaq, E., Hassan, S.H., Khan, M.R., Zaidi, D.A., & Baktashi, H., 2019. Correlation of Hand Dynamometry With Body Mass Index and Waist Hip Ratio in Medical Students. *Pak J Physiol*, 15, pp.19-21.
- Agtuahene, M.A., Quartey, J., & Kwakye, S., 2023. Influence of Hand Dominance, Gender, and Body Mass Index on Hand Grip Strength. *S Afr J Physiother*, 79, pp.1-6. doi: org/10.4102/sajp.v79i1.1923.
- Al-Asadi, J.N., 2018. Handgrip Strength in Medical Students: Correlation with Body Mass Index and Hand Dimensions. *Asian Journal of Medical Sciences*, 9, pp.21-26. doi: 10.3126/ajms.v9i1.18577.
- Badan Pusat Statistik Kota Administrasi Jakarta Barat., 2020. Kecamatan Cengkareng Dalam Angka, pp. 1-114.
- Chen, D., Zhu, Y., Ni, W., Li, Y., Yin, G., Shao, Z., & Zhu, J., 2023. Hand Grip Strength is Inversely Associated with Total Daily Insulin Dose Requirement in Patients with Type 2 Diabetes Mellitus: A Cross-Sectional Study. *PeerJ*, pp.1-13. doi: 10.7717/peerj.15761.
- Cooper, R., Tomlinson, D., Hamer, M., & Pereira, S.M.P., 2022. Lifetime Body Mass Index and Grip Strength at Age 46 Years: The 1970 British Cohort Study. *J Cachexia Sarcopenia Muscle*, 13, pp.1-10. doi: 10.1002/jcsm.12992.
- Fauzi, L., Handayani, O.W.K., Susilo, M.T., Kurnia, A.R., Rahayu, S.R., Irawan, F.A., Horng Lu, F.J., Lin, C., Lai, M.F., & Yu, Y.C., 2022. Obesity in Indonesian and Taiwanese Adolescents Related to Self Perception, Diet, Exercise and Body Image. *KEMAS*, 17(3):pp.453-461. doi: 10.15294/kemas.v17i3.34396
- Gamayanti, K.A.V., Aryana, I.G.P.S., & Gotera, W., 2023. Hubungan Antara Kekuatan Genggaman Tangan Dengan Kadar Gula Darah Sewaktu Pada Lansia Di Desa Melinggih, Kecamatan Payangan, Kabupaten Gianyar. *Intisari Sains Medis*, 14(2): pp.483-488.
- Giglio, B.M., Mota, J.F., Wall, B.T., & Pimentel, G.D., 2018. Low Handgrip Strength is Not Associated With Type 2 Diabetes Mellitus and Hyperglycemia: A Population-Based Study. *Clin Nutr Res*, 7(2): pp.112-116. doi: 10.7762/cnr.2018.7.2.112.

Commented [MA3]: - Daftar pustaka minimal 20.
- Minimal 80% daftar pustaka adalah ARTIKEL JURNAL INTERNASIONAL
- Pastikan daftar pustaka artikel ada: nama penulis, tahun, judul artikel, nama jurnal, no/volume
- tambahkan 5 sitasi dari jurnal kemas UNNES

Alvina:
-Sudah ditambahkan 5 sitasi dari jurnal KEMAS UNNES

Hammed, A.I., & Obaseki, C.O., 2017. Interdependence of Body Mass Index with Handgrip Strength and Endurance Among Apparently Healthy Teenagers. *Turk J Kin*, 4, pp.1-7.

Handayani, M.D.N., Sadewa, A.H., Farmawati, A., & Rochmah, W., 2018. Anthropometric Prediction Equations for Estimating Muscle Mass of Elderly Women. *KEMAS*, 14(2):pp.195-204. doi: 10.15294/kemas.v14i2.14073

Hardy, R., Cooper, R., Sayer, A.A., Shlomo, Y.B., Cooper, C., Deary, I.J., Demakakos, P., Gallacher, J., Martin, R.M., McNeill, G., et al., 2013. Body Mass Index, Muscle Strength and Physical Performance in Older Adults From Eight Cohort Studies: The Halcyon Programme. *PLoS One*, 8(2): pp.e56483. doi: 10.1371/journal.pone.0056483.

Heidy., Djuartina, T., & Irawan, R., 2019. Korelasi Kekuatan Genggaman Tangan Dengan Karakter Antropometri Lengan Bawah dan Tangan Serta Indeks Massa Tubuh. *Damianus Journal of Medicine*, 18, pp.1-7.

Ilmi, A.F., & Utari, D.M., 2020. Hubungan Lingkar Pinggang dan Rasio Lingkar Pinggang-Panggul (RLPP) Terhadap Kadar Glukosa Darah Puasa Pada Mahasiswa. *Journal of Nutrition College*, 9, pp.222-227.

Jaafar, M.H., Ismail, R., Ismail, N.H., Isa, Z.M., Tamil, A.M., Nasir, N.M., Keat, N.K., Razak, N.H., Abidin, N.Z., & Yusof, K.H., 2023. Normative Reference Values and Predicting Factors of Handgrip Strength For Dominant and Non-Dominant Hands Among Healthy Malay Adults in Malaysia. *BMC Musculoskelet Disord*, 24, pp. 1-9. doi: 10.1186/s12891-023-06181-8.

Joo, K.C., Son, D.H., & Park, J.M., 2022. Association Between Relative Handgrip Strength and Insulin Resistance in Korean Elderly Men Without Diabetes: Findings Of The 2015 Korea National Health Nutrition Examination Survey. *Korean J Fam Med*, 43, pp.199-205. doi:10.4082/kjfm.21.0138.

Krakauer, N.Y., & Krakauer, J.C., 2020. Association of Body Shape Index (ABSI) with Hand Grip Strength. *Int J Environ Res Public Health*, 17, pp.1-12. doi: doi:10.3390/ijerph17186797.

Kurniawan, A., Widjaja, D., & Lugito, N.P.H., 2018. Mean and Cutoff Value of Hand Grip Strength For Healthy Indonesian Young Adults. *World Congress on Osteoporosis, Osteoarthritis and Musculoskeletal Diseases*; Krakow, Poland.

Leong, D.P., Teo, K.K., Rangarajan, S., Kutty, R., Lanas, F., Hui, C., Quanyong, X., Zhenzhen, Q., Jinhua, T., Noorhassim, I., et al., 2016. Reference Ranges of Handgrip Strength From 125,462 Healthy Adults in 21 Countries: A Prospective Urban Rural Epidemiologic (PURE) Study. *J Cachexia Sarcopenia Muscle*, 7, pp.535-546. doi: 10.1002/jcsm.12112.

Mainous, A.G., Tanner, R.J., Anton, S.D., & Jo, A., 2015. Grip Strength as A Marker of Hypertension and Diabetes in Healthy Weight Adults. *Am J Prev Med*, 49(6): pp.850-858. doi: 10.1016/j.amepre.2015.05.025.

Manoharan, V.S., Sundaram, S.G., & Jason, J.I., 2015. Factors Affecting Hand Grip Strength and Its Evaluation: A Systemic Review. *Int J Physiother Res*, 3(6): pp.1288-1293. doi:10.16965/ijpr.2015.193.

McGrath, R., Johnson, N., Klawitter, L., Mahoney, S., Trautman, K., Carlson, C., Rockstad, E., & Hackney, K.J., 2020. What Are The Association Patterns Between Handgrip Strength and Adverse Health Conditions? A Topical Review. *SAGE Open Medicine*, 8, pp.1-12. doi:10.1177/2050312120910358.

Nakanishi, S., Shimoda, M., Kimura, T., Sanada, J., Fushimi, Y., Iwamoto, Y., Iwamoto, H., Dan, K., Mune, T., Kaku, K., et al., 2024. The Impact of Handgrip Strength and Waist Circumference on Glycemic Control: Prospective, Observational Study Using Outpatient Clinical Data in Japanese Patients With Type 2 Diabetes Mellitus. *J Diabetes Investig*, 15(7): pp.892-898. doi: 10.1111/jdi.14200.

Nathania, N.P.S., Nugraha, M.H.S., Primayanti, I.D.A., & Saraswati, P.A.S., 2023. Handgrip Strength is Associated With Balance in Elderly Women. *Majalah Ilmiah Fisioterapi Indonesia*, 11, pp.70-75.

Peraturan Gubernur Provinsi Daerah Khusus Ibukota Jakarta Nomor 7 tahun 2017 Tentang Penanganan Prasarana dan Sarana Umum Tingkat Kelurahan. Jakarta 2017.

Richter, E.A., & Hargreaves, M., 2013. Exercise, GLUT4, and Skeletal Muscle Glucose Uptake. *Physiol Rev*, 93(3): pp.993-1017. doi: 10.1152/physrev.00038.2012.

Rokhmah, F.D., Handayani, D., & Al-Rasyid, H., 2015. Korelasi Lingkar Pinggang dan Rasio Lingkar Pinggang-Panggul Terhadap Kadar Glukosa Plasma Menggunakan Tes Toleransi Glukosa Oral. *Jurnal Gizi Klinik Indonesia*, 12, pp.28-35.

Rondhianto., Ridla, A.Z., & Hasan, H., 2024. Sociodemographic Factors Affecting Diabetic Dietary Behavior in People with Type 2 Diabetes Mellitus. *KEMAS*, 19(3):pp.429-437. doi: 10.15294/kemas.v19i3.37856

Safitri, I., Setyarsih, L., Susanto, H., Suhartono., & Fitranti, D.Y., 2020. The Effect of Low and High Glycemic Load Diet on Muscle Fatigue of Young Soccer. *KEMAS*, 16(1):pp.111-120. doi: 10.15294/kemas.v16i1.23508

Schrager, M.A., Metter, E.J., Simonsick, E., Ble, A., Bandinelli, S., Lauretani, F., & Ferrucci, L., 2007. Sarcopenic Obesity and Inflammation in The Inchiatti Study. *J Appl Physiol*, 102(3): pp.919-925. doi: 10.1152/japplphysiol.00627.2006.

Sirajudeen, M.S., Shah, U.N., Pillai, P.S., Mohasin, N., & Shantaram, M., 2012. Correlation Between Grip Strength and Physical Factors in Men. *International Journal of Health and Rehabilitation Sciences*, 1, pp.58-63.

Soenyoto, T., Mollap, K., & Mungpong, J., 2017. Correlation of Energy and Protein Consumption Levels with Physical Endurance of Rhythmic Gymnast Athletes. KEMAS, 13(1):pp.131-136. doi:10.15294/kemas.v13i1.9888

Soraya, N., & Parwanto, E., 2023. The Controversial Relationship Between Body Mass Index and Handgrip Strength in The Elderly: An Overview. Malays J Med Sci, 30, pp.73-83.

Vaishya, R., Misra, A., Vaish, A., Ursino, N., & D'Ambrosi, R., 2024. Hand Grip Strength as A Proposed New Vital Sign Of Health: A Narrative Review of Evidences. Journal of Health, Population and Nutrition, 43, pp.1-14. doi:10.1186/s41043-024-00500-y.

Widjaja, D., Tjhai, K., Dermawan, K., Kalim, J., Tanzil, S., Pramusinta, A., Phenca, H.K., Kurniawan, A., 2018. Correlations Between Hand Grip Strength and Anthropometric Status in Young Indonesian Adults. World Congress on Osteoporosis, Osteoarthritis and Musculoskeletal Diseases; Krakow, Poland.

Zaccagni, L., Toselli, S., Bramanti, B., Gualdi-Russo, E., Mongillo, J., & Rinaldo, N., 2020. Handgrip Strength in Young Adults: Association With Anthropometric Variables and Laterality. Int J Environ Res Public Health, 17, pp.1-18. doi:10.3390/ijerph17124273.

4. Bukti konfirmasi review dan hasil review kedua (21 Juli 2025)



Alvina fk <dr.alvina@trisakti.ac.id>

Discussion (Submission)

2 messages

Oktia woro kasmini handayani <oktia2016@mail.unnes.ac.id>
To: Alvina Alvina <dr.alvina@trisakti.ac.id>

Mon, Jul 21, 2025 at 8:08 AM

Please enter your message.

Reply to this comment at [#25917 Alvina et al.](#) or [unsubscribe](#) from emails sent by [Jurnal Kesehatan Masyarakat](#).



Revision-(1)+Alvina-template-kemas+(Revisi)-Factor+associated+with+handgripstrength (1).docx
48K

Alvina fk <dr.alvina@trisakti.ac.id>

To: Oktia woro kasmini handayani <oktia2016@mail.unnes.ac.id>

Mon, Jul 21, 2025 at 12:35 PM

Dear,
Chief Editor Jurnal Kemas
I have sent revision my article.
Thank you.

Sincerely,
Alvina

[Quoted text hidden]

Factors Associated With Handgrip Strength of Public Works Personnel in West Jakarta

Alvina^{1,a)}, Pusparini^{2,b)}, Mario^{3,c)}, Yasmine Mashabi^{4,d)}

^{1,2,3,4} Department of Clinical Pathology, Faculty of Medicine, Universitas Trisakti

^{a)}Corresponding author: dr.alvina@trisakti.ac.id

^{b)}pusparini@trisakti.ac.id

^{c)}mario@trisakti.ac.id

^{d)} yasminedrp@gmail.com

Abstract

Public Works Personnel (PWP) in Jakarta perform their routine duties manually instead of using equipment. Handgrip strength (HGS) measures maximum hand strength as a quality indicator for muscle strength and mass. Factors affecting muscle strength are age, sex, body mass index (BMI), waist circumference (WC), and hip circumference (HC). Increased fasting blood glucose (FBG) is also associated with muscle quality, muscle strength, and physical performance. This study aimed to determine the factors associated with handgrip strength in PWP. A total of 192 male PWP from the Cengkareng district were recruited by simple random sampling. Collected data were demographics, BMI, FBG, WC, HC, and HGS. Statistical analysis was done using Pearson's correlation test and multiple regression analysis. There was a significant positive correlation of BMI, WC, and HC with HGS ($p=0.006$; $r=0.20$, $p=0.009$; $r=0.19$ and $p=0.005$; $r=0.20$) and a significant negative correlation between age and HGS ($p=0.008$; $r=-0.19$) but not between FBG and HGS ($p=0.847$). Multiple regression analysis showed height, BMI, and HC positively predicting HGS ($R^2=0.19$). Handgrip strength of public works personnel is associated with BMI, age, waist circumference, and hip circumference, but not with fasting blood glucose. Height, BMI, and hip circumference predict handgrip strength.

Keywords: Handgrip strength, Body mass index, Hip circumference, Waist circumference, Fasting blood glucose

INTRODUCTION

Personnel for the management of public infrastructure and facilities at the village level are workers who manage village public works for a given time period on the basis of a work order. The tasks of village-level public works personnel (PWP) consist of managing the infrastructure and facilities, such as streets, drains, and parks, including installed lights (Peraturan Gubernur DKI Jakarta, 2017). The work performed by village PWP consists mainly of intense physical activity involving the hands, such that handgrip strength is essential in their tasks. Lowering of handgrip strength may indicate reduced muscle strength, which may especially interfere with the daily activities of village PWP that consist mainly of manual labor.

Handgrip strength measures maximal hand strength and indicates muscle quality that reflects muscle strength and mass. Muscle strength refers to the ability of muscle groups to raise a given load. Low or poor grip strength is frequently associated with damaging health impacts, such as chronic disease, and is the initial sign of reduced muscle mass, limitations in physical functions, and mortality risk. Poor handgrip strength may also reduce a person's ability to perform activities of daily living and, therefore, impact quality of life (Soraya et al., 2023; Nathania et al., 2023). Muscle mass accounts for approximately 40% of total weight, such that knowledge of muscle mass conditions is essential in maintaining health and activities (Handayani et al., 2018).

Handgrip strength may represent the strength of all muscles in the body. Several factors possibly influencing an individual's muscle strength are age, sex, and body mass index (BMI) (Soraya et al., 2023). Obesity may also affect BMI. Obesity in adults is not only associated with genetic factors but also with environmental factors during development (Fauzi et al., 2022). Handgrip strength is also associated with chronic diseases such as diabetes, metabolic syndrome, and cardiovascular disease. Increased glucose concentration is also associated with disorders of muscle quality and strength, as well as physical performance (Chen et al., 2023). A reduction in muscle mass also affects an individual's strength and physical endurance. A combination of proteins is considered to be able to maintain muscle function, thereby building body strength and physical endurance (Soenyoto et al., 2017). In the present study, we included fasting blood glucose concentrations in the data collection in order to obtain more accurate measures of glucose concentrations that have not yet been influenced by food intake.

Handgrip strength may be a health indicator that provides information on a person's health (Vaishya et al., 2024). However, its clinical significance in determining the prevention and treatment of poor handgrip remains limited because of the still unclear etiology of muscle weakness, especially because the clinical outcomes associated with handgrip strength are extremely diverse (Mc Grath et al., 2020). The relationship between BMI and handgrip strength is still controversial, with several investigators finding a significant positive association and others reporting a non-significant relationship (Al Asadi, 2018).

There are few studies on handgrip strength, especially on workers in the sanitation sector such as public works personnel, such that the present investigators were interested in studying this topic. The aim of the present study was to determine the factors associated with handgrip strength in public works personnel.

METHOD

This cross-sectional study involving male public works personnel in Cengkareng District, West Jakarta, was conducted from September to October 2024. The study sample was selected by simple random sampling. Calculation of the sample size for a correlation with $r=0.2$, $Z\alpha=0.05$, and $Z\beta=0.1$, resulted in a sample of 192 subjects who met the inclusion criterion of being male and 25-60 years old, and the exclusion criteria of having an abnormality of the hands, heart disease, pulmonary disease, and malignancies.

Measurements were obtained on height in cm, weight in kg, as well as waist circumference and hip circumference in cm. Height and weight were determined using microtoise and digital scales, respectively. BMI was calculated using the formula of weight divided by height squared. Measurement of waist and hip circumferences was done using measuring tape. Waist circumference was determined with the subject in the upright position, at the end of a normal expiration, at the smallest abdominal circumference between the iliac crest and twelfth rib. The cut-off values for waist circumference in males and females are ≥ 90 cm and ≥ 80 cm, respectively. Hip circumference was measured using measuring tape that was applied around the hips at the level of the symphysis pubis and the maximal gluteal circumference (Rokhmah et al., 2015; Ilmi et al., 2020).

Fasting blood glucose (FBG) concentration was determined on venous blood by the hexokinase method after the respondents had fasted approximately 10 hours prior. Handgrip strength was measured with a handgrip strength dynamometer to the nearest kilogram as follows: The respondent was asked to sit down comfortably without reclining and with the feet completely relaxed. The hips and knees were positioned at a 90° angle, the gripping hand with the shoulder in adduction, the elbow flexed 90° , and the forearm and wrist in the neutral position. The gripping duration was 3 seconds. The respondent gripped the instrument with the dominant hand as hard as possible and the value indicated by the needle was recorded. In accordance with the protocol recommended by the American Society of Hand Therapists (ASHT), the recorded handgrip strength was the average of three handgrip strength tests (Nathania et al., 2023).

Prior to data analysis by means of the correlation test, the normality of the data distribution was determined using the Kolmogorov-Smirnov test. The Pearson correlation test was used for a normal data distribution, whereas for a non-normal data distribution, the Spearman correlation test was done, followed by a multiple regression analysis.

Ethics approval was issued by the Ethics Committee of the Faculty of Medicine, Universitas Trisakti, under No. 033/KER/FK/08/2024.

RESULT AND DISCUSSION

In this study the respondents comprised 192 male public works personnel of Cengkareng District that consisted of the villages (subdistricts) of Kedaung, Duri Kosambi, West Cengkareng, Kapuk, and Rawa Buaya. A total of 26% of the respondents were residents of Duri Kosambi, where Muslims comprised 98.4% of the population. The majority of the respondents at 77.6% had an educational level of senior high school, nearly all of the respondents, or 99%, were right-handed, using the right hand as the dominant one at work, and most were of the Betawi (Jakarta) ethnic group at 51.6% (Table 1).

Table 1. Demographic data of the respondents

Characteristic	N (%)
Village	
Kedaung	40 (20.8)
Duri Kosambi	50 (26)
Cengkareng Barat	33 (17.2)
Kapuk	43 (22.4)
Rawa Buaya	26 (13.5)
Religion	
Islamic	189 (98.4)
Christian	2 (1)
Catholic	1 (0.5)
Level of education	
Elementary school	7 (3.6)
Junior high school	29 (15.1)
Senior high school	149 (77.6)
Academy-University	7 (3.6)
Dominant hand	
Right hand	190 (99)
Left hand	2 (1)
Ethnic group	
Javanese	53 (27.6)
Betawi	99 (51.6)
Sundanese	25 (13)
Other	14 (7.3)

Legend: Other ethnicities comprised groups from Palembang, Lampung, Bima, and Kupang

Source: primary data 2024 (questionnaires and interviews with respondents)

Commented [MA1]: Dibawah setiap tabel sebutkan sumbernya. Silahkan mencermati aartikel yang sudah terbit di Jurnal KEMAS

Alvina: sudah ditambahkan sumber data dibawah setiap tabel

Commented [MA2R1]: Primari data, th.....

Alvina: sudah ditambahkan tahun primary data

Table 2. Respondents' characteristics

Variable	Mean $\pm$ SD
Age (years)	40.9 $\pm$ 8.9
Height (cm)	164.6 $\pm$ 6.7
Weight (kg)	64.7 $\pm$ 13.1
BMI (kg/m ²)	23.8 $\pm$ 4.3
Waist circumference (cm)	85.4 $\pm$ 10.4
Hip circumference (cm)	96.9 $\pm$ 8.4
HGS (kg)	33.1 $\pm$ 6.9
FBG (mg/dL)	87 (64-280)*

Legend: BMI: body mass index. *FBG: fasting blood glucose, expressed as Median (Min-Max).

HGS: handgrip strength

Source: primary data (respondents' measurements)

The bivariate analysis found a significant positive correlation of BMI, height, weight, waist circumference, and hip circumference with handgrip strength ($p=0.006$, $r=0.20$; $p<0.001$, $r=0.36$; $p<0.001$, $r=0.32$; $p=0.009$, $r=0.19$; $p=0.005$, $r=0.20$, respectively). This showed that the larger the BMI, height, weight, waist circumference, and hip circumference, the stronger the handgrip. However, there was no significant relationship between fasting glucose concentration and handgrip strength ($p=0.847$). This study also found that age had a significant negative correlation with handgrip strength ($p=0.011$; $r=-0.182$), signifying that the older a person, the lower the handgrip strength (Table 3). Results of multiple regression analysis on handgrip strength are listed in Table 4.

Table 3. Analysis of the relationship of age, height, weight, BMI, waist circumference, hip circumference, and fasting blood glucose concentration with handgrip strength

Variable	R	P value
Age	- 0.19	0.008*
Height	0.36	0.000*
Weight	0.32	0.000*
BMI	0.20	0.006*
Waist circumference	0.19	0.009*
Hip circumference	0.20	0.005*
FBG	0.02	0.847 [#]

*P value with Pearson's test, [#] Spearman's test

Source: primary data

Table 4. Multiple regression analysis on handgrip strength

Determinant	b	SE	p
Height	0.48	0.08	0.000
BMI	0.79	0.22	0.000
Hip circumference	- 0.29	0.12	0.015
Constant	- 36.98		

R²=0,19

Source: primary data

This study found that most of the public works personnel in Duri Kosambi village were Muslims, had a senior high school level of education, and were of Betawi ethnicity. This was in line with the demographic data in Cengkareng district, in that Duri Kosambi was the village with the greatest area, the majority of the residents were male, and the Islamic religion had the most believers (Badan Pusat Statistik Jakarta Barat, 2020). Among our respondents, more than ten percent were in the diabetes category. Socio-demographic factors capable of affecting dietary habits in patients with diabetes are level of education and type of family (Rondhianto et al., 2024). Among the respondents of this study, the dominant right hand was used for working. In the adult population, a total of 90% of the residents preferred to use the right hand for manual tasks, whereas around 10% used the left hand (Zaccagni et al., 2020). According to Agtuahene et al. (Agtuahene et al., 2023), the dominant hand is generally around 10% stronger than the non-dominant hand. Many activities in daily life need a high level of exertion of the muscles of the hand and forearm (Hammed et al., 2017). The frequent use of the dominant hand in daily living results in the muscles of the hand and forearm providing strength in gripping (Afaq et al., 2019).

The mean handgrip strength in this study was 33.1 ± 6.9 kg, which is relatively low if compared with the study results of Kurniawan et al. (Kurniawan et al., 2018). in healthy young Indonesian adults, where mean handgrip strength in males was 37.37 ± 8.29 kg and the lower limit of muscle strength in Indonesian males was 20.79 kg. The handgrip strength results in both studies are still below those of males in other Southeast Asian countries, according to the study results of Leong et al. (Leong et al., 2016). These investigators found that the mean handgrip strength in Southeast Asian males aged 35-40 years was 40 kg (34-44 kg); for the age range of 41-50 years, the handgrip strength was 38 kg (33-44 kg), for the age range of 51-60 years it was 34 kg (30-40 kg), and for ages 61-70 years 30 kg (24-34 kg). This may have been caused by ethnic differences mediated by genetic differences that also affect individual muscle strength. The highest handgrip strength is found in the European and North American populations, whereas the lowest handgrip strength is found in African, South Asian, and Southeast Asian populations. These variations in muscle strength in various countries may also be partly caused by differences in socio-economic status (Leong et al., 2016).

According to Agtuahene et al. (Agtuahene et al., 2023), handgrip strength is a physiological variable influenced by several factors, such as age, sex, and body measurements. In contrast, according to Zaccagni et al. (Zaccagni et al., 2020), handgrip strength, apart from being affected by age, may also be influenced by physical activity, grip measurement, height, and body mass. The present study found a significant positive correlation between BMI and handgrip strength. According to Hammed et al., this may be caused by the higher percentage of skeletal muscle mass compared to the percentage of fat mass, thereby resulting in a higher handgrip strength [13]. Our study's results are identical to those of Agtuahene et al. (Agtuahene et al., 2023), who found a

significant positive correlation ($r = 0.290$) of handgrip strength with BMI and height. This shows that taller individuals with higher BMI apparently have greater handgrip strength. Our study is also in line with the study of Krakauer in 2020, which showed a positive correlation between BMI and handgrip strength (Krakauer et al., 2020).

The study of Cooper et al. (Cooper et al., 2022) concluded that a high childhood BMI is associated with higher grip strength at age 46 years. This is because fat mass acts as a mechanical load that elicits an anabolic response that stimulates muscle growth and function. This anabolic response is usually greater in males than in females because of the higher testosterone concentration in males. However, it should also be noted that BMI does not differentiate between fat mass and lean mass as adipocyte indicators that may vary with sex. It should be noted that BMI is a measure of body fat that does not differentiate between weight changes as a result of an increase or decrease in muscle mass or body fat (Zaccagni et al., 2020). According to Jafar et al. (Jaafar et al., 2023), heavier individuals have a greater muscle mass, resulting in greater handgrip strength, than lighter individuals. In the study of Heidy et al. (Heidy et al., 2019), the investigators found a positive correlation between height and forearm length. The study by Sirajudeen et al. (Sirajudeen et al., 2012) found a positive correlation between hand length and handgrip strength. This shows that height affects the longer extremity or arm, thus approaching optimal muscle length that may result in greater grip strength. According to a prospective cohort study, handgrip strength in middle age appears to be associated with birth weight and the increase in height at puberty. Pubertal growth and handgrip strength possibly represent the development of muscle size and growth (Schrager et al., 2007).

Our study found no relationship between blood glucose concentration and handgrip strength. Our study agrees with the study of Giglio et al. (Giglio et al., 2018), which showed no significant relationship between muscle mass strength and blood glucose concentration. According to Giglio, this is because most respondents are individuals who routinely perform structured physical activity, even though they are in the light category. This is similar to our study respondents, who were public works personnel and who had performed physical activity such as street sweeping and cleaning storm drains. Glucose is the main fuel for muscle contraction, entering the muscle cells by diffusion through the GLUT4 glucose transporter, which, after muscle contraction, moves from its intracellular storage site to the plasma membrane and T-tubules (Richter et al., 2013). Glucose metabolism has an influence on handgrip strength through various mechanisms, such as disorders of glucose metabolism, particularly glycogenolysis, that may result in the loss of muscle strength. Insulin resistance in hyperglycemia may cause muscle degradation because hyperglycemia causes low muscle strength through effects on skeletal muscle mitochondria. The lowering of muscle strength by hyperglycemia occurs through Krüppel-like factor 15, which regulates skeletal muscle lipid flux. Hyperglycemia also increases pro-inflammatory cytokines, which may increase the catabolism that plays a role in the further decrease in muscle mass and quality (Gamayanti et al., 2023). The amount of carbohydrates also influences the glycemic response and post-prandial insulin level, impacting the supply of glycogen to the muscles (Safitri et al., 2020).

This study has a significant negative correlation between age and handgrip strength, which decreases with age (Manoharan et al., 2015). Handgrip strength initially experiences an increase with increasing age, reaches its peak between the age of 30 and 40 years, and decreases afterward (Zaccagni et al., 2020). The age-related changes in skeletal muscle composition, particularly the accumulation of lipids in skeletal muscle fibers, play a role in the poor quality of the muscles and may result in metabolic abnormalities such as insulin resistance. Apart from changes in muscle

quality, there is also a reduction in muscle mass that is consistently occurring during aging. The increase in oxidative stress and chronic inflammation that commonly occurs in the aging process apparently also plays a role in the development of decreased muscle mass and strength (Mainous et al., 2015). The decrease in muscle mass in the elderly corresponds to a 10% reduction in grip strength per decade, starting from the age of 40 years (Jaafar et al., 2023). Skeletal muscle plays an important role in glucose metabolism and has a significant influence on insulin sensitivity. Increased insulin resistance in older adults may be closely associated with skeletal muscle aging. A reduction in GLUT4 expression as a result of a decrease in muscle volume causes a reduction in insulin sensitivity in aging skeletal muscle (Joo et al., 2022).

The results of this study showed a correlation of waist circumference and hip circumference with handgrip strength. This is in line with the study of Widjaja et al. (Widjaja et al., 2018), who stated that there was a significant correlation of waist circumference and hip circumference with handgrip strength ($p < 0.001$, $r = 0.302$; and $p < 0.001$, $r = 0.296$). A greater waist circumference is associated with a greater handgrip strength in males (Hardy et al., 2013).

In the multiple regression analysis of HGS, we removed the data on body weight because of the occurrence of multicollinearity between weight and BMI, with a correlation coefficient of more than 0.9, indicating an extremely strong relationship between these two variables. Conceptually, only one of the variables is used in multivariate analysis. The multiple regression analysis included age, height, BMI, waist circumference, and hip circumference, with HGS as dependent variable in a stepwise dependent manner. After three iterations, only height, BMI, and hip circumference played a role in predicting HGS by means of the regression formula, which resulted in $R^2 = 0.19$. This signifies that for known values of height, BMI, and hip circumference, we can predict the HGS value 19% more accurately than without a knowledge of these data. For a clinical diagnosis, a minimum increase of 10% is needed. This study found that height, BMI, and hip circumference play a role in predicting handgrip strength through the regression formula, i.e., $\text{handgrip strength} = -36.98 + 0.48 \times \text{height} + 0.79 \times \text{BMI} - 0.29 \times \text{hip circumference}$. Handgrip strength and waist circumference are surrogate muscle strength and visceral adiposity markers, respectively (Nakanishi et al., 2024). Waist circumference is also more sensitive in determining body fat distribution, particularly in the abdomen. Waist circumference correlates better with abdominal fat distribution than BMI (Rokhmah et al., 2015). Waist circumference and hip circumference are associated with obesity, in which the fat content is higher, and the skeletal muscle and fascicular content is lower, thereby resulting in a lower contractile capacity of skeletal muscle (Afaq et al., 2019).

The limitation of this study is that it did not evaluate other factors that may affect handgrip strength, such as nutrient intake. From the anthropometric measures of the hands and feet in this study, the causes and effects of the decrease in handgrip strength could not be determined because this study used a cross-sectional design.

CONCLUSION

There are three factors that may be used to estimate handgrip strength in PWP, namely height, BMI, and hip circumference, whereas blood glucose concentration is not correlated with handgrip strength in PWP. It is recommended to pay attention to height, BMI, and hip circumference,

particularly in the acceptance criteria of PWP candidates, and to conduct further studies with consideration of other factors that affect handgrip strength, such as food intake and anthropometric features of the hands and arms.

ACKNOWLEDGMENTS

Financial support was provided by Universitas Trisakti for this research.

REFERENCES

Afaq, E., Hassan, S.H., Khan, M.R., Zaidi, D.A., & Baktashi, H., 2019. Correlation of Hand Dynamometry With Body Mass Index and Waist Hip Ratio in Medical Students. *Pak J Physiol*, 15, pp.19-21.

Aguahene, M.A., Quartey, J., & Kwakye, S., 2023. Influence of Hand Dominance, Gender, and Body Mass Index on Hand Grip Strength. *S Afr J Physiother*, 79, pp.1-6. doi: [org/10.4102/sajp.v79i1.1923](https://doi.org/10.4102/sajp.v79i1.1923).

Al-Asadi, J.N., 2018. Handgrip Strength in Medical Students: Correlation with Body Mass Index and Hand Dimensions. *Asian Journal of Medical Sciences*, 9, pp.21-26. doi: [10.3126/ajms.v9i1.18577](https://doi.org/10.3126/ajms.v9i1.18577).

Badan Pusat Statistik Kota Administrasi Jakarta Barat., 2020. Kecamatan Cengkareng Dalam Angka, pp. 1-114.

Chen, D., Zhu, Y., Ni, W., Li, Y., Yin, G., Shao, Z., & Zhu, J., 2023. Hand Grip Strength is Inversely Associated with Total Daily Insulin Dose Requirement in Patients with Type 2 Diabetes Mellitus: A Cross-Sectional Study. *PeerJ*, pp.1-13. doi: [10.7717/peerj.15761](https://doi.org/10.7717/peerj.15761).

Cooper, R., Tomlinson, D., Hamer, M., & Pereira, S.M.P., 2022. Lifetime Body Mass Index and Grip Strength at Age 46 Years: The 1970 British Cohort Study. *J Cachexia Sarcopenia Muscle*, 13, pp.1-10. doi: [10.1002/jcsm.12992](https://doi.org/10.1002/jcsm.12992).

Fauzi, L., Handayani, O.W.K., Susilo, M.T., Kurnia, A.R., Rahayu, S.R., Irawan, F.A., Horng Lu, F.J., Lin, C., Lai, M.F., & Yu, Y.C., 2022. Obesity in Indonesian and Taiwanese Adolescents Related to Self Perception, Diet, Exercise and Body Image. *KEMAS*, 17(3):pp.453-461. doi: [10.15294/kemas.v17i3.34396](https://doi.org/10.15294/kemas.v17i3.34396)

Gamayanti, K.A.V., Aryana, I.G.P.S., & Gotera, W., 2023. Hubungan Antara Kekuatan Genggaman Tangan Dengan Kadar Gula Darah Sewaktu Pada Lansia Di Desa Melinggih, Kecamatan Payangan, Kabupaten Gianyar. *Intisari Sains Medis*, 14(2): pp.483-488.

Giglio, B.M., Mota, J.F., Wall, B.T., & Pimentel, G.D., 2018. Low Handgrip Strength is Not Associated With Type 2 Diabetes Mellitus and Hyperglycemia: A Population-Based Study. *Clin Nutr Res*, 7(2): pp.112-116. doi: [10.7762/cnr.2018.7.2.112](https://doi.org/10.7762/cnr.2018.7.2.112).

Hammed, A.I., & Obaseki, C.O., 2017. Interdependence of Body Mass Index with Handgrip Strength and Endurance Among Apparently Healthy Teenagers. *Turk J Kin*, 4, pp.1-7.

Handayani, M.D.N., Sadewa, A.H., Farmawati, A., & Rochmah, W., 2018. Anthropometric Prediction Equations for Estimating Muscle Mass of Elderly Women. *KEMAS*, 14(2):pp.195-204. doi: 10.15294/kemas.v14i2.14073

Hardy, R., Cooper, R., Sayer, A.A., Shlomo, Y.B., Cooper, C., Deary, I.J., Demakakos, P., Gallacher, J., Martin, R.M., McNeill, G., et al., 2013. Body Mass Index, Muscle Strength and Physical Performance in Older Adults From Eight Cohort Studies: The Halcyon Programme. *PLoS One*, 8(2): pp.e56483. doi: 10.1371/journal.pone.0056483.

Heidy., Djuartina, T., & Irawan, R., 2019. Korelasi Kekuatan Genggaman Tangan Dengan Karakter Antropometri Lengan Bawah dan Tangan Serta Indeks Massa Tubuh. *Damianus Journal of Medicine*, 18, pp.1-7.

Ilmi, A.F., & Utari, D.M., 2020. Hubungan Lingkar Pinggang dan Rasio Lingkar Pinggang-Panggul (RLPP) Terhadap Kadar Glukosa Darah Puasa Pada Mahasiswa. *Journal of Nutrition College*, 9, pp.222-227.

Jaafar, M.H., Ismail, R., Ismail, N.H., Isa, Z.M., Tamil, A.M., Nasir, N.M., Keat, N.K., Razak, N.H., Abidin, N.Z., & Yusof, K.H., 2023. Normative Reference Values and Predicting Factors of Handgrip Strength For Dominant and Non-Dominant Hands Among Healthy Malay Adults in Malaysia. *BMC Musculoskelet Disord*, 24, pp. 1-9. doi: 10.1186/s12891-023-06181-8.

Joo, K.C., Son, D.H., & Park, J.M., 2022. Association Between Relative Handgrip Strength and Insulin Resistance in Korean Elderly Men Without Diabetes: Findings Of The 2015 Korea National Health Nutrition Examination Survey. *Korean J Fam Med*, 43, pp.199-205. doi:10.4082/kjfm.21.0138.

Krakauer, N.Y., & Krakauer, J.C., 2020. Association of Body Shape Index (ABSI) with Hand Grip Strength. *Int J Environ Res Public Health*, 17, pp.1-12. doi: doi:10.3390/ijerph17186797.

Kurniawan, A., Widjaja, D., & Lugito, N.P.H., 2018. Mean and Cutoff Value of Hand Grip Strength For Healthy Indonesian Young Adults. *World Congress on Osteoporosis, Osteoarthritis and Musculoskeletal Diseases*; Krakow, Poland.

Leong, D.P., Teo, K.K., Rangarajan, S., Kutty, R., Lanas, F., Hui, C., Quanyong, X., Zhenzhen, Q., Jinhua, T., Noorhassim, I., et al., 2016. Reference Ranges of Handgrip Strength From 125,462 Healthy Adults in 21 Countries: A Prospective Urban Rural Epidemiologic (PURE) Study. *J Cachexia Sarcopenia Muscle*, 7, pp.535-546. doi: 10.1002/jcsm.12112.

Mainous, A.G., Tanner, R.J., Anton, S.D., & Jo, A., 2015. Grip Strength as A Marker of Hypertension and Diabetes in Healthy Weight Adults. *Am J Prev Med*, 49(6): pp.850-858. doi: 10.1016/j.amepre.2015.05.025.

Manoharan, V.S., Sundaram, S.G., & Jason, J.I., 2015. Factors Affecting Hand Grip Strength and Its Evaluation: A Systemic Review. *Int J Physiother Res*, 3(6): pp.1288-1293. doi:10.16965/ijpr.2015.193.

McGrath, R., Johnson, N., Klawitter, L., Mahoney, S., Trautman, K., Carlson, C., Rockstad, E., & Hackney, K.J., 2020. What Are The Association Patterns Between Handgrip Strength and Adverse Health Conditions? A Topical Review. *SAGE Open Medicine*, 8, pp.1-12. doi:10.1177/2050312120910358.

Nakanishi, S., Shimoda, M., Kimura, T., Sanada, J., Fushimi, Y., Iwamoto, Y., Iwamoto, H., Dan, K., Mune, T., Kaku, K., et al., 2024. The Impact of Handgrip Strength and Waist Circumference on Glycemic Control: Prospective, Observational Study Using Outpatient Clinical Data in Japanese Patients With Type 2 Diabetes Mellitus. *J Diabetes Investig*, 15(7): pp.892-898. doi: 10.1111/jdi.14200.

Nathania, N.P.S., Nugraha, M.H.S., Primayanti, I.D.A., & Saraswati, P.A.S., 2023. Handgrip Strength is Associated With Balance in Elderly Women. *Majalah Ilmiah Fisioterapi Indonesia*, 11, pp.70-75.

Peraturan Gubernur Provinsi Daerah Khusus Ibukota Jakarta Nomor 7 tahun 2017 Tentang Penanganan Prasarana dan Sarana Umum Tingkat Kelurahan. Jakarta 2017.

Richter, E.A., & Hargreaves, M., 2013. Exercise, GLUT4, and Skeletal Muscle Glucose Uptake. *Physiol Rev*, 93(3): pp.993-1017. doi: 10.1152/physrev.00038.2012.

Rokhmah, F.D., Handayani, D., & Al-Rasyid, H., 2015. Korelasi Lingkar Pinggang dan Rasio Lingkar Pinggang-Panggul Terhadap Kadar Glukosa Plasma Menggunakan Tes Toleransi Glukosa Oral. *Jurnal Gizi Klinik Indonesia*, 12, pp.28-35.

Rondhianto., Ridla, A.Z., & Hasan, H., 2024. Sociodemographic Factors Affecting Diabetic Dietary Behavior in People with Type 2 Diabetes Mellitus. *KEMAS*, 19(3):pp.429-437. doi: 10.15294/kemas.v19i3.37856

Safitri, I., Setyarsih, L., Susanto, H., Suhartono., & Fitranti, D.Y., 2020. The Effect of Low and High Glycemic Load Diet on Muscle Fatigue of Young Soccer. *KEMAS*, 16(1):pp.111-120. doi: 10.15294/kemas.v16i1.23508

Schrager, M.A., Metter, E.J., Simonsick, E., Ble, A., Bandinelli, S., Lauretani, F., & Ferrucci, L., 2007. Sarcopenic Obesity and Inflammation in The Inchiati Study. *J Appl Physiol*, 102(3): pp.919-925. doi: 10.1152/japplphysiol.00627.2006.

Sirajudeen, M.S., Shah, U.N., Pillai, P.S., Mohasin, N., & Shantaram, M., 2012. Correlation Between Grip Strength and Physical Factors in Men. *International Journal of Health and Rehabilitation Sciences*, 1, pp.58-63.

Soenyoto, T., Mollap, K., & Mungpong, J., 2017. Correlation of Energy and Protein Consumption Levels with Physical Endurance of Rhythmic Gymnast Athletes. KEMAS, 13(1):pp.131-136. doi:10.15294/kemas.v13i1.9888

Soraya, N., & Parwanto, E., 2023. The Controversial Relationship Between Body Mass Index and Handgrip Strength in The Elderly: An Overview. Malays J Med Sci, 30, pp.73-83.

Vaishya, R., Misra, A., Vaish, A., Ursino, N., & D'Ambrosi, R., 2024. Hand Grip Strength as A Proposed New Vital Sign Of Health: A Narrative Review of Evidences. Journal of Health, Population and Nutrition, 43, pp.1-14. doi:10.1186/s41043-024-00500-y.

Widjaja, D., Tjhai, K., Dermawan, K., Kalim, J., Tanzil, S., Pramusinta, A., Phenca, H.K., Kurniawan, A., 2018. Correlations Between Hand Grip Strength and Anthropometric Status in Young Indonesian Adults. World Congress on Osteoporosis, Osteoarthritis and Musculoskeletal Diseases; Krakow, Poland.

Zaccagni, L., Toselli, S., Bramanti, B., Gualdi-Russo, E., Mongillo, J., & Rinaldo, N., 2020. Handgrip Strength in Young Adults: Association With Anthropometric Variables and Laterality. Int J Environ Res Public Health, 17, pp.1-18. doi:10.3390/ijerph17124273.

5. Bukti konfirmasi submit revisi kedua, respon
kepada reviewer, dan artikel yang diresubmit
(21 Juli 2025)



Alvina fk <dr.alvina@trisakti.ac.id>

Discussion (Submission)

2 messages

Oktia woro kasmini handayani <oktia2016@mail.unnes.ac.id>
To: Alvina Alvina <dr.alvina@trisakti.ac.id>

Mon, Jul 21, 2025 at 8:08 AM

Please enter your message.

Reply to this comment at [#25917 Alvina et al.](#) or [unsubscribe](#) from emails sent by [Jurnal Kesehatan Masyarakat](#).



Revision-(1)+Alvina-template-kemas+(Revisi)-Factor+associated+with+handgripstrength (1).docx
48K

Alvina fk <dr.alvina@trisakti.ac.id>

To: Oktia woro kasmini handayani <oktia2016@mail.unnes.ac.id>

Mon, Jul 21, 2025 at 12:35 PM

Dear,
Chief Editor Jurnal Kemas
I have sent revision my article.
Thank you.

Sincerely,
Alvina

[Quoted text hidden]

Factors Associated With Handgrip Strength of Public Works Personnel in West Jakarta

Alvina^{1,a)}, Pusparini^{2,b)}, Mario^{3,c)}, Yasmine Mashabi^{4,d)}

^{1,2,3,4} Department of Clinical Pathology, Faculty of Medicine, Universitas Trisakti

^{a)}Corresponding author: dr.alvina@trisakti.ac.id

^{b)}pusparini@trisakti.ac.id

^{c)}mario@trisakti.ac.id

^{d)} yasminedrp@gmail.com

Abstract

Public Works Personnel (PWP) in Jakarta perform their routine duties manually instead of using equipment. Handgrip strength (HGS) measures maximum hand strength as a quality indicator for muscle strength and mass. Factors affecting muscle strength are age, sex, body mass index (BMI), waist circumference (WC), and hip circumference (HC). Increased fasting blood glucose (FBG) is also associated with muscle quality, muscle strength, and physical performance. This study aimed to determine the factors associated with handgrip strength in PWP. A total of 192 male PWP from the Cengkareng district were recruited by simple random sampling. Collected data were demographics, BMI, FBG, WC, HC, and HGS. Statistical analysis was done using Pearson's correlation test and multiple regression analysis. There was a significant positive correlation of BMI, WC, and HC with HGS ($p=0.006$; $r=0.20$, $p=0.009$; $r=0.19$ and $p=0.005$; $r=0.20$) and a significant negative correlation between age and HGS ($p=0.008$; $r=-0.19$) but not between FBG and HGS ($p=0.847$). Multiple regression analysis showed height, BMI, and HC positively predicting HGS ($R^2=0.19$). Handgrip strength of public works personnel is associated with BMI, age, waist circumference, and hip circumference, but not with fasting blood glucose. Height, BMI, and hip circumference predict handgrip strength.

Keywords: Handgrip strength, Body mass index, Hip circumference, Waist circumference, Fasting blood glucose

INTRODUCTION

Personnel for the management of public infrastructure and facilities at the village level are workers who manage village public works for a given time period on the basis of a work order. The tasks of village-level public works personnel (PWP) consist of managing the infrastructure and facilities, such as streets, drains, and parks, including installed lights (Peraturan Gubernur DKI Jakarta, 2017). The work performed by village PWP consists mainly of intense physical activity involving the hands, such that handgrip strength is essential in their tasks. Lowering of handgrip strength may indicate reduced muscle strength, which may especially interfere with the daily activities of village PWP that consist mainly of manual labor.

Handgrip strength measures maximal hand strength and indicates muscle quality that reflects muscle strength and mass. Muscle strength refers to the ability of muscle groups to raise a given load. Low or poor grip strength is frequently associated with damaging health impacts, such as chronic disease, and is the initial sign of reduced muscle mass, limitations in physical functions, and mortality risk. Poor handgrip strength may also reduce a person's ability to perform activities of daily living and, therefore, impact quality of life (Soraya et al., 2023; Nathania et al., 2023). Muscle mass accounts for approximately 40% of total weight, such that knowledge of muscle mass conditions is essential in maintaining health and activities (Handayani et al., 2018).

Handgrip strength may represent the strength of all muscles in the body. Several factors possibly influencing an individual's muscle strength are age, sex, and body mass index (BMI) (Soraya et al., 2023). Obesity may also affect BMI. Obesity in adults is not only associated with genetic factors but also with environmental factors during development (Fauzi et al., 2022). Handgrip strength is also associated with chronic diseases such as diabetes, metabolic syndrome, and cardiovascular disease. Increased glucose concentration is also associated with disorders of muscle quality and strength, as well as physical performance (Chen et al., 2023). A reduction in muscle mass also affects an individual's strength and physical endurance. A combination of proteins is considered to be able to maintain muscle function, thereby building body strength and physical endurance (Soenyoto et al., 2017). In the present study, we included fasting blood glucose concentrations in the data collection in order to obtain more accurate measures of glucose concentrations that have not yet been influenced by food intake.

Handgrip strength may be a health indicator that provides information on a person's health (Vaishya et al., 2024). However, its clinical significance in determining the prevention and treatment of poor handgrip remains limited because of the still unclear etiology of muscle weakness, especially because the clinical outcomes associated with handgrip strength are extremely diverse (Mc Grath et al., 2020). The relationship between BMI and handgrip strength is still controversial, with several investigators finding a significant positive association and others reporting a non-significant relationship (Al Asadi, 2018).

There are few studies on handgrip strength, especially on workers in the sanitation sector such as public works personnel, such that the present investigators were interested in studying this topic. The aim of the present study was to determine the factors associated with handgrip strength in public works personnel.

METHOD

This cross-sectional study involving male public works personnel in Cengkareng District, West Jakarta, was conducted from September to October 2024. The study sample was selected by simple random sampling. Calculation of the sample size for a correlation with $r=0.2$, $Z\alpha=0.05$, and $Z\beta=0.1$, resulted in a sample of 192 subjects who met the inclusion criterion of being male and 25-60 years old, and the exclusion criteria of having an abnormality of the hands, heart disease, pulmonary disease, and malignancies.

Measurements were obtained on height in cm, weight in kg, as well as waist circumference and hip circumference in cm. Height and weight were determined using microtoise and digital scales, respectively. BMI was calculated using the formula of weight divided by height squared. Measurement of waist and hip circumferences was done using measuring tape. Waist circumference was determined with the subject in the upright position, at the end of a normal expiration, at the smallest abdominal circumference between the iliac crest and twelfth rib. The cut-off values for waist circumference in males and females are ≥ 90 cm and ≥ 80 cm, respectively. Hip circumference was measured using measuring tape that was applied around the hips at the level of the symphysis pubis and the maximal gluteal circumference (Rokhmah et al., 2015; Ilmi et al., 2020).

Fasting blood glucose (FBG) concentration was determined on venous blood by the hexokinase method after the respondents had fasted approximately 10 hours prior. Handgrip strength was measured with a handgrip strength dynamometer to the nearest kilogram as follows: The respondent was asked to sit down comfortably without reclining and with the feet completely relaxed. The hips and knees were positioned at a 90° angle, the gripping hand with the shoulder in adduction, the elbow flexed 90° , and the forearm and wrist in the neutral position. The gripping duration was 3 seconds. The respondent gripped the instrument with the dominant hand as hard as possible and the value indicated by the needle was recorded. In accordance with the protocol recommended by the American Society of Hand Therapists (ASHT), the recorded handgrip strength was the average of three handgrip strength tests (Nathania et al., 2023).

Prior to data analysis by means of the correlation test, the normality of the data distribution was determined using the Kolmogorov-Smirnov test. The Pearson correlation test was used for a normal data distribution, whereas for a non-normal data distribution, the Spearman correlation test was done, followed by a multiple regression analysis.

Ethics approval was issued by the Ethics Committee of the Faculty of Medicine, Universitas Trisakti, under No. 033/KER/FK/08/2024.

RESULT AND DISCUSSION

In this study the respondents comprised 192 male public works personnel of Cengkareng District that consisted of the villages (subdistricts) of Kedaung, Duri Kosambi, West Cengkareng, Kapuk, and Rawa Buaya. A total of 26% of the respondents were residents of Duri Kosambi, where Muslims comprised 98.4% of the population. The majority of the respondents at 77.6% had an educational level of senior high school, nearly all of the respondents, or 99%, were right-handed, using the right hand as the dominant one at work, and most were of the Betawi (Jakarta) ethnic group at 51.6% (Table 1).

Table 1. Demographic data of the respondents

Characteristic	N (%)
Village	
Kedaung	40 (20.8)
Duri Kosambi	50 (26)
Cengkareng Barat	33 (17.2)
Kapuk	43 (22.4)
Rawa Buaya	26 (13.5)
Religion	
Islamic	189 (98.4)
Christian	2 (1)
Catholic	1 (0.5)
Level of education	
Elementary school	7 (3.6)
Junior high school	29 (15.1)
Senior high school	149 (77.6)
Academy-University	7 (3.6)
Dominant hand	
Right hand	190 (99)
Left hand	2 (1)
Ethnic group	
Javanese	53 (27.6)
Betawi	99 (51.6)
Sundanese	25 (13)
Other	14 (7.3)

Legend: Other ethnicities comprised groups from Palembang, Lampung, Bima, and Kupang
Source: primary data 2024 (questionnaires and interviews with respondents)

Commented [MA1]: Dibawah setiap tabel sebutkan sumbernya. Silahkan mencermati aartikel yang sudah terbit di Jurnal KEMAS
Alvina: sudah ditambahkan sumber data dibawah setiap tabel
Commented [MA2R1]: Primari data, th.....
Alvina: sudah ditambahkan tahun primary data

Table 2. Respondents' characteristics

Variable	Mean $\pm$ SD
Age (years)	40.9 $\pm$ 8.9
Height (cm)	164.6 $\pm$ 6.7
Weight (kg)	64.7 $\pm$ 13.1
BMI (kg/m ²)	23.8 $\pm$ 4.3
Waist circumference (cm)	85.4 $\pm$ 10.4
Hip circumference (cm)	96.9 $\pm$ 8.4
HGS (kg)	33.1 $\pm$ 6.9
FBG (mg/dL)	87 (64-280)*

Legend: BMI: body mass index. *FBG: fasting blood glucose, expressed as Median (Min-Max).

HGS: handgrip strength

Source: primary data (respondents' measurements)

The bivariate analysis found a significant positive correlation of BMI, height, weight, waist circumference, and hip circumference with handgrip strength ($p=0.006$, $r=0.20$; $p<0.001$, $r=0.36$; $p<0.001$, $r=0.32$; $p=0.009$, $r=0.19$; $p=0.005$, $r=0.20$, respectively). This showed that the larger the BMI, height, weight, waist circumference, and hip circumference, the stronger the handgrip. However, there was no significant relationship between fasting glucose concentration and handgrip strength ($p=0.847$). This study also found that age had a significant negative correlation with handgrip strength ($p=0.011$; $r=-0.182$), signifying that the older a person, the lower the handgrip strength (Table 3). Results of multiple regression analysis on handgrip strength are listed in Table 4.

Table 3. Analysis of the relationship of age, height, weight, BMI, waist circumference, hip circumference, and fasting blood glucose concentration with handgrip strength

Variable	R	P value
Age	- 0.19	0.008*
Height	0.36	0.000*
Weight	0.32	0.000*
BMI	0.20	0.006*
Waist circumference	0.19	0.009*
Hip circumference	0.20	0.005*
FBG	0.02	0.847 [#]

*P value with Pearson's test, [#] Spearman's test

Source: primary data

Table 4. Multiple regression analysis on handgrip strength

Determinant	b	SE	p
Height	0.48	0.08	0.000
BMI	0.79	0.22	0.000
Hip circumference	- 0.29	0.12	0.015
Constant	- 36.98		

R²=0,19

Source: primary data

This study found that most of the public works personnel in Duri Kosambi village were Muslims, had a senior high school level of education, and were of Betawi ethnicity. This was in line with the demographic data in Cengkareng district, in that Duri Kosambi was the village with the greatest area, the majority of the residents were male, and the Islamic religion had the most believers (Badan Pusat Statistik Jakarta Barat, 2020). Among our respondents, more than ten percent were in the diabetes category. Socio-demographic factors capable of affecting dietary habits in patients with diabetes are level of education and type of family (Rondhianto et al., 2024). Among the respondents of this study, the dominant right hand was used for working. In the adult population, a total of 90% of the residents preferred to use the right hand for manual tasks, whereas around 10% used the left hand (Zaccagni et al., 2020). According to Agtuahene et al. (Agtuahene et al., 2023), the dominant hand is generally around 10% stronger than the non-dominant hand. Many activities in daily life need a high level of exertion of the muscles of the hand and forearm (Hammed et al., 2017). The frequent use of the dominant hand in daily living results in the muscles of the hand and forearm providing strength in gripping (Afaq et al., 2019).

The mean handgrip strength in this study was 33.1 ± 6.9 kg, which is relatively low if compared with the study results of Kurniawan et al. (Kurniawan et al., 2018). in healthy young Indonesian adults, where mean handgrip strength in males was 37.37 ± 8.29 kg and the lower limit of muscle strength in Indonesian males was 20.79 kg. The handgrip strength results in both studies are still below those of males in other Southeast Asian countries, according to the study results of Leong et al. (Leong et al., 2016). These investigators found that the mean handgrip strength in Southeast Asian males aged 35-40 years was 40 kg (34-44 kg); for the age range of 41-50 years, the handgrip strength was 38 kg (33-44 kg), for the age range of 51-60 years it was 34 kg (30-40 kg), and for ages 61-70 years 30 kg (24-34 kg). This may have been caused by ethnic differences mediated by genetic differences that also affect individual muscle strength. The highest handgrip strength is found in the European and North American populations, whereas the lowest handgrip strength is found in African, South Asian, and Southeast Asian populations. These variations in muscle strength in various countries may also be partly caused by differences in socio-economic status (Leong et al., 2016).

According to Agtuahene et al. (Agtuahene et al., 2023), handgrip strength is a physiological variable influenced by several factors, such as age, sex, and body measurements. In contrast, according to Zaccagni et al. (Zaccagni et al., 2020), handgrip strength, apart from being affected by age, may also be influenced by physical activity, grip measurement, height, and body mass. The present study found a significant positive correlation between BMI and handgrip strength. According to Hammed et al., this may be caused by the higher percentage of skeletal muscle mass compared to the percentage of fat mass, thereby resulting in a higher handgrip strength [13]. Our study's results are identical to those of Agtuahene et al. (Agtuahene et al., 2023), who found a

significant positive correlation ($r = 0.290$) of handgrip strength with BMI and height. This shows that taller individuals with higher BMI apparently have greater handgrip strength. Our study is also in line with the study of Krakauer in 2020, which showed a positive correlation between BMI and handgrip strength (Krakauer et al., 2020).

The study of Cooper et al. (Cooper et al., 2022) concluded that a high childhood BMI is associated with higher grip strength at age 46 years. This is because fat mass acts as a mechanical load that elicits an anabolic response that stimulates muscle growth and function. This anabolic response is usually greater in males than in females because of the higher testosterone concentration in males. However, it should also be noted that BMI does not differentiate between fat mass and lean mass as adipocyte indicators that may vary with sex. It should be noted that BMI is a measure of body fat that does not differentiate between weight changes as a result of an increase or decrease in muscle mass or body fat (Zaccagni et al., 2020). According to Jafar et al. (Jaafar et al., 2023), heavier individuals have a greater muscle mass, resulting in greater handgrip strength, than lighter individuals. In the study of Heidy et al. (Heidy et al., 2019), the investigators found a positive correlation between height and forearm length. The study by Sirajudeen et al. (Sirajudeen et al., 2012) found a positive correlation between hand length and handgrip strength. This shows that height affects the longer extremity or arm, thus approaching optimal muscle length that may result in greater grip strength. According to a prospective cohort study, handgrip strength in middle age appears to be associated with birth weight and the increase in height at puberty. Pubertal growth and handgrip strength possibly represent the development of muscle size and growth (Schrager et al., 2007).

Our study found no relationship between blood glucose concentration and handgrip strength. Our study agrees with the study of Giglio et al. (Giglio et al., 2018), which showed no significant relationship between muscle mass strength and blood glucose concentration. According to Giglio, this is because most respondents are individuals who routinely perform structured physical activity, even though they are in the light category. This is similar to our study respondents, who were public works personnel and who had performed physical activity such as street sweeping and cleaning storm drains. Glucose is the main fuel for muscle contraction, entering the muscle cells by diffusion through the GLUT4 glucose transporter, which, after muscle contraction, moves from its intracellular storage site to the plasma membrane and T-tubules (Richter et al., 2013). Glucose metabolism has an influence on handgrip strength through various mechanisms, such as disorders of glucose metabolism, particularly glycogenolysis, that may result in the loss of muscle strength. Insulin resistance in hyperglycemia may cause muscle degradation because hyperglycemia causes low muscle strength through effects on skeletal muscle mitochondria. The lowering of muscle strength by hyperglycemia occurs through Krüppel-like factor 15, which regulates skeletal muscle lipid flux. Hyperglycemia also increases pro-inflammatory cytokines, which may increase the catabolism that plays a role in the further decrease in muscle mass and quality (Gamayanti et al., 2023). The amount of carbohydrates also influences the glycemic response and post-prandial insulin level, impacting the supply of glycogen to the muscles (Safitri et al., 2020).

This study has a significant negative correlation between age and handgrip strength, which decreases with age (Manoharan et al., 2015). Handgrip strength initially experiences an increase with increasing age, reaches its peak between the age of 30 and 40 years, and decreases afterward (Zaccagni et al., 2020). The age-related changes in skeletal muscle composition, particularly the accumulation of lipids in skeletal muscle fibers, play a role in the poor quality of the muscles and may result in metabolic abnormalities such as insulin resistance. Apart from changes in muscle

quality, there is also a reduction in muscle mass that is consistently occurring during aging. The increase in oxidative stress and chronic inflammation that commonly occurs in the aging process apparently also plays a role in the development of decreased muscle mass and strength (Mainous et al., 2015). The decrease in muscle mass in the elderly corresponds to a 10% reduction in grip strength per decade, starting from the age of 40 years (Jaafar et al., 2023). Skeletal muscle plays an important role in glucose metabolism and has a significant influence on insulin sensitivity. Increased insulin resistance in older adults may be closely associated with skeletal muscle aging. A reduction in GLUT4 expression as a result of a decrease in muscle volume causes a reduction in insulin sensitivity in aging skeletal muscle (Joo et al., 2022).

The results of this study showed a correlation of waist circumference and hip circumference with handgrip strength. This is in line with the study of Widjaja et al. (Widjaja et al., 2018), who stated that there was a significant correlation of waist circumference and hip circumference with handgrip strength ($p < 0.001$, $r = 0.302$; and $p < 0.001$, $r = 0.296$). A greater waist circumference is associated with a greater handgrip strength in males (Hardy et al., 2013).

In the multiple regression analysis of HGS, we removed the data on body weight because of the occurrence of multicollinearity between weight and BMI, with a correlation coefficient of more than 0.9, indicating an extremely strong relationship between these two variables. Conceptually, only one of the variables is used in multivariate analysis. The multiple regression analysis included age, height, BMI, waist circumference, and hip circumference, with HGS as dependent variable in a stepwise dependent manner. After three iterations, only height, BMI, and hip circumference played a role in predicting HGS by means of the regression formula, which resulted in $R^2 = 0.19$. This signifies that for known values of height, BMI, and hip circumference, we can predict the HGS value 19% more accurately than without a knowledge of these data. For a clinical diagnosis, a minimum increase of 10% is needed. This study found that height, BMI, and hip circumference play a role in predicting handgrip strength through the regression formula, i.e., $\text{handgrip strength} = -36.98 + 0.48 \times \text{height} + 0.79 \times \text{BMI} - 0.29 \times \text{hip circumference}$. Handgrip strength and waist circumference are surrogate muscle strength and visceral adiposity markers, respectively (Nakanishi et al., 2024). Waist circumference is also more sensitive in determining body fat distribution, particularly in the abdomen. Waist circumference correlates better with abdominal fat distribution than BMI (Rokhmah et al., 2015). Waist circumference and hip circumference are associated with obesity, in which the fat content is higher, and the skeletal muscle and fascicular content is lower, thereby resulting in a lower contractile capacity of skeletal muscle (Afaq et al., 2019).

The limitation of this study is that it did not evaluate other factors that may affect handgrip strength, such as nutrient intake. From the anthropometric measures of the hands and feet in this study, the causes and effects of the decrease in handgrip strength could not be determined because this study used a cross-sectional design.

CONCLUSION

There are three factors that may be used to estimate handgrip strength in PWP, namely height, BMI, and hip circumference, whereas blood glucose concentration is not correlated with handgrip strength in PWP. It is recommended to pay attention to height, BMI, and hip circumference,

particularly in the acceptance criteria of PWP candidates, and to conduct further studies with consideration of other factors that affect handgrip strength, such as food intake and anthropometric features of the hands and arms.

ACKNOWLEDGMENTS

Financial support was provided by Universitas Trisakti for this research.

REFERENCES

Afaq, E., Hassan, S.H., Khan, M.R., Zaidi, D.A., & Baktashi, H., 2019. Correlation of Hand Dynamometry With Body Mass Index and Waist Hip Ratio in Medical Students. *Pak J Physiol*, 15, pp.19-21.

Aguahene, M.A., Quartey, J., & Kwakye, S., 2023. Influence of Hand Dominance, Gender, and Body Mass Index on Hand Grip Strength. *S Afr J Physiother*, 79, pp.1-6. doi: [org/10.4102/sajp.v79i1.1923](https://doi.org/10.4102/sajp.v79i1.1923).

Al-Asadi, J.N., 2018. Handgrip Strength in Medical Students: Correlation with Body Mass Index and Hand Dimensions. *Asian Journal of Medical Sciences*, 9, pp.21-26. doi: [10.3126/ajms.v9i1.18577](https://doi.org/10.3126/ajms.v9i1.18577).

Badan Pusat Statistik Kota Administrasi Jakarta Barat., 2020. Kecamatan Cengkareng Dalam Angka, pp. 1-114.

Chen, D., Zhu, Y., Ni, W., Li, Y., Yin, G., Shao, Z., & Zhu, J., 2023. Hand Grip Strength is Inversely Associated with Total Daily Insulin Dose Requirement in Patients with Type 2 Diabetes Mellitus: A Cross-Sectional Study. *PeerJ*, pp.1-13. doi: [10.7717/peerj.15761](https://doi.org/10.7717/peerj.15761).

Cooper, R., Tomlinson, D., Hamer, M., & Pereira, S.M.P., 2022. Lifetime Body Mass Index and Grip Strength at Age 46 Years: The 1970 British Cohort Study. *J Cachexia Sarcopenia Muscle*, 13, pp.1-10. doi: [10.1002/jcsm.12992](https://doi.org/10.1002/jcsm.12992).

Fauzi, L., Handayani, O.W.K., Susilo, M.T., Kurnia, A.R., Rahayu, S.R., Irawan, F.A., Horng Lu, F.J., Lin, C., Lai, M.F., & Yu, Y.C., 2022. Obesity in Indonesian and Taiwanese Adolescents Related to Self Perception, Diet, Exercise and Body Image. *KEMAS*, 17(3):pp.453-461. doi: [10.15294/kemas.v17i3.34396](https://doi.org/10.15294/kemas.v17i3.34396)

Gamayanti, K.A.V., Aryana, I.G.P.S., & Gotera, W., 2023. Hubungan Antara Kekuatan Genggaman Tangan Dengan Kadar Gula Darah Sewaktu Pada Lansia Di Desa Melinggih, Kecamatan Payangan, Kabupaten Gianyar. *Intisari Sains Medis*, 14(2): pp.483-488.

Giglio, B.M., Mota, J.F., Wall, B.T., & Pimentel, G.D., 2018. Low Handgrip Strength is Not Associated With Type 2 Diabetes Mellitus and Hyperglycemia: A Population-Based Study. *Clin Nutr Res*, 7(2): pp.112-116. doi: [10.7762/cnr.2018.7.2.112](https://doi.org/10.7762/cnr.2018.7.2.112).

Hammed, A.I., & Obaseki, C.O., 2017. Interdependence of Body Mass Index with Handgrip Strength and Endurance Among Apparently Healthy Teenagers. *Turk J Kin*, 4, pp.1-7.

Handayani, M.D.N., Sadewa, A.H., Farmawati, A., & Rochmah, W., 2018. Anthropometric Prediction Equations for Estimating Muscle Mass of Elderly Women. *KEMAS*, 14(2):pp.195-204. doi: 10.15294/kemas.v14i2.14073

Hardy, R., Cooper, R., Sayer, A.A., Shlomo, Y.B., Cooper, C., Deary, I.J., Demakakos, P., Gallacher, J., Martin, R.M., McNeill, G., et al., 2013. Body Mass Index, Muscle Strength and Physical Performance in Older Adults From Eight Cohort Studies: The Halcyon Programme. *PLoS One*, 8(2): pp.e56483. doi: 10.1371/journal.pone.0056483.

Heidy., Djuartina, T., & Irawan, R., 2019. Korelasi Kekuatan Genggaman Tangan Dengan Karakter Antropometri Lengan Bawah dan Tangan Serta Indeks Massa Tubuh. *Damianus Journal of Medicine*, 18, pp.1-7.

Ilmi, A.F., & Utari, D.M., 2020. Hubungan Lingkar Pinggang dan Rasio Lingkar Pinggang-Panggul (RLPP) Terhadap Kadar Glukosa Darah Puasa Pada Mahasiswa. *Journal of Nutrition College*, 9, pp.222-227.

Jaafar, M.H., Ismail, R., Ismail, N.H., Isa, Z.M., Tamil, A.M., Nasir, N.M., Keat, N.K., Razak, N.H., Abidin, N.Z., & Yusof, K.H., 2023. Normative Reference Values and Predicting Factors of Handgrip Strength For Dominant and Non-Dominant Hands Among Healthy Malay Adults in Malaysia. *BMC Musculoskelet Disord*, 24, pp. 1-9. doi: 10.1186/s12891-023-06181-8.

Joo, K.C., Son, D.H., & Park, J.M., 2022. Association Between Relative Handgrip Strength and Insulin Resistance in Korean Elderly Men Without Diabetes: Findings Of The 2015 Korea National Health Nutrition Examination Survey. *Korean J Fam Med*, 43, pp.199-205. doi:10.4082/kjfm.21.0138.

Krakauer, N.Y., & Krakauer, J.C., 2020. Association of Body Shape Index (ABSI) with Hand Grip Strength. *Int J Environ Res Public Health*, 17, pp.1-12. doi: doi:10.3390/ijerph17186797.

Kurniawan, A., Widjaja, D., & Lugito, N.P.H., 2018. Mean and Cutoff Value of Hand Grip Strength For Healthy Indonesian Young Adults. *World Congress on Osteoporosis, Osteoarthritis and Musculoskeletal Diseases*; Krakow, Poland.

Leong, D.P., Teo, K.K., Rangarajan, S., Kutty, R., Lanas, F., Hui, C., Quanyong, X., Zhenzhen, Q., Jinhua, T., Noorhassim, I., et al., 2016. Reference Ranges of Handgrip Strength From 125,462 Healthy Adults in 21 Countries: A Prospective Urban Rural Epidemiologic (PURE) Study. *J Cachexia Sarcopenia Muscle*, 7, pp.535-546. doi: 10.1002/jcsm.12112.

Mainous, A.G., Tanner, R.J., Anton, S.D., & Jo, A., 2015. Grip Strength as A Marker of Hypertension and Diabetes in Healthy Weight Adults. *Am J Prev Med*, 49(6): pp.850-858. doi: 10.1016/j.amepre.2015.05.025.

Manoharan, V.S., Sundaram, S.G., & Jason, J.I., 2015. Factors Affecting Hand Grip Strength and Its Evaluation: A Systemic Review. *Int J Physiother Res*, 3(6): pp.1288-1293. doi:10.16965/ijpr.2015.193.

McGrath, R., Johnson, N., Klawitter, L., Mahoney, S., Trautman, K., Carlson, C., Rockstad, E., & Hackney, K.J., 2020. What Are The Association Patterns Between Handgrip Strength and Adverse Health Conditions? A Topical Review. *SAGE Open Medicine*, 8, pp.1-12. doi:10.1177/2050312120910358.

Nakanishi, S., Shimoda, M., Kimura, T., Sanada, J., Fushimi, Y., Iwamoto, Y., Iwamoto, H., Dan, K., Mune, T., Kaku, K., et al., 2024. The Impact of Handgrip Strength and Waist Circumference on Glycemic Control: Prospective, Observational Study Using Outpatient Clinical Data in Japanese Patients With Type 2 Diabetes Mellitus. *J Diabetes Investig*, 15(7): pp.892-898. doi: 10.1111/jdi.14200.

Nathania, N.P.S., Nugraha, M.H.S., Primayanti, I.D.A., & Saraswati, P.A.S., 2023. Handgrip Strength is Associated With Balance in Elderly Women. *Majalah Ilmiah Fisioterapi Indonesia*, 11, pp.70-75.

Peraturan Gubernur Provinsi Daerah Khusus Ibukota Jakarta Nomor 7 tahun 2017 Tentang Penanganan Prasarana dan Sarana Umum Tingkat Kelurahan. Jakarta 2017.

Richter, E.A., & Hargreaves, M., 2013. Exercise, GLUT4, and Skeletal Muscle Glucose Uptake. *Physiol Rev*, 93(3): pp.993-1017. doi: 10.1152/physrev.00038.2012.

Rokhmah, F.D., Handayani, D., & Al-Rasyid, H., 2015. Korelasi Lingkar Pinggang dan Rasio Lingkar Pinggang-Panggul Terhadap Kadar Glukosa Plasma Menggunakan Tes Toleransi Glukosa Oral. *Jurnal Gizi Klinik Indonesia*, 12, pp.28-35.

Rondhianto., Ridla, A.Z., & Hasan, H., 2024. Sociodemographic Factors Affecting Diabetic Dietary Behavior in People with Type 2 Diabetes Mellitus. *KEMAS*, 19(3):pp.429-437. doi: 10.15294/kemas.v19i3.37856

Safitri, I., Setyarsih, L., Susanto, H., Suhartono., & Fitranti, D.Y., 2020. The Effect of Low and High Glycemic Load Diet on Muscle Fatigue of Young Soccer. *KEMAS*, 16(1):pp.111-120. doi: 10.15294/kemas.v16i1.23508

Schrager, M.A., Metter, E.J., Simonsick, E., Ble, A., Bandinelli, S., Lauretani, F., & Ferrucci, L., 2007. Sarcopenic Obesity and Inflammation in The Inchiati Study. *J Appl Physiol*, 102(3): pp.919-925. doi: 10.1152/japplphysiol.00627.2006.

Sirajudeen, M.S., Shah, U.N., Pillai, P.S., Mohasin, N., & Shantaram, M., 2012. Correlation Between Grip Strength and Physical Factors in Men. *International Journal of Health and Rehabilitation Sciences*, 1, pp.58-63.

Soenyoto, T., Mollap, K., & Mungpong, J., 2017. Correlation of Energy and Protein Consumption Levels with Physical Endurance of Rhythmic Gymnast Athletes. KEMAS, 13(1):pp.131-136. doi:10.15294/kemas.v13i1.9888

Soraya, N., & Parwanto, E., 2023. The Controversial Relationship Between Body Mass Index and Handgrip Strength in The Elderly: An Overview. Malays J Med Sci, 30, pp.73-83.

Vaishya, R., Misra, A., Vaish, A., Ursino, N., & D'Ambrosi, R., 2024. Hand Grip Strength as A Proposed New Vital Sign Of Health: A Narrative Review of Evidences. Journal of Health, Population and Nutrition, 43, pp.1-14. doi:10.1186/s41043-024-00500-y.

Widjaja, D., Tjhai, K., Dermawan, K., Kalim, J., Tanzil, S., Pramusinta, A., Phenca, H.K., Kurniawan, A., 2018. Correlations Between Hand Grip Strength and Anthropometric Status in Young Indonesian Adults. World Congress on Osteoporosis, Osteoarthritis and Musculoskeletal Diseases; Krakow, Poland.

Zaccagni, L., Toselli, S., Bramanti, B., Gualdi-Russo, E., Mongillo, J., & Rinaldo, N., 2020. Handgrip Strength in Young Adults: Association With Anthropometric Variables and Laterality. Int J Environ Res Public Health, 17, pp.1-18. doi:10.3390/ijerph17124273.

6. Bukti Accepted artikel
(1 Agustus 2025)



Alvina fk <dr.alvina@trisakti.ac.id>

Pemberitahuan Penyelesaian Administrasi

1 message

Jurnal Kemas <kemas@mail.unnes.ac.id>

Fri, Aug 1, 2025 at 4:04 PM

To: Alvina fk <dr.alvina@trisakti.ac.id>, pusparini@trisakti.ac.id, mario@trisakti.ac.id, yasminedrpk@gmail.com

Yth. Penulis di Jurnal KEMAS

Dengan hormat,
Kami beritahukan bahwa artikel Bapak/ Ibu dengan:

No. **25917** Judul: **Handgrip Strength of Public Works Personnel in West Jakarta**

akan segera kami proses untuk penerbitan di Jurnal KEMAS (volume 21, no.2, Oktober - Desember 2025).

Jika dalam 10 hari kerja
belum merespon email ini, maka redaksi berhak memundurkan penerbitan artikel ke edisi berikutnya

Sesuai dengan syarat administratif, bagi artikel yang akan dimuat:

1. Penulis dikenakan biaya publish dan translet/proofread sebesar Rp. 3.300.000,00 dengan ketentuan sebagai berikut:
 - a. Biaya tersebut dikirim ke Jurnal KEMAS ke no rek. BNI 0328641658 a.n Widya Hary Cahyati untuk mendapatkan surat LAYAK MUAT.
 - b. Bukti transfer harap dikirim ke email: kemas@mail.unnes.ac.id
2. Penulis menyertakan Surat Bebas Plagiat dan dikirim melalui email kemas@mail.unnes.ac.id (Form terlampir, mohon bubuhkan materai 10.000 pada form tersebut).

Demikian pemberitahuan kami, atas perhatian dan kerjasamanya kami ucapkan terimakasih.

Ketua Dewan Redaksi

Prof. Dr. dr. Oktia Woro K.H., M.Kes.

KEMAS Journal

F5 Building, 2nd Floor, Public Health Department, Sport Science Faculty, Semarang State University, Semarang, Central Java, Indonesia, 50229

<http://journal.unnes.ac.id/nju/index.php/kemas>

DISCLAIMER

This email may contain confidential or copyrighted information of UNNES. If you are not the intended recipient, please do not use or share this email. If received in error, please notify the sender and delete it. Check for viruses; UNNES is not liable for virus-related damages.



English_SURAT PERNYATAAN BEBAS PLAGIAT.doc

31K

7. Bukti konfirmasi proofreading
dan hasil proofreading
(23 September 2025)



Alvina fk <dr.alvina@trisakti.ac.id>

Dummy Article

3 messages

Jurnal Kemas <kemas@mail.unnes.ac.id>

Tue, Sep 23, 2025 at 6:53 PM

To: Alvina fk <dr.alvina@trisakti.ac.id>, pusparini@trisakti.ac.id, mario@trisakti.ac.id, yasminedrpk@gmail.com

Berikut kami kirimkan *Dummy* artikel Bapak/Ibu yang telah dilayout, yang akan diterbitkan di Jurnal KEMAS Vol. 21 No. 2

Mohon untuk mengecek dengan seksama artikel tersebut.

Jika Bapak/Ibu sudah setuju, mohon mengkonfirmasi persetujuan cetak. Jika masih ada kekeliruan mohon untuk menuliskan perbaikannya via email: kemas@mail.unnes.ac.id.

Kami tunggu konfirmasinya maksimal hari Sabtu, 27 September 2025 Jam 12.00 WIB.

Jika dalam waktu tersebut tidak ada konfirmasi, maka kami anggap bahwa Bapak/ Ibu telah menyetujui artikel tersebut.

Konfirmasi setelah tanggal dan jam tersebut tidak dapat dilayani.

Terima kasih

KEMAS Journal

F5 Building, 2nd Floor, Public Health Department, Sport Science Faculty, Semarang State University, Semarang, Central Java, Indonesia, 50229

<http://journal.unnes.ac.id/nju/index.php/kemas>

DISCLAIMER

This email may contain confidential or copyrighted information of UNNES. If you are not the intended recipient, please do not use or share this email. If received in error, please notify the sender and delete it. Check for viruses; UNNES is not liable for virus-related damages.

 **11_25917 Alvina 360 368.pdf**
152K**Alvina fk** <dr.alvina@trisakti.ac.id>

Thu, Sep 25, 2025 at 1:50 PM

To: Jurnal Kemas <kemas@mail.unnes.ac.id>

Kepada Yth,
Sekretariat Jurnal Kemas

Terima kasih atas kiriman dummy artikel, setelah saya periksa ada beberapa yang beda dengan naskah asli. Terlampir saya kirimkan koreksi dummy artikel dan artikel asli (dalam bentuk word).

Sudah saya tandai untuk dummy artikel yg berbeda dan di naskah asli sudah saya tandai juga dengan warna hijau.

Terima kasih.

Salam,

Alvina

[Quoted text hidden]

2 attachments

 **Input Koreksi 11_25917 Alvina 360 368.pdf**
141K

 **Alvina-Factor+associated+with+handgripstrength (1).docx**
43K

Jurnal Kemas <kemas@mail.unnes.ac.id>

To: Alvina fk <dr.alvina@trisakti.ac.id>

Sun, Sep 28, 2025 at 10:56 PM

Baik, akan kami sampaikan ke layouter.

Terima kasih

KEMAS Journal

F5 Building, 2nd Floor, Public Health Department, Sport Science Faculty, Semarang State University, Semarang, Central Java, Indonesia, 50229

<http://journal.unnes.ac.id/nju/index.php/kemas>

[Quoted text hidden]

[Quoted text hidden]



UNIVERSITAS TRISAKTI

"Is a one stop learning for sustainable development"

Kampus A, Jl. Kyai Tapa No.1, Grogol

Jakarta Barat 11440 - INDONESIA

www.trisakti.ac.id

(t) +62-21.566 3232, (f) +62-21.567 3001

[Quoted text hidden]



Handgrip Strength of Public Works Personnel in West Jakarta

Alvina^{1✉}, Pusparini¹, Mario¹, Yasmine Mashabi¹

¹Department of Clinical Pathology, Faculty of Medicine, Universitas Trisakti

Article Info

Article History:

Submitted: May 2025

Accepted: September 2025

Published: October 2025

Keywords:

Handgrip strength; Body mass index; Hip circumference; Waist circumference; Fasting blood glucose

DOI

<https://doi.org/10.15294/kemas.v21i2.25917>

Abstract

Public Works Personnel (PWP) in Jakarta perform their routine duties manually instead of using equipment. Handgrip strength (HGS) measures maximum hand strength as a quality indicator for muscle strength and mass. Factors affecting muscle strength are age, sex, body mass index (BMI), waist circumference (WC), and hip circumference (HC). Increased fasting blood glucose (FBG) is also associated with muscle quality, muscle strength, and physical performance. This study aimed to determine the factors associated with handgrip strength in PWP. A total of 192 male PWP from the Cengkareng district were recruited by simple random sampling. The collected data were demographics, BMI, FBG, WC, HC, and HGS. Statistical analysis was done using Pearson's correlation test and multiple regression analysis. There was a significant positive correlation of BMI, WC, and HC with HGS ($p=0.006$; $r=0.20$, $p=0.009$; $r=0.19$, and $p=0.005$; $r=0.20$) and a significant negative correlation between age and HGS ($p=0.008$; $r=-0.19$) but not between FBG and HGS ($p=0.847$). Multiple regression analysis showed height, BMI, and HC positively predicting HGS ($R^2=0.19$). Handgrip strength of public works personnel is associated with BMI, age, waist circumference, and hip circumference, but not with fasting blood glucose. Height, BMI, and hip circumference predict handgrip strength

Introduction

Personnel for the management of public infrastructure and facilities at the village level are workers who manage village public works for a given time period on the basis of a work order. The tasks of village-level public works personnel (PWP) consist of managing the infrastructure and facilities, such as streets, drains, and parks, including installed lights (Gubernur Provinsi Daerah Khusus Ibukota Jakarta, 2017). The work performed by village PWP consists mainly of intense physical activity involving the hands, such that handgrip strength is essential in their tasks. Lowering of handgrip strength may indicate reduced muscle strength, which may especially interfere with the daily activities of village PWP that consist mainly of manual labor.

Handgrip strength measures maximal hand strength and indicates muscle quality that reflects muscle strength and mass. Muscle

strength refers to the ability of muscle groups to raise a given load. Low or poor grip strength is frequently associated with damaging health impacts, such as chronic disease, and is the initial sign of reduced muscle mass, limitations in physical functions, and mortality risk. Poor handgrip strength may also reduce a person's ability to perform activities of daily living and, therefore, impact quality of life (Soraya *et al.*, 2023; Nathania *et al.*, 2023). Muscle mass accounts for approximately 40% of total weight, such that knowledge of muscle mass conditions is essential in maintaining health and activities (Handayani *et al.*, 2018).

Handgrip strength may represent the strength of all muscles in the body. Several factors possibly influencing an individual's muscle strength are age, sex, and body mass index (BMI) (Soraya *et al.*, 2023). Obesity may also affect BMI. Obesity in adults is not only associated with genetic factors but also with

✉ Correspondence Address:
Department of Clinical Pathology, Faculty of Medicine, Universitas Trisakti
Email: dr.alvina@trisakti.ac.id

environmental factors during development (Fauzi *et al.*, 2022). Handgrip strength is also associated with chronic diseases such as diabetes, metabolic syndrome, and cardiovascular disease. Increased glucose concentration is also associated with disorders of muscle quality and strength, as well as physical performance (Chen *et al.*, 2023). A reduction in muscle mass also affects an individual's strength and physical endurance. A combination of proteins can maintain muscle function, thereby building body strength and physical endurance (Soenyoto *et al.*, 2017). In the present study, we included fasting blood glucose concentrations in the data collection to obtain more accurate measures of glucose concentrations that have not yet been influenced by food intake.

Handgrip strength may be a health indicator that provides information on a person's health (Vaishya *et al.*, 2024). However, its clinical significance in determining the prevention and treatment of poor handgrip remains limited because of the still unclear etiology of muscle weakness, especially because the clinical outcomes associated with handgrip strength are extremely diverse (McGrath *et al.*, 2020). The relationship between BMI and handgrip strength is still controversial, with several investigators finding a significant positive association and others reporting a non-significant relationship (Al Asadi, 2018). There are few studies on handgrip strength, especially on workers in the sanitation sector, such as public works personnel, which is why the present investigators were interested in studying this topic. The present study aimed to determine the factors associated with handgrip strength in public works personnel.

Method

This cross-sectional study involving male public works personnel in Cengkareng District, West Jakarta, was conducted from September to October 2024. The study sample was selected by simple random sampling. Calculation of the sample size for a correlation with $r=0.2$, $Z\alpha=0.05$, and $Z\beta=0.1$ resulted in a sample of 192 subjects who met the inclusion criterion of being male and 25-60 years old, and the exclusion criteria of having an abnormality of the hands, heart disease, pulmonary disease,

and malignancies.

Measurements were obtained on height in cm, weight in kg, as well as waist circumference and hip circumference in cm. Height and weight were determined using microtoise and digital scales, respectively. BMI was calculated using the formula of weight divided by height squared. Measurement of waist and hip circumferences was done using a measuring tape. Waist circumference was determined with the subject in the upright position, at the end of a normal expiration, at the smallest abdominal circumference between the iliac crest and twelfth rib. The cut-off values for waist circumference in males and females are ≥ 90 cm and ≥ 80 cm, respectively. Hip circumference was measured using a measuring tape that was applied around the hips at the level of the symphysis pubis and the maximal gluteal circumference (Ilmi *et al.*, 2020).

Fasting blood glucose (FBG) concentration was determined on venous blood by the hexokinase method after the respondents had fasted for approximately 10 hours prior. Handgrip strength was measured with a handgrip strength dynamometer to the nearest kilogram as follows: The respondent was asked to sit down comfortably without reclining and with the feet completely relaxed. The hips and knees were positioned at a 90° angle, the gripping hand with the shoulder in adduction, the elbow flexed 90° , and the forearm and wrist in the neutral position. The gripping duration was 3 seconds. The respondent gripped the instrument with the dominant hand as hard as possible, and the value indicated by the needle was recorded. In accordance with the protocol recommended by the American Society of Hand Therapists (ASHT), the recorded handgrip strength was the average of three handgrip strength tests (Nathania *et al.*, 2023).

Before data analysis by means of the correlation test, the normality of the data distribution was determined using the Kolmogorov-Smirnov test. The Pearson correlation test was used for a normal data distribution, whereas for a non-normal data distribution, the Spearman correlation test was done, followed by a multiple regression analysis. Ethics approval was issued by the Ethics Committee of the Faculty of Medicine,

Universitas Trisakti, under No. 033/KER/FK/08/2024.

Result and Discussion

In this study, the respondents comprised 192 male public works personnel of Cengkareng District who consisted of the villages (subdistricts) of Kedaung, Duri Kosambi, West Cengkareng, Kapuk, and Rawa Buaya. A total of

26% of the respondents were residents of Duri Kosambi, where Muslims comprised 98.4% of the population. The majority of the respondents, at 77.6% had an educational level of senior high school. Nearly all of the respondents, or 99%, were right-handed, using the right hand as the dominant one at work, and most were of the Betawi (Jakarta) ethnic group at 51.6% (Table 1).

Table 1. Demographic Data of the Respondents

Characteristic	N (%)
Village	
Kedaung	40 (20.8)
Duri Kosambi	50 (26)
Cengkareng Barat	33 (17.2)
Kapuk	43 (22.4)
Rawa Buaya	26 (13.5)
Religion	
Islamic	189 (98.4)
Christian	2 (1)
Catholic	1 (0.5)
Level of education	
Elementary school	7 (3.6)
Junior high school	29 (15.1)
Senior high school	149 (77.6)
Academy-University	7 (3.6)
Dominant hand	
Right hand	190 (99)
Left hand	2 (1)
Ethnic group	
Javanese	53 (27.6)
Betawi	99 (51.6)
Sundanese	25 (13)
Other	14 (7.3)

Legend: Other ethnicities comprised groups from Palembang, Lampung, Bima, and Kupang
Source: primary data 2024 (questionnaires and interviews with respondents)

Table 2. Respondents' Characteristics

Variable	Mean ± SD
Age (years)	40.9 ± 8.9
Height (cm)	164.6 ± 6.7
Weight (kg)	64.7 ± 13.1
BMI (kg/m²)	23.8 ± 4.3
Waist circumference (cm)	85.4 ± 10.4
Hip circumference (cm)	96.9 ± 8.4

HGS (kg)	33.1 ± 6.9
FBG (mg/dL)	87 (64-280)*

Legend: BMI: body mass index. *FBG: fasting blood glucose, expressed as Median (Min-Max).

HGS: handgrip strength

Source: primary data (respondents' measurements)

Table 3. Analysis of the Relationship of Age, Height, Weight, BMI, Waist Circumference, Hip Circumference, and Fasting Blood Glucose Concentration with Handgrip Strength

Variable	R	P value
Age	- 0.19	0.008*
Height	0.36	0.000*
Weight	0.32	0.000*
BMI	0.20	0.006*
Waist circumference	0.19	0.009*
Hip circumference	0.20	0.005*
FBG	0.02	0.847 [#]

*P value with Pearson's test, [#] Spearman's test

Source: primary data

Table 4. Multiple Regression Analysis on Handgrip Strength

Determinant	b	SE	p
Height	0.48	0.08	0.000
BMI	0.79	0.22	0.000
Hip circumference	- 0.29	0.12	0.015
Constant	- 36.98		

R²=0,19

Source: primary data

The bivariate analysis found a significant positive correlation of BMI, height, weight, waist circumference, and hip circumference with handgrip strength ($p=0.006$, $r=0.20$; $p<0.001$, $r=0.36$; $p<0.001$, $r=0.32$; $p=0.009$, $r=0.19$; $p=0.005$; $r=0.20$, respectively). This showed that the larger the BMI, height, weight, waist circumference, and hip circumference, the stronger the handgrip. However, there was no significant relationship between fasting glucose concentration and handgrip strength ($p=0.847$). This study also found that age had a significant negative correlation with handgrip strength ($p=0.011$; $r=-0.182$), signifying that the older a person, the lower the handgrip strength (Table 3). Results of multiple regression analysis on handgrip strength are listed in Table 4.

This study found that most of the public works personnel in Duri Kosambi village were Muslims, had a senior high school level of education, and were of Betawi ethnicity. This was in line with the demographic data in the Cengkareng district, in that Duri Kosambi was

the village with the greatest area, most of the residents were male, and the Islamic religion had the most believers (Badan Pusat Statistik Jakarta Barat, 2020). Among our respondents, more than ten percent were in the diabetes category. Socio-demographic factors capable of affecting dietary habits in patients with diabetes are the level of education and the type of family (Rondhianto *et al.*, 2024). Among the respondents of this study, the dominant right hand was used for working. In the adult population, a total of 90% of the residents preferred to use the right hand for manual tasks, whereas around 10% used the left hand (Zaccagni *et al.*, 2020). According to Agtuahene *et al.* (Agtuahene *et al.*, 2023), the dominant hand is generally around 10% stronger than the non-dominant hand. Many activities in daily life need a high level of exertion of the muscles of the hand and forearm (Hammed *et al.*, 2017). The frequent use of the dominant hand in daily living results in the muscles of the hand and forearm providing strength in gripping (Afaq *et*

al., 2019).

The mean handgrip strength in this study was 33.1 ± 6.9 kg, which is relatively low if compared with the study results of Kurniawan *et al.* (Kurniawan *et al.*, 2018). In healthy young Indonesian adults, the mean handgrip strength in males was 37.37 ± 8.29 kg, and the lower limit of muscle strength in Indonesian males was 20.79 kg. The handgrip strength results in both studies are still below those of males in other Southeast Asian countries, according to the study results of Leong *et al.* (Leong *et al.*, 2016). These investigators found that the mean handgrip strength in Southeast Asian males aged 35-40 years was 40 kg (34-44 kg); for the age range of 41-50 years, the handgrip strength was 38 kg (33-44 kg), for the age range of 51-60 years it was 34 kg (30-40 kg), and for ages 61-70 years 30 kg (24-34 kg). This may have been caused by ethnic differences mediated by genetic differences that also affect individual muscle strength. The highest handgrip strength is found in the European and North American populations, whereas the lowest handgrip strength is found in African, South Asian, and Southeast Asian populations. These variations in muscle strength in various countries may also be partly caused by differences in socioeconomic status (Leong *et al.*, 2016).

According to Agtuahene *et al.* (Agtuahene *et al.*, 2023), handgrip strength is a physiological variable influenced by several factors, such as age, sex, and body measurements. In contrast, according to Zaccagni *et al.* (Zaccagni *et al.*, 2020), handgrip strength, apart from being affected by age, may also be influenced by physical activity, grip measurement, height, and body mass. The present study found a significant positive correlation between BMI and handgrip strength. According to Hammed *et al.*, this may be caused by the higher percentage of skeletal muscle mass compared to the percentage of fat mass, thereby resulting in a higher handgrip strength [13]. Our study's results are identical to those of Agtuahene *et al.* (Agtuahene *et al.*, 2023), who found a significant positive correlation ($r=0.290$) of handgrip strength with BMI and height. This shows that taller individuals with higher BMI apparently have greater handgrip strength. Our study is also in line with the study of Krakauer in 2020, which

showed a positive correlation between BMI and handgrip strength (Krakauer *et al.*, 2020).

The study of Cooper *et al.* (Cooper *et al.*, 2022) concluded that a high childhood BMI is associated with higher grip strength at age 46 years. This is because fat mass acts as a mechanical load that elicits an anabolic response that stimulates muscle growth and function. This anabolic response is usually greater in males than in females because of the higher testosterone concentration. However, it should also be noted that BMI does not differentiate between fat mass and lean mass as adipocyte indicators that may vary with sex. It should be noted that BMI is a measure of body fat that does not differentiate between weight changes because of an increase or decrease in muscle mass or body fat (Zaccagni *et al.*, 2020). According to Jafar *et al.* (Jaafar *et al.*, 2023), heavier individuals have a greater muscle mass, resulting in greater handgrip strength, than lighter individuals. In the study of Heidy *et al.* (Heidy *et al.*, 2019), the investigators found a positive correlation between height and forearm length. The study by Sirajudeen *et al.* (Sirajudeen *et al.*, 2012) found a positive correlation between hand length and handgrip strength. This shows that height affects the longer extremity or arm, thus approaching optimal muscle length that may result in greater grip strength. According to a prospective cohort study, handgrip strength in middle age appears to be associated with birth weight and the increase in height at puberty. Pubertal growth and handgrip strength possibly represent the development of muscle size and growth (Schrager *et al.*, 2007).

Our study found no relationship between blood glucose concentration and handgrip strength. Our study agrees with the study of Giglio *et al.* (Giglio *et al.*, 2018), which showed no significant relationship between muscle mass strength and blood glucose concentration. According to Giglio, this is because most respondents are individuals who routinely perform structured physical activity, even though they are in the light category. This is similar to our study respondents, who were public works personnel and who had performed physical activity such as street sweeping and cleaning storm drains. Glucose is the main fuel

for muscle contraction, entering the muscle cells by diffusion through the GLUT4 glucose transporter, which, after muscle contraction, moves from its intracellular storage site to the plasma membrane and T-tubules (Richter *et al.*, 2013). Glucose metabolism has an influence on handgrip strength through various mechanisms, such as disorders of glucose metabolism, particularly glycogenolysis, that may result in the loss of muscle strength. Insulin resistance in hyperglycemia may cause muscle degradation because hyperglycemia causes low muscle strength through effects on skeletal muscle mitochondria. The lowering of muscle strength by hyperglycemia occurs through Krüppel-like factor 15, which regulates skeletal muscle lipid flux. Hyperglycemia also increases pro-inflammatory cytokines, which may increase the catabolism that plays a role in the further decrease in muscle mass and quality (Gamayanti *et al.*, 2023). The amount of carbohydrates also influences the glycemic response and post-prandial insulin level, impacting the supply of glycogen to the muscles (Safitri *et al.*, 2020).

This study has a significant negative correlation between age and handgrip strength, which decreases with age (Manoharan *et al.*, 2015). Handgrip strength initially experiences an increase with increasing age, reaches its peak between the ages of 30 and 40 years, and decreases afterward (Zaccagni *et al.*, 2020). The age-related changes in skeletal muscle composition, particularly the accumulation of lipids in skeletal muscle fibers, play a role in the poor quality of the muscles and may result in metabolic abnormalities such as insulin resistance. Apart from changes in muscle quality, there is also a reduction in muscle mass that consistently occurs during aging. The increase in oxidative stress and chronic inflammation that commonly occurs in the aging process apparently also plays a role in the development of decreased muscle mass and strength (Mainous *et al.*, 2015). The decrease in muscle mass in the elderly corresponds to a 10% reduction in grip strength per decade, starting from the age of 40 years (Jaafar *et al.*, 2023). Skeletal muscle plays an important role in glucose metabolism and has a significant influence on insulin sensitivity. Increased

insulin resistance in older adults may be closely associated with skeletal muscle aging. A reduction in GLUT4 expression as a result of a decrease in muscle volume causes a reduction in insulin sensitivity in aging skeletal muscle (Joo *et al.*, 2022). The results of this study showed a correlation of waist circumference and hip circumference with handgrip strength. This is in line with the study of Widjaja *et al.* (Widjaja *et al.*, 2018), who stated that there was a significant correlation of waist circumference and hip circumference with handgrip strength ($p < 0.001$, $r = 0.302$; and $p < 0.001$, $r = 0.296$). A greater waist circumference is associated with a greater handgrip strength in males (Hardy *et al.*, 2013).

In the multiple regression analysis of HGS, we removed the data on body weight because of the occurrence of multicollinearity between weight and BMI, with a correlation coefficient of more than 0.9, indicating an extremely strong relationship between these two variables. Conceptually, only one of the variables is used in multivariate analysis. The multiple regression analysis included age, height, BMI, waist circumference, and hip circumference, with HGS as the dependent variable in a stepwise manner. After three iterations, only height, BMI, and hip circumference played a role in predicting HGS by means of the regression formula, which resulted in $R^2 = 0.19$. This signifies that for known values of height, BMI, and hip circumference, we can predict the HGS value 19% more accurately than without knowledge of these data. For a clinical diagnosis, a minimum increase of 10% is needed. This study found that height, BMI, and hip circumference play a role in predicting handgrip strength through the regression formula, i.e., $\text{handgrip strength} = -36.98 + 0.48 \times \text{height} + 0.79 \times \text{BMI} - 0.29 \times \text{hip circumference}$. Handgrip strength and waist circumference are surrogate muscle strength and visceral adiposity markers, respectively (Nakanishi *et al.*, 2024). Waist circumference is also more sensitive in determining body fat distribution, particularly in the abdomen. Waist circumference correlates better with abdominal fat distribution than BMI (Rondhianto *et al.*, 2024). Waist circumference and hip circumference are associated with obesity, in which the fat content is higher, and

the skeletal muscle and fascicular content is lower, thereby resulting in a lower contractile capacity of skeletal muscle (Afaq *et al.*, 2019). The limitation of this study is that it did not evaluate other factors that may affect handgrip strength, such as nutrient intake. From the anthropometric measures of the hands and feet in this study, the causes and effects of the decrease in handgrip strength could not be determined because this study used a cross-sectional design.

Conclusion

Three factors may be used to estimate handgrip strength in PWP, namely height, BMI, and hip circumference, whereas blood glucose concentration is not correlated with handgrip strength in PWP. It is recommended to pay attention to height, BMI, and hip circumference, particularly in the acceptance criteria of PWP candidates, and to conduct further studies with consideration of other factors that affect handgrip strength, such as food intake and anthropometric features of the hands and arms.

Acknowledgments

Financial support was provided by Universitas Trisakti for this research.

References

- Afaq, E., Hassan, S.H., Khan, M.R., Zaidi, D.A., & Baktashi, H., 2019. Correlation of Hand Dynamometry With Body Mass Index and Waist-Hip Ratio in Medical Students. *Pak J Physiol.*, 15, pp.19-21.
- Agtuahene, M.A., Quartey, J., & Kwakye, S., 2023. Influence of Hand Dominance, Gender, and Body Mass Index on Hand Grip Strength. *S Afr J Physiother.*, 79, pp.1-6.
- Al-Asadi, J.N., 2018. Handgrip Strength in Medical Students: Correlation with Body Mass Index and Hand Dimensions. *Asian Journal of Medical Sciences*, 9, pp.21-26.
- Badan Pusat Statistik Kota Administrasi Jakarta Barat., 2020. *Kecamatan Cengkareng Dalam Angka*, pp.1-114.
- Chen, D., Zhu, Y., Ni, W., Li, Y., Yin, G., Shao, Z., & Zhu, J., 2023. Hand Grip Strength is Inversely Associated with Total Daily Insulin Dose Requirement in Patients with Type 2 Diabetes Mellitus: A Cross-Sectional Study. *PeerJ.*, pp.1-13.
- Cooper, R., Tomlinson, D., Hamer, M., & Pereira, S.M.P., 2022. Lifetime Body Mass Index and Grip Strength at Age 46 Years: The 1970 British Cohort Study. *J Cachexia Sarcopenia Muscle*, 13, pp.1-10.
- Fauzi, L., Handayani, O.W.K., Susilo, M.T., Kurnia, A.R., Rahayu, S.R., Irawan, F.A., Horng Lu, F.J., Lin, C., Lai, M.F., & Yu, Y.C., 2022. Obesity in Indonesian and Taiwanese Adolescents Related to Self Perception, Diet, Exercise, and Body Image. *KEMAS*, 17(3), pp.453-461.
- Gamayanti, K.A.V., Aryana, I.G.P.S., & Gotera, W., 2023. Hubungan Antara Kekuatan Genggaman Tangan Dengan Kadar Gula Darah Sewaktu Pada Lansia Di Desa Melinggih, Kecamatan Payangan, Kabupaten Gianyar. *Intisari Sains Medis*, 14(2), pp.483-488.
- Giglio, B.M., Mota, J.F., Wall, B.T., & Pimentel, G.D., 2018. Low Handgrip Strength is Not Associated With Type 2 Diabetes Mellitus and Hyperglycemia: A Population-Based Study. *Clin Nutr Res.*, 7(2), pp.112-116.
- Hammed, A.I., & Obaseki, C.O., 2017. Interdependence of Body Mass Index with Handgrip Strength and Endurance Among Apparently Healthy Teenagers. *Turk J Kin.*, 4, pp.1-7.
- Handayani, M.D.N., Sadewa, A.H., Farmawati, A., & Rochmah, W., 2018. Anthropometric Prediction Equations for Estimating Muscle Mass of Elderly Women. *KEMAS*, 14(2), pp.195-204.
- Hardy, R., Cooper, R., Sayer, A.A., Shlomo, Y.B., Cooper, C., Deary, I.J., Demakakos, P., Gallacher, J., Martin, R.M., McNeill, G., Starr, J.M., Steptoe, A., Syddall, H., & Kuh, D., 2013. Body Mass Index, Muscle Strength, and Physical Performance in Older Adults From Eight Cohort Studies: The Halcyon Programme. *PLoS One*, 8(2), pp.e56483.
- Heidy., Djuartina, T., & Irawan, R., 2019. Korelasi Kekuatan Genggaman Tangan Dengan Karakter Antropometri Lengan Bawah dan Tangan Serta Indeks Massa Tubuh. *Damianus Journal of Medicine*, 18, pp.1-7.
- Ilmi, A.F., & Utari, D.M., 2020. Hubungan Lingkar Pinggang dan Rasio Lingkar Pinggang-Panggul (RLPP) Terhadap Kadar Glukosa Darah Puasa Pada Mahasiswa. *Journal of Nutrition College*, 9, pp.222-227.
- Jaafar, M.H., Ismail, R., Ismail, N.H., Isa, Z.M., Tamil, A.M., Nasir, N.M., Keat, N.K., Razak, N.H., Abidin, N.Z., & Yusof, K.H., 2023. Normative Reference Values and Predicting Factors of Handgrip Strength For Dominant

- and Non-Dominant Hands Among Healthy Malay Adults in Malaysia. *BMC Musculoskeletal Disord.*, 24, pp.1-9.
- Joo, K.C., Son, D.H., & Park, J.M., 2022. Association Between Relative Handgrip Strength and Insulin Resistance in Korean Elderly Men Without Diabetes: Findings Of The 2015 Korea National Health Nutrition Examination Survey. *Korean J Fam Med.*, 43, pp.199-205.
- Krakauer, N.Y., & Krakauer, J.C., 2020. Association of Body Shape Index (ABSI) with Hand Grip Strength. *Int J Environ Res Public Health*, 17, pp.1-12.
- Kurniawan, A., Widjaja, D., & Lugito, N.P.H., 2018. *Mean and Cutoff Value of Hand Grip Strength For Healthy Indonesian Young Adults*. World Congress on Osteoporosis, Osteoarthritis and Musculoskeletal Diseases; Krakow, Poland.
- Leong, D.P., Teo, K.K., Rangarajan, S., Kutty, R., Lanas, F., Hui, C., Quanyong, X., Zhenzhen, Q., Jinhua, T., Noorhassim, I., AlHabib, K.F., Moss, S.J., Rosengren, A., Akalin, A.A., Rahman, O., Chifamba, J., Orlandini, A., Kumar, R., Yeates, K., Gupta, R., Yusufali, A., Dans, A., Avezum, A., Lopez-Jaramillo, P., Poirier, P., Heidari, H., Zatonska, K., Iqbal, R., Khatib, R., & Yusuf, S., 2016. Reference Ranges of Handgrip Strength From 125,462 Healthy Adults in 21 Countries: A Prospective Urban Rural Epidemiologic (PURE) Study. *J Cachexia Sarcopenia Muscle*, 7, pp.535-546.
- Mainous, A.G., Tanner, R.J., Anton, S.D., & Jo, A., 2015. Grip Strength as A Marker of Hypertension and Diabetes in Healthy Weight Adults. *Am J Prev Med*, 49(6), pp.850-858.
- Manoharan, V.S., Sundaram, S.G., & Jason, J.I., 2015. Factors Affecting Hand Grip Strength and Its Evaluation: A Systemic Review. *Int J Physiother Res.*, 3(6), pp.1288-1293.
- McGrath, R., Johnson, N., Klawitter, L., Mahoney, S., Trautman, K., Carlson, C., Rockstad, E., & Hackney, K.J., 2020. What Are The Association Patterns Between Handgrip Strength and Adverse Health Conditions? A Topical Review. *SAGE Open Medicine*, 8, pp.1-12.
- Nakanishi, S., Shimoda, M., Kimura, T., Sanada, J., Fushimi, Y., Iwamoto, Y., Iwamoto, H., Dan, K., Mune, T., Kaku, K., & Kaneto, H., 2024. The Impact of Handgrip Strength and Waist Circumference on Glycemic Control: Prospective, Observational Study Using Outpatient Clinical Data in Japanese Patients With Type 2 Diabetes Mellitus. *J Diabetes Investig.*, 15(7), pp.892-898.
- Nathania, N.P.S., Nugraha, M.H.S., Primayanti, I.D.A., & Saraswati, P.A.S., 2023. Handgrip Strength is Associated With Balance in Elderly Women. *Majalah Ilmiah Fisioterapi Indonesia*, 11, pp.70-75.
- Gubernur Provinsi Daerah Khusus Ibukota Jakarta., 2017. *Peraturan Gubernur Provinsi Daerah Khusus Ibukota Jakarta Nomor 7 tahun 2017 Tentang Penanganan Prasarana dan Sarana Umum Tingkat Kelurahan*. Jakarta 2017.
- Richter, E.A., & Hargreaves, M., 2013. Exercise, GLUT4, and Skeletal Muscle Glucose Uptake. *Physiol Rev.*, 93(3), pp.993-1017.
- Rondhianto., Ridla, A.Z., & Hasan, H., 2024. Sociodemographic Factors Affecting Diabetic Dietary Behavior in People with Type 2 Diabetes Mellitus. *KEMAS*, 19(3), pp.429-437.
- Safitri, I., Setyarsih, L., Susanto, H., Suhartono., & Fitranti, D.Y., 2020. The Effect of Low and High Glycemic Load Diet on Muscle Fatigue of Young Soccer Athletes. *KEMAS*, 16(1), pp.111-120.
- Schrager, M.A., Metter, E.J., Simonsick, E., Ble, A., Bandinelli, S., Lauretani, F., & Ferrucci, L., 2007. Sarcopenic Obesity and Inflammation in The Inchiati Study. *J Appl Physiol.*, 102(3), pp.919-925.
- Sirajudeen, M.S., Shah, U.N., Pillai, P.S., Mohasin, N., & Shantaram, M., 2012. Correlation Between Grip Strength and Physical Factors in Men. *International Journal of Health and Rehabilitation Sciences*, 1, pp.58-63.
- Soenyoto, T., Mollap, K., & Mungpong, J., 2017. Correlation of Energy and Protein Consumption Levels with Physical Endurance of Rhythmic Gymnast Athletes. *KEMAS*, 13(1), pp.131-136.
- Soraya, N., & Parwanto, E., 2023. The Controversial Relationship Between Body Mass Index and Handgrip Strength in The Elderly: An Overview. *Malays J Med Sci.*, 30, pp.73-83.
- Vaishya, R., Misra, A., Vaish, A., Ursino, N., & D'Ambrosi, R., 2024. Hand Grip Strength as A Proposed New Vital Sign Of Health: A Narrative Review of Evidences. *Journal of Health, Population and Nutrition*, 43, pp.1-14.
- Widjaja, D., Tjhai, K., Dermawan, K., Kalim, J., Tanzil, S., Pramusinta, A., Phenca, H.K., Kurniawan, A., 2018. *Correlations Between Hand Grip Strength and Anthropometric Status in Young Indonesian Adults*. World

Congress on Osteoporosis, Osteoarthritis
and Musculoskeletal Diseases; Krakow,
Poland.

Zaccagni, L., Toselli, S., Bramanti, B., Gualdi-Russo,
E., Mongillo, J., & Rinaldo, N., 2020. Handgrip
Strength in Young Adults: Association With
Anthropometric Variables and Laterality. *Int
J Environ Res Public Health*, 17, pp.1-18.

8. Bukti konfirmasi perbaikan proofreading
dan hasil perbaikan proofreading
(28 September 2025)



Alvina fk <dr.alvina@trisakti.ac.id>

Dummy Article

3 messages

Jurnal Kemas <kemas@mail.unnes.ac.id>

Tue, Sep 23, 2025 at 6:53 PM

To: Alvina fk <dr.alvina@trisakti.ac.id>, pusparini@trisakti.ac.id, mario@trisakti.ac.id, yasminedrpk@gmail.com

Berikut kami kirimkan *Dummy* artikel Bapak/Ibu yang telah dilayout, yang akan diterbitkan di Jurnal KEMAS Vol. 21 No. 2

Mohon untuk mengecek dengan seksama artikel tersebut.

Jika Bapak/Ibu sudah setuju, mohon mengkonfirmasi persetujuan cetak. Jika masih ada kekeliruan mohon untuk menuliskan perbaikannya via email: kemas@mail.unnes.ac.id.

Kami tunggu konfirmasinya maksimal hari Sabtu, 27 September 2025 Jam 12.00 WIB.

Jika dalam waktu tersebut tidak ada konfirmasi, maka kami anggap bahwa Bapak/ Ibu telah menyetujui artikel tersebut.

Konfirmasi setelah tanggal dan jam tersebut tidak dapat dilayani.

Terima kasih

KEMAS Journal

F5 Building, 2nd Floor, Public Health Department, Sport Science Faculty, Semarang State University, Semarang, Central Java, Indonesia, 50229

<http://journal.unnes.ac.id/nju/index.php/kemas>

DISCLAIMER

This email may contain confidential or copyrighted information of UNNES. If you are not the intended recipient, please do not use or share this email. If received in error, please notify the sender and delete it. Check for viruses; UNNES is not liable for virus-related damages.

 **11_25917 Alvina 360 368.pdf**
152K**Alvina fk** <dr.alvina@trisakti.ac.id>

Thu, Sep 25, 2025 at 1:50 PM

To: Jurnal Kemas <kemas@mail.unnes.ac.id>

Kepada Yth,
Sekretariat Jurnal Kemas

Terima kasih atas kiriman dummy artikel, setelah saya periksa ada beberapa yang beda dengan naskah asli. Terlampir saya kirimkan koreksi dummy artikel dan artikel asli (dalam bentuk word).

Sudah saya tandai untuk dummy artikel yg berbeda dan di naskah asli sudah saya tandai juga dengan warna hijau.

Terima kasih.

Salam,

Alvina

[Quoted text hidden]

2 attachments



Input Koreksi 11_25917 Alvina 360 368.pdf
141K



Alvina-Factor+associated+with+handgripstrength (1).docx
43K

Jurnal Kemas <kemas@mail.unnes.ac.id>
To: Alvina fk <dr.alvina@trisakti.ac.id>

Sun, Sep 28, 2025 at 10:56 PM

Baik, akan kami sampaikan ke layouter.

Terima kasih

KEMAS Journal

F5 Building, 2nd Floor, Public Health Department, Sport Science Faculty, Semarang State University, Semarang, Central Java, Indonesia, 50229

<http://journal.unnes.ac.id/nju/index.php/kemas>

[Quoted text hidden]

[Quoted text hidden]



UNIVERSITAS TRISAKTI

"Is a one stop learning for sustainable development"

Kampus A, Jl. Kyai Tapa No.1, Grogol

Jakarta Barat 11440 - INDONESIA

www.trisakti.ac.id

(t) +62-21.566 3232, (f) +62-21.567 3001

[Quoted text hidden]



Handgrip Strength of Public Works Personnel in West Jakarta

Alvina^{1✉}, Pusparini¹, Mario¹, Yasmine Mashabi¹

¹Department of Clinical Pathology, Faculty of Medicine, Universitas Trisakti

Article Info

Article History:

Submitted: May 2025

Accepted: September 2025

Published: October 2025

Keywords:

Handgrip strength; Body mass index; Hip circumference; Waist circumference; Fasting blood glucose

DOI

<https://doi.org/10.15294/kemas.v21i2.25917>

Abstract

Public Works Personnel (PWP) in Jakarta perform their routine duties manually instead of using equipment. Handgrip strength (HGS) measures maximum hand strength as a quality indicator for muscle strength and mass. Factors affecting muscle strength are age, sex, body mass index (BMI), waist circumference (WC), and hip circumference (HC). Increased fasting blood glucose (FBG) is also associated with muscle quality, muscle strength, and physical performance. This study aimed to determine the factors associated with handgrip strength in PWP. A total of 192 male PWP from the Cengkareng district were recruited by simple random sampling. The collected data were demographics, BMI, FBG, WC, HC, and HGS. Statistical analysis was done using Pearson's correlation test and multiple regression analysis. There was a significant positive correlation of BMI, WC, and HC with HGS ($p=0.006$; $r=0.20$, $p=0.009$; $r=0.19$, and $p=0.005$; $r=0.20$) and a significant negative correlation between age and HGS ($p=0.008$; $r=-0.19$) but not between FBG and HGS ($p=0.847$). Multiple regression analysis showed height, BMI, and HC positively predicting HGS ($R^2=0.19$). Handgrip strength of public works personnel is associated with BMI, age, waist circumference, and hip circumference, but not with fasting blood glucose. Height, BMI, and hip circumference predict handgrip strength

Introduction

Personnel for the management of public infrastructure and facilities at the village level are workers who manage village public works for a given time period on the basis of a work order. The tasks of village-level public works personnel (PWP) consist of managing the infrastructure and facilities, such as streets, drains, and parks, including installed lights (Gubernur Provinsi Daerah Khusus Ibukota Jakarta, 2017). The work performed by village PWP consists mainly of intense physical activity involving the hands, such that handgrip strength is essential in their tasks. Lowering of handgrip strength may indicate reduced muscle strength, which may especially interfere with the daily activities of village PWP that consist mainly of manual labor.

Handgrip strength measures maximal hand strength and indicates muscle quality that reflects muscle strength and mass. Muscle

strength refers to the ability of muscle groups to raise a given load. Low or poor grip strength is frequently associated with damaging health impacts, such as chronic disease, and is the initial sign of reduced muscle mass, limitations in physical functions, and mortality risk. Poor handgrip strength may also reduce a person's ability to perform activities of daily living and, therefore, impact quality of life (Soraya *et al.*, 2023; Nathania *et al.*, 2023). Muscle mass accounts for approximately 40% of total weight, such that knowledge of muscle mass conditions is essential in maintaining health and activities (Handayani *et al.*, 2018).

Handgrip strength may represent the strength of all muscles in the body. Several factors possibly influencing an individual's muscle strength are age, sex, and body mass index (BMI) (Soraya *et al.*, 2023). Obesity may also affect BMI. Obesity in adults is not only associated with genetic factors but also with

✉ Correspondence Address:
Department of Clinical Pathology, Faculty of Medicine, Universitas Trisakti
Email: dr.alvina@trisakti.ac.id

environmental factors during development (Fauzi *et al.*, 2022). Handgrip strength is also associated with chronic diseases such as diabetes, metabolic syndrome, and cardiovascular disease. Increased glucose concentration is also associated with disorders of muscle quality and strength, as well as physical performance (Chen *et al.*, 2023). A reduction in muscle mass also affects an individual's strength and physical endurance. A combination of proteins can maintain muscle function, thereby building body strength and physical endurance (Soenyoto *et al.*, 2017). In the present study, we included fasting blood glucose concentrations in the data collection to obtain more accurate measures of glucose concentrations that have not yet been influenced by food intake.

Handgrip strength may be a health indicator that provides information on a person's health (Vaishya *et al.*, 2024). However, its clinical significance in determining the prevention and treatment of poor handgrip remains limited because of the still unclear etiology of muscle weakness, especially because the clinical outcomes associated with handgrip strength are extremely diverse (McGrath *et al.*, 2020). The relationship between BMI and handgrip strength is still controversial, with several investigators finding a significant positive association and others reporting a non-significant relationship (Al Asadi, 2018). There are few studies on handgrip strength, especially on workers in the sanitation sector, such as public works personnel, which is why the present investigators were interested in studying this topic. The present study aimed to determine the factors associated with handgrip strength in public works personnel.

Method

This cross-sectional study involving male public works personnel in Cengkareng District, West Jakarta, was conducted from September to October 2024. The study sample was selected by simple random sampling. Calculation of the sample size for a correlation with $r=0.2$, $Z\alpha=0.05$, and $Z\beta=0.1$ resulted in a sample of 192 subjects who met the inclusion criterion of being male and 25-60 years old, and the exclusion criteria of having an abnormality of the hands, heart disease, pulmonary disease,

and malignancies.

Measurements were obtained on height in cm, weight in kg, as well as waist circumference and hip circumference in cm. Height and weight were determined using microtoise and digital scales, respectively. BMI was calculated using the formula of weight divided by height squared. Measurement of waist and hip circumferences was done using a measuring tape. Waist circumference was determined with the subject in the upright position, at the end of a normal expiration, at the smallest abdominal circumference between the iliac crest and twelfth rib. The cut-off values for waist circumference in males and females are ≥ 90 cm and ≥ 80 cm, respectively. Hip circumference was measured using a measuring tape that was applied around the hips at the level of the symphysis pubis and the maximal gluteal circumference (Ilmi *et al.*, 2020).

Fasting blood glucose (FBG) concentration was determined on venous blood by the hexokinase method after the respondents had fasted for approximately 10 hours prior. Handgrip strength was measured with a handgrip strength dynamometer to the nearest kilogram as follows: The respondent was asked to sit down comfortably without reclining and with the feet completely relaxed. The hips and knees were positioned at a 90° angle, the gripping hand with the shoulder in adduction, the elbow flexed 90° , and the forearm and wrist in the neutral position. The gripping duration was 3 seconds. The respondent gripped the instrument with the dominant hand as hard as possible, and the value indicated by the needle was recorded. In accordance with the protocol recommended by the American Society of Hand Therapists (ASHT), the recorded handgrip strength was the average of three handgrip strength tests (Nathania *et al.*, 2023).

Before data analysis by means of the correlation test, the normality of the data distribution was determined using the Kolmogorov-Smirnov test. The Pearson correlation test was used for a normal data distribution, whereas for a non-normal data distribution, the Spearman correlation test was done, followed by a multiple regression analysis. Ethics approval was issued by the Ethics Committee of the Faculty of Medicine,

Universitas Trisakti, under No. 033/KER/FK/08/2024.

Result and Discussion

In this study, the respondents comprised 192 male public works personnel of Cengkareng District who consisted of the villages (subdistricts) of Kedaung, Duri Kosambi, West Cengkareng, Kapuk, and Rawa Buaya. A total of

26% of the respondents were residents of Duri Kosambi, where Muslims comprised 98.4% of the population. The majority of the respondents, at 77.6% had an educational level of senior high school. Nearly all of the respondents, or 99%, were right-handed, using the right hand as the dominant one at work, and most were of the Betawi (Jakarta) ethnic group at 51.6% (Table 1).

Table 1. Demographic Data of the Respondents

Characteristic	N (%)
Village	
Kedaung	40 (20.8)
Duri Kosambi	50 (26)
Cengkareng Barat	33 (17.2)
Kapuk	43 (22.4)
Rawa Buaya	26 (13.5)
Religion	
Islamic	189 (98.4)
Christian	2 (1)
Catholic	1 (0.5)
Level of education	
Elementary school	7 (3.6)
Junior high school	29 (15.1)
Senior high school	149 (77.6)
Academy-University	7 (3.6)
Dominant hand	
Right hand	190 (99)
Left hand	2 (1)
Ethnic group	
Javanese	53 (27.6)
Betawi	99 (51.6)
Sundanese	25 (13)
Other	14 (7.3)

Legend: Other ethnicities comprised groups from Palembang, Lampung, Bima, and Kupang
Source: primary data 2024 (questionnaires and interviews with respondents)

Table 2. Respondents' Characteristics

Variable	Mean ± SD
Age (years)	40.9 ± 8.9
Height (cm)	164.6 ± 6.7
Weight (kg)	64.7 ± 13.1
BMI (kg/m²)	23.8 ± 4.3
Waist circumference (cm)	85.4 ± 10.4
Hip circumference (cm)	96.9 ± 8.4

HGS (kg)	33.1 ± 6.9
FBG (mg/dL)	87 (64-280)*

Legend: BMI: body mass index. *FBG: fasting blood glucose, expressed as Median (Min-Max).

HGS: handgrip strength

Source: primary data (respondents' measurements)

Table 3. Analysis of the Relationship of Age, Height, Weight, BMI, Waist Circumference, Hip Circumference, and Fasting Blood Glucose Concentration with Handgrip Strength

Variable	R	P value
Age	- 0.19	0.008*
Height	0.36	0.000*
Weight	0.32	0.000*
BMI	0.20	0.006*
Waist circumference	0.19	0.009*
Hip circumference	0.20	0.005*
FBG	0.02	0.847 [#]

*P value with Pearson's test, [#] Spearman's test

Source: primary data

Table 4. Multiple Regression Analysis on Handgrip Strength

Determinant	b	SE	p
Height	0.48	0.08	0.000
BMI	0.79	0.22	0.000
Hip circumference	- 0.29	0.12	0.015
Constant	- 36.98		

R²=0,19

Source: primary data

The bivariate analysis found a significant positive correlation of BMI, height, weight, waist circumference, and hip circumference with handgrip strength ($p=0.006$, $r=0.20$; $p<0.001$, $r=0.36$; $p<0.001$, $r=0.32$; $p=0.009$, $r=0.19$; $p=0.005$; $r=0.20$, respectively). This showed that the larger the BMI, height, weight, waist circumference, and hip circumference, the stronger the handgrip. However, there was no significant relationship between fasting glucose concentration and handgrip strength ($p=0.847$). This study also found that age had a significant negative correlation with handgrip strength ($p=0.011$; $r=-0.182$), signifying that the older a person, the lower the handgrip strength (Table 3). Results of multiple regression analysis on handgrip strength are listed in Table 4.

This study found that most of the public works personnel in Duri Kosambi village were Muslims, had a senior high school level of education, and were of Betawi ethnicity. This was in line with the demographic data in the Cengkareng district, in that Duri Kosambi was

the village with the greatest area, most of the residents were male, and the Islamic religion had the most believers (Badan Pusat Statistik Jakarta Barat, 2020). Among our respondents, more than ten percent were in the diabetes category. Socio-demographic factors capable of affecting dietary habits in patients with diabetes are the level of education and the type of family (Rondhianto *et al.*, 2024). Among the respondents of this study, the dominant right hand was used for working. In the adult population, a total of 90% of the residents preferred to use the right hand for manual tasks, whereas around 10% used the left hand (Zaccagni *et al.*, 2020). According to Agtuahene *et al.* (Agtuahene *et al.*, 2023), the dominant hand is generally around 10% stronger than the non-dominant hand. Many activities in daily life need a high level of exertion of the muscles of the hand and forearm (Hammed *et al.*, 2017). The frequent use of the dominant hand in daily living results in the muscles of the hand and forearm providing strength in gripping (Afaq *et*

al., 2019).

The mean handgrip strength in this study was 33.1 ± 6.9 kg, which is relatively low if compared with the study results of Kurniawan *et al.* (Kurniawan *et al.*, 2018). In healthy young Indonesian adults, the mean handgrip strength in males was 37.37 ± 8.29 kg, and the lower limit of muscle strength in Indonesian males was 20.79 kg. The handgrip strength results in both studies are still below those of males in other Southeast Asian countries, according to the study results of Leong *et al.* (Leong *et al.*, 2016). These investigators found that the mean handgrip strength in Southeast Asian males aged 35-40 years was 40 kg (34-44 kg); for the age range of 41-50 years, the handgrip strength was 38 kg (33-44 kg), for the age range of 51-60 years it was 34 kg (30-40 kg), and for ages 61-70 years 30 kg (24-34 kg). This may have been caused by ethnic differences mediated by genetic differences that also affect individual muscle strength. The highest handgrip strength is found in the European and North American populations, whereas the lowest handgrip strength is found in African, South Asian, and Southeast Asian populations. These variations in muscle strength in various countries may also be partly caused by differences in socioeconomic status (Leong *et al.*, 2016).

According to Agtuahene *et al.* (Agtuahene *et al.*, 2023), handgrip strength is a physiological variable influenced by several factors, such as age, sex, and body measurements. In contrast, according to Zaccagni *et al.* (Zaccagni *et al.*, 2020), handgrip strength, apart from being affected by age, may also be influenced by physical activity, grip measurement, height, and body mass. The present study found a significant positive correlation between BMI and handgrip strength. According to Hammed *et al.*, this may be caused by the higher percentage of skeletal muscle mass compared to the percentage of fat mass, thereby resulting in a higher handgrip strength [13]. Our study's results are identical to those of Agtuahene *et al.* (Agtuahene *et al.*, 2023), who found a significant positive correlation ($r=0.290$) of handgrip strength with BMI and height. This shows that taller individuals with higher BMI apparently have greater handgrip strength. Our study is also in line with the study of Krakauer in 2020, which

showed a positive correlation between BMI and handgrip strength (Krakauer *et al.*, 2020).

The study of Cooper *et al.* (Cooper *et al.*, 2022) concluded that a high childhood BMI is associated with higher grip strength at age 46 years. This is because fat mass acts as a mechanical load that elicits an anabolic response that stimulates muscle growth and function. This anabolic response is usually greater in males than in females because of the higher testosterone concentration. However, it should also be noted that BMI does not differentiate between fat mass and lean mass as adipocyte indicators that may vary with sex. It should be noted that BMI is a measure of body fat that does not differentiate between weight changes because of an increase or decrease in muscle mass or body fat (Zaccagni *et al.*, 2020). According to Jafar *et al.* (Jaafar *et al.*, 2023), heavier individuals have a greater muscle mass, resulting in greater handgrip strength, than lighter individuals. In the study of Heidy *et al.* (Heidy *et al.*, 2019), the investigators found a positive correlation between height and forearm length. The study by Sirajudeen *et al.* (Sirajudeen *et al.*, 2012) found a positive correlation between hand length and handgrip strength. This shows that height affects the longer extremity or arm, thus approaching optimal muscle length that may result in greater grip strength. According to a prospective cohort study, handgrip strength in middle age appears to be associated with birth weight and the increase in height at puberty. Pubertal growth and handgrip strength possibly represent the development of muscle size and growth (Schrager *et al.*, 2007).

Our study found no relationship between blood glucose concentration and handgrip strength. Our study agrees with the study of Giglio *et al.* (Giglio *et al.*, 2018), which showed no significant relationship between muscle mass strength and blood glucose concentration. According to Giglio, this is because most respondents are individuals who routinely perform structured physical activity, even though they are in the light category. This is similar to our study respondents, who were public works personnel and who had performed physical activity such as street sweeping and cleaning storm drains. Glucose is the main fuel

for muscle contraction, entering the muscle cells by diffusion through the GLUT4 glucose transporter, which, after muscle contraction, moves from its intracellular storage site to the plasma membrane and T-tubules (Richter *et al.*, 2013). Glucose metabolism has an influence on handgrip strength through various mechanisms, such as disorders of glucose metabolism, particularly glycogenolysis, that may result in the loss of muscle strength. Insulin resistance in hyperglycemia may cause muscle degradation because hyperglycemia causes low muscle strength through effects on skeletal muscle mitochondria. The lowering of muscle strength by hyperglycemia occurs through Krüppel-like factor 15, which regulates skeletal muscle lipid flux. Hyperglycemia also increases pro-inflammatory cytokines, which may increase the catabolism that plays a role in the further decrease in muscle mass and quality (Gamayanti *et al.*, 2023). The amount of carbohydrates also influences the glycemic response and post-prandial insulin level, impacting the supply of glycogen to the muscles (Safitri *et al.*, 2020).

This study has a significant negative correlation between age and handgrip strength, which decreases with age (Manoharan *et al.*, 2015). Handgrip strength initially experiences an increase with increasing age, reaches its peak between the ages of 30 and 40 years, and decreases afterward (Zaccagni *et al.*, 2020). The age-related changes in skeletal muscle composition, particularly the accumulation of lipids in skeletal muscle fibers, play a role in the poor quality of the muscles and may result in metabolic abnormalities such as insulin resistance. Apart from changes in muscle quality, there is also a reduction in muscle mass that consistently occurs during aging. The increase in oxidative stress and chronic inflammation that commonly occurs in the aging process apparently also plays a role in the development of decreased muscle mass and strength (Mainous *et al.*, 2015). The decrease in muscle mass in the elderly corresponds to a 10% reduction in grip strength per decade, starting from the age of 40 years (Jaafar *et al.*, 2023). Skeletal muscle plays an important role in glucose metabolism and has a significant influence on insulin sensitivity. Increased

insulin resistance in older adults may be closely associated with skeletal muscle aging. A reduction in GLUT4 expression as a result of a decrease in muscle volume causes a reduction in insulin sensitivity in aging skeletal muscle (Joo *et al.*, 2022). The results of this study showed a correlation of waist circumference and hip circumference with handgrip strength. This is in line with the study of Widjaja *et al.* (Widjaja *et al.*, 2018), who stated that there was a significant correlation of waist circumference and hip circumference with handgrip strength ($p < 0.001$, $r = 0.302$; and $p < 0.001$, $r = 0.296$). A greater waist circumference is associated with a greater handgrip strength in males (Hardy *et al.*, 2013).

In the multiple regression analysis of HGS, we removed the data on body weight because of the occurrence of multicollinearity between weight and BMI, with a correlation coefficient of more than 0.9, indicating an extremely strong relationship between these two variables. Conceptually, only one of the variables is used in multivariate analysis. The multiple regression analysis included age, height, BMI, waist circumference, and hip circumference, with HGS as the dependent variable in a stepwise manner. After three iterations, only height, BMI, and hip circumference played a role in predicting HGS by means of the regression formula, which resulted in $R^2 = 0.19$. This signifies that for known values of height, BMI, and hip circumference, we can predict the HGS value 19% more accurately than without knowledge of these data. For a clinical diagnosis, a minimum increase of 10% is needed. This study found that height, BMI, and hip circumference play a role in predicting handgrip strength through the regression formula, i.e., $\text{handgrip strength} = -36.98 + 0.48 \times \text{height} + 0.79 \times \text{BMI} - 0.29 \times \text{hip circumference}$. Handgrip strength and waist circumference are surrogate muscle strength and visceral adiposity markers, respectively (Nakanishi *et al.*, 2024). Waist circumference is also more sensitive in determining body fat distribution, particularly in the abdomen. Waist circumference correlates better with abdominal fat distribution than BMI (Rondhianto *et al.*, 2024). Waist circumference and hip circumference are associated with obesity, in which the fat content is higher, and

the skeletal muscle and fascicular content is lower, thereby resulting in a lower contractile capacity of skeletal muscle (Afaq *et al.*, 2019). The limitation of this study is that it did not evaluate other factors that may affect handgrip strength, such as nutrient intake. From the anthropometric measures of the hands and feet in this study, the causes and effects of the decrease in handgrip strength could not be determined because this study used a cross-sectional design.

Conclusion

Three factors may be used to estimate handgrip strength in PWP, namely height, BMI, and hip circumference, whereas blood glucose concentration is not correlated with handgrip strength in PWP. It is recommended to pay attention to height, BMI, and hip circumference, particularly in the acceptance criteria of PWP candidates, and to conduct further studies with consideration of other factors that affect handgrip strength, such as food intake and anthropometric features of the hands and arms.

Acknowledgments

Financial support was provided by Universitas Trisakti for this research.

References

- Afaq, E., Hassan, S.H., Khan, M.R., Zaidi, D.A., & Baktashi, H., 2019. Correlation of Hand Dynamometry With Body Mass Index and Waist-Hip Ratio in Medical Students. *Pak J Physiol.*, 15, pp.19-21.
- Agtuahene, M.A., Quartey, J., & Kwakye, S., 2023. Influence of Hand Dominance, Gender, and Body Mass Index on Hand Grip Strength. *S Afr J Physiother.*, 79, pp.1-6.
- Al-Asadi, J.N., 2018. Handgrip Strength in Medical Students: Correlation with Body Mass Index and Hand Dimensions. *Asian Journal of Medical Sciences*, 9, pp.21-26.
- Badan Pusat Statistik Kota Administrasi Jakarta Barat., 2020. *Kecamatan Cengkareng Dalam Angka*, pp.1-114.
- Chen, D., Zhu, Y., Ni, W., Li, Y., Yin, G., Shao, Z., & Zhu, J., 2023. Hand Grip Strength is Inversely Associated with Total Daily Insulin Dose Requirement in Patients with Type 2 Diabetes Mellitus: A Cross-Sectional Study. *PeerJ.*, pp.1-13.
- Cooper, R., Tomlinson, D., Hamer, M., & Pereira, S.M.P., 2022. Lifetime Body Mass Index and Grip Strength at Age 46 Years: The 1970 British Cohort Study. *J Cachexia Sarcopenia Muscle*, 13, pp.1-10.
- Fauzi, L., Handayani, O.W.K., Susilo, M.T., Kurnia, A.R., Rahayu, S.R., Irawan, F.A., Horng Lu, F.J., Lin, C., Lai, M.F., & Yu, Y.C., 2022. Obesity in Indonesian and Taiwanese Adolescents Related to Self Perception, Diet, Exercise, and Body Image. *KEMAS*, 17(3), pp.453-461.
- Gamayanti, K.A.V., Aryana, I.G.P.S., & Gotera, W., 2023. Hubungan Antara Kekuatan Genggaman Tangan Dengan Kadar Gula Darah Sewaktu Pada Lansia Di Desa Melinggih, Kecamatan Payangan, Kabupaten Gianyar. *Intisari Sains Medis*, 14(2), pp.483-488.
- Giglio, B.M., Mota, J.F., Wall, B.T., & Pimentel, G.D., 2018. Low Handgrip Strength is Not Associated With Type 2 Diabetes Mellitus and Hyperglycemia: A Population-Based Study. *Clin Nutr Res.*, 7(2), pp.112-116.
- Hammed, A.I., & Obaseki, C.O., 2017. Interdependence of Body Mass Index with Handgrip Strength and Endurance Among Apparently Healthy Teenagers. *Turk J Kin.*, 4, pp.1-7.
- Handayani, M.D.N., Sadewa, A.H., Farmawati, A., & Rochmah, W., 2018. Anthropometric Prediction Equations for Estimating Muscle Mass of Elderly Women. *KEMAS*, 14(2), pp.195-204.
- Hardy, R., Cooper, R., Sayer, A.A., Shlomo, Y.B., Cooper, C., Deary, I.J., Demakakos, P., Gallacher, J., Martin, R.M., McNeill, G., Starr, J.M., Steptoe, A., Syddall, H., & Kuh, D., 2013. Body Mass Index, Muscle Strength, and Physical Performance in Older Adults From Eight Cohort Studies: The Halcyon Programme. *PLoS One*, 8(2), pp.e56483.
- Heidy., Djuartina, T., & Irawan, R., 2019. Korelasi Kekuatan Genggaman Tangan Dengan Karakter Antropometri Lengan Bawah dan Tangan Serta Indeks Massa Tubuh. *Damianus Journal of Medicine*, 18, pp.1-7.
- Ilmi, A.F., & Utari, D.M., 2020. Hubungan Lingkar Pinggang dan Rasio Lingkar Pinggang-Panggul (RLPP) Terhadap Kadar Glukosa Darah Puasa Pada Mahasiswa. *Journal of Nutrition College*, 9, pp.222-227.
- Jaafar, M.H., Ismail, R., Ismail, N.H., Isa, Z.M., Tamil, A.M., Nasir, N.M., Keat, N.K., Razak, N.H., Abidin, N.Z., & Yusof, K.H., 2023. Normative Reference Values and Predicting Factors of Handgrip Strength For Dominant

- and Non-Dominant Hands Among Healthy Malay Adults in Malaysia. *BMC Musculoskeletal Disord.*, 24, pp.1-9.
- Joo, K.C., Son, D.H., & Park, J.M., 2022. Association Between Relative Handgrip Strength and Insulin Resistance in Korean Elderly Men Without Diabetes: Findings Of The 2015 Korea National Health Nutrition Examination Survey. *Korean J Fam Med.*, 43, pp.199-205.
- Krakauer, N.Y., & Krakauer, J.C., 2020. Association of Body Shape Index (ABSI) with Hand Grip Strength. *Int J Environ Res Public Health*, 17, pp.1-12.
- Kurniawan, A., Widjaja, D., & Lugito, N.P.H., 2018. *Mean and Cutoff Value of Hand Grip Strength For Healthy Indonesian Young Adults*. World Congress on Osteoporosis, Osteoarthritis and Musculoskeletal Diseases; Krakow, Poland.
- Leong, D.P., Teo, K.K., Rangarajan, S., Kutty, R., Lanas, F., Hui, C., Quanyong, X., Zhenzhen, Q., Jinhua, T., Noorhassim, I., AlHabib, K.F., Moss, S.J., Rosengren, A., Akalin, A.A., Rahman, O., Chifamba, J., Orlandini, A., Kumar, R., Yeates, K., Gupta, R., Yusufali, A., Dans, A., Avezum, A., Lopez-Jaramillo, P., Poirier, P., Heidari, H., Zatonska, K., Iqbal, R., Khatib, R., & Yusuf, S., 2016. Reference Ranges of Handgrip Strength From 125,462 Healthy Adults in 21 Countries: A Prospective Urban Rural Epidemiologic (PURE) Study. *J Cachexia Sarcopenia Muscle*, 7, pp.535-546.
- Mainous, A.G., Tanner, R.J., Anton, S.D., & Jo, A., 2015. Grip Strength as A Marker of Hypertension and Diabetes in Healthy Weight Adults. *Am J Prev Med*, 49(6), pp.850-858.
- Manoharan, V.S., Sundaram, S.G., & Jason, J.I., 2015. Factors Affecting Hand Grip Strength and Its Evaluation: A Systemic Review. *Int J Physiother Res.*, 3(6), pp.1288-1293.
- McGrath, R., Johnson, N., Klawitter, L., Mahoney, S., Trautman, K., Carlson, C., Rockstad, E., & Hackney, K.J., 2020. What Are The Association Patterns Between Handgrip Strength and Adverse Health Conditions? A Topical Review. *SAGE Open Medicine*, 8, pp.1-12.
- Nakanishi, S., Shimoda, M., Kimura, T., Sanada, J., Fushimi, Y., Iwamoto, Y., Iwamoto, H., Dan, K., Mune, T., Kaku, K., & Kaneto, H., 2024. The Impact of Handgrip Strength and Waist Circumference on Glycemic Control: Prospective, Observational Study Using Outpatient Clinical Data in Japanese Patients With Type 2 Diabetes Mellitus. *J Diabetes Investig.*, 15(7), pp.892-898.
- Nathania, N.P.S., Nugraha, M.H.S., Primayanti, I.D.A., & Saraswati, P.A.S., 2023. Handgrip Strength is Associated With Balance in Elderly Women. *Majalah Ilmiah Fisioterapi Indonesia*, 11, pp.70-75.
- Gubernur Provinsi Daerah Khusus Ibukota Jakarta., 2017. *Peraturan Gubernur Provinsi Daerah Khusus Ibukota Jakarta Nomor 7 tahun 2017 Tentang Penanganan Prasarana dan Sarana Umum Tingkat Kelurahan*. Jakarta 2017.
- Richter, E.A., & Hargreaves, M., 2013. Exercise, GLUT4, and Skeletal Muscle Glucose Uptake. *Physiol Rev.*, 93(3), pp.993-1017.
- Rondhianto., Ridla, A.Z., & Hasan, H., 2024. Sociodemographic Factors Affecting Diabetic Dietary Behavior in People with Type 2 Diabetes Mellitus. *KEMAS*, 19(3), pp.429-437.
- Safitri, I., Setyarsih, L., Susanto, H., Suhartono., & Fitranti, D.Y., 2020. The Effect of Low and High Glycemic Load Diet on Muscle Fatigue of Young Soccer Athletes. *KEMAS*, 16(1), pp.111-120.
- Schrager, M.A., Metter, E.J., Simonsick, E., Ble, A., Bandinelli, S., Lauretani, F., & Ferrucci, L., 2007. Sarcopenic Obesity and Inflammation in The Inchiati Study. *J Appl Physiol.*, 102(3), pp.919-925.
- Sirajudeen, M.S., Shah, U.N., Pillai, P.S., Mohasin, N., & Shantaram, M., 2012. Correlation Between Grip Strength and Physical Factors in Men. *International Journal of Health and Rehabilitation Sciences*, 1, pp.58-63.
- Soenyoto, T., Mollap, K., & Mungpong, J., 2017. Correlation of Energy and Protein Consumption Levels with Physical Endurance of Rhythmic Gymnast Athletes. *KEMAS*, 13(1), pp.131-136.
- Soraya, N., & Parwanto, E., 2023. The Controversial Relationship Between Body Mass Index and Handgrip Strength in The Elderly: An Overview. *Malays J Med Sci.*, 30, pp.73-83.
- Vaishya, R., Misra, A., Vaish, A., Ursino, N., & D'Ambrosi, R., 2024. Hand Grip Strength as A Proposed New Vital Sign Of Health: A Narrative Review of Evidences. *Journal of Health, Population and Nutrition*, 43, pp.1-14.
- Widjaja, D., Tjhai, K., Dermawan, K., Kalim, J., Tanzil, S., Pramusinta, A., Phenca, H.K., Kurniawan, A., 2018. *Correlations Between Hand Grip Strength and Anthropometric Status in Young Indonesian Adults*. World

Congress on Osteoporosis, Osteoarthritis
and Musculoskeletal Diseases; Krakow,
Poland.

Zaccagni, L., Toselli, S., Bramanti, B., Gualdi-Russo,
E., Mongillo, J., & Rinaldo, N., 2020. Handgrip
Strength in Young Adults: Association With
Anthropometric Variables and Laterality. *Int
J Environ Res Public Health*, 17, pp.1-18.

9. Bukti konfirmasi publikasi artikel
dan artikel yang dipublikasi
(6 Oktober 2025)



KEMAS JOURNAL EDITORIAL
DEPARTMENT OF PUBLIC HEALTH SCIENCES
MEDICAL FACULTY
UNIVERSITAS NEGERI SEMARANG
Kelud Campuss, Gajah Mungkur, Semarang 50237
Telp (024) 8440516
Laman: <http://journal.unnes.ac.id/nju/index.php/kemas>,
e-mail: kemas@mail.unnes.ac.id

No. : 2/KEMAS/VIII/2025
Attachment : -
Subject : **Eligible Article Notification**

To:

Mr./Mrs.: *Alvina, Pusparini, Mario, Yasmine Mashabi*

Based on your article which submitted to KEMAS Journal with title:

Handgrip Strength of Public Works Personnel in West Jakarta

Herewith we inform you about the result of the editorial board assessment, your article deserves to be published in KEMAS journal Vol. 21 No. 2 (October – December 2025).

Thank you for your attention and good cooperation.

Semarang, August 8th 2025

Ketua Dewan Redaksi,

Prof. Dr. dr. Oktia Woro K.H., M.Kes.
NIP. 19591001 198703 2 001



Handgrip Strength of Public Works Personnel in West Jakarta

Alvina^{1✉}, Pusparini¹, Mario¹, Yasmine Mashabi¹

¹Department of Clinical Pathology, Faculty of Medicine, Universitas Trisakti

Article Info

Article History:

Submitted: May 2025

Accepted: September 2025

Published: October 2025

Keywords:

Handgrip strength; Body mass index; Hip circumference; Waist circumference; Fasting blood glucose

DOI

<https://doi.org/10.15294/kemas.v21i2.25917>

Abstract

Public Works Personnel (PWP) in Jakarta perform their routine duties manually instead of using equipment. Handgrip strength (HGS) measures maximum hand strength as a quality indicator for muscle strength and mass. Factors affecting muscle strength are age, sex, body mass index (BMI), waist circumference (WC), and hip circumference (HC). Increased fasting blood glucose (FBG) is also associated with muscle quality, muscle strength, and physical performance. This study aimed to determine the factors associated with handgrip strength in PWP. A total of 192 male PWP from the Cengkareng district were recruited by simple random sampling. The collected data were demographics, BMI, FBG, WC, HC, and HGS. Statistical analysis was done using Pearson's correlation test and multiple regression analysis. There was a significant positive correlation of BMI, WC, and HC with HGS ($p=0.006$; $r=0.20$, $p=0.009$; $r=0.19$, and $p=0.005$; $r=0.20$) and a significant negative correlation between age and HGS ($p=0.008$; $r=-0.19$) but not between FBG and HGS ($p=0.847$). Multiple regression analysis showed height, BMI, and HC positively predicting HGS ($R^2=0.19$). Handgrip strength of public works personnel is associated with BMI, age, waist circumference, and hip circumference, but not with fasting blood glucose. Height, BMI, and hip circumference predict handgrip strength

Introduction

Personnel for the management of public infrastructure and facilities at the village level are workers who manage village public works for a given time period on the basis of a work order. The tasks of village-level public works personnel (PWP) consist of managing the infrastructure and facilities, such as streets, drains, and parks, including installed lights (Gubernur Provinsi Daerah Khusus Ibukota Jakarta, 2017). The work performed by village PWP consists mainly of intense physical activity involving the hands, such that handgrip strength is essential in their tasks. Lowering of handgrip strength may indicate reduced muscle strength, which may especially interfere with the daily activities of village PWP that consist mainly of manual labor.

Handgrip strength measures maximal hand strength and indicates muscle quality that reflects muscle strength and mass. Muscle

strength refers to the ability of muscle groups to raise a given load. Low or poor grip strength is frequently associated with damaging health impacts, such as chronic disease, and is the initial sign of reduced muscle mass, limitations in physical functions, and mortality risk. Poor handgrip strength may also reduce a person's ability to perform activities of daily living and, therefore, impact quality of life (Soraya *et al.*, 2023; Nathania *et al.*, 2023). Muscle mass accounts for approximately 40% of total weight, such that knowledge of muscle mass conditions is essential in maintaining health and activities (Handayani *et al.*, 2018).

Handgrip strength may represent the strength of all muscles in the body. Several factors possibly influencing an individual's muscle strength are age, sex, and body mass index (BMI) (Soraya *et al.*, 2023). Obesity may also affect BMI. Obesity in adults is not only associated with genetic factors but also with

✉ Correspondence Address:
Department of Clinical Pathology, Faculty of Medicine, Universitas Trisakti
Email: dr.alvina@trisakti.ac.id

environmental factors during development (Fauzi *et al.*, 2022). Handgrip strength is also associated with chronic diseases such as diabetes, metabolic syndrome, and cardiovascular disease. Increased glucose concentration is also associated with disorders of muscle quality and strength, as well as physical performance (Chen *et al.*, 2023). A reduction in muscle mass also affects an individual's strength and physical endurance. A combination of proteins can maintain muscle function, thereby building body strength and physical endurance (Soenyoto *et al.*, 2017). In the present study, we included fasting blood glucose concentrations in the data collection to obtain more accurate measures of glucose concentrations that have not yet been influenced by food intake.

Handgrip strength may be a health indicator that provides information on a person's health (Vaishya *et al.*, 2024). However, its clinical significance in determining the prevention and treatment of poor handgrip remains limited because of the still unclear etiology of muscle weakness, especially because the clinical outcomes associated with handgrip strength are extremely diverse (McGrath *et al.*, 2020). The relationship between BMI and handgrip strength is still controversial, with several investigators finding a significant positive association and others reporting a non-significant relationship (Al Asadi, 2018). There are few studies on handgrip strength, especially on workers in the sanitation sector, such as public works personnel, which is why the present investigators were interested in studying this topic. The present study aimed to determine the factors associated with handgrip strength in public works personnel.

Method

This cross-sectional study involving male public works personnel in Cengkareng District, West Jakarta, was conducted from September to October 2024. The study sample was selected by simple random sampling. Calculation of the sample size for a correlation with $r=0.2$, $Z\alpha=0.05$, and $Z\beta=0.1$ resulted in a sample of 192 subjects who met the inclusion criterion of being male and 25-60 years old, and the exclusion criteria of having an abnormality of the hands, heart disease, pulmonary disease,

and malignancies.

Measurements were obtained on height in cm, weight in kg, as well as waist circumference and hip circumference in cm. Height and weight were determined using microtoise and digital scales, respectively. BMI was calculated using the formula of weight divided by height squared. Measurement of waist and hip circumferences was done using a measuring tape. Waist circumference was determined with the subject in the upright position, at the end of a normal expiration, at the smallest abdominal circumference between the iliac crest and twelfth rib. The cut-off values for waist circumference in males and females are ≥ 90 cm and ≥ 80 cm, respectively. Hip circumference was measured using a measuring tape that was applied around the hips at the level of the symphysis pubis and the maximal gluteal circumference (Rokhmah *et al.*, 2015).

Fasting blood glucose (FBG) concentration was determined on venous blood by the hexokinase method after the respondents had fasted for approximately 10 hours prior. Handgrip strength was measured with a handgrip strength dynamometer to the nearest kilogram as follows: The respondent was asked to sit down comfortably without reclining and with the feet completely relaxed. The hips and knees were positioned at a 90° angle, the gripping hand with the shoulder in adduction, the elbow flexed 90° , and the forearm and wrist in the neutral position. The gripping duration was 3 seconds. The respondent gripped the instrument with the dominant hand as hard as possible, and the value indicated by the needle was recorded. In accordance with the protocol recommended by the American Society of Hand Therapists (ASHT), the recorded handgrip strength was the average of three handgrip strength tests (Nathania *et al.*, 2023).

Before data analysis by means of the correlation test, the normality of the data distribution was determined using the Kolmogorov-Smirnov test. The Pearson correlation test was used for a normal data distribution, whereas for a non-normal data distribution, the Spearman correlation test was done, followed by a multiple regression analysis. Ethics approval was issued by the Ethics Committee of the Faculty of Medicine,

Universitas Trisakti, under No. 033/KER/FK/08/2024.

Result and Discussion

In this study, the respondents comprised 192 male public works personnel of Cengkareng District who consisted of the villages (subdistricts) of Kedaung, Duri Kosambi, West Cengkareng, Kapuk, and Rawa Buaya. A total of

26% of the respondents were residents of Duri Kosambi, where Muslims comprised 98.4% of the population. The majority of the respondents, at 77.6% had an educational level of senior high school. Nearly all of the respondents, or 99%, were right-handed, using the right hand as the dominant one at work, and most were of the Betawi (Jakarta) ethnic group at 51.6% (Table 1).

Table 1. Demographic Data of the Respondents

Characteristic	N (%)
Village	
Kedaung	40 (20.8)
Duri Kosambi	50 (26)
Cengkareng Barat	33 (17.2)
Kapuk	43 (22.4)
Rawa Buaya	26 (13.5)
Religion	
Islamic	189 (98.4)
Christian	2 (1)
Catholic	1 (0.5)
Level of education	
Elementary school	7 (3.6)
Junior high school	29 (15.1)
Senior high school	149 (77.6)
Academy-University	7 (3.6)
Dominant hand	
Right hand	190 (99)
Left hand	2 (1)
Ethnic group	
Javanese	53 (27.6)
Betawi	99 (51.6)
Sundanese	25 (13)
Other	14 (7.3)

Legend: Other ethnicities comprised groups from Palembang, Lampung, Bima, and Kupang
Source: primary data 2024 (questionnaires and interviews with respondents)

Table 2. Respondents' Characteristics

Variable	Mean ± SD
Age (years)	40.9 ± 8.9
Height (cm)	164.6 ± 6.7
Weight (kg)	64.7 ± 13.1
BMI (kg/m²)	23.8 ± 4.3
Waist circumference (cm)	85.4 ± 10.4
Hip circumference (cm)	96.9 ± 8.4

HGS (kg)	33.1 ± 6.9
FBG (mg/dL)	87 (64-280)*

Legend: BMI: body mass index. *FBG: fasting blood glucose, expressed as Median (Min-Max).

HGS: handgrip strength

Source: primary data (respondents' measurements)

Table 3. Analysis of the Relationship of Age, Height, Weight, BMI, Waist Circumference, Hip Circumference, and Fasting Blood Glucose Concentration with Handgrip Strength

Variable	R	P value
Age	- 0.19	0.008*
Height	0.36	0.000*
Weight	0.32	0.000*
BMI	0.20	0.006*
Waist circumference	0.19	0.009*
Hip circumference	0.20	0.005*
FBG	0.02	0.847 [#]

*P value with Pearson's test, [#] Spearman's test

Source: primary data

Table 4. Multiple Regression Analysis on Handgrip Strength

Determinant	b	SE	p
Height	0.48	0.08	0.000
BMI	0.79	0.22	0.000
Hip circumference	- 0.29	0.12	0.015
Constant	- 36.98		

R²=0,19

Source: primary data

The bivariate analysis found a significant positive correlation of BMI, height, weight, waist circumference, and hip circumference with handgrip strength ($p=0.006$, $r=0.20$; $p<0.001$, $r=0.36$; $p<0.001$, $r=0.32$; $p=0.009$, $r=0.19$; $p=0.005$; $r=0.20$, respectively). This showed that the larger the BMI, height, weight, waist circumference, and hip circumference, the stronger the handgrip. However, there was no significant relationship between fasting glucose concentration and handgrip strength ($p=0.847$). This study also found that age had a significant negative correlation with handgrip strength ($p=0.011$; $r=-0.182$), signifying that the older a person, the lower the handgrip strength (Table 3). Results of multiple regression analysis on handgrip strength are listed in Table 4.

This study found that most of the public works personnel in Duri Kosambi village were Muslims, had a senior high school level of education, and were of Betawi ethnicity. This was in line with the demographic data in the Cengkareng district, in that Duri Kosambi was

the village with the greatest area, most of the residents were male, and the Islamic religion had the most believers (Badan Pusat Statistik Jakarta Barat, 2020). Among our respondents, more than ten percent were in the diabetes category. Socio-demographic factors capable of affecting dietary habits in patients with diabetes are the level of education and the type of family (Rondhianto *et al.*, 2024). Among the respondents of this study, the dominant right hand was used for working. In the adult population, a total of 90% of the residents preferred to use the right hand for manual tasks, whereas around 10% used the left hand (Zaccagni *et al.*, 2020). According to Agtuahene *et al.* (Agtuahene *et al.*, 2023), the dominant hand is generally around 10% stronger than the non-dominant hand. Many activities in daily life need a high level of exertion of the muscles of the hand and forearm (Hammed *et al.*, 2017). The frequent use of the dominant hand in daily living results in the muscles of the hand and forearm providing strength in gripping (Afaq *et*

al., 2019).

The mean handgrip strength in this study was 33.1 ± 6.9 kg, which is relatively low if compared with the study results of Kurniawan *et al.* (Kurniawan *et al.*, 2018). In healthy young Indonesian adults, the mean handgrip strength in males was 37.37 ± 8.29 kg, and the lower limit of muscle strength in Indonesian males was 20.79 kg. The handgrip strength results in both studies are still below those of males in other Southeast Asian countries, according to the study results of Leong *et al.* (Leong *et al.*, 2016). These investigators found that the mean handgrip strength in Southeast Asian males aged 35-40 years was 40 kg (34-44 kg); for the age range of 41-50 years, the handgrip strength was 38 kg (33-44 kg), for the age range of 51-60 years it was 34 kg (30-40 kg), and for ages 61-70 years 30 kg (24-34 kg). This may have been caused by ethnic differences mediated by genetic differences that also affect individual muscle strength. The highest handgrip strength is found in the European and North American populations, whereas the lowest handgrip strength is found in African, South Asian, and Southeast Asian populations. These variations in muscle strength in various countries may also be partly caused by differences in socioeconomic status (Leong *et al.*, 2016).

According to Agtuahene *et al.* (Agtuahene *et al.*, 2023), handgrip strength is a physiological variable influenced by several factors, such as age, sex, and body measurements. In contrast, according to Zaccagni *et al.* (Zaccagni *et al.*, 2020), handgrip strength, apart from being affected by age, may also be influenced by physical activity, grip measurement, height, and body mass. The present study found a significant positive correlation between BMI and handgrip strength. According to Hammed *et al.*, this may be caused by the higher percentage of skeletal muscle mass compared to the percentage of fat mass, thereby resulting in a higher handgrip strength. Our study's results are identical to those of Agtuahene *et al.* (Agtuahene *et al.*, 2023), who found a significant positive correlation ($r=0.290$) of handgrip strength with BMI and height. This shows that taller individuals with higher BMI apparently have greater handgrip strength. Our study is also in line with the study of Krakauer in 2020, which

showed a positive correlation between BMI and handgrip strength (Krakauer *et al.*, 2020).

The study of Cooper *et al.* (Cooper *et al.*, 2022) concluded that a high childhood BMI is associated with higher grip strength at age 46 years. This is because fat mass acts as a mechanical load that elicits an anabolic response that stimulates muscle growth and function. This anabolic response is usually greater in males than in females because of the higher testosterone concentration. However, it should also be noted that BMI does not differentiate between fat mass and lean mass as adipocyte indicators that may vary with sex. It should be noted that BMI is a measure of body fat that does not differentiate between weight changes because of an increase or decrease in muscle mass or body fat (Zaccagni *et al.*, 2020). According to Jafar *et al.* (Jaafar *et al.*, 2023), heavier individuals have a greater muscle mass, resulting in greater handgrip strength, than lighter individuals. In the study of Heidy *et al.* (Heidy *et al.*, 2019), the investigators found a positive correlation between height and forearm length. The study by Sirajudeen *et al.* (Sirajudeen *et al.*, 2012) found a positive correlation between hand length and handgrip strength. This shows that height affects the longer extremity or arm, thus approaching optimal muscle length that may result in greater grip strength. According to a prospective cohort study, handgrip strength in middle age appears to be associated with birth weight and the increase in height at puberty. Pubertal growth and handgrip strength possibly represent the development of muscle size and growth (Schrager *et al.*, 2007).

Our study found no relationship between blood glucose concentration and handgrip strength. Our study agrees with the study of Giglio *et al.* (Giglio *et al.*, 2018), which showed no significant relationship between muscle mass strength and blood glucose concentration. According to Giglio, this is because most respondents are individuals who routinely perform structured physical activity, even though they are in the light category. This is similar to our study respondents, who were public works personnel and who had performed physical activity such as street sweeping and cleaning storm drains. Glucose is the main fuel

for muscle contraction, entering the muscle cells by diffusion through the GLUT4 glucose transporter, which, after muscle contraction, moves from its intracellular storage site to the plasma membrane and T-tubules (Richter *et al.*, 2013). Glucose metabolism has an influence on handgrip strength through various mechanisms, such as disorders of glucose metabolism, particularly glycogenolysis, that may result in the loss of muscle strength. Insulin resistance in hyperglycemia may cause muscle degradation because hyperglycemia causes low muscle strength through effects on skeletal muscle mitochondria. The lowering of muscle strength by hyperglycemia occurs through Krüppel-like factor 15, which regulates skeletal muscle lipid flux. Hyperglycemia also increases pro-inflammatory cytokines, which may increase the catabolism that plays a role in the further decrease in muscle mass and quality (Gamayanti *et al.*, 2023). The amount of carbohydrates also influences the glycemic response and post-prandial insulin level, impacting the supply of glycogen to the muscles (Safitri *et al.*, 2020).

This study has a significant negative correlation between age and handgrip strength, which decreases with age (Manoharan *et al.*, 2015). Handgrip strength initially experiences an increase with increasing age, reaches its peak between the ages of 30 and 40 years, and decreases afterward (Zaccagni *et al.*, 2020). The age-related changes in skeletal muscle composition, particularly the accumulation of lipids in skeletal muscle fibers, play a role in the poor quality of the muscles and may result in metabolic abnormalities such as insulin resistance. Apart from changes in muscle quality, there is also a reduction in muscle mass that consistently occurs during aging. The increase in oxidative stress and chronic inflammation that commonly occurs in the aging process apparently also plays a role in the development of decreased muscle mass and strength (Mainous *et al.*, 2015). The decrease in muscle mass in the elderly corresponds to a 10% reduction in grip strength per decade, starting from the age of 40 years (Jaafar *et al.*, 2023). Skeletal muscle plays an important role in glucose metabolism and has a significant influence on insulin sensitivity. Increased

insulin resistance in older adults may be closely associated with skeletal muscle aging. A reduction in GLUT4 expression as a result of a decrease in muscle volume causes a reduction in insulin sensitivity in aging skeletal muscle (Joo *et al.*, 2022). The results of this study showed a correlation of waist circumference and hip circumference with handgrip strength. This is in line with the study of Widjaja *et al.* (Widjaja *et al.*, 2018), who stated that there was a significant correlation of waist circumference and hip circumference with handgrip strength ($p < 0.001$, $r = 0.302$; and $p < 0.001$, $r = 0.296$). A greater waist circumference is associated with a greater handgrip strength in males (Hardy *et al.*, 2013).

In the multiple regression analysis of HGS, we removed the data on body weight because of the occurrence of multicollinearity between weight and BMI, with a correlation coefficient of more than 0.9, indicating an extremely strong relationship between these two variables. Conceptually, only one of the variables is used in multivariate analysis. The multiple regression analysis included age, height, BMI, waist circumference, and hip circumference, with HGS as the dependent variable in a stepwise manner. After three iterations, only height, BMI, and hip circumference played a role in predicting HGS by means of the regression formula, which resulted in $R^2 = 0.19$. This signifies that for known values of height, BMI, and hip circumference, we can predict the HGS value 19% more accurately than without knowledge of these data. For a clinical diagnosis, a minimum increase of 10% is needed. This study found that height, BMI, and hip circumference play a role in predicting handgrip strength through the regression formula, i.e., $\text{handgrip strength} = -36.98 + 0.48 \times \text{height} + 0.79 \times \text{BMI} - 0.29 \times \text{hip circumference}$. Handgrip strength and waist circumference are surrogate muscle strength and visceral adiposity markers, respectively (Nakanishi *et al.*, 2024). Waist circumference is also more sensitive in determining body fat distribution, particularly in the abdomen. Waist circumference correlates better with abdominal fat distribution than BMI (Rokhmah *et al.*, 2015). Waist circumference and hip circumference are associated with obesity, in which the fat

content is higher, and the skeletal muscle and fascicular content is lower, thereby resulting in a lower contractile capacity of skeletal muscle (Afaq *et al.*, 2019). The limitation of this study is that it did not evaluate other factors that may affect handgrip strength, such as nutrient intake. From the anthropometric measures of the hands and feet in this study, the causes and effects of the decrease in handgrip strength could not be determined because this study used a cross-sectional design.

Conclusion

Three factors may be used to estimate handgrip strength in PWP, namely height, BMI, and hip circumference, whereas blood glucose concentration is not correlated with handgrip strength in PWP. It is recommended to pay attention to height, BMI, and hip circumference, particularly in the acceptance criteria of PWP candidates, and to conduct further studies with consideration of other factors that affect handgrip strength, such as food intake and anthropometric features of the hands and arms.

Acknowledgments

Financial support was provided by Universitas Trisakti for this research.

References

- Afaq, E., Hassan, S.H., Khan, M.R., Zaidi, D.A., & Baktashi, H., 2019. Correlation of Hand Dynamometry With Body Mass Index and Waist-Hip Ratio in Medical Students. *Pak J Physiol.*, 15, pp.19-21.
- Agtuahene, M.A., Quartey, J., & Kwakye, S., 2023. Influence of Hand Dominance, Gender, and Body Mass Index on Hand Grip Strength. *S Afr J Physiother.*, 79, pp.1-6.
- Al-Asadi, J.N., 2018. Handgrip Strength in Medical Students: Correlation with Body Mass Index and Hand Dimensions. *Asian Journal of Medical Sciences*, 9, pp.21-26.
- Badan Pusat Statistik Kota Administrasi Jakarta Barat., 2020. *Kecamatan Cengkareng Dalam Angka*, pp.1-114.
- Chen, D., Zhu, Y., Ni, W., Li, Y., Yin, G., Shao, Z., & Zhu, J., 2023. Hand Grip Strength is Inversely Associated with Total Daily Insulin Dose Requirement in Patients with Type 2 Diabetes Mellitus: A Cross-Sectional Study. *PeerJ.*, pp.1-13.
- Cooper, R., Tomlinson, D., Hamer, M., & Pereira, S.M.P., 2022. Lifetime Body Mass Index and Grip Strength at Age 46 Years: The 1970 British Cohort Study. *J Cachexia Sarcopenia Muscle*, 13, pp.1-10.
- Fauzi, L., Handayani, O.W.K., Susilo, M.T., Kurnia, A.R., Rahayu, S.R., Irawan, F.A., Horng Lu, F.J., Lin, C., Lai, M.F., & Yu, Y.C., 2022. Obesity in Indonesian and Taiwanese Adolescents Related to Self Perception, Diet, Exercise, and Body Image. *KEMAS*, 17(3), pp.453-461.
- Gamayanti, K.A.V., Aryana, I.G.P.S., & Gotera, W., 2023. Hubungan Antara Kekuatan Genggaman Tangan Dengan Kadar Gula Darah Sewaktu Pada Lansia Di Desa Melinggih, Kecamatan Payangan, Kabupaten Gianyar. *Intisari Sains Medis*, 14(2), pp.483-488.
- Giglio, B.M., Mota, J.F., Wall, B.T., & Pimentel, G.D., 2018. Low Handgrip Strength is Not Associated With Type 2 Diabetes Mellitus and Hyperglycemia: A Population-Based Study. *Clin Nutr Res.*, 7(2), pp.112-116.
- Hammed, A.I., & Obaseki, C.O., 2017. Interdependence of Body Mass Index with Handgrip Strength and Endurance Among Apparently Healthy Teenagers. *Turk J Kin.*, 4, pp.1-7.
- Handayani, M.D.N., Sadewa, A.H., Farmawati, A., & Rochmah, W., 2018. Anthropometric Prediction Equations for Estimating Muscle Mass of Elderly Women. *KEMAS*, 14(2), pp.195-204.
- Hardy, R., Cooper, R., Sayer, A.A., Shlomo, Y.B., Cooper, C., Deary, I.J., Demakakos, P., Gallacher, J., Martin, R.M., McNeill, G., Starr, J.M., Steptoe, A., Syddall, H., & Kuh, D., 2013. Body Mass Index, Muscle Strength, and Physical Performance in Older Adults From Eight Cohort Studies: The Halcyon Programme. *PLoS One*, 8(2), pp.e56483.
- Heidy., Djuartina, T., & Irawan, R., 2019. Korelasi Kekuatan Genggaman Tangan Dengan Karakter Antropometri Lengan Bawah dan Tangan Serta Indeks Massa Tubuh. *Damianus Journal of Medicine*, 18, pp.1-7.
- Ilmi, A.F., & Utari, D.M., 2020. Hubungan Lingkar Pinggang dan Rasio Lingkar Pinggang-Panggul (RLPP) Terhadap Kadar Glukosa Darah Puasa Pada Mahasiswa. *Journal of Nutrition College*, 9, pp.222-227.
- Jaafar, M.H., Ismail, R., Ismail, N.H., Isa, Z.M., Tamil, A.M., Nasir, N.M., Keat, N.K., Razak, N.H., Abidin, N.Z., & Yusof, K.H., 2023. Normative Reference Values and Predicting Factors of Handgrip Strength For Dominant

- and Non-Dominant Hands Among Healthy Malay Adults in Malaysia. *BMC Musculoskeletal Disord.*, 24, pp.1-9.
- Joo, K.C., Son, D.H., & Park, J.M., 2022. Association Between Relative Handgrip Strength and Insulin Resistance in Korean Elderly Men Without Diabetes: Findings Of The 2015 Korea National Health Nutrition Examination Survey. *Korean J Fam Med.*, 43, pp.199-205.
- Krakauer, N.Y., & Krakauer, J.C., 2020. Association of Body Shape Index (ABSI) with Hand Grip Strength. *Int J Environ Res Public Health*, 17, pp.1-12.
- Kurniawan, A., Widjaja, D., & Lugito, N.P.H., 2018. *Mean and Cutoff Value of Hand Grip Strength For Healthy Indonesian Young Adults*. World Congress on Osteoporosis, Osteoarthritis and Musculoskeletal Diseases; Krakow, Poland.
- Leong, D.P., Teo, K.K., Rangarajan, S., Kutty, R., Lanas, F., Hui, C., Quanyong, X., Zhenzhen, Q., Jinhua, T., Noorhassim, I., AlHabib, K.F., Moss, S.J., Rosengren, A., Akalin, A.A., Rahman, O., Chifamba, J., Orlandini, A., Kumar, R., Yeates, K., Gupta, R., Yusufali, A., Dans, A., Avezum, A., Lopez-Jaramillo, P., Poirier, P., Heidari, H., Zatonska, K., Iqbal, R., Khatib, R., & Yusuf, S., 2016. Reference Ranges of Handgrip Strength From 125,462 Healthy Adults in 21 Countries: A Prospective Urban Rural Epidemiologic (PURE) Study. *J Cachexia Sarcopenia Muscle*, 7, pp.535-546.
- Mainous, A.G., Tanner, R.J., Anton, S.D., & Jo, A., 2015. Grip Strength as A Marker of Hypertension and Diabetes in Healthy Weight Adults. *Am J Prev Med*, 49(6), pp.850-858.
- Manoharan, V.S., Sundaram, S.G., & Jason, J.I., 2015. Factors Affecting Hand Grip Strength and Its Evaluation: A Systemic Review. *Int J Physiother Res.*, 3(6), pp.1288-1293.
- McGrath, R., Johnson, N., Klawitter, L., Mahoney, S., Trautman, K., Carlson, C., Rockstad, E., & Hackney, K.J., 2020. What Are The Association Patterns Between Handgrip Strength and Adverse Health Conditions? A Topical Review. *SAGE Open Medicine*, 8, pp.1-12.
- Nakanishi, S., Shimoda, M., Kimura, T., Sanada, J., Fushimi, Y., Iwamoto, Y., Iwamoto, H., Dan, K., Mune, T., Kaku, K., & Kaneto, H., 2024. The Impact of Handgrip Strength and Waist Circumference on Glycemic Control: Prospective, Observational Study Using Outpatient Clinical Data in Japanese Patients With Type 2 Diabetes Mellitus. *J Diabetes Investig.*, 15(7), pp.892-898.
- Nathania, N.P.S., Nugraha, M.H.S., Primayanti, I.D.A., & Saraswati, P.A.S., 2023. Handgrip Strength is Associated With Balance in Elderly Women. *Majalah Ilmiah Fisioterapi Indonesia*, 11, pp.70-75.
- Gubernur Provinsi Daerah Khusus Ibukota Jakarta., 2017. *Peraturan Gubernur Provinsi Daerah Khusus Ibukota Jakarta Nomor 7 tahun 2017 Tentang Penanganan Prasarana dan Sarana Umum Tingkat Kelurahan*. Jakarta 2017.
- Richter, E.A., & Hargreaves, M., 2013. Exercise, GLUT4, and Skeletal Muscle Glucose Uptake. *Physiol Rev.*, 93(3), pp.993-1017.
- Rokhmah, F.D., Handayani, D., & Al-Rasyid, H., 2015. Korelasi Lingkar Pinggang dan Rasio Lingkar Pinggang-Panggul Terhadap Kadar Glukosa Plasma Menggunakan Tes Toleransi Glukosa Oral. *Jurnal Gizi Klinik Indonesia*, 12, pp.28-35.
- Rondhianto., Ridla, A.Z., & Hasan, H., 2024. Sociodemographic Factors Affecting Diabetic Dietary Behavior in People with Type 2 Diabetes Mellitus. *KEMAS*, 19(3), pp.429-437.
- Safitri, I., Setyarsih, L., Susanto, H., Suhartono., & Fitranti, D.Y., 2020. The Effect of Low and High Glycemic Load Diet on Muscle Fatigue of Young Soccer Athletes. *KEMAS*, 16(1), pp.111-120.
- Schrager, M.A., Metter, E.J., Simonsick, E., Ble, A., Bandinelli, S., Lauretani, F., & Ferrucci, L., 2007. Sarcopenic Obesity and Inflammation in The InChianti Study. *J Appl Physiol.*, 102(3), pp.919-925.
- Sirajudeen, M.S., Shah, U.N., Pillai, P.S., Mohasin, N., & Shantaram, M., 2012. Correlation Between Grip Strength and Physical Factors in Men. *International Journal of Health and Rehabilitation Sciences*, 1, pp.58-63.
- Soenyoto, T., Mollap, K., & Mungpong, J., 2017. Correlation of Energy and Protein Consumption Levels with Physical Endurance of Rhythmic Gymnast Athletes. *KEMAS*, 13(1), pp.131-136.
- Soraya, N., & Parwanto, E., 2023. The Controversial Relationship Between Body Mass Index and Handgrip Strength in The Elderly: An Overview. *Malays J Med Sci.*, 30, pp.73-83.
- Vaishya, R., Misra, A., Vaish, A., Ursino, N., & D'Ambrosi, R., 2024. Hand Grip Strength as A Proposed New Vital Sign Of Health: A Narrative Review of Evidences. *Journal of Health, Population and Nutrition*, 43, pp.1-

14.

Widjaja, D., Tjhai, K., Dermawan, K., Kalim, J., Tanzil, S., Pramusinta, A., Phenca, H.K., Kurniawan, A., 2018. *Correlations Between Hand Grip Strength and Anthropometric Status in Young Indonesian Adults*. World Congress on Osteoporosis, Osteoarthritis and Musculoskeletal Diseases; Krakow, Poland.

Zaccagni, L., Toselli, S., Bramanti, B., Gualdi-Russo, E., Mongillo, J., & Rinaldo, N., 2020. Handgrip Strength in Young Adults: Association With Anthropometric Variables and Laterality. *Int J Environ Res Public Health*, 17, pp.1-18.