

Advances in Social Science, Education and Humanities Research
Editor-in-Chief: Wadim Strielkowski

Zulfikar Bagus Pambuko · Muji Setiyo ·
Chrisna Bagus Edhita Praja ·
Agus Setiawan · Fitriana Yuliasuti ·
Lintang Muliawanti · Veni Soraya Dewi *Editors*

Proceedings of the 4th Borobudur International Symposium on Humanities and Social Science 2022 (BIS-HSS 2022) · Volume 778

OPEN ACCESS

Editors

Zulfikar Bagus Pambuko
Islamic Economics Law
Universitas Muhammadiyah Magelang
Magelang, Indonesia

Muji Setiyo
Mechanical Engineering
Universitas Muhammadiyah Magelang
Magelang, Indonesia

Chrisna Bagus Edhita Praja
Law Science
Universitas Muhammadiyah Magelang
Magelang, Indonesia

Agus Setiawan
Informatics Engineering
Universitas Muhammadiyah Magelang
Magelang, Indonesia

Fitriana Yuliasuti
Pharmacy Science
Universitas Muhammadiyah Magelang
Magelang, Indonesia

Lintang Muliawanti
Communications Science
Universitas Muhammadiyah Magelang
Magelang, Indonesia

Veni Soraya Dewi
Accounting
Universitas Muhammadiyah Magelang
Magelang, Indonesia



ISSN 2731-8060

ISSN 2352-5398 (electronic)

Advances in Social Science, Education and Humanities Research

ISBN 978-2-38476-117-3

ISBN 978-2-38476-118-0 (eBook)

<https://doi.org/10.2991/978-2-38476-118-0>

© The Editor(s) (if applicable) and The Author(s) 2024, corrected publication 2024. This book is an open access publication.

Open Access This book is licensed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits any noncommercial use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this book are included in the book's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the book's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.

This work is subject to copyright. All commercial rights are reserved by the author(s), whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed. Regarding these commercial rights a non-exclusive license has been granted to the publisher.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors, and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, expressed or implied, with respect to the material contained herein or for any errors or omissions that may have been made. The publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This Atlantis Press imprint is published by the registered company Atlantis Press S.A.R.L.,
part of Springer Nature

The registered company address is: 22 Rue de Palestro 75002 Paris France

Paper in this product is recyclable.

Contents

Peer-Review Statements	1
<i>Zulfikar Bagus Pambuko, Muji Setiyo, Chrisna Bagus Edhita Praja, Agus Setiawan, Fitriana Yuliasuti, Lintang Muliawanti, and Veni Soraya Dewi</i>	
The Effectiveness of Using Cup Board Hoop Media on Children's Self-Confidence	4
<i>Rina Syafrida, Ega Trisna Rahayu, Ervina, and Nursamsi</i>	
Keisya Levronka's "Tak Ingin Usai" Phenomenon in the Context of the Music Industry	11
<i>Ni Wayan Ardini</i>	
Community Empowerment in Magelang City through the Interaction of Innovation	18
<i>Andjar Prasetyo, Dewi Gartika, Heri Wahyudianto, Agustinus Hartopo, Beki Putri Harwijayanti, and Juliana Carolina Kilmanun</i>	
Community-Based Education Quality Management Model	30
<i>Undang Ruslan Wahyudin, Hinggil Permana, and Ella Nurlailasari</i>	
Cleanse and Protect: Harnessing the Antibacterial Power of Guava Leaves in Liquid Soap Antiseptic Formulation	43
<i>Nuniek Nizmah Fajriyah, Khusna Santika Rahmasari, Urmatul Waznah, and Herni Rejeki</i>	
TikTok as a Promotional Media for Post COVID-19 Art Exhibitions	56
<i>Yessy Bella Anggraeni and Mohammad Isa Pramana Koesoemadinata</i>	
Clustering Patterns of Food Crops to Increase Community Income in Papua Province	63
<i>Andjar Prasetyo, Agustinus Hartopo, Jack Syauta, Heri Wahyudianto, Albert Soplanit, and Beki Putri Harwijayanti</i>	
The Validity of the Pro-Based Caturalis Learning Model to Improve Elementary School Students' Scientific Literacy	74
<i>Ari Suryawan, Suyanta, and Insih Wilujeng</i>	

Policy Adjustment Towards Green Economy in Customary Territory after the Expansion of Papua Province	81
<i>Andjar Prasetyo, Heri Wahyudianto, Payung Layuk, Siswanto Siswanto, Maryam Nurdin, Dewi Gartika, Agustinus Hartopo, and Niki Elistus Lewaherilla</i>	
Using Video Physical Assessment to Enhance Nursing Student's Skills	94
<i>Fajar Agung Nugroho, Dadi Santoso, Podo Yuwono, and Nur Indarwati Septiriana</i>	
Analysis of Three Levels of Product Potential of Rural Tourism Areas as A Buffer for the Borobudur National Strategic Area	103
<i>Andjar Prasetyo, Tri Weda Raharjo, Herrukmi Septa Rinawati, Abdul Hamid, Heri Wahyudianto, and Agustinus Hartopo</i>	
A Model of Harmony Village in Magelang District: The Perspective of Cross-Religious Leaders	117
<i>Eko Kurniasih Pratiwi, Nasitotul Janah, and Yusuf Ardianto</i>	
Ghost Kitchen Concept: An Alternative Culinary Business Method with a Minimum Space	126
<i>Toddy Hendrawan Yupardhi</i>	
Feasibility Study of Salt Tourism as Role Model and Branding of Madura Halal Destination	132
<i>Dinara Maya Julijanti and Fachrur Rozi</i>	
Formulation and Antibacterial Tests of Serum Preparation of Ethanol Extract of Guava Leaves (<i>Psidium Guajava</i> L) as an Anti-Acne	143
<i>Neni Sri Gunarti, Shintia, Farhamzah, Eko Sri Wahyuningsih, and Putri Agustina</i>	
Effect of Mahabbah Package to Increase Breast Milk Production in Postpartum Mothers after The Covid-19 Pandemic: A Case Study in Indonesia	151
<i>Maryatun, Indarwati, Endang Sri Wahyuni, Fida' Husain, Dewi Kartika Sari, and Tri Susilowati</i>	
The Influence of Product Packaging/Packaging Design, Health Consciousness and Advertising on Purchase Decision Through Attitude and Intention on You C-1000 Consumers in Samarinda City	160
<i>Muhammad Nawawi, Amiril Azizah, Annisa Rasyidin, and Hanifah Ekawati</i>	

Actualization of Social Values in The Implementation of The Mombowa Tumpe Ceremony in The Batui Indigenous Communities, Banggai District	166
<i>Falimu and Karmila P. Lamadang</i>	
Database Management Infrastructure, Human Resources, Financing, and Equipment in The Sustainable Management of The Covid-19 Pandemic	174
<i>Andjar Prasetyo, Abdul Hamid, Bekti Putri Harwijayanti, Herrukmi Septa Rinawati, Tri Weda Raharjo, Heri Wahyudianto, and Agustinus Hartopo</i>	
Mindful Tourism: Management of Tourist Destinations as an Effort to Counter Stereotypes of the Madurese People	186
<i>Nikmah Suryandari and Farida Nurul Rahmawati</i>	
Comparative and Collaborative Data Management as a Driver for Regional Competitiveness Efficiency in the Economic Development Region	196
<i>Andjar Prasetyo, Trisnani Trisnani, Heri Wahyudianto, Agustinus Hartopo, Joula Olvy Maya Sondakh, Juliana Carolina Kilmanun, and Ndoheba Kenda</i>	
Development of Educational Quality Management Model Based on Industrial Technology 4.0 at Vocational High School	209
<i>Undang Ruslan Wahyudin, Hinggil Permana, Ella Nurlailasari, and Endin Nasrudin</i>	
Visual Communication Elements and Meaning of Hatten Wines Bali Label	222
<i>Made Arini Hanindharputri, I Wayan Mudra, Ni Putu Emilika Budi Lestari, and A. A. Sagung Intan Pradnyanita</i>	
Branding of Islamic Boarding Schools as New Edutourism in Madura	229
<i>Fachrur Rozi and Nadya Poernamasari</i>	
The <i>Peteng Bulan</i> Dance Choreography Values	237
<i>Ni Made Ruastiti, Ni Wayan Karmini, and I Made Sidia</i>	
EYL Students' Engagement in Flipped English Classroom through Interactive Digital Text	244
<i>Nina Puspitaloka, Evi Karlina Ambarwati, Indah Purnama Dewi, and Putri Kamalia Hakim</i>	

Study Narrative of The Role Communication to Kiai Madura as A Leader and Therapist of People with Mental Disorders	252
<i>Sri Wahyuningsih and Mohammad Arief Wahyudi</i>	
Evaluation of UI/UX Usability in Augmented Reality Application of Balinese Shadow Puppet <i>Panca Pandawa</i>	261
<i>I Komang Try Adi Stanaya, I Wayan Mudra, and I Gst Ngr Dwijaksana</i>	
Health Assessment of Women in Reproductive Age as Expectant Mothers Through Body Mass Index Examination	268
<i>Emi Nurlaela and Dian Kartikasari</i>	
Induction <i>Kelentangan</i> as a Pre-Therapy Psychobiophysics Media	274
<i>Ketut Sumerjana and Ary Nugraha Wijayanto</i>	
Digital Financial Transformation in The Financial Inclusion Program and Its Impact on Income Inequality: The Case of Middle-Income Countries	281
<i>Siti Aisyah and Tika Pratika</i>	
Consistency of Blitar City Policy towards the Area of Acceleration of Economic Development in the Perspective of Regional Competitive	288
<i>Andjar Prasetyo, Herrukmi Septa Rinawati, Abdul Hamid, Tri Weda Raharjo, Trisnani Trisnani, Heri Wahyudianto, and Agustinus Hartopo</i>	
The Influence of The Human Development Index and Unemployment on Poverty: Zakat as Moderation	299
<i>Ellyana Amran, Aekram Faisal, and Dian Octaviani</i>	
Semiotics Perspective on Representation of Patriarchal Culture in Films	307
<i>Ni Kadek Dwiyani, I Made Saryana, and Ni Luh Gede Liswahyuningsih</i>	
MSMEs Sustainability: Application of GSM-DE as a Business Growth Model for MSME	314
<i>Nia Kurniati Bachtiar, Faqiatul Mariya Waharini, and Nur Laila Yuliani</i>	
Figures of Speech in The Dance Theatre The Cry of Sita	323
<i>Ni Ketut Dewi Yulianti</i>	

Digitalization of Madura Tourism Branding in the New Normal	431
<i>Bani Eka Dartiningsih, Mas'amah, and Surokim</i>	
From the Genetic Mutation to the Specific Pathologies	439
<i>Salsabila Salma Zahrah, Ratna Wijayatri, Imron Wahyu Hidayat, Alfian Syarifuddin, and Setiyo Budi Santoso</i>	
The Values of the <i>Lebaran</i> Tradition in Surakarta: An Ethnographic Study	448
<i>Sigit Haryanto and Andi Haris Prabawa</i>	
Improving SMEs Performance through Omni-channel Strategy	455
<i>Kosasih, Abdul Yusuf and Maulana Rifai</i>	
Creating Behavioral Intention Through Online Experience on Virtual Tourism	463
<i>Abdul Yusuf and Rina Maria Hendriyani</i>	
Cultural Commodification Analysis Based on Local Wisdom in an Effort to Strengthen Madura's Potential as a Wellness Tourism Destination	468
<i>Farida Nurul Rahmawati, Nikmah Suryandari, and Windiani</i>	
Post Covid-19 Student's Burnout Level in Boarding School's Non-Academic Programs	475
<i>Ulfah Nafi'ah, Fajar Ciptandi, and Runik Machfiroh</i>	
The Reasons Behind the Forms of Cyberbullying Teenagers Choose of Senior High School in Surakarta	481
<i>Siti Nurina Hakim and Dara Febrian Chita Prastiwi</i>	
Post-Covid-19 Tourism Recovery and Innovation Methods: Practical Evidence from Indonesia	489
<i>Sabrina Mawar Rahmah and Lintang Muliawanti</i>	
Bibliometric Analysis of Global Research on Technology Information in Antimicrobial Stewardship using Scopus Database	496
<i>Heni Lutfiyati, Nanang Munif Yasin, Jarir At Thobari, Zullies Ikawati, and Prasojo Pribadi</i>	
Husband and Family Support for Breastfeeding Mothers with COVID-19: The Mother's Perspective	507
<i>Nur Intan Kusuma, Rini Kristiyanti, and Milatun Khanifah</i>	
Changes in Priority Scale Between Needs vs Wants in Customer Perspective During and After the Covid-19 Pandemic Using Fuzzy AHP	517
<i>Muhamad Ali Pahmi, Ahmad Faisal Ayob, and Gendut Suprayitno</i>	

Relationship of Arm Muscle Strength and Hand-Eye Coordination with Pointing Throwing Ability in Petanque Game of Banggai Regency Athletes	606
<i>Muhammad Salahuddin, Hariadi Said, Abu Bakar, Ardiansyah Nur, and Frischa Priscillia Mongguwi</i>	
Socioeconomic Status and Protein Intake Adequacy in Elementary Children in Surakarta	613
<i>Setyaningrum Rahmawaty, Afrianus Karo, Dian Nazikha Khusna, Ufairah Maliha Shofwah, Ari Murti Nindiyanti, Niki Dwi Astuti, and Jannike Øyen</i>	
Prevalence Analysis of Intestinal Worm Infection in Pregnant Women	621
<i>Sulastri, Dian Hudiyawati, Siti Arifah, Winarsih Nur Ambarwati, Widya Putri Rahmawati, and Anindyati Fatikhul Jannah</i>	
Is Social Support a Mediator in the Relationship between Gratitude and Resilience of Covid-19 Survivors?	630
<i>Rini Lestari, Juliani Prasetyaningrum, Daliman, and Aqilah Larasati Imadanty</i>	
Countering Transnational Shadow Economy Crime: Mutual Legal Assistance Mechanism	637
<i>Wita Setyaningrum, Ridhayana Andi Baso, Muhammad Nur, and Mufti Khakim</i>	
Trends in Mental Health Research during Covid-19	648
<i>Sugeng Mashudi, Filia Icha Sukamto, Lukman Handoko, Eky Okviana Armyati, Nurul Sri Wahyuni, and Naylil Mawadda Rohma</i>	
The Effect of the Sharia Credit Payment System on Increasing Income (Study Gold Store “Mas-Masan”)	655
<i>Nugraheni Fitroh Rezqi Syakarna, Fatkhur Rohman Albanjari, Agus Mujiyono, Nurul Abidin, and Sigit Dwi Laksana</i>	
The Effectiveness of the Home Rehabilitation Program (HRP) after the Application of Transcranial Magnetic Stimulation (TMS) Post-Stroke	661
<i>Umi Budi Rahayu, Sakinah Sakinah, and Santri Ramida</i>	
Development of Movement Quality Instruments for Physical Fitness Activities Based on Developmental Psychology of Children Aged 13-18 Years	672
<i>Resty Gustiawati, Terbit Surya, Alwi Sihab, and Muhammad Surur</i>	

Summary of Factors Affecting Indonesian Islamic Banking Efficiency in 2014-2022	527
<i>Abdullah Hasan Alifuddin and Nur Fatwa</i>	
Strengthening the Role of Communities in the Prevention of Sexual Violence in Higher Education: Opportunities and Challenges	532
<i>Khaerul Umam Noer, Titiek Kartika Hendrastiti, Lidwina Inge Nurtjahyo, and Varinia Pura Damaiyanti</i>	
Improving Youth Moral Knowledge Through Classical Guidance: Empirical Study	541
<i>Purwati Purwati, Muhammad Japar, Dewi Lianasari, and Filda Khoirun Nikmah</i>	
Teachers' Preparations in Teaching English through Online Learning	548
<i>Mukmin, Nurhaida Lakuana, Marhana Rullu, Armin Haluti, and Iswath Al Mawaddath</i>	
The Cultural Meaning of <i>Para-Para</i> for the Dondai Villager, Papua	559
<i>S. I Wayan Rai, Ni Made Ruastiti, Ni Nyoman Lia Susanti, Ni Wayan Masyuni Sujayanti, Gede Yoga Kharisma Pradana, and Yunus Wafom</i>	
The Contribution of Parental Attachment to Adolescent Moral Intelligence	566
<i>Puspa Rahayu Utami Rahman, Wina Lova Riza, and Regi Ramadan</i>	
Teacher Ability in Designing Entrepreneurship-Based Mathematics Learning	574
<i>Alpha Galih Adirakasiwi, Attin Warmi, and Achmad Nawawi</i>	
The Effect of Toxic Workplace Climate Dimensions on Millennial Employees in Jakarta	584
<i>Netania Emilisa, Karina Ganadi, Lidia Wahyuni, Beta Oki Baliartati, and Shafrani Dizar</i>	
Precision Journalism: Digitalization and Research Approaches in Journalism Practices in Indonesian Online Media	594
<i>Ansar Suherman, Hasrullah, Hafied Cangara, and Sudirman Karnay</i>	
The Implementation of Mural Thematic and Individual Whitespace as an Interaction Concept of Innovative Learning Space for Elementary School Students in Bali	600
<i>I Nyoman Larry Julianto, I Wayan Swandi, and I Wayan Agus Eka Cahyadi</i>	

Relationship of Arm Muscle Strength and Hand-Eye Coordination with Pointing Throwing Ability in Petanque Game of Banggai Regency Athletes	606
<i>Muhammad Salahuddin, Hariadi Said, Abu Bakar, Ardiansyah Nur, and Frischa Priscillia Mongguwi</i>	
Socioeconomic Status and Protein Intake Adequacy in Elementary Children in Surakarta	613
<i>Setyaningrum Rahmawaty, Afrianus Karo, Dian Nazikha Khusna, Ufairoh Maliha Shofwah, Ari Murti Nindiyanti, Niki Dwi Astuti, and Jannike Øyen</i>	
Prevalence Analysis of Intestinal Worm Infection in Pregnant Women	621
<i>Sulastri, Dian Hudiawati, Siti Arifah, Winarsih Nur Ambarwati, Widya Putri Rahmawati, and Anindyati Fatikhul Jannah</i>	
Is Social Support a Mediator in the Relationship between Gratitude and Resilience of Covid-19 Survivors?	630
<i>Rini Lestari, Juliani Prasetyaningrum, Daliman, and Aqilah Larasati Imadanty</i>	
Countering Transnational Shadow Economy Crime: Mutual Legal Assistance Mechanism	637
<i>Wita Setyaningrum, Ridhayana Andi Baso, Muhammad Nur, and Mufti Khakim</i>	
Trends in Mental Health Research during Covid-19	648
<i>Sugeng Mashudi, Filia Icha Sukamto, Lukman Handoko, Eky Okviana Armyati, Nurul Sri Wahyuni, and Naylil Mawadda Rohma</i>	
The Effect of the Sharia Credit Payment System on Increasing Income (Study Gold Store “Mas-Masan”)	655
<i>Nugraheni Fitroh Rezqi Syakarna, Fatkhur Rohman Albanjari, Agus Mujiyono, Nurul Abidin, and Sigit Dwi Laksana</i>	
The Effectiveness of the Home Rehabilitation Program (HRP) after the Application of Transcranial Magnetic Stimulation (TMS) Post-Stroke	661
<i>Umi Budi Rahayu, Sakinah Sakinah, and Santri Ramida</i>	
Development of Movement Quality Instruments for Physical Fitness Activities Based on Developmental Psychology of Children Aged 13-18 Years	672
<i>Resty Gustiawati, Terbit Surya, Alwi Sihab, and Muhammad Surur</i>	

The Effect of Acupressure on Length of Labor: A Literature Review	679
<i>Aziza Vira Arunita, Heni Setyowati Esti Rahayu, and Kartika Wijayanti</i>	
Systematic Literature Review of Satisfaction Model and Interest In Adoption of E-Government Services	692
<i>Erick Sorongan, Danar Retno Sari, and Hendra Sanjaya Kusno</i>	
Adapting Content-Based Instruction Principle in Teaching Culinary Arts Vocabulary at Vocational High School	703
<i>Hardianti and Yusniati Nadir Sabata</i>	
Waqf Core Principles Implementation at Muhammadiyah <i>Nazhir</i> Waqf: An Analytic Network Process Approach	710
<i>Adilnia Fifi Susanti, Berliana Florensia, Habib Abdul Kholid Al Huzain, Ilyas Syah, and Fahmi Medias</i>	
The Learning Paradigm of Democracy-Based Islamic Education in Fostering Students' Tolerance	719
<i>Siti Wardah Hanafie Das, Abdul Halik, and S. Muhammad Nasir</i>	
Linguistics Factors as Speaking Obstacle of the EFL Students in Classroom	729
<i>Yusniati N. Sabata, Riya Mulyanti, Abdul Rabbi Arrasul, Sitti Hardianti, and Desriani Nggolaon</i>	
Mapping About Bonding Relations in the Communication Network of Madura Salt Production Workers	737
<i>Imam Sofyan, Moch. Imron Rosyidi, and R. Bambang Moertijoso</i>	
An Economic Added Value Analysis of Ready-To-Drink Prebiotic Candidate Milk	744
<i>Silvia Oktavia Nur Yudiastuti, Saiful Anwar, Yossi Wibisono, Agung Wahyono, Anna Maria Handayani, Annisa'u Choirun, Resti Pranata Putri, and Findi Citra Kusumasari</i>	
Analysis of Needs for Fitness Activity Tools in Physical Education Learning in Schools	755
<i>Fahrudin, Tedi Purbangkara, Resty Gustiawati, and Muhammad Mury Syafei</i>	
The Effect of Self-Control on Impulsive Buying of In-Game Virtual Goods among Mobile Legends Bang Bang Players	764
<i>Maya Mardi Yanti, Aftina Nurul Husna, and Laili Qomariyah</i>	

Photography as a Branding process on MSME Products	772
<i>Imam Budi Sumarna, Ira Wirasari, and Dandi Yunidar</i>	
Evaluation of Thematic Learning Curriculum 2013 in Madrasah Ibtidaiyah	779
<i>Umri Nur Safitri, Ahwy Oktradiksa, and Norma Dewi Shalikhah</i>	
Mathematical Understanding: Learning Number Operation Using Media in The Context of Futsal	789
<i>Kiki Nia Sania Effendi, Dany Aulia, Indrie Noor Aini, and Rina Marlina</i>	
Community-Based Education Financing Management in Karawang	796
<i>Hinggil Permana, Undang Ruslan Wahyudin, Muhamad Erihadiana, and Yayat Herdiana</i>	
Green Human Resources Management, Green Environmental: Influence on Sustainability Performance Mediated by Digital Metaverse Technology in the Industry 4.0 Era	809
<i>Retno Purwani Setyaningrum and Anna Wulandari</i>	
The Development of Tourist Object and Attraction Digitization in Surakarta, Indonesia	819
<i>Argyo Demartoto</i>	
Health Seeking Behavior for the Non-Communicable Diseases and Mental Disorder	826
<i>Arif Widodo, Mustika Adelia, Reny Novia Karlina, and Wiwin Renny Rahmawati</i>	
The Role of Genetic Mutation on Schizophrenia: A Basic Review Prior to Pharmacogenomics	835
<i>Dhea Nur Hikmah, Alfian Syarifuddin, Setiyo Budi Santoso, Ratna Wijayatri, and Imron Wahyu Hidayat</i>	
Improving Grade XI Students' Vocabulary Through Extensive Reading in Senior High School 4 Kendari	848
<i>Suharjuddin, Wa Saripah, and Yazid Husain</i>	
Independent Village Agrotourism Based on Regional Innovation System Governance in Jayapura Regency, Papua Province	856
<i>Andjar Prasetyo, Bekti Putri Harwijayanti, Syahrudin Kadir, Heri Wahyudianto, Dewi Gartika, and Mursid Zuhri</i>	

Phenomenological Study on Sexual Harassment of Women at the Workplace in the Karawang Region	870
<i>Arif Rahman Hakim, Linda Mora, Cempaka Putrie Dimala, and Dinda Aisha</i>	
Utilization of Prime-Legal Drafting (Prime-LD) Application in The Preparation of Legislation for The Government in Indonesia	877
<i>Dyah Adriantini Sintha Dewi, Habib Muhsin Safingi, Suharso, Dili Trisna Noviasari, and dan Wafda Vivid Izziyana</i>	
Determinants of Profit Efficiency among Small and Medium-Sized Enterprises in Indonesia	884
<i>Muhamad Wahid Ibrahim and Ravindra Ardiana Darmadi</i>	
The Implementation of Multiple Intelligence-Based Soft Skill Learning Model for Students	898
<i>Hasrat Abu Bakar Aimang</i>	
Analysis of the Implementation of the Independent Curriculum in Mathematics Subjects in Elementary Schools	906
<i>Ayu Fitri, Tatang Herman, Udin Saefudin Sa'ud, Tarpan Suparman, Rahman, and Yulistina Nur DS</i>	
Millennial Investor Analysis of Cryptocurrency Investments	916
<i>Alfiana Yogi Saputri and Marlina Kurnia</i>	
The Initiation of Madura's Content Creator, Digital Community, and Citizen Media Development as the Strengthened Strategy of Marketing Communication in the Madura Region within the Digital Communication Era	922
<i>Surokim, Bani Eka Dartiningsih, and Luthfi Awwalia</i>	
Analysis of Financial Resilience in Magelang to Develop Effective Policies	931
<i>Naufal Afif and Nur Hidayah</i>	
STEM-Based Biology Instruction Using an Inquiry-Based Learning Approach to Foster Students' Creative Thinking	936
<i>R. Nur Zakiyah, Ibrohim, and Andi Jusman Tharihk</i>	
Visual Structure of the Digital Comic Banggaber with the Theme of Parents and Children	943
<i>Syarip Hidayat, I Wayan Mudra, and Ranti Rachmawanti</i>	

Teachers Public Speaking Ability in Teaching English at Islamic Boarding School in Karawang Regency	950
<i>Yousef Bani Ahmad, Mobit, Andriamella Elfarissyah, Hasna Afifah, Dinda Annisa Syahida, and Diajeng Nada Vici</i>	
Middlemen and The Debt bondage System in Salt Production at Madura	957
<i>Prasetyono, Emi Rahmawati, Fitri Agustin, Muhammad Syam Kusufi, Ach Fawaid As'ad, Sultan Syah, and Muhammad Akbars</i>	
Heritage Tourism Branding Strategy (Comparative Study of Heritage Tourism Branding Strategies in Madura, Indonesia and Malaysia)	969
<i>Ahmad Cholil, Tatag Handaka, and Hamid Busthami Noer</i>	
Local Government Spending on Information and Communication Technology	976
<i>Ratna Wulaningrum, Venti Eka Satya, and Dwi Cahyadi</i>	
Social Interaction in Local Government Financial Reporting	981
<i>Hendrawan Santosa Putra, Andriana, and Siti Maria Wardayati</i>	
Cultural Values in the Toponymy of Medalsari Village and a Conservation Model to Support Tourism Based on Local Wisdom in Karawang District	987
<i>Suntoko, Abduloh, and Tedi Purbangkara</i>	
Engineering for Making Commercial Compost Fertilizer of Oyster Mushroom Waste Baglog Using Molasses Activator in Efforts to Implement Green Economy	997
<i>Abel Malvin Putra Hia, Suparni Setyowati Rahayu, and Muchlis</i>	
Determinants of Whistleblowing Intentions in Regional Organization	1006
<i>Miladia Ismi Khoirunisa, Nur Laila Yuliani, Pranita Siska Utami, and Wahyu Anggit Prasetya</i>	
Innovation in Learning Financial Literacy Through the Development of Animation Film for Elementary School Students	1017
<i>Attin Warmi, Alpha Galih Adirakasiwi, and Ahmad Nawawi</i>	
Perceptions of Karang Taruna Management in Karawang Regency on Mastery of English Skills in the Globalization Era	1026
<i>Yousef Bani Ahmad, Hawignyo, and Liya Megawati</i>	
The Effect of Al-Qur'an <i>Murottal</i> Therapy on Anxiety in Third Trimester Pregnant Women: Study Literature Reviews	1033
<i>Nia Afifatul Ulya, Heni Setyowati Esti Rahayu, and Dwi Sulistyono</i>	

The Effectiveness of Rope Ladder Physical Activity Media on Physical Motor Activity in Early Childhood	1043
<i>Ega Trisna Rahayu, Rina Syafrida, Ine Nirmala, Ervina Istanti, and Nursyamsi</i>	
A Combination Supplement of Bitter Melon Extract (<i>momordica charantia</i> L.) with Snakehead Fish (<i>channa striata</i>) Powder has no Effect as an Anti-glycation Agent in type 2 Diabetes Mellitus	1055
<i>Fahrur Nur Rosyid, Muhtadi, Andi Suhendi, and Ahmad Fadhlur Rahman</i>	
An Investigation of Students' Barriers, Hopes, and Short-Term Plan toward Their Speaking Skill Improvements	1064
<i>Ali Wira Rahman and Rizka Indahyanti</i>	
Stigma Analysis of HIV/AIDS at SMK Negeri 1 Cilacap	1076
<i>Dwi Maryanti, Choirur Rizqiya, and Sohimah</i>	
Gender-Based Disaster Risk Reduction: A Case Study of Strengthening Women's Capacity in Pacitan District	1083
<i>Nur Izzaturrahmah, Sayidah Ummul Solihah, Firgi Pradipta Firdaus, and Satria Iman Prasetyo</i>	
Community Based Tourism on Natural Tourism Destination	1095
<i>Abdul Yusuf and Kosasih</i>	
Effectiveness of Group Counseling with Digital Mind Mapping (DMM) Techniques to Improve Self-Regulation	1102
<i>Hijrah Eko Putro, Astiwi Kurniati, and Lutfi Fahrurozi</i>	
Utilization of Herbal Supplements to Improve Health Quality after Suffering from Covid-19 Infection in Elderly Posyandu Partners	1111
<i>Safira Bela Annisa, Febrianti Nur Antika, Amalia Siti Choerun Nisa, Reza Rohmatun Zakiyah, Fahrur Nur Rosyid, Guntur Nurcahyanto, and Muhtadi</i>	
Entrepreneurial Marketing Strategy and Government Policy to Improve MSMEs Performance During the Covid-19 Pandemic	1120
<i>Danang Kusnanto, Solihin Sidik, Novian Ekawaty, Hana Salsabila Kastanya, Lida Arlini, and Alya Farah Azzahra</i>	
The Role of Science in Islamic Religious Education in the Modern Era	1129
<i>Anip Dwi Saputro, Sudarmadi, Adib Khusnul Rois, Syamsul Arifin, Ja'far Arifin, and Maylina Wulandari</i>	

Implication of Technology in Traditional Sculpture Production in Batubulan Village Sukawati Bali	1235
<i>I Wayan Suardana, I Ketut Muka, and I Made Ruta</i>	
Problematics of Intolerance Conflict Between Religious People in Islamic Law Politics in the Digitalization Era	1242
<i>Dewi Iriani, Arief Budiono, Muhammad Fauzan, Layyin Mahfiana, Esti Ningrum, and Ely Masykuroh</i>	
Utilization of Antidiabetic Herbal Medicines Research Results as Adjuvant Treatment of Patients with Diabetes Mellitus Vulnerable to Covid-19 at the Elderly Posyandu, Sukoharjo Regency	1251
<i>Muhtadi, E. M. Sutrisna, Ihwan Susila, Andi Suhendi, Safira Bela Annisa, Febrianti Nur Antika, Amalia Siti Choerun Nisa, and Reza Rohmatun Zakiyah</i>	
Mapping Farmer's Digital Competencies Using The Agrowing Digital Platforms	1260
<i>Ahsin Daroini and Adisty Riska Hardianti</i>	
Sangku as Indonesian Cultural Property	1268
<i>I Wayan Mudra, I Nyoman Larry Julianto, I P. Ketut Muka, and I Wayan Swandi</i>	
Bibliometric Analysis of Iron Overload (IO) of Thalassemia Disease	1275
<i>Titi Pudji Rahayu, S. Yuliani, and H. Susanti</i>	
Streamlining the Bureaucracy for First-Time Voters in Indonesian General Elections	1281
<i>Anom Wahyu Asmorojati, Aliz Zulis Al Hurni, and Danang Rizky Fadilla</i>	
Training for Competency Improvement of Widyaiswara in Writing Scientific Writings in the Field of Training and Human Resources Development Agency (BKPSDM) Karawang District	1290
<i>Solehudin and Dayat Hidayat</i>	
The Effectiveness of Kinds of Rubber Rope Games Media to Improve Fine Motor Skills for Early Childhood	1303
<i>Ega Trisna Rahayu, Rina Syafrida, Nurunnabilah, and Haliza Syahnurmala</i>	
Implementation of Inclusive Education at Elementary School Level in Surakarta	1313
<i>Ardheila Setya Yudhani and Honest Umami Kaltsum</i>	

Implication of Technology in Traditional Sculpture Production in Batubulan Village Sukawati Bali	1235
<i>I Wayan Suardana, I Ketut Muka, and I Made Ruta</i>	
Problematics of Intolerance Conflict Between Religious People in Islamic Law Politics in the Digitalization Era	1242
<i>Dewi Iriani, Arief Budiono, Muhammad Fauzan, Layyin Mahfiana, Esti Ningrum, and Ely Masykuroh</i>	
Utilization of Antidiabetic Herbal Medicines Research Results as Adjuvant Treatment of Patients with Diabetes Mellitus Vulnerable to Covid-19 at the Elderly Posyandu, Sukoharjo Regency	1251
<i>Muhtadi, E. M. Sutrisna, Ihwan Susila, Andi Suhendi, Safira Bela Annisa, Febrianti Nur Antika, Amalia Siti Choerun Nisa, and Reza Rohmatun Zakiyah</i>	
Mapping Farmer's Digital Competencies Using The Agrowing Digital Platforms	1260
<i>Ahsin Daroini and Adisty Riska Hardianti</i>	
Sangku as Indonesian Cultural Property	1268
<i>I Wayan Mudra, I Nyoman Larry Julianto, I P. Ketut Muka, and I Wayan Swandi</i>	
Bibliometric Analysis of Iron Overload (IO) of Thalassemia Disease	1275
<i>Titi Pudji Rahayu, S. Yuliani, and H. Susanti</i>	
Streamlining the Bureaucracy for First-Time Voters in Indonesian General Elections	1281
<i>Anom Wahyu Asmorojati, Aliz Zulis Al Hurni, and Danang Rizky Fadilla</i>	
Training for Competency Improvement of Widyaiswara in Writing Scientific Writings in the Field of Training and Human Resources Development Agency (BKPSDM) Karawang District	1290
<i>Solehudin and Dayat Hidayat</i>	
The Effectiveness of Kinds of Rubber Rope Games Media to Improve Fine Motor Skills for Early Childhood	1303
<i>Ega Trisna Rahayu, Rina Syafrida, Nurunnabilah, and Haliza Syahnurmala</i>	
Implementation of Inclusive Education at Elementary School Level in Surakarta	1313
<i>Ardheila Setya Yudhani and Honest Ummi Kaltsum</i>	

Formalin and Borax Content of Chicken Sempol Marketed in Purworejo Regency, Indonesia	1322
<i>Roisu Eny Mudawaroch, Rinawidiastuti, and Muhammad Azziz</i>	
Development of Teaching Materials Based on Blended Learning for Elementary School Students	1330
<i>Nita Hidayati, Lessa Roesdiana, and Betha Nurina Sari</i>	
A Systematic Literature Review on Video Media: Application to Mathematics Learning	1336
<i>Tria Mardiana and Sardin</i>	
Investigation of the Role of Entrepreneurship Characteristics in Moderating Creativity and the Business Environment towards Business Sustainability	1350
<i>Rochiyati Murniningsih, Herning Indriastuti, and Siti Noor Khikmah</i>	
A Model of Freshwater Fisheries Development Strategy in the Context of Economic Resilience and Environmental Sustainability - evidence in Indonesia	1365
<i>Retno Rusdijati, Veni Soraya Dewi, and Barkah Susanto</i>	
Investigation of Firm Value in Transportation Sector: The Impact of Covid 19 in Indonesia	1380
<i>Lilik Andriyani, Veni Soraya Dewi, and Catur Dewi Larasati</i>	
Health and Halal of Sea Products: Legal Perspective Halal Product Assurance	1389
<i>Arief Budiono, Dewi Iriani, Sofyan Wimbo Agung Pradnyawan, Ayesha Hendriana Ngestiningrum, Abdullah al Mamun, and Muhammad Amin Hanafi</i>	
Model of Tourism Village Development in Karangasem Village, Bulu District, Sukoharjo Regency as Family Tourist Destinations Based on Local Wisdom	1397
<i>Ratna Susanti, Suci Purwandari, and Basnendar Herry Prilosadoso</i>	
Creation Process and Visual Aesthetics at Jesus and Mary Images in Balinese Wayang Characters	1403
<i>Andreas James Darmawan, I Wayan Mudra, and I Ketut Suteja</i>	

Socialization of SAK EMKM, Perceptions of MSME Actors, and Education Level of MSME Owners on the Implementation of SAK EMKM in Financial Reports for MSMEs in Ponorogo District	1414
<i>Vianita Diah Safitri, Khusnatul Zulfa Wafirotin, Iin Wijayanti, Slamet Santoso, and Nurul Hidayah</i>	
Innovative Wayang as an Educational Means to Prevent the Spread of Covid-19 “Sang Kala Korona”	1424
<i>I Kadek Widnyana, Sulistyani, and Rinto Widyanto</i>	
The Effect of Gender and High School Origin towards English Grammar Mastery	1431
<i>Johan Arief Budiman and Harris Gadih Pratomo</i>	
Investigative Audit and Forensic Accounting from the Perspective of Criminal Procedural Law	1438
<i>Ramadhana Anindyajati Bachry, Andika Surya, and Noviard Rizky</i>	
Cianjur Earthquake Disaster Emergency Management: Muhammadiyah Disaster Management Center Sitrep Study	1445
<i>Margono</i>	
Motives, Messages, and Media in the Process of Child Marriage in Madura	1452
<i>Netty Dyah Kurniasari, Emy Susanti, and Yuyun Wahyu Izzati Surya</i>	
Implementation of Education Quality Assurance Management in the Process Blended Learning	1458
<i>Bambang Ismaya, Astuti Darmiyanti, Abduloh, and Selis</i>	
Thyroid Disorders as a Factor Related for Stunting: A Literature Review	1469
<i>Septi Wardani and Dwi Sulistyono</i>	
The Influence of Servant Leadership and Self-Efficacy on Employee Performance with Job Satisfaction as Intervening Variables (Empirical Study on Rural Banks in the Magelang Region)	1475
<i>Atina Harin Hayuni and Muhdiyanto</i>	
Analysis of Fulfillment of the Rights of Students with Special Needs in Inclusive Schools (Case Study of SD Lauzardi Kamila Global Islamic School)	1487
<i>Andre Eksaputra Nugraha and Arief Cahyo Utomo</i>	

Indonesian Multiethnic of the Dinasti Matahari Music Video in the Iconography Context	1494
<i>Nyoman Lia Susanthi, Ketut Hery Budiayana, Ni Wayan Masyuni Sujayanthi, and Igor de Almaida Amanajas</i>	
Soft Innovation Strategy of Batik MSMEs in Indonesia	1509
<i>Dita Andansari, Khairul Hisyam Kamarudin, and Darius Shyafary</i>	
The Influence of the Coordination Pattern of the Election Supervisory Board (Bawaslu) with the Luwuk District Supervisory Committee (Panwascam) on the Implementation of Elections in Banggai District in 2020	1514
<i>Erwin Nursin, Risno Mina, and Gisela Luigi Septian</i>	
Mobile Learning to Improve Mathematical Communication Skills	1522
<i>Adi Ihsan Imami and Fitria Febriyanti</i>	
How is HOTS Applied in Elementary Schools? A Review	1529
<i>Agrissto Bintang Aji Pradana, Aditia Eska Wardana, Putri Meinita Triana, and Fitra Amaldi</i>	
The Culture of Packaging Traditional Balinese Snacks with Banana Leaves	1538
<i>Ni Luh Desi In Diana Sari, Ni Ketut Pande Sarjani, and Eldiana Tri Narulita</i>	
The Effect of Learning Independence on Students' Self-Adjustment after the Covid-19 Pandemic at SMPN 1 Secang	1548
<i>Widya Maharani, Hermahayu, and Aftina Nurul Husna</i>	
Implementation of Character and Multicultural Education in Elementary Schools through Extracurricular Activities in SD Kartika XII-1	1553
<i>Puji Rahmawati and Aditia Eska Wardhana</i>	
The Effect of Peer Conformity on Intention to Buy In-Game Virtual Goods in Adolescent Free Fire Online Game Players	1562
<i>Yollawati Arifi, Aftina Nurul Husna, and Laili Qomariyah</i>	
Islamic Corporate Governance and Sharia Compliance on Financial Performance	1571
<i>Farida and Muji Mranani</i>	
Punden Berundak of Pura Penulisan Supporting the Bali Tourism Industry	1578
<i>I Gede Mugi Raharja</i>	

Legal Standing of Criminal Confiscation Against General Confiscation Based on Indonesian Bankruptcy Procedural Law	1585
<i>Ramadhana Anindyajati Bachry</i>	
The Role of Management Control System in Non-Financial Performance	1591
<i>Yanti, Tubagus Ismail, Imam Abu Hanifah, and Munawar Muchlish</i>	
Environmental and Financial Performance of Publicly Traded Indonesian Companies	1601
<i>Ratna Wulaningrum, Venti Eka Satya, Amiril Azizah, and Dwi Yana Amalia Sari Fala</i>	
Improving Marketing Performance in Construction Companies as a Significant Impact from Value Co-Creation and Business Process Agility	1608
<i>Redi Nusantara, Vincent Didiek Wiet Aryanto, and Yohan Wismantoro</i>	
Application of Financial Technology and Increasing Literacy Finance for MSME Business Strengthening Strategy	1619
<i>Arisyahidin, Adisty Riska Hardianti, and Aji Seto Arifianto</i>	
Correction to: The Power of Ergo-Iconic Values Applied to the Management of Scientific Seminar Implementation to Improve Service Quality	C1
<i>Andriyansah, Kartono, Ni Made Suwitri Parwati, Lukman Samboteng, N. S. Jurana, Siti Umamah Naili Muna, and Hadian Pratama Hamzah</i>	



Effect of Extraction Solvent on Extraction Yield, Cytotoxic Activity and Bioactive Compound in *Zingiber officinale* Roscoe var rubrum

Kirana Anggraini¹✉, Dyah Ayu Woro Setyaningrum¹, Laela Wulansari^{1,2}, Hening Tyas Andayani³, Laviany Putri Shihran¹, and Isra Fauziyyah¹

¹Universitas Trisakti, Jakarta, Indonesia
kirana_anggraeni@trisakti.ac.id

²Tropical Biopharmaca Research Center, Institute for Research and Community Service (LPPM) IPB University, Bogor, Indonesia

³Fachärztin der Innere Medizin, Städtische Kliniken, Mönchengladbach, Germany

Abstract. Red ginger (*Zingiber officinale* Roscoe var rubrum) is a commonly used spice in Indonesia. It contains various chemical constituents, such as phenolic compounds, terpenes, polysaccharides, lipids, organic acids, and fiber. The health benefits of ginger are mainly related to its phenolic compounds, such as gingerols and shogaols. 6-Gingerol has various biological properties including anticancer, antioxidant, anti-inflammatory, anti-platelet aggregation, and anti-fungal. The method used to obtain the red ginger extract was macerated using 30%, 70%, and 96% ethanols, n-hexane, ethyl acetate, and water. The research was started by testing the phytochemicals and water content of the simplicia. Furthermore, toxicity tests were carried out on each extract on shrimp larvae, and the determination of gingerol and shogaol levels was by High-Performance Liquid Chromatography. The results showed that the highest yield of red ginger extract was 12.1% in a water solvent. Phytochemical testing of red ginger simplicia contains saponins. The 96% ethanol extract had the highest cytotoxic activity with an LC50 value of -79.516 ppm. The highest contents of 6-gingerol, 8-gingerol, 10-gingerol, and 6-shogaol were found in n-hexane solvents, respectively 123.07 mg/g; 26.39 mg/g; 61.40 mg/g and 22.41 mg/g.

Keywords: Extraction Solvent, Cytotoxic Activity, Bioactive Compound, *Zingiber officinale* Roscoe var rubrum

1 Introduction

Ginger is one of the most widely used herbal plants in Asian countries across generations since hundreds of years ago in the medicinal and culinary fields. This plant belongs to the Zingiberaceae family and the Zingiberales order. The Zingiberaceae family has 50 genera with 1,300 species, some of which are widely found and used in Indonesia. The main ginger-producing countries in the world are India, China, Indonesia, and Nigeria. The scientific name for red ginger is *Zingiber officinale* Roscoe var. Rubrum,

© The Author(s) 2023

M. Setiyo et al. (eds.), *Proceedings of the 4th Borobudur International Symposium on Science and Technology 2022 (BIS-STE 2022)*, Advances in Engineering Research 225,

https://doi.org/10.2991/978-94-6463-284-2_86

known as a variant of ginger is different from other varieties, especially in its rhizome which is layered in orange to red [1].

It is common to use spice plants as medicine in Indonesia because they are easy to obtain, easy to process, and have been used for generations. One of the most used spices is a ginger rhizome. Red ginger extraction is a separation process that is carried out to obtain certain desired components from the starting material. The selection of methods, solvents, and steps greatly affects the quality of the resulting product. The bioactive compounds in red ginger have certain biological activities when red ginger is consumed or extracted [2]. Extraction is the process of soluble chemical compounds withdrawal, so that they are separated from materials that are insoluble in liquid solvents. Different solvents type will affect secondary metabolite compound produced from red ginger rhizome extraction process. Secondary metabolite is an inessential for organism growth, which can be found in different or unique types on each species. The function of secondary metabolite is to defend the plants from endangering environmental condition, for example to defend themselves from pests or diseases, to attract pollinators, and as signaling molecule [3]. The identification of secondary metabolite is an important initial stage to discover bioactive compound from natural materials, which can be made as precursor for novel synthetic medication or certain activity drug prototype generation [4]. The chemical content of ginger rhizome is volatile components (camphene, β -phellandrene, curcumene, cineole, geranyl acetate, terpineol, borneol, geraniol, limonene, β -elemene, zingiberol, linalool, α -zingiberene, β -sesquiphellandrene, β -bisabolene, zingiberenol and α -Farnesene) and non-volatile components which consists of biologically active components namely gingerols (6, 8 and 10), shogaol (6, 8, and 10), paradol and zingerone [5]. Ginger contains pungent phenolic compounds known as gingerols. One of them, 6-gingerol (1-[4'-hydroxy-3'-methoxyphenyl]-5-hydroxy-3-decanone), is the main pharmacologically active component of ginger [6, 7], and the active part of the molecule is part of the aliphatic chain containing the hydroxyl [8]. 6-Gingerol has been reported to have various biological properties including anticancer, antioxidant, anti-inflammatory, anti-platelet aggregation, and anti-fungal [9–11].

The composition of chemical compounds from plants (phytochemicals) is essential for determining the quality of herbal products because it determines their safety and effectiveness. The effectiveness of an extraction process of a compound by solvent is highly depended on the solubility of the compound with the solvent, according to the principle of like dissolve like, where a compound will be dissolved on solvent with similar natures, based on the polarity of the compound in the solvent during extraction process. Polar compound can only be dissolved on polar solvents such as ethanol, methanol, butane, and water. Non-polar compounds can only be dissolved on non-polar solvents such as ether, chloroform, and n-hexane [12]. This study is aimed to acknowledge the influence of different types of solvents (water, ethanol 30%, ethanol 70%, ethanol 96%, n-hexane, and ethyl acetate) on yield rate, cytotoxic activity, and bioactive compound content (6-gingerol, 8-gingerol, 10-gingerol and 6-shagol) on red ginger rhizome. Also to identify the most suitable solvent for the extraction process to isolate bioactive component and to gain cytotoxic activity of red ginger rhizome.

2 Method

2.1 Place, Time of Sampling, and Determination of Samples

The research material was the fruit of the red ginger plant taken from the Biopharmaca Cultivation and Conservation Unit (BCCU) of the Tropical Biopharmaceutical Research Center, LPPM IPB University which was obtained from Java and harvested at the age of 9 months. The determination was carried out at the Biopharmaca Cultivation and Conservation Unit (BCCU).

2.2 Making Simplicia

Three kilograms of fresh red ginger rhizome were weighed and then washed and dried in direct sunlight for 4-5 days. Then, the dry sorting was done and mashed with a blender. The simplicia powder obtained was sieved using an 80 mesh sieve and then weighed. After that, it was stored in a clean, dry container and protected from sunlight for the next extraction process.

2.3 Preparation of Red Ginger Extract

Extract preparation and testing were carried out at the Laboratory of the Tropical Biopharmaca Research Center, LPPM IPB University. Samples of red ginger simplicia were weighed for extraction with 10 grams of various solvents for each, then solvents were added namely 96%, 70%, 30% ethanol, 500 mL of water, ethyl acetate, and n-hexane. Maceration was carried out 2 x 24 hours with several times of stirring, then filtering. The collected filtrate was concentrated using a vacuum rotary evaporator at 45-50°C to obtain a viscous ethanol extract of 96%, 70%, 30%, water extract, ethyl acetate extract, and n-hexane extract.

2.4 Water content

Two grams of red ginger simplicia were weighed in a container with a constant weight. Then, red ginger simplicia was heated in an oven at a temperature of +105 degrees Celsius for 3 hours. After being heated and then cooled in a desiccator, it was weighed until it reached a constant weight [13].

2.5 Analysis of Red Ginger Extract Yield

The yield of red ginger extract was calculated by comparing the weight of the red ginger extract with the weight of the red ginger simplicia used for extraction.

2.6 Phytochemical Screening

Phytochemical screening in this study includes the tests for alkaloids, flavonoids, tannins, saponins, triterpenoids, and steroids based on the method of Harborne 1987.

2.7 Alkaloids

0.5 grams of condensed extract or 1 gram of simplicia was dripped with 3-5 drops of ammonia. Then, 5 mL of chloroform was added. After that, it was homogenized and filtered. The filtrate obtained was added with 2M Sulfuric acid reagent, then it was homogenized. The top layer was taken and used as an experimental solution which was treated as follows: 1) Experimental solvent 1 was added 2 drops of Mayer's reagent, a positive result with the formation of a white precipitate. 2) Experimental solvent 2 was added 2 drops of Dragendorff reagent, positive result with the formation of an orange or orange precipitate. 3) Experimental solvent 3 was added 2 drops of Wagner's reagent, positive result with the formation of a brown precipitate.

2.8 Flavonoids, Tannins, and Saponins

0.5 grams of condensed extract or 5 grams of simplicia was dissolved in distilled water and then heated for 5 minutes. Then, the solvent was filtered and divided into 3 parts. To test the flavonoid filtrate, Mg and HCl powder:Ethanol (1:1) were added, then amyl alcohol was added. The color formed on the amyl alcohol layer was observed, if a yellow, orange or red color is formed then it contains flavonoids. For the tannin filtrate test, 3 drops of 10% FeCl₃ were added, if a greenish-black color is formed then it contains tannins. For the saponin test, the filtrate was shaken for 10 seconds. A positive result is indicated by the formation of stable foam for over 2 minutes.

2.9 Triterpenoids and Steroids

A sample of 0.5 grams of condensed extract or 1 gram of simplicia was dissolved in ethanol and heated for 5 minutes, then the sample was filtered into a porcelain dish. Then, the filtrate was heated to dryness, and 1 mL of diethyl ether, 1 drop of acetic anhydrous, and 1 drop of concentrated sulfuric acid were added. A positive reaction is indicated by the formation of a red/purple solution for triterpenoids and blue or green for steroids.

2.10 Quinone

0.5 gram of condensed extract or 1 gram of simplicia was added with methanol and then heated, after that it was filtered. The filtrate results are added 3 drops of 10% NaOH. A positive reaction is indicated by the formation of a red color for hydroquinone [14].

2.11 Cytotoxic Activity Test on Shrimp Larvae

The cytotoxic activity test by determining the LC50 value was carried out using *Artemia salina* shrimp eggs. *A. salina* used for the toxicity test was obtained from hatching using seawater with the help of an aerator to meet dissolved oxygen levels. The extract toxicity test was carried out using *A. salina* shrimp larvae. The shrimp larvae used were aged 48 hours after the shrimp larvae hatched. *A. salina* cysts of as much as ± 50 mg were put into a container containing seawater that had been filtered and equipped with an aerator. The cysts were left for 48 hours under light to hatch completely. After hatching, 10 *A. salina* larvae were put into a 2 ml vial, then it was added a stock extract solution with a concentration of 4000 ppm and adjusted the volume with seawater so that the final concentration of the extract was 0, 10, 100, and 1000 ppm. After 24 hours, the number of dead larvae was counted. The lethal concentration (LC) value was determined by the probit analysis method with a 95% confidence interval [15].

2.12 Determination of Gingerol content with High-Performance Liquid Chromatography (HPLC)

Standard and sample preparation. Standards of 6.8 and 10 gingerols and 6 shogaols dissolved in methanol were made at concentrations of 50 ppm, 25 ppm, 50 ppm, and 50 ppm respectively. Each extract was weighed as much as 0.1 gram and then added 8 mL of methanol solvent, was then sonication was done for 1 hour. Then the sample solvent was filtered into a 10 mL flask and then calibrated with methanol up to 10 mL, then filtered with 0.45 micrometer Whatman filter paper and then 20 μ L injected into the HPLC.

Identification by HPLC. The mobile phase used was Acetonitrile and aqua bides with a composition comparison in Table 1. The wavelength used was 280 nm, while the flow of the mobile phase was 1 mL/minute. Table 1. Comparison of the composition of Acetonitrile and Aqua bides:

Table 1. The composition of the mobile phase in the determination of gingerol content

Time (minutes)	Acetonitrile (%)	Aqua bides (%)
0	40	60
10	40	60
40	10	90
40.5	0	100
45	0	100
45.5	60	40
50	60	40

3 Result and Discussion

3.1 Plant Determination

The test sample was identified at the Biopharmaca Conservation & Cultivation Station (BCCS) Tropical Biopharmaceutical Study Center, Institute for Research and Community Service (LPPM) IPB University, showing a sample of Red Ginger (*Zingiber Officinale* Roscoe var *rubrum*) from the Zingiberaceae tribe.

3.2 Phytochemical compounds

Tests for the content of phytochemical compounds were carried out on *simplicia*, which contained saponin compounds. The positive reactions in the flavonoids, saponins, and tannins indicate the presence of phenol groups.

3.3 Water Content and Extraction

The water content of red ginger *simplicia* obtained was 10%, fulfilling the quality requirements so that it can be used for further analysis. Removing the water content up to a certain amount is useful for extending the durability of *simplicia*. Water content that is too high will become a medium for the growth of microorganisms that cause damage to the *simplicia* [16]. The viscous extract obtained is blackish-brown in color, and has a distinctive aroma of ginger. The color of red ginger extract produced in n-hexane, ethyl acetate, and ethanol solvents is a dark brown liquid and slightly viscous.

3.4 Impact of Extraction Solvents on the Yield of the Red Ginger Extract

The yield of the red ginger extract can be seen in Table 2. The highest yield was found in an aqueous extract. This extract was then tested for its toxicity to shrimp larvae, and a test was carried out to determine the levels of 6-gingerol, 8-gingerol, 10-gingerol, and 6-shogaol by HPLC.

Table 2. The yield of red ginger rhizome extract

	ETOH 30%	ETOH 70%	ETOH 96%	Water	Ethyl acetate	N hexane
Extract (g)	11.02	10.06	9.1	12.1	5.9	2.5

Impact of Extraction Solvents on Shrimp Larvae Cytotoxic Activity Test. Shrimp larvae cytotoxic activity tests were carried out to observe the potential bioactivity and toxicity of each extract so that a safe extract concentration could be determined for further testing. A plant extract will be bioactive if it has an LC50 value of less than 1000 ppm [15]. Based on Table 3 it can be seen that all red ginger rhizome extracts have the potential as bioactive compounds and can be used as medicine. It was because each extract produced an LC50 of less than 1000 ppm so that at low concentrations it

was able to kill 50% of the population of *A. salina* shrimp larvae. The extract that has the highest bioactive potential and is toxic was 96% ethanol extract. It is because the 96% ethanol extract of red ginger rhizome has the lowest LC₅₀ value, namely -79.516 ppm, which means that at a small concentration, this extract can kill half the population of *A. salina* shrimp larvae. The LC₅₀ value is the highest concentration limit for determining various extract concentrations in subsequent tests.

Table 3. LC₅₀ value of red ginger rhizome extract against *A. Salina* larvae

	ETOH 30%	ETOH 70%	ETOH 96%	Water	Ethyl acetate	N hexan
LC ₅₀ (ppm)	208.62	365.39	-79.516	290.16	191.47	365.89

3.5 Impact of Extraction Solvents on Content of Gingerol and Shogaol.

The results of calculating the concentration of gingerol and shogaol compounds in each extract can be seen in Figure 1-4. The highest 6-gingerol compound is found on n-hexane solvent, followed by ethyl acetate, water, and ethanol 96% respectively at 123.07 mg/g, 81.79 mg/g, 31.15 mg/g, and 12.34 mg/g. The highest 10-gingerol content was discovered on n-hexane solvent, followed by ethyl acetate, and ethanol 96% respectively at 61.40 mg/g, 36.28 mg/g, and 5.10 mg/g. The highest 8-gingerol content was discovered in n-hexane solvent, followed by ethyl acetate, and ethanol 96% respectively at 26.39 mg/g, 16.70 mg/g, and 2.68 mg/g. The highest 6-shagol content was discovered on n-hexane solvent, followed by ethyl acetate, and ethanol 96% respectively at 24.41 mg/g, 14.70 mg/g, and 2.73 mg/g. It shows that the compounds 6-gingerol, 8-gingerol, 10-gingerol, and 6-shogaol can be extracted higher with non-polar solvents. In general, the bioactive components in red ginger are non-polar, characterized by more bioactive components that dissolve in hexane, then ethyl acetate and ethanol [17]. It is also caused by ethyl acetate, which is a semi polar solvent that has the ability to attract both polar and non-polar compounds, meanwhile ethanol 96% is a universal solvent that can attract polar, non-polar, or semi polar compounds [18].

The content of 6-gingerol, 8-gingerol, and 10-gingerol in each red ginger extract in this study was higher than in previous studies [19, 20]. The difference in the high content of gingerol in this study and the previous study could be due to the different ecological conditions of the ginger plant, the variety of ginger used, the age of ginger harvest, the method of making ginger simplicia, and the extraction method used.




Fig. 1. Graph of 6-gingerol levels of various red ginger extracts




Fig. 2. Graph of 10-gingerol levels of various red ginger extracts




Fig. 3. Graph of 8-gingerol levels of various red ginger extracts




Fig. 4. Graph of 6-shogaol levels of various red ginger extracts

4 Conclusion

The different type of solvent will produce different amount of rhizome, cytotoxic activities, and gingerol and shagol bioactive compounds. The highest yield of red ginger extract was 12.1% in a water solvent. The secondary metabolites in red ginger simplicia are saponins. The extract that has the highest bioactive potential and cytotoxic activity is 96% ethanol extract. The highest content of 6-gingerol was found in n-hexane solvent of 123.07 mg/g. The highest content of 8-gingerol was found in the n-hexane solvent of 26.39 mg/g. The highest content of 10-gingerol was found in n-hexane solvent of 61.40 mg/g. The highest content of 6-shogaol was found in n-hexane solvent of 22.41 mg/g.

Acknowledgements. We thank Universitas Trisakti for funding this research through the Faculty Research grant (PUF) Number: 322/A.1/LPT/USAKTI/II/2022 on behalf of dr. Kirana Anggraini MKM.

References

1. F. Nurdyansyah and D. Widyastuti, *Jahe Merah: Senyawa Bioaktif, Manfaat, dan Metode Analisisnya*. Bandung: Widina Bhakti Persada Bandung, 2022.
2. M. Adiyasa and M. Meiyanti, "Pemanfaatan obat tradisional di Indonesia: Distribusi dan faktor demografis yang berpengaruh," *J. Biomedika dan Kesehat.*, vol. 4, no. 3, pp. 130–138, 2021, doi: 10.18051/JBiomedKes.2021.v4.130-138.
3. R. Verpoorte, *Secondary Metabolism BT - Metabolic Engineering of Plant Secondary Metabolism*. Dordrecht: Springer Netherlands, 2000.
4. J. D. Phillipson, "Phytochemistry and medicinal plants," *Phytochemistry*, vol. 56, no. 3, pp. 237–243, 2001, doi: [https://doi.org/10.1016/S0031-9422\(00\)00456-8](https://doi.org/10.1016/S0031-9422(00)00456-8).
5. V. Govindarajan and D. Connell, "Ginger—chemistry, technology, and quality evaluation: part 1," *Crit. Rev. Food Sci. Nutr.*, vol. 17, no. 1, pp. 1–96, 1983.

6. A. M. Bode, W.-Y. Ma, Y.-J. Surh, and Z. Dong, "Inhibition of Epidermal Growth Factor-induced Cell Transformation and Activator Protein 1 Activation by [6]-Gingerol," *Cancer Res.*, vol. 61, no. 3, pp. 850–853, Feb. 2001.
7. Y.-J. Surh, "Cancer chemoprevention with dietary phytochemicals," *Nat. Rev. Cancer*, vol. 3, no. 10, pp. 768–780, 2003, doi: 10.1038/nrc1189.
8. G. Yang et al., "Genotoxic effect of 6-gingerol on human hepatoma G2 cells," *Chem. Biol. Interact.*, vol. 185, no. 1, pp. 12–17, 2010, doi: <https://doi.org/10.1016/j.cbi.2010.02.017>.
9. Q.-Y. Wei, J.-P. Ma, Y.-J. Cai, L. Yang, and Z.-L. Liu, "Cytotoxic and apoptotic activities of diarylheptanoids and gingerol-related compounds from the rhizome of Chinese ginger," *J. Ethnopharmacol.*, vol. 102, no. 2, pp. 177–184, 2005, doi: <https://doi.org/10.1016/j.jep.2005.05.043>.
10. E. Tjendraputra, V. H. Tran, D. Liu-Brennan, B. D. Roufogalis, and C. C. Duke, "Effect of Ginger Constituents and Synthetic Analogues on Cyclooxygenase-2 Enzyme in Intact Cells," *Bioorg. Chem.*, vol. 29, no. 3, pp. 156–163, 2001, doi: <https://doi.org/10.1006/bioo.2001.1208>.
11. C. Ficker et al., "Bioassay-guided isolation and identification of antifungal compounds from ginger," *Phyther. Res.*, vol. 17, no. 8, pp. 897–902, Sep. 2003, doi: <https://doi.org/10.1002/ptr.1335>.
12. W. Leksono, R. Pramesti, G. Santosa, and W. Setyati, "Jenis pelarut metanol dan N-Heksana terhadap aktivitas antioksidan ekstrak rumput laut gelidium sp. dari pantai drini Gunungkidul–Yogyakarta," *J. Kelaut. Trop.*, vol. 21, no. 1, pp. 9–16, 2018.
13. Kemenkes RI, *Farmakope Herbal Indonesia*. Jakarta: Kementerian Kesehatan, Republik Indonesia, 2008.
14. A. Harborne, *Phytochemical methods a guide to modern techniques of plant analysis*. Berlin, Heidelberg: Springer Dordrecht, 1998.
15. B. N. F. Meyer, N. R. Putnam, J. E. Jacobsen, L. B. Nichols, D. E. McLaughlin, J. L., "Brine Shrimp: A Convenient General Bioassay for Active Plant Constituents," *Planta Med.*, vol. 45, no. 05, pp. 31–34, 1982, doi: 10.1055/s-2007-971236.
16. E. Yuslianti, B. Bachtari, D. Suniarti, and A. Sutjiatmo, "Standardisasi farmasitikal bahan alam menuju fitofarmaka untuk pengembangan obat tradisional Indonesia," *Dentika Dent. J.*, vol. 19, no. 2, pp. 179–185, 2016.
17. K. Ahmed, G. Shaheen, and H. Asif, "Zingiber officinale Roscoe (pharmacological activity)," *J. Med. Plants Res.*, vol. 5, no. 3, pp. 344–348, 2011.
18. J. Santoso, S. Anwarayah, R. O. Rumiantin, A. P. Putri, N. Ukhty, and Y. Y. - Stark, "Phenol Content, Antioxidant Activity and Fibers Profile of Four Tropical Seagrasses From Indonesia," *J. Coast. Zo. Manag.*, vol. 15, no. 2, pp. 189–196, 2012.
19. F. Tririzqi, "Ekstraksi Senyawa Gingerol dari Rimpang Jahe dengan Metode Maserasi Bertingkat," IPB University, 2013.
20. S. Srikandi, M. Humaeroh, and R. Sutamihardja, "Kandungan gingerol dan shogaol dari ekstrak jahe merah (Zingiber officinale Roscoe) dengan metode maserasi bertingkat," *Al-Kimiya J. Ilmu Kim. dan Terap.*, vol. 7, no. 2, pp. 75–81, 2020.

Open Access This chapter is licensed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits any noncommercial use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this chapter are included in the chapter's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the chapter's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.





Influence of Extraction Solvents on Capsaicin Compound and Cytotoxic Activity of *Piper retrofractum Vahl*

Kirana Anggraini¹ (✉), Dyah Ayu Woro Setyaningrum¹, Laela Wulansari^{1,2}, Hening Tyas Andayani³, Laviyany Putri Shihran¹, and Isra Fauziyyah¹

¹Universitas Trisakti, Jakarta, Indonesia
kirana_anggraeni@trisakti.ac.id

²Tropical Biopharmaca Research Center, Institute for Research and Community Service (LPPM) IPB University, Bogor, Indonesia

³Fachärztin der Innere Medizin, Städtische Kliniken, Mönchengladbach, Germany

Abstract. Capsaicin (8-Methyl-N-vanillyl-trans-6-nonenamide) is an active compound found in plants from the genus *Capsicum*. Genus *Capsicum* possesses different species of red chilli (*Capsicum annuum Linn*), green chilli (*Capsicum annuum var. annuum*) and other varieties that are familiar in Indonesia. However, the plants were not originally from Indonesia but from American Continent. Indonesian ancestor utilized chilli to add spiciness on foods which refers to *Piper retrofractum Vahl* or Javanese chilli. Capsaicin has positive effects on health such as analgesic, cardioprotective, anorexigenic, chemopreventive, chemotherapy, and cardioprotective effects. In addition, capsaicin has beneficial effects in maintaining glucose levels, insulin homeostasis, reducing itching and symptoms in Non-Allergic Rhinitis (NAR), and an alternative therapy for neurogenic bladder. This research is aimed to acknowledge if capsaicin is also discovered in Javanese chilli. The method used to obtain the extract was by maceration of Java chillies using 30%, 70%, and 96% ethanol, n-hexane, ethyl acetate, and water as solvents. The research was started by conducting phytochemical and moisture content tests on *simplicia*. Then, each extract was tested for cytotoxic activity on shrimp larvae and the determination of capsaicin levels by High Performance Liquid Chromatography (HPLC). The results showed that the highest yield of Java chili extract was 51.14% in ethanol 70%. Phytochemical testing of Java chili *simplicia* contains flavonoids and saponins. The ethanol 70% extract had the highest cytotoxic activity with an LC50 value of 205.21 ppm. The highest concentration of capsaicin was found in ethyl acetate solvent of 785.58 mg/g followed by ethanol 70% at 486.06mg/g.

Keywords: Extraction Solvents, Capsaicin Compound, Cytotoxic Activity, *Piper retrofractum Vahl*

1 Introduction

Capsaicin (8-Methyl-N-vanillyl-trans-6-nonenamide) is the active compound found in hot peppers that gives chillies their hot taste [1]. Capsaicin was first purified in 1876 but

its structure began to be described in 1919 [2]. Due to its chemical structure, capsaicin is well absorbed up to 94% when administered topically or orally [3]. The use of spice plants is also influenced by the level of knowledge, age, education level, economic status, environmental factors, and sources of information/information media [4]. Studies demonstrate the efficacy of capsaicin as an analgesic. Treatment with capsaicin is effective in various types of pain conditions such as complex regional pain syndromes, neuropathic pain postsurgical neuropathic pain [5] post-herpetic neuralgia [6] and painful diabetic peripheral neuropathy [7]. Research has also shown capsaicin to be effective in weight loss and obesity improvement [8]. Capsaicin also has beneficial effects on glucose levels, insulin homeostasis, and diabetes [9].

The evidence showed the beneficial effects of capsaicin on the cardiovascular system [10]. Capsaicin activates TRPV1 which can stimulate the release of CGRP, which is the most powerful vasodilator that regulates blood pressure in both physiological and pathophysiological conditions, producing a blood pressure-lowering effect [11]. Capsaicin has been shown to inhibit platelet aggregation [12], which may also provide protection against cardiovascular disease [13]. Long-term activation of TRPV1 by capsaicin will inhibit the formation of foam cells so that it will reduce lipid storage and atherosclerotic lesions in the aortic sinus and thoracoabdominal aorta of rats, and ultimately slow down the process of atherosclerosis. The antioxidant properties of capsaicin contribute to its protective effect on the cardiovascular system. LDL oxidation is an early factor for the formation and development of atherosclerosis [14].

Capsaicin has been shown to have chemopreventive and chemotherapeutic effects [15]. Capsaicin also works on itching with nostalgia paresthetica [16], and neuropathic pruritus [17]. Capsaicin has a gastroprotective effect by modulating sensory neurons [18]. Capsaicin has been studied as an alternative therapy for symptomatic relief of neurogenic bladder, a urological disorder that seriously affects patients' quality of life [19]. The clinical use of capsaicin is also successful in patients with detrusor bladder hyperreflexia with multiple sclerosis, as well as after spinal cord injury [20].

Capsaicin is discovered in various type of capsium genus plants such as red chilli (*Capsicum annum* Linn), green chilli (*Capsicum annum* var. *annuum*) and other varieties known in Indonesia. Capsicum is not originally from Nusantara, instead it comes from American Continent. Indonesian ancestors used chilli to add spiciness on food currently refers to *Piper retrofractum* Vahl or green chilli. This research was conducted to determine the capsaicin content in Javanese chilies, as well as to find out the best solvent in the extraction process from Javanese chilies which produced the highest capsaicin levels.

2 Method

2.1 Place, Collecting, and Determining Samples

The research material was the fruit of the Javanese chili plant taken from the Biopharmaca Cultivation Conservation Unit (BCCU) of the Tropical Biopharmaca Research Center, LPPM IPB, which was obtained from Java which was harvested at the age of

3-4 months after flowering. The determination was carried out at the Biopharmaca Cultivation Conservation Unit (BCCU).

2.2 Simplicia Making

Three kilograms of fresh Java chilies were weighed, then washed and dried in direct sunlight for 4-5 days. After drying, the dry sorting is carried out, and mashed with a blender. The simplicia powder obtained was sieved using an 80-mesh sieve and then weighed. The simplicia powder is then stored in a clean, dry container and protected from sunlight for the next extraction process.

2.3 Production of Java Chilli Extract

Extract preparation and testing were carried out at the Laboratory of the Tropical Biopharmaca Research Center, Institute for Research and Community Service (LPPM), IPB University. Samples of Java chili simplicia were weighed for extraction with 10 grams of various solvents each, then added solvents namely 96%, 70%, 30% ethanol, 500 mL of water, ethyl acetate, and n-hexane. Maceration was carried out 2 x 24 hours with several times of stirring, then it was filtered. The collected filtrate was concentrated using a vacuum rotary evaporator at 45-50°C to obtain a viscous ethanol extract of 96%, 70%, 30%, water, ethyl acetate, and n-hexane extract.

2.4 Water content

Two grams of Javanese chili simplicia were weighed in a container with a constant weight. Then, it was heated in an oven at + 105 degrees Celsius for 3 hours. After that, it was cooled in a desiccator and weighed until it reached a constant weight.

2.5 Yield Analysis of Javanese Chilli Extract

The yield of Javanese chili extract was calculated by comparing the weight of the Javanese chili extract with the weight of the Javanese chili simplicia used for extraction.

2.6 Phytochemical Screening

Phytochemical screenings were carried out for alkaloids, flavonoids, tannins, saponins, triterpenoids, and steroids based on the method of Harborne 1987.

2.7 Alkaloids

A total of 0.5 grams of condensed extract or 1 gram of simplicial was dripped with 3-5 drops of ammonia, then 5 mL of chloroform was added, then it was homogenized and filtered. The filtrate obtained was added with 2M Sulfuric acid reagent, then homogenized. The top layer was taken and used as an experimental solvent which was then

treated as follows: 1) Experimental solvent 1 was added with 2 drops of Mayer's reagent, a positive result indicated by the formation of a white precipitate. 2) Experiment solvent 2 was added with 2 drops of Dragendorff reagent, and a positive result was indicated by the formation of an orange precipitate. 3) Experiment solvent 3 was added with 2 drops of Wagner reagent, and a positive result was indicated by the formation of a brown precipitate.

2.8 Flavonoids, Tannins, and Saponins

0.5 grams of condensed extract or 5 grams of simplicial was dissolved in distilled water and heated for 5 minutes. Then the solvent was filtered and the filtrate was divided into 3 parts. To test the flavonoid filtrate, Mg powder and HCl : Ethanol (1:1) were added, followed by amyl alcohol. The color formed on the amyl alcohol layer was observed, if a yellow, orange or red coloris formed then it consists of flavonoids. For the tannin filtrate test, 3 drops of 10% FeCl₃ were added, if a greenish-black color is formed then it consists of tannins. For the saponin test, the filtrate was shaken for 10 seconds. A positive result is indicated by the formation of stable foam for over 2 minutes.

2.9 Triterpenoids and Steroids

0.5 grams sample of condensed extract or 1 gram of simplicial was dissolved in ethanol and heated for 5 minutes, then the sample was filtered into a porcelain dish. The filtrate was heated to dryness and then 1 mL of diethyl ether, 1 drop of acetic anhydrous, and 1 drop of concentrated sulfuric acid were added. A positive reaction is indicated by the formation of a red/purple solvent for triterpenoids and blue or green for steroids.

2.10 Quinone

0.5 grams of condensed extract or 1 gram of simplicial was added to methanol then it was heated and filtered. The filtrate results were added with 3 drops of 10% NaOH. A positive reaction is indicated by the formation of a red color for hydroquinone [21].

2.11 Cytotoxic Activity Test on Shrimp Larvae

The cytotoxic activity test by determining the LC₅₀ value was carried out using *Artemia salina* shrimp eggs. *A. salina* used for the toxicity test was obtained from hatching using seawater with the help of an aerator to meet dissolved oxygen levels. The extract toxicity test was carried out using *A. salina* shrimp larvae. The shrimp larvae used were aged 48 hours after the shrimp larvae hatched. *A. salina* cysts of as much as ± 50 mg were put into a container containing seawater that had been filtered and equipped with an aerator. The cysts were left for 48 hours under light to hatch completely. After hatching, 10 *A. salina* larvae were put into a 2 ml vial, then it was added a stock extract solution with a concentration of 4000 ppm and adjusted the volume with seawater so that the final concentration of the extract was 0, 10, 100, and 1000 ppm. After 24 hours,

the number of dead larvae was counted. The lethal concentration (LC) value was determined by the probit analysis method with a 95% confidence interval [22].

2.12 Determination of Capsaicin content with High Performance Liquid Chromatography (HPLC)

Standard and sample preparation. Standard capsaicin was dissolved in methanol to make a concentration of 100 ppm. Each extract was weighed as much as 0.1 gram and then 8 mL of methanol solvent was added and sonicated for 1 hour. Then the sample solution was filtered into a 10 mL flask and calibrated with methanol up to 10 mL. After that, it was filtered with 0.45 micrometer Whatman filter paper and injected into HPLC as much as 20 μ L.

Identification by HPLC. The mobile phase used was methanol and aquabides with a composition ratio of 80:20. The wavelength was 235 nm, while the flow of the mobile phase was 1 mL/minute.

3 Result and Discussion

3.1 Plant Determination

The test sample was identified at the Biopharmaca Cultivation and Conservation Unit (BCCU) of the Tropical Biopharmaca Research Center, Institute for Research and Community Service (LPPM) IPB University, it showed that a sample of *Piper retrofractum* Vahl was from the Piperaceae tribe.

3.2 Phytochemical compounds

Tests for the content of phytochemical compounds were carried out on simplicia, which contained saponin and flavonoid. The positive reactions in the flavonoids, saponins, and tannins indicate the presence of phenol groups

3.3 Water Content and Extraction

The water content results of the Javanese chili simplicia were 8.75%, fulfilling the quality requirements and can be used for further analysis. Removing the water content up to a certain amount is useful for extending the durability of simplicia. Extremely high water content can become a medium for the growth of microorganisms that cause damage to the simplicia [23].

3.4 Rhizome Content on Different Types of Extract

The yield of Javanese chili extract is presented in Table 1, where the highest yield was found in 70% ethanol extract.

Table 1. The yield of Javanese chili extract

	ETOH 30%	ETOH 70%	ETOH 96%	Water	Ethyl acetate	N hexane
Extract(g)	7.59	51.14	8.44	11.85	6.33	8.30

3.5 Shrimp Larvae Cytotoxic Activity Test on Different Types of Extracting Solvent

Shrimp larvae toxicity tests were carried out to observe the potential bioactivity and cytotoxic activity of each extract so that a safe extract concentration could be determined for further testing. In addition, extracts that exhibit toxic properties can be developed as anti-cancer drugs [24]. A plant extract will be bioactive and have anti-cancer potential if it has an LC_{50} value of less than 1000 ppm. Based on Table 2, it can be seen that all Javanese chili extracts have the potential as bioactive compounds and can be used as medicine. It is due to the fact that each extract produced an LC_{50} of less than 1000 ppm so that, at low concentrations, it was able to kill 50% of the population of *A. salina* shrimp larvae. The highest bioactive potential and the toxic extract was 70% ethanol extract because it had the lowest LC_{50} value, which was 205.21 ppm, which means that, at a small concentration, this extract can kill half the population of *A. salina* shrimp larvae. The LC_{50} value is the highest concentration limit for determining various extract concentrations in subsequent tests [21].

Table 2. LC_{50} value of Javanese chili extract on *A. Salina* larvae

	ETOH 30%	ETOH 70%	ETOH 96%	Water	Ethyl acetate	Water
LC_{50} (ppm)	434.57	205.21	334.32	650.186	443.03	292.78

3.6 Capsaicin Content on Different Types of Extract

The results of the calculation of the capsaicin compound concentration in each extract can be seen in Fig 1. The highest capsaicin content was found in ethyl acetate solvent of 785.53mg/g followed by 70% ethanol of 426.34mg/g and the lowest in water solvent of 6.55mg/g. It is due to the fact that capsaicin is a non-polar phenolic compound so it cannot dissolve in water. The main solvents used to extract and maintain the properties of capsaicin are nonpolar solvents such as ether, benzene, dimethyl sulfoxide, and acetone, as well as ethanol due to their mixed nature [3].

The content of capsaicin in each extract of Javanese chilies in this study was higher than in previous studies conducted in Nepal and Saudi Arabia. In Nepal, research conducted on 16 types of chilies (*Capsium* fruit) found capsaicin levels ranging from 2.19 to 19.73 mg/g [25]. While research conducted in Saudi Arabia on hot chili, red chili, green chili, green pepper, red pepper, and yellow pepper found capsaicin levels range from 0.001 to 4.24 mg/g [26]. The difference related to the content of capsaicin

in this study and the previous study could be due to differences in plant ecological conditions, varieties, harvesting ages, simplicia manufacturing methods, and extraction methods used.




Fig. 1. Graph of capsaicin levels of various Javanese chili extracts

4 Conclusion

Capsaicin is discovered on *Piper retrofractum* Vahl fruits. The secondary metabolites in Javanese chili simplicia are flavonoids and saponins. The highest yield of extract was 11.85 g in a water solvent. The highest bioactive potential and cytotoxic activity extract is a 70% ethanol extract with an LC₅₀ of 205.21 ppm. The highest content of capsaicin was found in ethyl acetate solvent of 785.53mg/g followed by ethanol 70% at 486.06mg/g.

Acknowledgment. We thank Trisakti University for funding this research through the Faculty Research grant (PUF) Number: 322/A.1/LPT/USAKTI/II/2022 on behalf of dr. Kirana Anggraini MKM

References

1. E. K. Nelson and L. E. Dawson, "The constitution of capsaicin, the pungent principle of capsicum (III)," *J. Am. Chem. Soc.*, vol. 45, no. 9, pp. 2179–2181, Sep. 1923, doi: 10.1021/ja01662a023.
2. E. K. Nelson, "The constitution of capsaicin, the pungent principle of capsicum," *J. Am. Chem. Soc.*, vol. 41, no. 7, pp. 1115–1121, Jul. 1919, doi: 10.1021/ja02228a011.

3. P. Suresh and K. Srinivasan, "Tissue distribution & elimination of capsaicin, piperine & curcumin following oral intake in rats," *Indian J. Med. Res.*, vol. 131, pp. 682–691, May 2010.
4. M. Adiyasa and M. Meiyanti, "Pemanfaatan obat tradisional di Indonesia: Distribusi dan faktor demografis yang berpengaruh," *J. Biomedika dan Kesehat.*, vol. 4, no. 3, pp. 130–138, 2021, doi: 10.18051/JBiomedKes.2021.v4.130-138.
5. W. S. Kingery, "A critical review of controlled clinical trials for peripheral neuropathic pain and complex regional pain syndromes," *Pain*, vol. 73, no. 2, pp. 123–139, 1997, doi: [https://doi.org/10.1016/S0304-3959\(97\)00049-3](https://doi.org/10.1016/S0304-3959(97)00049-3).
6. P. Zis, A. Apsokardos, C. Isaia, P. Sykioti, and A. Vadalouca, "Posttraumatic and postsurgical neuropathic pain responsive to treatment with capsaicin 8% topical patch," *Pain Physician*, vol. 17, no. 2, pp. E213–E218, 2014.
7. J. Kiani, F. Sajedi, S. Nasrollahi, and F. Esna-Ashari, "A randomized clinical trial of efficacy and safety of the topical clonidine and capsaicin in the treatment of painful diabetic neuropathy," *J. Res. Med. Sci. Off. J. Isfahan Univ. Med. Sci.*, vol. 20, no. 4, pp. 359–363, 2015.
8. Q. Yu *et al.*, "Expression of TRPV1 in rabbits and consuming hot pepper affects its body weight," *Mol. Biol. Rep.*, vol. 39, no. 7, pp. 7583–7589, 2012, doi: 10.1007/s11033-012-1592-1.
9. L.-J. Yuan *et al.*, "Capsaicin-containing chili improved postprandial hyperglycemia, hyperinsulinemia, and fasting lipid disorders in women with gestational diabetes mellitus and lowered the incidence of large-for-gestational-age newborns," *Clin. Nutr.*, vol. 35, no. 2, pp. 388–393, 2016, doi: <https://doi.org/10.1016/j.clnu.2015.02.011>.
10. N. Harada and K. Okajima, "Effects of Capsaicin and Isoflavone on Blood Pressure and Serum Levels of Insulin-Like Growth Factor-I in Normotensive and Hypertensive Volunteers with Alopecia," *Biosci. Biotechnol. Biochem.*, vol. 73, no. 6, pp. 1456–1459, Jun. 2009, doi: 10.1271/bbb.80883.
11. P.-Y. Deng and Y.-J. Li, "Calcitonin gene-related peptide and hypertension," *Peptides*, vol. 26, no. 9, pp. 1676–1685, 2005, doi: <https://doi.org/10.1016/j.peptides.2005.02.002>.
12. M. J. Adams, K. D. K. Ahuja, and D. P. Geraghty, "Effect of capsaicin and dihydrocapsaicin on *in vitro* blood coagulation and platelet aggregation," *Thromb. Res.*, vol. 124, no. 6, pp. 721–723, Dec. 2009, doi: 10.1016/j.thromres.2009.05.001.
13. D. L. Bhatt and E. J. Topol, "Scientific and therapeutic advances in antiplatelet therapy," *Nat. Rev. Drug Discov.*, vol. 2, no. 1, pp. 15–28, 2003, doi: 10.1038/nrd985.
14. R. Stocker and J. F. Keaney, "Role of Oxidative Modifications in Atherosclerosis," *Physiol. Rev.*, vol. 84, no. 4, pp. 1381–1478, Oct. 2004, doi: 10.1152/physrev.00047.2003.
15. R. Zhang, I. Humphreys, R. P. Sahu, Y. Shi, and S. K. Srivastava, "In vitro and in vivo induction of apoptosis by capsaicin in pancreatic cancer cells is mediated through ROS generation and mitochondrial death pathway," *Apoptosis*, vol. 13, no. 12, pp. 1465–1478, 2008, doi: 10.1007/s10495-008-0278-6.
16. H. H. Andersen, C. Sand, and J. Elberling, "Considerable Variability in the Efficacy of 8% Capsaicin Topical Patches in the Treatment of Chronic Pruritus in 3 Patients with Notalgia Paresthetica," *Ann Dermatol*, vol. 28, no. 1, pp. 86–89, Feb. 2016, [Online]. Available: <https://doi.org/10.5021/ad.2016.28.1.86>.
17. L. Misery *et al.*, "Successful treatment of refractory neuropathic pruritus with capsaicin 8% patch: a bicentric retrospective study with long-term follow-up," *Acta Derm. Venereol.*, vol. 95, no. 7, pp. 864–865, 2015, doi: 10.2340/00015555-2085.
18. P. Holzer and W. Sametz, "Gastric mucosal protection against ulcerogenic factors in the rat mediated by capsaicin-sensitive afferent neurons," *Gastroenterology*, vol. 91, no. 4, pp. 975–981, 1986, doi: [https://doi.org/10.1016/0016-5085\(86\)90702-X](https://doi.org/10.1016/0016-5085(86)90702-X).

19. H. Foster and A. Lake, "Use of Vanilloids in Urologic Disorders," *Prog. Drug Res.*, vol. 68, pp. 307–317, Jun. 2014, doi: 10.1007/978-3-0348-0828-6-13.
20. M. de Sèze, L. Wiart, P.-A. Joseph, J.-P. Dosque, J.-M. Mazaux, and M. Barat, "Capsaicin and neurogenic detrusor hyperreflexia: A double-blind placebo-controlled study in 20 patients with spinal cord lesions," *Neurol. Urodyn.*, vol. 17, no. 5, pp. 513–523, Jan. 1998, doi: [https://doi.org/10.1002/\(SICI\)1520-6777\(1998\)17:5<513::AID-NAU7>3.0.CO;2-G](https://doi.org/10.1002/(SICI)1520-6777(1998)17:5<513::AID-NAU7>3.0.CO;2-G).
21. A. Harborne, *Phytochemical methods a guide to modern techniques of plant analysis*. Berlin, Heidelberg: Springer Dordrecht, 1998.
22. B. N. F. Meyer N R; Putnam, J E; Jacobsen, L B; Nichols, D E; McLaughlin, J L, "Brine Shrimp: A Convenient General Bioassay for Active Plant Constituents," *Planta Med.*, vol. 45, no. 05, pp. 31–34, 1982, doi: 10.1055/s-2007-971236.
23. E. Yuslianti, B. Bachtiar, D. Suniarti, and A. Sutjiatmo, "Standardisasi farmasitikal bahan alam menuju fitofarmaka untuk pengembangan obat tradisional Indonesia," *Dentika Dent. J.*, vol. 19, no. 2, pp. 179–185, 2016.
24. J. L. Carballo, Z. L. Hernández-Inda, P. Pérez, and M. D. García-Grávalos, "A comparison between two brine shrimp assays to detect in vitro cytotoxicity in marine natural products," *BMC Biotechnol.*, vol. 2, no. 1, p. 17, 2002, doi: 10.1186/1472-6750-2-17.
25. B. Thapa, N. Skalko-Basnet, A. Takano, K. Masuda, and P. Basnet, "High-Performance Liquid Chromatography Analysis of Capsaicin Content in 16 Capsicum Fruits from Nepal," *J. Med. Food.*, vol. 12, pp. 908–913, Sep. 2009, doi: 10.1089/jmf.2008.0187.
26. Z. Alothman, A. Badjah-Hadj-Ahmed, M. Habila, and A. Ghafar, "Determination of Capsaicin and Dihydrocapsaicin in Capsicum Fruit Samples using High Performance Liquid Chromatography," *Molecules*, vol. 16, pp. 8919–8929, Dec. 2011, doi: 10.3390/molecules16108919.

Open Access This chapter is licensed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits any noncommercial use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this chapter are included in the chapter's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the chapter's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.





Certificate

To certify that :
Kirana Anggraini

has contributed as

Presenter

in The 4th Borobudur International Symposium 2022
"The Innovation Chain: A Contribution to Society and Industry"

Rector
of Universitas Muhammadiyah Magelang



Dr. Lilik Andriyani, SE., M.Si

Chairman of 4th BIS 2022



Prof. Dr. Muji Setiyo, ST., MT.

Organized by : Co-host:



December 21st, 2022
Magelang, Central of Java, Indonesia





BIS 2022

The 4th Borobudur International Symposium
Online Virtual Conference, 21 December 2022

Website: <https://bis.unimma.ac.id/2022>

Email: bis@unimma.ac.id

Date: 21 June 2023

Letter of Acceptance for Full Paper

Dear Authors: Kirana Anggraini*, Dyah Ayu Woro Setyaningrum, Laviany Putri Shihran, Isra Fauziyyah, Laela Wulansari, Hening Tyas Andayani

We are pleased to inform you that your paper, entitled:

**"Effect of Different Solvent Types on Red Ginger Rhizome Extraction Process
(*Zingiber officinale* Roscoe var *rubrum*) on Gingerol and Shagol Levels"**

has been reviewed and accepted to be presented at BIS 2022 conference to be held on 21 December 2022 in Magelang, Indonesia.

Please make the payment for registration fee before the deadlines, visit our website for more information.

Thank You.

Best regards,



Prof. Dr. Muji Setiyo, S.T., M.T.
BIS 2022 Chairperson





BIS 2022

The 4th Borobudur International Symposium
Online Virtual Conference, 21 December 2022

Website: <https://bis.unimma.ac.id/2022>

Email: bis@unimma.ac.id

Date: 21 June 2023

Letter of Acceptance for Full Paper

Dear Authors: Kirana Anggraini*, Dyah Ayu Woro Setyaningrum, Lavianny Putri Shihran, Isra Fauziyyah, Laela Wulansari, Hening Tyas Andayani

We are pleased to inform you that your paper, entitled:

"The Effect of Different Types of Solvents in The Extraction Process of Javanese Chilies Fruits (*Piper retrofractum* Vahl) on Capsaicin Levels"

has been reviewed and accepted to be presented at BIS 2022 conference to be held on 21 December 2022 in Magelang, Indonesia.

Please make the payment for registration fee before the deadlines, visit our website for more information.

Thank You,

Best regards,

Prof. Dr. Muji Setiyo, S.T., M.T.
BIS 2022 Chairperson

