



**A Quantitative Survey of Antibiotic Use at a Hospital in Jambi Province Indonesia in Three-month Before and After Implementation of Antimicrobial Resistance Control Program by RASPRO Concept**

RONALD IRWANTO NATADIDJAJA<sup>1,2</sup>, RUDI ASMAJAYA<sup>2</sup>, HERMINA BASRIE<sup>2</sup>, HADI SUMARSONO<sup>2</sup>

<sup>1</sup>Department of Internal Medicine, Faculty of Medicine, Universitas Trisakti, Jakarta, 11440, Indonesia  
<sup>2</sup>Pelita RASPRO Indonesia Foundation, Jakarta, 11470, Indonesia



**INTRODUCTION**

Based on Decree of Minister of Health Number 8/2015 in article 11 concerning quality indicators of Antimicrobial Resistance Control Program (ARCP)/ Program Pengendalian Resistensi Antimikroba (PPRA) implementation in hospitals, it has been known that reduced quantity of antimicrobial use has become one of those indicators.


**OBJECTIVES**

This survey is a descriptive study using secondary data retrieved between July and September 2019 (3 months before implementation of RASPRO concept) as well as between October and December 2019 (3 months after the implementation), which was aimed to evaluate impacts on implementing *Regulasi Antimikroba Sistem Prospektif (RASPRO)* concept at a hospital in Jambi province, Indonesia.


**METHODS**

The survey was carried out by calculating the expenditure of 3 antibiotic classes, which were the most commonly used and usually given by injection in hospitals and Intensive Care Units (ICU), i.e. the beta-lactam, quinolones and carbapenem.

**RESULT**



We found reduced use of Ceftriaxone as many as 890 ampules (37.11%), for Cefotaxime the reduction was 580 ampules (67.13%); while the use of Cefoperazone reduced as many as 76 ampules (47.50%) and Cefazidime reduced as many as 10 ampules (7.14%). The use of Ciprofloxacin reduced as many as 327 ampules (71.40%), but there was a drastic increase in the use of Levofloxacin as many as 59 ampules (>100%). The use of Carbapenems increased, which included 79 ampules (34.20%) for Meropenem; while the use of Impenem increased as many as 9 ampules (100%).



In three months after the implementation of RASPRO concept, 92.58% prophylaxis antibiotic had been given for appropriate indication and the antibiotic use of Cefazolin 71.31%. Within three months before and after the implementation of RASPRO concept, there was a total reduction of antibiotic use, which reached 1736 ampules (40.57%).

**CONCLUSION**

The implementation of RASPRO concept can be executed as an effort to reduce the quantity of antimicrobial use in hospitals. However, larger studies and longer monitoring are required in order to identify the impact of implementation of RASPRO concepts at a hospital.

**Acknowledgement**

We thank Pelita RASPRO Indonesia Foundation that has facilitated the operation of our study and the hospital that has been willing to provide data for this study.

Natadidjaja RI, Heary T, Adiani H, Arsyani A, Bur B. Antibiotic usage at a private hospital in Central Java: results of implementing the Indonesian Regulation on the Prospective Antimicrobial System (Regulasi Antimikroba Sistem Prospektif Indonesia (RASPRO)). *Int J Infect Control* [Internet]. 2021;17(1):1-10. Available from: <https://doi.org/10.3396/ijcs17-2041>

Natadidjaja RI, Eliza Y, Sarayu YB, Matatala A, Sundariningrum BV. Decreasing the Broad Spectrum Antibiotics Use With the Prospective Antimicrobial Stewardship of RASPRO INDONESIA Model. *Journal of Antimicrobial Resistance and Infection Control* [Internet]. Geneva, Switzerland; 2019. p. 147-8. Available from: <https://arj.ijournal.biomedcentral.com/articles/10.1186/s13758-019-0050-7>

