

# **2021 IEEE International Conference on Health, Instrumentation & Measurement, and Natural Sciences (InHeNce 2021)**

**Medan, Indonesia  
14 – 16 July 2021**

**IEEE Catalog Number: CFP21AM3-POD  
ISBN: 978-1-6654-4182-7**



# Conference Program



2021

## **ORGANIZING COMMITTEE (OC)**

Advisory Council	:	Nyoman Ehrich Lister, Florenly, Chrismis Novalinda Ginting, Arjon Turnip, Endra Joelianto
General Chair	:	Refi Ikhtiari
Secretary1	:	Abdi Dharma
Secretary2	:	Santy Deasy Siregar
Finance Chair	:	Ermi Girsang
Technical Program Chair	:	Florenly
Session Panel Chair	:	Linda Chiuman
Publication Chair	:	Arjon Turnip

## **INTERNATIONAL PROGRAM COMMITTEE (IPC)**

José Aranguren	:	University of Madrid, Spain
Young Ho Kim	:	Chungnam National University
Wen-Tao Liu	:	Nanjing Medical University, China
Kuwat Triyana	:	Universitas Gadjah Mada, Indonesia
Hwee Ming Cheng	:	University of Malaya, Malaysia
Azizon Binti Othman	:	Sultanah Maliha Hospital, Langkawi, Malaysia
Adang Bachtiar	:	Universitas Indonesia, Indonesia
Muhammad Hadi	:	Universitas Muhammadiyah Jakarta, Indonesia
Ermita Isfandiary Ibrahim Ilyas	:	Universitas Indonesia, Indonesia
Gusbakti Rusip	:	Universitas Muhammadiyah Sumatera Utara, Indonesia
Rahmiana Zein	:	Universitas Andalas, Indonesia
Titania Tj. Nugroho	:	Universitas Riau, Indonesia
Nurul Taufiqu Rochman	:	Lembaga Ilmu Pengetahuan Indonesia, Indonesia
Wahyu Widowati	:	Universitas Kristen Maranatha, Indonesia
Arjon Turnip	:	Universitas Padjajaran, Indonesia
Endra Joelianto	:	Institut Teknologi Bandung Indonesia
Refi Ikhtiari	:	Universitas Prima Indonesia, Indonesia
Hairus Abdullah	:	National Taiwan University Science & Technology, Taiwan

Dessy Novita : Universitas Padjadjaran, Indonesia

Erwin Handoko : Universitas Prima Indonesia, Indonesia

Ihsan Iswaldi : Universitas Prasetiya Mulya, Indonesia

Masteria Yunovilsa Putra : Lembaga Ilmu Pengetahuan Indonesia, Indonesia

Ario Betha Juanssilfero : Lembaga Ilmu Pengetahuan Indonesia, Indonesia

Rudy Syahputra : Universitas Islam Indonesia, Indonesia

Fotarisman Zalukhu : Dinas Kesehatan Kota Gunungsitoli, Indonesia

Haripin Togap Sinaga : Poltekkes Medan GIzi Lubuk Pakam, Indonesia

Hartono : IBBI Medan, Indonesia

Edmi Edison : Universitas Muhammadiyah Prof. Dr. HAMKA, Indonesia

Albert Hutapea : Universitas Advent Indonesia, Indonesia

Suyitno : Universitas Sebelas Maret, Indonesia

Sutikno : Universitas Negeri Semarang, Indonesia

Sovia Lenny : Universitas Sumatera Utara, Indonesia

Lee Yun Li : Sunway University, Malaysia

Lakshmi Selvaratnam : Monash University Malaysia

Narendra Pamidi : Monash University Malaysia

Monica Dwi Hartanti : Universitas Trisakti, Indonesia

Yenny : Universitas Trisakti, Indonesia

Dian Ratih Laksmiawati : Universitas Pancasila, Indonesia

Patricia B Liman : Universitas Trisakti, Indonesia

Ratheesh Kumar Meleppat : University of California Davis, USA

**2021 International Conference on Health, Instrumentation & Measurement,  
And Natural Science (InHeNce)**

**Table of Contents**

S. No	Paper ID	Paper Title	Page No
1	1570713128	Phytochemical and Hypoglycemia Effect Test Using Extract of Barangan Banana Peel with OGTT in Male White Rats Induced with Sucrose	1
2	1570713154	Isolation and Molecular Identification of Inulinase Producing Thermophilic Bacteria from Geothermal Hot Spring	6
3	1570713196	Analysis of Mosquito Genetic with PCR-RAPD Approach	12
4	1570715206	Aloe Vera Protective Effect on Lipopolysaccharide- Induced RAW 264.7 Inflamed Cells	18
5	1570715694	The Effectiveness of Red Spinach Extract Ointment ( <i>Amaranthus Tricolor L.</i> ) Against Healing Second Degree Burns in Wistar Rat	24
6	1570716179	Antioxidant Potency of <i>Kaempferia Galanga</i> Linn and <i>Zingiber Officinale</i> Var. <i>Rubra</i> Rhizomes	30
7	1570719716	Effect of Andaliman Fruit Extract on Cervical Cancer Rat's Histology	36
8	1570720267	The Effect of Extract Ethanol <i>Rosa Damascena</i> to the Interleukin-6 Level of Obese Rats	41
9	1570720314	Improvement of Liver Function from Lemon Pepper Fruit Ethanol Extract in Streptozotocin-Induced Wistar Rats	46
10	1570720672	Effectivity of Nano Herbal Andaliman ( <i>Zanthoxylum Acanthopodium</i> ) to the Vascular Endothelial Growth Factor (VEGF) Expression in Burn Wound in Diabetic Rats	51
11	1570720816	Progress of Zn(O, S) Based Nanoparticles for Hydrogen Evolution Reaction and Its Application for Hydrogenation Reaction	56
12	1570720917	Synthesis of Photocatalyst Fe <sub>3</sub> O <sub>4</sub> -CuO/Activated Carbon for Removal of Aniline in Water	62

13	1570722523	Liver Protection Activity of Sunkist Orange ( <i>Citrus Sinensis</i> L. Osbeck) Peels in Cyclophosphamide Induced Male Wistar Rat	67
14	1570724089	Relationship Between Cervical Vertebrae Maturity Levels with Mandibular Growth at the Age of 8-20 Years in Batak Tribe	72
15	1570712960	Antioxidant Activity of Green Tea Extract and Myricetin	77
16	1570724959	Potential Polymorphism of BMP-2 rs235768 in North Sumatra Sub-Population with Mandibular Asymmetry	82
17	1570726833	Scheduling for Healthcare Centre for COVID-19: Deep Learning and Genetic Algorithmic Approach	87
18	1570727467	Potential of Human Wharton's Jelly Mesenchymal Stem Cells (hWJMSCs) Secretome for COVID-19 Adjuvant Therapy Candidate	92
19	1570729413	Animals and Technology in Space: A Perspective from Aerospace Engineering to Veterinary Medicine	98
20	1570729583	Stronger Antibacterial Efficacy of Red Beetroots Compared to Red Dragon Fruit Peels Extract on <i>Streptococcus Mutans</i>	104
21	1570729769	HIPAA-Based Analysis on the Awareness Level of Medical Personnel in Indonesia to Secure Electronic Protected Health Information (ePHI)	109
22	1570729868	Measurement of Gas Bubbles Distribution on Electroflotation Process Using Titanium and Stainless Steel Electrode with DinoCapture 2.0	115
23	1570730336	Artificial Intelligence in Medicine: A Review of Challenges in Implementation and Disparity	120
24	1570730698	Antioxidant, Total Phenol, Total Flavonoid, and LC-MS/MS Analysis of <i>Pometia Pinnata</i> Ethanol Extract	126
25	1570731907	A Promising Anti-Inflammatory Drugs from <i>Citrus Amblycarpa</i> (Hassk.) Ochse Seeds	131
26	1570731909	Extracted <i>Passiflora Edulis</i> Pulp to Reduce Inflammation in LPS-Activated Macrophage Cell Line: RAW 264.7	137
27	1570731935	Analysis of Bubbles Size Produced in Electroflotation Using Graphite and Stainless Steel Electrode with DinoCapture 2.0	142

28	1570732244	Development of a Heart Wave Transmitter Between Android Devices Using iLBC Coding over the Internet	147
29	1570732245	Anti-Aging Effectiveness of Red Spinach Extract Ointment ( <i>Amaranthus Tricolor</i> L.) Against Collagen, Elasticity, Hydration, Sebum, and Pigment Levels in Wistar Rats	151
30	1570732683	Potential of <i>Gnetum Gnemon</i> L. Seed Extract Against Insulin Levels and PDX-1 Expression in Diabetes Mellitus Rats Model	157
31	1570732695	Antioxidant Properties of Soybean ( <i>Glycine Max</i> L.) Extract and Isoflavone	162
32	1570732818	<i>Syzygium Polyanthum</i> Ethanol Extract Ameliorates Benzene-Induced Nephrotoxicity in Rats	168
33	1570732913	Information Systems of School Financial Management with Digital Signature Recognition Using MobileNet Algorithm	174
34	1570732925	The Protective Effect of Ethanolic Extract of <i>Syzygium Polyanthum</i> on the Pancreas of Benzene-Induced Rats	180
35	1570733008	Hematological Profiles of Benzene-Induced Rats Treated with Ethanolic Extract of <i>Syzygium Polyanthum</i>	186
36	1570733310	Role of Platelet-Rich Plasma to Sperm Quality in Male Partners Undergoing Infertility Treatment	192
37	1570733384	Antioxidant Activity, Total Phenol, and Total Flavonoid of <i>Syzygium Polyanthum</i>	197
38	1570733494	Anti-Hyperuricemia of Avocado Leaves Ethanol Extract in Potassium Oxonate Induced-Rats	202
39	1570733677	Evaluation of Honey Effectivity on Burned-Wound Contraction in <i>Rattus Norvegicus</i>	207
40	1570733739	The Hepatoprotective Effect of White Turmeric Ethanol Extract on Cyclophosphamide-Induced Male Wistar Rats	212
41	1570733846	Identification of Chemical Compounds to Predict in Silico Toxicity Using <i>Syzygium Polyanthum</i> Ethanol Extract	217
42	1570733976	The Impact of Red Dragon Fruit ( <i>Hylocereus Polyrhizus</i> ) Consumption on Improving Cell Function in Mitochondria in Delaying Fatigue During Strenuous Exercise	221

43	1570734017	Ethanollic Extract of Syzygium Polyanthum Effect in Reduce Biomarkers Levels to Prevent Rats Liver Damage	226
44	1570734019	Reduction of Tropinin Levels in Rats Through Cardioprotective Activity of Syzygium Polyanthum Ethanol Extract	231
45	1570735239	Papaya Latex for Healing the Second Degree of Burn Wound in Male Mice	236
46	1570735240	The Effect of Administration Fig Leaf Tea to Reduce Blood Glucose Levels in Diabetes Mellitus Patients	242
47	1570735427	Pancreas Protective Effect Ethanol Extract of Curcuma on Rats Induced Doxorubicin	247
48	1570735482	Neuroprotective Effect of Syzygium Polyanthum Extract on TNF Alpha and Dopamine Levels in Benzene-Induced Rats	252
49	1570735756	Anti-Diabetic Activity of the Rose Petal Methanolic Extract in Alloxan-Diabetic Rats	257
50	1570735759	Exploration of Analgesic-Antipyretic Effect from Lemon Peppers Methanol Extract	262
51	1570735763	Anti-Dyslipidemia Activity from Lemon Pepper in PTU and High Fat Diet Induced Rats	267
52	1570735868	Association Between Body Mass Index (BMI) and DNA Fragmentation Index	273
53	1570736227	Microencapsulation of Lemongrass Leaves Effect on Reactive Oxygen Species (ROS) Fibroblasts	279
54	1570736234	Synthesis of Silver Nanoparticles from Lemongrass Leaves Induced Wound Healing by Reduction ROS Fibroblasts	284
55	1570736339	Effectiveness of Reducing Lead Levels on Feather Shells with Various Types of Filtrates	289
56	1570736358	Antibiofilm Activity of Parabiotic Reuterin on Acrylic Resin Plates	294
57	1570736359	Goji Berry Extract (Lycium Barbarum L.) Efficacy on Oral Pathogen Biofilms	300
58	1570736360	A Comparison of Commercial Mouthwashes Towards Oral Pathogens Biofilms in Vitro	306
59	1570736386	Association Rules Analysis Using Algorithm Apriori and Fuzzy Normalization	311
60	1570736389	Adaboost Hybrid Algorithm with Color Space Transformation and Lighting Compensation Method to Detect Face Based on Emotion	316



61	1570736738	Effects of Chili Sauce on the Absorbency and Diametral Tensile Strength of Nanocomposite	322
62	1570736785	Effect of Moringa Oleifera Leaves on Human Blood Coagulation Process	327
63	1570737020	Increase of Fibroblast Proliferation by Composite Membrane (Polyvinyl Alcohol - Collagen - Hydroxyapatite)	332
64	1570737125	Smart Container Design for Transporting Virus Sample in Remote Areas	337
65	1570737580	Performance of Spectral, Autocorrelation and Peak Count Based PR Estimation Methods Under Normal/Abnormal PPG for Wearable Devices	343
66	1570737907	Nephroprotective Effect from Rose Petals ( <i>Rosa Damascena</i> ) Extract Against Cadmium Toxicity	349
67	1570737933	In Silico Automatic Annotation of Phenolamides in Plants by Tandem Mass Spectra	355
68	1570738172	Analysis Accuracy of Complementary Label for Multi-Class Classification: Case Study on Chest X-Ray Images	360
69	1570738296	Demographic Factors and Total Muscle Mass are Associated with Handgrip Strength in Selected Indonesian Adults	365
70	1570738606	The Effectiveness of Pelvic Rocking and Birthing Ball Exercise on Labor Pain During First Stage of Active Phase	370
71	1570738687	Phaleria Macrocarpa Crude Extract as Antidote for Cd (II) Contamination in Kidney Organs	374
72	1570740335	Phytochemical and Antibacterial Analysis of Mangkokan Leaf Extract Against <i>Salmonella Typhimurium</i> Bacteria	379
73	1570741503	Lightweight Compressed Sensing (CS) and Partial DCT Based Compression Schemes for Energy-Efficient Wearable PPG Monitoring Devices	384
74	1570741572	Coexistence of Hemoglobin E Beta Thalassemia and Southeast Asian Ovalocytosis & RBC Changes Conferring Malarial Protection	390
75	1570741948	Performance of True Transfer Learning Using CNN DenseNet121 for COVID-19 Detection from Chest X-Ray Image	395

76	1570760328	Wound Healing Activity of Avocado Peel Ointment Against Second Degree Burn Wound	400
77	1570760403	Antihyperlipidemia Activity of Syzygium Polyanthum Ethanol Extract on Benzene Induced Rats	406
78	1570760973	Sentiment Analysis of Public Reaction to COVID19 in Twitter Media Using Naïve Bayes Classifier	411
79	1570760979	Adaptive Land Management for Climate-Smart Agriculture	415
80	1570760984	Automatic Vertebra Fracture & Classification using Dual Energy X-Ray Absorptiometry	422
81	1570761237	Development of Medical Robot Covid-19 Based 2D Mapping LIDAR and IMU Sensors	426
82	1570762388	Classification of Brain Signal Activities Before and After Consuming Soy Peptide Using Fuzzy Logic	430

### Schedule at Glance

Time	Activities	Description
<b>Wednesday, July 14<sup>th</sup></b>		
08:00-10:00 WIB	Committee preparation	Organizing Committee
10:00-15:00 WIB	Registration Pre conference	Registration Form for Participants from UNPRI <a href="https://bit.ly/inhenceforunpri">bit.ly/inhenceforunpri</a>  Registration Form for Non-UNPRI Participants <a href="https://bit.ly/inhencenonunpri">https://bit.ly/inhencenonunpri</a>  Registration Form for Presenter <a href="https://bit.ly/inhencepresenter">https://bit.ly/inhencepresenter</a>  Registration Form for Speaker <a href="https://bit.ly/inhencespeaker">https://bit.ly/inhencespeaker</a>
<b>Thursday, July 15<sup>th</sup></b>		
<b>08:00 - 08:05 WIB</b>	<b>Opening Ceremony</b>	<b>Master of Ceremony</b> (Herbert Wau M.P.H dan Risya Yoela Sinaga)
08:05 - 08:10 WIB	National Anthem	Indonesia Raya
08:10 - 08:15 WIB	Welcoming Message	General Chair of InHeNce 2021 <b>Refi Ikhtiari, Ph.D.</b>
08:15 - 08:20 WIB	Opening Remarks	IEEE Indonesia Section IMS/ITS Joint Chapters <b>Endra Joeliando Ph.D.</b>
08:20 - 08:30 WIB	Opening Remarks	Rector of Universitas Prima Indonesia <b>Prof. Dr. Chrismis Novalinda Ginting, M.Kes., AIFO.</b>
<b>08:30-12:00 WIB</b>	<b>Speaker session</b>	<b>Moderator (Frans Judea Samosir, M.P.H.)</b>
<b>08:30 - 09:10 WIB</b> (09:30 - 10:10 Nanjing Time)	<b>Keynote Speaker 1</b>	<b>Prof. Wen-Tao Liu, Ph.D.</b> Institute of Translational Medicine, Nanjing Medical University, China
09:10 - 09:20 WIB	Question and Answer	Moderator (Frans Judea Samosir, M.P.H.)
<b>09:20 - 10:00 WIB</b>	<b>Keynote Speaker 2</b>	<b>Prof. Dr. Eng. Kuwat Triyana, M.Si.</b> Inventor of GeNoSe Universitas Gajah Mada, Indonesia
10:00 - 10:10 WIB	Question and Answer	Moderator (Frans Judea Samosir, M.P.H.)
<b>10:10 WIB</b>	<b>Photo Session (Master of Ceremony)</b>	
10:10 - 10:20 WIB	Transition from Main Room to Break out Room	
<b>10:20 - 12:00 WIB</b>	<b>Parallel Session</b>	<b>Room 1 -14</b>

<b>Track: Instrumentation and Measurement (20 papers)</b> <b>2 Room Parallel Session</b>		
<b>Room 1</b>	<b>Invited Talk 1 (10:20—10.50 WIB)</b>  Question and Answer (10.50-11.00 WIB)  Oral Presentation (11:00-12:00 WIB)	<b>Endra Joelianto, Ph.D.</b> SMIEEE, Chair of IEEE IS IMS/ITS Join Chapter Institut Teknologi Bandung, Indonesia  Session Chair  4 Papers (@15 minutes)
<b>Room 2</b>	<b>Invited Talk 2 (10:20—10.50 WIB)</b>  Question and Answer (10.50-11.00 WIB)  Oral Presentation (11:00-12:00 WIB)	<b>Arjon Turnip, Ph.D.</b> Chair of IEEE IS CSS/RAS Joint Chapter Department of Electrical Engineering, Universitas Padjajaran, Indonesia  Session Chair  4 Papers (@15 minutes)
<b>Track: Natural Sciences (65 papers)</b> <b>7 Room Parallel Session</b>		
<b>Room 3</b>	<b>Invited Talk 3 (10:20—10.50 WIB)</b>  Question and Answer (10.50-11.00 WIB)  Oral Presentation (11:00-12:00 WIB)	<b>Prof. Dr. Nurul Taufiqu Rochman, M.Eng.</b> Professor of Nanomaterials LIPI&CEO Nano Centre Indonesia Recipient of Habibie Technology Award 2014  Session Chair  4 Papers (@15 minutes)
<b>Room 4</b>	<b>Invited Talk 4 (10:20—10.50 WIB)</b>  Question and Answer (10.50-11.00 WIB)  Oral Presentation (11:00-12:00 WIB)	<b>Prof. Dr. Titania Tj. Nugroho, M.Si.</b> Professor of Biochemistry, FMIPA Universitas Riau, Indonesia  Session Chair  4 Papers (@15 minutes)
<b>Room 5</b>	<b>Invited Talk 5 (10:20—10.50 WIB)</b>  Question and Answer (10.50-11.00 WIB)  Oral Presentation (11:00-12:00 WIB)	<b>Dr. Wahyu Widowati, M.Si.</b> Universitas Kristen Maranatha, Indonesia President of PT Aretha Medika Utama  Session Chair  4 Papers (@15 minutes)
<b>Room 6</b>	<b>No Invited Talk</b>  Oral Presentation (10:20-12:00 WIB)	<b>Host</b>  6 Papers (@15 minutes)
<b>Room 7</b>	<b>No Invited Talk</b>  Oral Presentation (10:20-12:00 WIB)	<b>Host</b>  6 Papers (@15 minutes)

<b>Room 8</b>	<b>No Invited Talk</b> Oral Presentation (10:20-12:00 WIB)	<b>Host</b> 6 Papers (@15 minutes)
<b>Room 9</b>	<b>Invited Talk 9 (10:20—10.50 WIB)</b>  Question and Answer (10.50-11.00 WIB)  Oral Presentation (11:00-12:00 WIB)	<b>Prof. Dr. Gusbakti Rusip, M.Sc., P.K.K., AIFM.</b> Chairman of Indonesia Physiological Society of North Sumatera Universitas Muhammadiyah Sumatera Utara, Indonesia  Session Chair  4 Papers (@15 minutes)
<b>Track: Health (53 papers) 5 Room parallel session</b>		
<b>Room 10</b>	<b>Invited Talk 10 (10:20—10.50 WIB)</b> 11:20 - 12:50 Malaysia Time  Question and Answer (10.50-11.00 WIB)  Oral Presentation (11:00-12:00 WIB)	<b>Prof. Dr. Hwee Ming Cheng</b> University of Malaya, Malaysia  Session Chair  4 Papers (@15 minutes)
<b>Room 11</b>	<b>Invited Talk 11 (10:20—10.50 WIB)</b>  Question and Answer (10.50-11.00 WIB)  Oral Presentation (11:00-12:00 WIB)	<b>Adang Bachtiar, dr. MPH., D.Sc</b> Universitas Indonesia, Indonesia  Session Chair  4 Papers (@15 minutes)
<b>Room 12</b>	<b>No Invited Talk</b> Oral Presentation (10:20-12:00 WIB)	<b>Host</b> 6 Papers (@15 minutes)
<b>Room 13</b>	<b>Invited Talk 13 (10:20—10.50 WIB)</b> 11:20 - 12:50 Malaysia Time  Question and Answer (10.50-11.00 WIB)  Oral Presentation (11:00-12:00 WIB)	<b>Dato' Dr. Azizon Binti Othman,</b> Sultanah Maliha Hospital, Langkawi, Malaysia  Session Chair  4 Papers (@15 minutes)
<b>Room 14</b>	<b>Invited Talk 14 (10:20—10.50 WIB)</b>  Question and Answer (10.50-11.00 WIB)  Oral Presentation (11:00-12:00 WIB)	<b>Dr. Muhammad Hadi, M.Kep.</b> Chairman of Association of Indonesia Nurse Education Centre (AINEC) Universitas Muhammadiyah Jakarta, Indonesia  Session Chair  4 Papers (@15 minutes)
<b>12:00 - 13:00 WIB</b>	<b>Lunch Break</b>	<b>Master of Ceremony</b> (Herbert Wau M.P.H dan Risya Yoela Sinaga)

<b>13:00 - 13:40 WIB</b> (08:00-08:40 Spain Time)	<b>Keynote Speaker 3</b>	<b>Prof. José Aranguren, D.D.S., M.S.</b> Professor in Endodontics, Rey Juan Carlos University of Madrid, Spain
13:40 - 13:50 WIB	Question and Answer	Moderator (Frans Judea Samosir, M.P.H.)
13:50 - 14:00 WIB	Transition from Main Room to Break out Room	
<b>Continued Parallel Session</b> <b>(14:00-16:00 WIB)</b>	<b>Track Instrumentation and Measurements (20 papers)</b> <b>2 Room Parallel Session</b>	
<b>Room 1</b>	Oral Presentation (14:00-16:00 WIB)	6 Papers (@15 minutes)
<b>Room 2</b>	Oral Presentation (14:00-16:00 WIB)	6 Papers (@15 minutes)
<b>Continued Parallel Session</b>	<b>Track: Natural Sciences (65 papers)</b> <b>7 Room Parallel Session</b>	
<b>Room 3</b>	Oral Presentation (14:00-16:00 WIB)	6 Papers (@15 minutes)
<b>Room 4</b>	Oral Presentation (14:00-16:00)	6 Papers (@15 minutes)
<b>Room 5</b>	Oral Presentation (14:00-16:00 WIB)	6 Papers (@15 minutes)
<b>Room 6</b>	Oral Presentation (14:00-16:00 WIB)	4 Papers (@15 minutes)
<b>Room 7</b>	Oral Presentation (14:00-16:00 WIB)	4 Papers (@15 minutes)
<b>Room 8</b>	Oral Presentation (14:00-16:00 WIB)	4 Papers (@15 minutes)
<b>Room 9</b>	Oral Presentation (14:00-16:00 WIB)	6 Papers (@15 minutes)
<b>Continued Parallel Session</b>	<b>Track: Health (53 papers)</b> <b>5 Room parallel session</b>	
<b>Room 10</b>	Oral Presentation (14:00-16:00 WIB)	6 Papers (@15 minutes)
<b>Room 11</b>	Oral Presentation (14:00-16:00 WIB)	6 Papers (@15 minutes)
<b>Room 12</b>	Oral Presentation (14:00-16:00 WIB)	4 Papers (@15 minutes)
<b>Room 13</b>	Oral Presentation (14:00-16:00 WIB)	6 Papers (@15 minutes)
<b>Room 14</b>	Oral Presentation (14:00-16:00 WIB)	6 Papers (@15 minutes)
<b>16:00 WIB</b>	<b>Time off</b>	Master of Ceremony (Herbert Wau M.P.H. and Risya Yoela Sinaga)
<b>Friday, July 16<sup>th</sup></b>		
<b>08:00 - 08:05 WIB</b>	<b>Opening Day-2</b>	<b>Master of Ceremony</b> (Herbert Wau M.P.H and Risya Yoela Sinaga) Moderator Frans Judea
<b>08:05 - 08:45 WIB</b> (10:05 - 10:45 Korean Time)	<b>Keynote Speaker 4</b>	<b>Prof. Young Ho Kim, Ph.D.</b> Chungnam National University, Korea

08:45 - 08:55 WIB	Question and Answer	Moderator (Frans Judea Samosir, M.P.H.)
08:55 - 09:00 WIB	<b>Announcement of Best Presenter Awards (Certificate and Photo Session)</b>	Master of Ceremony (Herbert Wau M.P.H and Risya Yoela Sinaga)
<b>09:00 - 09:05 WIB</b>	<b>Closing Remark</b>	Master of Ceremony (Herbert Wau M.P.H and Risya Yoela Sinaga) <b>IEEE Indonesia Section CSS/RAS Chapters Arjon Turnip, Ph.D.</b>



# KEYNOTE SPEAKERS

 **INTERNATIONAL  
CONFERENCE**

*ON HEALTH, INSTRUMENTATION & MEASUREMENT, AND NATURAL SCIENCES*

**MEDAN, JULY 14-16<sup>TH</sup> 2021**



## KEYNOTE SPEAKERS



**Prof. José Aranguren, DDS, MS**  
Professor in Endodontics, Rey Juan  
Carlos University of Madrid, Spain

**Prof. Young Ho Kim Ph.D**  
Professor, Dept. of Natural Product Chemistry,  
College of Pharmacy, Chungnam National  
University, Korea.



**Prof. Dr. Wen-Tao Liu**  
Professor, Vice President of Institute of  
Translational Medicine, Nanjing Medical  
University, China.

**Prof. Dr. Eng. Kuwat Triyana, M.Si.**  
Professor in Physics of Material & Instrumentation, Dept  
of Physics, Universitas Gadjah Mada, Indonesia (Inventor  
of GeNose®, a COVID-19 Quick Detection).



# Goji Berry Extract (*Lycium barbarum* L.) Efficacy on Oral Pathogen Biofilms

Sheila Soesanto  
Pharmacology Department Faculty of  
Dentistry  
Universitas Trisakti  
Jakarta, Indonesia  
mass9977@yahoo.com

Ricky Chandra Jaya Soen  
Undergraduate Program Faculty of  
Dentistry  
Universitas Trisakti  
Jakarta, Indonesia  
ricksoen@gmail.com

Reynalda Oktaviani  
Undergraduate Program Faculty of  
Dentistry  
Universitas Trisakti  
Jakarta, Indonesia  
reynaldaoktaviani@gmail.com

Armelia Sari Widyarman\*  
Microbiology Department Faculty of  
Dentistry, Universitas Trisakti  
Jakarta, Indonesia  
armeliasari@trisakti.ac.id

**Abstract**— *Lycium barbarum* L. fruit, which contains flavonoids and phenolic acids, has antibacterial properties that are expected to inhibit bacterial growth. The objective of this study is to determine the antibacterial and the antibiofilm effects of *L. barbarum* L. fruit ethanol extract towards *S. mutans* and *P. gingivalis*. An in-vitro laboratory experiment was performed with a post-test control group design. The extract of *L. barbarum* L. fruit was obtained by maceration technique using 96% ethanol as a solvent. The antibacterial assay was performed by microdilution and plate count methods. The antibiofilm effect was performed using a biofilm-assay method. The results of microdilution and plate count methods showed that the most effective antibacterial concentration against *S. mutans* and *P. gingivalis* was 100 µg/mL when compared with negative control ( $p < 0.05$ ). In the biofilm assay, the most effective concentration against *S. mutans* was 100 µg/mL at the 3-hours incubation time, while for *P. gingivalis*, the most effective concentration was 100 µg/mL at 24-hours incubation time when compared with negative control ( $p < 0.05$ ). These results indicate that ethanol extract of *L. barbarum* L. fruit was demonstrated to have antibacterial and antibiofilm effects against oral pathogens *S. mutans* and *P. gingivalis* with 87.15% and 97.73% of biofilms reduction respectively.

**Keywords**— antibacterial, antibiofilm, *Lycium barbarum* L., *Porphyromonas gingivalis*, *Streptococcus mutans*.

## I. INTRODUCTION

Oral health positively affects the appearance, physical, mental, and interpersonal well-being, of an individual. Oral health is a part of overall health, contributes to the quality of life [1]. National Basic Health Research Data (Riskesdas 2018) showed dental and oral health problems in 57.6% of the Indonesian population. The prevalence of dental caries in Indonesia in 2018 was 88.8%, with an average DMF-T index of 7.1, which is a very high severity of dental caries. Moreover, 74.1% of the Indonesian population experienced periodontitis [2].

Caries are the process of the demineralization of inorganic material and the dissolution of organic material, leading to bacterial invasion through the dentin layer until it reaches the pulp [3,4]. The process of dental caries depends on the presence of fermentable sugars (substrates), the type of tooth and saliva (host), cariogenic microbial flora (biofilm), and time [5]. Periodontitis is a disease caused by microorganisms that cause inflammation of tooth-supporting tissue and causes progressive destruction of periodontal ligament and alveolar bone. The sign of periodontitis is the

formation of pockets, recessions, or both [6]. Periodontitis in adults caused by numerous local factors, such as biofilms or calculus, is classified as chronic periodontitis [7].

The formation of biofilms begins when microorganisms in the planktonic state merge into bacterial colonies and wrap themselves in a self-produced extracellular polymer matrix [8]. In the initial phase of biofilm formation, there is an increase in Gram-positive cocci activity, one of which is *Streptococcus mutans*, which is able to attach to tooth surface through the formation of extracellular polysaccharides that cause biofilm matrix to have a gelatin-like consistency that facilitates attachment of bacteria to the tooth surface [4,9,10]. *Porphyromonas gingivalis*, which is a secondary bacterium, is an anaerobic Gram-negative bacterium found in periodontal pockets that causes chronic periodontitis. Various virulence factors of *P. gingivalis* such as gingipains, fimbriae, and lipopolysaccharides, play important role in periodontal disease progression and induce dysbiosis in biofilms [11,12].

Chlorhexidine mouthwash is used to prevent caries and treat periodontitis and is considered as the gold standard for controlling dental plaque and gingivitis due to its efficacy against a wide variety of microorganisms. However, chlorhexidine has various side effects, including taste disturbances, discoloration of teeth and mucosa, mucosal desquamation, salivary stone formation, irritation, dry oral cavity, and allergic reactions, such as contact stomatitis. The World Health Organization (WHO) recommends finding new natural ingredients to overcome the side effects of chemical agents [13,14].

The use of natural ingredients as antimicrobial agents has become an alternative because of their low cost and lower toxicity [15]. According to WHO, traditional medicine has been used globally and can be a major source of health for millions of people and sometimes the only source of care and is also culturally acceptable, affordable, and trusted by community [16]. Goji berry (*Lycium barbarum* L.) has been widely used as a traditional medicine by people in Asia, especially in the northwestern part of China, for more than 2000 years. Recently, *L. barbarum* L. has been gaining popularity as a highly nutritious food used to improve health in North America, Europe, and Asia [17]. *Lycium barbarum* L. has a red, oblong fruit with a length of 6–20 mm and a diameter of 3–10 mm. *Lycium barbarum* L. fruit is harvested when it is ripe and is then dried for later use [18]. The fruit, roots, tree bark, and flowers of *L. barbarum* L. are used as medicine [19].

The polysaccharides of *L. barbarum* L. exhibited properties that improve eye health and reproductive system; reduce fat and blood sugar; regulate immunity. It's also anticancer, anti-tumor, antioxidant, anti-fatigue, antiviral, anti-aging, hepatoprotective, neuroprotective, and cardioprotective properties [17,20]. The flavonoids and phenolic acids of *L. barbarum* L. have potential as antioxidants and antimicrobials [21].

*Lycium barbarum* L. fruit is effective against Gram-negative bacteria (e.g. *Escherichia coli*) and Gram-positive bacteria (e.g. *Staphylococcus aureus*)[22]. However, there have been no studies regarding the antibacterial effect of *L. barbarum* L. fruit against *S. mutans* and *P. gingivalis* as causing bacteria of dental caries and chronic periodontitis. Thus, this is the first study that analyzed the antibacterial and antibiofilm efficacy of goji berry (*L. barbarum* L.) ethanol extract towards *S. mutans* and *P. gingivalis* as oral pathogens.

## II. MATERIALS AND METHODS

### A. Ethanol extract production from *L. barbarum* L. fruit

Dried *L. barbarum* L. fruit (100 g) from Chinese medicine store "Lancar Jaya" at Teluk Gong Raya No. 43, Jakarta Utara (produced in Zhongning, Ningxia, China) was ground in a blender until it became powder. It was then immersed in 96% ethanol with a ratio of 1:8 for 72 hours, stirring every 15 minutes. Furthermore, filtration was performed using Whatman No. 1 filter paper and evaporated with a rotary evaporator at 40°C temperature, 60 rpm speed, and 20 atm pressure so a thick and solvent-free extract was obtained with a concentration of 100 µg/mL. Furthermore, extracts were diluted using sterile distilled water until concentrations of 50, 25, 12.5, and 6.25 µg/mL were obtained.

### B. Phytochemical Assay

Phytochemical assays were performed qualitatively to determine whether the ethanol extract of *L. barbarum* L. fruit contained flavonoids, phenols, quinones, steroids, terpenoids, and alkaloids. The qualitative results are expressed as (+) for the presence and (-) for the absence of phytochemicals.

### C. Bacterial Cultures

*Streptococcus mutans* ATCC 25175 and *Porphyromonas gingivalis* ATCC 33277 from Microbiology Center of Research and Education (MiCORE) Laboratory, Faculty of Dentistry, Trisakti University, were cultured on brain heart infusion (BHI) (Oxoid, Hampshire) broth medium and incubated at 37°C for 24 hours in anaerobic atmosphere. Furthermore, the absorbance measurements were performed to obtain the McFarland standard of  $0.5 = 1.5 \times 10^8$  CFU/mL (OD<sub>600</sub> = 0.132).

### D. Microdilution

Each well of a 96-well plate was distributed 100 µL of either an *S. mutans* or *P. gingivalis* culture. Subsequently, 100 µL of the following solutions was used as a treatment: ethanol extracts of *L. barbarum* L. fruit at 100, 50, 25, 12.5, and 6.25 µg/mL concentrations, 0.2% chlorhexidine gluconate as a positive control, and sterile distilled water as a negative control. The measurement of bacterial cell density was performed using microplate reader at 600 nm wavelength before and after the 96-well plates were

incubated for 24 hours. All treatments were done in triplicate.

### E. Total Plate Count

Bacterial growth was measured by re-diluting contents in 96-well plates for 10,000 times and cultured on BHI agar medium and incubated for 24 hours at 37°C.

### F. Biofilm Assay

Bacterial culture (200 µL) was dispensed into each 96-well plate and incubated at 37°C for 48 hours in an anaerobic atmosphere. Furthermore, the supernatant was removed until a thin layer of biofilm was left on the bottom surface of the well. Then, wells were rinsed with a solution of phosphate-buffered saline (PBS). The ethanol extracts of *L. barbarum* L. fruit at a concentration of 100, 50, 25, 12.5, and 6.25 µg/mL, 0.2% chlorhexidine gluconate as positive control, and sterile distilled water as negative control were added 200 µL to wells using a micropipette and incubated at 37°C for 1, 3, and 24 h in an anaerobic atmosphere. The well was rinsed twice using PBS and then fixated over a flame. Crystal violet dye (200 µL; 0.05% w/v) was added to each well and left for 15 minutes. The well was rinsed twice using PBS and left for 15 minutes. Then, 200 µL of 96% ethanol was inserted, and optical density was measured using a microplate reader (SAFAS MP96, Monaco) at 595 nm wavelength.

### G. Statistical Analysis

Data were processed by using Statistical Product and Service Solution (SPSS) software version 25.0 and the normality test was performed by using the Shapiro-Wilk method. Normally distributed data ( $p > 0.05$ ) was analyzed by one-way analysis of variance (ANOVA) test. Significant data ( $p < 0.05$ ) were analyzed with a posthoc test using Tukey's test with a significance level of  $P < 0.05$  to determine which groups were significantly different.

## III. RESULTS

### A. Phytochemical Assay Results

The phytochemical test qualitatively showed that the ethanol extract of *L. barbarum* L. fruit contained flavonoids, phenols, steroids, and terpenoids (Table I).

Table I . THE QUALITATIVE PHYTOCHEMICAL TEST RESULTS OF ETHANOL EXTRACT OF *LYCIUM BARBARUM* L. FRUIT

Extract	Test	Result
Ethanol extract of <i>L. barbarum</i> L. fruit	Flavonoids	+
	Phenols	+
	Quinones	-
	Steroids	+
	Terpenoids	+
	Alkaloids	-

### B. Microdilution Results and Total Plate Count Results

Results of this study showed that ethanol extract of *L. barbarum* L. fruit has antibacterial and antibiofilm effects against *S. mutans* and *P. gingivalis*. The most effective antibacterial effect was at 100 µg/mL concentration, with an optical density value of  $0.358 \pm 0.002$  (Fig. 1) and the total number of *S. mutans* colonies of  $3 \pm 3.46 \times 10^6$  CFU/mL (Fig.

3). The optical density value of *P. gingivalis* was  $0.458 \pm 0.024$  (Fig. 2) with a total number of *P. gingivalis* colonies of  $41 \pm 4.58 \times 10^6$  CFU/mL (Fig. 4).




Fig. 1. *Streptococcus mutans* (optical density) concentration response curves for treatment with ethanol extract of *Lycium barbarum* L. fruit in different concentration (100 µg/mL, 50 µg/mL, 25 µg/mL, 12.5 µg/mL, and 6.25 µg/mL). Chlorhexidine gluconate (0.2%) was used as a positive control and sterile distilled water was used as a negative control. \*\*Significant difference at  $p < 0.01$ .




Fig. 2. *Porphyromonas gingivalis* (optical density) concentration response curves for treatment with ethanol extract of *Lycium barbarum* L. fruit in different concentration (100 µg/mL, 50 µg/mL, 25 µg/mL, 12.5 µg/mL, and 6.25 µg/mL). Chlorhexidine gluconate (0.2%) was used as a positive control and sterile distilled water was used as a negative control. \*\*Significant difference at  $p < 0.01$ .





Fig. 3. *Streptococcus mutans* (colony forming unit) concentration response for treatment with ethanol extract of *Lycium barbarum* L. fruit in different concentration. Chlorhexidine gluconate (0.2%) was used as a positive control and sterile distilled water was used as a negative control. A. Distilled water as negative control; B. Concentration of 6.25 µg/mL; C. Concentration of 12.5 µg/mL; D. Concentration of 25 µg/mL; E. Concentration of 50 µg/mL; F. Concentration of 100 µg/mL; G. Chlorhexidine gluconate (0.2%) as positive control. \*\*Significant difference at  $p < 0.01$  and \*  $p < 0.05$ .





Fig. 4. *Porphyromonas gingivalis* (colony forming unit) concentration response for treatment with ethanol extract of *Lycium barbarum* L. fruit in different concentration. Chlorhexidine gluconate (0.2%) was used as a positive control and sterile distilled water was used as a negative control. A. Distilled water as a negative control; B. Concentration of 6.25 µg/mL; C. Concentration of 12.5 µg/mL; D. Concentration of 25 µg/mL; E. Concentration of 50 µg/mL; F. Concentration of 100 µg/mL; G. Chlorhexidine gluconate (0.2%) as a positive control. \*\*Significant difference at  $p < 0.01$ .

### C. Biofilm Assay Results

In biofilm assay, the concentration of 100  $\mu\text{g/mL}$  with 3 hours of incubation was the most effective in inhibiting the formation of *S. mutans* biofilm with an optical density value of  $0.042 \pm 0.002$  (Fig. 6), whereas for *P. gingivalis* biofilm, the concentration of 100  $\mu\text{g/mL}$  with 24 hours of incubation was the most effective (optical density value:  $0.007 \pm 0.003$ ; Fig. 10). Statistical analysis showed that all ethanol extract concentrations of *L. barbarum* L. fruit were significantly different from negative control (Fig. 5-10) ( $p < 0.05$ ).




Fig. 5. *Streptococcus mutans* (optical density) concentration response curves for treatment with ethanol extract of *Lycium barbarum* L. fruit in different concentration (100  $\mu\text{g/mL}$ , 50  $\mu\text{g/mL}$ , 25  $\mu\text{g/mL}$ , 12.5  $\mu\text{g/mL}$ , and 6.25  $\mu\text{g/mL}$ ) after 1 h incubation time. Chlorhexidine gluconate (0.2%) was used as a positive control and sterile distilled water was used as a negative control. \*\*Significant difference at  $p < 0.01$ .




Fig. 6. *Streptococcus mutans* (optical density) concentration response curves for treatment with ethanol extract of *Lycium barbarum* L. fruit in different concentration (100  $\mu\text{g/mL}$ , 50  $\mu\text{g/mL}$ , 25  $\mu\text{g/mL}$ , 12.5  $\mu\text{g/mL}$ , and 6.25  $\mu\text{g/mL}$ ) after 3 h incubation time. Chlorhexidine gluconate (0.2%) was used as a positive control and sterile distilled water was used as a negative control. \*\*Significant difference at  $p < 0.01$ .




Fig. 7. *Streptococcus mutans* (optical density) concentration response curves for treatment with ethanol extract of *Lycium barbarum* L. fruit in different concentration (100  $\mu\text{g/mL}$ , 50  $\mu\text{g/mL}$ , 25  $\mu\text{g/mL}$ , 12.5  $\mu\text{g/mL}$ , and 6.25  $\mu\text{g/mL}$ ) after 24 h incubation time. Chlorhexidine gluconate (0.2 %) was used as a positive control and sterile distilled water was used as a negative control. \*\*Significant difference at  $p < 0.01$ .




Fig. 8. *Porphyromonas gingivalis* (optical density) concentration response curves for treatment with ethanol extract of *Lycium barbarum* L. fruit in different concentration (100  $\mu\text{g/mL}$ , 50  $\mu\text{g/mL}$ , 25  $\mu\text{g/mL}$ , 12.5  $\mu\text{g/mL}$ , and 6.25  $\mu\text{g/mL}$ ) after 1 h incubation time. Chlorhexidine gluconate (0.2%) was used as a positive control and sterile distilled water was used as a negative control. \*\*Significant difference at  $p < 0.01$ .




Fig. 9. *Porphyromonas gingivalis* (optical density) concentration response curves for treatment with ethanol extract of *Lycium barbarum* L. fruit in different concentration (100  $\mu\text{g/mL}$ , 50  $\mu\text{g/mL}$ , 25  $\mu\text{g/mL}$ , 12.5  $\mu\text{g/mL}$ , and 6.25  $\mu\text{g/mL}$ ) after 3 h incubation time. Chlorhexidine gluconate (0.2%) was used as a positive control and sterile distilled water was used as a negative control. \*Significant difference at  $p < 0.05$ ; \*\*Significant difference at  $p < 0.01$ .




Fig. 10. *Porphyromonas gingivalis* (optical density) concentration response curves for treatment with ethanol extract of *Lycium barbarum* L. fruit in different concentration (100  $\mu\text{g/mL}$ , 50  $\mu\text{g/mL}$ , 25  $\mu\text{g/mL}$ , 12.5  $\mu\text{g/mL}$ , and 6.25  $\mu\text{g/mL}$ ) after 24 h incubation time. Chlorhexidine gluconate (0.2%) was used as a positive control and sterile distilled water was used as a negative control. \*\*Significant difference at  $p < 0.01$ .

### IV. DISCUSSION

The ethanol extract of *L. barbarum* L. fruit has various secondary metabolite such as flavonoids, phenols, steroids, and terpenoids. These metabolites have important role in inhibiting the bacterial growth of *S. mutans* and *P. gingivalis* biofilms. Flavonoid compounds can damage bacterial cell walls by removing bacterial proteins, nucleic acids, and nucleotides so that bacterial cell lysis occurs [23]. Flavonoids also interfere with the quorum sensing mechanism, which inhibits bacterial adhesion and biofilms formation on tooth

surface. The formation of biofilms is inhibited by the reduction of glucans, which are a medium for bacterial attachment, due to the inactivity of the glucosyltransferase enzyme by flavonoids [24]. The ability of bacterial cell protein denaturation by phenol compounds through the formation of bonds between phenols and proteins causes damage to protein structures. The cell wall and the cytoplasmic membrane are composed of these proteins. The disruption of permeability in the cell wall and cytoplasmic membrane causes irreversible damage and leads to lysis of the bacterial cell [25,26].

Steroid compounds interaction with cell phospholipid membranes are also capable of causing lysosome leakage for the lysis of bacterial cells. Terpenoids are lipophilic and can bind to carbohydrates and fats, causing disruption of the permeability of bacterial cell walls, denaturation of cytoplasmic proteins, and inactivation of cellular enzymes, causing lysis of bacterial cells [25,27,28].

There were fewer studies about the concentration of *L. barbarum* L. as antibacterial and antibiofilm in vitro. Therefore, as a preliminary study, the concentration of *L. barbarum* L. was used from 6.25 µg/mL to 100 µg/mL. Chlorhexidine gluconate (0.2%), a broad-spectrum antimicrobial biguanide, was used as a positive control due to its potent antiplaque agent. Chlorhexidine leads to cell death by penetrating into the cell and causes leakage of intracellular component. It is considered as a gold standard mouth rinse against gingivitis and periodontitis. Chlorhexidine is indicated to reduce pocket depth in periodontitis as an adjunct therapy to dental scaling and root planning procedure.

In microdilution method, smaller OD value corresponds to higher antibacterial activity. Results of this study showed that ethanol extract of *L. barbarum* L. in all concentration were significantly lower than negative control. Total plate count results also showed reduction of total bacteria number as the concentration of the extract increased. It means that ethanol extract of *L. barbarum* L. has antibacterial effect against *S. mutans* and *P. gingivalis*. Ethanol extract of *L. barbarum* L. fruit at 100 µg/mL concentration was the most effective concentration in inhibiting *S. mutans* and *P. gingivalis* bacteria and biofilms. The results of this study are supported with research by Alassadi et al (2015) towards *L. barbarum* L. fruit, which showed that alcohol group (-OH) in flavonoid structure increased the ability of the extract to inhibit microbial growth by increasing the permeability of bacterial cell membranes, and the highest concentration, at 100 µg/mL, possessed the most effective antibacterial activity compared to other concentrations, due to less flavonoid content at lower concentrations [18]. The results of this study are following with previous study regarding inhibition effect of *L. barbarum* L. extracts towards *S. aureus* and *E. coli* using disc diffusion method. Based on the results of these studies, there is an antibacterial effect against *E. coli* [29]. The results of other studies using the well diffusion method have stated that the ethanol extract of *L. barbarum* L. fruit at concentrations of 10 µg/mL and 20 µg/mL had inhibitory effects against *S. aureus* and *E. coli* [22]

This study used three different incubation times in the biofilm assay, namely 1, 3, and 24 hours, to determine the most effective phase for inhibiting *S. mutans* and *P. gingivalis* biofilms. The difference in incubation time was

similar to the biofilm formation phase, starting with the pellicle formation phase in a few minutes to 1 hour, the initial adhesion phase at 2 to 4 hours, and the maturation phase after 24 hours [30].

Biofilm assay results against *S. mutans* showed that all extract concentration in 1, 3, and 24-hours incubation time had significantly lower OD than negative control. It means that ethanol extract of *L. barbarum* L. can inhibit the formation of *S. mutans* biofilm starts from 6.25 µg/mL to 100 µg/mL. In 1 and 24-hours incubation time, 100 µg/mL extract had significantly higher OD than the positive control. Hence, the antibiofilm effect of 100 µg/mL extract was more effective than positive control. This result is similar to extract concentration of 25 µg/mL, 50 µg/mL, and 100 µg/mL in 3-hours incubation.

The formation of *P. gingivalis* biofilm was inhibited from concentration of 12.5 µg/mL to 100 µg/mL in 3 and 24-hours incubation time and all concentration in 1 hour incubation time. Therefore, the antibiofilm effect of 50 µg/mL and 100 µg/mL extract in 3 and 24-hours incubation were more effective than positive control. This result is similar to extract concentration of 25 µg/mL, 50 µg/mL, and 100 µg/mL in 1-hour incubation.

The results of the antibiofilm assay showed that the most effective incubation time for inhibiting the formation of *S. mutans* biofilms was at 3-hours of incubation time, and for *P. gingivalis*, it was at 24-hours of incubation time. The most effective times for inhibiting biofilm formation were at the initial adhesion and maturation phases, respectively. The antibiofilm effect depends on the inhibition of polymer matrix formation and quorum sensing, or communication, between bacterial cells in biofilms by inhibiting autoinducer peptides, signaling molecules in Gram-positive bacteria, and acylhomoserine lactones (AHLs) in Gram-negative bacteria so bacterial virulence factors and biofilm development may be inhibited [31].

This is proven by the lowest optical density value found at a 100 µg/mL concentration in *S. mutans* ( $0.042 \pm 0.002$ ) and *P. gingivalis* ( $0.007 \pm 0.003$ ). This antibiofilm assay also showed that a concentration 100 µg/mL had a lower optical density value and was significantly different from the positive control, which means that at a 100 µg/mL concentration, the antibiofilm effect was more effective than the positive control.

## V. CONCLUSION

The ethanol extract of *L. barbarum* L. fruit, containing flavonoids, phenols, steroids, and terpenoids, have antibacterial and antibiofilm effects against *S. mutans* and *P. gingivalis*. The most effective concentration is 100 µg/mL for both bacteria with 87.15% and 97.73% of biofilms reduction for *S. mutans* and *P. gingivalis*, respectively. *L. barbarum* L. might be a promising natural-therapeutic agent as an alternative therapy. However, further research using toxicity, preclinical, and clinical tests is needed to determine whether *L. barbarum* L. fruit ethanol extract can be used as alternative mouthwash for preventing caries and treating chronic periodontitis.

## ACKNOWLEDGMENT

The authors thank Faculty of Dentistry, Trisakti University, for invaluable support in this study. Authors also

would like to thank Mario Richi, S.Si from the Microbiology Center of Research and Education (MiCORE) laboratory and Sheila Sutanto, S.Si from Biological Collaborative Research and Education (BioCORE) laboratory for their laboratory assistances.

#### REFERENCES

- [1] F. Katge, B. Rusawat, A. Shitoot, M. Poojari, T. Pammi, and D. Patil, "DMFT index assessment, plaque pH, and microbiological analysis in children with special health care needs, India," *Journal of International Society of Preventive and Community Dentistry*, vol. 5, pp. 383–388, 2015
- [2] Ministry of Health of the Republic of Indonesia. "Report on result of national basic health research (Riskesmas) 2018", Jakarta: Ministry of Health of the Republic of Indonesia pp. 1–582, 2018
- [3] C. Heng, "Tooth decay is the most prevalent disease," *Federal Practitioner*, vol. 33(10), pp. 31–33, 2016
- [4] S. Chenicheri, U. R. Ramachandran, V. Thomas, and A. Wood, "Insight into oral biofilm: primary, secondary and residual caries and phyto-challenged solutions," *Open Dentistry Journal*, vol. 11, pp. 312–333, 2017
- [5] G. Conrads and I. About, "Pathophysiology of dental caries," *Monograph in Oral Science*, vol. 27, pp. 1–10, 2018
- [6] J. E. Hinrichs and G. Kotsakis, "Classification of diseases and conditions affecting the periodontium," in Carranza, 2015, pp. 51–55.
- [7] S. Kumar and J. Sengupta, "Endodontic treatment for mandibular molars using ProTaper," *Medical Journal Armed Forces India*, vol. 67, no. 4, pp. 377–379, 2011
- [8] D. Sharma, L. Misba, and A. U. Khan. "Antibiotics versus biofilm: an emerging battleground in microbial communities," *Antimicrobial Resistance and Infection Control*, vol. 8, pp. 1–10, 2019
- [9] A. S. K. J. Arul and P. Palanivelu. "Biofilm forming ability of a new bacterial isolate from dental caries: An atomic force microscopic study," *Journal of Natural Science, Biology, and Medicine*, vol. 5, no. 2, pp. 278–283, 2014
- [10] J. Abranches, L. Zeng, J. K. Kajfasz, S. R. Palmer, B. Chakraborty, Z. T. Wen et al., "Biology of oral Streptococci," *Microbiology Spectrum*, vol. 6, no. 5, 2018
- [11] K. Bao, G. N. Belibasakis, T. Thurnheer, J. Aduse- Opoku, M. A. Curtis, and N. Bostanci, "Role of Porphyromonas gingivalis gingipains in multi-species biofilm formation," *BioMed Central Microbiology*, vol. 14, p. 258, 2014
- [12] L. Jia, N. Han, J. Du, L. Guo, Z. Luo, and Y. Liu, "Pathogenesis of Important Virulence Factors of Porphyromonas gingivalis via Toll-Like Receptors," *Frontier in Cellular and Infection Microbiology*, vol. 9, p. 262, 2019
- [13] Z. L. S. Brukes, R. Bescos, L. A. Belfield, K. Ali, and A. Robert, "Current uses of chlorhexidine for management of oral disease: a narrative review," *Journal of Dentistry*, vol. 103, p. 103497, 2020
- [14] N. Jeddy, S. Ravi, T. Radhika, and L. J. S. Lakshmi, "Comparison of the efficacy of herbal mouth rinse with commercially available mouth rinses: A clinical trial," *Journal of Oral Maxillofacial Pathology*, vol. 22, no. 3, pp. 332–334, 2018
- [15] C. C. Martínez, M. D. Gómez, and M. S. Oh. "Use of traditional herbal medicine as an alternative in dental treatment in mexican dentistry: A review," *Pharmaceutical Biology*, vol. 55 (1), pp. 1992–1998, 2017
- [16] World Health Organization. "WHO traditional medicine strategy: 2014-2023", Geneva: World Health Organization, 2013; Available from: [https://www.who.int/medicines/publications/traditional/trm\\_strategy14\\_23/en/](https://www.who.int/medicines/publications/traditional/trm_strategy14_23/en/)
- [17] Z. F. Ma, H. Zhang, S. S. Teh, C. W. Wang, Y. Zhang, F. Hayford et al., "Goji berries as a potential natural antioxidant medicine: An insight into their molecular mechanisms of action," *Oxidative Medicine and Cellular Longevity*, p. e2437397, 2019
- [18] I. J. Alassadi, F. S. Sabah, and L. A. Alrubaie. "Isolation of flavonoid compound from iraqi awsaj plant (Lycium Barbarum L.) Fruits and the Study of Its Antibacterial Activity," *European Scientific Journal*, vol 11, no. 24, pp. 268–276, 2015
- [19] S-E. Byambasuren, J. Wang, and G. Gaudel, "Medicinal value of wolfberry (Lycium barbarum L.)," *Journal of Medical Plants Studies*, vol. 7, no. 4, pp. 90–97, 2019
- [20] Z. F. Ma, H. Zhang, S. S. Teh, C. W. Wang, Y. Zhang, F. Hayford et al., "Goji berries as a potential natural antioxidant medicine: An insight into their molecular mechanisms of action," *Hindawi*, pp. 1–9, 2019
- [21] P. Skenderidis, D. Lampakis, I. Giavasis, S. Leontopoulos, K. Petrotos, C. Hadjichristodoulou et al., "Chemical properties, fatty-acid composition, and antioxidant activity of Goji berry (Lycium barbarum L. and Lycium chinese mill.) fruits," *Antioxidants*, vol. 8, no. 3, p. 60, 2019
- [22] P. Skenderidis, D. Lampakis, I. Giavasis, S. Leontopoulos, K. Petrotos, C. Hadjichristodoulou et al., "The in vitro antimicrobial activity assessment of ultrasound assisted Lycium barbarum fruit extracts and pomegranate fruit peels," *Journal of Food Measurement and Characterization*, vol. 13, no 5, pp. 1-15, 2019
- [23] Z. Y. Dewi, A. Nur, and T. Hertriani, "Efek antibakteri dan penghambatan biofilm ekstrak sereh (Cymbopogon nardus L.) terhadap bakteri Streptococcus mutans," *Majalah Kedokteran Gigi Indonesia*, vol 1, no. 2, p. 136, 2015
- [24] S. Loresta, S. Murwani, and P. Trisunuwati "Efek ekstrak etanol daun kelor (Moringa oleifera) terhadap pembentukan biofilm Staphylococcus aureus secara in vitro, pp. 20–25, Available from: <http://library1.nida.ac.th/termpaper6/sd/2554/19755.pdf>
- [25] S. Bontjura, O. A. Waworuntu, and K. V. Siagia. "Uji efek antibakteri ekstrak daun leilem (Clerodendrum Minahassae L.) terhadap bakteri Streptococcus mutans," *Pharmacon*, vol. 4, no. 4, p. 6, 2015
- [26] L. Bouarab-Chibane, V. Forquet, P. Lantéri, Y. Clément, L. Léonard-Akkari, N. Oulahal et al., "Antibacterial properties of polyphenols: characterization and QSAR (quantitative structure-activity relationship) models," *Frontiers in Microbiology*, vol. 10, p. 829, 2019
- [27] Y. Mulya, "Phytochemical analysis and antibacterial properties of some selected Indian medicinal plants," *International Journal of Current Microbiology and Applied Sciences*, pp. 228–235, 2015
- [28] A. Ludwiczuk, K. Skalicka-Woźniak, and M. I. Georgiev, "Terpenoids," in *Pharmacognosy: Fundamentals, Applications and Strategy*, p. 251, 2017
- [29] N. I. Fit, F. Chirilă, G. Nadăș, E. Pall, and R. Mureșan, "Comparative testing of antimicrobial activity of aqueous extracts of Aloe vera and Lycium barbarum," *Bulletin of University of Agricultural Sciences and Veterinary Medicine of Cluj*, vol. 70, no. 1, pp. 72–76, 2013
- [30] A. S. Widyarman and N. K. E. Lazaroni, "Persistent endodontics pathogens biofilm inhibited by Lactobacillus reuteri Indonesian strain," *Journal of Dentistry Indonesia*, vol. 26, no. 3, pp. 160–164, 2019
- [31] L. Lu, W. Hu, Z. Tian, D. Yuan, G. Yi, Y. Zhou Y et al., "Developing natural products as potential anti-biofilm agents," *Chinese Medicine*, p. 11, 2019

# CERTIFICATE

This Certificate is awarded to :

*drg. Sheila Soesanto, MKG*

for the contribution as a Participant in :



July, 14-16<sup>th</sup> 2021  
Medan - Indonesia

Refi Ikhtiari, Ph.D  
General Chair

Organized by :





# Goji Berry Extract (*Lycium barbarum* L.) Efficacy on Oral Pathogen Biofilms

*by* Sheila Soesanto FKG

---

**Submission date:** 22-Feb-2024 01:49PM (UTC+0700)

**Submission ID:** 2301393628

**File name:** xtract\_Lycium\_barbarum\_L.\_Efficacy\_on\_Oral\_Pathogen\_Biofilms.pdf (3.06M)

**Word count:** 4980

**Character count:** 27091

# Goji Berry Extract (*Lycium barbarum* L.) Efficacy on Oral Pathogen Biofilms

Sheila Soesanto  
Pharmacology Department Faculty of  
Dentistry  
Universitas Trisakti  
Jakarta, Indonesia  
mass9977@yahoo.com

Ricky Chandra Jaya Soen  
Undergraduate Program Faculty of  
Dentistry  
Universitas Trisakti  
Jakarta, Indonesia  
ricksoen@gmail.com

Reynalda Oktaviani  
Undergraduate Program Faculty of  
Dentistry  
Universitas Trisakti  
Jakarta, Indonesia  
reynaldaoktaviani@gmail.com

Armelia Sari Widyarman\*  
Microbiology Department Faculty of  
Dentistry, Universitas Trisakti  
Jakarta, Indonesia  
armeliasari@trisakti.ac.id

**Abstract**— *Lycium barbarum* L. fruit, which contains flavonoids and phenolic acids, has antibacterial properties that are expected to inhibit bacterial growth. The objective of this study is to determine the antibacterial and the antibiofilm effects of *L. barbarum* L. fruit ethanol extract towards *S. mutans* and *P. gingivalis*. An in-vitro laboratory experiment was performed with a post-test control group design. The extract of *L. barbarum* L. fruit was obtained by maceration technique using 96% ethanol as a solvent. The antibacterial assay was performed by microdilution and plate count methods. The antibiofilm effect was performed using a biofilm-assay method. The results of microdilution and plate count methods showed that the most effective antibacterial concentration against *S. mutans* and *P. gingivalis* was 100 µg/mL when compared with negative control ( $p < 0.05$ ). In the biofilm assay, the most effective concentration against *S. mutans* was 100 µg/mL at the 3-hours incubation time, while for *P. gingivalis*, the most effective concentration was 100 µg/mL at 24-hours incubation time when compared with negative control ( $p < 0.05$ ). These results indicate that ethanol extract of *L. barbarum* L. fruit was demonstrated to have antibacterial and antibiofilm effects against oral pathogens *S. mutans* and *P. gingivalis* with 87.15% and 97.73% of biofilms reduction respectively.

**Keywords**— antibacterial, antibiofilm, *Lycium barbarum* L., *Porphyromonas gingivalis*, *Streptococcus mutans*.

## I. INTRODUCTION

Oral health positively affects the appearance, physical, mental, and interpersonal well-being, of an individual. Oral health is a part of overall health, contributes to the quality of life [1]. National Basic Health Research Data (Riskesdas 2018) showed dental and oral health problems in 57.6% of the Indonesian population. The prevalence of dental caries in Indonesia in 2018 was 88.8%, with an average DMF-T index of 7.1, which is a very high severity of dental caries. Moreover, 74.1% of the Indonesian population experienced periodontitis [2].

Caries are the process of the demineralization of inorganic material and the dissolution of organic material, leading to bacterial invasion through the dentin layer until it reaches the pulp [3,4]. The process of dental caries depends on the presence of fermentable sugars (substrates), the type of tooth and saliva (host), cariogenic microbial flora (biofilm), and time [5]. Periodontitis is a disease caused by microorganisms that cause inflammation of tooth-supporting tissue and causes progressive destruction of periodontal ligament and alveolar bone. The sign of periodontitis is the

formation of pockets, recessions, or both [6]. Periodontitis in adults caused by numerous local factors, such as biofilms or calculus, is classified as chronic periodontitis [7].

The formation of biofilms begins when microorganisms in the planktonic state merge into bacterial colonies and wrap themselves in a self-produced extracellular polymer matrix [8]. In the initial phase of biofilm formation, there is an increase in Gram-positive cocci activity, one of which is *Streptococcus mutans*, which is able to attach to tooth surface through the formation of extracellular polysaccharides that cause biofilm matrix to have a gelatin-like consistency that facilitates attachment of bacteria to the tooth surface [4,9,10]. *Porphyromonas gingivalis*, which is a secondary bacterium, is an anaerobic Gram-negative bacterium found in periodontal pockets that causes chronic periodontitis. Various virulence factors of *P. gingivalis* such as gingipains, fimbriae, and lipopolysaccharides, play important role in periodontal disease progression and induce dysbiosis in biofilms [11,12].

Chlorhexidine mouthwash is used to prevent caries and treat periodontitis and is considered as the gold standard for controlling dental plaque and gingivitis due to its efficacy against a wide variety of microorganisms. However, chlorhexidine has various side effects, including taste disturbances, discoloration of teeth and mucosa, mucosal desquamation, salivary stone formation, irritation, dry oral cavity, and allergic reactions, such as contact stomatitis. The World Health Organization (WHO) recommends finding new natural ingredients to overcome the side effects of chemical agents [13,14].

The use of natural ingredients as antimicrobial agents has become an alternative because of their low cost and lower toxicity [15]. According to WHO, traditional medicine has been used globally and can be a major source of health for millions of people and sometimes the only source of care and is also culturally acceptable, affordable, and trusted by community [16]. Goji berry (*Lycium barbarum* L.) has been widely used as a traditional medicine by people in Asia, especially in the northwestern part of China, for more than 2000 years. Recently, *L. barbarum* L. has been gaining popularity as a highly nutritious food used to improve health in North America, Europe, and Asia [17]. *Lycium barbarum* L. has a red, oblong fruit with a length of 6–20 mm and a diameter of 3–10 mm. *Lycium barbarum* L. fruit is harvested when it is ripe and is then dried for later use [18]. The fruit, roots, tree bark, and flowers of *L. barbarum* L. are used as medicine [19].

The polysaccharides of *L. barbarum* L. exhibited properties that improve eye health and reproductive system; reduce fat and blood sugar; regulate immunity. It's also anticancer, anti-tumor, antioxidant, anti-fatigue, antiviral, anti-aging, hepatoprotective, neuroprotective, and cardioprotective properties [17,20]. The flavonoids and phenolic acids of *L. barbarum* L. have potential as antioxidants and antimicrobials [21].

*Lycium barbarum* L. fruit is effective against Gram-negative bacteria (e.g. *Escherichia coli*) and Gram-positive bacteria (e.g. *Staphylococcus aureus*) [22]. However, there have been no studies regarding the antibacterial effect of *L. barbarum* L. fruit against *S. mutans* and *P. gingivalis* as causing bacteria of dental caries and chronic periodontitis. Thus, this is the first study that analyzed the antibacterial and antibiofilm efficacy of goji berry (*L. barbarum* L.) ethanol extract towards *S. mutans* and *P. gingivalis* as oral pathogens.

## II. MATERIALS AND METHODS

### A. Ethanol extract production from *L. barbarum* L. fruit

Dried *L. barbarum* L. fruit (100 g) from Chinese medicine store "Lancar Jaya" at Teluk Gong Raya No. 43, Jakarta Utara (produced in Zhongning, Ningxia, China) was ground in a blender until it became powder. It was then immersed in 96% ethanol with a ratio of 1:8 for 72 hours, stirring every 15 minutes. Furthermore, filtration was performed using Whatman No. 1 filter paper and evaporated with a rotary evaporator at 40°C temperature, 60 rpm speed, and 20 atm pressure so a thick and solvent-free extract was obtained with a concentration of 100 µg/mL. Furthermore, extracts were diluted using sterile distilled water until concentrations of 50, 25, 12.5, and 6.25 µg/mL were obtained.

### B. Phytochemical Assay

Phytochemical assays were performed qualitatively to determine whether the ethanol extract of *L. barbarum* L. fruit contained flavonoids, phenols, quinones, steroids, terpenoids, and alkaloids. The qualitative results are expressed as (+) for the presence and (-) for the absence of phytochemicals.

### C. Bacterial Cultures

*Streptococcus mutans* ATCC 25175 and *Porphyromonas gingivalis* ATCC 33277 from Microbiology Center of Research and Education (MiCORE) Laboratory, Faculty of Dentistry, Trisakti University, were cultured on brain heart infusion (BHI) (Oxoid, Hampshire) broth medium and incubated at 37°C for 24 hours in anaerobic atmosphere. Furthermore, the absorbance measurements were performed to obtain the McFarland standard of  $0.5 = 1.5 \times 10^8$  CFU/mL ( $OD_{600} = 0.132$ ).

### D. Microdilution

Each well of a 96-well plate was distributed 100 µL of either an *S. mutans* or *P. gingivalis* culture. Subsequently, 100 µL of the following solutions was used as a treatment: ethanol extracts of *L. barbarum* L. fruit at 100, 50, 25, 12.5, and 6.25 µg/mL concentrations, 0.2% chlorhexidine gluconate as a positive control, and sterile distilled water as a negative control. The measurement of bacterial cell density was performed using microplate reader at 600 nm wavelength before and after the 96-well plates were

incubated for 24 hours. All treatments were done in triplicate.

### E. Total Plate Count

Bacterial growth was measured by re-diluting contents in 96-well plates for 10,000 times and cultured on BHI agar medium and incubated for 24 hours at 37°C.

### F. Biofilm Assay

Bacterial culture (200 µL) was dispensed into each 96-well plate and incubated at 37°C for 48 hours in an anaerobic atmosphere. Furthermore, the supernatant was removed until a thin layer of biofilm was left on the bottom surface of the well. Then, wells were rinsed with a solution of phosphate-buffered saline (PBS). The ethanol extracts of *L. barbarum* L. fruit at a concentration of 100, 50, 25, 12.5, and 6.25 µg/mL, 0.2% chlorhexidine gluconate as positive control, and sterile distilled water as negative control were added 200 µL to wells using a micropipette and incubated at 37°C for 1, 3, and 24 h in an anaerobic atmosphere. The well was rinsed twice using PBS and then fixed over a flame. Crystal violet dye (200 µL; 0.05% w/v) was added to each well and left for 15 minutes. The well was rinsed twice using PBS and left for 15 minutes. Then, 200 µL of 96% ethanol was inserted, and optical density was measured using a microplate reader (SAFAS MP96, Monaco) at 595 nm wavelength.

### G. Statistical Analysis

Data were processed by using Statistical Product and Service Solution (SPSS) software version 25.0 and the normality test was performed by using the Shapiro-Wilk method. Normally distributed data ( $p > 0.05$ ) was analyzed by one-way analysis of variance (ANOVA) test. Significant data ( $p < 0.05$ ) were analyzed with a posthoc test using Tukey's test with a significance level of  $P < 0.05$  to determine which groups were significantly different.

## III. RESULTS

### A. Phytochemical Assay Results

The phytochemical test qualitatively showed that the ethanol extract of *L. barbarum* L. fruit contained flavonoids, phenols, steroids, and terpenoids (Table I.).

Table I . THE QUALITATIVE PHYTOCHEMICAL TEST RESULTS OF ETHANOL EXTRACT OF *LYCIUM BARBARUM* L. FRUIT

Extract	Test	Result
Ethanol extract of <i>L. barbarum</i> L. fruit	Flavonoids	+
	Phenols	+
	Quinones	-
	Steroids	+
	Terpenoids	+
	Alkaloids	-

### B. Microdilution Results and Total Plate Count Results

Results of this study showed that ethanol extract of *L. barbarum* L. fruit has antibacterial and antibiofilm effects against *S. mutans* and *P. gingivalis*. The most effective antibacterial effect was at 100 µg/mL concentration, with an optical density value of  $0.358 \pm 0.002$  (Fig. 1) and the total number of *S. mutans* colonies of  $3 \pm 3.46 \times 10^6$  CFU/mL (Fig.

3). The optical density value of *P. gingivalis* was  $0.458 \pm 0.024$  (Fig. 2) with a total number of *P. gingivalis* colonies of  $41 \pm 4.58 \times 10^6$  CFU/mL (Fig. 4).




Fig. 1. Streptococcus mutans (optical density) concentration response curves for treatment with ethanol extract of Lycium barbarum L. fruit in different concentration (100  $\mu\text{g/mL}$ , 50  $\mu\text{g/mL}$ , 25  $\mu\text{g/mL}$ , 12.5  $\mu\text{g/mL}$ , and 6.25  $\mu\text{g/mL}$ ). Chlorhexidine gluconate (0.2%) was used as a positive control and sterile distilled water was used as a negative control. \*\*Significant difference at  $p < 0.01$ .




Fig. 2. Porphyromonas gingivalis (optical density) concentration response curves for treatment with ethanol extract of Lycium barbarum L. fruit in different concentration (100  $\mu\text{g/mL}$ , 50  $\mu\text{g/mL}$ , 25  $\mu\text{g/mL}$ , 12.5  $\mu\text{g/mL}$ , and 6.25  $\mu\text{g/mL}$ ). Chlorhexidine gluconate (0.2%) was used as a positive control and sterile distilled water was used as a negative control. \*\*Significant difference at  $p < 0.01$ .





Fig. 3. Streptococcus mutans (colony forming unit) concentration response for treatment with ethanol extract of Lycium barbarum L. fruit in different concentration. Chlorhexidine gluconate (0.2%) was used as a positive control and sterile distilled water was used as a negative control. A. Distilled water as negative control; B. Concentration of 6.25  $\mu\text{g/mL}$ ; C. Concentration of 12.5  $\mu\text{g/mL}$ ; D. Concentration of 25  $\mu\text{g/mL}$ ; E. Concentration of 50  $\mu\text{g/mL}$ ; F. Concentration of 100  $\mu\text{g/mL}$ ; G. Chlorhexidine gluconate (0.2%) as positive control. \*\*Significant difference at  $p < 0.01$  and \*  $p < 0.05$ .





Fig. 4. Porphyromonas gingivalis (colony forming unit) concentration response for treatment with ethanol extract of Lycium barbarum L. fruit in different concentration. Chlorhexidine gluconate (0.2%) was used as a positive control and sterile distilled water was used as a negative control. A. Distilled water as a negative control; B. Concentration of 6.25  $\mu\text{g/mL}$ ; C. Concentration of 12.5  $\mu\text{g/mL}$ ; D. Concentration of 25  $\mu\text{g/mL}$ ; E. Concentration of 50  $\mu\text{g/mL}$ ; F. Concentration of 100  $\mu\text{g/mL}$ ; G. Chlorhexidine gluconate (0.2%) as positive control. \*\*Significant difference at  $p < 0.01$ .

### C. Biofilm Assay Results

In biofilm assay, the concentration of 100  $\mu\text{g/mL}$  with 3 hours of incubation was the most effective in inhibiting the formation of *S. mutans* biofilm with an optical density value of  $0.042 \pm 0.002$  (Fig. 6), whereas for *P. gingivalis* biofilm, the concentration of 100  $\mu\text{g/mL}$  with 24 hours of incubation was the most effective (optical density value:  $0.007 \pm 0.003$ ; Fig. 10). Statistical analysis showed that all ethanol extract concentrations of *L. barbarum* L. fruit were significantly different from negative control (Fig. 5-10) ( $p < 0.05$ ).




Fig. 5. *Streptococcus mutans* (optical density) concentration response curves for treatment with ethanol extract of *Lycium barbarum* L. fruit in different concentration (100  $\mu\text{g/mL}$ , 50  $\mu\text{g/mL}$ , 25  $\mu\text{g/mL}$ , 12.5  $\mu\text{g/mL}$ , and 6.25  $\mu\text{g/mL}$ ) after 1 h incubation time. Chlorhexidine gluconate (0.2%) was used as a positive control and sterile distilled water was used as a negative control. \*\*Significant difference at  $p < 0.01$ .




Fig. 6. *Streptococcus mutans* (optical density) concentration response curves for treatment with ethanol extract of *Lycium barbarum* L. fruit in different concentration (100  $\mu\text{g/mL}$ , 50  $\mu\text{g/mL}$ , 25  $\mu\text{g/mL}$ , 12.5  $\mu\text{g/mL}$ , and 6.25  $\mu\text{g/mL}$ ) after 3 h incubation time. Chlorhexidine gluconate (0.2%) was used as a positive control and sterile distilled water was used as a negative control. \*\*Significant difference at  $p < 0.01$ .




Fig. 7. *Streptococcus mutans* (optical density) concentration response curves for treatment with ethanol extract of *Lycium barbarum* L. fruit in different concentration (100  $\mu\text{g/mL}$ , 50  $\mu\text{g/mL}$ , 25  $\mu\text{g/mL}$ , 12.5  $\mu\text{g/mL}$ , and 6.25  $\mu\text{g/mL}$ ) after 24 h incubation time. Chlorhexidine gluconate (0.2%) was used as a positive control and sterile distilled water was used as a negative control. \*\*Significant difference at  $p < 0.01$ .




Fig. 8. *Porphyromonas gingivalis* (optical density) concentration response curves for treatment with ethanol extract of *Lycium barbarum* L. fruit in different concentration (100  $\mu\text{g/mL}$ , 50  $\mu\text{g/mL}$ , 25  $\mu\text{g/mL}$ , 12.5  $\mu\text{g/mL}$ , and 6.25  $\mu\text{g/mL}$ ) after 1 h incubation time. Chlorhexidine gluconate (0.2%) was used as a positive control and sterile distilled water was used as a negative control. \*\*Significant difference at  $p < 0.01$ .




Fig. 9. *Porphyromonas gingivalis* (optical density) concentration response curves for treatment with ethanol extract of *Lycium barbarum* L. fruit in different concentration (100  $\mu\text{g/mL}$ , 50  $\mu\text{g/mL}$ , 25  $\mu\text{g/mL}$ , 12.5  $\mu\text{g/mL}$ , and 6.25  $\mu\text{g/mL}$ ) after 3 h incubation time. Chlorhexidine gluconate (0.2%) was used as a positive control and sterile distilled water was used as a negative control. \*Significant difference at  $p < 0.05$ ; \*\*Significant difference at  $p < 0.01$ .




Fig. 10. *Porphyromonas gingivalis* (optical density) concentration response curves for treatment with ethanol extract of *Lycium barbarum* L. fruit in different concentration (100  $\mu\text{g/mL}$ , 50  $\mu\text{g/mL}$ , 25  $\mu\text{g/mL}$ , 12.5  $\mu\text{g/mL}$ , and 6.25  $\mu\text{g/mL}$ ) after 24 h incubation time. Chlorhexidine gluconate (0.2%) was used as a positive control and sterile distilled water was used as a negative control. \*\*Significant difference at  $p < 0.01$ .

### IV. DISCUSSION

The ethanol extract of *L. barbarum* L. fruit has various secondary metabolite such as flavonoids, phenols, steroids, and terpenoids. These metabolites have important role in inhibiting the bacterial growth of *S. mutans* and *P. gingivalis* biofilms. Flavonoid compounds can damage bacterial cell walls by removing bacterial proteins, nucleic acids, and nucleotides so that bacterial cell lysis occurs [23]. Flavonoids also interfere with the quorum sensing mechanism, which inhibits bacterial adhesion and biofilms formation on tooth

surface. The formation of biofilms is inhibited by the reduction of glucans, which are a medium for bacterial attachment, due to the inactivity of the glucosyltransferase enzyme by flavonoids [24]. The ability of bacterial cell protein denaturation by phenol compounds through the formation of bonds between phenols and proteins causes damage to protein structures. The cell wall and the cytoplasmic membrane are composed of these proteins. The disruption of permeability in the cell wall and cytoplasmic membrane causes irreversible damage and leads to lysis of the bacterial cell [25,26].

Steroid compounds interaction with cell phospholipid membranes are also capable of causing lysosome leakage for the lysis of bacterial cells. Terpenoids are lipophilic and can bind to carbohydrates and fats, causing disruption of the permeability of bacterial cell walls, denaturation of cytoplasmic proteins, and inactivation of cellular enzymes, causing lysis of bacterial cells [25,27,28].

There were fewer studies about the concentration of *L. barbarum* L. as antibacterial and antibiofilm in vitro. Therefore, as a preliminary study, the concentration of *L. barbarum* L. was used from 6.25 µg/mL to 100 µg/mL. Chlorhexidine gluconate (0.2%), a broad-spectrum antimicrobial biguanide, was used as a positive control due to its potent antiplaque agent. Chlorhexidine leads to cell death by penetrating into the cell and causes leakage of intracellular component. It is considered as a gold standard mouth rinse against gingivitis and periodontitis. Chlorhexidine is indicated to reduce pocket depth in periodontitis as an adjunct therapy to dental scaling and root planning procedure.

In microdilution method, smaller OD value corresponds to higher antibacterial activity. Results of this study showed that ethanol extract of *L. barbarum* L. in all concentration were significantly lower than negative control. Total plate count results also showed reduction of total bacteria number as the concentration of the extract increased. It means that ethanol extract of *L. barbarum* L. has antibacterial effect against *S. mutans* and *P. gingivalis*. Ethanol extract of *L. barbarum* L. fruit at 100 µg/mL concentration was the most effective concentration in inhibiting *S. mutans* and *P. gingivalis* bacteria and biofilms. The results of this study are supported with research by Alassadi et al (2015) towards *L. barbarum* L. fruit, which showed that alcohol group (-OH) in flavonoid structure increased the ability of the extract to inhibit microbial growth by increasing the permeability of bacterial cell membranes, and the highest concentration, at 100 µg/mL, possessed the most effective antibacterial activity compared to other concentrations, due to less flavonoid content at lower concentrations [18]. The results of this study are following with previous study regarding inhibition effect of *L. barbarum* L. extracts towards *S. aureus* and *E. coli* using disc diffusion method. Based on the results of these studies, there is an antibacterial effect against *E. coli* [29]. The results of other studies using the well diffusion method have stated that the ethanol extract of *L. barbarum* L. fruit at concentrations of 10 µg/mL and 20 µg/mL had inhibitory effects against *S. aureus* and *E. coli* [22]

This study used three different incubation times in the biofilm assay, namely 1, 3, and 24 hours, to determine the most effective phase for inhibiting *S. mutans* and *P. gingivalis* biofilms. The difference in incubation time was

similar to the biofilm formation phase, starting with the pellicle formation phase in a few minutes to 1 hour, the initial adhesion phase at 2 to 4 hours, and the maturation phase after 24 hours [30].

Biofilm assay results against *S. mutans* showed that all extract concentration in 1, 3, and 24-hours incubation time had significantly lower OD than negative control. It means that ethanol extract of *L. barbarum* L. can inhibit the formation of *S. mutans* biofilm starts from 6.25 µg/mL to 100 µg/mL. In 1 and 24-hours incubation time, 100 µg/mL extract had significantly higher OD than the positive control. Hence, the antibiofilm effect of 100 µg/mL extract was more effective than positive control. This result is similar to extract concentration of 25 µg/mL, 50 µg/mL, and 100 µg/mL in 3-hours incubation.

The formation of *P. gingivalis* biofilm was inhibited from concentration of 12.5 µg/mL to 100 µg/mL in 3 and 24-hours incubation time and all concentration in 1 hour incubation time. Therefore, the antibiofilm effect of 50 µg/mL and 100 µg/mL extract in 3 and 24-hours incubation were more effective than positive control. This result is similar to extract concentration of 25 µg/mL, 50 µg/mL, and 100 µg/mL in 1-hour incubation.

The results of the antibiofilm assay showed that the most effective incubation time for inhibiting the formation of *S. mutans* biofilms was at 3-hours of incubation time, and for *P. gingivalis*, it was at 24-hours of incubation time. The most effective times for inhibiting biofilm formation were at the initial adhesion and maturation phases, respectively. The antibiofilm effect depends on the inhibition of polymer matrix formation and quorum sensing, or communication, between bacterial cells in biofilms by inhibiting autoinducer peptides, signaling molecules in Gram-positive bacteria, and acylhomoserine lactones (AHLs) in Gram-negative bacteria so bacterial virulence factors and biofilm development may be inhibited [31].

This is proven by the lowest optical density value found at a 100 µg/mL concentration in *S. mutans* ( $0.042 \pm 0.002$ ) and *P. gingivalis* ( $0.007 \pm 0.003$ ). This antibiofilm assay also showed that a concentration 100 µg/mL had a lower optical density value and was significantly different from the positive control, which means that at a 100 µg/mL concentration, the antibiofilm effect was more effective than the positive control.

## V. CONCLUSION

The ethanol extract of *L. barbarum* L. fruit, containing flavonoids, phenols, steroids, and terpenoids, have antibacterial and antibiofilm effects against *S. mutans* and *P. gingivalis*. The most effective concentration is 100 µg/mL for both bacteria with 87.15% and 97.73% of biofilms reduction for *S. mutans* and *P. gingivalis*, respectively. *L. barbarum* L. might be a promising natural-therapeutic agent as an alternative therapy. However, further research using toxicity, preclinical, and clinical tests is needed to determine whether *L. barbarum* L. fruit ethanol extract can be used as alternative mouthwash for preventing caries and treating chronic periodontitis.

## ACKNOWLEDGMENT

The authors thank Faculty of Dentistry, Trisakti University, for invaluable support in this study. Authors also

would like to thank Mario Richi, S.Si from the Microbiology Center of Research and Education (MiCORE) laboratory and Sheila Sutanto, S.Si from Biological Collaborative Research and Education (BiCORE) laboratory for their laboratory assistances.

#### REFERENCES

- [1] F. Katge, B. Rusawat, A. Shitoot, M. Poojari, T. Pammi, and D. Patil, "DMFT index assessment, plaque pH, and microbiological analysis in children with special health care needs, India," *Journal of International Society of Preventive and Community Dentistry*, vol. 5, pp. 383–388, 2015
- [2] Ministry of Health of the Republic of Indonesia. "Report on result of national basic health research (Riskesdas) 2018", Jakarta: Ministry of Health of the Republic of Indonesia pp. 1–582, 2018
- [3] C. Heng, "Tooth decay is the most prevalent disease," *Federal Practitioner*, vol. 33(10), pp. 31–33, 2016
- [4] S. Chenicheri, U. R. R. Ramachandran, V. Thomas, and A. Wood, "Insight into oral biofilm: primary, secondary and residual caries and phyto-challenged solutions," *Open Dentistry Journal*, vol. 11, pp. 312–333, 2017
- [5] G. Conrads and I. About, "Pathophysiology of dental caries," *Monograph in Oral Science*, vol. 27, pp. 1–10, 2018
- [6] J. E. Hinrichs and G. Kotsakis, "Classification of diseases and conditions affecting the periodontium," in *Carranza*, 2015, pp. 51–55.
- [7] S. Kumar and J. Sengupta, "Endodontic treatment for mandibular molars using ProTaper," *Medical Journal Armed Forces India*, vol. 67, no. 4, pp. 377–379, 2011
- [8] D. Sharma, L. Misba, and A. U. Khan. "Antibiotics versus biofilm: an emerging battleground in microbial communities," *Antimicrobial Resistance and Infection Control*, vol. 8, pp. 1–10, 2019
- [9] A. S. K. J. Arul and P. Palanivelu. "Biofilm forming ability of a new bacterial isolate from dental caries: An atomic force microscopic study," *Journal of Natural Science, Biology, and Medicine*, vol. 5, no. 2, pp. 278–283, 2014
- [10] J. Abranches, L. Zeng, J. K. Kajfász, S. R. Palmer, B. Chakraborty, Z. T. Wen et al., "Biology of oral Streptococci," *Microbiology Spectrum*, vol. 6, no. 5, 2018
- [11] K. Bao, G. N. Belibasakis, T. Thumheer, J. Aduse- Opoku, M. A. Curtis, and N. Bostanci, "Role of Porphyromonas gingivalis gingipains in multi-species biofilm formation," *BioMed Central Microbiology*, vol. 14, p. 258, 2014
- [12] L. Jia, N. Han, J. Du, L. Guo, Z. Luo, and Y. Liu, "Pathogenesis of Important Virulence Factors of Porphyromonas gingivalis via Toll-Like Receptors," *Frontier in Cellular and Infection Microbiology*, vol. 9, p. 262, 2019
- [13] Z. L. S. Brukes, R. Bescos, L. A. Belfield, K. Ali, and A. Robert, "Current uses of chlorhexidine for management of oral disease: a narrative review," *Journal of Dentistry*, vol. 103, p. 103497, 2020
- [14] N. Jeddy, S. Ravi, T. Radhika, and L. J. S. Lakshmi, "Comparison of the efficacy of herbal mouth rinse with commercially available mouth rinses: A clinical trial," *Journal of Oral Maxillofacial Pathology*, vol. 22, no. 3, pp. 332–334, 2018
- [15] C. C. Martínez, M. D. Gómez, and M. S. Oh. "Use of traditional herbal medicine as an alternative in dental treatment in mexican dentistry: A review," *Pharmaceutical Biology*, vol. 55 (1), pp. 1992–1998, 2017
- [16] World Health Organization. "WHO traditional medicine strategy: 2014-2023", Geneva: World Health Organization, 2013; Available from: [https://www.who.int/medicines/publications/traditional/tm\\_strategy14\\_23/en/](https://www.who.int/medicines/publications/traditional/tm_strategy14_23/en/)
- [17] Z. F. Ma, H. Zhang, S. S. Teh, C. W. Wang, Y. Zhang, F. Hayford et al., "Goji berries as a potential natural antioxidant medicine: An insight into their molecular mechanisms of action," *Oxidative Medicine and Cellular Longevity*, p. e2437397, 2019
- [18] I. J. Alassadi, F. S. Sabah, and L. A. Alrubaie. "Isolation of flavonoid compound from iraqi awsaj plant (Lycium Barbarum L.) Fruits and the Study of Its Antibacterial Activity," *European Scientific Journal*, vol. 11, no. 24, pp. 268–276, 2015
- [19] S-E. Byambasuren, J. Wang, and G. Gaudel, "Medicinal value of wolfberry (Lycium barbarum L.)," *Journal of Medical Plants Studies*, vol. 7, no. 4, pp. 90–97, 2019
- [20] Z. F. Ma, H. Zhang, S. S. Teh, C. W. Wang, Y. Zhang, F. Hayford et al., "Goji berries as a potential natural antioxidant medicine: An insight into their molecular mechanisms of action," *Hindawi*, pp. 1–9, 2019
- [21] P. Skenderidis, D. Lampakis, I. Giavasis, S. Leontopoulos, K. Petrotos, C. Hadjichristodoulou et al., "Chemical properties, fatty-acid composition, and antioxidant activity of Goji berry (Lycium barbarum L. and Lycium chinese mill.) fruits," *Antioxidants*, vol. 8, no. 3, p. 60, 2019
- [22] P. Skenderidis, D. Lampakis, I. Giavasis, S. Leontopoulos, K. Petrotos, C. Hadjichristodoulou et al., "The in vitro antimicrobial activity assessment of ultrasound assisted Lycium barbarum fruit extracts and pomegranate fruit peels," *Journal of Food Measurement and Characterization*, vol. 13, no 5, pp. 1–15, 2019
- [23] Z. Y. Dewi, A. Nur, and T. Hertriani, "Efek antibakteri dan penghambatan biofilm ekstrak serih (Cymbopogon nardus L.) terhadap bakteri Streptococcus mutans," *Majalah Kedokteran Gigi Indonesia*, vol 1, no. 2, p. 136, 2015
- [24] S. Loresta, S. Murwani, and P. Trisunuwati "Efek ekstrak etanol daun kelor (Moringa oleifera) terhadap pembentukan biofilm Staphylococcus aureus secara in vitro, pp. 20–25, Available from: <http://library1.nida.ac.th/tempaper6/sd/2554/19755.pdf>
- [25] S. Bontjura, O. A. Waworuntu, and K. V. Siagia. "Uji efek antibakteri ekstrak daun leilem (Clerodendrum Minahassae L.) terhadap bakteri Streptococcus mutans," *Pharmakon*, vol. 4, no. 4, p. 6, 2015
- [26] L. Bouarab-Chibane, V. Forquet, P. Lanteri, Y. Clément, L. Léonard-Akkari, N. Oulahal et al., "Antibacterial properties of polyphenols: characterization and QSAR (quantitative structure-activity relationship) models," *Frontiers in Microbiology*, vol. 10, p. 829, 2019
- [27] Y. Mulay, "Phytochemical analysis and antibacterial properties of some selected Indian medicinal plants," *International Journal of Current Microbiology and Applied Sciences*, pp. 228–235, 2015
- [28] A. Ludwiczuk, K. Skalicka-Woźniak, and M. I. Georgiev, "Terpenoids," in *Pharmacognosy: Fundamentals, Applications and Strategy*, p. 251, 2017
- [29] N. I. Fit, F. Chirilă, G. Nadăș, E. Pall, and R. Mureșan, "Comparative testing of antimicrobial activity of aqueous extracts of Aloe vera and Lycium barbarum," *Bulletin of University of Agricultural Sciences and Veterinary Medicine of Cluj*, vol. 70, no. 1, pp. 72–76, 2013
- [30] A. S. Widyarman and N. K. E. Lazaroni, "Persistent endodontics pathogens biofilm inhibited by Lactobacillus reuteri Indonesian strain," *Journal of Dentistry Indonesia*, vol. 26, no. 3, pp. 160–164, 2019
- [31] L. Lu, W. Hu, Z. Tian, D. Yuan, G. Yi, Y. Zhou Y et al., "Developing natural products as potential anti-biofilm agents," *Chinese Medicine*, p. 11, 2019

# Goji Berry Extract (*Lycium barbarum* L.) Efficacy on Oral Pathogen Biofilms

## ORIGINALITY REPORT

9%

SIMILARITY INDEX

10%

INTERNET SOURCES

8%

PUBLICATIONS

9%

STUDENT PAPERS

## PRIMARY SOURCES

1

[www.nature.com](http://www.nature.com)

Internet Source

3%

2

Submitted to National University of Singapore

Student Paper

2%

3

Submitted to Coventry University

Student Paper

2%

4

[www.contempclindent.org](http://www.contempclindent.org)

Internet Source

2%

Exclude quotes  On

Exclude bibliography  On

Exclude matches  < 2%



# Goji Berry Extract (*Lycium barbarum* L.) Efficacy on Oral Pathogen Biofilms

---

GRADEMARK REPORT

---

FINAL GRADE

GENERAL COMMENTS

**/0**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

mail.yahoo.com/d/folders/70/messages/17187?guce\_referrer=aHR0cHM6Ly9sb2dpbi55VWhvby5jb20v&guce\_referrer\_sig...

USAKTI - SIS SIMPPM SS

HOME MAIL NEWS FINANCE SPORT CELEBRITY STYLE WEATHER MORE... y!mail+ Upgrade now

yahoo!mail Inhence Search in inhence... Advanced

Compose

Back Archive Move Delete Spam

[InHence 2021] Setting up your EDAS account password

From: inhence2021\_chairs@edes.info  
To: Sheila Soesanto

Sat, 19 Jun 2021 at 02:14

Dear Sheila Soesanto,

An EDAS publication management account has been created for you for one of the following reasons:

- you are a co-author of a paper;
- you are a technical program committee member;
- you will be asked to review a paper;
- you will be asked to chair a session.

The account was created for InHence 2021 by Armelia Sari Widayaman with the information:  
Faculty of Dentistry, Universitas Trisakti  
Indonesia

Your EDAS user name is mass9977@yahoo.com, your EDAS ID number 1874015  
Your initial password can be set at  
<https://www.edas.info/5ef0e49e0f0a1132a0baa560151e092>  
You can log in at <https://edas.info>

Your EDAS account can be used for all EDAS-managed conferences and journals. You should not create a new account for each conference.

Regards,  
The EDAS manager [help@edas.info](mailto:help@edas.info)

Folders: Hide  
New folder  
Alyssa (8)  
Alyssa\_art (4)  
ariel (1)  
Goethe  
etc  
Inhence  
kampus  
Karate  
Lampiris  
MAHENDRA IND... (1)  
PDGI  
Relathan Biomec... (2)  
PKM  
sung  
Thesis

mail.yahoo.com/d/folders/70/messages/17188?guce\_referrer=aHR0cHM6Ly9sb2dpbi55VWhvby5jb20v&guce\_referrer\_sig...

USAKTI - SIS SIMPPM SS

HOME MAIL NEWS FINANCE SPORT CELEBRITY STYLE WEATHER MORE... y!mail+ Upgrade now

yahoo!mail Inhence Search in inhence... Advanced

Compose

Back Archive Move Delete Spam

[InHence 2021] Information about paper #1570736359 (Antibiofilm Efficacy of Goji Berry Ethanol Extract (Lycium Barbarum L) on Streptococcus Mutans and Porphyromonas Gingivalis Biofilms) has been changed

From: inhence2021\_chairs@edes.info  
To: Armelia Sari Widayaman, Sheila Soesanto

Sat, 19 Jun 2021 at 09:14

Dear Dr. Armelia Widayaman,

Information about your paper #1570736359 (Antibiofilm Efficacy of Goji Berry Ethanol Extract (Lycium Barbarum L) on Streptococcus Mutans and Porphyromonas Gingivalis Biofilms) for InHence 2021 was changed by Armelia Sari Widayaman ():

Sheila Soesanto added as author

No further action is required from you.

If you have already submitted your manuscript, you can change it at any time before the deadline by [web form upload](#).

You can [view all your submissions](#), using the EDAS user name. From there, you can see the current status of the papers, whether a manuscript has been submitted and can edit the paper information.

You can [directly view information about your paper](#).

Once you update your manuscript, you will receive another email confirmation.

Regards: The conference chairs

Folders: Hide  
New folder  
Alyssa (8)  
Alyssa\_art (4)  
ariel (1)  
Goethe  
etc  
Inhence  
kampus  
Karate  
Lampiris  
MAHENDRA IND... (1)  
PDGI  
Relathan Biomec... (2)  
PKM  
sung

Browser tabs: Fw: [InHeNce 2021] Your paper #1... (15 unread) - mass9977@yahoo.com

Address bar: mail.yahoo.com/d/folders/70/messages/17189?guce\_referrer=aHR0cHM6Ly9sb2dpbi55YWhvby5jb20v&guce\_referrer\_sig...

Navigation: HOME MAIL NEWS FINANCE SPORT CELEBRITY STYLE WEATHER MORE... y!mail Upgrade now

Search: Inhence Search in inhence... Advanced

Compose

Folders: Hide  
New folder  
Alyssa  
Alyssa\_art  
ariel  
Goethe  
etc  
Inhence  
kampus  
Karate  
Lampikes  
MAHENDRA IN...  
FDGI  
Rebahan Biomo...  
PKM  
sung

Back Archive Move Delete Spam

[InHeNce 2021] Information about paper #1570736359 (Antibiofilm Efficacy of Goji Berry Ethanol Extract (Lycium Barbarum L.) on Streptococcus Mutans and Porphyromonas Gingivalis Biofilms) has been changed. Sheila/Inhence

From: inhence2021-chairs@edas.info  
To: Armelia Sari Widayman, Sheila Soesanto, Ricky Soen

Sat, 19 Jun 2021 at 09:16

Dear Dr. Armelia Widayman:

Information about your paper #1570736359 (Antibiofilm Efficacy of Goji Berry Ethanol Extract (Lycium Barbarum L.) on Streptococcus Mutans and Porphyromonas Gingivalis Biofilms) for InHeNce 2021 was changed by Armelia Sari Widayman ():

Ricky Soen added as author

No further action is required from you.

If you have already submitted your manuscript, you can change it at any time before the deadline by [web form upload](#).

You can [see all your submissions](#), using the EDAS user name. From there, you can see the current status of the papers, whether a manuscript has been submitted and can edit the paper information.

You can [directly view information about your paper](#).

Once you update your manuscript, you will receive another email confirmation.

Regards: The conference chairs

Browser tabs: Fw: [InHeNce 2021] Your paper #1... (15 unread) - mass9977@yahoo.com

Address bar: mail.yahoo.com/d/folders/70/messages/17190?guce\_referrer=aHR0cHM6Ly9sb2dpbi55YWhvby5jb20v&guce\_referrer\_sig...

Navigation: HOME MAIL NEWS FINANCE SPORT CELEBRITY STYLE WEATHER MORE... y!mail Upgrade now

Search: Inhence Search in inhence... Advanced

Compose

Folders: Hide  
New folder  
Alyssa  
Alyssa\_art  
ariel  
Goethe  
etc  
Inhence  
kampus  
Karate  
Lampikes  
MAHENDRA IN...  
FDGI  
Rebahan Biomo...  
PKM  
sung  
Thesis

Back Archive Move Delete Spam

[InHeNce 2021] Information about paper #1570736359 (Antibiofilm Efficacy of Goji Berry Ethanol Extract (Lycium Barbarum L.) on Streptococcus Mutans and Porphyromonas Gingivalis Biofilms) has been changed. Sheila/Inhence

From: inhence2021-chairs@edas.info  
To: Armelia Sari Widayman, Sheila Soesanto, Ricky Soen, Reynaida Oktavianri

Sat, 19 Jun 2021 at 09:17

Dear Dr. Armelia Widayman:

Information about your paper #1570736359 (Antibiofilm Efficacy of Goji Berry Ethanol Extract (Lycium Barbarum L.) on Streptococcus Mutans and Porphyromonas Gingivalis Biofilms) for InHeNce 2021 was changed by Armelia Sari Widayman ():

Reynaida Oktavianri added as author

No further action is required from you.

If you have already submitted your manuscript, you can change it at any time before the deadline by [web form upload](#).

You can [see all your submissions](#), using the EDAS user name. From there, you can see the current status of the papers, whether a manuscript has been submitted and can edit the paper information.

You can [directly view information about your paper](#).

Once you update your manuscript, you will receive another email confirmation.

Regards: The conference chairs

Browser tabs: Fw: [InHeNce 2021] Your paper #1 (15 unread) - mass9977@yahoo.com

Address bar: mail.yahoo.com/d/folders/70/messages/17310?guce\_referrer=aHR0cHM6Ly9sb2dpbi55YWhvby5jb20v&guce\_referrer\_sig=AQ...

Navigation: HOME MAIL NEWS FINANCE SPORT CELEBRITY STYLE WEATHER MORE... Upgrade now

Search: Inhence x Search in inhence... Advanced

Left sidebar: Compose, ariel (1), Goethe, etc, Inhence, kampus, Karate, Lamptkes, MAHENDRA..., PDGI, Pelatihan Bio..., PKM, sung, Thesis

Message header:
 

- Sent: Tue, 29 Jun 2021 at 12:17
- Subject: [InHeNce 2021] Your paper #1570736359 (Antibiofilm Efficacy of Goji Berry Ethanol Extract (Lyclum Barbarum L.) on Streptococcus Mutans and Porphyromonas Gingivalis Biofilms)

Dear Dr. Sheila Soesanto,

Congratulations - your paper #1570736359 (Antibiofilm Efficacy of Goji Berry Ethanol Extract (Lyclum Barbarum L.) on Streptococcus Mutans and Porphyromonas Gingivalis Biofilms) for InHeNce 2021 has been accepted and will be presented in the session titled ...

The reviews are below or can be found at <https://edas.info/showPaper.php?m=1570736359>.

**Review 1**

Originality: New or Novel contribution  
Accept (8)

Significance of Topic: Relating to knowledge contribution  
Accept (8)

Presentation: Clarity and Organisation of Content  
Accept (8)

Browser tabs: Fw: [InHeNce 2021] Your paper #1 (15 unread) - mass9977@yahoo.com

Address bar: mail.yahoo.com/d/folders/70/messages/17310?guce\_referrer=aHR0cHM6Ly9sb2dpbi55YWhvby5jb20v&guce\_referrer\_sig=AQ...

Navigation: HOME MAIL NEWS FINANCE SPORT CELEBRITY STYLE WEATHER MORE... Upgrade now

Search: Inhence x Search in inhence... Advanced

Left sidebar: Compose, ariel (1), Goethe, etc, Inhence, kampus, Karate, Lamptkes, MAHENDRA..., PDGI, Pelatihan Bio..., PKM, sung, Thesis

Message content:
 

- Accept (8)
- Recommendation: Overall view and recommendation**  
Accept (8)
- Strengths/Weakness: What are the major reasons to accept/reject the paper? [Be brief]**

The manuscript contains all part of the article such as title, abstract, introduction, materials and methods, results, discussion, conclusion and references. The manuscript's topic sounds interesting but the authors need to revise the article. It is better if the article is provided an English editing service.

**Contributions & Detailed comments: What are the major issues addressed in the paper? Do you consider them important? Comment on the degree of novelty, creativity and technical depth in the paper. Please provide detailed comments that will be helpful to the TPC for assessing the paper, as well as feedback to the authors.**

- Please delete the IEEE template in the affiliation part
- The title of the table should be capital
- Please increase the size of the word in the table to be 8 pt
- Please explain the base of the dosage that the author choose in this paper
- It's beter if the author makes 2 figures in fig. 3 to be one (not separate) so the reader will be easy understand
- In the conclusion part, please show wwhich dose is most appropriate to use based on research

**Review 2**

Originality: New or Novel contribution

Browser tabs: Fw: [InHence 2021] your paper #1 | (15 unread) - mass9977@yahoo.com

Address bar: mail.yahoo.com/d/folders/70?guce\_referrer=aHR0cHM6Ly9sb2dpbi55YWVhby5jb20v&guce\_referrer\_sig=AQAAAFwIABh927iPL...

Navigation: HOME MAIL NEWS FINANCE SPORT CELEBRITY STYLE WEATHER MORE... Upgrade now

Search: Inhence x Search in inhence... Advanced

Compose

- ▼ ariel (1)
- Goethe
- etc
- Inhence
- kampus
- Karate
- Lampitkes
- MAHENDRA ... (1)
- PDGI
- pelatihan Bio... (2)
- PKM
- sung
- Thesis

Actions: Back, Archive, Move, Delete, Spam

Weak Accept (6)

**Significance of Topic: Relating to knowledge contribution**

Weak Accept (6)

**Presentation: Clarity and Organisation of Content**

Neutral (5)

**Recommendation: Overall view and recommendation**

Weak Accept (7)

**Strengths/Weakness: What are the major reasons to accept/reject the paper? [Be brief.]**

this study was to determine antibacterial and antibiofilm effects of goji berry (*L. barbarum* L.) ethanol extract towards *S. mutans* and *P. gingivalis*. This study is interesting. However please revise as follows

**Contributions & Detailed comments: What are the major issues addressed in the paper? Do you consider them important? Comment on the degree of novelty, creativity and technical depth in the paper. Please provide detailed comments that will be helpful to the TPC for assessing the paper, as well as feedback to the authors.**

1. Please make the authorship according to the template, one author one column
2. English needs to be rechecked

Browser tabs: Fw: [InHence 2021] your paper #1 | (15 unread) - mass9977@yahoo.com

Address bar: mail.yahoo.com/d/folders/70?guce\_referrer=aHR0cHM6Ly9sb2dpbi55YWVhby5jb20v&guce\_referrer\_sig=AQAAAFwIABh927iPL...

Navigation: HOME MAIL NEWS FINANCE SPORT CELEBRITY STYLE WEATHER MORE... Upgrade now

Search: Inhence x Search in inhence... Advanced

Compose

- ▼ ariel (1)
- Goethe
- etc
- Inhence
- kampus
- Karate
- Lampitkes
- MAHENDRA ... (1)
- PDGI
- pelatihan Bio... (2)
- PKM
- sung
- Thesis

Actions: Back, Archive, Move, Delete, Spam

**Contributions & Detailed comments: What are the major issues addressed in the paper? Do you consider them important? Comment on the degree of novelty, creativity and technical depth in the paper. Please provide detailed comments that will be helpful to the TPC for assessing the paper, as well as feedback to the authors.**

1. Please make the authorship according to the template, one author one column
2. English needs to be rechecked
3. describe the implication of the study
4. Title is too long, please have 15 words.
5. Tables and figures need to follow the template.
6. Make the most of space in the paper

Regards,  
The conference chairs

Navigation: Back, Forward, Reply, Reply all or Forward

mail.yahoo.com/d/folders/70/messages/17360?gucereferer=aHR0cHM6Ly9sb2dpbi55YWhvby5jb20v&gucereferer\_sl...

USAKTI - SIS SIMPPM SS

HOME MAIL NEWS FINANCE SPORT CELEBRITY STYLE WEATHER MORE... Upgrade now

yahoo/mail Inhence Search in inhence... Advanced

Compose

- ariel
- Goethe
- etc
- Inhence
- kampus
- Karate
- Lampukes
- MAHENDRA ...
- PDGI
- Pelatihan Bio...
- PKM
- sung
- Thesis

Back Archive Move Delete Spam

has been changed

inhence2021-chairs@edas.info Fri, 2 Jul 2021 at 15:16

From: inhence2021-chairs@edas.info  
To: Sheila Soesanto, Reynalda Oktaviani, Ricky Soen, Armelia Sari Widjarmen

Dear Dr. Sheila Soesanto:

Information about your paper #1570736359 (Antibiofilm Efficacy of Gaji Berry Ethanol Extract (*Lycium Barbarum L.*) on *Streptococcus Mutans* and *Porphyromonas Gingivalis* Biofilms) for InHeNce 2021 was changed by Refi Ikhtilari ()

Track changed from Health to Natural Sciences.

No further action is required from you.

If you have already submitted your manuscript, you can change it at any time before the deadline by [web form upload](#)

You can [see all your submissions](#) using the EDAS user name mass9977@yahoo.com. From there, you can see the current status of the papers, whether a manuscript has been submitted and can edit the paper information.

You can [directly view information](#) about your [paper](#)

Once you update your manuscript, you will receive another email confirmation.

Regards, The conference chairs

mail.yahoo.com/d/folders/70/messages/17430?gucereferer=aHR0cHM6Ly9sb2dpbi55YWhvby5jb20v&gucereferer\_slg=AQ...

USAKTI - SIS SIMPPM SS

HOME MAIL NEWS FINANCE SPORT CELEBRITY STYLE WEATHER MORE... Upgrade now

yahoo/mail Inhence Search in inhence... Advanced

Compose

- ariel
- Goethe
- etc
- Inhence
- kampus
- Karate
- Lampukes
- MAHENDRA ...
- PDGI
- Pelatihan Bio...
- PKM
- sung
- Thesis

Back Archive Move Delete Spam

[[InHeNce 2021] #1570741082 has been uploaded

Sheila/inhence Wed, 7 Jul 2021 at 22:57

inhence2021-chairs@edas.info

From: inhence2021-chairs@edas.info  
To: Sheila Soesanto, Reynalda Oktaviani, Ricky Soen, Armelia Sari Widjarmen

Dear Mr. Frans Samosir:

Thank you for uploading your paper 1570741082 (*Antibiofilm Efficacy of Gaji Berry Ethanol Extract (Lycium Barbarum L.) on Streptococcus Mutans and Porphyromonas Gingivalis Biofilms*) to 2021 IEEE International Conference on Health, Instrumentation & Measurement, and Natural Sciences (InHeNce). The paper is of type application/vnd.openxmlformats-officedocument.wordprocessingml.document and has a length of 1245012 bytes.

You can modify your paper at <https://edas.info/showPaper.php?m=1570741082> and see all your submissions at <https://edas.info/index.php?c=28172> using the EDAS identifier fransjudeasamosir@unprindn.ac.id

Regards,  
The conference chairs

Browser tabs: Fw: [inHence 2021] your paper #1 | [15 unread] - mass9977@yahoo.com

Address bar: mail.yahoo.com/d/folders/70/messages/17790?guce\_referrer=aHR0cHM6Ly9sb2dpbi55YWhvby5jb20v&guce\_referrer\_sig=AQ...

Navigation: HOME MAIL NEWS FINANCE SPORT CELEBRITY STYLE WEATHER MORE...

Search: Inhence Search in inhence... Advanced

Compose

- aniel
- Goethe
- etc
- Inhence**
- kampus
- Karate
- Lampfkes
- MAHENDRA ...
- PDGI
- Pelatihan Bio...
- PKM
- sung
- Thesis

Actions: Back Archive Move Delete Spam

Message: INHENCE FINAL REV 3 Sheila/Inhence

From: inhence@unprimdn.ac.id  
To: mass9977@yahoo.com, Ricky Chandra Jaya Soen, Armella Sari

Wed, 25 Aug 2021 at 22:47

Dear Author,

Please revise your FINAL manuscript in the attachment file (DOWNLOAD and REVISE) according to some feedback from Editor below:

1570736359-Sheila Soesanto	This paper has not been well revised such as no discussions but literature review.
----------------------------	--

Your paper has strong potential for publication according to IEEE after you revise as above. Please notice that you need to follow up and send back the manuscript to this email by 27 August 2021 at 23:59 WIB.

We thank you and hope to give you the best service.

InHence 2021  
[inhence@unprimdn.ac.id](mailto:inhence@unprimdn.ac.id)

## Effect of goji berry ethanol extract- (*Lycium barbarum* L.) on *Streptococcus mutans* and *Porphyromonas gingivalis* biofilms

### ABSTRACT

**Background:** Caries and periodontitis are commonly found in the Indonesian population. *Streptococcus mutans* and *Porphyromonas gingivalis* in the form of biofilms play a major role in causing caries and chronic periodontitis. Chlorhexidine mouthwash can be used to prevent and treat periodontitis; however, due to its many side effects, an alternative treatment, using natural ingredients that have antibacterial effects, is needed. *Lycium barbarum* L. fruit, which contains flavonoids and phenolic acids, has antibacterial properties that are expected to inhibit bacterial growth and the formation of *S. mutans* and *P. gingivalis* biofilms. **Objective:** To determine the antibacterial and antibiofilm effects of *L. barbarum*-fruit ethanol extract against *S. mutans* and *P. gingivalis*. **Methods:** An in vitro laboratory experiment was performed with a post-test control group design. The extract of *L. barbarum*-fruit was obtained by maceration using 96% ethanol as a solvent. The test solutions were *L. barbarum*-fruit ethanol extract at a concentration of 100%, 50%, 25%, 12.5%, and 6.25%, chlorhexidine gluconate 0.2% as a positive control, and sterile distilled water as a negative control. The antibacterial assay was performed by microdilution and plate count methods. The antibiofilm effect was performed using a biofilm assay method. **Result:** The results of the microdilution and plate count methods showed that the most effective concentration with antibacterial properties against *S. mutans* and *P. gingivalis* was 100% when compared with the negative control ( $p < 0.05$ ). In the biofilm assay, the most effective concentration against *S. mutans* was 100% at the 3-hour incubation time, while for *P. gingivalis*, the most effective concentration was 100% at the 24-hour incubation time when compared with the negative control ( $p < 0.05$ ). **Conclusion:** The ethanol extract of *L. barbarum*-fruit was demonstrated to have antibacterial and antibiofilm effects against *S. mutans* and *P. gingivalis*.

Keyword:



## 1. Introduction

According to the World Health Organization (WHO), optimal dental and oral health is free of dental caries, periodontal disease, oral cancer, infections and sores in the mouth, ~~noma~~ cleft lip and palate, tooth decay, tooth loss, and diseases that cause biting disorders, all of which negatively impact chewing, smiling, talking, and psychosocial well-being (WHO, 2020). Oral health positively affects the appearance, as well as the physical, mental, and interpersonal well-being, of an individual. Thus, oral health, which is part of overall health, contributes to quality of life (Katge et al., 2015).

Formatted: Indent: First line: 0"

Basic Health Research Data (Rikesdas Rikesdas, 2018) shows that 57.6% of the Indonesian population experiences dental and oral health problems. The prevalence of dental caries in Indonesia in 2018 was 88.8%, with an average DMF-T index of 7.1, which is a very high severity of dental caries. Periodontitis is experienced by 74.1% of the Indonesian population (Kemenkes, 2018).

Commented [.1]: How do these numbers compare to other regions?

Caries is the process of the demineralization of inorganic material and the dissolution of organic material, leading to bacterial invasion through the dentin layer until it reaches the pulp (Heng, 2016; Chenicheri et al., 2017). The process of dental caries depends on the presence of fermentable sugars (substrates), the type of tooth and saliva (host), cariogenic microbial flora (biofilm), and time (Conrads and About, 2018). Periodontitis is a disease caused by inflammation of the tooth-supporting tissue, caused by microorganisms, that causes progressive destruction of the periodontal ligament and alveolar bone, with the formation of pockets, recessions, or both (Hinrichs and Kotsakis, 2015). Periodontitis in adults is caused by numerous local factors, such as biofilms or calculus, classified as chronic periodontitis (Kumar and Sengupta, 2011).

Formatted: Indent: First line: 0"

The formation of biofilms begins when microorganisms in the planktonic state merge into bacterial colonies and wrap themselves in a self-produced extracellular polymer matrix (Ollie Yira-Yu et al., 2017). At the beginning of the formation of biofilms, there is an increase in the activity of

Formatted: Indent: First line: 0"

Commented [.2]: Please check that only the author's surname is being used here.

~~G~~ram-positive *cocci*, one of which is *Streptococcus mutans*, which is able to adhere to the tooth surface through the formation of extracellular polysaccharides that cause the biofilm matrix to have a gelatin-like consistency that facilitates the attachment of bacteria to the tooth's surface (Arul and Palanivelu, 2014; Chenicheri et al., 2017; Abranches et al., 2018). *Porphyromonas gingivalis*, which is a secondary bacterium, is an anaerobic ~~G~~ram-negative bacterium found in periodontal pockets that causes chronic periodontitis. ~~Various virulence factor of P. gingivalis such as gingipains, fimbriae, and lipopolysaccharides,~~ plays an important role in ~~regulating virulence,~~ ~~such as gingipains, fimbriae, and lipopolysaccharides, which furthers~~ the development of periodontal disease and induces dysbiosis in biofilms (Bao et al., 2014; Mysak et al., 2014).

Chlorhexidine mouthwash is used to prevent caries and treat periodontitis and is considered the gold standard for controlling dental plaque and gingivitis due to its efficacy against a wide variety of bacteria, fungi, and viruses. However, chlorhexidine has various side effects, including taste disturbances, discoloration of the teeth and mucosa, mucosal desquamation, salivary stone formation, irritation, dry oral cavity, and allergic reactions, such as contact stomatitis. The WHO recommends finding new natural ingredients to overcome the side effects of chemical agents (Rezaei et al., 2016; Jeddy et al., 2018).

The use of natural ingredients as antimicrobial agents has become an alternative because of their low cost and lower toxicity (Martinez et al., 2017). Various herbal mouthwashes that have been ~~tested successfully used by the community~~ include mimba (*Azadirachta indica*), aloe vera (*Aloe perfoliata* var. *Vera L.*), and tea tree oil (*Melaleuca alternifolia*) (Manipal et al., 2016).

Goji berry (*Lycium barbarum L.*) ~~(L.)~~ has been widely used as a traditional medicine by people in Asia, especially in the northwestern part of China, for more than 2000 years. Recently, *L. barbarum* has been gaining popularity ~~as and is referred to as a superfruit, which is a~~ highly nutritious food used to improve health in North America, Europe, and Asia (Ma et al., 2019). *L. barbarum* has a red, oblong fruit with a length of 6–20 mm and a diameter of 3–10 mm. ~~*L. yeium barbarum L.*~~ fruit is harvested when it is ripe and is then dried for later use (Alassadi-Fatima-Sabah and Alrubaie et al., 2015). The fruit, roots, tree bark, and flowers of *L. barbarum* have been shown to be used as medicine (Byambasuren et al., 2019).

**Commented [.3]:** According to journal guidelines, when multiple sources are used in an in-text citation, they should be ordered chronologically. This has been adjusted here and throughout, where applicable.

**Formatted:** Field Code Changed

**Commented [.4]:** What does this bacterium regulate the virulence of? Your meaning here is not quite clear. Please clarify.

**Commented [.5]:** A parenthetical citation should be given before the period, when being placed at the end of the sentence.

**Formatted:** Indent: First line: 0"

**Formatted:** Indent: First line: 0"

**Commented [.6]:** According to journal guidelines, when a source has 3 or more authors, the first author's surname should be used followed by et al. in in-text citations.

**Commented [.7]:** Are these specific to Indonesia? Additionally, who is considered 'the community' in this sentence.

**Commented [.8]:** The L after the scientific name only needs to be used in its first appearance in the abstract and main text.

**Formatted:** Indent: First line: 0"

**Formatted:** Font: Not Italic

**Commented [.9]:** Double quotation marks should only be used for direct quotes or titles. Use single quotation marks for emphasis.

**Commented [.10]:** Please confirm that only the authors' surnames are being used in this citation.

**Commented [.11]:** The full genus name only needs to be provided upon first mention in the abstract, main text, and in each table and figure legend. Thereafter, the abbreviation can be used.

The polysaccharides of *L. barbarum* have exhibited properties that improve eye health and reproductive system health, reduce fat and blood sugar, and regulate immunity; they have also been shown to have anticancer, anti-tumor, antioxidant, anti-fatigue, antiviral, anti-aging, hepatoprotective, neuroprotective, and cardioprotective properties (Cheng et al., 2015; Ma et al., 2019). The flavonoids and phenolic acids of *L. barbarum* have potential as antioxidants and antimicrobials (Skenderidis, Mitsagga, et al., 2019).

Formatted: Highlight

Commented [.12]: Please clarify which of these is the surname.

*Lycium barbarum* fruit has been shown to be effective in inhibiting gram-negative bacteria, such as *Escherichia coli*, and gram-positive bacteria, such as *Staphylococcus aureus* (Skenderidis, Mitsagga, et al., 2019). However, there have been no studies regarding the antibacterial effect of *L. barbarum* fruit against *S. mutans* and *P. gingivalis* as bacteria that cause caries and chronic periodontitis. Thus, the aim of this study was to determine the antibacterial and antibiofilm effects of goji berry (*L. barbarum*) ethanol extract against *S. mutans* and *P. gingivalis*.

Formatted: Indent: First line: 0"

Commented [.13]: Use commas around specific examples when they are not essential. For example, I like different kinds of cheese, such as sharp cheddar and feta, as well as other dairy products. In this sentence, 'such as sharp cheddar and feta' are not necessary for me to convey that I like different kinds of cheese.

Commented [.14]: Please clarify which of these is the surname.

## 2. Material and methods

### 2.1 Ethanol extract of *L. barbarum* fruit

Dried *L. barbarum* fruit (100 g) from Chinese medicine store "Lancar Jaya" at Teluk Gong Raya No. 43, Jakarta Utara (produced in Zhongning, Ningxia, China) was ground in a blender until it became a powder. It was then immersed in 96% ethanol with a ratio of 1:8 for 72 hours, stirring every 15 minutes. Furthermore, filtration was performed using Whatman No. 1 filter paper and evaporated with a rotary evaporator at 40°C temperature, a speed of 60 rpm, and a pressure of 20 atm so that a thick and solvent-free extract was obtained with a concentration of 100%. The extracts were then diluted using sterile distilled water until concentrations of 50, 25, 12.5, and 6.25% were obtained.

Commented [.15]: Presumably, different sources of the fruit could have different extract profiles. I recommend including the source of the fruit used in this study.

Formatted: Indent: First line: 0 ch

Formatted: Font: 12 pt

Formatted: Font: 12 pt

### 2.2 Phytochemical assay

Phytochemical assays were performed qualitatively to determine whether the ethanol extract of *L. barbarum* fruit contained flavonoids, phenols, quinones, steroids, terpenoids, and alkaloids.

Commented [.16]: This is not repeatable. Please provide more detail on how these assays were performed.

Formatted: Indent: First line: 0 ch

### 2.3 Bacterial cultures

*Streptococcus mutans* ATCC 25175 and *P. gingivalis* ATCC 33277 from MiCORE Laboratory, Faculty of Dentistry, Trisakti University, were cultured on BHI-B medium and incubated at 37°C for 24 hours in an anaerobic atmosphere. Furthermore, the absorbance measurements were performed to reach the McFarland standard of  $0.5 = 1.5 \times 10^8$  CFU/mL ( $OD_{600} = 0.132$ ).

Formatted: Indent: First line: 0 ch

Commented [.17]: Please provide the recipe for this medium or a reference to where the recipe can be obtained.

#### 2.4 Microdilution

Each well of a 96-well plate was distributed 100 µL of either an *S. mutans* or *P. gingivalis* culture. Subsequently, 100 µL of the following solutions was used as a treatment: ethanol extracts of *L. barbarum* fruit at 100, 50, 25, 12.5, and 6.25% concentrations, 0.2% chlorhexidine gluconate as a positive control, and sterile distilled water as a negative control. The measurement of bacterial cell density was performed using a microplate reader at a 600 nm wavelength before and after the 96-well plates were incubated for 24 hours.

Formatted: Indent: First line: 0 ch

Commented [.18]: Presumably, each bacterium and extract was included separately in its own well. Please confirm that the intended meaning was maintained in this paragraph.

Commented [.19]: How many replications and runs of the experiment were there?

#### 2.5 Plate count

The microdiluted contents in the 96-well plates were re-diluted 10,000 times and cultured on BHI-A medium and incubated for 24 hours at 37°C to measure bacterial growth. The total bacterial number was calculated by the following formula:

Formatted: Indent: First line: 0 ch

#### 2.6 Biofilm assay

Bacterial culture (200 µL) was dispensed into each well of a 96-well plate and incubated at 37°C for 48 hours in an anaerobic atmosphere. Furthermore, the supernatant was removed until a thin layer of biofilm was left on the bottom surface of the well. The wells were rinsed with a solution of phosphate-buffered saline (PBS). The ethanol extracts of *L. barbarum* fruit at a concentration of 100, 50, 25, 12.5, and 6.25%, 0.2% chlorhexidine gluconate as a positive control, and sterile distilled water as a negative control were added 200 µL to the wells, as much as 200 µL, using a micropipette and incubated at 37°C for 1, 3, or 24 h in an anaerobic atmosphere. The well was rinsed twice using PBS and then fixated over a flame. Crystal violet dye (200 µL; 0.05% w/v) was added to each well and left for 15 minutes. The well was rinsed twice using PBS and left for 15 minutes. Then, 200 µL of 96% ethanol was inserted, and OD measurements were performed using a microplate reader at a 595 nm wavelength.

Commented [.20]: When a sentence starts with a number, the number should be spelled out; however, it is better to avoid starting a sentence with a number.

Commented [.21]: Please provide the exact volume of each solution that was added per well.

Commented [.22]: Please provide the exact volume of each solution that was added per well.

## 2.7 Statistical analyses

Statistical Product and Service Solution (SPSS) software version 25.0 was used to process the collected data. The Shapiro-Wilk method was used to test normality. Data that was normally distributed ( $P > 0.05$ ) was analyzed by a one-way analysis of variance (ANOVA) test. Significant data ( $P < 0.05$ ) were analyzed with a post-hoc test using Tukey's test ~~HSD~~ with a significance level of  $P < 0.05$  to determine which groups were significantly different.

Formatted: Indent: First line: 0 ch

Commented [.23]: All abbreviations should be defined the first time they appear in the main text of the manuscript.

## 3. Results

The phytochemical test qualitatively showed that the ethanol extract of *L. barbarum* fruit contained flavonoids, phenols, steroids, and terpenoids.

Formatted: Indent: First line: 0 ch

The results of this study indicated that the ethanol extract of *L. barbarum* fruit has antibacterial and antibiofilm effects against *S. mutans* and *P. gingivalis*. The ethanol extract of *L. barbarum* fruit with a concentration of 100% had the most effective antibacterial effect against *S. mutans* and *P. gingivalis*, with a total number of *S. mutans* colonies of  $3 \pm 3.46 \times 10^6$  CFU/mL (Figure 1) and an OD value of  $0.358 \pm 0.002$  (Figure 3). The total number of *P. gingivalis* colonies was  $41 \pm 4.58 \times 10^6$  CFU/mL (Figure 2), with an OD value of  $0.458 \pm 0.024$  (Figure 4).

Formatted: Indent: Left: 0", First line: 0 ch

Commented [.24]: This does not make sense because 3 - 3.46 would be a negative number. Also this number is much lower than that found for *P. gingivalis*. Please check that the reported number is correct.




Figure 1. Optical Density (OD) value of *Streptococcus mutans* based on the concentration of ethanol extract of *Lycium barbarum* fruit. \*\*Significant difference at  $P < 0.01$ .

Commented [.25]: All abbreviations should be defined anew in each table and figure caption.

Commented [.26]: Genus names should be spelled out upon first use in each table and figure caption.




Figure 2. Optical Density (OD) value of *Porphyromonas gingivalis* based on the concentration of ethanol extract of *Lycium barbarum* fruit. \*\*Significant difference at  $P < 0.01$ .

**Commented [.27]:** Consider combining Figures 1 and 2 into one figure. The same could be done for Figures 3 and 4, Figures 5, 6, and 7, and Figures 8, 9, and 10.

**Commented [.28]:** All abbreviations should be defined anew in each table and figure caption.

**Formatted:** Not Highlight

**Formatted:** Font: Not Italic




Figure 3. Total number of *Streptococcus mutans* based on the ethanol extract concentration of *Lycium barbarum* fruit. \*Significant difference at  $P < 0.05$ ; \*\*Significant difference at  $P < 0.01$ .

**Formatted:** Font: Not Italic




Figure 4. Total number of *Porphyromonas gingivalis* based on the concentration of ethanol extract of *Lycium barbarum* L. fruit. *P. gingivalis* based on the ethanol extract concentration of *L. barbarum* L. fruit. \*\*Significant difference at  $P < 0.01$ .

In the biofilm assay, the 100% concentration at a 3 h incubation time was the most effective in inhibiting the formation of *S. mutans* biofilm with an OD value of  $0.042 \pm 0.002$  (Figure 6), whereas for *P. gingivalis* biofilm, the 24 h incubation time at a concentration of 100% was the most effective (OD value:  $0.007 \pm 0.003$ ; Figure 10). Statistical analysis showed that all ethanol extract concentrations of *L. barbarum* fruit were significantly different from the negative control ( $P < 0.05$ ).

Formatted: Indent: First line: 0 ch




Figure 5. Optical Density (OD) value of *Streptococcus mutans* biofilm at a 1 h incubation time. \*\*Significant difference at P < 0.01.




Figure 6. Optical Density (OD) value of *Streptococcus mutans* biofilm at a 3 h incubation time. \*\*Significant difference at P < 0.01.






Figure 7. *Optical Density (OD) value of Streptococcus mutans* ~~OD value of *S. mutans*~~ biofilm at a 24 h incubation time. \*\*Significant difference at P < 0.01.

Formatted: Font: Italic




Figure 8. *Optical Density (OD) value of Porphyromonas gingivalis* ~~OD value of *P. gingivalis*~~ biofilm at a 1 h incubation time. \*\*Significant difference at P < 0.01.

Formatted: Font: Italic




Figure 9. Optical Density (OD) value of *Porphyromonas gingivalis* biofilm at a 3 h incubation time. \*Significant difference at  $P < 0.05$ ; \*\*Significant difference at  $P < 0.01$ .

Formatted: Font: Italic




Figure 10. Optical Density (OD) value of *Porphyromonas gingivalis* biofilm after 24 h of incubation. \*\*Significant difference at  $P < 0.01$ .

Formatted: Font: Italic

~~Statistical analysis showed that all ethanol extract concentrations of *L. barbarum* fruit were significantly different from the negative control ( $P < 0.05$ ).~~

**Commented [.29]:** Please include this sentence in another paragraph. Short paragraphs (1 or 2 sentences) can disrupt the flow of the manuscript.

**Formatted:** Indent: Left: 0", First line: 0 ch

#### 4. Discussion

The ethanol extract of *L. barbarum* fruit has various secondary metabolites, such as flavonoids, phenols, steroids, and terpenoids, which play a role in inhibiting bacterial growth and biofilm formation of *S. mutans* and *P. gingivalis*. Flavonoid compounds are able to damage bacterial cell walls by removing substances such as proteins, nucleic acids, and nucleotides so that bacterial cell lysis occurs (Dewi et al., 2015). Flavonoids can also interfere with the quorum sensing mechanism, causing inhibition of bacterial adhesion and biofilm formation on the tooth's surface. The formation of biofilms is inhibited by the reduction of glucans, which are a medium for bacterial attachment, due to the inactivity of the glucosyltransferase enzyme by flavonoids (Loresta Sonya Loresta and Sri Murwani et al., 2015).

**Formatted:** Indent: First line: 0 ch

**Commented [.30]:** Please confirm that only the authors' surnames are being used in this citation.

The ability of bacterial cell protein denaturation by phenol compounds through the formation of bonds between phenols and proteins causes damage to protein structures. The disruption of permeability in the cell wall and cytoplasmic membrane, which is composed of these proteins, causes irreversible damage and leads to lysis of the bacterial cell (Bontjura et al., 2015; Bouarab-Chibane et al., 2019).

**Formatted:** Indent: First line: 0 ch

**Field Code Changed**

Steroid compounds, through their interaction with cell phospholipid membranes, are also capable of causing lysosome leakage for the lysis of bacterial cells. Terpenoids are lipophilic and can bind to carbohydrates and fats, causing disruption of the permeability of bacterial cell walls, denaturation of cytoplasmic proteins, and inactivation of cellular enzymes, causing lysis of bacterial cells (Bontjura et al., 2015; Shinde and Mulay, 2015; Ludwiczuk et al., 2017).

**Field Code Changed**

The results of this study are in accordance with previous studies regarding the inhibition effect of *L. barbarum* extracts against *S. aureus* and *E. coli* by the disc diffusion method. Based on the results of these studies, there is an antibacterial effect against *E. coli* (Fiğ et al., 2013). The results of other studies using the well diffusion method have stated that the ethanol extract of *L. barbarum* fruit at concentrations of 10% and 20% had inhibitory effects against *S. aureus* and *E. coli* (Skenderidis et al., 2019).

**Formatted:** Indent: First line: 0 ch

**Commented [.31]:** Why was this different from your results? What factors could have contributed to these differences?

This study used three different incubation times in the biofilm assay, namely 1, 3, and 24 hours, to determine the most effective phase for inhibiting *S. mutans* and *P. gingivalis* biofilms. The difference in incubation time was similar to the biofilm formation phase, starting with the pellicle formation phase in a few minutes to 1 hour, the initial adhesion phase at 2 to 4 hours, and the maturation phase after 24 hours (Widyarman and Lazaroni, 2019).

Formatted: Indent: First line: 0 ch

The ethanol extract of *L. barbarum* fruit at a concentration of 100% was the most effective concentration in inhibiting *S. mutans* and *P. gingivalis* bacteria and biofilms. The results of this study are in accordance with the research of ~~Alassadi et al (2015) et al.~~ toward *L. barbarum* fruit, which showed that the presence of the alcohol group (-OH) in the flavonoid structure increased the ability of the extract to inhibit microbial growth by increasing the permeability of bacterial cell membranes, and the highest concentration, at 100%, was the most effective antibacterial compared to other concentrations, due to less flavonoid content at lower concentrations (Alassadi et al (Alassadi Fatima S Sabah and Alrubaie, 2015)).

Formatted: Indent: First line: 0 ch

Commented [.32]: When a narrative in-text citation is introduced, the year of publication should immediately follow the author's surname.

Formatted: Not Highlight

Formatted: Highlight

Commented [.33]: Please confirm that only the authors' surnames are being used in this citation.

The results of the antibiofilm assay showed that the most effective incubation time for inhibiting the formation of *S. mutans* biofilms was at 3 hours of incubation time, and for *P. gingivalis*, it was at a 24-hour incubation time. The most effective times for inhibiting biofilm formation were at the initial adhesion and maturation phases, respectively. The antibiofilm effect depends on the inhibition of the polymer matrix formation and quorum sensing, or communication, between bacterial cells in biofilms by inhibiting autoinducer peptides, signaling molecules in gram-positive bacteria, and acylhomoserine lactones (AHLs) in gram-negative bacteria so bacterial virulence factors and biofilm development may be inhibited (Lu et al., 2019).

Formatted: Indent: First line: 0 ch

This is proven by the lowest OD value found at a 100% concentration in *S. mutans* ( $0.042 \pm 0.002$ ) and *P. gingivalis* ( $0.007 \pm 0.003$ ). This antibiofilm assay also showed that a 100% concentration had a lower OD value and was significantly different from the positive control, which means that at a 100% concentration, the antibiofilm effect was more effective than the positive control.

## 5. Conclusions

The ethanol extract of *L. barbarum* fruit, containing flavonoids, phenols, steroids, and terpenoids, had antibacterial and antibiofilm effects against *S. mutans* and *P. gingivalis*. However, further research is needed, using toxicity, preclinical, and clinical tests, to determine if *L. barbarum* fruit ethanol extract can be used as an alternative mouthwash in preventing caries and treating chronic periodontitis.

## Reference

- Abranches, J., Zeng, L., Kajfasz, J.K., SR Palmer, S.R., Chakraborty, B., Wen, Z.T., Richards, V.P., Brady L.J., Lemos, J.A. 2018. Biology of Oral Streptococci. *Microbiol Spectr.* 6(5), GPP3-0042-2018. doi: 10.1128/microbiolspec.GPP3-0042-2018
- Alassadi, I. J., Fatima S-Sabah, F.S, I. J. and Al-Rubaie, L. A. (2015). Isolation of Flavonoid Compound From Iraqi Awsaj Plant (*Lycium Barbarum L.*) Fruits and the Study of Its Antibacterial Activity. *Eur. Sci. J.*; 11(24), p.268–276.
- Arul, A.S.K.J., Palanivelu, P. 2014. Biofilm forming ability of a new bacterial isolate from dental caries: An atomic force microscopic study. *J Nat Sci Biol Med.* 5(2), 278–283. doi: 10.4103/0976-9668.136162
- Bao, K., Belibasakis, G.N., Thurnheer, T., Opoku, J.A., Curtis, M.A., Bostanci, N. 2014. Role of *Porphyromonas gingivalis* gingipains in multi-species biofilm formation. *BMC Microbiol.* 14, 258. doi:10.1186/s12866-014-0258-7
- Bontjura, S., Waworuntu, O. A., and Siagian, K. V. (2015). Uji Efek Antibakteri Ekstrak Daun Jelema (*Clorodendrum Minahassae L.*) Terhadap Bakteri *Streptococcus Mutans*. *Pharmacol.* 4(4), p. 6. doi: 10.35799/pha.4.2015.10198
- Byambasuren, S.-E., Wang, J., Gaudel, G. et al. (2019). Medicinal value of wolfberry (*Lycium barbarum L.*). *J. Med. Plants. Stud.*; *Journal of Medicinal Plants Studies*, 7(4), p. 90–97.
- Chenicheri, S., R. U., Ramachandran, R., Thomas, V., Wood, A., et al. (2017). Insight into Oral Biofilm: Primary, Secondary and Residual Caries and Phyto-Challenged Solutions. *Open Dent. J. The Open Dentistry Journal*, 11, p. 312–333. doi: 10.2174/1874210601711010312
- Cheng, J., Zhou, Z., Sheng, H., He, L., Fan, X., He, Z., Sun, T., Zhang, X., Zhao, R., Gu, L., Cao, C., Zhou, S. 2015. An evidence-based update on the pharmacological activities and possible molecular targets of *Lycium barbarum* polysaccharides. *Drug Des Devel Ther.* 9, 33–78. doi: 10.2147/DDDT.S72892
- Chibane, L.B., Forquet, V., Lantéri, P., Clément, Y., Léonard, L., Oulahal, N., Degraeve, P., Bordes, C. 2019. Antibacterial properties of polyphenols: Characterization and QSAR (Quantitative structure-activity relationship) models. *Front. Microbiol.* 10, 829. doi: 10.3389/fmicb.2019.00829
- Conrads, G., and About, I. (2018). Pathophysiology of Dental Caries. *Monographs in Oral Science*, 27, pp. 1–10. doi: 10.1159/000487826
- Dewi, Z. Y., Nur, A., and Hertriani, T. (2015). Efek Antibakteri dan Penghambatan Biofilm Ekstrak Sereh (*Cymbopogon nardus L.*) terhadap Bakteri *Streptococcus mutans*. *Maj. kedokt. gigi Indones.*; *Majalah Kedokteran Gigi Indonesia*, 20(2), p. 136. doi: 10.22146/majkedgind.9120
- Fitri, N. I., Chirila, F., Nadas, G., Pall, E. et al. (2013). Comparative testing of antimicrobial activity of aqueous extracts of *Aloe vera* and *Lycium barbarum*. *Bull Univ Agric Sci Vet Med Cluj Napoca*; *Bulletin of University of Agricultural Sciences and Veterinary Medicine Cluj*

Formatted: Indent: First line: 0 ch

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold

Formatted: Indent: Left: 0", Hanging: 0.25"

Field Code Changed

Formatted: Indent: Left: 0", Hanging: 0.25"

Formatted: Font: Not Italic

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Font color: Text 1

Formatted: Check spelling and grammar

Formatted: Font color: Text 1

Formatted: Font: (Default) Times New Roman, 12 pt, Font color: Text 1

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Indent: Left: 0", Hanging: 0.25"

Formatted: Font: Not Italic

Formatted

Formatted: German (Germany)

Formatted

Formatted

Formatted

Formatted: Font color: Text 1



E. C. M. L. and C. H. C. (2017) 'Dental Biofilm and Laboratory Microbial Culture Models for Cariology Research', *Dentistry Journal*, 5(2), p. 21.

Sari Widyarman, A. and Kiky Elysia Lazaroni, N. (2019) 'Persistent Endodontics Pathogens Biofilm Inhibited by Lactobacillus reuteri Indonesian Strain Lactobacillus reuteri Indonesian', *Journal of Dentistry Indonesia*, 26(3), pp. 160-164.

Skenderidis, P., Lampakis, D., Lampakis, D., Giavasis, I., Leontopoulos, S., Petrotos, K., Hadjichristodoulou, C., Tsakalof, A., et al. (2019) 'Chemical properties, fatty-acid composition, and antioxidant activity of Goji berry (Lycium barbarum L. and Lycium Chinense mill.) fruits', *Antioxidants (Basel)*, 8(3), p. 60. doi: 10.3390/antiox8030060.

Skenderidis, P., Mitsagga, C., Giavasis, I., Petrotos, K., Lampakis, D., Leontopoulos, S., Hadjichristodoulou, C., Tsakalof, A. 2019. et al. (2019) 'The in vitro antimicrobial activity assessment of ultrasound assisted Lycium barbarum fruit extracts and pomegranate fruit peels', *J. Food Meas. Charact.*, *Journal of Food Measurement and Characterization*. Springer US, 13(35), p. 1-15 2017-2031. doi:10.1007/s11694-019-00123-6

Shinde, A.B., Mulay, Y.R. 2015. Phytochemical Analysis and Antibacterial Properties of Some Selected Indian Medicinal Plants. *Int J Curr Microbiol Appl Sci*. 4(3), 228-235.

Widyarman, A.S., Kiky Elysia Lazaroni, N.K.E. 2019. Persistent Endodontics Pathogens Biofilm Inhibited by Lactobacillus reuteri Indonesian Strain Lactobacillus reuteri Indonesian. *J Dent Indones*. 26(3), 160-164. doi:10.14693/jdi.v26i3.1113

Sonya Loresta, Sri Murwani, P. T. (2015) 'Efek Ekstrak Etanol Daun Kelor (Moringa oleifera) Terhadap Pembentukan Biofilm Staphylococcus aureus Secara In Vitro', 1(2), p. 20-25.

WHO. (2020) Oral Health. Available at: <https://www.who.int/news-room/fact-sheets/detail/oral-health>.

Yu, O.Y., Zhao, I. S., Mei, M.L., Lo E.C., Chu, C. 2017. Dental Biofilm and Laboratory Microbial Culture Models for Cariology Research. *Dent J (Basel)*. 5(2), 21. doi: 10.3390/dj5020021

Formatted: Font color: Text 1

Formatted: Indent: Left: 0", Hanging: 0.25"

Formatted: Font: Not Italic, Font color: Text 1

Formatted: Font color: Text 1

Formatted: Font color: Text 1

Formatted: Indent: Left: 0", Hanging: 0.25"

Formatted: Font color: Text 1

Formatted: Font: Not Italic, Font color: Text 1

Formatted: Font color: Text 1

Commented [A34]: Penulisan Dapfus ini sudah kuperbaiki, mohon bantuan koreksi untuk dapfus :

Hinrich, JE  
Kemenkes  
Loresta  
Ludwiczuk

Formatted: Indent: Hanging: 0.25"

Formatted: Indent: Left: 0", Hanging: 0.25"

Formatted: Indent: Left: 0", Space After: 8 pt,  
Line spacing: Multiple 1.08 li