

# Efficacy of *Tagetes erecta* Linn. leaf extract cream on rat dermal wound healing

*By* Mauritius Lambertus Edy Parwanto

## Efficacy of *Tagetes erecta* Linn. leaf extract cream on rat dermal wound healing

---

### ABSTRACT

Dermal wound of the human body are a problem that must be cured immediately. Therefore it is necessary to look for new herbal medicine options. The purpose of this study was to determine the effectiveness of the dose of *Tagetes erecta* Linn. leaf extract cream on rat dermal wound healing. This study is an animal treatment in Wistar rats. Rats were divided into 3 groups, namely groups I, II and III. All rats in the treatment groups were excision on the back skin. Group I (control) after excision was given the drug nitrofurazone cream, group II was given *T. erecta* Linn. leaf extract cream 3% and group III was given *T. erecta* Linn. leaf extract cream 4%. *T. erecta* Linn. leaf extract cream 3% and 4% play a role in increasing the strength of wound contraction, the percentage of wound contraction and accelerating the epithelialization period compared with the positive control group (nitrofurazone). *T. erecta* Linn. leaf extract cream 3% and 4% better as antibacterial than positive control (nitrofurazone). *T. erecta* Linn. leaf extract cream 3% and 4% increase protein and DNA content in the wound area compared with the positive control group (nitrofurazone). *T. erecta* Linn. leaf extract cream 3% and 4% can be followed by clinical trials in humans so that it can be an option for treating on dermal wounds healing.

**Key words:** *Tagetes erecta* Linn. leaf extract cream, wound healing, epithelialization period.

### INTRODUCTION

Dermal wound on the human body are a problem that must be cured immediately. Therefore it is necessary to look for new natural treatment options, both sourced from animals and plants. For this purpose we have tried to examine the contents of natural materials derived from animals and plants. We have tried to show 8 proteins fraction of earth worms (Parwanto *et al*, 2016). In the future, we hope that earthworm extract can be used for dermal wound healing. Beside that it has been demonstrated a formulation of *Lantana camara* Linn. leaf extract ointment (Parwanto *et al*, 2013) and look for a doses effectiveness for wound healing in rats (Parwanto, 2017). After the injury, there will be growth of dermal tissue to cover the wound. To accelerate wound healing, new drug preparations should be sought. The search for new drug preparations can be done by involving various extract of natural ingredients, including *Tagetes erecta* Linn. leaf extract.

The results of previous studies showed that isolation of steroids and phenolic acids from *T. erecta* Linn. leaf extract was carried out using column chromatography (Jayavant *et al*, 2020). The other study showed that 19 phytochemicals were contained by leaf extract of *L. camara* Linn. The main bioactive compounds are tetra decanoic acid, 2,6,10- trimethyl 14-ethylene-14-pentadecme, N-hexadecanmic acid, 15-hydroxy penta decanoic acid and stigmasterol (Devika and Justin, 2014). In addition, has been reported that qualitative phytochemical screening of *T. erecta* Linn. oil. The screening results showed that *T. erecta* Linn. contained terpenoids, flavonoids, alkaloids, quinones,

phenols, coumanins, carbohydrates and tannins. The screening shows that *T. erecta* Linn. contains terpenoids, flavonoids, alkaloids, quinones, phenols, coumanins, carbohydrates and tannins (Rajvanshi and Dwivedi, 2017). Furthermore, it has been proven quantitatively that lipophilic extract from *T. erecta plena* Linn. var. Hawaii contains carotenoids (up to  $80.51 \pm 1.15$  mg% equivalent to  $\beta$ -carotene) and flavonoids (up to  $0.7120 \pm 0.0060$ % equivalent to patuletin) (Maliuhina *et al*, 2020). Beside that, *T. erecta* Linn. also contains such as quercetin and its glycoside derivatives, ellagic acid, chlorogenic acid, kaempferol, myricitrin and rutin (Marrelli *et al*, 2013). Another study has also been carried out to evaluate the chemical content of *T. erecta* Linn. leaf extract were cultivated at Bandung, Central Java, Indonesia. Gas chromatography-mass spectrometry (GC-MS) was used to perform 17 active compounds with a retention time of 31,284 to 50,614. The three compounds with the highest content are neophytadine 43.88%, 9,12,15-acid-methyl ester Octadecadienoic 13.45%, and hexadecanoic acid-methyl ester 13.24% (Edy *et al*, 2017). There have been pharmacological studies of *T. erecta* Linn. leaf extract.

*T. erecta* Linn. extract has various pharmacological effects, including anti-bacterial, wound healing, antioxidant, antidiabetic, antihyperlipidemic, hepatoprotective, original and repellent, nematosidal, insecticidal, analgesic and cytotoxic (Karwani & Sisoidia, 2015). Research on wound healing in white rats has been carried out with cream extract of *T. erecta* Linn. 5% and 10%. The results showed that *T. erecta* Linn. extract cream was more beneficial to wound healing compared with *C. asiatica* extract (Chartterjee, 2011). Other studies have shown the effect of *T. erecta* Linn. 2.5% extract gel on wound healing in white rats (Kiranmai *et al*, 2011). In addition, the use of *T. erecta* Linn. for traditional treatment has been carried out.

Utilization of *T. erecta* Linn. has been carried out among others as an antiseptic ingredient and to overcome kidney problems, muscle aches and ulcers. Part of *T. erecta* Linn. flower is used to cure fever, epilepsy, stomach ache, scabies, liver disease, eye pain, rheumatism and bronchitis (Karwani & Sisoidia, 2015). It has been reported that methanol extract of *T. erecta* Linn. could protect human skin against photo-aging by attenuating oxidative damage, suppressing matrix metalloproteinase expression and/or activity as well as by stimulating collagen synthesis (Kang *et al*, 2018). Previous studies have reported that aqueous and alcoholic extracts of *T. erecta* Linn. has anxiolytic effect (Manisha *et al*, 2013). In addition, that each part of the plant of *T. erecta* Linn. has a certain phytochemical content. The main constituents of each part of the plant of *T. erecta* Linn. have different pharmacological effects such as anti-nociceptive, anti-inflammatory, anti-fungal effects, insecticides, larvicidal, hepatoprotective, antipyretic, wound healing, antibacterial, antimicrobial, antiepileptic, and antifungal properties (Singh *et al*, 2020). Another studies have shown that *T. erecta* Linn. flower extract is proven to have the ability of central nervous system

stimulant, antidepressant, and antipyretic activity (Shetty *et al.*, 2009). Based on the description of *T. erecta* Linn. in health, we would like focus on the use of this plants for dermal wound healing.

Wound healing is a process that occurs if there is physical interference with cells and tissues. So that wound healing is more effective and does not cause hypersensitive effects, it can be done by providing potential plant extract. Data shows that many plants have the potential for wound healing. Therefore, ethnobotany experts collect data about plants that have the potential to wound healing (Arun *et al.*, 2013). *T. erecta* Linn. also has wide ethno medicinal importance attributed to the bioactive compounds and essential oil found in leaves and having commercial application as medicine (Mir *et al.*, 2019). Furthermore was demonstrated the healing activity of the lipophilic extract of *T. erecta* Linn. (Maliuhina *et al.*, 2020). Related to this, it turns out DNA and protein play a role in wound healing.

Increased DNA content in wound healing indicates cellular hyperplasia. Along with this, there is an increase in total protein that shows active synthesis and deposition of matrix proteins in granulation tissue. The wound healing process depends on the regulation of biosynthesis, deposition and maturation of collagen. Increased content and deposition of collagen accelerate wound healing. Increasing the amount of collagen can increase the total cell count as a result of increased cell division (Sumitra *et al.*, 2005). A previous study has reported that various extract of *T. erecta* Linn. contains carotenoid lutein. Carotenoid lutein in *T. erecta* Linn. acts as an antioxidant thereby reducing the level of cell damage. In addition, carotenoid lutein also has the effect of reducing inflammation and immunosuppressants (Hajare *et al.*, 2013). Decreasing the level of cell damage can be done in the process of dermal wound healing.

In addition of dermal wound healing, the level of cell damage is also influenced by the infecting bacteria. Previous study has revealed the role of *T. erecta* Linn. extract as an antibacterial. *T. erecta* Linn. extract also has antibacterial effects against *Escherichia coli*, *Proteus vulgaris*, *Proteus mirabilis*, *Aeromonas sobria*, *Aeromonas hydrophila*, *Plesiomonas shigelloides* and *Salmonella enterica*. In addition, extract of *T. erecta* Linn. also have antibacterial effects against *Bacillus subtilis*, *Bacillus cereus*, *Bacillus circulans* and *Staphylococcus aureus* (Jain *et al.*, 2012). *T. erecta* Linn. 2.5% extract gel has also been shown to be antibacterial. The inhibition zone diameter of the extract against *S. aureus* was 17.87 mm, whereas for *E. coli* it was 15.33 mm (Edy *et al.*, 2019). Gel preparations have been proven stable after being tested using a climate chamber (Memmert) with a temperature of 45 °C±2 °C and a relative humidity of 75%±5% for 90 days (Edy *et al.*, 2017). As a consequence, increasing the number of bacteria inhibits the wound healing process. Therefore it is necessary to use *T. erecta* Linn. extract in various preparations for wound healing.

The formula of *T. erecta* Linn. leaf extract cream needs to be carefully optimized. This is based on our experience in making *L. camara* Linn. leaf extract ointment 20% and 24%. The *L. camara* Linn. leaf extract ointment conforms to the parameters of organoleptic, homogeneity and pH, but does not meet the spread power ointment requirements (Parwanto *et al*, 2013). In addition, *T. erecta* Linn. extract has also been successfully formulated into various pharmaceutical forms such as gel, cream, anti-mosquito lotion and hair dye (Edy & Parwanto, 2019). A previous study has proven the effect of *T. erecta* Linn. 2.5% leaf extract cream (Gopi *et al*, 2012) or *T. erecta* Linn. 5% extract gel on wound healing (Kiranmai *et al*, 2011). The purpose of this study was to determine the effect of *T. erecta* Linn. leaf extract cream 3% and 4% on dermal wound healing in Wistar rats.

## **MATERIALS AND METHODS**

### **Study area**

The research project was divided into 3 stages, namely the first stage is plant cultivation of *T. erecta* Linn., the second stage is making *T. erecta* Linn. leaf extract cream, while the third stage is treatment of *T. erecta* Linn. leaf extract cream in Wistar rats. Cultivation of *T. erecta* Linn. carried out on agricultural land in the Bandung District, Semarang Regency, Central Java Province, Indonesia. The cultivation time is from July to November 2017. The making of *T. erecta* Linn. leaf extract cream was carried out at the Biomedical Laboratory, Faculty of Medicine, Universitas Trisakti, Indonesia from December 2017 to March 2018. The treatment of *T. erecta* Linn. leaf extract cream on the Wistar rat was carried out from April 2018 to August 2018 in Animal Laboratory, Faculty of Veterinary Medicine, IPB University, Bogor, Indonesia.

### **Ethical clearance approval**

The study was only done after the recommendation approval by “Komisi Etik Riset, Fakultas Kedokteran, Universitas Trisakti, Indonesia” No. 90/KER/FK/V/2016.

### **Formulation of *T. erecta* Linn. leaf extract cream**

*T. erecta* Linn. identified beforehand to determine the species of plants. Leaves of *T. erecta* Linn. obtained by cultivated at Bandung, Semarang, Central Java, Indonesia. Leaves of *T. erecta* Linn. covered with a black cloth and then dried in the sun to dry. The dried leaves are then weighed and made powder. Extraction of *T. erecta* Linn. powder using ethanol 70% and then filtered to yield a viscous supernatant as crude extract. Aliquot of the extract lyophilized and the results are weighed. *T. erecta* Linn. leaf extract is used as an active ingredient of cream. Composition of the

cream base ingredient has been optimized. Cream base ingredient consist of 14.2 grams stearic acid, 10 grams glycerol, 0.25 grams borax, 1 grams triethanolamine, 0.1 grams nipagin, 0.05 grams nipasol, 9.5 mL mono sodium phosphate (NaH<sub>2</sub>PO<sub>4</sub>) 2.55%, 0.5 mL NaH<sub>2</sub>PO<sub>4</sub> 1.85% and aquabidestilata add to 100 mL. *T. erecta* Linn. leaf extract cream 3% and 4% are made by adding 3 grams and 4 grams of *T. erecta* Linn. leaf extract into each the cream base ingredients.

### **Animal maintenance**

In this study involve 24 of Wistar male rats (*Rattus novergicus*, Wistar starin), aged between 3-4 months with average weight of 150-200 gram were used. The animals were made acclimatization in the cage for 1 week before treatment. Rats were fed and drunk of libitum according to standards. Treatment rooms are equipped with air conditioners with temperatures of 22±3 °C, relative humidity of 55±5%, and artificial fluorescent lamps (12:12 hours, light and dark cycles).

### **Experimental design**

Randomly, Wistar rats are grouped into 3 groups. Group I: Wistar rats were dermal back excision and treated with nitrofurazone (positive control). Group II: Wistar rats were dermal back excision and treated with *T. erecta* Linn. leaf extract cream 3%. Group III: Wistar rats were dermal back excision and treated with *T. erecta* Linn. leaf extract cream 4%. Combination of ketamine (50 mg/kg) and xylazine (10 mg/kg) intraperitoneally were used for rats anesthetized before dermal back excision. Administration of nitrofurazone and *T. erecta* Linn. leaf extract cream daily, for 16 days after back skin excision. Granulation tissue was taken on days 4, 8, 12 and 16 for analysis.

### **Measurement of dermal wound healing**

Graphical planimetry methods was used to measurement of rat wound on days 0, 4, 8, 12 and 16. The size of the wound is marked on all edges of the wound using paper with ink markers. The paper is then attached to graph paper, then measured manually in units of mm<sup>2</sup>. The strength of wound contraction (SWC) is the difference of wound on the measurement at day 0 compared with the specified day. The percentage of wound contractions (PWC) is calculated using the formula as follows  $PWC = \{(\text{area on day 0} - \text{area on final day}) / \text{area on day 0}\} \times 100$  (Dwivedi and Chaudhary, 2012). Epithelialization period is determined by the number of days needed for epithelialization at the site of the wound (Sumitra *et al*, 2005).

### **Calculation of bacteria colonies number**

In this study we used 2 media, phosphate-buffered saline (PBS) and Azide blood agar (ABA). The composition of PBS consisted of 8 grams sodium chloride, 0.2 grams kalium chloride, 1.42 grams disodium phosphate and 0.42 grams potassium phosphate monobasic. Final pH of PBS is 7.4, stored at 25 °C. The composition of ABA media consists of 3 grams meat extract, 10 grams tryptose, 5 grams sodium chloride, 0.2 grams sodium azide, 15 grams agar powder, deionized water is added to 1000 mL. Final pH of ABA is 7.3, stored at 25 °C. The number of bacteria in the granulation tissue was calculated on the 6<sup>th</sup> day of treatment (Hirsch *et al*, 2008). One hundred  $\mu$ L homogenate granulation tissue was taken to count bacterial colonies. The solution series was inoculated in the PBS for isolation, while the selective culture medium was ABA. Petri was incubated aerobically at 37 °C for 24 hours. Bacteria colonies number expressed as log of 10 colony-forming units (CFU) per gram of tissue. Number of bacteria colonies  $>1 \times 10^5$  declared granulation tissue infected with bacteria (O'Meara *et al*, 2006).

### Biochemical measurements

DNA and protein content in the granulation tissue were extracted in 5% trichloroacetic acid (trichloroacetic acid=TCA). Ten mL of TCA (5%) added to the tissue (100 mg dry weight of the tissue) was heated at 90 °C for 30 minutes in a water bath to extract DNA and proteins. The solution is centrifuged and the supernatant is used to measure DNA and protein levels. Measurement of DNA content using the Burton method, whereas protein measurement using the Lowry method, (Sumitra *et al*, 2005).

### Histology analysis

At the end of the treatment phase, rats were anesthetized using a combination of xylazine (10 mg/kg) and ketamine (100 mg/kg), and skin tissue (0.8 $\times$ 0.5 cm) were collected into a 10% formalin solution for histological analysis. After the skin tissue is fixed, then wash with running water. The next phase is dehydrated with multilevel alcohol, and then washed with xylene. After the dehydration and washing phase, skin tissue is wrapped in paraffin (melting point 56 °C). Hematoxylin and Eosin (H&E) are used to paint the tissue. Tissue histology observations were carried out using a microscope and documented with photomicrographs.

### Statistical analysis

Data of all groups were estimated by one-way analysis of variance (ANOVA) and declared significance if  $p < 0.05$ . If there are differences between groups, followed by the least significant difference (LSD) test.

## RESULTS

*T. erecta* Linn. were used in this study as a result of plant cultivation on agricultural land in Bandungan District, Semarang Regency, Central Java Province, Indonesia (Fig. 1). The geographical location of Bandungan District which is on the southern slopes of Mount Ungaran. Based on Urchin Tracking Module (UTM) coordinates, Bandungan District is located between 425275 - 435093 mE and 9197640 - 9205676 mN.

There are differences in the effect of *T. erecta* Linn. leaf extract cream compared with nitrofurazone on wound healing (Fig. 2). *T. erecta* Linn. leaf extract cream 4% is faster healing process, followed by the *T. erecta* Linn. leaf extract cream 3% and the nitrofurazone group (positive control). The speed of wound healing is characterized by a decrease in wound size (Table 1). Wound size on 4<sup>th</sup> day showed that group I>II (P=0.022), I>III (P=0.000), II>III (P=0.024). Wound size on 8<sup>th</sup> day showed that group I>II, III(0.000), while group II=III (0.059). Wound size on 12<sup>th</sup> day showed that group I>II (0.006), group I>III (0.000) and group II>III (0.022). Wound size on 16<sup>th</sup> day showed that group I>II>III (0.000).

Strength of wound contraction is a healing response to reduce of wound size, then the damaged tissue is completely repaired. Strength of wound contraction between groups I, II and III at day 0 of treatment as initial treatment day, as well as a base line data. Strength of wound contraction between groups I, II and III at 4<sup>th</sup>, 8<sup>th</sup> and 16<sup>th</sup> days were significantly different (P=0.000). The results of this study showed that *T. erecta* Linn. leaf extract cream 3% and 4% at 4<sup>th</sup>, 8<sup>th</sup> and 16<sup>th</sup> days had better of strength wound contraction compared with positive controls (nitrofurazone) (P=0.000) (Table 2). Strength of wound contraction on 4<sup>th</sup> day showed that group I<II (P=0.022), I<III (P=0.000), II<III (P=0.024). Strength of wound contraction on 8<sup>th</sup> day showed that group I<II, III(P=0.000), while group II=III (P=0.059). Strength of wound contraction on 12<sup>th</sup> day showed that group I<II (P=0.006), group I<III (P=0.000) and group II<III (P=0.022). Strength of wound contraction on 16<sup>th</sup> day showed that group I<II<III (P=0.000).

Percentage of wound contractions between groups I, II and III at day 0 of treatment is initial treatment day, as well as a base line data. Percentage of wound contractions between groups I, II and III at 4<sup>th</sup>, 8<sup>th</sup>, 12<sup>th</sup> and 16<sup>th</sup> days were significantly different (P=0.000). The percentage of wound contraction in group II and III are stronger than group I (Table 3). *T. erecta* Linn. leaf extract cream increase the percentage of rat dermal wound healing compared with nitrofurazone. *T. erecta* Linn. leaf extract cream 3% and 4% at 4<sup>th</sup>, 8<sup>th</sup> and 16<sup>th</sup> days had better on percentage of wound contraction compared with positive controls (nitrofurazone).



We used microscopic observation for determine of epithelialization. Base on the result of epithelialization period in this study, we are divided into 3 stages of the wound healing process, namely early, middle and advanced stages (Fig. 3). In the early stages, only 1 cutaneous muscle is seen in the field of view on a microscope with a magnification of 40X (Fig. 3. A). In the middle stage, two cutaneous muscles appear in the field of view on a microscope. The distance between the two cutaneous muscles is getting closer which shows progress in wound healing (Fig. 3. B, C and D). At an advanced stage, two cutaneous muscles appear close together (Fig. 3. E). In the advanced stages of wound healing, epithelialization appears on the surface of the wound healing area (Fig. 3. F).

Number of bacterial colonies (NBC) at 6<sup>th</sup> day in this study showed that in group I the highest, then group III, while group II was lowest (P=0.000) (Fig. 5). DNA levels at 8<sup>th</sup>, 12<sup>th</sup> and 16<sup>th</sup> showed that group I was lower than group II and III (P<0.05). The all groups showed that the DNA content at the 8<sup>th</sup> day was highest and decreased at the 12<sup>th</sup> day and the lowest at the 16<sup>th</sup> day. At the 16<sup>th</sup> day, groups II and III did not differ in their DNA content (P=0.457), both groups had higher DNA content compared with group I (P=0.000)(Table 4). Proteins levels at 8<sup>th</sup>, 12<sup>th</sup> and 16<sup>th</sup> showed that group I was lowest compared with group II and III (P<0.05). The all groups showed that the proteins levels at the 8<sup>th</sup> day was lowest and increased at the 12<sup>th</sup> day and then decreased at the 16<sup>th</sup> day (P<0.05). The proteins levels at 8<sup>th</sup> day and 12<sup>th</sup> day observation showed that group I was lowest compared with group II, while group III was higher (P=0.000). The proteins levels at 16<sup>th</sup> day observation showed that group I was lowest compared with group II and III (P=0.000), likewise group II was lower compared with group III (P=0.013) (Table 5).

## DISCUSSION

Identification of *T. erecta* Linn. was carried out according to instructions in the book *Flora of Java* (specifically for Spermatophyta) (Backer CA and van den Brink B, 1963-1968). The result of *T. erecta* Linn. extraction are used as active ingredients of the cream. Furthermore, the cream containing *T. erecta* Linn. leaf extract was tested for its effect on wound healing in Wistar rats. We have experience extracting another specimen namely *L. camara* Linn. and has been tested as an ointment for rat dermal wound healing (Parwanto *et al*, 2013, 2017). Next, we will to examine effects of *T. erecta* Linn. extract cream compared with *L. camara* Linn. extract cream in rat dermal wound healing.

The base cream were used in this study has been simulated. *T. erecta* Linn. leaf extract cream contains active ingredients namely *T. erecta* Linn. leaf extract. *T. erecta* Linn. 3% and 4% leaf extract cream in this study were semi-solid, with the distinctive aroma of *T. erecta* Linn.

extract, slightly black green, pH 5, homogeneous and not thick, spread power of 5.2 cm. The characteristics of the cream fulfill as topical preparations that are suitable for use. In addition, bacterial media is has been prepared and easy to use for isolation and culture.

Base on wound size, *T. erecta* Linn. leaf extract cream 3% and 4% are faster healing process in Wistar rats compared to nitrofurazone. In addition to the reduction of wound size, strength of wound contraction and percentage of wound contraction can also be used as an indicator of the speed of wound healing. Base on strength of wound contraction, *T. erecta* Linn. leaf extract cream 3% and 4% are stronger on healing process in Wistar rats compared with nitrofurazone. Based on these results, we expect that *T. erecta* Linn. 3% and 4% leaf extract cream can be used as a healing agent for dermal wounds. The results of this study according to the results of previous studies which showed the effectiveness of *T. erecta* Linn. on rat dermal wound healing (Kiranmai et al, 2011). Furthermore demonstrated that strength of wound contraction in control rats 21.6% to 68.3% from 4<sup>th</sup> day to 12<sup>th</sup> day and 80.6% to 98.1% from 14<sup>th</sup> day to 20<sup>th</sup> day (Murthy et al, 2013). In addition, recent research was conducted to examine the effect of anti-ulcer activity of *T. erecta* Linn. leaf extract gel 2.5% and 5% compared with a standard drug, triamcinolone. The results showed that *T. erecta* Linn. leaf extract gel showed significant oral ulcer protective activity (83.6%) compared with triamcinolone. This fact demonstrated that the *T. erecta* Linn. leaf extract gel has better potential against oral ulcers (Lakshana et al, 2020).

*T. erecta* Linn. leaf extract cream 3% and 4% at 4<sup>th</sup>, 8<sup>th</sup> and 16<sup>th</sup> days had better on percentage of wound contraction compared with positive controls (nitrofurazone). This fact is consistent with studies of *T. erecta* Linn. extract in rat dermal wound healing (Kiranmai et al, 2011). The results of this study are consistent with other studies, which show that ethanol extract of *T. erecta* Linn. have different percentage of wound contraction between 4<sup>th</sup>, 8<sup>th</sup> and 16<sup>th</sup> days (Charterjee et al, 2011). If we pay attention, the data about strength of wound contraction is related to the percentage of wound contraction. Base on data in this study, *T. erecta* Linn. leaf extract cream 3 % and 4% showed strength of wound contraction and percentage of wound contraction are better than nitrofurazone, then clinical trials need to be done on human wound healing.

The result of epithelialization period analysis between groups I, II and III were significant different ( $P < 0.05$ ). Effect of *T. erecta* Linn. leaf extract cream on epithelialization period of dermal wound healing in Wistar rats showed that group I > II, III ( $P = 0.000$ ), while group II = III ( $P = 0.232$ ) (Fig. 4). In the rat dermal wound healing, the effect of *T. erecta* Linn. leaf extract cream 3% on epithelialization period was not significantly different compare with *T. erecta* Linn. leaf extract cream 4%. The effect on epithelialization period of the two creams is shorter than the nitrofurazone cream. Because *T. erecta* Linn. cream is faster in wound healing than nitrofurazone cream,

subsequently *T. erecta* Linn. leaf extract cream 3% and 4% have <sup>42</sup> the potential to be developed as a wound healing <sup>8</sup> drug. Previous studies have shown that epithelialization period of *T. erecta* Linn. leaf extract gel 2.5% was faster than *T. erecta* Linn. leaf extract cream 3% and 4% in this study, however *T. erecta* Linn. leaf extract cream 3% and 4% can be developed as an alternative to wound healing. The results of other studies showed that epithelialization period of ethanol extract of *T. erecta* Linn. 5% (w/w) was significantly different compared with *T. erecta* Linn. 10% (w/w). Epithelialization period of ethanol extract of *T. erecta* Linn. 5% is  $18.00 \pm 0.51$  days, while ethanol extract of *T. erecta* Linn. 10% is  $14.16 \pm 0.75$  days (Charterjee et al, 2011). Beside that, complete epithelialization and healing in control rats were observed on 24<sup>th</sup> day (Murthy et al, 2013). Apparently the epithelialization period of dermal wound healing in Wistar rats in this study is different from mice. The results of studies on the injured mice, then treated with nitrofurazone showed that epithelialization period was around 14.67 days (Belachev et al, 2020). The evidence shows that different epithelialization periods are caused by different types of experimental animals.

Base on data in this study indicated that the *T. erecta* Linn. leaf extract cream 3% and 4% has a better role as an antibacterial than a <sup>27</sup> positive control (nitrofurazone). The results of this study are consistent with studies showing that the ethyl acetate fraction of *T. erecta* Linn. has activity against *S. aureus* (Trinh et al, 2020). Beside that, it has been demonstrated that *T. erecta* Linn. had antimicrobial activity inhibiting the growth of *S. agalactiae* and suggesting that they may be useful in the treatment of *S. agalactiae* infections in humans and animals (Mekvimol et al, 2020).

Data in this study indicated that the process of wound healing involves the role of DNA. Indications of wound healing involving DNA in this studies in accordance with the statement that <sup>41</sup> wound healing is a complex process with stages that are not linear, but there is an overlapping process involving 1,000 genes to achieve closure of dermal wounds (Cooper et al, 2005). The dynamics of changes in DNA levels on wound healing in this study are similar to the results of our previous study of the efficacy <sup>1</sup> of *L. camara* Linn. leaf extracts ointment on dermal wound healing (Parwanto, 2017). We have not found references to the effects of *T. erecta* Linn. on DNA levels in wound healing. Nevertheless, a previous studies in rats showed the effect of *Potentilla fulgens* on increasing DNA levels up to 60.0% on the 16<sup>th</sup> day (Kundu et al, 2016). Furthermore, the ointment consists of *Cordia obliqua* Willd. and *Vigna radiata* Linn. extract proven to increase DNA synthesis, thereby increasing tissue regeneration and accelerating wound contraction (Kadhiravan et al, 2017).

## CONCLUSION

*T. erecta* Linn. leaf extract cream 3% and 4% better act as antibacterial than positive control (nitrofurazone). *T. erecta* Linn. leaf extract cream 3% and 4% play a role in increasing the strength of wound contraction, the percentage of wound contraction and accelerating the time of epithelialization compared with the positive control group (nitrofurazone). *T. erecta* Linn. leaf extract cream 3% and 4% increased the DNA and protein content in the injured area compared with the positive control group (nitrofurazone). We recommend that *T. erecta* Linn. leaf extract cream 3% and 4% can be continued to clinical trials on human dermal wound healing.

### **SIGNIFICANCE STATEMENT**

This study discovered that *T. erecta* Linn. leaf extract cream 3% and 4% were faster than nitrofurazone cream on the wound healing process in Wistar rat. Scientific evidence from this study in the future can be followed by clinical trials in humans so that it can be an option for treating on dermal wounds healing.

### **AUTHOR'S CONTRIBUTIONS**

MLEP and DT: Schemed and designed experiment. MLEP and HJE: Collecting and interpretation of the results. All authors analyzed the data and wrote the manuscript. All authors read and approved the final manuscript.

### **ACKNOWLEDGEMENTS**

This work was supported by the Faculty of Medicine, Universitas Trisakti, Indonesia. This research was funded in the budget year 2017-2018.

### **REFERENCES**

- Arun M, Satish S and Anima P. Herbal Boon for Wounds. *Int J Pharm Pharm Sci*, 2013, 5(2):1-12.
- Backer CA and Bakhuizen van den Brink Jr RC. *Flora of Java (Spermatophytes only)*. Groningen: P. Noordhoff, 1963-1968.
- Belachew TF, Asrade S, Geta M, and Fentahun E. In Vivo Evaluation of Wound Healing and Anti-Inflammatory Activity of 80% Methanol Crude Flower Extract of *Hagenia abyssinica* (Bruce) J.F. Gmel in Mice. *Evid Based Complementary Altern Med*, 2020, Article ID 9645792: 1-12. <https://doi.org/10.1155/2020/9645792>.

Chartterjee S, Prakash T, Kotrsha D, Rao NR and Goli D. Comparative efficacy of *Tagetes erecta* and *Centella asiatica* extracts on wound healing in albino rat. *Chin Med*, 2011; 2:138-142. doi:10.4236/cm.2011.24023.

Cooper L, Johnson C, Burslem F, Martin P. Wound healing and inflammation genes revealed by array analysis of 'macrophageless' PU.1 null mice. *Genome Biol*, 2005; 6(1):R5:1-17.

Devika R, Justin K. Screening and evaluation of bioactive components of *Tagetes erecta* L. by gc-ms analysis. *Asian J Pharm Clin Res*, 2014; 7(Suppl 2):58-60.

Dwivedi VK and Chaudhary M. Comparative wound healing efficacy of ampuicare and becaplermin in diabetic rat. *Afr J Pharm Pharmacol*, 2012; 6(12):883-892.

Edy HJ, Marchaban, Wahyuono S, Nugroho AE. Pengujian Aktivitas Antibakteri Hidrogel Ekstrak Etanol Daun *Tagetes erecta* L. *Jurnal MIPA*, 2019; 8 (3):96-98.

Edy HJ, Wahyuono S, Nugroho AE, Marchaban. Characterization and Evaluation of Bioactive Compounds of Extract Ethanol *Tagetes Erecta* L. by GC-MS. *Int J Chemtech Res*, 2017; 10:172-175.

Edy HJ, Wahyuono S, Nugroho AE, Marchaban. Formulation and Evaluation of Hydrogel Containing *Tagetes erecta* L. Leaves Etanolic Extract. *IJCIR*, 2017; 3:627-630.

Edy HJ, Parwanto MLE. Pemanfaatan tanaman *Tagetes erecta* Linn. dalam kesehatan. *JBK*, 2020, 2(2): 77-80.

Gopi G, Elumalai A, Jayasri P. A concise review on *Tagetes erecta*. *Int J of Phytopharm Res*, 2012; 3(1):16-19.

Gurtner GC, Werner S, Barrandon Y, Longaker MT. Wound repair and regeneration. *Nature*, 2008; 453:314–321.

Hajare R, Ray A, Shreya S, Tharachand C, Avadhani MMN, Selvaraj C. Extraction and Quantification of Antioxidant Lutein from Various Plant Sources. *Int J Pharm Sci Rev Res*, 2013; 22(1):152-157.

Hirsch T, Spielmann M, Zuhaili B, Koehler T, Fossum M, Steinau HU, Yao F, Steinstraesser L, Onderdonk AB and Eriksson E. Enhanced susceptibility to infections in a diabetic wound healing model. *BMC Surgery*, 2008; 8(5):1-5. **DOI:** 10.1186/1471-2482-8-5.

Jain R, Katare N, Kumar V, Samanta AK, Goswami S, Shrotri CK. *In Vitro* Anti Bacterial Potential of Different Extracts of *Tagetes Erecta* and *Tagetes Patula*. J Nat Sci Res, 2012; 2(5):84-90.

Jayavant TK, Shirote PJ, Kadam AS, Deshmane B, Dhanashree RK, Tejaswini YS, Nivedita PH. Isolation and investigation of phytochemicals and pharmacological screening of *Tagetes erecta* L. leaves extract. AJPRD, 2018;6 (4):39-44. <http://ajprd.com/index.php>. DOI: <http://dx.doi.org/10.22270/ajprd.v6.i4.360>.

Kadhiravan M, Keerthana K, Shobana G, Jothi G, Radhika J. Healing potential of a polyherbal ointment on excision wound in normal and diabetes-induced albino rats. Asian J Pharm Clin Res, 2017; 10(4):41-45.

Karwani G and Sisodia SS. *Tagetes erecta* plant: Review with significant pharmacological activities. World J Pharm Sci, 2015; 3(6):1180-1183.

Kiranmai M, Kazim SM and Ibrahim M. Combined wound healing activity of *Gymnema sylvestere* and *Tagetes erecta* Linn. IJPA, 2011; 2 (2):135-140.

Kundu A, Ghosh A, Singh NK, Singh GK, Seth A, Maurya SK, Hemalatha S and Laloo D. Wound healing activity of the ethanol root extract and polyphenolic rich fraction from *Potentilla fulgens*. Pharm Biol, 2016; 54(11):2383-2393. <http://dx.doi.org/10.3109/13880209.2016.1157192>.

Lakshana S, Vijayalakshmi S, Dinakar J, Asok Kumar K. Effects of *Tagetes erecta* gel on experimentally induced oral ulcer model in rats. Int J Res Pharm Sci, 2020; 11(2):1844-1848. DOI: <https://doi.org/10.26452/ijrps.v11i2.2090>.

Maliuhina OO, Smoilovska HP, Bielenichev IF, Mazulin OV, Khortetska TV. Wound healing activity of the lipophilic extract of *Tagetes erecta* L. Zaporozhye Med J, 2019; 21(2):253-257. UDC: 615.322:582.998.16]:616.5-001.4-08. <http://zmmj.zsmu.edu.ua>.

Manisha RL, Riyaz Shaik, B Satyanarayana, Nazma SK, Nadeem SK, Vidhyadhararao B, Vijay Kumar J, Venkateswara Rao I, Sadhik SK. Evaluation of Anxiolytic Activity of Flower Extracts of *Tagetes Erecta* Linn (Asteraceae) in Rats. JAPS, 2013; 39(12):75-82. DOI: <10.7324/JAPS.2013.31214>.

Marrelli M, Loizzo MR, Nicoletti M, Menichini F, Confort F. Inhibition of key enzymes linked to obesity by preparations from Mediterranean dietary plants: effects on  $\alpha$ -amylase and pancreatic lipase activities. Plant Food Hum Nutr, 2013; 68:340-346.

Mekvimol T, Poonthong G, Chaipunna C and Pumipuntu N. Antimicrobial activity of marigold (*Tagetes erecta*), mulberry (*Morus indica*), and red shallot (*Allium ascalonicum*) extracts against *Streptococcus agalactiae*. *Int J One Health*, 2020; 6(1):56-60. [www.onehealthjournal.org/Vol.6/No.1/10.pdf](http://www.onehealthjournal.org/Vol.6/No.1/10.pdf).

Mir RA, Ahanger MA, Agarwal RM. Marigold: From Mandap to Medicine and from Ornamentation to Remediation. *Am J Plant Sci*, 2019; 10:309-338. DOI: 10.4236/ajps.2019.102024. <http://www.scirp.org/journal/ajps>.

Murthy S, Gautam MK, Goel S, Purohit V, Sharma H, and Goel RK. Evaluation of in vivo wound healing activity of *Bacopa monniera* on different wound model in rats. *BioMed Res In*, 2013, Article ID 972028:1-9. <http://dx.doi.org/10.1155/2013/972028>.

O'Meara S, Nelson EA, Golder S, Dalton JE, Craig D, Iglesias C. Systematic review of methods to diagnose infection in foot ulcers in diabetes. *Diabet Med*, 2006; 23(4):341-347.

Parwanto EML, Senjaya H, Edy HJ. Formulasi salep antibakteri ekstrak etanol daun *L camara* (*Lantana camara* L). *Pharmacon*, 2013; 2 (3):104-108.

Parwanto MLE. Efficacy of *Lantana camara* Linn. leaf extracts ointment on dermal wound healing were infected with *Staphylococcus epidermidis*. *Int J Basic Clin Pharmacol*, 2017; 6:503-510.

Parwanto MLE, Mahyunis, Senjaya H, Edy HJ, Syamsurizal. Fractionation and Characterization of Proteins in *Lumbricus rubellus* Powders. *IJPCR*, 2016; 8(1):15-21. [www.ijpcr.com](http://www.ijpcr.com).

Rajvanshi SK and Dwivedi DH. Phytochemical screening studies of bioactive compounds of African marigold (*Tagetes erecta* L.). *J Pharmacogn Phytochem*, 2017; 6(4):524-527.

Shetty LJ, Harikiran H, Fernandes J. Pharmacological evaluation of ethanolic extract of flowers of *Tagetes erecta* on epilepsy. *J Pharm Res*, 2009; 2(6):1035-1038.

Shao K, Han B, Gao J, Jiang Z, Liu W, Liu W, and Liang Y. Fabrication and feasibility study of an absorbable diacetyl chitin surgical suture for wound healing. *J Biomed Mater Res B Appl Biomater*, 2016; 104:116-125.

Singh Y, Gupta A and Kannoja P. *Tagetes erecta* (Marigold)-A review on its phytochemical and medicinal properties. *Curr Med Drug Res*, 2020; 4(1):1-6.

Sumitra M, Manikandana P, Suguna L. Efficacy of *Butea monosperma* on dermal wound healing in rats. *Int J Biochem Cell Biol*, 2005; 37(3):566-573. doi: 10.1016/j.biocel.2004.08.003.

Trinh PC, Thao LTT, Ha HTV, and Nguyen TA. DPPH-Scavenging and Antimicrobial Activities of Asteraceae Medicinal Plants on Uropathogenic Bacteria. *Evid Based Complementary Altern*, 2020: Article ID 7807026:1-9. <https://doi.org/10.1155/2020/7807026>.

Table 1. Effect of *Tagetes erecta* Linn. leaf extract cream on wound size in Wistar rats.

Variables	Group	Days of observation				
		0 day (Means ± SD)	4 <sup>th</sup> day Means ± SD	8 <sup>th</sup> day Means ± SD	12 <sup>th</sup> day Means ± SD	16 <sup>th</sup> day Means ± SD
WS	I	400.00 ± 0.00	344.59 ± 6.33	276.56 ± 10.29	85.20 ± 1.73	29.48 ± 2.71



(mm <sup>2</sup> )	II	400.00 ± 0.00	334.77 ± 10.04	244.50 ± 6.86	80.10 ± 2.55	22.26 ± 1.27
	III	400.00 ± 0.00	325.97 ± 6.98	234.02 ± 13.35	75.03 ± 4.93	16.76 ± 2.65
	P	-	I>II (0.022) I>III (0.000) II>III(0.024)	I>II, III(0.000) II=III(0.059)	I>II (0.006) I>III (0.000) II>III (0.022)	I>II>III (0.000)

Abbreviations: WS=wound size, group I=Wistar rats were back skin incision<sup>1</sup> and treated with nitrofurazone (positive control), group II=Wistar rats were back skin incision and treated with *T. erecta* Linn.leaf extract cream 3%, group III=Wistar rats were back skin incision and treated with *T. erecta* Linn.leaf extract cream 4%, mm<sup>2</sup>=millimeter square, SD=standard of deviation, P=significance value.

Table 2. Effect of *Tagetes erecta* Linn. leaf extract cream on strength of wound contraction in Wistar rats.

Variables	Group	Days of observation				
		Δ WS	Δ WS	Δ WS	Δ WS	Δ WS
		0 day – 0 day (Means ± SD)	0 day – 4 <sup>th</sup> day Means ± SD	0 day – 8 <sup>th</sup> day Means ± SD	0 day – 12 <sup>th</sup> day Means ± SD	0 day – 16 <sup>th</sup> day Means ± SD
SWC (mm <sup>2</sup> )	I	0.00 ± 0.00	55.41 ± 6.33	123.44 ± 10.29	314.80 ± 1.73	370.52 ± 2.71
	II	0.00 ± 0.00	65.23 ± 10.04	155.50 ± 6.86	319.90 ± 2.55	377.74 ± 1.27
	III	0.00 ± 0.00	74.93 ± 6.98	165.98 ± 13.35	324.07 ± 4.93	383.24 ± 2.65
P	-	-	I<II (0.022) I<III (0.000) II<III (0.024)	I<II, III (0.000) II=III (0.059)	I<II (0.006) I<III (0.000) II<III (0.022)	I<II<III (0.000)

Abbreviations: SWC=strength of wound contraction, Δ WS= difference of wound size on the measurement at 0 day observation compared with the specified day, group I=Wistar rats were back skin incision and treated with nitrofurazone (positive control), group II=Wistar rats were back skin incision and treated with *T. erecta* Linn.leaf extract cream 3%, group III=Wistar rats were back skin incision and treated with *T. erecta* Linn.leaf extract cream 4%, mm<sup>2</sup>=millimeter square, SD=standard of deviation, P=significance value.

Table 3. Effect of *Tagetes erecta* Linn. leaf extract cream on percentage of wound contraction in Wistar rats.

Variables	Group	Days of observation				
		0 day	4 <sup>th</sup> day	8 <sup>th</sup> day	12 <sup>th</sup> day	16 <sup>th</sup> day
		(Means ± SD)	Means ± SD	Means ± SD	Means ± SD	Means ± SD
PWC	I	0.00 ± 0.00	13.85 ± 1.58	30.86 ± 2.57	78.70 ± 0.43	92.63 ± 0.68

(%)	II	0.00 ± 0.00	16.31 ± 2.51	38.88 ± 1.72	79.98 ± 0.64	94.44 ± 0.32
	III	0.00 ± 0.00	18.73 ± 1.74	41.49 ± 3.34	81.02 ± 1.23	95.81 ± 0.66
	P	-	I<II (0.022)	I<II, III (0.000)	I<II (0.006)	I<II<III (0.000)
			I<III (0.000)	II=III (0.059)	I<III (0.000)	
			II<III (0.023)		II<III (0.021)	

Abbreviations: PWC=percentage of wound contraction, group I=Wistar rats were back skin incision and treated with nitrofurazone (positive control), group II=Wistar rats were back skin incision and treated with *T. erecta* Linn.leaf extract cream 3%, group III=Wistar rats were back skin incision and treated with *T. erecta* Linn.leaf extract cream 4%, mm<sup>2</sup>=millimeter square, SD=standard of deviation, P=significance value.

Table 4. Effect of *Tagetes erecta* Linn. leaf extract cream on DNA levels of wound healing in Wistar rats.

Variables	Groups	Days of observation		
		8	12	16
		Means ± SD	Means ± SD	Means ± SD
DNA Levels (mg/mL)	I	38.37 ± 0.44	29.33 ± 0.84	23.74 ± 0.71
	II	54.35 ± 0.47	42.15 ± 1.21	32.59 ± 1.00
	III	55.85 ± 0.85	45.94 ± 0.75	32.29 ± 0.76
	P	I<II<III (0.000)	I<II<III (0.000)	I<II, III (0.000)
				II=III (0.473)

Abbreviations: DNA=deoxyribo nucleic acid, Group I=Wistar rats were back skin incision and treated with nitrofurazone (positive control), group II=Wistar rats were back skin incision and treated with *T. erecta* Linn.leaf extract cream 3%, group III=Wistar rats were back skin incision and treated with *T. erecta* Linn.leaf extract cream 4%, SD=standard of deviation, mg/dL=milligram per deciliter, P=significance value.

Table 5. Effect of *Tagetes erecta* Linn. leaf extract cream on protein levels of wound healing in Wistar rats.

Variables	Groups	Days of observation		
		8	12	16
		Means ± SD	Means ± SD	Means ± SD
Protein	I	10.58 ± 0.42	14.39 ± 0.41	13.29 ± 0.43

levels (mg/mL)	II	16.66 ± 0.52	26.37 ± 0.58	21.53 ± 0.95
	III	18.19 ± 0.36	28.64 ± 0.41	22.34 ± 0.51
	P	I<II<III (0.000)	I<II<III (0.000)	I<II, III (0.000) II<III (0.013)

Abbreviations: Group I=Wistar rats were back skin incision and treated with nitrofurazone (positive control), group II=Wistar rats were back skin incision and treated with *T. erecta* Linn.leaf extract cream 3%, group III=Wistar rats were back skin incision and treated with *T. erecta* Linn.leaf extract cream 4%, SD=standard of deviation, mg/dL=milligram per deciliter, P=significance value.

Figure 1. Cultivation of *Tagetes erecta* Linn. in Bandungan District, Semarang Regency, Central Java province, Indonesia. A. *T. erecta* Linn. seedlings in a poly bag. B. *T. erecta* Linn. at a young age (flowers are still fresh). C. *T. erecta* Linn. with fresh flowers. D. *T. erecta* Linn. in old age (flowers have dried). Photographed by Hosea Jaya Edy on October 3, 2017.

Figure 2. Effect of *T. erecta* Linn.leaf extract cream on wound healing process in Wistar rats. Group I=Wistar rats were back skin incision and treated with nitrofurazone (positive control), group II=Wistar rats were back skin incision and treated with *T. erecta* Linn.leaf extract cream 3%, group III=Wistar rats were back skin incision and treated with *T. erecta* Linn.leaf extract cream 4%.

Figure 3. Histological observation of dermal wound healing process in Wistar rats (hematoxylin and eosin painting, magnification 40X). A. Wound healing in the early stage. B, C, and D. Wound healing in the middle stage. E. Wound healing in the advanced stage. F. Wound healing area (lime arrow).

Figure 4. Effect of *T. erecta* Linn. leaf extract cream on epithelialization period (EP) of dermal wound healing in Wistar rats. Group I=Wistar rats were back skin incision and treated with nitrofurazone (positive control), group II=Wistar rats were back skin incision and treated with *T. erecta* Linn.leaf extract cream 3%, group III=Wistar rats were back skin incision and treated with *T. erecta* Linn.leaf extract cream 4%, SD=standard of deviation, p=significance value, group I>II, III (P=0.000), while group II=III (P=0.232).

Figure 5. Effect of *T. erecta* Linn. leaf extract cream on the number of bacterial colonies (NBC) on the 6<sup>th</sup> day of wound healing in Wistar rats. Group I=Wistar rats were back skin incision and treated with nitrofurazone (positive control), group II=Wistar rats were back skin incision and treated with *T. erecta* Linn.leaf extract cream 3%, group III=Wistar rats were back skin incision and treated with *T. erecta* Linn.leaf extract cream 4%. NBC at the 6<sup>th</sup> day on wound healing in Wistar rats showed that group I>III>II (P=0.000)

# Efficacy of *Tagetes erecta* Linn. leaf extract cream on rat dermal wound healing

ORIGINALITY REPORT

# 18%

SIMILARITY INDEX

## PRIMARY SOURCES

1	<a href="http://www.ijbcp.com">www.ijbcp.com</a> Internet	216 words — 3%
2	<a href="http://www.apicareonline.com">www.apicareonline.com</a> Internet	57 words — 1%
3	<a href="http://academicjournals.org">academicjournals.org</a> Internet	46 words — 1%
4	Sumitra, M.. "Efficacy of <i>Butea monosperma</i> on dermal wound healing in rats", <i>International Journal of Biochemistry and Cell Biology</i> , 200503 Crossref	41 words — 1%
5	<a href="http://repository-tnmgrmu.ac.in">repository-tnmgrmu.ac.in</a> Internet	39 words — 1%
6	N. Duran, A. Koc, H. Oksuz, C. Tamer, Y. Akaydin, T. Kozlu, M. Çelik. "The protective role of topical propolis on experimental keratitis via nitric oxide levels in rabbits", <i>Molecular and Cellular Biochemistry</i> , 2006 Crossref	37 words — 1%
7	Ren-Chuen Tzou, Hsi-Peng Lu. "Exploring the emotional, aesthetic, and ergonomic facets of innovative product on fashion technology acceptance model", <i>Behaviour &amp; Information Technology</i> , 2009 Crossref	36 words — 1%
8	<a href="http://www.hindawi.com">www.hindawi.com</a> Internet	35 words — 1%

- 9 Sung In Kim, Seung Hyuk Yim, Dong Wan Sohn, Seung-Ju Lee, U-Syn Ha, Chang Hee Han, Yong-Hyun Cho. "Preventive Effect on Inflammation and Anti-Inflammatory and Anti-Microbial Effects of Garlic Extract in a Rat Model of Chronic Bacterial Prostatitis", Korean Journal of Urology, 2009  
Crossref 33 words — 1%
- 
- 10 [www.sphinxesai.com](http://www.sphinxesai.com)  
Internet 31 words — < 1%
- 
- 11 Miriyala Sumitra. "<i>Emblica officinalis</i> exerts wound healing action through up-regulation of collagen and extracellular signal-regulated kinases (ERK1/2)", Wound Repair and Regeneration, 01/2009  
Crossref 30 words — < 1%
- 
- 12 K. C. Ogbanya, C. A. Eze, S. V. O. Shoyinka. "Hematological and serum biochemistry comparison between cotton hair thread and silk suture on surgical skin wounds of rabbits", Comparative Clinical Pathology, 2013  
Crossref 24 words — < 1%
- 
- 13 [iiste.org](http://iiste.org)  
Internet 23 words — < 1%
- 
- 14 [www.scirp.org](http://www.scirp.org)  
Internet 23 words — < 1%
- 
- 15 [www.toxicolres.org](http://www.toxicolres.org)  
Internet 23 words — < 1%
- 
- 16 [dyuthi.cusat.ac.in](http://dyuthi.cusat.ac.in)  
Internet 22 words — < 1%
- 
- 17 [japsonline.com](http://japsonline.com)  
Internet 22 words — < 1%
- 
- 18 [www.mdpi.com](http://www.mdpi.com)  
Internet 21 words — < 1%
- 
- 19 Weiyou Wang, Honggao Xu, Hua Chen, Kedong Tai, Fuguo Liu,

Yanxiang Gao. "In vitro antioxidant, anti-diabetic and antilipemic potentials of quercetagenin extracted from marigold (*Tagetes erecta* L.) inflorescence residues", *Journal of Food Science and Technology*, 2016

Crossref

18 words — < 1 %

20 [biotechnologyforbiofuels.biomedcentral.com](http://biotechnologyforbiofuels.biomedcentral.com)

Internet

17 words — < 1 %

21 Hosea Jaya Edy, ML Edy Parwanto. "Pemanfaatan tanaman *Tagetes erecta* Linn. dalam kesehatan", *Jurnal Biomedika dan Kesehatan*, 2019

Crossref

17 words — < 1 %

22 [www.tandfonline.com](http://www.tandfonline.com)

Internet

17 words — < 1 %

23 Debjit Ghosh, Sumanta Mondal, K. Ramakrishna. "A topical ointment formulation containing leaves extract of *Aegialitis rotundifolia* Roxb., accelerates excision, incision and burn wound healing in rats", *Wound Medicine*, 2019

Crossref

16 words — < 1 %

24 [www.nbp.gov.pl](http://www.nbp.gov.pl)

Internet

15 words — < 1 %

25 Hakan Berument. "Inflation Dynamics and its Sources in the Ottoman Empire: 1586-1913", *International Review of Applied Economics*, 2007

Crossref

14 words — < 1 %

26 Miñones, Jose, Montserrat Muñoz, Jose Miñones Trillo, Isabel Haro, Maria Antonia Busquets, and M<sup>a</sup>Asunción Alsina Esteller. "Miscibility and Langmuir studies of the interaction of E2 (279-298) peptide sequence of Hepatitis G Virus/GB Virus-C with Dipalmitoyl Phosphatidyl Choline and Dimiristoyl Phosphatidyl Choline phospholipids.", *Langmuir*

Crossref

14 words — < 1 %

27 T. K. Lim. "Edible Medicinal And Non-Medicinal Plants", Springer

Science and Business Media LLC, 2014

Crossref

14 words — < 1%

28 [pharmascope.org](http://pharmascope.org)

Internet

13 words — < 1%

29 Muthukumar, Thangavelu, Kannan Anbarasu, Dharmalingam Prakash, and Thotapalli Parvathaleswara Sastry. "Effect of growth factors and pro-inflammatory cytokines by the collagen biocomposite dressing material containing *Macrotyloma uniflorum* plant extract—In vivo wound healing", *Colloids and Surfaces B Biointerfaces*, 2014.

Crossref

13 words — < 1%

30 Hosea Jaya Edy, Marchaban Marchaban, Subagus Wahyuono, Agung Endro Nugroho. "Pengujian Aktivitas Antibakteri Hidrogel Ekstrak Etanol Daun *Tagetes erecta* L", *Jurnal MIPA*, 2019

Crossref

13 words — < 1%

31 [oa.upm.es](http://oa.upm.es)

Internet

12 words — < 1%

32 Mojtaba Kalantar, Mehdi Goudarzi, Hossein Foruozandeh, Amir Siahpoosh, Mohammad Javad Khodayar, Sara Mahmoodi Koshkghazi. "The Topical Effect of *Capparis spinosa* L. Extract on Burn Wound Healing", *Jundishapur Journal of Natural Pharmaceutical Products*, 2016

Crossref

12 words — < 1%

33 [hdl.handle.net](http://hdl.handle.net)

Internet

11 words — < 1%

34 [jpet.aspetjournals.org](http://jpet.aspetjournals.org)

Internet

11 words — < 1%

35 [apsjournals.apsnet.org](http://apsjournals.apsnet.org)

Internet

10 words — < 1%

36 Mohammad Yaseen Khan, Saleh Abbas Ali, Kilambi Pundarikakshudu. " Wound healing activity of extracts derived from resin ", *Pharmaceutical Biology*, 2015

10 words — < 1%

- 
- 37 [docobook.com](http://docobook.com) 10 words — < 1%  
Internet
- 
- 38 Pengde Kang, Bin Shen, Jing Yang, Jingqiu Cheng, Fuxing Pei. "Repairing Defect and Preventing Collapse of Canine Femoral Head Using Titanium Implant Enhanced by Autogenous Bone Graft and rhBMP-2", *Connective Tissue Research*, 2009 10 words — < 1%  
Crossref
- 
- 39 Marcio Fronza, Guilherme F. Caetano, Marcel N. Leite, Claudia S. Bitencourt et al. "Hyaluronidase Modulates Inflammatory Response and Accelerates the Cutaneous Wound Healing", *PLoS ONE*, 2014 8 words — < 1%  
Crossref
- 
- 40 [ujpr.org](http://ujpr.org) 8 words — < 1%  
Internet
- 
- 41 Tobias Hirsch, Simin Limoochi-Deli, Armin Lahmer, Frank Jacobsen et al. "Antimicrobial Activity of Clinically Used Antiseptics and Wound Irrigating Agents in Combination with Wound Dressings", *Plastic and Reconstructive Surgery*, 2011 8 words — < 1%  
Crossref
- 
- 42 Jasvidianto C. Kotta, Agatha B. S. Lestari, Damiana S. Candrasari, Maywan Hariono. "Medicinal Effect, In Silico Bioactivity Prediction, and Pharmaceutical Formulation of *Ageratum conyzoides* L.: A Review", *Scientifica*, 2020 8 words — < 1%  
Crossref
- 
- 43 [www.ijppsjournal.com](http://www.ijppsjournal.com) 8 words — < 1%  
Internet
- 
- 44 [journals.usm.ac.id](http://journals.usm.ac.id) 8 words — < 1%  
Internet
- 
- 45 A. Ruiz-Extremera, E. Cervilla, P. Munoz-Rueda, E. Ocete, A. Palacios, M.A. Lopez-Garrido, J. Salmeron. 8 words — < 1%



"Importance of HGV and TTV coinfection in the vertical transmission of HCV", Journal of Hepatology, 2003

Crossref

46 P. Marie Arockianathan, S. Sekar, S. Sankar, B. Kumaran, T.P. Sastry. "Evaluation of biocomposite films containing alginate and sago starch impregnated with silver nano particles", Carbohydrate Polymers, 2012

8 words — < 1%

Crossref

47 Sara E. Bianchi, Barbara E.K. Machado, Marília G.C. da Silva, Michelle M.A. da Silva et al. "Coumestrol/hydroxypropyl-β-cyclodextrin association incorporated in hydroxypropyl methylcellulose hydrogel exhibits wound healing effect: in vitro and in vivo study", European Journal of Pharmaceutical Sciences, 2018

8 words — < 1%

Crossref

48 [estudogeral.sib.uc.pt](http://estudogeral.sib.uc.pt)

Internet

8 words — < 1%

49 [ukaazpublications.com](http://ukaazpublications.com)

Internet

8 words — < 1%

50 María G. Peralta-Sánchez, Fernando C. Gómez-Merino, Olga Tejeda-Sartorius, Gabriel Alcántar-González et al. "Nitrogen supply and shading affect morphology and composition of the essential oil in marigold (Tagetes erecta L.)", Folia Horticulturae, 2020

8 words — < 1%

Crossref

51 E. G. Karapanagioti, A. N. Assimopoulou. "Naturally Occurring Wound Healing Agents: An Evidence-Based Review", Current Medicinal Chemistry, 2016

7 words — < 1%

Crossref

52 Yuri Akishima. "Role of macrophage and smooth muscle cell apoptosis in association with oxidized low-density lipoprotein in the atherosclerotic development", Modern Pathology, 03/2005

6 words — < 1%

Crossref

---

EXCLUDE QUOTES

ON

EXCLUDE MATCHES

OFF

EXCLUDE  
BIBLIOGRAPHY

ON