

Influence of Extraction Solvents on Capsaicin Compound and Cytotoxic Activity of Piper retrofractum Vahl

by Cek Turnitin

Submission date: 16-Mar-2023 01:06PM (UTC+0800)

Submission ID: 2038341351

File name: Cabe_Jawa_Revisi.docx (60K)

Word count: 3270

Character count: 18876

Influence of Extraction Solvents on Capsaicin Compound and Cytotoxic Activity of *Piper retrofractum Vahl*

Kirana Anggraini^{1,a)}, Dyah Ayu Woro Setyaningrum^{2,b)}, Laela Wulansari^{3,c)},
Hening Tyas Andayani⁴⁾, Laviany Putri Shihran⁵⁾, Isra Fauziyyah⁵⁾

14

Author Affiliations

¹Department of Obstetrics and Gynecology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

²Department of Anatomical Pathology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

³Tropical Biopharmaceutical Research Center, Institute for Research and Community Service (LPPM)
IPB University, Indonesia

⁴Fachärztin der Innere Medizin, Städtische Kliniken Mönchengladbach, Germany

⁵Medical Science Study Program, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

^{a)} Corresponding author: kirana_anggraeni@trisakti.ac.id

^{b)} dyahayu94@trisakti.ac.id

^{c)} laela2002@gmail.com

Abstract. Capsaicin (8-Methyl-N-vanillyl-trans-6-nonenamide) is an active compound found in plants from the genus *Capsicum*. Genus *Capsicum* possesses different species of red chilli (*Capsicum annuum Linn*), green chilli (*Capsicum annuum var. annuum*) and other varieties that are familiar in Indonesia. However, the plants were not originally from Indonesia but from American Continent. Indonesian ancestor utilized chilli to add spiciness on foods which refers to *Piper retrofractum Vahl* or Javanese chilli. Capsaicin has positive effects on health such as analgesic, cardioprotective, anorexigenic, chemopreventive, chemotherapy, and cardioprotective effects. In addition, capsaicin has beneficial effects in maintaining glucose levels, insulin homeostasis, reducing itching and symptoms in Non-Allergic Rhinitis (NAR), and an alternative therapy for neurogenic bladder. This research is aimed to acknowledge if capsaicin is also discovered in Javanese chilli. The method used to obtain the extract was by maceration of Java chillies using 30%, 70%, and 96% ethanol, n-hexane, ethyl acetate, and water as solvents. The research was started by conducting phytochemical and moisture content tests on simplicia. Then, each extract was tested for cytotoxic activity on shrimp larvae and the determination of capsaicin levels by High Performance Liquid Chromatography (HPLC). The results showed that the highest yield of Java chili extract was 51.14% in ethanol 70%. Phytochemical testing of Java chili simplicia contains flavonoids and saponins. The ethanol 70% extract had the highest

cytotoxic activity with an LC50 value of 205.21 ppm. The highest concentration of capsaicin was found in ethyl acetate solvent of 785.58 mg/g followed by ethanol 70% at 486.06mg/g.

7 INTRODUCTION

Capsaicin (8-Methyl-N-vanillyl-trans-6-nonenamide) is the active compound found in hot peppers that gives chilies their hot taste¹. Capsaicin was first purified in 1876 but its structure began to be described in 1919². Due to its chemical structure, capsaicin is well absorbed up to 94% when administered topically or orally³. The use of spice plants is also influenced by the level of knowledge, age, education level, economic status, environmental factors, and sources of information/information media⁴. Studies demonstrate the efficacy of capsaicin as an analgesic. Treatment with capsaicin is effective in various types of pain conditions such as complex regional pain syndromes, neuropathic pain postsurgical neuropathic pain⁵ post-herpetic neuralgia⁶ and painful diabetic peripheral neuropathy⁷. Research has also shown capsaicin to be effective in weight loss and obesity improvement⁸. Capsaicin also has beneficial effects on glucose levels, insulin homeostasis, and diabetes⁹.

The evidence showed the beneficial effects of capsaicin on the cardiovascular system¹⁰. Capsaicin activates TRPV1 which can stimulate the release of CGRP, which is the most powerful vasodilator that regulates blood pressure in both physiological and pathophysiological conditions, producing a blood pressure-lowering effect¹¹. Capsaicin has been shown to inhibit platelet aggregation¹², which may also provide protection against cardiovascular disease¹³. Long-term activation of TRPV1 by capsaicin will inhibit the formation of foam cells so that it will reduce lipid storage and atherosclerotic lesions in the aortic sinus and thoracoabdominal aorta of rats, and ultimately slow down the process of atherosclerosis. The antioxidant properties of capsaicin contribute to its protective effect on the cardiovascular system. LDL oxidation is an early factor for the formation and development of atherosclerosis¹⁴.

Capsaicin has been shown to have chemopreventive and chemotherapeutic effects¹⁵. Capsaicin also works on itching with nostalgia paresthetica¹⁶, and neuropathic pruritus¹⁷. Capsaicin has a gastroprotective effect by modulating sensory neurons¹⁸. Capsaicin has been studied as an alternative therapy for symptomatic relief of neurogenic bladder, a urological disorder that seriously affects patients' quality of life¹⁹. The clinical use of capsaicin is also

successful in patients with detrusor bladder hyperreflexia with multiple sclerosis, as well as after spinal cord injury²⁰.

Capsaicin is discovered in various type of capsium genus plants such as red chilli (*Capsicum annuum Linn*), green chilli (*Capsicum annuum var.annuum*) and other varieties known in Indonesia. Capsicum is not originally from Nusantara, instead it comes from American Continent. Indonesian ancestors used chilli to add spiciness on food currently refers to Piper retrofractum Vahl or green chilli. This research was conducted to determine the capsaicin content in Javanese chilies, as well as to find out the best solvent in the extraction process from Javanese chilies which produced the highest capsaicin levels.

METHOD

Place, Collecting, and Determining Samples. The research material was the fruit of the Javanese chili plant taken from the Biopharmaca Cultivation Conservation Unit (BCCU) of the Tropical Biopharmaca Research Center, LPPM IPB, which was obtained from Java which was harvested at the age of 3-4 months after flowering. The determination was carried out at the Biopharmaca Cultivation Conservation Unit (BCCU).

Simplicia Making. Three kilograms of fresh Java chilies were weighed, then washed and dried in direct sunlight for 4-5 days. After drying, the dry sorting is carried out, and mashed with a blender. The simplicia powder obtained was sieved using an 80-mesh sieve and then weighed. The simplicia powder is then stored in a clean, dry container and protected from sunlight for the next extraction process.

Production of Java Chilli Extract. Extract preparation and testing were carried out at the Laboratory of the Tropical Biopharmaca Research Center, Institute for Research and Community Service (LPPM), IPB University. Samples of Java chili simplicia were weighed for extraction with 10 grams of various solvents each, then added solvents namely 96%, 70%, 30% ethanol, 500 mL of water, ethyl acetate, and n-hexane. Maceration was carried out 2 x 24 hours with several times of stirring, then it was filtered. The collected filtrate was concentrated using a vacuum rotary evaporator at 45-50°C to obtain a viscous ethanol extract of 96%, 70%, 30%, water, ethyl acetate, and n-hexane extract.

Water content. Two grams of Javanese chili simplicia were weighed in a container with a constant weight. Then, it was heated in an oven at + 105 degrees Celsius for 3 hours. After that, it was cooled in a desiccator and weighed until it reached a constant weight.

Yield Analysis of Javanese Chilli Extract. The yield of Javanese chili extract was calculated by comparing the weight of the Javanese chili extract with the weight of the Javanese chili simplicia used for extraction.

Phytochemical Screening. Phytochemical screenings were carried out for alkaloids, flavonoids, tannins, saponins, triterpenoids, and steroids based on the method of Harborne 1987.

Alkaloids. A total of 0.5 grams of condensed extract or 1 gram of simplicia was dripped with 3-5 drops of ammonia, then 5 mL of chloroform was added, then it was homogenized and filtered. The filtrate obtained was added with 2M Sulfuric acid reagent, then homogenized. The top layer was taken and used as an experimental solvent which was then treated as follows: 1) Experimental solvent 1 was added with 2 drops of Mayer's reagent, a positive result indicated by the formation of a white precipitate. 2) Experiment solvent 2 was added with 2 drops of Dragendorf reagent, and a positive result was indicated by the formation of an orange precipitate. 3) Experiment solvent 3 was added with 2 drops of Wagner reagent, and a positive result was indicated by the formation of a brown precipitate.

Flavonoids, Tannins, and Saponins. 0.5 grams of condensed extract or 5 grams of simplicia was dissolved in distilled water and heated for 5 minutes. Then the solvent was filtered and the filtrate was divided into 3 parts. To test the flavonoid filtrate, Mg powder and HCl:Ethanol (1:1) were added, followed by amyl alcohol. The color formed on the amyl alcohol layer was observed, if a yellow, orange or red color is formed then it consists of flavonoids. For the tannin filtrate test, 3 drops of 10% FeCl₃ were added, if a greenish-black color is formed then it consists of tannins. For the saponin test, the filtrate was shaken for 10 seconds. A positive result is indicated by the formation of stable foam for over 2 minutes.

Triterpenoids and Steroids. 0.5 grams sample of condensed extract or 1 gram of simplicia was dissolved in ethanol and heated for 5 minutes, then the sample was filtered into a porcelain dish. The filtrate was heated to dryness and then 1 mL of diethyl ether, 1 drop of acetic anhydrous, and 1 drop of concentrated sulfuric acid were added. A positive reaction is indicated by the formation of a red/purple solvent for triterpenoids and blue or green for steroids.

Quinone. 0.5 grams of condensed extract or 1 gram of simplicia was added to methanol then

it was heated and filtered. The filtrate results were added with 3 drops of 10% NaOH. A positive reaction is indicated by the formation of a red color for hydroquinone.²¹

Cytotoxic Activity Test on Shrimp Larvae. The cytotoxic activity test by determining the LC₅₀ value was carried out using *Artemia salina* shrimp eggs. *A. salina* used for the toxicity test was obtained from hatching using seawater with the help of an aerator to meet dissolved oxygen levels. The extract toxicity test was carried out using *A. salina* shrimp larvae. The shrimp larvae used were aged 48 hours after the shrimp larvae hatched. *A. salina* cysts of as much as ± 50 mg were put into a container containing seawater that had been filtered and equipped with an aerator. The cysts were left for 48 hours under light to hatch completely. After hatching, 10 *A. salina* larvae were put into a 2 ml vial, then it was added a stock extract solution with a concentration of 4000 ppm and adjusted the volume with seawater so that the final concentration of the extract was 0, 10, 100, and 1000 ppm. After 24 hours, the number of dead larvae was counted. The lethal concentration (LC) value was determined by the probit analysis method with a 95% confidence interval.²²

Determination of Capsaicin content with High Performance Liquid Chromatography (HPLC)

Standard and sample preparation. Standard capsaicin was dissolved in methanol to make a concentration of 100 ppm. Each extract was weighed as much as 0.1 gram and then 8 mL of methanol solvent was added and sonicated for 1 hour. Then the sample solution was filtered into a 10 mL flask and calibrated with methanol up to 10 mL. After that, it was filtered with 0.45 micrometer Whatman filter paper and injected into HPLC as much as 20 µL.

Identification by HPLC. The mobile phase used was methanol and aquabides with a composition ratio of 80:20. The wavelength was 235 nm, while the flow of the mobile phase was 1 mL/minute.

RESULT AND DISCUSSION

Plant Determination. The test sample was identified at the Biopharmaca Cultivation and Conservation Unit (BCCU) of the Tropical Biopharmaca Research Center, Institute for Research and Community Service (LPPM) IPB University, it showed that a sample of *Piper retrofractum* Vahl was from the Piperaceae tribe.

Phytochemical compounds. Tests for the content of phytochemical compounds were carried out on simplicia, which contained saponin and flavonoid. The positive reactions in the flavonoids, saponins, and tannins indicate the presence of phenol groups

Water Content and Extraction. The water content results of the Javanese chili simplicia were 8.75%, fulfilling the quality requirements and can be used for further analysis. Removing the water content up to a certain amount is useful for extending the durability of simplicia. Extremely high water content can become a medium for the growth of microorganisms that cause damage to the simplicia.²³

Rhizome Content on Different Types of Extract. The yield of Javanese chili extract is presented in table 1, where the highest yield was found in 70% ethanol extract.

Table 1. The yield of Javanese chili extract

	ETOH 30%	ETOH 70%	ETOH 96%	Water	Ethyl acetate	N hexane
Extract(g)	7.59	51.14	8.44	11.85	6.33	8.30

Shrimp Larvae Cytotoxic Activity Test on Different Types of Extracting Solvent. Shrimp larvae toxicity tests were carried out to observe the potential bioactivity and cytotoxic activity of each extract so that a safe extract concentration could be determined for further testing. In addition, extracts that exhibit toxic properties can be developed as anti-cancer drugs²⁴. A plant extract will be bioactive and have anti-cancer potential if it has an LC₅₀ value of less than 1000 ppm. Based on Table 2, it can be seen that all Javanese chili extracts have the potential as bioactive compounds and can be used as medicine. It is due to the fact that each extract produced an LC₅₀ of less than 1000 ppm so that, at low concentrations, it was able to kill 50% of the population of *A. salina* shrimp larvae. The highest bioactive potential and the toxic extract was 70% ethanol extract because it had the lowest LC₅₀ value, which was 205.21 ppm, which means that, at a small concentration, this extract can kill half the population of *A. salina* shrimp larvae. The LC₅₀ value is the highest concentration limit for determining various extract concentrations in subsequent tests.²¹

Table 2. LC₅₀ value of Javanese chili extract on A. Salina larvae

	ETOH 30%	ETOH 70%	ETOH 96%	Water	Ethyl acetate	Water
LC ₅₀ (ppm)	434.57	205.21	334.32	650.18 6	443.03	292.78

Capsaicin Content on Different Types of Extract. The results of the calculation of the capsaicin compound concentration in each extract can be seen in Figure 1. The highest capsaicin content was found in ethyl acetate solvent of 785.53mg/g followed by 70% ethanol of 426.34mg/g and the lowest in water solvent of 6.55mg/g. It is due to the fact that capsaicin is a non polar phenolic compound so it cannot dissolve in water. The main solvents used to extract and maintain the properties of capsaicin are nonpolar solvents such as ether, benzene, dimethyl sulfoxide, and acetone, as well as ethanol due to their mixed nature²⁵.

The content of capsaicin in each extract of Javanese chilies in this study was higher than in previous studies conducted in Nepal and Saudi Arabia. In Nepal, research conducted on 16 types of chilies (Capsium fruit) found capsaicin levels ranging from 2.19-19.73 mg/g²⁶. While research conducted in Saudi Arabia on hot chili, red chili, green chili, green pepper, red pepper, and yellow pepper found capsaicin levels range from 0-4.24mg/g²⁷. The difference related to the content of capsaicin in this study and the previous study could be due to differences in plant ecological conditions, varieties, harvesting ages, simplicia manufacturing methods, and extraction methods used.




Figure 1. Graph of capsaicin levels of various Javanese chili extracts

CONCLUSION

Capsaicin is discovered on *Piper retrofractum Vahl* fruits. The secondary metabolites in Javanese chili simplicia are flavonoids and saponins. The highest yield of extract was 11.85 g in a water solvent. The highest bioactive potential and cytotoxic activity extract is a 70% ethanol extract with an LC50 of 205.21 ppm. The highest content of capsaicin was found in ethyl acetate solvent of 785.53mg/g followed by ethanol 70% at 486.06mg/g.

ACKNOWLEDGMENT

We thank Trisakti University for funding this research through the Faculty Research grant (PUF) Number: 322/A.1/LPT/USAKTI/II/2022 on behalf of dr. KiranaAnggraini MKM

REFERENCES

1. Nelson, E. K., & Dawson, L. E. (1923). The constitution of capsaicin, the pungent principle of Capsicum. III. *Journal of the American Chemical Society*, 45(9), 2179-2181.
2. Nelson, E. K. (1919). The constitution of capsaicin, the pungent principle of capsicum. *Journal of the American Chemical Society*, 41(7), 1115-1121.
3. Suresh, D., & Srinivasan, K. (2010). Tissue distribution & elimination of capsaicin, piperine & curcumin following oral intake in rats. *Indian Journal of Medical Research*, 131(5), 682-691.
4. Adiyasa, M. R., & Meiyanti, M. (2021). Pemanfaatan obat tradisional di Indonesia: distribusi dan faktor demografis yang berpengaruh. *Jurnal Biomedika Dan Kesehatan*, 4(3), 130-138.
5. Kingery, W. S. (1997). A critical review of controlled clinical trials for peripheral neuropathic pain and complex regional pain syndromes. *Pain*, 73(2), 123-139.

6. Zis, P., Apsokardos, A., Isaia, C., Sykioti, P., & Vadalouca, A. (2014). Posttraumatic and postsurgical neuropathic pain responsive to treatment with capsaicin 8% topical patch. *Pain Physician*, *17*(2), E213.
7. Kiani, J., Sajedi, F., Nasrollahi, S. A., & Esna-Ashari, F. (2015). A randomized clinical trial of efficacy and safety of the topical clonidine and capsaicin in the treatment of painful diabetic neuropathy. *Journal of Research in Medical Sciences: the Official Journal of Isfahan University of Medical Sciences*, *20*(4), 359.
8. Yu, Q., Wang, Y., Yu, Y., Li, Y., Zhao, S., Chen, Y., ... & Liu, E. (2012). Expression of TRPV1 in rabbits and consuming hot pepper affects its body weight. *Molecular biology reports*, *39*, 7583-7589.
9. Yuan, L. J., Qin, Y., Wang, L., Zeng, Y., Chang, H., Wang, J., ... & Mi, M. T. (2016). Capsaicin-containing chili improved postprandial hyperglycemia, hyperinsulinemia, and fasting lipid disorders in women with gestational diabetes mellitus and lowered the incidence of large-for-gestational-age newborns. *Clinical nutrition*, *35*(2), 388-393.
10. Harada, N., & Okajima, K. (2009). Effects of capsaicin and isoflavone on blood pressure and serum levels of insulin-like growth factor-I in normotensive and hypertensive volunteers with alopecia. *Bioscience, biotechnology, and biochemistry*, *73*(6), 1456-1459.
11. Deng, P. Y., & Li, Y. J. (2005). Calcitonin gene-related peptide and hypertension. *Peptides*, *26*(9), 1676-1685.
12. Adams, M. J., Ahuja, K. D., & Geraghty, D. P. (2009). Effect of capsaicin and dihydrocapsaicin on in vitro blood coagulation and platelet aggregation. *Thrombosis research*, *124*(6), 721-723.
13. Bhatt, D. L., & Topol, E. J. (2003). Scientific and therapeutic advances in antiplatelet therapy. *Nature Reviews Drug Discovery*, *2*(1), 15-28.
14. Stocker, R., & Keane Jr, J. F. (2004). Role of oxidative modifications in atherosclerosis. *Physiological reviews*, *84*(4), 1381-1478.
15. Zhang, R., Humphreys, I., Sahu, R. P., Shi, Y., & Srivastava, S. K. (2008). In vitro and in vivo induction of apoptosis by capsaicin in pancreatic cancer cells is mediated through ROS generation and mitochondrial death pathway. *Apoptosis*, *13*, 1465-1478.
16. Andersen, H. H., Sand, C., & Elberling, J. (2016). Considerable variability in the efficacy of 8% capsaicin topical patches in the treatment of chronic pruritus in 3 patients with notalgia paresthetica. *Annals of Dermatology*, *28*(1), 86-89.
17. Misery, L., Erfan, N., Castela, E., Brenaut, E., Lantéri-Minet, M., Lacour, J. P., & Passeron, T. (2015). Successful treatment of refractory neuropathic pruritus with capsaicin 8% patch: a bicentric retrospective study with long-term follow-up. *Acta dermato-venereologica*, *95*(7), 864-865.
18. Holzer, P., & Sametz, W. (1986). Gastric mucosal protection against ulcerogenic factors in the rat mediated by capsaicin-sensitive afferent neurons. *Gastroenterology*, *91*(4), 975-981.
19. Foster, H. E., & Lake, A. G. (2014). Use of vanilloids in urologic disorders. *Capsaicin as a therapeutic molecule*, 307-317.
20. de Sèze, M., Wiart, L., Joseph, P. A., Dosque, J. P., Mazaux, J. M., & Barat, M. (1998). Capsaicin and neurogenic detrusor hyperreflexia: a double-blind placebo-controlled study in 20 patients with spinal cord lesions. *Neurourology and Urodynamics: Official Journal of the International Continence Society*, *17*(5), 513-523.
21. Harborne, A. J. (1998). *Phytochemical methods a guide to modern techniques of plant analysis*. springer science & business media.
22. Meyer, B. N., Ferrigni, N. R., Putnam, J. E., Jacobsen, L. B., Nichols, D. J., & McLaughlin, J. L. (1982). Brine shrimp: a convenient general bioassay for active plant constituents. *Planta medica*, *45*(05), 31-34.

23. Yuslianti, E. R., Bachtiar, B. M., Suniarti, D. F., & Sutjiatmo, A. B. (2016). Standardisasi farmasitikal bahan alam menuju fitofarmaka untuk pengembangan obat tradisional indonesia. *Dentika dental journal*, *19*(2), 179-185.
24. Carballo, J. L., Hernández-Inda, Z. L., Pérez, P., & García-Grávalos, M. D. (2002). A comparison between two brine shrimp assays to detect in vitro cytotoxicity in marine natural products. *BMC biotechnology*, *2*(1), 1-5.
25. Suresh, D., & Srinivasan, K. (2010). Tissue distribution & elimination of capsaicin, piperine & curcumin following oral intake in rats. *Indian Journal of Medical Research*, *131*(5), 682-691.
26. Thapa, B., Skalko-Basnet, N., Takano, A., Masuda, K., & Basnet, P. (2009). High-performance liquid chromatography analysis of capsaicin content in 16 Capsicum fruits from Nepal. *Journal of Medicinal Food*, *12*(4), 908-913.
27. Othman, Z. A. A., Ahmed, Y. B. H., Habila, M. A., & Ghafar, A. A. (2011). Determination of capsaicin and dihydrocapsaicin in Capsicum fruit samples using high performance liquid chromatography. *Molecules*, *16*(10), 8919-8929.

Influence of Extraction Solvents on Capsaicin Compound and Cytotoxic Activity of Piper retrofractum Vahl

ORIGINALITY REPORT

15%

SIMILARITY INDEX

13%

INTERNET SOURCES

10%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|---|----|
| 1 | www.ncbi.nlm.nih.gov
Internet Source | 4% |
| 2 | Submitted to Iowa State University
Student Paper | 1% |
| 3 | Muhammad A. Kurnianto, Harsi D. Kusumaningrum, Hanifah N. Lioe, Ekowati Chasanah. "Partial Purification and Characterization of Bacteriocin-Like Inhibitory Substances Produced by Streptomyces sp. Isolated from the Gut of Chanos chanos", BioMed Research International, 2021
Publication | 1% |
| 4 | www.pubfacts.com
Internet Source | 1% |
| 5 | Dwi Atty Mardiana, Burhanudinnur, R. S. Trijana Kartoatmodjo. "Analysis of extensive use of variable split components on flexible gross split scheme", AIP Publishing, 2020
Publication | 1% |

6	Submitted to Fundação Escola Americana de Lisboa/Carlucci American International School of Lisbon Student Paper	1 %
7	Karl-Erik Andersson, Delphine Behr-Roussel, Pierre Denys, Francois Giuliano. "Acute Intravesical Capsaicin for the Study of TRPV1 in the Lower Urinary Tract: Clinical Relevance and Potential for Innovation", Medical Sciences, 2022 Publication	<1 %
8	Zhang, Ming-Jie, Yan-Wei Yin, Bing-Hu Li, Yun Liu, Shao-Qiong Liao, Chang-Yue Gao, Jing-Cheng Li, and Li-Li Zhang. "The role of TRPV1 in improving VSMC function and attenuating hypertension", Progress in Biophysics and Molecular Biology, 2015. Publication	<1 %
9	journal.uad.ac.id Internet Source	<1 %
10	www.researchgate.net Internet Source	<1 %
11	www.science.gov Internet Source	<1 %
12	Submitted to The College of New Jersey Student Paper	<1 %

13	www.omicsonline.org Internet Source	<1 %
14	librarysearch.aut.ac.nz Internet Source	<1 %
15	fst.ipb.ac.id Internet Source	<1 %
16	eprints.ugd.edu.mk Internet Source	<1 %
17	Adam Tobolka, Tereza Škorpilová, Zuzana Dvořáková, Eloy Fernández Cusimamani, Aleš Rajchl. "Determination of capsaicin in hot peppers (<i>Capsicum</i> spp.) by direct analysis in real time (DART) method", <i>Journal of Food Composition and Analysis</i> , 2021 Publication	<1 %
18	biblio.ugent.be Internet Source	<1 %
19	dk.um.si Internet Source	<1 %
20	eprints.utas.edu.au Internet Source	<1 %
21	www.mdpi.com Internet Source	<1 %
22	www.researchsquare.com Internet Source	<1 %

23

Fang Sun, Shiqiang Xiong, Zhiming Zhu.
"Dietary Capsaicin Protects Cardiometabolic
Organs from Dysfunction", *Nutrients*, 2016
Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On