

High-density Lipoprotein Cholesterol as a Risk Factor of Health-Related Quality of Life in 50–70-Year-Old Community-Dwelling Women

By Yenny Yenny



High-density Lipoprotein Cholesterol as a Risk Factor of Health-Related Quality of Life in 50–70-Year-Old Community-Dwelling Women

Raditya Wratsangka^{1*}, Elly Herwana², Yenny Yenny², Endrico Xavieres³, Aditya Krishna⁴

¹Department of Obstetrics and Gynecology, Faculty of Medicine, Universitas Trisakti, West Jakarta, Indonesia; ²Department of Pharmacology and Therapeutics, Faculty of Medicine, Universitas Trisakti, West Jakarta, Indonesia; ³Department of Biochemistry, Faculty of Medicine, Universitas Trisakti, West Jakarta, Indonesia; ⁴Department of Physiology, Faculty of Medicine, Universitas Trisakti, West Jakarta, Indonesia

Abstract

Citation: Wratsangka R, Herwana E, Yenny Y, Xavieres E, Krishna A. High-density Lipoprotein Cholesterol as a Risk Factor of Health-Related Quality of Life in 50–70-Year-Old Community-Dwelling Women. Open Access Maced J Med Sci. 2021 Nov 06; 9(E):1092-1096. <https://doi.org/10.3889/oamjms.2021.7466>

Keywords: Community-dwelling; High-density lipoprotein cholesterol; Health-related quality of life; 50–70-year-old women

***Correspondence:** Raditya Wratsangka, Department of Obstetrics and Gynecology, Faculty of Medicine, Universitas Trisakti, Indonesia. E-mail: raditya@trisakti.ac.id
Received: 28-Sep-2021
Revised: 24-Oct-2021
Accepted: 27-Oct-2021

Copyright: © 2021 Raditya Wratsangka, Elly Herwana, Yenny Yenny, Endrico Xavieres, Aditya Krishna

Funding: This study was wholly funded by the research budget of Universitas Trisakti (2019-2020)

Competing Interests: The authors have declared that no competing interests exist

Open Access: This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

BACKGROUND: The prevalence of dyslipidemia, a risk factor of cardiovascular disease, is at present sufficiently high, particularly in the elderly. Health-related quality of life (HRQoL) is also an important outcome in the treatment of dyslipidemia, which is currently more targeted at lowering the low-density lipoprotein cholesterol (LDL-C) concentration.

AIM: The aim of this study was to determine the relationship between the high-density lipoprotein cholesterol (HDL-C) concentration and HRQoL in community-dwelling women aged 50–70 years.

METHODS: A cross-sectional study involving women aged 50–70 years who underwent blood sampling for determining the concentrations of hemoglobin, total cholesterol (TC), LDL-C, and HDL-C. A questionnaire was given to all subjects regarding data on age, menopausal status, level of education, and also another question of SF-36 for evaluation HRQoL. Bivariate and multivariate logistic regression analysis was used to identify risk factors associated with HRQoL.

RESULTS: In bivariate logistic regression analysis, age, menopausal status, education level, anemia, TC, and LDL-C were found to have non-statistically significant association with HRQoL, but HDL-C was significantly associated with HRQoL (odds ratio = 0.44; 95% Confidence interval [CI] = 0.20–0.97; $p = 0.042$). In multivariate analysis, menopausal status and education level have a non-statistically significant association with HRQoL, but those normal level HDL cholesterol was found 0.18 times less likely to be associated with poor HRQoL (Adjusted odds ratio 0.18; 95% CI = 0.03–0.91; $p = 0.038$).

CONCLUSION: This study showed that HDL-C levels were risk factors of HRQoL in women 50–70 years of age.

Introduction

The health-related quality of life (HRQoL) is currently an important outcome that is reported by patients during or after treatment of a disorder, in addition to increasing the life expectancy in such chronic conditions as hypertension, cardiac failure, and diabetes [1]. Various health problems may occur with increasing age of an individual, and the life style changes that occur in old age may result in dyslipidemia. The prevalence of dyslipidemia in subjects aged >60 years is around 56.8%, with high total cholesterol (TC), high low-density lipoprotein cholesterol (LDL-C), low high-density lipoprotein cholesterol (HDL-C), and high triglycerides prevalence rates of 8.4%, 13.9%, 23.1%, and 11.4%, respectively [2].

Dyslipidemia, especially of LDL-C, has a causal relationship with atherosclerotic cardiovascular disease according to genetic, observational, and clinical

outcome studies [3], [4]. Several epidemiological studies have also demonstrated that a low plasma HDL-C concentration is a risk factor of cardiovascular disease. The HDL-C concentration may predict cardiovascular incidence, even in patients who have been treated with statins, but clinical studies on the relationship of increased HDL-C concentration with cardiovascular protection are not convincing [5], [6].

The results of the previous studies demonstrate a still inconsistent relationship between HDL-C concentration and quality of life. Studies conducted on a group of male and female subjects aged 50–79 years showed that a decrease in LDL-C concentration and an increase in HDL-C concentration are associated with increased HRQoL [7]. However, inconsistent results were found in a study involving subjects aged 100 years and over, showing that HDL-C concentration is not significantly associated with quality of life [8]. Therefore, a study was performed to determine plasma HDL-C concentration as a risk factor of HRQoL in

community-dwelling women aged between 50 and 70 years.

Material and Methods

21

Study design and participants

This was a cross-sectional study conducted in five villages (*kelurahan*) of Mampang Prapatan Sub-district, South Jakarta, from March to July 2020. The sample size was calculated based on a correlation between HDL-C concentration and HRQoL of 0.34 (data from a preliminary study using 40 women aged 50–70 years), $\alpha = 0.05$ and $\beta = 0.1$, from which a sample size of 113 subjects was found. With an anticipated study drop-out rate of 10.5%, the optimal sample size was found to be 130. The inclusion criteria were women aged between 50 and 70 years, agreeing to undergo a complete blood examination, capable of good communication, and agreeing to sign informed consent. Women with neurological disorders (stroke, epilepsy, Parkinson's disease, and dementia), and psychiatric disorders (major depression and psychosis) were excluded from the study. Selection of the study subjects was by consecutive non-random sampling.

Informed consent was signed by the subjects after having received oral and written information on the general outlines of the study. For the protocol of this study, ethical clearance was obtained from the Research Ethics Commission, Faculty of Medicine, Universitas Trisakti under no. 160/KER/FK/II/2020.

Questionnaire

To all subjects, a questionnaire was given regarding data on socio-demographic characteristics comprising age, menopausal status, and level of education. The variable of age, menopause status, and education level was each categorized into two groups: (1) Age: (a) <60 years and (b) ≥60 years; (2) menopause status: (a) Premenopause and (b) postmenopause; and (3) education level: (a) Low and (b) moderate/high.

HRQoL

Evaluation of the HRQoL was performed using the SF-36 questionnaire consisting of 36 questions or statements. The filling-in of the questionnaire was done with the assistance of the enumerators in interviews with the subjects. There was also an evaluation of the physical component summary (PCS) with four subcomponents (physical functioning, physical limitations, bodily pain, and general health); and a mental component summary (MCS) with four subcomponents (vitality, social functioning, emotional

27

limitations, and mental health). The quality of life (HRQoL) of the subjects was considered to be good if the SF-36 score was above the median value, and to be poor if the score was under the median value for SF-36 total score, SF-36 PCS, and SF-36 MCS as well as each of the subcomponents [9], [10], [11].

Laboratory analysis

The subjects were required to fast for 10–12 h from 20:00 h of the preceding night. On the following day, 14 mL of venous blood was drawn from the right cubital vein of each of the subjects. The hemoglobin (Hb) concentration was determined by the cyanide-free sodium lauryl sulfate Hb detection method using an automated hematological analyzer (type KX-21N, Sysmex, Kobe, Japan). TC and triglyceride concentrations were determined by means of an enzymatic-colorimetric method, while measurement of LDL-C and HDL-C concentrations was by means of the homogenous enzymatic-colorimetric method using the Cobas 311 instrument (Roche Products Ltd, Basel, Switzerland). These variables of laboratory results were each also categorized into two groups: (1) Hb concentration (g/dL): (a) Normal: ≥12 and (b) low (anemia): <12; (2) TC (mg/dL): (a) Normal: <200 and (b) high: ≥200; (3) LDL-C concentration (mg/dL): (a) Normal: <100 and (b) high ≥100; and (4) HDL-C concentration (mg/dL): (a) Normal: ≥40 and (b) low: <40.

Quality control

All the enumerators were trained on interview technique. The questionnaire was collected on the spot after completion and uniformly numbered in the order of site and recall questionnaire. The laboratory examination was performed by laboratory technicians (who were blinded as to the identity of the subjects to whom the blood samples belonged) at the Prodia Clinical Laboratory–Jakarta that had obtained the SNI certification ISO 15189:2012 as a medical laboratory with regard to quality requirements and competence. The data were checked to avoid inconsistency.

28

Statistical analysis

The SPSS version 25 was used for data processing. Descriptive analysis was presented frequency for categorical data. Bivariate logistic regression analysis was used to find associations between the independent variables and the HRQoL. All the variables showing a significant association in a bivariate analysis at $p < 0.25$ were entered into a multivariate logistic regression model to determine risk factors associated with the HRQoL. $p < 0.05$ was considered statistically significant.

Results

Age and clinical characteristics

The number of subjects aged <60 years and ≥60 years is almost evenly matched, and the majority of the subjects were already in menopause (94.2%), had a low level of education (64.9%). A total of 24 (17.5%) subjects had a low Hb concentration (anemia); 82 (55.9%) subjects had a high TC, 119 (86.9%) subjects had a high LDL-C, and 11 (8%) subjects had a low HDL-C. The number of subjects with poor and good HRQoL is almost balanced, both for total HRQoL, HRQoL PCS, and MCS (Table 1).

Table 1: Distribution of subject based on socio-demographic factor, laboratory findings, and HRQoL (n = 137)

Variables	n (%)
Age (years)	
- <60	70 (51.1)
- ≥60	67 (48.9)
Menopause status	
- Premenopause	8 (5.8)
- Postmenopause	129 (94.2)
Education level	
- Low	89 (64.9)
- Moderate/High	48 (35.1)
Anemia status	
- No	113 (82.5)
- Yes	24 (17.5)
Total cholesterol	
- Normal	55 (40.1)
- High	82 (55.9)
LDL cholesterol	
- Normal	18 (13.1)
- High	119 (86.9)
HDL cholesterol	
- Normal	126 (92.0)
- Low	11 (8.0)
HRQoL (total)	
- Poor	69 (50.4)
- Good	68 (49.6)
HRQoL – Physical Component	
- Poor	67 (48.9)
- Good	70 (51.1)
HRQoL – Mental Component	
- Poor	68 (49.6)
- Good	69 (50.4)

HRQoL: Health-related quality of life, HDL: High-density lipoprotein, LDL: Low-density lipoprotein

Risk factors associated with HRQoL

Age and several risk factors associated with HRQoL were explored using bivariate logistic regression analysis and subsequent multivariate logistic regression analyses. In bivariate logistic regression analysis, age, menopausal status, education level, anemia, TC, and LDL cholesterol were found to have non-statistically significant association with HRQoL (Table 2). However, HDL cholesterol was significantly associated with HRQoL (odds ratio = 0.44; 95% Confidence interval [CI] = 0.20–0.97; p = 0.042) (Table 2). Normal HDL cholesterol level was found 0.44 times less likely to be associated with poor HRQoL.

Table 2: Bivariate logistic regression between risk factors and HRQoL in women 50–70 years of age

Risk factor	OR	95% CI	p
Age (years)			
- <60	1.37	0.70–2.76	0.349
- ≥60	1		
Menopausal status			
- Premenopause	0.12	0.01–1.01	0.055
- Postmenopause	1		
Education level			
- Low	0.51	0.25–1.04	0.065
- Moderate/High	1		
Anemia status			
- No	0.65	0.26–1.59	0.350
- Yes	1		
Total cholesterol			
- Normal	0.96	0.48–1.91	0.917
- High	1		
LDL-cholesterol			
- Normal	0.78	0.29–2.19	0.637
- High	1		
HDL-cholesterol			
- Normal	0.44	0.20–0.97	0.042
- Low	1		

HRQoL: Health-related quality of life, OR: Odds ratio, CI: Confidence interval, HDL: High-density lipoprotein, LDL: Low-density lipoprotein.

In multivariate analysis, menopausal status and education level have not a statistically significant association with HRQoL (Table 3). Those normal level of HDL cholesterol was found 0.18 times less likely to be associated with poor HRQoL (Adjusted odds ratio = 0.18; 95% CI = 0.03–0.91; p = 0.038) (Table 3). It can be concluded that HDL-C levels were a risk factors of HRQoL.

Discussion

In the present study, we found that the majority of subjects were in postmenopausal. However, menopausal status was not a risk factor of HRQoL. Moreover, this result was in-line with a population-based survey based on 1140 Greek middle-age women found no effect of menopause on the HRQoL [12]. Another study showed different results compared to our study, women experiencing menopause may have impaired HRQoL [13]. Menopause is also the stage of life that is marked by changes in the role of the menopausal woman, such as the cessation of child rearing and child care, or even the prospect of her being abandoned by the children. Although the biological impact of menopause (such as bone loss) is known with regard to its characteristics, the impact of menopause on the functioning and welfare of women (HRQoL) is still unclear [14], [15]. From the results of observations spanning a period of 5 years, Hess *et al.* [16] found that menopause had a negative impact on several subcomponents of HRQoL, apart from the symptoms of menopause; even the quality of life of the postmenopausal women did not improve after the symptoms of menopause had disappeared.

Table 3: Multiple logistic regression between risk factors and HRQoL in women 50–70 years of age

Risk factor	AOR	95% CI	p
Menopausal status			
• Premenopause	0.15	0.01–1.39	0.091
• Postmenopause	1		
Education level			
• Low	0.53	0.25–1.11	0.094
• Moderate/High	1		
HDL cholesterol			
• Normal	0.18	0.03–0.91	0.038
• Low	1		

AOR: Adjusted odds ratio, CI: confidence interval, HDL: High-density lipoprotein, LDL: Low-density lipoprotein

Subjects with anemia had a lower quality of life (HRQoL) as compared to subjects with normal Hb concentration (without anemia), but not statistically significant. The impact of anemia on HRQoL has been studied in patients from a number of different populations, and most studies found a relationship between HRQoL and anemia in elderly individuals, which on the subscale level was particularly associated with physical health. Various models of multivariate analysis have concluded that anemia is an independent risk factor of decreased HRQoL in the elderly group. In South Korea, data from the Korean National Health and Nutrition Examination Survey, that used the EQ-5D questionnaire, showed that the overall incidence of anemia in individuals aged 10 years and above was around 8.2% in 2005–2015 [17].

In this study, it was found that LDL-C was not a risk factor of HRQoL. Dyslipidemia, especially of LDL-C, has a causal relationship with atherosclerotic cardiovascular disease according to genetic, observational, and clinical outcome studies. A number of study results also showed that dyslipidemia impacts on quality of life (QoL), in that patients with dyslipidemia have a low HRQoL when compared with patients without dyslipidemia [4], [5]. In the multivariate logistic regression on HRQoL involving various risk factors of HRQoL (menopausal status, educational status, and HDL-C), it may be observed that the subjects with low HDL-C had a significantly lower quality of life in comparison with the group of subjects with normal HDL-C concentration. From the results of this study, it is also apparent that a low HDL-C concentration was a risk factor of HRQoL. The previous study results showed a still inconsistent relationship between HDL cholesterol concentration and quality of life. The Chinese Hypertension Intervention Efficacy randomized controlled trial conducted on a group of 13,542 men and women aged 50–79 years showed that decreased LDL-C concentration and increased HDL-C concentration were associated with increased HRQoL [14]. The inconsistent results found in study involving subjects aged 100 years and over showed that HDL-C concentration was not significantly associated with quality of life [8]. Among the oldest elderly (100 years or over), it was found that increased lipid profile was not a risk factor for health and longevity, but in contrast became a protective factor, which differs from traditional reasoning about the impact of blood lipids on health [18], [19].

There is strong evidence for the relationship between high LDL-C concentration and cardiovascular events based on studies of clinical outcomes [20], [21], such that LDL-C constitutes the main target in the management of dyslipidemia. A number of epidemiological studies have also demonstrated that low plasma HDL-C concentration is a risk factor of cardiovascular disease. Similarly, several large-scale randomized clinical trials have proven that administration of HDL-C-increasing medications to patients with low HDL-C and high triglyceride concentrations may yield beneficial clinical effects. However, up to now the causal relationship between HDL-C concentration and cardiovascular disease cannot be explained. HDL-C may predict cardiovascular events even in patients on statin treatment, but clinical studies on the relationship between increased HDL-C concentration and cardiovascular protection are not convincing [8], [22]. HDL-C is at present used for estimating cardiovascular risk and is not a therapeutic target because the clinical studies are not convincing, even in high-risk patients with the LDL-C concentration meeting the therapeutic target. An HDL-C concentration of >40 mg/dL in males and >50 mg/dL in women only indicates a lower level of risk.

Limitation

This observational study with cross-sectional approach cannot prove a causal relationship between HDL-C concentration and HRQoL. Data collected through interview of study participants were subjected to recall bias, particularly in questions related to past events.

Conclusions and Recommendation

This study demonstrated that HDL-C concentrations were significantly a risk factor of HRQoL in women aged 50–70-years. The effect of lipid profile on health status needs to be considered and further verified in women aged 50–70 years. As a follow-up to this study, there is a need for conducting a study with a better design, such as a cohort study, that can determine a cause-and-effect relationship between HDL-C concentration and quality of life in women aged 50–70 years.

Acknowledgments

We are grateful to Head, Staff, and health cadres from the Mampang Prapatan Sub-district

Puskesmas, South Jakarta, for the research area and supporting data facility. We are indebted to all respondents who participated in this study.

References

- Martino G, Catalano A, Bellone F, Langher V, Lasco C, Penna A, *et al.* Quality of life in postmenopausal women: Which role for Vitamin D? *Mediterr J Clin Psychol.* 2018;6:1-14. <https://doi.org/10.6092/2282-1619/2018.6.1875>
- Lin HQ, Wu JY, Chen ML, Chen FQ, Liao YJ, Wu YT, *et al.* Prevalence of dyslipidemia and prediction of 10-year CVD risk among older adults living in southeast coastal regions in China: A cross-sectional study. *Clin Interv Aging.* 2019;14:1119-29. <https://doi.org/10.2147/CIA.S207665>
PMid:31354254
- Gidding SS, Allen NB. Cholesterol and atherosclerotic cardiovascular disease: A lifelong problem. *J Am Heart Assoc.* 2019;8(11):e012924. <https://doi.org/10.1161/JAHA.119.01292>
PMid:31137996
- Park H. The impact of dyslipidemia on the health-related quality of life of Korean females aged 50 years and older. *Iran J Public Health.* 2019;48(3):556-8. <https://doi.org/10.18502/ijph.v48i3.903>
PMid:31223587
- Ko DT, Alter DA, Guo H, Koh M, Lau G, Austin PC, *et al.* High-density lipoprotein cholesterol and cause-specific mortality in individuals without previous cardiovascular conditions: The CANHEART study. *J Am Coll Cardiol.* 2016;68(19):2073-83. <https://doi.org/10.1016/j.jacc.2016.08.038>
PMid:27810046
- Li YH, Tseng WK, Yin WH, Lin FJ, Wu YW, Hsieh IC, *et al.* Prognostic effect of high-density lipoprotein cholesterol level in patients with atherosclerotic cardiovascular disease under statin treatment. *Sci Rep.* 2020;10(1):21835. <https://doi.org/10.1038/s41598-020-78828-8>
PMid:33318590
- Yan R, Gu HQ, Wang W, Ma L, Li W. Health-related quality of life in blood pressure control and blood lipid-lowering therapies: Results from the CHIEF randomized controlled trial. *Hypertens Res.* 2019;42(10):1561-71. <https://doi.org/10.1038/s41440-019-0281-z>
PMid:31197264
- Wang S, Yang S, Jia W, Cao W, Han K, Liu M, *et al.* Relationships of lipids profile with health-related quality of life in Chinese centenarians. *J Nutr Heal Aging.* 2020;24:404-11. <https://doi.org/10.1007/s12603-020-1340-5>
PMid:32242208
- Karimi M, Brazil J. Health, health-related quality of life, and quality of life: What is the difference? *Pharmacoeconomics.* 2016;34(7):645-9. <https://doi.org/10.1007/s40273-016-0389-9>
PMid:26892973
- Lins L, Carvalho FM. SF-36 total score as a single measure of health-related quality of life: scoping review. *SAGE Open Med.* 2016;4:205031211667172. <https://doi.org/10.1177/2050312116671725>
PMid:27757230
- Jörling M, Rutzner S, Hecht M, Fietkau R, Distel LV. Deterioration of health-related quality of life scores under treatment predicts longer survival. *Biomed Res Int.* 2020;2020:3565238.
- Giannouli P, Zervas I, Armeni E, Koundi K, Spyropoulou A, Alexandrou A, *et al.* Determinants of quality of life in Greek middle-age women: A population survey. *Maturitas.* 2012;71(2):154-61. <https://doi.org/10.1016/j.maturitas.2011.11.013>
PMid:22177978
- Poomalar GK, Arounassalame B. The quality of life during and after menopause among rural women. *J Clin Diagnostic Res.* 2013;7(1):135-9. <https://doi.org/10.7860/JCDR/2012/4910.2688>
PMid:23450244
- Whiteley J, Dibonaventura MD, Wagner JS, Alvir J, Shah S. The impact of menopausal symptoms on quality of life, productivity, and economic outcomes. *J Womens Health.* 2013;22:983-90. <https://doi.org/10.1089/jwh.2012.3719>
PMid:24083674
- Molander JM, Aalto AM, Raitanen J, Hemminki E, Aro AR, Luoto R. Physical activity and change in quality of life during menopause -an 8-year follow-up study. *Health Qual Life Outcomes.* 2012;10:1-7. <https://doi.org/10.1186/1477-7525-10-8>
- Hess R, Thurston RC, Hays RD, Chang CC, Dillon SN, Ness RB, *et al.* The impact of menopause on health-related quality of life: Results from the STRIDE longitudinal study. *Qual Life Res.* 2012;21(3):535-44. <https://doi.org/10.1007/s11136-011-9959-7>
PMid:21755412
- Kim YJ, Han KD, Cho KH, Kim YH, Park YG. Anemia and health-related quality of life in South Korea: Data from the Korean national health and nutrition examination survey 2008-2016. *BMC Public Health.* 2019;19(1):735. <https://doi.org/10.1186/s12889-019-6930-y>
PMid:31196013
- Ravnskov U, Diamond DM, Hama R, Hamazaki T, Hammarskjöld B, Hynes N, *et al.* Lack of an association or an inverse association between low-density-lipoprotein cholesterol and mortality in the elderly: A systematic review. *BMJ Open.* 2016;6(6):e010401. <https://doi.org/10.1136/bmjopen-2015-010401>
PMid:27292972
- Takata Y, Ansai T, Soh I, Awano S, Nakamichi I, Akifusa S, *et al.* Serum total cholesterol concentration and 10-year mortality in an 85-year-old population. *Clin Interv Aging.* 2014;9:293-300. <https://doi.org/10.2147/CIA.S53754>
PMid:24611005
- Baigent C, Blackwell L, Emberson J, Holland LE, Reith C, Bhalra N, *et al.* Efficacy and safety of more intensive lowering of LDL cholesterol: A meta-analysis of data from 170 000 participants in 26 randomised trials. *Lancet.* 2010;376(9753):1670-81. [https://doi.org/10.1016/S0140-6736\(10\)61350-5](https://doi.org/10.1016/S0140-6736(10)61350-5)
PMid:21067804
- Penson PE, Pirro M, Banach M. LDL-C: Lower is better for longer even at low risk. *BMC Med.* 2020;18:1-6. <https://doi.org/10.1186/s12916-020-01792-7>
- Voight BF, Peloso GM, Orho-Melander M, Frikke-Schmidt R, Barbalic M, Jensen MK, *et al.* Plasma HDL cholesterol and risk of myocardial infarction: A mendelian randomisation study. *Lancet.* 2012;380(9841):572-80. [https://doi.org/10.1016/S0140-6736\(12\)60312-2](https://doi.org/10.1016/S0140-6736(12)60312-2)
PMid:22607825

High-density Lipoprotein Cholesterol as a Risk Factor of Health-Related Quality of Life in 50–70-Year-Old Community- Dwelling Women

ORIGINALITY REPORT

18%

SIMILARITY INDEX

PRIMARY SOURCES

1	Şenay Topuz, Nuriye BÜYÜKKAYACI DUMAN, Gülzade Kuşoğlu Uysal, Dilek Öcalan. "Breastfeeding self-efficacy and related factors during early postpartum period", <i>Universa Medicina</i> , 2021 <small>Crossref</small>	40 words — 1%
2	www.sciencegate.app <small>Internet</small>	28 words — 1%
3	pure.gustaveroussy.fr <small>Internet</small>	26 words — 1%
4	www.jstage.jst.go.jp <small>Internet</small>	22 words — 1%
5	www.uspharmacist.com <small>Internet</small>	22 words — 1%
6	businessdocbox.com <small>Internet</small>	19 words — 1%
7	Cannon, C.P.. "High-density lipoprotein cholesterol and residual cardiometabolic risk in metabolic syndrome", <i>Clinical Cornerstone</i> , 2007 <small>Crossref</small>	18 words — 1%
8	Jon R. Webb. "Handbook of Spirituality, Health, and Well-Being - A Psychological Perspective", Routledge,	18 words — 1%

-
- 9 ejournal.uika-bogor.ac.id 18 words — 1 %
Internet
-
- 10 João G. Ribeiro de Lima, Gabriela F. Abud, Ellen C. de Freitas, Carlos R. Bueno Júnior. "Effects of the COVID-19 pandemic on the global health of women aged 50 to 70years", *Experimental Gerontology*, 2021 17 words — 1 %
Crossref
-
- 11 Zhi-Hao Li, Yue-Bin Lv, Wen-Fang Zhong, Xiang Gao et al. "High-Density Lipoprotein Cholesterol and All-Cause and Cause-Specific Mortality Among the Elderly", *The Journal of Clinical Endocrinology & Metabolism*, 2019 17 words — 1 %
Crossref
-
- 12 Mark Hamer, Gary O'Donovan, Emmanuel Stamatakis. "High-Density Lipoprotein Cholesterol and Mortality", *Arteriosclerosis, Thrombosis, and Vascular Biology*, 2018 15 words — < 1 %
Crossref
-
- 13 S. Wang, S. Yang, W. Jia, W. Cao, K. Han, Miao Liu, Y. He. "Relationships of Lipids Profile with Health-Related Quality of Life in Chinese Centenarians", *The journal of nutrition, health & aging*, 2020 15 words — < 1 %
Crossref
-
- 14 stacks.cdc.gov 15 words — < 1 %
Internet
-
- 15 www.utupub.fi 15 words — < 1 %
Internet
-
- 16 "Cardiovascular Disease in the Elderly", Springer Science and Business Media LLC, 2023 14 words — < 1 %
Crossref

-
- 17 www.hindawi.com 14 words — < 1 %
Internet
-
- 18 pubmed.ncbi.nlm.nih.gov 13 words — < 1 %
Internet
-
- 19 Anan S. Jarab, Eman A. Alefishat, Walid Al-Qerem, Tareq L. Mukattash, Lina Abu-Zaytoun. "Variables associated with poor health-related quality of life among patients with dyslipidemia in Jordan", Quality of Life Research, 2021 12 words — < 1 %
Crossref
-
- 20 Chunzi Wang, Yunwei Zhang, Shuangcheng Wang, Yi Wang, Yifei Shen, Hansheng Ding. "Identifying risk factors associated with the health-related quality of life for coronary heart diseases elderly using association rule mining", Scientific Reports, 2025 11 words — < 1 %
Crossref
-
- 21 Jingyu Zhao, Zhexiang Kuang, Jing Xu, Xiao Yu, Jin Dong, Juan Li, Liyun Li, Yanjie Liu, Xintong He, Chun Xu, Xia Li, Jun Shi. "Association of lifestyle behaviors with health-related quality of life among patients with hematologic diseases", Quality of Life Research, 2023 11 words — < 1 %
Crossref
-
- 22 Yoshiko, N.. "A long term survey of plasma and tissue ferritin in mice fed on iron deficient diets", Comparative Biochemistry and Physiology -- Part A: Physiology, 1988 11 words — < 1 %
Crossref
-
- 23 Natthawadi Thimthong, Apichat Photi-a, Chanchai Traivaree, Dollapak Apipongrat, Oytip Nathalang. "DIAGNOSIS OF IRON DEFICIENCY ANEMIA IN THAI FEMALE ADOLESCENTS USING RETICULOCYTE HEMOGLOBIN EQUIVALENT", Journal of Southeast Asian Medical Research, 2022 10 words — < 1 %

24 Yoshida, H.. "Effects of simvastatin 20 mg/d on serum lipid profiles in Japanese hyperlipidemic patients: A prospective, open-label pilot study", *Current Therapeutic Research*, 200511/12

Crossref

10 words — < 1 %

25 etd.aau.edu.et

Internet

10 words — < 1 %

26 ouci.dntb.gov.ua

Internet

10 words — < 1 %

27 Aida Giloyan, Tsovinar Harutyunyan, Ani Babayan, Varduhi Petrosyan. "Factors associated with health-related quality of life among people with visual impairments living in nursing homes in Armenia: a cross-sectional study", *Disability and Rehabilitation*, 2023

Crossref

9 words — < 1 %

28 Mitra Darbandi, Farid Najafi, Yahya Pashar, Shahab Rezaeian. "Structural equation model analysis for the evaluation of factors associated with overweight and obesity in menopausal women in RaNCD cohort study", *Menopause*, 2020

Crossref

9 words — < 1 %

29 Qiaochun Chen, Jing Zhou, Na Li, Luming Liu, Yixuan Li, Wenfang Long, Ziyue Luo, Yunru Liu, Sha Xiao. "Factors influencing changes in the quality of life of the Hainan migratory population with hypertension: a survey of the Chengmai mangrove bay community", *BMC Public Health*, 2025

Crossref

9 words — < 1 %

30 Raheb Ghorbani, Mohammad Nassaji, Akram Shahbazi, Bemane Rostami, Maryam Taheri. "Association between quality of life, menopausal status, and sociodemographic factors among middle-aged women in Iran", *Journal of the Egyptian Public Health Association*, 2015

9 words — < 1 %

-
- 31 Seung Hee Kim, Ki Young Son. "Association of Lipoprotein Cholesterol With Future Cardiovascular Disease and Mortality in Older Adults: A Korean Nationwide Longitudinal Study", Research Square Platform LLC, 2020
9 words — < 1%
Crossref Posted Content
-
- 32 Ying-Ying Zheng, Ting-Ting Wu, Xian-Geng Hou, Yi Yang et al. "Low HDL Cholesterol Associated with Reduced Bleeding Risk in Patients Who Underwent PCI: Findings from the PRACTICE Study", Thrombosis and Haemostasis, 2023
9 words — < 1%
Crossref
-
- 33 repository-tnmgrmu.ac.in
Internet
9 words — < 1%
-
- 34 Jemai, H.. "Hypolipidimic and antioxidant activities of oleuropein and its hydrolysis derivative-rich extracts from Chemlali olive leaves", Chemico-Biological Interactions, 20081125
8 words — < 1%
Crossref
-
- 35 Ruohua Yan, Hong-Qiu Gu, Wen Wang, Liyuan Ma, Wei Li. "Health-related quality of life in blood pressure control and blood lipid-lowering therapies: results from the CHIEF randomized controlled trial", Hypertension Research, 2019
8 words — < 1%
Crossref
-
- 36 Shioh-Ru Chang, Cheng Fang Yang, Kuang-Ho Chen. "Relationships between body image, sexual dysfunction, and health-related quality of life among middle-aged women: A cross-sectional study", Maturitas, 2019
8 words — < 1%
Crossref
-
- 37 Ying-Chang Tung, Tsung-Han Tsai, Yu-Jui Hsieh, Tzyy-Jer Hsu, Fu-Chih Hsiao, Chia-Pin Lin, Pao-
8 words — < 1%

Hsien Chu. "Lipid Variability and the Risk of Incident Peripheral Artery Disease", Circulation Journal, 2025

Crossref

38	jurnal.unimus.ac.id Internet	8 words — < 1%
39	new.clinicalsocialwork.eu Internet	8 words — < 1%
40	preprod.aviesan.fr Internet	8 words — < 1%
41	repository.ju.edu.et Internet	8 words — < 1%
42	www.springermedizin.de Internet	8 words — < 1%

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES < 8 WORDS
EXCLUDE MATCHES OFF