

Archives

[Home](#) » Archives

2026

ISSN: 2454-6615

WORLD WIDE JOURNAL OF MULTIDISCIPLINARY RESEARCH AND DEVELOPMENT

VOLUME 12 ISSUE 1

January 2026

Open



World Wide Journal of Multidisciplinary Research and Development

Sr. No.	Title and Authors Name
1	Data Protection of Children and Adolescents in Electronic Health Records in Brazilian Hospitals: Implications of the LGPD and the Statute of the Child and Adolescent Carolina Sturm Trindade Country : Brasil Subject : Health Sciences View Details
2	Internet of Battlefield Things (IoBT): A Systematic Review Dr Anjan K Sinha Country : India Subject : View Details
3	Interethnic marriages and the question of food heritage: the case of Nkam nationals residing in the city of Ngaoundéré (Adamawa-Cameroon) Patrice Wam Mandeng Country : Cameroun Subject : History View Details
4	Community health in rural areas in Ngaoundéré (Adamawa-Cameroon): Gada-Dang case Patrice WAM MANDENG Country : Cameroun Subject : History View Details
5	Ancient healing techniques and methods in Cameroon: a local heritage of majestic rank for different communities. Patrice WAM MANDENG Country : Cameroon Subject : History View Details
6	Establishment of the Railway station in Ngaoundéré (Adamawa-Cameroon) and its side effects: Case study of the spread of sexually transmitted infections (HIV/AIDS) from 1988 to 2004 Patrice Wam Mandeng Country : Cameroon Subject : History View Details

7

Informal Commodity and Derivatives Markets in the CEMAC Region: An Institutional Analysis of the Plantain Banana and Freshwater Fish Value Chains in the Republic of Congo

Claude Ornella SAMBA ZITOU

Country : Democratic Republic of the Congo

Subject : Economic Sciences

[View Details](#)

8

Globalization and the Economic Transformation of Panama

Arthur Kraft

Country : USA

Subject : Business and Economics

[View Details](#)

9

Revolutionizing Entertainment: The AI Generated Tools and its Impact on Mass Media and Entertainment

Dr. Vikranth K

Country : India

Subject : Computer Science

[View Details](#)

10

The System of Graded Intervention for Juvenile Delinquents in China: Challenges and Solutions

Chao Wang

Country : China

Subject : Law

[View Details](#)

11

Between Promised Cooperation and Practiced Unilateralism: The Paradox of Health Policies in Mercosur Countries During the COVID-19 Pandemic

Antonio Neres Norberg

Country : Brazil

Subject : Parasitology

[View Details](#)

12

Severe Preeclampsia with Bell's Palsy in Pregnancy: A Rare Case Report

Diani Nazma

Country : Indonesia

Subject : Anesthesiology

[View Details](#)

Editor-in-Chief

Prof.(Dr.) P. K. Upadhyay

✉ drpku.2010@yahoo.com, dr.prempku@gmail.com, dean.agri@madhavuniversity.edu.in
M.Sc., Ph.D, FISGPB, FIBS, FSRDA, FSSR, FISGBRD & FSBSD, Former Head Genetics and Plant
Breeding RBS College (DrBRA University Agra) Presently working as Professor & Dean College of
Agriculture, Madhav University, Rajasthan, India



				
Dr. K. Leelavathy ✉ priyaleelavathy@gmail.com Assistant professor PG and Research Department of Commerce, Bon Secours College for women, Thanjavur, India	Dr. V. Raghu Raman ✉ drraghuraman@yahoo.com M.Com. PGDBM MBA. Ph.D, Sr.Faculty ,Business Studies Department, Ibra College of Technology, Ibra, Al Sharqiyah North Governorate, Sultanate of Oman	Jibin Shrestha ✉ jibanshrestha@narc.gov.np Scientist, Nepal Agricultural Research Council, National Maize Research Programme, Rampur, Chitwan, Nepal	Dr. Mahdi Zowghi ✉ mahdizoughi@gmail.com Industrial and System Engineering, Management and Soft Computing, Manchester Universial Academy, London, UK	Dr. Wasin Charerntantanakul ✉ wasin@mju.ac.th Associate Professor Program of Biotechnology, Faculty of Science, Maejo University, Thailand 63 M.4 Samsai Chiang Mai, Thailand
				

**WORLD WIDE JOURNAL OF
MULTIDISCIPLINARY RESEARCH AND
DEVELOPMENT**

WWJMRD 2026; 12(01): 79-82

www.wwjmr.com

International Journal

Peer Reviewed Journal

Refereed Journal

Indexed Journal

Impact Factor SJIF 2017:

5.182 2018: 5.51, (ISI) 2020-

2021: 1.361

E-ISSN: 2454-6615

Christian Soesilo

Department of Anesthesiology,
Faculty of Medicine,
Universitas Trisakti, Jakarta,
Indonesia.

Diani Nazma

Department of Anesthesiology,
Faculty of Medicine,
Universitas Trisakti, Jakarta,
Indonesia.

Lira Panduwyat

Department of Anesthesiology,
Faculty of Medicine,
Universitas Trisakti, Jakarta,
Indonesia.

Karlina Mahardewi¹

Department of Anesthesiology,
Faculty of Medicine,
Universitas Trisakti, Jakarta,
Indonesia.

Antin Trilaksni

Department of Anesthesiology,
Faculty of Medicine,
Universitas Trisakti, Jakarta,
Indonesia.

Correspondence:

Diani Nazma

Department of Anesthesiology,
Faculty of Medicine,
Universitas Trisakti, Jakarta,
Indonesia.

Severe Preeclampsia with Bell's Palsy in Pregnancy: A Rare Case Report

Christian Soesilo, Diani Nazma, Lira Panduwyat, Karlina Mahardewi, Antin Trilaksni

Abstract

Introduction: Preeclampsia remains a challenging problem in pregnant mothers. Complications may arise with bell's palsy being one of them. With Bells' palsy, clinicians must be able to exclude acute stroke as a differential diagnosis and face the challenge of deciding the appropriate treatment regimen for the pregnant mother. Therefore, this case report presents a case of a pregnant patient with severe preeclampsia and Bell's palsy.

Case Presentation: A 31-year-old G3P2A0 38-week gestation pregnant female patient presented with acute left-sided facial weakness. The patient had been diagnosed with severe preeclampsia and was receiving oral nifedipine. Physical examination was significant for elevated blood pressure of 161/96 mmHg and left facial asymmetry with forehead involvement indicating peripheral facial nerve palsy. Brain CT-scan showed mild edema at the bilateral subcortical parietooccipital region. Obstetric assessment showed viable term fetus without labor signs. Emergency cesarean section was performed with spinal anesthesia. Both the mother and the neonate had favorable outcome with no perioperative and postoperative complication.

Conclusion: Bell's palsy in patients with severe preeclampsia presents a clinical challenge with multiple management considerations. Early diagnosis, particularly to exclude acute cerebrovascular disease, and timely management are crucial to ensure maternal safety and a favorable neonatal outcome.

Keywords: bell's palsy, preeclampsia, pregnancy.

1. Introduction

Problems during pregnancy is a concerning maternal health problem. Maternal mortality rate (MMR) reaches up to 260,000 women during and after pregnancy with 20% accounted to Southeast Asia. Concerningly, around 92% of all maternal mortality rate occurs in low and lower-middle income countries. Of these maternal mortality rate, around 26% accounted to preeclampsia (PE) in low-income countries.¹ In Indonesia, the maternal mortality rate due to PE reaches up to 173 out of 100,000 livebirths, making Indonesia the fourth highest MMR in Southeast Asia and above the region's MMR of 140 per 100,000 livebirths.² Furthermore, PE contributes as the second most common cause of maternal mortality following postpartum hemorrhage.³

Not only does it cause mortality towards the mother and the fetus, PE may also cause various comorbidities. Among them, Bell's palsy is a rare comorbidity that may cooccur with PE in pregnancy. The incidence of Bell's palsy is higher in pregnant women compared to non-pregnant women with an incidence rate of 40 per 100,000 women in pregnancy. The incidence occurs mostly during the third trimester and during the postpartum period where physiological adaptation, including hypercoagulable state, hormonal changes, relative immunosuppression that may amplify viral inflammation, and edema in preeclampsia predispose to the development of Bell's palsy.⁴

Most patients improve with treatment completely with symptoms appearing for 2-3 days without permanent neurological paralyses. Nevertheless, the treatment of Bell's palsy in pregnancy remains complicated with fewer treatment options compared to non-pregnant women.⁴ Thus, this case report would like to present a rare case of Bell's palsy with

preeclampsia in pregnancy diagnosed during the third trimester and its birth management.

2. Case Presentation

A 31-year-old G3P2A0 female patient at 38 weeks of gestation presented to the emergency department with left-sided facial weakness since 8 hours prior to presentation. The patient's fetal movement was active and denied contraction. Patient had been diagnosed with severe preeclampsia 1 week prior and was treated with oral nifedipine. The patient was alert, vitals were stable except for blood pressure of 161/96 mmHg. On physical examination, the face was asymmetric with drooping of the left side and inability to wrinkle forehead suggesting CN

VII lesion (figure 1). Motoric and sensory examination was within normal limits for both side of the body. The patient's heart and lung examination were also within normal limits. Obstetric examination showed uterine fundal height of 34 cm, fetal heart rate of 148x/minute, no cervical dilatation and a thick rigid cervix. Her supporting examinations showed sinus rhythm on ECG, leukocytosis on complete blood count, and proteinuria in urinalysis. The patient's CT scan result showed slight hypodense at the bilateral subcortical parietooccipital region with a suspicion of posterior reversible encephalopathy syndrome (figure 2). With the confirmation diagnosis of Bell's palsy, the patient was given mecabalamin 500 mg thrice daily and ibuprofen 400 mg twice daily.



Fig. 1: A. Patient's left-sided facial drooping; B. Patient's inability to wrinkle forehead on the left side.




Fig. 2: Patient's CT scan showing slight hypodense at the bilateral subcortical parietooccipital.

The patient was scheduled for an emergency cesarean section. Preoperatively, the patient received nifedipine, magnesium sulfate and was prepared for spinal anesthesia performed in sitting condition under standard aseptic conditions. Spinal anesthesia was performed using a 27-gauge spinal needle at the L2-L3 intervertebral space. The intrathecal anesthetic regimen consisted of hyperbaric bupivacaine 10 mg combined with fentanyl 25 mcg and

morphine 100 mcg, administered over approximately 15 seconds. Monitoring included ECG, heart rate, peripheral oxygen saturation, and respiratory rate. Foley catheter has previously been inserted in the emergency room. Adequate anesthesia was achieved, with sensory blockade up to the T8 dermatome and motor blockade of the lower extremities, evidenced by inability to raise both legs and prick test at level T6 showed no pain.

The cesarean section was performed successfully, and the neonate was delivered with APGAR score of 7 and 8 at 1 and 5 minutes, respectively. Intraoperatively, the fluid was restricted with less than 250 mL of Ringer's lactate and 20 IU of oxytocin drip was given in a 20 mL

syringe instead of regular 500 mL dilution. There were no complications with stable hemodynamics (table 1), and the estimated blood loss was within normal limits for cesarean delivery with no blood transfusion required.

Table 1: Patient's intraoperative hemodynamic stability during cesarean section.

Time (minutes)	Heart rate	SpO2	SBP	DBP	MAP	Respiratory rate
0	106	99	-	-	-	-
5	95	99	177	106	119	-
10	93	99	150	102	118	19
15	88	98	152	84	110	25
20	106	99	144	97	109	22
25	71	99	145	78	93	25

Note: DBP, diastolic blood pressure; MAP, mean arterial pressure; SBP, systolic blood pressure.

Postoperatively, the patient was transferred to the intensive care unit (ICU) for observation. Management included oxygen supplementation via nasal cannula at 2-3 L/min, IV oxytocin 20 IU every 8 hours, 3x1 gram of IV paracetamol, maintenance IV magnesium sulfate, IV furosemide 20 mg, and maintenance fluid of ringer lactate 1000 mL/day. Patient was still continued on mecabalamin and ibuprofen for Bell's palsy. The transversus abdominis plane (TAP) block was also performed for postoperative analgesia. During observation, patient remained hemodynamically stable with no episodes of hypotension, bradycardia, or respiratory depression. Blood pressure and heart rate were maintained within acceptable ranges throughout the procedure. No nausea, vomiting, or neurological deficits was observed in the patient. Neurological evaluation revealed gradual regression of sensory and motor blockade with no abnormal findings. The patient remained alert and oriented. The following day, the patient's symptoms of Bell's palsy subsided. With the stabilization of blood pressure and continued neurological recovery, both maternal and neonatal conditions were satisfactory, with the neonate transferred to routine newborn care.

3. Discussion

Bell's palsy in severe preeclampsia remains a challenging problem in pregnancy. The management remains limited and further complicates childbirth. Patients that experience symptoms and signs of Bell's palsy with preeclampsia needs to be suspected of acute maternal stroke as the first differential diagnosis. Thus, evaluation and physical examination of unilateral weakness or other focal neurological deficits are mandatory. To further exclude or confirm suspicions on acute stroke, brain CT-scan is warranted.^{4,5}

As for the current patient, neurological assessment showed a peripheral facial nerve palsy indicating Bell's palsy. To further confirm the findings and exclude acute stroke, brain CT-scan showed no signs of intracranial hemorrhage or ischemia. Given that stroke is a known complication of severe preeclampsia, it must be carefully considered as a differential diagnosis.⁶ However, the CT-scan only showed subtle hypodense at the bilateral subcortical parietooccipital region which indicated a suspected posterior reversible encephalopathy syndrome due to severe preeclampsia.⁷ In the absence of radiological findings indicative of acute stroke, a diagnosis of Bell's palsy in the setting of severe preeclampsia was reasonably established.

In severe preeclampsia, delivery remains the definitive

treatment. The timing and mode of delivery should be individualized based on gestational age, fetal condition, cervical status, and patient's overall stability. Patients with severe features should undergo expedited delivery to prevent progression to eclampsia, HELLP syndrome, and other neurological complications. Nevertheless, blood pressure control through antihypertensive therapy, in this case oral nifedipine, should still be given to maintain blood pressure, as well as magnesium sulfate for seizure prophylaxis.⁸

Several anesthetic options may be considered for cesarean section, either through general anesthesia or spinal anesthesia. Spinal anesthesia is generally the preferred technique in cesarean section due to minimum fetal drug exposure; however, it still carries a risk for hemodynamic instability and possible change of intracranial pressure (ICP). On the other hand, general anesthesia may be indicated when neuraxial techniques are contraindicated, such as in patients with suspected increased ICP. In our case, the patient was obese which posed potential airway challenges and would have necessitated rapid sequence intubation, with an associated increased risk of perioperative complications. Therefore, after increased ICP was excluded by brain CT-scan, spinal anesthesia was safely performed with appropriate precautions and monitoring.^{9,10}

Postoperatively, the patient was managed in the ICU for continued close observation. Continued magnesium sulfate therapy, strict hemodynamic monitoring, and multimodal analgesia, in our case added TAP block, contributed to a stable postoperative recovery without respiratory depression, neurological deterioration, nor other complications. Improvement of Bell's palsy symptoms occurred concurrently with blood pressure normalization and resolution of pregnancy-related pathophysiological changes. Although the parieto-occipital subcortical edema observed on imaging is unlikely to directly explain the peripheral facial nerve palsy, both findings may reflect systemic endothelial dysfunction and edema associated with severe preeclampsia. Both maternal and neonatal outcome were favorable, underscoring the importance of multidisciplinary management.^{11,12}

Overall, this case highlights the need for early recognition of neurological symptoms in severe preeclampsia, prompt exclusion of obstetric and cerebrovascular emergencies, and individualized anesthesia planning. With appropriate evaluation and monitoring, spinal anesthesia remains a safe and effective option for cesarean delivery in complex

patients with severe preeclampsia, even in the presence of uncommon neurological complications, in this case, Bell's palsy.

4. Conclusion

Patients with severe preeclampsia present significant clinical challenges, which are further compounded by uncommon complications, such as Bell's palsy. Prompt recognition and thorough exclusion of cerebrovascular and obstetric emergencies are essential. Timely, multidisciplinary management is crucial to ensure maternal safety and achieve favorable neonatal outcomes.

5. Acknowledgments

The author would like to thank the Faculty of Medicine, Universitas Trisakti for their support for this case report. The author would also like to thank dr. Otniel Labobar, SpAn and dr. Tjanya Aryasa, SpAn-KAO for their continuous support. We would like to thank dr. Grady for data collection and summary.

References

1. Feng J, Huang J, Wu Q, Wei K, Bin Q. Maternal mortality ratios in Southeast Asia (1990–2021): trends, disparities, and projections through 2050 based on the global burden of disease 2021, with reference to SDG 3.1. *BMC Pregnancy Childbirth.* 2025 Nov 25;25(1):1264.
2. Syairaji M, Nurdjati DS, Wiratama BS, Prüst ZD, Bloemenkamp KWM, Verschueren KJC. Trends and causes of maternal mortality in Indonesia: a systematic review. *BMC Pregnancy Childbirth.* 2024 Jul 30;24(1):515.
3. Pribadi A. Zero Mother Mortality Preeclampsia Program: Opportunity for a Rapid Acceleration in the Decline of Maternal Mortality Rate in Indonesia. *International Journal of Women's Health and Reproduction Sciences.* 2021 May 2;9(3):160–3.
4. Dorjey Y. Bell's palsy with preeclampsia in pregnancy. *Clin Case Rep.* 2022 May;10(5):e05918.
5. Miller EC. Preeclampsia and Cerebrovascular Disease. *Hypertension.* 2019 Jul;74(1):5–13.
6. Bushnell C, Chireau M. Preeclampsia and Stroke: Risks during and after Pregnancy. *Stroke Res Treat.* 2011 Jan 20;2011:858134.
7. Mai H, Liang Z, Chen Z, Liu Z, Xu Y, Chen X, et al. MRI characteristics of brain edema in preeclampsia/eclampsia patients with posterior reversible encephalopathy syndrome. *BMC Pregnancy Childbirth.* 2021 Dec 3;21(1):669.
8. Sharma DD, Chandresh NR, Javed A, Girgis P, Zeeshan M, Fatima SS, et al. The Management of Preeclampsia: A Comprehensive Review of Current Practices and Future Directions. *Cureus.* 2024 Jan 2;
9. Ring L, Landau R, Delgado C. The Current Role of General Anesthesia for Cesarean Delivery. *Curr Anesthesiol Rep.* 2021 Feb 24;11(1):18–27.
10. Ariyo P, Trelles M, Helmand R, Amir Y, Hassani GH, Mftavyanka J, et al. Providing Anesthesia Care in Resource-limited Settings. *Anesthesiology.* 2016 Mar;124(3):561–9.
11. Macones GA, Caughey AB, Wood SL, Wrench IJ, Huang J, Norman M, et al. Guidelines for postoperative care in cesarean delivery: Enhanced Recovery After Surgery (ERAS) Society recommendations (part 3). *Am J Obstet Gynecol.* 2019 Sep;221(3):247.e1-247.e9.
12. Hammer ES, Cipolla MJ. Cerebrovascular Dysfunction in Preeclamptic Pregnancies. *Curr Hypertens Rep.* 2015 Aug 1;17(8):64.

Severe Preeclampsia with Bell's Palsy in Pregnancy: A Rare Case Report

by diani nazma

Submission date: 29-Jan-2026 07:51AM (UTC+0700)

Submission ID: 2735388778

File name: 1178.1.pdf (330.02K)

Word count: 2364

Character count: 13785



WWJMRD 2026; 12(01): 00-00
www.wwjmr.com
International Journal
Peer Reviewed Journal
Referred Journal
Indexed Journal
Impact Factor SJIF 2017: 5.182 2018: 5.51, (ISI) 2020: 2021: 1.361
E-ISSN: 2454-6615

Christian Soesilo
Department of Anesthesiology,
Faculty of Medicine,
Universitas Trisakti, Jakarta,
Indonesia.

Diani Nazma
Department of Anesthesiology,
Faculty of Medicine,
Universitas Trisakti, Jakarta,
Indonesia.

Lira Panduwaty
Department of Anesthesiology,
Faculty of Medicine,
Universitas Trisakti, Jakarta,
Indonesia.

Karlina Mahardewi¹
Department of Anesthesiology,
Faculty of Medicine,
Universitas Trisakti, Jakarta,
Indonesia.

Antin Trilaksmi
Department of Anesthesiology,
Faculty of Medicine,
Universitas Trisakti, Jakarta,
Indonesia.

Correspondence:
Diani Nazma
Department of Anesthesiology,
Faculty of Medicine,
Universitas Trisakti, Jakarta,
Indonesia.

9 Severe Preeclampsia with Bell's Palsy in Pregnancy: A Rare Case Report

Christian Soesilo, Diani Nazma, Lira Panduwaty, Karlina Mahardewi, Antin Trilaksmi

Abstract

Introduction: Preeclampsia remains a challenging problem in pregnant mothers. Complications may arise with bell's palsy being one of them. With Bell's palsy, clinicians must be able to exclude acute stroke as a differential diagnosis and face the challenge of deciding the appropriate treatment regimen for the pregnant mother. Therefore, this case report presents a case of a pregnant patient with severe preeclampsia and Bell's palsy.

Case Presentation: A 31-year-old G3P2A0 38-week gestation pregnant female patient presented with acute left-sided facial weakness. The patient had been diagnosed with severe preeclampsia and was receiving oral nifedipine. Physical examination was significant for elevated blood pressure of 161/96 mmHg and left facial asymmetry with forehead involvement indicating peripheral facial nerve palsy. Brain CT-scan showed mild edema at the bilateral subcortical parietooccipital region. Obstetric assessment showed viable term fetus without labor signs. Emergency cesarean section was performed with spinal anesthesia. Both the mother and the neonate had favorable outcome with no perioperative and postoperative complication.

Conclusion: Bell's palsy in patients with severe preeclampsia presents a clinical challenge with multiple management considerations. Early diagnosis, particularly to exclude acute cerebrovascular disease, and timely management are crucial to ensure maternal safety and a favorable neonatal outcome.

Keywords: bell's palsy, preeclampsia, pregnancy.

1. Introduction

Problems during pregnancy is a concerning maternal health problem. Maternal mortality rate (MMR) reaches up to 260,000 women during and after pregnancy with 20% accounted to Southeast Asia. Concerningly, around 92% of all maternal mortality rate occurs in low and lower-middle income countries. Of these maternal mortality rate, around 26% accounted to preeclampsia (PE) in low-income countries.¹ In Indonesia, the maternal mortality rate due to PE reaches up to 173 out of 100,000 livebirths, making Indonesia the fourth highest MMR in Southeast Asia.² Above the region's MMR of 140 per 100,000 livebirths.² Furthermore, PE contributes as the second most common cause of maternal mortality following postpartum hemorrhage.³

Not only does it cause mortality towards the mother and the fetus, PE may also cause various comorbidities. Among them, Bell's palsy is a rare comorbidity that may cooccur with PE in pregnancy. The incidence of Bell's palsy is higher in pregnant women compared to non-pregnant women with an incidence rate of 40 per 100,000 women in pregnancy. The incidence occurs mostly during the third trimester and during the postpartum period where physiological adaptation, including hypercoagulable state, hormonal changes, relative immunosuppression that may amplify viral inflammation, and edema in preeclampsia predispose to the development of Bell's palsy.⁴

Most patients improve with treatment completely with symptoms appearing for 2-3 days without permanent neurological paralyses. Nevertheless, the treatment of Bell's palsy in pregnancy remains complicated with fewer treatment options compared to non-pregnant women.⁴ Thus, this case report would like to present a rare case of Bell's palsy with

preeclampsia in pregnancy diagnosed during the third trimester and its birth management.

2. Case Presentation

A 31-year-old G3P2A0 female patient at 38 weeks of gestation presented to the emergency department with left-sided facial weakness since 8 hours prior to presentation. The patient's fetal movement was active and denied contraction. Patient had been diagnosed with severe preeclampsia 1 week prior and was treated with oral nifedipine. The patient was alert, vitals were stable except for blood pressure of 161/96 mmHg. On physical examination, the face was asymmetric with drooping of the left side and inability to wrinkle forehead suggesting CN

VII lesion (figure 1). Motoric and sensory examination was within normal limits for both side of the body. The patient's heart and lung examination were also within normal limits. Obstetric examination showed uterine fundal height of 34 cm, fetal heart rate of 148x/minute, no cervical dilatation and a thick rigid cervix. Her supporting examinations showed sinus rhythm on ECG, leukocytosis on complete blood count, and proteinuria in urinalysis. The patient's CT scan result showed slight hypodense at the bilateral subcortical parietooccipital region with a suspicion of posterior reversible encephalopathy syndrome (figure 2). With the confirmation diagnosis of Bell's palsy, the patient was given mecabalamin 500 mg thrice daily and ibuprofen 400 mg twice daily.



Fig. 1: A. Patient's left-sided facial drooping; B. Patient's inability to wrinkle forehead on the left side.




Fig. 2: Patient's CT scan showing slight hypodense at the bilateral subcortical parietooccipital.

The patient was scheduled for an emergency cesarean section. Preoperatively, the patient received nifedipine, magnesium sulfate and was prepared for spinal anesthesia performed in sitting condition under standard aseptic conditions. Spinal anesthesia was performed using a 27-gauge spinal needle at the L2–L3 intervertebral space. The intrathecal anesthetic regimen consisted of hyperbaric bupivacaine 10 mg combined with fentanyl 25 mcg and

morphine 100 mcg, administered over approximately 15 seconds. Monitoring included ECG, heart rate, peripheral oxygen saturation, and respiratory rate. Foley catheter has previously been inserted in the emergency room. Adequate anesthesia was achieved, with sensory blockade up to the T8 dermatome and motor blockade of the lower extremities, evidenced by inability to raise both legs and prick test at level T6 showed no pain.

The cesarean section was performed successfully, and the neonate was delivered with APGAR score of 7 and 8 at 1 and 5 minutes, respectively. Intraoperatively, the fluid was restricted with less than 250 mL of Ringer's lactate and 20 IU of oxytocin drip was given in a 20 mL

syringe instead of regular 500 mL dilution. There were no complications with stable hemodynamics (table 1), and the estimated blood loss was within normal limits for cesarean delivery with no blood transfusion required.

Table 1: Patient's intraoperative hemodynamic stability during cesarean section.

Time (minutes)	Heart rate	SpO2	SBP	DBP	MAP	Respiratory rate
0	106	99	-	-	-	-
5	95	99	177	106	119	-
10	93	99	150	102	118	19
15	88	98	152	84	110	25
20	106	99	144	97	109	22
25	71	99	145	78	93	25

Note: DBP, diastolic blood pressure; MAP, mean arterial pressure; SBP, systolic blood pressure.

Postoperatively, the patient was transferred to the intensive care unit (ICU) for observation. Management included oxygen supplementation via nasal cannula at 2-3 L/min IV oxytocin 20 IU every 8 hours, 3x1 gram of IV paracetamol, maintenance IV magnesium sulfate, IV furosemide 20 mg, and maintenance fluid of ringer lactate 1000 mL/day. Patient was still continued on mecabalamin and ibuprofen for Bell's palsy. The transversus abdominis plane (TAP) block was also performed for postoperative analgesia. During observation, patient remained hemodynamically stable with no episodes of hypotension, bradycardia, or respiratory depression. Blood pressure and heart rate were maintained within acceptable ranges throughout the procedure. No nausea, vomiting, or neurological deficits was observed in the patient. Neurological evaluation revealed gradual regression of sensory and motor blockade with no abnormal findings. The patient remained alert and oriented. The following day, the patient's symptoms of Bell's palsy subsided. With the stabilization of blood pressure and continued neurological recovery, both maternal and neonatal conditions were satisfactory, with the neonate transferred to routine newborn care.

3. Discussion

Bell's palsy in severe preeclampsia remains a challenging problem in pregnancy. The management remains limited and further complicates childbirth. Patients that experience symptoms and signs of Bell's palsy with preeclampsia needs to be suspected of acute maternal stroke as the first differential diagnosis. Thus, evaluation and physical examination of unilateral weakness or other focal neurological deficits are mandatory. To further exclude or confirm suspicions on acute stroke, brain CT-scan is warranted.^{4,5}

As for the current patient, neurological assessment showed a peripheral facial nerve palsy indicating Bell's palsy. To further confirm the findings and exclude acute stroke, brain CT-scan showed no signs of intracranial hemorrhage or ischemia. Given that stroke is a known complication of severe preeclampsia, it must be carefully considered as a differential diagnosis.⁶ However, the CT-scan only showed subtle hypodense at the bilateral subcortical parietooccipital region which indicated a suspected posterior reversible encephalopathy syndrome due to severe preeclampsia.⁷ In the absence of radiological findings indicative of acute stroke, a diagnosis of Bell's palsy in the setting of severe preeclampsia was reasonably established.

In severe preeclampsia, delivery remains the definitive

treatment. The timing and mode of delivery should be individualized based on gestational age, fetal condition, cervical status, and patient's overall stability. Patients with severe features should undergo expedited delivery to prevent progression to eclampsia, HELLP syndrome, and other neurological complications. Nevertheless, blood pressure control through antihypertensive therapy, in this case oral nifedipine, should still be given to maintain blood pressure, as well as magnesium sulfate for seizure prophylaxis.⁸

Several anesthetic options may be considered for cesarean section, either through general anesthesia or spinal anesthesia. Spinal anesthesia is generally the preferred technique in cesarean section due to minimum fetal drug exposure; however, it still carries a risk for hemodynamic instability and possible change of intracranial pressure (ICP). On the other hand, general anesthesia may be indicated when neuraxial techniques are contraindicated, such as in patients with suspected increased ICP. In our case, the patient was obese which posed potential airway challenges and would have necessitated rapid sequence intubation, with an associated increased risk of perioperative complications. Therefore, after increased ICP was excluded by brain CT-scan, spinal anesthesia was safely performed with appropriate precautions and monitoring.^{9,10}

Postoperatively, the patient was managed in the ICU for continued close observation. Continued magnesium sulfate therapy, strict hemodynamic monitoring, and multimodal analgesia, in our case added TAP block, contributed to a stable postoperative recovery without respiratory depression, neurological deterioration, nor other complications. Improvement of Bell's palsy symptoms occurred concurrently with blood pressure normalization and resolution of pregnancy-related pathophysiological changes. Although the parieto-occipital edema observed on imaging is unlikely to directly explain the peripheral facial nerve palsies, both findings may reflect systemic endothelial dysfunction and edema associated with severe preeclampsia. Both maternal and neonatal outcome were favorable, underscoring the importance of multidisciplinary management.^{11,12}

Overall, this case highlights the need for early recognition of neurological symptoms in severe preeclampsia, prompt exclusion of obstetric and cerebrovascular emergencies, and individualized anesthesia planning. With appropriate evaluation and monitoring, spinal anesthesia remains a safe and effective option for cesarean delivery in complex

patients with severe preeclampsia, even in the presence of uncommon neurological complications, in this case, Bell's palsy.

4. Conclusion

Patients with severe preeclampsia present significant clinical challenges, which are further compounded by uncommon complications, such as Bell's palsy. Prompt recognition and thorough exclusion of cerebrovascular and obstetric emergencies are essential. Timely, multidisciplinary management is crucial to ensure maternal safety and achieve favorable neonatal outcomes.

5. Acknowledgments

The author would like to thank the Faculty of Medicine, Universitas Trisakti for their support for this case report. The author would also like to thank dr. Otniel Labobar, SpAn and dr. Tjanya Aryasa, SpAn-KAO for their continuous support. We would like to thank dr. Grady for data collection and summary.

References

- Recovery After Surgery (ERAS) Society recommendations (part 3). *Am J Obstet Gynecol.* 2019 Sep;221(3):247.e1-247.e9.
- Hammer ES, Cipolla MJ. Cerebrovascular Dysfunction in Preeclamptic Pregnancies. *Curr Hypertens Rep.* 2015 Aug 1;17(8):64.
- Feng J, Huang J, Wu Q, Wei K, Bin Q. Maternal mortality ratios in Southeast Asia (1990–2021): trends, disparities, and projections through 2050 based on the global burden of disease 2021, with reference to SDG 3.1. *BMC Pregnancy Childbirth.* 2025 Nov 25;25(1):1264.
- Syairaji M, Nurdiani DS, Wiratama BS, Prüst ZD, Bloemenkamp KWM, Verschueren KJC. Trends and causes of maternal mortality in Indonesia: a systematic review. *BMC Pregnancy Childbirth.* 2024 Jul 30;24(1):515.
- Pribadi A. Zero Mother Mortality Preeclampsia Program: Opportunity for Rapid Acceleration in the Decline of Maternal Mortality Rate in Indonesia. *International Journal of Women's Health and Reproduction Sciences.* 2021 May 2;9(3):160–3.
- Dorjey Y. Bell's palsy with preeclampsia in pregnancy. *Clin Case Rep.* 2022 May;10(5):e05918.
- Miller EC. Preeclampsia and Cerebrovascular Disease. *Hypertension.* 2019 Jul;74(1):5–13.
- Bushnell C, Chireau M. Preeclampsia and Stroke: Risks during and after Pregnancy. *Stroke Res Treat.* 2011 Jan 20;2011:858134.
- Mai H, Liang Z, Chen Z, Liu Z, Xu Y, Chen X, et al. MRI characteristics of brain edema in preeclampsia/eclampsia patients with posterior reversible encephalopathy syndrome. *BMC Pregnancy Childbirth.* 2021 Dec 3;21(1):669.
- Sharma DD, Chandresh NR, Javed A, Gergis P, Zeeshan M, Fatima SS, et al. The Management of Preeclampsia: A Comprehensive Review of Current Practices and Future Directions. *Cureus.* 2024 Jan 2;
- Ring L, Landau R, Delgado C. The Current Role of General Anesthesia for Cesarean Delivery. *Curr Anesthesiol Rep.* 2021 Feb 24;11(1):18–27.
- Ariyo P, Trelles M, Helmand R, Amir Y, Hassani GH, Mftavyanka J, et al. Providing Anesthesia Care in Resource-limited Settings. *Anesthesiology.* 2016 Mar;124(3):561–9.
- Macones GA, Caughey AB, Wood SL, Wrench JJ, Huang J, Norman M, et al. Guidelines for postoperative care in cesarean delivery: Enhanced

Severe Preeclampsia with Bell's Palsy in Pregnancy: A Rare Case Report

ORIGINALITY REPORT

13%
SIMILARITY INDEX

9%
INTERNET SOURCES

8%
PUBLICATIONS

7%
STUDENT PAPERS

PRIMARY SOURCES

1	doaj.org Internet Source	3%
2	wwjmr.com Internet Source	2%
3	Submitted to New England Institute of Technology Student Paper	1%
4	Yeshey Dorjey. "Bell's palsy with preeclampsia in pregnancy", Clinical Case Reports, 2022 Publication	1%
5	"Principles and Practice of Maternal Critical Care", Springer Science and Business Media LLC, 2020 Publication	1%
6	Submitted to University of Sydney Student Paper	1%
7	www.coursehero.com Internet Source	1%
8	Submitted to Heriot-Watt University Student Paper	1%
9	R. M. T. Chathuranga, L. D. W. Gunawardane, R. K. G. N. S. De Silva. "Bell's palsy associated with preeclampsia in late pregnancy", Sri Lanka Journal of Obstetrics and Gynaecology, 2025 Publication	1%
10	www.mdpi.com Internet Source	1%

11 J. Moodley, G. Kalane. "A Review of the Management of Eclampsia: Practical Issues", *Hypertension in Pregnancy*, 2009

Publication

1 %

12 scholar.unair.ac.id

Internet Source

1 %

Exclude quotes Off
Exclude bibliography On

Exclude matches < 10 words