

# Hasil Uji similaritas The Comparison of Aspirin

*by Erica Kholinne*

---

**Submission date:** 08-Feb-2025 10:07PM (UTC+0700)

**Submission ID:** 2582059779

**File name:** 10-1055-s-0044-1785510.pdf (456.59K)


**Word count:** 3672

**Character count:** 20018



# The Comparison of Aspirin and Direct Oral Anticoagulant as Thromboprophylaxis Following Total Knee Replacement: A Retrospective Study

## *Comparação de ácido acetilsalicílico e anticoagulante oral direto como tromboprolaxia após artroplastia total do joelho: Um estudo retrospectivo*

Endrotomo Sumargono<sup>1,2</sup> Maria Anastasia<sup>2</sup> Leonard Christianto Singjie<sup>2</sup> Ifran Saleh<sup>1,3</sup>  
Ira Juliet Anestessia<sup>4</sup> Erica Kholinne<sup>1,4</sup> 

<sup>1</sup> Department of Orthopedic Surgery, St. Carolus Hospital, Jakarta, Indonesia

<sup>2</sup> Faculty of Medicine, Universitas Katolik Indonesia Atma Jaya, Indonesia

<sup>3</sup> Department of Orthopedics and Traumatology, Faculty of Medicine, Universitas Indonesia, Jakarta, Indonesia

<sup>4</sup> Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Address for correspondence: Erica Kholinne, MD, PhD, Department of Orthopedic Surgery, St. Carolus Hospital, Jakarta, Indonesia (e-mail: erica\_kholinne@yahoo.com).

Rev Bras Ortop 2024;59(6):e883–e887.

### Abstract

**Objective** Venous thromboembolism (VTE) is still a major challenge after major orthopaedic surgery, including total knee replacement (TKR). The aim of this study was to estimate the risk of VTE with aspirin-only pharmacologic prophylaxis following primary TKR surgery versus direct oral anticoagulant (DOAC).

**Methods** The study included 476 patients who underwent primary TKR from 2016 to 2020. All patients received thromboprophylaxis with DOAC (DOAC group) ( $n = 267$ ) or aspirin (aspirin group) ( $n = 209$ ). Clinical outcomes were evaluated and compared between those who received DOAC and aspirin. The primary outcome was the incidence of VTE. The secondary outcome was wound complications.

**Result** Aspirin and DOAC were comparable in preventing VTE in patients who underwent primary TKR. The incidence of deep vein thrombosis was similar in the aspirin (10%) and factor Xa inhibitor groups (10.1%), ( $p = 0.98$ ) with zero case of pulmonary emboli in both groups. There was no significant difference between the aspirin (1.4%) and DOAC groups (1.5%) regarding wound complication ( $p = 0.95$ ).

### Keywords

- ▶ anticoagulants
- ▶ arthroplasty, replacement, knee
- ▶ venous thromboembolism

Work carried out the Department of Orthopedic Surgery, St. Carolus Hospital, Jakarta, Indonesia.

received  
March 4, 2023  
accepted  
July 21, 2023

DOI <https://doi.org/10.1055/s-0044-1785510>.  
ISSN 0102-3616.

© 2024. The Author(s).  
This is an open access article published by Thieme under the terms of the Creative Commons Attribution 4.0 International License, permitting copying and reproduction so long as the original work is given appropriate credit (<https://creativecommons.org/licenses/by/4.0/>).  
Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

**Conclusion** Postoperative thromboprophylaxis with aspirin only was not associated with a higher risk of postoperative VTE compared with DOAC following TKR. Considering the wide availability and cost-effectiveness, aspirin may serve as a promising alternative to DOAC for VTE prophylaxis.

## Resumo

**Objetivo** O tromboembolismo venoso (TEV) ainda é um grande desafio após grandes cirurgias ortopédicas, incluindo a artroplastia total do joelho (ATJ). O objetivo deste estudo foi estimar o risco de TEV com profilaxia farmacológica apenas com ácido acetilsalicílico ou anticoagulante oral direto (AOD) após a cirurgia primária de ATJ.

**Métodos** O estudo incluiu 476 pacientes submetidos a ATJ primária de 2016 a 2020. Todos os pacientes receberam trombopprofilaxia com AOD (grupo AOD) ( $n = 267$ ) ou ácido acetilsalicílico (AAS) (grupo AAS) ( $n = 209$ ). Os desfechos clínicos foram avaliados e comparados entre aqueles que receberam AOD e AAS. O desfecho primário foi a incidência de TEV. O desfecho secundário foi a ocorrência de complicações da ferida.

**Resultado** O AAS e o AOD foram comparáveis na prevenção de TEV em pacientes submetidos a ATJ primária. A incidência de trombose venosa profunda foi semelhante nos grupos AAS (10%) e AOD (10,1%,  $p = 0,98$ ), sem nenhum caso de embolia pulmonar em ambos os grupos. Não houve diferença significativa entre os grupos AAS (1,4%) e AOD (1,5%) em relação às complicações da ferida ( $p = 0,95$ ).

**Conclusão** A trombopprofilaxia pós-operatória apenas com AAS não foi associada a um maior risco de TEV pós-operatório em comparação à AOD após ATJ. Considerando a ampla disponibilidade e custo-benefício, o AAS pode ser uma alternativa promissora ao AOD para a profilaxia de TEV.

## Palavras-chave

- ▶ anticoagulantes
- ▶ artroplastia do joelho
- ▶ tromboembolia venosa

## Introduction

Total knee replacement (TKR) has been one of the most successful orthopaedic interventions which have an impact on improving the quality of life. Projection studies suggest that the demand for primary TKR will steadily increase in the years to come.<sup>1</sup> However, the TKR procedure also has some postoperative complications. One of the most common yet serious complications is venous thromboembolic disease (VTE). Venous thromboembolism is a known complication after major orthopaedic surgery, including knee or hip replacement.<sup>2,3</sup>

Several studies have examined the efficacy of several antithrombotic drugs over their years as well as their side effects profiles, including prolonged wound leakage, bleeding, and infection.<sup>4,5</sup> The following agents that are usually used for thromboprophylaxis are antiplatelet (aspirin) or direct oral anticoagulant (DOAC), such as factor Xa inhibitor (rivaroxaban or edoxaban). However, it is currently recognized that aspirin is preferable over other anticoagulants because of its low risk of inducing major bleeding.<sup>6,7</sup> Aspirin may serve as an alternative thromboprophylaxis agent following primary knee or hip replacement, particularly in patients at high risk of bleeding. Other advantages of aspirin over DOACs include its ease of administration, lack of monitoring needs, and low cost.<sup>8</sup>

Indonesia is one of the lower-middle income countries according to the World Bank.<sup>9</sup> Therefore, there is a need to find the most effective and safe thromboprophylaxis with the most reasonable cost. The current study aims to assess


the clinical effectiveness and safety of aspirin compared with DOAC for VTE prophylaxis after TKR in a single institution. We hypothesized that there was no significant difference between aspirin and DOAC in preventing VTE after TKR procedure.

## Methods

### Data Source and Study Cohort

This Institutional Review Board (IRB) of our institution approved the present study under the number (Number 940/RSSC-5B/AL/IX/DIRUT/2022). Retrospective data of 476 primary TKRs was extracted from the 5 years (2016–2020) of annual reports of TKR registry data in our institution. The inclusion criteria were as follows: (1) patients who underwent primary TKR surgery; and (2) patients who received aspirin or DOAC for VTE prophylaxis. The exclusion criteria were as follows: (1) Prior hemorrhagic disease; (2) history of hypercoagulation state; (3) history of malignancy; (4) history of venous insufficiency; (5) bilateral TKR; (6) non-metal back tibial component. The sample selection of the study is shown in ▶ Fig. 1.

The primary outcome was postoperative symptomatic venous thromboembolism (VTE) occurring during the index hospitalization, or responsible for readmission within 30 days, or an outpatient visit within 90 days after discharge, because an increased thromboembolic risk persists up to 3 months postoperatively. The secondary outcome was wound complication. All patients underwent the same



**Fig. 1** Flow chart of study sample selection.

surgical technique (medial parapatellar approach and application of a tourniquet 150 mmHg above diastolic blood pressure) and perioperative protocol (perioperative pain medication, 2 g of intravenous cefazoline as prophylactic antibiotics. The mechanical prophylaxis was periodic ankle pumping, started 6 hours postoperatively, along with the application of an intermittent pneumatic compression device (IPCD). All patients received chemical thromboprophylaxis, which were aspirin 160 mg aspirin group) or factor Xa inhibitors (DOAC group), which was either rivaroxaban 10 mg (Xarelto; Bayer Pharmaceutical, Leverkusen, Germany) or edoxaban 30 mg (Lixiana; Kalbe Farma, Jakarta, Indonesia) 12 hours following surgery and continued for 14 days. All of the implants used in this study were PFC Sigma or Attune (DePuy Synthes, Raynham, MA, USA). Investigations for deep vein thrombosis (DVT) were initiated on symptomatic patients in the routine follow-up examination. The diagnosis of VTE was made by confirming thrombosis of the deep vein involving the popliteal vein or more proximal leg veins (including the femoral, common femoral, and iliac veins) with a sonogram by a radiologist. Wound complication was defined as patients requiring irrigation and debridement, with or without component exchange. All data were

collected from medical records by two medical doctors posted in the orthopedic service who were not involved in the surgery.

### Statistical Analysis

All of the experimental data were statistically analyzed using SPSS Statistics for Windows version 26.0 software (IBM Corp, Armonk, NY, USA). Bivariate analysis was used to test the association between individual thromboprophylaxis and the major clinical outcomes of postoperative VTE using the Chi-squared test. Statistical significance was set at  $p < 0.05$ .

### Result

The demographic data in **Table 1** showed that there were no significant demographic differences between the two groups. The gender majority in both groups was female (80% versus 81%). More than 2/3 of the patients in both groups had comorbidities. Both groups showed similar comorbidities profiles, with hypertension being the most common comorbidity (45,4% and 51,3% in the aspirin and the DOAC groups, respectively). The length of stay in the aspirin and DOAC groups was equivalent (4.4 versus 4.6 days).

There were 21 DVT cases (10%) in the aspirin group compared with 27 cases (10.1%) in those in DOAC group ( $p = 0.98$ ). For wound complications, there were 3 cases in the aspirin group (1.4%) and 4 cases in the DOAC group (1.5%); ( $p = 0.95$ ). Only 2 cases required component exchanges because of prosthetic joint infection, 1 from each group. The rest were treated with irrigation and debridement. **Fig. 2** showed that there was a similar number of DVT incidence and wound complications after TKR between the aspirin and DOAC groups.


### Discussion

From our study, we found that aspirin and the DOAC (rivaroxaban or edoxaban) were comparable in preventing VTE in

**Table 1** Patients' d data

Characteristic	Aspirin group N = 209 (45%)	DOAC group N = 267 (55%)	P-value
Mean age (years)	68.3	67.9	0.32
Gender: male/female (%)	43/166 (20/80)	53/214 (19/81)	0.06
Comorbidities			0.94
Hypertension (%)	95 (45,4)	137 (51,3)	
Diabetes mellitus type 2 (%)	32 (15,3)	39 (14,6)	
Cardiovascular disease (%)	29 (13,8)	28 (10,4)	
Kidney disease (%)	7 (3,3)	2 (0,7)	
Cerebrovascular disease (%)	3 (1,4)	3 (1,1)	
Liver disease (%)	3 (1,4)	3 (1,1)	
Mean length of stay (SD)	4.4 (2.7)	4.6 (2.1)	0.09

Abbreviation: DOAC, direct oral anticoagulant.



**Fig. 2** Incidence of deep vein thrombosis (DVT) & wound complication in the Aspirin and direct oral anticoagulant (DOAC) groups.

patients who underwent TKR. The incidence of DVT was similar in the aspirin and factor Xa inhibitor groups (10% versus 10.1%, respectively), with no occurrence of pulmonary emboli (PE) in either group. There was no association between the type of VTE prophylaxis agent with the risk of wound complication in post-TKR patients because the incidence in the aspirin and factor Xa inhibitor groups (1.4% versus 1.5%, respectively) was also similar.

The exact incidence of DVT and pulmonary emboli remains controversial.<sup>10</sup> Without any prophylaxis protocol, incidence rates greater than 50% following joint replacement have been reported.<sup>11</sup> The incidence has dramatically decreased nowadays, with early mobilization, mechanical compression devices, and chemical prophylaxis.<sup>11</sup> There has been substantial debate regarding the preferred chemical prophylactic agent among aspirin, enoxaparin, warfarin, and DOAC when weighing VTE prophylaxis efficacy versus bleeding risk.<sup>12</sup> Direct oral anticoagulants have a better VTE prophylaxis compared with other anticoagulants, with a lower rate of complications, such as the need for blood transfusion and prolonged wound drainage.<sup>13</sup> Aspirin is commonly used for primary or secondary prevention of cardiovascular diseases, as it inhibits the aggregation of platelets by the irreversible acetylation of cyclooxygenase, thus interfering with the ability of platelet to produce prostaglandin. Aspirin also provided a comparable VTE prophylaxis to that of DOACs.<sup>12,14</sup> Recently, some studies recommended the use of aspirin as another option for VTE prophylaxis, in addition to DOAC or low-molecular-weight heparin (LMWH) for elective joint replacement surgery, including TKR, combined with mechanical prophylaxis until hospital discharge.<sup>15</sup>

The current study showed that the VTE incidence was not influenced by the type of thromboprophylaxis used is supported by recent literatures. Several studies also concluded that the use of aspirin following TKR was associated with low rates of thromboembolic events, similar to the rates of those treated with DOAC.<sup>14,15</sup> Aspirin has been found to have a low VTE rate, with no difference when compared with enoxaparin following TKR procedure.<sup>16</sup> Venker et al.<sup>17</sup> showed that DOAC was superior to enoxaparin in preventing DVT. Other randomized clinical trials by Le et al.<sup>18</sup> and Xu et al.<sup>19</sup> reported that there was no difference between DOACs (rivar-

oxaban) and aspirin when used as VTE prophylaxis following TKR, which supported the finding of the current study. Administration of VTE prophylaxis also comes with several drawbacks, such as postoperative bleeding and prolonged wound drainage.<sup>19</sup> Postoperative bleeding is one of the major concerns of VTE prophylaxis. The more potent the pharmacologic agent, the higher the risk it poses.<sup>20</sup> Aspirin has been found to have a lower bleeding risk compared with other VTE prophylaxis agents, as reported by Wilson et al.<sup>21</sup> Similarly, DOAC also has a low bleeding risk compared with enoxaparin and warfarin, especially if given in a low dose.<sup>13</sup> Another disadvantage of VTE prophylaxis is postoperative wound complication. A study conducted by Lassen et al.<sup>22</sup> reported that administration of rivaroxaban increased wound complications following TKR, although not as severe as those found in patients using low molecular weight Heparin (LMWH), as reported by Kulsrestha et al.<sup>23</sup> Singh et al.<sup>24</sup> have demonstrated that the use of aspirin lowers the rate of prolonged wound drainage in post-TKR patients, hence, lower the risk of developing any possible wound complication, including superficial and deep wound infection. Comparably, another study by Garfinkel et al.<sup>25</sup> reported that patients who received DOAC, such as apixaban, for VTE prophylaxis following joint replacement had a significantly higher rate of wound complications and bleeding. Interestingly, the current study found that there was no statistical difference between aspirin and DOAC in terms of wound complications.

Aspirin is a generic, inexpensive, and widely-available anti-platelet drug. Hence, the use of aspirin as a VTE prophylaxis can be such a cost-effective choice.<sup>26</sup> As previously stated, Indonesia still comes under the lower-middle income countries, as reported by the World Health Organization. According to the e-catalogue from the Ministry of Health, the cost of aspirin is Rp 116,-/tablet (0.79€); meanwhile, the price of rivaroxaban is Rp 23.500,-/tablet (\$1.39).<sup>27</sup> Therefore, given the comparable effect to DOAC as a thromboprophylaxis agent, aspirin remains the most cost-effective thromboprophylaxis agent, especially when compared with factor Xa inhibitor.

There are a few limitations to this study. First, concerning the nature of a retrospective study, the strength of evidence is still limited. The findings still need to be confirmed with future prospective trials with a higher level of evidence. Second, there are inherent selections and observer biases as patients were from a single hospital. Despite these limitations, our findings are clinically important. The samples in our study are large; therefore, they can be adequate to show aspirin is not inferior compared with rivaroxaban. Future multi-center trials are required to reduce the biases.

## Conclusion

Patients undergoing major orthopedic surgery, such as TKR, are at the highest risk for developing VTE during and after hospitalization. The current study showed that there was no significant difference between aspirin and DOAC in preventing VTE following TKR. When choosing the VTE prophylaxis

in patients undergoing TKR surgery, given the safety, wide availability, and cost-effectiveness, aspirin may be considered as a promising alternative to DOAC for the VTE prophylaxis.

#### Author Contributions

All authors contributed to the study conception and design. Material preparation, data collection and analysis were performed by Maria Anastasia and Leonard Christianto. All authors read and approved the final manuscript.

#### Financial Support

The authors declare that this research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

#### Conflict of Interests

The authors declare that there is no conflict of interest regarding the publication of this paper.

#### References

- Skou ST, Roos EM, Laursen MB, et al. A randomized, controlled trial of total knee replacement. *N Engl J Med* 2015;373(17):1597–1606
- Sloan M, Premkumar A, Sheth NP. Projected volume of primary total joint arthroplasty in the U.S., 2014 to 2030. *J Bone Joint Surg Am* 2018;100(17):1455–1460
- Healy WL, Della Valle CJ, Iorio R, et al. Complications of total knee arthroplasty: standardized list and definitions of the Knee Society. *Clin Orthop Relat Res* 2013;471(01):215–220
- Lieberman JR, Pensak MJ. Prevention of venous thromboembolic disease after total hip and knee arthroplasty. *J Bone Joint Surg Am* 2013;95(19):1801–1811
- Sasaki H, Ishida K, Shibamura N, et al. Retrospective comparison of three thromboprophylaxis agents, edoxaban, fondaparinux, and enoxaparin, for preventing venous thromboembolism in total knee arthroplasty. *Int Orthop* 2014;38(03):525–529
- Lewis CG, Inneh IA, Schutzer SF, Grady-Benson J. Evaluation of the first-generation AAOS clinical guidelines on the prophylaxis of venous thromboembolic events in patients undergoing total joint arthroplasty: experience with 3289 patients from a single institution. *J Bone Joint Surg Am* 2014;96(16):1327–1332
- Matharu GS, Kunutsor SK, Judge A, Blom AW, Whitehouse MR. Clinical Effectiveness and safety of aspirin for venous thromboembolism prophylaxis after total hip and knee replacement: A systematic review and meta-analysis of randomized clinical trials. *JAMA Intern Med* 2020;180(03):376–384
- Raphael IJ, Tischler EH, Huang R, Rothman RH, Hozack WJ, Parvizi J. Aspirin: an alternative for pulmonary embolism prophylaxis after arthroplasty? *Clin Orthop Relat Res* 2014;472(02):482–488
- Worldbank. Indonesia | Data. (n.d.). Available from: <https://data.worldbank.org/country/ID>. [Accessed October 17, 2022]
- The ICM-VTE Hip & Knee Delegates. Recommendations from the ICM-VTE: Hip & Knee. [published correction appears in *J Bone Joint Surg Am*. 2022 Aug 3;104(15):e70] *J Bone Joint Surg Am* 2022;104(Suppl 1):180–231
- Etzioni DA, Lessow C, Bordeianou LG, et al. Venous Thromboembolism after Inpatient Surgery in Administrative Data vs NSQIP: A Multi-Institutional Study. *J Am Coll Surg* 2018;226(05):796–803
- Baumgartner C, Maselli J, Auerbach AD, Fang MC. Aspirin compared with anticoagulation to prevent venous thromboembolism after knee or hip arthroplasty: a large retrospective cohort study. *J Gen Intern Med* 2019;34(10):2038–2046
- Sun G, Wu J, Wang Q, et al. Factor Xa inhibitors and direct thrombin inhibitors versus low-molecular-weight heparin for thromboprophylaxis after total hip or total knee arthroplasty: A systematic review and meta-analysis. *J Arthroplasty* 2019;34(04):789–800.e6
- Deirmengian GK, Heller S, Smith EB, Maltenfort M, Chen AF, Parvizi J. Aspirin can be used as prophylaxis for prevention of venous thromboembolism after revision hip and knee arthroplasty. *J Arthroplasty* 2016;31(10):2237–2240
- Anderson DR, Dunbar M, Murnaghan J, et al. Aspirin or rivaroxaban for VTE prophylaxis after hip or knee arthroplasty. *N Engl J Med* 2018;378(08):699–707
- Farey JE, An VVG, Sidhu V, Karunaratne S, Harris IA. Aspirin versus enoxaparin for the initial prevention of venous thromboembolism following elective arthroplasty of the hip or knee: A systematic review and meta-analysis. *Orthop Traumatol Surg Res* 2021;107(01):102606
- Venker BT, Ganti BR, Lin H, Lee ED, Nunley RM, Gage BF. Safety and efficacy of new anticoagulants for the prevention of venous thromboembolism after hip and knee arthroplasty: A meta-analysis. *J Arthroplasty* 2017;32(02):645–652
- Le G, Yang C, Zhang M, et al. Efficacy and safety of aspirin and rivaroxaban for venous thromboembolism prophylaxis after total hip or knee arthroplasty: A protocol for meta-analysis. *Medicine (Baltimore)* 2020;99(49):e23055
- Xu J, Kanagaratnam A, Cao JY, Chaggar GS, Bruce W. A comparison of aspirin against rivaroxaban for venous thromboembolism prophylaxis after hip or knee arthroplasty: A meta-analysis. *J Orthop Surg (Hong Kong)* 2020;28(01):2309499019896024
- Jenny JY, Bulaid Y, Boisrenoult P, et al; French Society of Orthopaedic Surgery, Traumatology (SofCOT) Bleeding and thromboembolism risk of standard antithrombotic prophylaxis after hip or knee replacement within an enhanced recovery program. *Orthop Traumatol Surg Res* 2020;106(08):1533–1538
- Wilson DG, Poole WE, Chauhan SK, Rogers BA. Systematic review of aspirin for thromboprophylaxis in modern elective total hip and knee arthroplasty. *Bone Joint J* 2016;98-B(08):1056–1061
- Lassen MR, Gent M, Kakkar AK, et al. The effects of rivaroxaban on the complications of surgery after total hip or knee replacement: results from the RECORD programme. *J Bone Joint Surg Br* 2012;94(11):1573–1578
- Kulshrestha V, Kumar S. DVT prophylaxis after TKA: routine anticoagulation vs risk screening approach - a randomized study. *J Arthroplasty* 2013;28(10):1868–1873
- Singh V, Shahi A, Saleh U, Tarabichi S, Oliashirazi A. Persistent wound drainage among total joint arthroplasty patients receiving aspirin vs coumadin. *J Arthroplasty* 2020;35(12):3743–3746
- Garfinkel JH, Gladnick BP, Roland N, Romness DW. Increased incidence of bleeding and wound complications with Factor-Xa inhibitors after total joint arthroplasty. *J Arthroplasty* 2018;33(02):533–536
- Parvizi J, Huang R, Restrepo C, et al. Low-dose aspirin is effective chemoprophylaxis against clinically important venous thromboembolism following total joint arthroplasty: A preliminary analysis. *J Bone Joint Surg Am* 2017;99(02):91–98
- Peraturan Menteri Kesehatan Republik Indonesia. (n.d.). Available from: [www.inaproc.lkpp.go.id](http://www.inaproc.lkpp.go.id); [accessed October 17, 2022]

# Hasil Uji similaritas The Comparison of Aspirin

---

## ORIGINALITY REPORT

---

18%

SIMILARITY INDEX

11%

INTERNET SOURCES

14%

PUBLICATIONS

8%

STUDENT PAPERS

---

## MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

---

1%

★ Daniel C. Santana, Ahmed K. Emara, Melissa N. Orr, Alison K. Klika et al. "An Update on Venous Thromboembolism Rates and Prophylaxis in Hip and Knee Arthroplasty in 2020", Medicina, 2020

Publication

---

Exclude quotes  On

Exclude matches  < 10 words

Exclude bibliography  On