

HOME / ARCHIVES / Vol 13 No 7 (2023): Volume 13, Issue 7, July 2023

Published: 2023-07-15

RESEARCH

PREVENTIVE EFFECT OF TOTAL ETHANOLIC EXTRACT OF BUTEA MONOSPERMA ON BONE LOSS IN OSTEOPENIC RATS

Sonu Khanka, Amardeep Lakra, Ritika Gupta, Kriti Sharma, Rakesh Maurya, Naibedy Chattopadhyay, Divya Singh Pages 6-14

[Abstract](#) [Download PDF](#) [View PDF](#) DOI [10.22270/jddt.v13i7.5887](https://doi.org/10.22270/jddt.v13i7.5887) [View HTML](#)

FORMULATION AND EVALUATION OF BUCCOADHESIVE TABLET OF VERAPAMIL HYDROCHLORIDE FOR THE TREATMENT OF HYPERTENSION

Sohan Sukaldas Thipe, Amol V Sawale, Shraddha Dilip Muneshwar, Shiwani A Gulhane, Sakshi S Gahukar Pages 15-21

[Abstract](#) [Download PDF](#) [View PDF](#) DOI [10.22270/jddt.v13i7.6122](https://doi.org/10.22270/jddt.v13i7.6122) [View HTML](#)

PHENOLIC PROFILE AND ANTIOXIDANT POTENTIAL OF AQUEOUS EXTRACTS OF SELECTED TRADITIONAL ANTI-COUGH PLANTS

Olachi Lilian Osuagwu, Chidi Uzoma Igwe, Chiamaka Perpetua Nzebude, Onyinyechi Chijioko Njoku, Juliet Chioma Ejiolor Pages 22-29

[Abstract](#) [Download PDF](#) [View PDF](#) DOI [10.22270/jddt.v13i7.6124](https://doi.org/10.22270/jddt.v13i7.6124) [View HTML](#)

EMERGING PROBLEM EVALUATION OF HYPOCALCEMIA IN SUPER VASMOL HAIR DYE POISONING PATIENTS: CLINICAL STUDY IN RIMS, KADAPA

K. Leela Prasad Babu, Shaik Kareemulla, M. Ramani, L. Nagamani, B. Abhishek, K. Sravanthi Bai Pages 30-33

[Abstract](#) [Download PDF](#) [View PDF](#) DOI [10.22270/jddt.v13i7.6116](https://doi.org/10.22270/jddt.v13i7.6116) [View HTML](#)

CAN MORINGA OLEIFERA LEAF ETHYL ACETATE EXTRACT INHIBIT CANDIDA ALBICANS PLANKTONIC CELL GROWTH AND BIOFILM FORMATION IN VITRO?

Herastuti Sulistyani, Siti Sulastri, Dewi Risnawati, Dewi Agustina Pages 34-37

[Abstract](#) [Download PDF](#) [View PDF](#) DOI [10.22270/jddt.v13i7.6146](https://doi.org/10.22270/jddt.v13i7.6146) [View HTML](#)

EXPLORING DRUG PRESCRIPTION PATTERNS IN THE NEUROLOGY DEPARTMENT OF A TERTIARY CARE HOSPITAL: A COMPREHENSIVE ANALYSIS

R Nithya, Solanky K Neesha, UDHAYAKUMAR Narmadha, R. R. Redlin, A Ramya, B Manisha Pages 38-45

[Abstract](#) [Download PDF](#) [View PDF](#) DOI [10.22270/jddt.v13i7.6139](https://doi.org/10.22270/jddt.v13i7.6139) [View HTML](#)

FORMULATION AND EVALUATION OF POLYHERBAL TABLET BY USING NEEM, TULSI, TURMERIC AND GINGER EXTRACT

Souvik Paul, Tanmoy Dey, Prabin Koirala, Sidhant Tamang, Soumik Bhattacharya, Rajat Das Pages 46-51

[Abstract](#) [Download PDF](#) [View PDF](#) DOI [10.22270/jddt.v13i7.5895](https://doi.org/10.22270/jddt.v13i7.5895) [View HTML](#)

FACTORS ASSOCIATED WITH PLASMA MALONDIALDEHYDE LEVELS IN PEOPLE OVER 40 YEARS

Meiyanti Meiyanti, Yohana, Eveline Margo, Juni Chudri, Pusparini, Meutia Atika Faradilla Pages 52-56

[Abstract](#) [Download PDF](#) [View PDF](#) DOI [10.22270/jddt.v13i7.6142](https://doi.org/10.22270/jddt.v13i7.6142) [View HTML](#)

ANALYSIS OF THE FACTORS THAT INFLUENCE THE FAMILY IN THE USE OF TRADITIONAL MEDICINE AS SELF MEDICATION IN ILIR TIMUR II DISTRICT

Hairun Niza, Sarah Hanifah Az-Zahra Anwar, Wahyudi Pages 57-61

[Abstract](#) [Download PDF](#) [View PDF](#) DOI [10.22270/jddt.v13i7.6149](https://doi.org/10.22270/jddt.v13i7.6149) [View HTML](#)

EVALUATION OF THE POTENTIAL FOR MEDICATION ERROR IN OUTPATIENT ELDERLY PATIENTS AT SITI FATIMAH AZ-ZAHRA REGIONAL HOSPITAL

Hairun Niza, Sarah Hanifah Az-Zahra Anwar, Sindi Utami Pages 62-64

[Abstract](#) [Download PDF](#) [View PDF](#) DOI [10.22270/jddt.v13i7.6153](https://doi.org/10.22270/jddt.v13i7.6153) [View HTML](#)

STRUCTURE-BASED DRUG DESIGN, ADME AND MOLECULAR DOCKING ANALYSES OF ANTI-VIRAL AGENTS AGAINST SARS-COV-2 VIRUS, ZIKA VIRUS AND HEPATITIS C VIRUS

Peter Solo, Sangdintuile Zeliang, Muluvelu Lohe, Avünü Neikha, Akumsunep Jamir Pages 65-74

[Abstract](#) [Download PDF](#) [View PDF](#) DOI [10.22270/jddt.v13i7.5909](https://doi.org/10.22270/jddt.v13i7.5909) [View HTML](#)

POTENTIAL ACTIVITY OF STEVIA REBAUDIANA BERT. IN INHIBITING CYCLOOXYGENASE AND LIPOXYGENASE ENZYMES AS ANTI-INFLAMMATORY CANDIDATES: A MOLECULAR DOCKING STUDY AND ADMET PREDICTION

Meida Adelia, Arif Setiawansyah, Rina SE Sitindaon Pages 75-86



Online ISSN: 2250-1177

MAKE A SUBMISSION

## Editorial Board

### EDITOR-IN-CHIEF

**Dr. Ram C Dhakar**, Hospital Pharmacy, SRG Hospital and Medical College Jhalawar, India-326001, India. Email: [editor.jddt@gmail.com](mailto:editor.jddt@gmail.com) Phone: +91-9783920207 [Google Scholar](#) [ResearchGate](#) [ORCID](#) [SCOPUS](#) Research fields: Novel Drug delivery, Nanotechnology, Microspheres, TDDS

### ASSOCIATE EDITORS

**Dr. Aamir Jalal Al-Mosawi**, Advisor And Expert Trainer, Baghdad Medical City And The National Training And Development Center, Iraqi Ministry Of Health, Baghdad, Iraq. [ResearchGate](#) [Google Scholar](#) [SCOPUS](#) [ORCID](#) [NCBI-NLM](#) Email: [almosawiaj@yahoo.com](mailto:almosawiaj@yahoo.com). Research Fields: Pediatrics, pediatric psychiatry, nephrology, pediatric neurology, genetics

**Dr. S K Prajapati**, Professor & Head, Institute of Pharmacy, Bundelkhand University, Jhansi (U.P.), India, India. Email: [drsuniilprajapati@gmail.com](mailto:drsuniilprajapati@gmail.com) [ResearchGate](#) [Website](#) [Google Scholar](#) Research Area: Novel Drug Delivery, Microspheres, Nanoparticles

**Dr. Rajesh Ramasamy**, Associate Professor, Head of Department, Department of Pathology, Faculty Medicine and Health Science, Universiti Putra Malaysia, Malaysia Phone: +603 8947 2377 Mobile: +6012 393 6444 Fax: +603 8941 2787 Email: [rajesh@upm.edu.my](mailto:rajesh@upm.edu.my) [Google Scholar](#) [ResearchGate](#) [Website](#) SCOPUS ID: 26024442700 Research Fields: Stem Cells, Immunology, Diabetes, Medicinal Plant Research, Sidha Medicine

**Dr. Syed Mohd Danish Rizvi**, Associate Professor, Department of Pharmaceutics, College of Pharmacy, University of Hail, Hail, Saudi Arabia. Email: syedrizvi10@yahoo.com Official Email: [sm.danish@uoh.edu.sa](mailto:sm.danish@uoh.edu.sa) | [ORCID ID](#) | [Google Scholar](#) | [Scopus](#) | [Research Gate](#) | [Website](#) | Research fields: Microspheres, Nanoparticles, Cancer

Therapy

### SECTION EDITORS

**Dr. Priyanka Bhatt**, Department of Pharmaceutical Sciences, College of Pharmacy, University of South Florida, 12901 Bruce B. Downs Blvd., MDC30, Tampa, FL 33612, Email: [priyankabhatter88@yahoo.com](mailto:priyankabhatter88@yahoo.com) [psbhatt@health.usf.edu](mailto:psbhatt@health.usf.edu) [Google Scholar](#) [ResearchGate](#) [Website](#)

**Dr. Siddhartha Pati**, Research Associate, Centre of Excellence, Khallikote University, Odisha, India. +91-8895577750/7008251372 [Patisiddhartha@gmail.com](mailto:Patisiddhartha@gmail.com) [ResearchGate](#) [ORCID](#)

**Dr. Iqrar A. Ansari**, Assistant Professor, Department of Pharmaceutical Chemistry, Prof. Ravindra Nikam College of Pharmacy, Gondur, Dhule, 424002, Maharashtra, India. Mobile No.+91 9284541386, Email: [ansariiqrar50@gmail.com](mailto:ansariiqrar50@gmail.com), [Google Scholar](#), [Researchgate](#), [LinkedIn](#) Research Fields: [EGFR Inhibitors](#), [CADD](#), [Docking](#), [MD Simulation](#)

**Dr. Prakash Doddapattar**, Carver College of Medicine, University of Iowa, USA. Research Fields: Molecular Biology · Immunology · Cardiovascular Biology. [ResearchGate](#) [LinkedIn](#) [ORCID ID](#) Email: [prakashdpbt@gmail.com](mailto:prakashdpbt@gmail.com)

**Dr. Ashok K Datusalia**, Assistant Professor, National Institute of Pharmaceutical Education and Research, Raebareilly, Uttar Pradesh, 226002, India Email: [ashokdatusalia@gmail.com](mailto:ashokdatusalia@gmail.com) [Google Scholar](#) [ResearchGate](#) [Website](#) Research Interests: Neuroscience, Pharmacology and Toxicology

**Dr. Dinesh Nyavanandi**, Formulation Scientist Continuus Pharmaceuticals 25 Olympia Ave, Woburn, Massachusetts, USA. 01801. (P): +1(662) 550-9555 (E): [ndinesh624@gmail.com](mailto:ndinesh624@gmail.com); [dnyavanandi@continuuspharma.com](mailto:dnyavanandi@continuuspharma.com) *Research Experience:* Oral drug delivery, Lipid based drug delivery, Strategies for enhancement of solubility and bioavailability, Gastro-retentive Drug delivery, Ocular Drug Delivery, Topical Drug Delivery [LinkedIn](#) [Google Scholar](#) [ORCID](#) [SCOPUS](#)

## INTERNATIONAL EDITORIAL MEMBERS

**Prof. Dr. rer. Nat. Hesham A. El Enshasy**, Director, Institute of Bioproduct Development (IBD), Universiti Teknologi Malaysia (UTM), 81310 UTM, Johor Bahru, Malaysia. Tel: 006-07-5531573 (office), Fax: 006-07-5569706, E-mail: [henshasy@ibd.utm.my](mailto:henshasy@ibd.utm.my)  
[ResearchGate](#) [Google Scholar](#)

**Dr. Arif Nur Muhammad Ansori**, Assistant Lecturer, Faculty of Science and Technology, Universitas Airlangga, Surabaya, Indonesia. Email: [arif.nma-17@fkh.unair.ac.id](mailto:arif.nma-17@fkh.unair.ac.id) [ansori.anm@gmail.com](mailto:ansori.anm@gmail.com) [Website](#) [Google Scholar](#) [ORCID](#) [LiveDNA](#) [LinkedIn](#)

**Dr. Sitansu Sekhar Nanda**, Department of Chemistry, Myongji University, Yongin, South Korea, Email: [nandasitansusekhar@gmail.com](mailto:nandasitansusekhar@gmail.com) [Google Scholar](#) [ResearchGate](#) [Website](#)

**Dr. Kishor Mazumder**, Associate Professor At Jashore University Of Science And Technology, Chattogram, Bangladesh Email: [Mazumderkis2002@Yahoo.Com](mailto:Mazumderkis2002@Yahoo.Com) [LinkedIn](#) [Google Scholar](#) [ResearchGate](#)

**Dr. Priya Bharate**, Analytical Scientist, Continuus Pharmaceuticals, MA, USA. [Google Scholar](#) [ResearchGate](#) [Website](#) Email: [pbharate@continuuspharma.com](mailto:pbharate@continuuspharma.com) Interested fields: Carbohydrates synthesis, Chemical biology, Supramolecular chemistry, Analytical method development and qualifications of drug molecules, Nanomaterials, and organic chemistry

**Dr. Ahmed Salih Sahib**, Professor, Dept of Pharmacology & Toxicology, College of Pharmacy, University of Kerbala, Iraq, Email: [ahmedsalih73@yahoo.com](mailto:ahmedsalih73@yahoo.com) [ahmed.alkhezaly@uokerbala.edu.iq](mailto:ahmed.alkhezaly@uokerbala.edu.iq) Tel : +964-7901585579 [Google Scholar](#)  
[ResearchGate](#) [Website](#) Interested fields: Pharmacology, Clinical Pharmacology, Pharmacogenomics

**Prof. Fahad I. Al-Saikhan**, Department of Clinical Pharmacy, Prince Sattam Bin Abdulaziz University, College of Pharmacy, Kingdom of Saudi Arabia Email: [f.alsaikhan@psau.edu.sa](mailto:f.alsaikhan@psau.edu.sa) Telephone: +966115886074 [ResearchGate](#) [Website](#) Interested fields: Primary care/Ambulatory care, health outcome, pharmacogenomics

**Dr. Nahlah Elkudssiah Ismail**, Professor, Department of Clinical Pharmacy, Malaysian Academy of Pharmacy, Malaysia. [researchgate.net](#) [orcid.org](#) [LinkedIn](#) Email: [elkudssiah77@yahoo.com](mailto:elkudssiah77@yahoo.com) Research interest: Pharmaceutics (drug delivery; nanotechnology), Pharmaceutical Analysis (method development and validation), Clinical Pharmacy, Pharmacy Education, Pharmacy Practice & Public Health.

**Dr. Vijai Kumar Reddy Tangadanchu**, Ph. D. Postdoctoral Research Associate, Mallinckrodt Institute of Radiology, Washington University School of Medicine Saint Louis, Missouri-63110, USA E-mail: [tvjajikumarreddy@wustl.edu](mailto:tvjajikumarreddy@wustl.edu) [tvkreddyict@outlook.com](mailto:tvkreddyict@outlook.com) Mobile: +1 314-665-8974. [Google Scholar](#) [Publons](#)

**Dr. Solmaz Ghaffari**, Department of Pharmaceutics, Faculty of Pharmacy, Islamic Azad University, Tehran, Iran; Email: [soligh@yahoo.com](mailto:soligh@yahoo.com) [Google Scholar](#) [researchgate.net](http://researchgate.net) Tel: +9891252721 Fax: +982122600091

**Dr. Mostafa Madmoli**, Emergency Medical Technician, Dezful University of Medical Sciences, Dezful, Iran. Email: [mostafamadmoli10@yahoo.com](mailto:mostafamadmoli10@yahoo.com) Phone: [989370176590] Interested fields: Diabetes, Alzheimer's Disease,

**Dr. Vijay Kumar Siripuram**, Laboratory of Molecular and Cellular Neuroscience The Rockefeller University, New York, NY, USA. Email: [vijay.aleru@gmail.com](mailto:vijay.aleru@gmail.com) ; Phone: (+1) 646-238-5620. Interested fields: NMR, FTIR, HPLC, GCMS/LCMS, column chromatography, crystallization [Google Scholar](#) [Publons](#)

**Dr Saket Patel**, Research Associate at Memorial Sloan Kettering Cancer Center, Department of Radiology, New York, 10065, USA. Interested fields: Organic chemistry, new molecule design, developing new methods for hyperpolarization via dissolution DNP, and real time metabolism (in vivo/ in vitro). Phone: +1 929-553-4960 Email: [saketpatel32@gmail.com](mailto:saketpatel32@gmail.com) [patels24@mskcc.org](mailto:patels24@mskcc.org)  
[Google Scholar](#) [Website](#)

**Dr Arun Butreddy**, Ph.D., Scientist, Protein Pharmaceutical Development, Biogen Inc, Massachusetts, USA. [ResearchGate](#) [Google Scholar](#) [LinkedIn](#) Email: [butreddyaran@gmail.com](mailto:butreddyaran@gmail.com) Mobile: +1 (662)-380-0558

**Dr. Madhukiran Parvathaneni**, Manager - Global Regulatory Affairs, CRC Pharma, New Jersey, USA. Interested Fields: Regulatory Affairs, Clinical Research, Pharmaceuticals and Medical Devices. Ph: +1 (609)582-6496 Email: [madhukiran.parvathaneni@gmail.com](mailto:madhukiran.parvathaneni@gmail.com) [Website](#) [ResearchGate](#) [Google Scholar](#) [LinkedIn](#)

**Praveen Kolimi**, Department of Pharmaceutics and Drug Delivery, University of Mississippi, Faser Hall, MA, 38677, USA  
Mail: [praveenkoliminiper@gmail.com](mailto:praveenkoliminiper@gmail.com) [Google Scholar](#) [ResearchGate](#) [LinkedIn](#)

#### EXECUTIVE EDITORS

**Dr. Sheo Datta Maurya**, Sarvodaya College of Pharmacy, Azamgarh, India Email: [meetshiv23@gmail.com](mailto:meetshiv23@gmail.com) [Google Scholar](#) [ResearchGate](#) Interested fields: Pharmacy (Novel Drug delivery, Liposomes, Ethosomes,)

#### ASSISTANT EDITORS

**Dr. Dito Anurogo, M.D., M.Sc.**, International PhD Program for Cell Therapy and Regeneration Medicine (IPCTRM), College of Medicine, Taipei Medical University Taiwan and Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Makassar, Makassar, Indonesia. Mail: [dito.anurogo@med.unismuh.ac.id](mailto:dito.anurogo@med.unismuh.ac.id) [d151109004@tmu.edu.tw](mailto:d151109004@tmu.edu.tw) Phone: +62-81224422693. [Google Scholar](#) [Publons](#) [LinkedIn](#) Research Interest: drugs and delivery, pharmacology, regenerative medicine, oncohematology, neuroscience, molecular biology & medicine, bioinformatics, nanomedicine.

**Dr. Aktsar Roskiana Ahmad**, Faculty of Pharmacy, Universitas Muslim Indonesia, Urip Sumoharjo Km. 5, 90231, Indonesia. Email: [aktsar.roskiana@umi.ac.id](mailto:aktsar.roskiana@umi.ac.id) Phone: +6281355184404 [Google Scholar](#) [ResearchGate](#)

**Dr. Sandeep K. Singh**, Postdoctoral Research Associate, Department of Pharmacy Practice and Science, University of Nebraska Medical Center, Omaha, NE, USA. [Google Scholar](#) [sasinghlko@gmail.com](mailto:sasinghlko@gmail.com)

**Dr. Mousumi Kar**, Professor and Head, Faculty of Pharmacy Medi-Caps University, Indore, INDIA, India. Email: [karmousumi@hotmail.com](mailto:karmousumi@hotmail.com) [Website](#) [Google Scholar](#) [ResearchGate](#) [LinkedIn](#) [Website](#) Interested fields: Pharmaceutics, Novel Drug Delivery

**Mr. Vaibhav Patel**, Director (Quality Assurance), Elucida Oncology Inc., 1 Deer Park Drive, Suite E Monmouth Junction, NJ 08852, USA. Website Email: [vpatel0212@gmail.com](mailto:vpatel0212@gmail.com) [Google Scholar](#) Phone No: (551) 208-5652

**Mr. Girish Nihalani**, Research Fellow of Drug Product Development at Teva Pharmaceuticals Inc. (Actavis) based at Florida, USA. e-mail: [girlani679@gmail.com](mailto:girlani679@gmail.com) [girish9122@gmail.com](mailto:girish9122@gmail.com) Contact: +1- 954-647-7740 [LinkedIn](#) [Google Scholar](#) Interested fields: Pharmaceutics, Novel Drug Delivery

## Indexing

1. [EBSCO](#) (Elsevier's Database)
2. [Sherpa Romeo](#)
3. [Global Health](#)
4. [CAB Abstracts](#)
5. [Index Copernicus, ICV 2022:100](#)
6. [Scilit](#)
7. [SCITE](#)
8. [EXALY](#)
9. [ScienceGate](#)
10. [NLM \(ID:101656626\)](#)
11. [ROAD](#)
12. [CAS Source Index](#) (CASSI, American Chemical Society)
13. [Publons](#)
14. [Google Scholar](#)
15. [Geneva Foundation of Medical Education & Research](#) (GFMER)
16. [WorldCat](#) (ID: 879704190)
17. [MIAR](#) (Information Matrix for analysis of Journals)
18. [SCISPACE](#)
19. [Open Academic Journal Index](#) (OAJI)
20. [Bielefeld Academic Search Engine](#) (BASE)
21. [Veterinary Science Database](#)
22. [Review of Medical and Veterinary Entomology](#)
23. [Review of Aromatic and Medicinal Plants](#)
24. [Open University of Malaysia Library](#) OUM Control No: [000330909](#)

## Membership

1. [ICMJE](#)
2. [Portico](#)
3. [CrossRef](#)

Available online on 15.07.2023 at <http://jddtonline.info>

# Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

Copyright © 2023 The Author(s): This is an open-access article distributed under the terms of the CC BY-NC 4.0 which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited



Open Access Full Text Article



Check for updates

Research Article

## Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years

Meiyanti\*<sup>ORCID</sup>, Yohana<sup>ORCID</sup>, Eveline Margo<sup>ORCID</sup>, Juni Chudri, Pusparini<sup>ORCID</sup>, Meutia Atika Faradilla<sup>ORCID</sup>

Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

### Article Info:

### Abstract



#### Article History:

Received 08 May 2023  
Reviewed 11 June 2023  
Accepted 24 June 2023  
Published 15 July 2023

#### Cite this article as:

Meiyanti, Yohana, Margo E, Chudri J, Pusparini, Faradilla MA, Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years, Journal of Drug Delivery and Therapeutics. 2023; 13(7):52-56

DOI: <http://dx.doi.org/10.22270/jddt.v13i7.6142>

#### \*Address for Correspondence:

Meiyanti, Department of Pharmacology and Medical Pharmacy, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Aging is a progressive process caused by a physiological decline in body functions. It is known that several factors can accelerate aging, namely age, lifestyle, oxidative stress, lipid peroxidation, and mitochondrial damage. Various pollutants such as cigarette smoke, motor vehicle fumes, industrial waste, and exposure to sunlight / ultraviolet light will form free radicals. Malondialdehyde (MDA) level is one of the indicators used to measure oxidative stress. Increased MDA levels are found in various degenerative diseases. This study aims to determine plasma MDA levels in the age group of 40 years and over and factors that can affect plasma MDA levels. This study was a cross-sectional study by including 42 subjects. Data collection was carried out using a questionnaire to obtain subject demographic data and physical activity, as well as physical examinations complemented by laboratory examinations and MDA levels. The mean age of the subjects was 52.7 years, and most of the 32 (76.2%) subjects were categorized as obese and had high activity. Pearson correlation test obtained body mass index statistically significantly correlated with plasma MDA levels with a value of  $p = 0.039$ , as well as physical activity statistically significantly correlated with a negative correlation, with a value of  $r = -0.26$  and  $p = 0.043$ . Conclusion Body mass index and activity are significantly correlated with plasma MDA levels.

**Keywords:** malondialdehyde, stress, oxidative, radicals

## INTRODUCTION

Aging is a progressive process caused by a physiological decline in body function.<sup>1</sup> A decrease in body homeostasis is accompanied by an increased risk of various chronic diseases including diabetes mellitus, cancer, neurodegenerative diseases, and heart disease. It is known that several factors can accelerate aging, namely age, lifestyle, oxidative stress, lipid peroxidation, and mitochondrial damage. Previous studies have shown that changes in sphingomyelin metabolism in the central nervous system and age are associated with neurodegenerative diseases.<sup>2</sup> In the US, aging is one of the most important factors in diabetes mellitus and heart disease. In 2015, an estimated 12% of the population had diabetes mellitus and an undetected 24% had diabetes mellitus. Heart disease is the highest contributor to mortality in the population aged 65-84 years.<sup>3</sup> Recent studies have shown that non-diabetic populations whose fasting glucose levels increase in the blood will increase the risk of coronary heart disease. But on the other hand, the data on fasting blood glucose increase is still debated in the population of cardiovascular disease patients who do not have diabetes mellitus. A recent study in North Sumatra showed a significant relationship between MDA levels in obese and non-obese populations.<sup>4</sup>

Free radicals are formed through two processes, namely

exogenous and endogenous processes. Various pollutants such as cigarette smoke, motor vehicle fumes, industrial waste, and exposure to sunlight/ultraviolet will form free radicals. Illness, stress, and excessive exercise/physical activity are endogenous factors produced by the human body that can form free radicals. The formation of free radicals occurs due to oxidation processes and metabolic processes that take place during exercise or activity, inflammatory processes, consumption of certain foods, and others. Free radicals will begin to activate with molecules in the surrounding cells to obtain electrons to stabilize the condition of one's body. The electrons that have been taken in the cell molecules will trigger the occurrence of free radicals. If this continues for a long time, it can cause oxidative stress which can cause inflammation, and DNA disease can also trigger the disease. Oxidative stress conditions characterized by increased production of Reactive Oxygen Species (ROS) have implications for various diseases such as hypertension, atherosclerosis, diabetes mellitus, stroke, chronic renal failure, heart failure, and other chronic diseases.<sup>5-7</sup>

Oxidative stress is an imbalance between oxidants (free radicals) and antioxidants. Free radicals that commonly cause oxidative damage are reactive oxygen species (ROS).<sup>8</sup> ROS compounds are formed through aerobic metabolism. These compounds can cause extensive damage through catalytic reactions. They are responsible for damage at the cellular level

by oxidizing proteins, lipids, DNA (Deoxyribonucleic acid), and other biomolecules. Various compounds can be used as markers of oxidative stress such as malondialdehyde (MDA), and protein carbonylation. 4-hydroxy-2-nonenal (4HNE), and 8-hydroxy-deoxyguanosine (8OHdG).<sup>9</sup>

Malondialdehyde (MDA) is a result of lipid peroxidation in the body. The concentration of MDA indicates an oxidation process that occurs in the body's cell membranes due to metabolic processes. One of the factors that cause aging is the production of excess reactive oxygen species (ROS). ROS are generated through the process of cell respiration in the mitochondria and the phagocytosis process of foreign body elimination. To stabilize the amount of ROS, the body can use endogenous and exogenous antioxidants. Disruption of the balance between ROS and antioxidants can cause irreversible damage to cell organelle components up to the DNA level.<sup>10,11</sup>

MDA levels are strongly influenced by many factors such as age, illness, stress, various pollutants, food consumption, antioxidant supplements and excessive physical activity or exercise, and others. The other study found that more than 30% of subjects had high MDA levels, and concluded that there was a significant relationship between eating habits, body mass index, glucose levels with MDA levels, as well as smoking habits significantly related to MDA levels.<sup>12</sup> Recent studies have shown that plasma MDA levels in populations aged 70 years and 60 years are not significantly different but there is an increase in the specific activity of the catalase enzyme that increases with age.<sup>10</sup> In another study, there was a significant difference in plasma MDA levels in the obesity group without metabolic syndrome compared to the obesity group with metabolic syndrome.<sup>4</sup>

There are still variations in MDA data in various groups with different ages and different disease groups, so this study aims to determine plasma MDA levels in the age group 40 years and over and factors that can affect plasma MDA levels.

## METHOD

This observational analytic study with a cross-sectional design approach was conducted from November 2022 to February 2023 in Angke village, West Jakarta, Indonesia. The inclusion criteria for this study were: men and women aged  $\geq 40$  years, willing to participate in the study and sign informed consent. Meanwhile, the exclusion criteria were having a history of malignancy, kidney failure, a history of autoimmune disease, taking drugs related to psychiatric conditions, and having limitations in mobility. This study used the calculation of sample size with the formula  $N / 1 + N.\alpha^2$ , and obtained the minimum number of samples required was 40 subjects. Demographic data and physical activity were obtained from distributed questionnaires. Blood chemistry examination (complete blood, glucose, total cholesterol, liver enzymes (SGOT and SGPT) using an enzymatic method. Examination of malondialdehyde levels was carried out using the spectrophotometric method. This research has obtained approval from the ethics committee of the Faculty of Medicine, Trisakti University, Number: 178/KER/FK/1X/2022.

## RESULTS

A total of 42 subjects participated in this study, most of the subjects were female 27 (64.3%), and the mean age of the subjects was 52.7 years. Anthropometric data of the subjects are as follows body weight  $67.53 \pm 13.64$  and height  $156.02 \pm 8.73$  cm, while for abdominal circumference the average is 93.09 cm. Most of the subjects 59.5% fell into the normotensive category, with an average systolic blood pressure of 141.62 mmHg and an average diastolic blood pressure of 82.45 mmHg. A total of 32 (76.2%) subjects were

categorized as obese with BMI  $\geq 25.0$ , and most subjects had high activity levels. (Table 1)

Table 1: Distribution of Subject Characteristics (n=42)

Characteristics	n (%)	Mean $\pm$ SD
Gender		
Male	15 (35.7)	
Female	27 (64.3)	
Blood pressure (mmHg)		
Normal (< 140/90)	25 (59.5)	
Hypertension ( $\geq 140/90$ )	17 (40.5)	
Physical Activity		
Low (< 600 Mets)	11 (26.2)	
High ( $\geq 600$ Mets)	31 (73.8)	
Body Mass Index (kg/m <sup>2</sup> )		27.69 $\pm$ 4.90
Not obese (< 25.0)	10 (23.8)	
Obese ( $\geq 25.0$ )	32 (76.2)	
Age (years)		52.7 $\pm$ 10.60
Body weight (kg)		67.53 $\pm$ 13.64
Height (cm)		156.02 $\pm$ 8.73
Abdominal circumference (cm)		93.09 $\pm$ 10.54
Blood pressure (mmHg)		
Systole		141.62 $\pm$ 20.77
Diastole		82.45 $\pm$ 12.70

The examination of the subject's plasma malondialdehyde levels obtained a mean of 0.77 with the highest value of 1.03 and the lowest of 0.51. Hemoglobin levels averaged 13.16 g/dL. Fasting blood glucose level was  $95.06 \pm 41.01$  mg/dL and total cholesterol was  $179.83 \pm 31.85$  mg/dL. Laboratory examinations to determine liver function were carried out by measuring SGPT and SGOT levels, obtaining mean values of 24.21 U/L and 26.31 U/L respectively. (Table 2)

Table 2: Laboratory examination results of subjects (n=42)

Variable	Mean $\pm$ SD
Hemoglobin (g/dL)	13.16 $\pm$ 1.56
Fasting blood glucose (mg/dL)	95.06 $\pm$ 41.01
Total cholesterol (mg/dL)	179.83 $\pm$ 31.85
SGPT (U/L)	24.21 $\pm$ 13.50
SGOT (U/L)	26.31 $\pm$ 11.51
Plasma malondialdehyde	0.77 $\pm$ 0.26

Determining the type of correlation test to be used by first conducting a data normality test. The normality test used was Shapiro-Wilk because the number of subjects was less than 50 subjects. The correlation test used to assess the correlation between blood glucose, total cholesterol, SGPT, SGOT levels, age, body mass index and physical activity with plasma MDA levels was the Pearson correlation test. The test results are presented in Table 3. Blood glucose, total cholesterol, SGPT,

SGOT, and age did not have a significant correlation with MDA levels. Body mass index has a significant correlation with plasma MDA ( $p = 0.039$ ), with a positive correlation direction, and the strength of the correlation is statistically weak with a value of  $r = 0.396$  ( $<0.4$ ). The physical activity variable has a significant correlation with plasma MDA levels ( $p=0.043$ ) with a negative correlation direction but with a statistically weak correlation strength ( $r=-0.26$ ).

Table 3: Correlation Test of Age, Gender, Glucose, SGOT, SGPT, BMI, Physical Activity with Plasma MDA Levels (n=42)

	Plasma MDA Level	
	r	p
Age	0.098	0.756
Gender	0.124	0.108
Glucose	0.330	0.835
SGOT	0.104	0.513
SGPT	0.008	0.614
Body Mass Index	0.396	0.039*
Physical activity	-0.260	0.043*

\*  $p < 0.05$

## DISCUSSION

The level of oxidative stress can be measured using Malondialdehyde (MDA) concentration. MDA is the result of lipid peroxidation, which will cause cell damage. Oxidant compounds such as free radicals or non-radical species attack lipids that contain carbon-carbon double bonds, especially unsaturated fatty acids (PUFA). MDA is one of the biomarkers widely used to assess oxidative stress. The mean serum MDA in this study 0.77 was found to be lower than the previous study<sup>10</sup>, this is because the age difference of the subjects in this study was younger than in the previous study. In contrast, a study conducted in Iraq on hypertensive patients concluded that the increase in MDA was associated with an increase in age, gender, and duration of disease.<sup>13</sup>

Free radicals such as ROS are the etiology of most chronic diseases, as it is known that active free radicals are the result of metabolic reactions. Damage to body tissue cells occurs because active free electrons remain looking for other electrons to settle, if no other electrons are available then active free electrons will attack cells resulting in cell damage. Although free radicals naturally have benefits for the body, when the level of free radicals exceeds the normal value, it will cause harm to the body.<sup>11,12</sup>

Measurement of oxidative stress can be done by measuring plasma malondialdehyde (MDA) levels. MDA is a secondary result of the lipid peroxidation process, producing various active compounds that will damage cells. The process of MDA formation through enzymatic and non-enzymatic processes. Currently, MDA is one of the biomarkers used to assess oxidative stress.<sup>14</sup> Some previous researchers concluded that higher serum MDA was found in patients with hyperlipidemia. The condition of increased fat content in the body is in line with increased oxidative stress. Hyperlipidemia is associated with physical changes in cell membranes that facilitate the release of free radicals from the mitochondrial electron transport chain or activation of NADPH oxidase. In patients with diabetes mellitus (DM) there is also an increase in MDA levels, malondialdehyde interacts irreversibly and reversibly with proteins and phospholipids to have a profound effect, especially on the collagen of the cardiovascular system. MDA levels remain within normal limits due to the antioxidant

effect, but in DM, this protective effect is impaired resulting in increased MDA levels.<sup>1,15</sup>

Increased lipid peroxidation is considered a consequence of oxidative stress that occurs when the dynamic balance between prooxidants and antioxidants is disrupted. The statin class of drugs for hyperlipidemia therapy is proven to be anti-atherogenic, can improve endothelial function, inhibit vascular inflammation, and stabilize atherosclerotic plaques. In addition to statins, non-steroidal anti-inflammatory drugs (NSAIDs) can affect MDA concentration. Previous studies have found that NSAIDs have significant antioxidant properties to scavenge free radicals.<sup>16,17</sup>

In this study, there was no significant correlation between age and MDA levels. Similar results were also obtained by other researchers, cell damage due to oxidative stress is not related to increasing age but is associated with habits, behavior patterns, and physiological changes due to age. Various habitual patterns such as smoking, fatty food habits, lack of activity, and inflammatory processes can lead to increased oxidative stress. The habit of taking vitamin C (ascorbic acid) has a beneficial impact due to its antioxidant effect so cell damage and DNA damage are lower in this group.<sup>18</sup> MDA levels were significantly higher among all levels of smokers. MDA levels were lower in light smokers compared to heavy or moderate smokers. Higher vitamin C levels were found in the non-smoker group. All body tissues are susceptible to free radical damage, but based on location, the respiratory epithelium is particularly vulnerable to cigarette smoke. Oxygen radicals generated close to the cell membrane oxidize membrane phospholipids which will result in lipid peroxidation with the end products of this reaction being malondialdehyde, ethane, and pentane. this causes an increase in MDA levels in smokers.<sup>19</sup> Another study conducted in Klaten on foundry workers concluded that type of work, body mass index and smoking habits did not affect MDA levels, but length of work and marital status were associated with MDA levels. Long and continuous exposure to metallic materials will result in the accumulation of these materials in the body and will increase the oxidative process of polyunsaturated fats in cell membranes and worsen the level of lipid peroxidase, characterized by an increase in MDA levels as a reflection of the intensity and rate of lipid peroxidase reactions of damage to body cells.<sup>20</sup>

Hardiany's study also concluded that there was no significant difference in plasma MDA levels in the age population, but increasing age along with an increase in catalase-specific activity.<sup>10</sup> This increase in catalase-specific activity is a protective effect especially for elderly women to overcome high levels of oxidative damage. The role of endogenous antioxidant catalase is to suppress oxidative damage.<sup>18,21</sup> Oxidative stress plays an important role in the incidence of cardiovascular diseases including coronary artery disease. Oxidative stress is considered an important mechanism for the development of cardiovascular diseases especially atherosclerosis. Hypertension, cardiac hypertrophy, and ischemia-reperfusion injury are three other major cardiovascular conditions for which gender is a risk factor. Gender is also associated with differences in oxidative stress. Under physiological conditions, women appear to be less susceptible to oxidative stress. This may be due to the antioxidant properties of estrogen, sex differences in NADPH-oxidase activity or other mechanisms yet to be determined. Sex hormones including estrogen are globally present with their receptors in the heart so sex hormones can exert effects on the cardiovascular system. Estrogen can improve heart function and reduce injury to the heart. In addition, estrogen has antioxidant effects so that it can function as a cardioprotective.<sup>22,23</sup>



This study found a significant correlation between body mass index and MDA levels, the higher the BMI value, the higher the MDA levels. Similar results were also obtained in a study in India that concluded that in metabolic syndrome patients with one of the signs is an increase in BMI, there is an increase in MDA levels compared to the control group. In obese subjects, the level of MDA is higher than normal IMT. In a healthy body, there is redox hemostasis that occurs in cells, if an imbalance arises it will cause oxidative stress. Obese patients will have an increased metabolic load and mechanical load on the myocardium, increased body mass, and excessive food intake will cause the formation of lipid peroxidation, and free radicals.<sup>24</sup> The increase in some of these things will stimulate antioxidant enzymes in the long run, this will cause antioxidant enzyme reserves to be depleted and unable to overcome and balance the increase in oxidative stress.<sup>25,26</sup>

These types of training activities are exhausting, long exercise, overtraining syndrome, various types of exercise will induce a significant response to oxidative stress. Exercise activities such as light exercise, low-intensity exercise, and prolonged training will have an impact on improving endogenous antioxidant status. Reactive oxygen species play an important role in cell signaling and in regulating antioxidant gene expression. Exertion results in hyperregulation of nuclear factor kappa B and mitogen-activated protein kinases that activate gene expression of a number of enzymes and proteins with important roles in maintaining intracellular oxidative or antioxidant homeostasis.<sup>1,6,27</sup> The limitation of this study is that some factors that may affect MDA levels such as consumption of vitamin supplements, diet, and length of illness suffered were not examined in this study.

## CONCLUSIONS

Body mass index and activity were significantly correlated with MDA levels, while age, gender, blood glucose levels, and liver enzymes were not significantly correlated with MDA levels.

## Conflict of Interest:

The investigators declare there is no conflict of interest in this study.

## Acknowledgments

The authors thank all the subjects who participated in this study, as well as the laboratory staff who assisted in the implementation of this study.

## REFERENCES

- Sharifi-Rad M, Anil Kumar N V., Zucca P, Varoni EM, Dini L, Panzarini E, et al. Lifestyle, Oxidative Stress, and Antioxidants: Back and Forth in the Pathophysiology of Chronic Diseases. *Front Physiol.* 2020; 11(July):1-21. <https://doi.org/10.3389/fphys.2020.00694>
- Park MH, Jin HK, Bae J sung. Potential therapeutic target for aging and age-related neurodegenerative diseases: the role of acid sphingomyelinase. *Exp Mol Med [Internet].* 2020; 52(3):380-9. <https://doi.org/10.1038/s12276-020-0399-8>
- Całyniuk B, Grochowska-Niedworok E, Walkiewicz K, Kawecka S, Popiołek E, Fatyga E. Malondialdehyde (MDA) - product of lipid peroxidation as marker of homeostasis disorders and aging. *Ann Acad Medicae Silesiensis.* 2016; 70:224-8. <https://doi.org/10.18794/aams/65697>
- Savira M, Rusdiana ., Widjaja SS, Syahputra M. Comparison Malondialdehyde (MDA) Level between Obesity Non Metabolic Syndrome and Obesity with Metabolic Syndrome Patients. 2020; (5):644-7.
- Yekti R, Bukhari A, Jafar N, Thaha AR. Measurement of malondialdehyde (MDA) as a good indicator of lipid peroxidation.

- Int J Allied Med Sci Clin [Internet]. 2018; 6(4):6-8. Available from: [www.ijamsc.com](http://www.ijamsc.com)
- Denise Mafra ME. Influence of Physical Exercise on Oxidative Stress and Inflammation in Hemodialysis Patients. *J Mol Biomark Diagn.* 2015; 06(03). <https://doi.org/10.4172/2155-9929.1000e120>
- Mao C, Yuan JQ, Lv Y Bin, Gao X, Yin ZX, Kraus VB, et al. Associations between superoxide dismutase, malondialdehyde and all-cause mortality in older adults: A community-based cohort study. *BMC Geriatr.* 2019; 19(1):1-9. <https://doi.org/10.1186/s12877-019-1109-z>
- Ayala A, Muñoz MF, Argüelles S. Lipid peroxidation: Production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. *Oxid Med Cell Longev.* 2014; 2014. <https://doi.org/10.1155/2014/360438>
- Situmorang N, Zulham Z. Malondialdehyde (Mda) (Zat Oksidan Yang Mempercepat Proses Penuaan). *J Keperawatan Dan Fisioter.* 2020; 2(2):117-23. <https://doi.org/10.35451/jkf.v2i2.338>
- Hardiany NS, Sucitra S, Paramita R. Profile of malondialdehyde (MDA) and catalase specific activity in plasma of elderly woman. *Heal Sci J Indones.* 2020; 10(2):132-6. <https://doi.org/10.22435/hsji.v12i2.2239>
- Hardiany NS, Sadikin M, Siregar N, Wanandi SI. The suppression of manganese superoxide dismutase decreased the survival of human glioblastoma multiforme T98G cells. *Med J Indones.* 2017; 26(1):19-25. <https://doi.org/10.13181/mji.v26i1.1511>
- 439-445\_nova.pdf.
- A. Hassan A, Sayyah SG. Oxidative Stress Marker Malondialdehyde and Glutathione Antioxidant in Hypertensive Patients. *Eur J Biomed Res.* 2023; 2(1):31-6. <https://doi.org/10.24018/ejbiomed.2023.2.1.47>
- Khoubnasab Jafari M, Ansarin K, Jouyban A. Comments on "use of malondialdehyde as a biomarker for assessing oxidative stress in different disease pathologies: A review." *Iran J Public Health.* 2015; 44(5):714-5.
- Shalash M, Badra M, Imbaby S, ElBanna E. Malondialdehyde in Type 2 Diabetics and Association With Cardiovascular Risk Factors. *J Med Res Inst.* 2020; 41(2):21-30. <https://doi.org/10.21608/jmalexu.2020.147116>
- Oesterle A, Laufs U, Liao JK. Pleiotropic Effects of Statins on the Cardiovascular System. *Circ Res.* 2017; 120(1):229-43. <https://doi.org/10.1161/CIRCRESAHA.116.308537>
- Sekunda KN, Ariosta A, Limantoro C, Setiawan AA. Difference in Profiles of Oxidative Stress Marker (MDA) in STEMI and NSTEMI. *Diponegoro Med J (Jurnal Kedokt Diponegoro).* 2021; 10(2):145-50. <https://doi.org/10.14710/dmj.v10i2.29673>
- Block G, Dietrich M, Norkus EP, Morrow JD, Hudes M, Caan B, et al. Factors associated with oxidative stress in human populations. *Am J Epidemiol.* 2002; 156(3):274-85. <https://doi.org/10.1093/aje/kwf029>
- Kubihal C V., Naik HD. Effect of smoking on vitamin C and MDA: a cross sectional comparative study. *Int J Res Med Sci.* 2019; 7(3):746. <https://doi.org/10.18203/2320-6012.ijrms20190511>
- Ari Suwondo, Ahmad Mohan, Melyana Nurul Widyawati M. Factors Related to Malondialdehyde (MDA) Levels in Metal Casting Industry Workers. *Preprints.* 2020; 1(October):1-8. <https://doi.org/10.21203/rs.3.rs-99597/v1>
- Schöttker B, Brenner H, Jansen EHJM, Gardiner J, Peasey A, Kubínová R, et al. Evidence for the free radical/oxidative stress theory of ageing from the CHANCES consortium: A meta-analysis of individual participant data. *BMC Med.* 2015; 13(1):1-15. <https://doi.org/10.1186/s12916-015-0537-7>
- Kander MC, Cui Y, Liu Z. Gender difference in oxidative stress: a new look at the mechanisms for cardiovascular diseases. *J Cell Mol Med.* 2017; 21(5):1024-32. <https://doi.org/10.1111/jcmm.13038>
- Bell JR, Bernasochi GB, Varma U, Raaijmakers AJA, Delbridge LMD. Sex and sex hormones in cardiac stress - Mechanistic insights. J

- Steroid Biochem Mol Biol. 2013; 137:124-35.  
<https://doi.org/10.1016/j.jsbmb.2013.05.015>
24. P G, Curcio F, Bulli G, Aran L, Della-morte D, Testa G, et al. Oxidative Stress and Diseases. *Oxidative Stress Dis.* 2012; 757-72.
25. Sabitha K, Venugopal B, Rafi M, V Ramana K. Role of Antioxidant Enzymes in Glucose and Lipid Metabolism in Association with Obesity and Type 2 Diabetes. *Am J Med Sci Med.* 2014; 2(1):21-4.  
<https://doi.org/10.12691/ajmsm-2-1-5>
26. Manna P, Jain SK. Obesity, Oxidative Stress, Adipose Tissue Dysfunction, and the Associated Health Risks: Causes and Therapeutic Strategies. *Metab Syndr Relat Disord.* 2015; 13(10):423-44. <https://doi.org/10.1089/met.2015.0095>
27. Lewis NA, Howatson G, Morton K, Hill J, Pedlar CR. Alterations in Redox Homeostasis in the Elite Endurance Athlete. *Sport Med.* 2015; 45(3):379-409. <https://doi.org/10.1007/s40279-014-0276-5>

# Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years

*by Meiyanti Meiyanti*

---

**Submission date:** 28-Jun-2023 08:09PM (UTC+0700)

**Submission ID:** 2123934068

**File name:** aldehyde\_Levels\_in\_People\_over\_40\_Years\_JDDT\_Meiyanti\_et\_al.docx (61.09K)

**Word count:** 3703

**Character count:** 20873

# Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years

Meiyanti\*, Yohana, Eveline Margo, Juni Chudri, Pusparini, Meutia Atika Faradilla

Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Corresponding Email: [meiyanti@trisakti.ac.id](mailto:meiyanti@trisakti.ac.id)

3

Address for correspondence: Meiyanti, Department of Pharmacology and Medical Pharmacy, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

## Acknowledgments 13

The authors thank all the subjects who participated in this study, as well as the laboratory staff who assisted in the implementation of this study.

**Abstract.** Aging is a progressive process caused by a physiological decline in body functions. It is known that several factors can accelerate aging, namely age, lifestyle, oxidative stress, lipid peroxidation, and mitochondrial damage. Various pollutants such as cigarette smoke, motor vehicle fumes, industrial waste, and exposure to sunlight / ultraviolet light will form free radicals. Malondialdehyde (MDA) level is one of the indicators used to measure oxidative stress. Increased MDA levels are found in various degenerative diseases. This study aims to determine plasma MDA levels in the age group of 40 years and over and factors that can affect plasma MDA levels. This study was a cross-sectional study by including 42 subjects. Data collection was carried out using a questionnaire to obtain subject demographic data and physical activity, as well as physical examinations complemented by laboratory examinations and MDA levels. The mean age of the subjects was 52.7 years, and most of the 32 (76.2%) subjects were categorized as obese and had high activity. Pearson correlation test obtained body mass index statistically significantly correlated with plasma MDA levels with a value of  $p = 0.039$ , as well as physical activity statistically significantly correlated with a negative correlation, with a value of  $r = -0.26$  and  $p = 0.043$ . Conclusion Body mass index and activity are significantly correlated with plasma MDA levels.

**Keywords:** malondialdehyde, stress, oxidative, radicals

## INTRODUCTION

Aging is a progressive process caused by a physiological decline in body function. <sup>(1)</sup> A decrease in body homeostasis is accompanied by an increased risk of various chronic diseases including diabetes mellitus, cancer, neurodegenerative diseases, and heart disease. It is known that several factors can accelerate aging, namely age, lifestyle, oxidative stress, lipid peroxidation, and mitochondrial damage. Previous studies have shown that changes in sphingomyelin metabolism in the central nervous system and age are associated with neurodegenerative diseases. <sup>(2)</sup> In the US, aging is one of the most important factors in diabetes mellitus and heart disease. In 2015, an estimated 12% of the population had diabetes mellitus and an undetected 24% had diabetes mellitus. Heart disease is the highest contributor to mortality in the population aged 65-84 years. <sup>(3)</sup> Recent studies have shown that non-diabetic populations whose fasting glucose levels increase in the blood will increase the risk of coronary heart disease. But on the other hand, the data on fasting blood glucose increase is still debated in the population of cardiovascular disease patients who do not have diabetes mellitus. A recent study in North Sumatra showed a significant relationship between MDA levels in obese and non-obese populations. <sup>(4)</sup>

Free radicals are formed through two processes, namely exogenous and endogenous processes. Various pollutants such as cigarette smoke, motor vehicle fumes, industrial waste, and exposure to sunlight/ultraviolet will form free radicals. Illness, stress, and excessive exercise/physical activity are endogenous factors produced by the human body that can form free radicals. The formation of free radicals occurs due to oxidation processes and metabolic processes that take place during exercise or activity, inflammatory processes, consumption of certain foods, and others. Free radicals will begin to activate with molecules in the surrounding cells to obtain electrons to stabilize the condition of one's body. The electrons that have been taken in the cell molecules will trigger the

occurrence of free radicals. If this continues for a long time, it can cause oxidative stress which can cause inflammation, and DNA damage can also trigger the disease. Oxidative stress conditions characterized by increased production of Reactive Oxygen Species (ROS) have implications for various diseases such as hypertension, atherosclerosis, diabetes mellitus, stroke, chronic renal failure, heart failure, and other chronic diseases.<sup>(5-7)</sup>

Oxidative stress is an imbalance between oxidants (free radicals) and antioxidants. Free radicals that commonly cause oxidative damage are reactive oxygen species (ROS).<sup>(8)</sup> ROS compounds are formed through aerobic metabolism. These compounds can cause extensive damage through catalytic reactions. They are responsible for damage at the cellular level by oxidizing proteins, lipids, DNA (Deoxyribonucleic acid), and other biomolecules. Various compounds can be used as markers of oxidative stress such as malondialdehyde (MDA), and protein carbonylation. 4-hydroxy-2-nonenal (4HNE), and 8-hydroxy-deoxyguanosine (8OHdG).<sup>(9)</sup>

Malondialdehyde (MDA) is a result of lipid peroxidation in the body. The concentration of MDA indicates an oxidation process that occurs in the body's cell membranes due to metabolic processes. One of the factors that cause aging is the production of excess reactive oxygen species (ROS). ROS are generated through the process of cell respiration in the mitochondria and the phagocytosis process of foreign body elimination. To stabilize the amount of ROS, the body can use endogenous and exogenous antioxidants. Disruption of the balance between ROS and antioxidants can cause irreversible damage to cell organelle components up to the DNA level.<sup>(10,11)</sup>

MDA levels are strongly influenced by many factors such as age, illness, stress, various pollutants, food consumption, antioxidant supplements and excessive physical activity or exercise, and others. The other study found that more than 30% of subjects had high MDA levels, and concluded that there was a significant relationship between eating habits, body mass index, glucose levels with MDA levels, as well as smoking habits significantly related to MDA levels.<sup>(12)</sup> Recent studies have shown that plasma MDA levels in populations aged 70 years and 60 years are not significantly different but there is an increase in the specific activity of the catalase enzyme that increases with age.<sup>(10)</sup> In another study, there was a significant difference in plasma MDA levels in the obesity group without metabolic syndrome compared to the obesity group with metabolic syndrome.<sup>(4)</sup>

There are still variations in MDA data in various groups with different ages and different disease groups, so this study aims to determine plasma MDA levels in the age group 40 years and over and factors that can affect plasma MDA levels.

## **METHOD**

This observational analytic study with a cross-sectional design approach was conducted from November 2022 to February 2023 in Angke village, West Jakarta, Indonesia. The inclusion criteria for this study were: men and women aged  $\geq 40$  years, willing to participate in the study and sign informed consent. Meanwhile, the exclusion criteria were having a history of malignancy, kidney failure, a history of autoimmune disease, taking drugs related to psychiatric conditions, and having limitations in mobility. This study used the calculation of sample size with the formula  $N / 1 + N \cdot \alpha^2$ , and obtained the minimum number of samples required was 40 subjects. Demographic data and physical activity were obtained from distributed questionnaires. Blood chemistry examination (complete blood, glucose, total cholesterol, liver enzymes (SGOT and SGPT) using an enzymatic method. Examination of malondialdehyde levels was carried out using the spectrophotometric method. This research has obtained approval from the ethics committee of the Faculty of Medicine, Trisakti University, Number: 178/KER/FK/1X/2022.

## **RESULTS**

A total of 42 subjects participated in this study, most of the subjects were female 27 (64.3%), and the mean age of the subjects was 52.7 years. Anthropometric data of the subjects are as follows body weight  $67.53 \pm 13.64$  and height  $156.02 \pm 8.73$  cm, while for abdominal circumference the average is 93.09 cm. Most of the subjects 59.5% fell into the normotensive category, with an average systolic blood pressure of 141.62 mmHg and an average diastolic blood pressure of 82.45 mmHg. A total of 32 (76.2%) subjects were categorized as obese with BMI  $\geq 25.0$ , and most subjects had high activity levels. (Table 1)

Table 1. Distribution of Subject Characteristics (n=42)

Characteristics	n (%)	Mean ± SD
Gender		
Male	15 (35.7)	
Female	27 (64.3)	
Blood pressure (mmHg)		
Normal (< 140/90)	25 (59.5)	
Hypertension (≥ 140/90)	17 (40.5)	
Physical Activity		
Low (< 600 Mets)	11 (26.2)	
High (≥ 600 Mets)	31 (73.8)	
Body Mass Index (kg/m <sup>2</sup> )		27.69 ± 4.90
Not obese (< 25.0)	10 (23.8)	
Obese (≥ 25.0)	32 (76.2)	
Age (years)		52.7 ± 10.60
Body weight (kg)		67.53 ± 13.64
Height (cm)		156.02 ± 8.73
Abdominal circumference (cm)		93.09 ± 10.54
Blood pressure (mmHg)		
Systole		141.62 ± 20.77
Diastole		82.45 ± 12.70

The examination of the subject's plasma malondialdehyde levels obtained a mean of 0.77 with the highest value of 1.03 and the lowest of 0.51. Hemoglobin levels averaged 13.16 g/dL. Fasting blood glucose level was 95.06 ± 41.01 mg/dL and total cholesterol was 179.83 ± 31.85 mg/dL. Laboratory examinations to determine liver function were carried out by measuring SGPT and SGOT levels, obtaining mean values of 24.21 U/L and 26.31 U/L respectively. (Table 2)

Table 2. Laboratory examination results of subjects (n=42)

Variable	Mean ± SD
Hemoglobin (g/dL)	13.16 ± 1.56
Fasting blood glucose (mg/dL)	95.06 ± 41.01
Total cholesterol (mg/dL)	179.83 ± 31.85
SGPT (U/L)	24.21 ± 13.50
SGOT (U/L)	26.31 ± 11.51
Plasma malondialdehyde	0.77 ± 0.26

Determining the type of correlation test to be used by first conducting a data normality test. The normality test used was Shapiro-Wilk because the number of subjects was less than 50 subjects. The correlation test used to assess the correlation between blood glucose, total cholesterol, SGPT, SGOT levels, age, body mass index and physical activity with plasma MDA levels was the Pearson correlation test. The test results are presented in Table 3. Blood glucose, total cholesterol, SGPT, SGOT, and age did not have a significant correlation with MDA levels. Body mass index has a significant correlation with plasma MDA ( $p = 0.039$ ), with a positive correlation direction, and the strength of the correlation is statistically weak with a value of  $r = 0.396$  ( $< 0.4$ ). The physical activity variable has a significant correlation with plasma MDA levels ( $p = 0.043$ ) with a negative correlation direction but with a statistically weak correlation strength ( $r = -0.26$ ).

Table 3. Correlation Test of Age, Gender, Glucose, SGOT, SGPT, BMI, Physical Activity with Plasma MDA Levels (n=42)

	Plasma MDA Level	
	r	p
Age	0.098	0.756
Gender	0.124	0.108
Glucose	0.330	0.835
SGOT	0.104	0.513
SGPT	0.008	0.614
Body Mass Index	0.396	0.039*
Physical activity	0.260	0.043*

\* p < 0.05

### Discussion

The level of oxidative stress can be measured using Malondialdehyde (MDA) concentration. MDA is the result of lipid peroxidation, which will cause cell damage. Oxidant compounds such as free radicals or non-radical species attack lipids that contain carbon-carbon double bonds, especially unsaturated fatty acids (PUFA). MDA is one of the biomarkers widely used to assess oxidative stress. The mean serum MDA in this study 0.77 was found to be lower than the previous study<sup>(10)</sup>, this is because the age difference of the subjects in this study was younger than in the previous study. In contrast, a study conducted in Iraq on hypertensive patients concluded that the increase in MDA was associated with an increase in age, gender, and duration of disease.<sup>(13)</sup>

Free radicals such as ROS are the etiology of most chronic diseases, as it is known that active free radicals are the result of metabolic reactions. Damage to body tissue cells occurs because active free electrons remain looking for other electrons to settle, if no other electrons are available then active free electrons will attack cells resulting in cell damage. Although free radicals naturally have benefits for the body, when the level of free radicals exceeds the normal value, it will cause harm to the body.<sup>(11,12)</sup>

Measurement of oxidative stress can be done by measuring plasma malondialdehyde (MDA) levels. MDA is a secondary result of the lipid peroxidation process, producing various active compounds that will damage cells. The process of MDA formation through enzymatic and non-enzymatic processes. Currently, MDA is one of the biomarkers used to assess oxidative stress.<sup>(14)</sup> Some previous researchers concluded that higher serum MDA was found in patients with hyperlipidemia. The condition of increased fat content in the body is in line with increased oxidative stress. Hyperlipidemia is associated with physical changes in cell membranes that facilitate the release of free radicals from the mitochondrial electron transport chain or activation of NADPH oxidase. In patients with diabetes mellitus (DM) there is also an increase in MDA levels, malondialdehyde interacts irreversibly and reversibly with proteins and phospholipids to have a profound effect, especially on the collagen of the cardiovascular system. MDA levels remain within normal limits due to the antioxidant effect, but in DM, this protective effect is impaired resulting in increased MDA levels.<sup>(1,15)</sup>

Increased lipid peroxidation is considered a consequence of oxidative stress that occurs when the dynamic balance between prooxidants and antioxidants is disrupted. The statin class of drugs for hyperlipidemia therapy is proven to be anti-atherogenic, can improve endothelial function, inhibit vascular inflammation, and stabilize atherosclerotic plaques. In addition to statins, non-steroidal anti-inflammatory drugs (NSAIDs) can affect MDA concentration. Previous studies have found that NSAIDs have significant antioxidant properties to scavenge free radicals.<sup>(16,17)</sup>

In this study, there was no significant correlation between age and MDA levels. Similar results were also obtained by other researchers, cell damage due to oxidative stress is not related to increasing age but is associated with habits, behavior patterns, and physiological changes due to age. Various habitual patterns such as smoking, fatty food habits, lack of activity, and inflammatory processes can lead to increased oxidative stress. The habit of taking vitamin C (ascorbic acid) has a beneficial impact due to its antioxidant effect so cell damage and DNA damage are lower in this group.<sup>(18)</sup> MDA levels

5  
were significantly higher among all levels of smokers. MDA levels were lower in light smokers compared to heavy or moderate smokers. Higher vitamin C levels were found in the non-smoker group. All body tissues are susceptible to free radical damage, but based on location, the respiratory epithelium is particularly vulnerable to cigarette smoke. Oxygen radicals generated close to the cell membrane oxidize membrane phospholipids which will result in lipid peroxidation with the end products of this reaction being malondialdehyde, ethane, and pentane. This causes an increase in MDA levels in smokers.<sup>(19)</sup> Another study conducted in Klaten on foundry workers concluded that type of work, body mass index and smoking habits did not affect MDA levels, but length of work and marital status were associated with MDA levels. Long and continuous exposure to metallic materials will result in the accumulation of these materials in the body and will increase the oxidative process of polyunsaturated fats in cell membranes and worsen the level of lipid peroxidase, characterized by an increase in MDA levels as a reflection of the intensity and rate of lipid peroxidase reactions of damage to body cells.<sup>(20)</sup>

Hardiany's study also concluded that there was no significant difference in plasma MDA levels in the age population, but increasing age along with an increase in catalase-specific activity.<sup>(10)</sup> This increase in catalase-specific activity is a protective effect especially for elderly women to overcome high levels of oxidative damage. The role of endogenous antioxidant catalase is to suppress oxidative damage.<sup>(18,21)</sup> Oxidative stress plays an important role in the incidence of cardiovascular diseases including coronary artery disease. Oxidative stress is considered an important mechanism for the development of cardiovascular diseases especially atherosclerosis. Hypertension, cardiac hypertrophy, and ischemia-reperfusion injury are three other major cardiovascular conditions for which gender is a risk factor. Gender is also associated with differences in oxidative stress. Under physiological conditions, women appear to be less susceptible to oxidative stress. This may be due to the antioxidant properties of estrogen, sex differences in NADPH-oxidase activity or other mechanisms yet to be determined. Sex hormones including estrogen are globally present with their receptors in the heart so sex hormones can exert effects on the cardiovascular system. Estrogen can improve heart function and reduce injury to the heart. In addition, estrogen has antioxidant effects so that it can function as a cardioprotective.<sup>(22,23)</sup>

This study found a significant correlation between body mass index and MDA levels, the higher the BMI value, the higher the MDA levels. Similar results were also obtained in a study in India that concluded that in metabolic syndrome patients with one of the signs is an increase in BMI, there is an increase in MDA levels compared to the control group. In obese subjects, the level of MDA is higher than normal IMT. In a healthy body, there is redox hemostasis that occurs in cells, if an imbalance arises it will cause oxidative stress. Obese patients will have an increased metabolic load and mechanical load on the myocardium, increased body mass, and excessive food intake will cause the formation of lipid peroxidation, and free radicals.<sup>(24)</sup> The increase in some of these things will stimulate antioxidant enzymes in the long run, this will cause antioxidant enzyme reserves to be depleted and unable to overcome and balance the increase in oxidative stress.<sup>(25,26)</sup>

These types of training activities are exhausting, long exercise, overtraining syndrome, various types of exercise will induce a significant response to oxidative stress. Exercise activities such as light exercise, low-intensity exercise, and prolonged training will have an impact on improving endogenous antioxidant status. Reactive oxygen species play an important role in cell signaling and in regulating antioxidant gene expression. Exertion results in hyperregulation of nuclear factor kappa B and mitogen-activated protein kinases that activate gene expression of a number of enzymes and proteins with important roles in maintaining intracellular oxidative or antioxidant homeostasis.<sup>(1,6,27)</sup> The limitation of this study is that some factors that may affect MDA levels such as consumption of vitamin supplements, diet, and length of illness suffered were not examined in this study.

## Conclusions

Body mass index and activity were significantly correlated with MDA levels, while age, gender, blood glucose levels, and liver enzymes were not significantly correlated with MDA levels.



**Conflict of Interest:**

The investigators declare there is no conflict of interest in this study.

**References**

1. Sharifi-Rad M, Anil Kumar N V., Zucca P, Varoni EM, Dini L, Panzarini E, et al. Lifestyle, Oxidative Stress, and Antioxidants: Back and Forth in the Pathophysiology of Chronic Diseases. *Front Physiol.* 2020;11(July):1–21.
2. Park MH, Jin HK, Bae J sung. Potential therapeutic target for aging and age-related neurodegenerative diseases: the role of acid sphingomyelinase. *Exp Mol Med [Internet].* 2020;52(3):380–9. Available from: <http://dx.doi.org/10.1038/s12276-020-0399-8>
3. Całyniuk B, Grochowska-Niedworok E, Walkiewicz K, Kawecka S, Popiołek E, Fatyga E. Malondialdehyde (MDA) – product of lipid peroxidation as marker of homeostasis disorders and aging. *Ann Acad Medicae Silesiensis.* 2016;70:224–8.
4. Savira M, Rusdiana ., Widjaja SS, Syahputra M. Comparison Malondialdehyde (MDA) Level between Obesity Non Metabolic Syndrome and Obesity with Metabolic Syndrome Patients. 2020;(5):644–7.
5. Yekti R, Bukhari A, Jafar N, Thaha AR. Measurement of malondialdehyde (MDA) as a good indicator of lipid peroxidation. *Int J Allied Med Sci Clin [Internet].* 2018;6(4):6–8. Available from: [www.ijamsc.com](http://www.ijamsc.com)
6. Denise Mafra ME. Influence of Physical Exercise on Oxidative Stress and Inflammation in Hemodialysis Patients. *J Mol Biomark Diagn.* 2015;06(03).
7. Mao C, Yuan JQ, Lv Y Bin, Gao X, Yin ZX, Kraus VB, et al. Associations between superoxide dismutase, malondialdehyde and all-cause mortality in older adults: A community-based cohort study. *BMC Geriatr.* 2019;19(1):1–9.
8. Ayala A, Muñoz MF, Argüelles S. Lipid peroxidation: Production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. *Oxid Med Cell Longev.* 2014;2014.
9. Situmorang N, Zulham Z. Malondialdehyde (Mda) (Zat Oksidan Yang Mempercepat Proses Penuaan). *J Keperawatan Dan Fisioter.* 2020;2(2):117–23.
10. Hardiany NS, Sucitra S, Paramita R. Profile of malondialdehyde (MDA) and catalase specific activity in plasma of elderly woman. *Heal Sci J Indones.* 2020;10(2):132–6.
11. Hardiany NS, Sadikin M, Siregar N, Wanandi SI. The suppression of manganese superoxide dismutase decreased the survival of human glioblastoma multiforme T98G cells. *Med J Indones.* 2017;26(1):19–25.
12. 439-445\_nova.pdf.
13. A. Hassan A, Syyah SG. Oxidative Stress Marker Malondialdehyde and Glutathione Antioxidant in Hypertensive Patients. *Eur J Biomed Res.* 2023;2(1):31–6.
14. Khoubnasab Jafari M, Ansarin K, Jouyban A. Comments on “use of malondialdehyde as a biomarker for assessing oxidative stress in different disease pathologies: A review.” *Iran J Public Health.* 2015;44(5):714–5.
15. Shalash M, Badra M, Imbaby S, ElBanna E. Malondialdehyde in Type 2 Diabetics and Association With Cardiovascular Risk Factors. *J Med Res Inst.* 2020;41(2):21–30.
16. Oesterle A, Laufs U, Liao JK. Pleiotropic Effects of Statins on the Cardiovascular System. *Circ Res.* 2017;120(1):229–43.
17. Sekunda KN, Ariosta A, Limantoro C, Setiawan AA. Difference in Profiles of Oxidative Stress Marker (MDA) in STEMI and NSTEMI. *Diponegoro Med J (Jurnal Kedokt Diponegoro).* 2021;10(2):145–50.
18. Block G, Dietrich M, Norkus EP, Morrow JD, Hudes M, Caan B, et al. Factors associated with oxidative stress in human populations. *Am J Epidemiol.* 2002;156(3):274–85.
19. Kubihal C V., Naik HD. Effect of smoking on vitamin C and MDA: a cross sectional comparative study. *Int J Res Med Sci.* 2019;7(3):746.
20. Ari Suwondo, Ahmad Mohan, Melyana Nurul Widyawati M. Factors Related to Malondialdehyde (MDA) Levels in Metal Casting Industry Workers. Preprints.

- 2020;1(October):1–8.
21. Schöttker B, Brenner H, Jansen EHJM, Gardiner J, Peasey A, Kubínová R, et al. Evidence for the free radical/oxidative stress theory of ageing from the CHANCES consortium: A meta-analysis of individual participant data. *BMC Med.* 2015;13(1):1–15.
  22. Kander MC, Cui Y, Liu Z. Gender difference in oxidative stress: a new look at the mechanisms for cardiovascular diseases. *J Cell Mol Med.* 2017;21(5):1024–32.
  23. Bell JR, Bernasochi GB, Varma U, Raaijmakers AJA, Delbridge LMD. Sex and sex hormones in cardiac stress - Mechanistic insights. *J Steroid Biochem Mol Biol.* 2013;137:124–35.
  24. P G, Curcio F, Bulli G, Aran L, Della-morte D, Testa G, et al. Oxidative Stress and Diseases. *Oxidative Stress Dis.* 2012;757–72.
  25. Sabitha K, Venugopal B, Rafi M, V Ramana K. Role of Antioxidant Enzymes in Glucose and Lipid Metabolism in Association with Obesity and Type 2 Diabetes. *Am J Med Sci Med.* 2014;2(1):21–4.
  26. Manna P, Jain SK. Obesity, Oxidative Stress, Adipose Tissue Dysfunction, and the Associated Health Risks: Causes and Therapeutic Strategies. *Metab Syndr Relat Disord.* 2015;13(10):423–44.
  27. Lewis NA, Howatson G, Morton K, Hill J, Pedlar CR. Alterations in Redox Homeostasis in the Elite Endurance Athlete. *Sport Med.* 2015;45(3):379–409.

# Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years

## ORIGINALITY REPORT

18%

SIMILARITY INDEX

17%

INTERNET SOURCES

13%

PUBLICATIONS

8%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="https://onlinelibrary.wiley.com">onlinelibrary.wiley.com</a> Internet Source	3%
2	<a href="http://ejournal3.undip.ac.id">ejournal3.undip.ac.id</a> Internet Source	3%
3	<a href="http://jddtonline.info">jddtonline.info</a> Internet Source	2%
4	<a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a> Internet Source	2%
5	<a href="http://innovpub.org">innovpub.org</a> Internet Source	1%
6	<a href="http://www.researchgate.net">www.researchgate.net</a> Internet Source	1%
7	M. Shalash, M. Badra, S. Imbaby, E. ElBanna. "MALONDIALDEHYDE IN TYPE 2 DIABETICS AND ASSOCIATION WITH CARDIOVASCULAR RISK FACTORS", Journal of the Medical Research Institute, 2020 Publication	1%

8	Submitted to Universitas Airlangga Student Paper	1 %
9	Submitted to Trisakti University Student Paper	1 %
10	<a href="https://repository.stikim.ac.id">repository.stikim.ac.id</a> Internet Source	1 %
11	<a href="https://mdpi-res.com">mdpi-res.com</a> Internet Source	1 %
12	<a href="https://www.pubmedcentral.nih.gov">www.pubmedcentral.nih.gov</a> Internet Source	1 %
13	Anna Thomson, Ellen Christensen, Maria Wiklund, Monica Christianson. "A safe place – Adolescents' and young adults' perceptions of youth clinics in northern Sweden", <i>Sexual &amp; Reproductive Healthcare</i> , 2022 Publication	1 %
14	<a href="https://www.researchsquare.com">www.researchsquare.com</a> Internet Source	1 %
15	Warisya Miftah Amanda, Chandra Isabella Hostanida Purba, Ristina Mirwanti. "DIFFERENCE OF BOWEL SOUND RETURN TIME AMONG POST-LAPARATOMY SURGERY PATIENTS AFTER CHEWING GUM", <i>Belitung Nursing Journal</i> , 2019 Publication	<1 %

---

Exclude quotes Off

Exclude matches < 15 words

Exclude bibliography On

# Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years

---

GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---