

**AIMS Public Health**

ISSN 2327-8994

[Contact by email](#)

AIMS Public Health is Open Access and an international quarterly publication devoted to publishing peer-reviewed, high quality, original papers in the field of public health. We publish the following article types: original research articles, reviews, editorials, letters, and conference reports.

All published papers will be indexed in Web of Science (ESCI), Scopus and PMC.

Impact Factor
2.7**Citescore**
5.3[Advanced Search](#)

Author/Keywords/DOI

[Journal Home](#)[About **▼**](#)[Contribute **▼**](#)[Articles **▼**](#)[Submit a Paper](#)[Reviewer Login](#)[Email Notifications](#)**Special Issues Open For Submission**

International Evidence from Research and Practice to Foster Healthy Lifestyles and Mental Health in Adolescence and Emerging Adulthood

Guest Editors: Diego Gomez-Baya, Margarida Gaspar de Matos

[Journal Home](#)[About **▼**](#)[Contribute **▼**](#)[Articles **▼**](#)**AIMS Public Health**

Keith Elder (Founding Editor in Chief)



President, Saint Xavier University

Hui Wang (Co-Editor in Chief)



School of Public Health, Shanghai Jiao Tong University School of Medicine, Shanghai, China

Dahai Zhao (Associate Editor in Chief)



School of International and Public Affairs, Shanghai Jiao Tong University, Shanghai, 200025, China

Shixue Li (Associate Editor in Chief)



Dean and Professor School of Public Health, Cheeloo College of Medicine, Shandong University, Jinan, China

Stefano Rizza (Associate Editor in Chief)



System Medicine Department, University of Rome Tor Vergata, Via Montpellier 1, 00133, Rome

Wenjun Li (Associate Editor in Chief)



Professor of Public Health, Director of Center for Health Statistics, Zuckerberg School of Health Sciences, University of Massachusetts Lowell, 61 Wilder St, O'Leary 540U, Lowell, MA 01854, USA

Io Hong Cheong (Associate Editor in Chief)



School of Public Health, Shanghai Jiao Tong University School of Medicine, Shanghai, China

Shengjiao Pang (Managing Editor)



AIMS Press, Anli Road, Chaoyang District, Beijing 100101, China

**AIMS Public Health****Impact Factor**
2.7**Citescore**
5.3[Submit a Paper](#)[Reviewer Login](#)[Email Alert](#)

Editorial Board

Alessandro Leite Cavalcanti	✉ Department of Dentistry, State University of Paraiba, Brazil.
Andreas Sönnichsen	✉ Department of General Practice and Family Medicine, Center for Public Health, Medical University of Vienna, Austria
Behdin Nowrouzi-Kia	✉ Department of Occupational Science and Occupational Therapy, University of Toronto
Bo Chen	✉ Department of Nutrition and Food Hygiene, School of Public Health, Fudan University, Shanghai, China
Carla M. A. Pinto	✉ School of Engineering, Polytechnic of Porto, Portugal
Carmine Finelli	✉ Department of Internal Medicine, ASL Napoli 3 Sud, Torre del Greco (Naples), Italy and Professor of Internal Medicine at the Federico II University Medical School in Naples, Nursing School, Learning Centre of Nola (Naples)
Clemens Drenowatz	✉ Division of Sport, Physical Activity and Health; University of Education Upper Austria, Austria
Daglia Maria	✉ Full Professor of Food Chemistry at the Department of Pharmacy, University of Naples Federico II, Via Domenico Montesano 49, 80131 Naples – Italy
Donald B. Giddon	✉ Faculty of Medicine, Harvard University, 188 Longwood Ave, Boston, MA 02115, USA
Emanuele Amadio	✉ U.O. Servizio Epidemiologia e Programmazione Dipartimento PAC, Agenzia di Tutela della Salute (ATS) della Brianza, Corso Carlo Alberto 120, Lecco, Italy
Fradelos C. Evangelos	✉ Department of Nursing, University of Thessaly, Greece



Galanis Petros	✉ Department of Nursing, National & Kapodistrian University of Athens
Gang Xu	✉ Department of Community Health and Behavioral Medicine, School of Public Health, Shanghai Jiaotong University, Shanghai, China
Gareth Morgan	✉ Department of Planning and Primary Care, Hywel Dda Health Board, Heol Cropin, Llanelli, Carmarthenshire SA14 8QW, NHS Wales, Wales, UK
Giuseppe Murdaca	✉ Di.M.I.-University of Genova Viale Benedetto XV, n. 6-16132 Genova, Italy
Giustino Varrassi	✉ Paolo Procacci Foundation, Former Professor of Anesthesiology and Pain Medicine, University of L'Aquila, 67100 L'Aquila, Italy
Haijiang Lin	✉ Taizhou City Center for Disease Control and Prevention, Zhejiang, China
Haslyn E. R. Hunte	✉ School of Public Health, Robert C Byrd Health Sciences Center, West Virginia University, PO Box 9190, Morgantown, WV 26506-9190, USA
Hell Markus	✉ Head of division of Medical Microbiology, Institute of Laboratory Medicine – Paracelsus Medical University, Salzburg, University Hospital, Salzburg, Austria
Hilary Karasz	✉ Public Health – Seattle & King County, 401 5th Avenue, Suite 1300, Seattle, WA 98104, USA
Hsin-Hung Wu	✉ Department of Business Administration, College of Management, National Changhua University of Education, Changhua, Taiwan, China
Huabin Luo	✉ Department of Public Health, Brody School of Medicine, East Carolina University, Greenville, NC 27834, USA
Hualiang Lin	✉ School of Public Health, Sun Yat-sen University, Guangzhou, China



I-Shiang Tzeng	1. Department of Research, Taipei Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation, New Taipei City, Taiwan, China; 2. Department of Statistic, National Taipei University, Taipei, Taiwan, China; 3. Department of Applied Mathematics; Department of Exercise and Health Promotion, Chinese Culture University, Taipei, Taiwan, China
James A. Swartz	Jane Addams College of Social Work, University of Illinois at Chicago, 1040 W. Harrison Street, MC 309 Chicago, IL 60607, USA
Jangho Yoon	School of Social and Behavioral Health Sciences, College of Public Health and Human Sciences, Oregon State University, Corvallis, Oregon, USA
Jinhui Liu	Department of Gynecology, The First Affiliated Hospital of Nanjing Medical University, Nanjing, China
Junfeng Jiao	School of Architecture, The University of Texas at Austin, Austin, TX, 78759, USA
Liang Zhou	School of Public Health, Shanghai Jiao Tong University School of Medicine, Shanghai, China
Liping Li	School of Public Health, Shantou University, Shantou 515041, China
Luca Testarelli	Department of Maxillo and Oro-Facial Sciences, University of Rome La Sapienza, Rome, Italy
Maciej Henneberg	Emeritus Wood Jones Professor of Anthropological and Comparative Anatomy, School of Biomedicine, The University of Adelaide, Australia
Mark Menniti Stecker	Fresno Institute of Neuroscience, 1558 East Shadow Creek Drive Fresno, CA 93730, USA
Marco Tatullo	University of Bari-Italy
María Auxiliadora Robles-Bello	Department of Psychology, Universidad de Jaén, Jaén, Spain



Mario Coccia	National Research Council of Italy, Italy
Martha Slay Wingate	Martha Slay Wingate, University of Alabama at Birmingham, 1665 University Boulevard, Birmingham, Alabama, USA
Martin Burtscher	University of Innsbruck, Dept. of Sport Science, Innsbruck Austria
Michelle G. Dresser	Bureau of Chronic Disease Prevention & Tobacco Control, NYC Department of Health and Mental Hygiene, Gotham Center, CN-18 42-09 28th Street, 9th Floor, Queens, NY 11101, USA
Miguel Viñas Ciordia	Dept of Pathology&Experimental Therapeutics Medical School, University of Barcelona, Catalonia, Spain
Nicola Magnavita	Postgraduate School of Occupational Health, Università Cattolica del Sacro Cuore, Largo F. Vito 1, 00168 Roma, Italy
Quan Zheng	Center for Single-Cell Omics, School of Public Health, Shanghai Jiao Tong University School of Medicine, Shanghai, China
Roy G. Beran	Western Sydney University, School of Medicine, 80 Goldsmith Avenue, Campbelltown, Sydney, Australia
Russell Kabir	School of Allied Health, Faculty of Health, Education, Medicine & Social Care, Anglia Ruskin University, Bishop Hall Lane Chelmsford, UK
Samuel Asumadu Sarkodie	Nord University, Norway
Sandra Racionero-Plaza	Department of Psychology, University of Barcelona, Barcelona, Spain
Shahab S. Band	Department of IAI, National Yunlin University Science and Technology, Taiwan
Shariful Islam	Institute for Physical Activity and Nutrition, School of Exercise and Nutrition Sciences, Faculty of Health, Deakin University, Australia



Shervin Assari	 	Department of Urban Public Health, Charles R Drew University of Medicine and Science, Los Angeles, CA, USA
Sónia F. Dias	 	Instituto de Higiene e Medicina Tropical, Universidade Nova de Lisboa, Lisbon, Portugal
Stefano Campostrini	 	Department of Statistics, University Ca Foscari of Venice, Venice, 30121, Italy
S. R. Pandi-Perumal	 	Somnogen Canada Inc., College Street, Toronto, ON M6H 1C5, Canada
Vijay Kumar Chattu	 	Department of Medicine, Temerty Faculty of Medicine, University of Toronto, Toronto- Canada
Wendy Cozen	 	Department of Preventive Medicine, USC/Norris Comprehensive Cancer Center, 1441 Eastlake Ave. MC 9175, Los Angeles, California, USA
Xiaojun Sun	 	School of Psychology, Central China Normal University, Wuhan, China
Yang Yang	 	Department of Food Safety and Toxicology, School of Public Health, Shanghai Jiaotong University, Shanghai, China
Yueliang Zhao	 	School of Public Health, Shanghai Jiao Tong University School of Medicine, Shanghai, China
Zhuohui Zhao	 	Department of Environmental Health, School of Public Health, Fudan University, Shanghai, China

www.aimspress.com/aimsph/article/2025/4/archive-articles

Home Journals 

Journal Home About Contribute Articles

2025 Volume 12, Issue 4

[Research article](#) | [Special issue](#)
 psychological first aid in a cross-border medical emergency drill: Detected barriers, perceptions, and opportunities for improvement
 Olga Mata, Andrés Cuartero, Laia Reñés, Judith Cattivelli, Yurt Lázaro
 2025, Volume 12, Issue 4: 999-1004, doi: 10.3934/publichealth.2025049
 [Abstract](#)  

[Research article](#) | [Special issue](#)
 Associations between modifiable risk factors and limitation in activities of daily living among individuals aged ≥ 45 years: Evidence from the China health and retirement longitudinal study (CHARLS)
 Yanheng Li, Jian Gao, Wenzhu Song, Xiaodan Liang, Junhao He, Fuliang Yi, Wenhao Song, Dongliang Yin
 2025, Volume 12, Issue 4: 1005-1025, doi: 10.3934/publichealth.2025050
 [Abstract](#)  

[Research article](#)
 Comparison of physical fitness in youth with post-COVID-19: A study of individuals with and without symptoms
 Parshwarya Amgut, Simina Wongphon
 2025, Volume 12, Issue 4: 1026-1034, doi: 10.3934/publichealth.2025051
 [Abstract](#)  

[Research article](#) | [Special issue](#)
 Exploring digital interaction and positive youth development in a sample of Spanish undergraduates
 Esther López-Bermúdez, Gina Tomé, Diego Gómez-Baya
 2025, Volume 12, Issue 4: 1035-1054, doi: 10.3934/publichealth.2025052
 [Abstract](#)  

[Research article](#) | [Special issue](#)
 Significant risk of developing asbestos-related diseases in Japan's industries: An analysis of workers' compensation
 Lea Hess-Indrayati, Masanobu Etoku, Naoto Arai, J.-P. Taro Tamura, Narufumi Suganuma
 2025, Volume 12, Issue 4: 1055-1068, doi: 10.3934/publichealth.2025053
 [Abstract](#)  

[Research article](#) | [Special issue](#)
 Association between nursing work environment, burnout, and turnover intention: A cross-sectional study in Greece
 Ioanna Mousouli, Aglaia Kotsopoulos, Evangelos C. Fradelos, Ioanna V. Papathanasiou, Theodoros Parasikas, Ioanna Prasi, Maria Katsikas, Alkis Kotsopoulos, Maria Chorti, Petros Galanis
 2025, Volume 12, Issue 4: 1069-1083, doi: 10.3934/publichealth.2025054
 [Abstract](#)  

The association between $PM_{2.5}$ level and respiratory tract infections among children: A cross-sectional study
 Hani Krimanou, Pumaravadee Tjhin
 2025, Volume 12, Issue 4: 1084-1114, doi: 10.3934/publichealth.2025055
 [Abstract](#)  

[About AIMS Press](#) | [Open Access Policy](#) | [Contact us](#)
 Copyright © AIMS Press

www.scopus.com/sources.uri

Title

Title: AIMS Public Health

CiteScore 2024 has been released. [View CiteScore methodology >](#)

Filter refine list [Clear filters](#)

1 result [Learn more about Scopus Source List](#)

All

View metrics for year:

Display options

Source title ↓	CiteScore ↓	Highest percentile ↓	Citations 2021-24 ↓	Documents 2021-24 ↓	% Cited ↓
1 AIMS Public Health Open Access	5.3	75% 169/687 Public Health, Environmental and Occupational Health	1.104	207	68

Research article

The association between PM_{2.5} level and respiratory tract infections among children: A cross-sectional study

Hari Krismanuel* and Purnamawati Tjhin


Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

* Correspondence: Email: hari_krismanuel@trisakti.ac.id.

Abstract: *Background:* PM_{2.5} is a key air pollutant that contributes to respiratory morbidity, especially in children. In Jakarta, Indonesia, PM_{2.5} levels often exceed safe thresholds. This study contributes local evidence from Indonesia, where research on the health effects of PM_{2.5} in children remains limited. To address this gap in the existing literature, particularly within the Indonesian context, this study offers novel insights by specifically investigating the association between ambient PM_{2.5} exposure and respiratory tract infections (RTIs) in school-aged children and further exploring this association within male and female subgroups, an aspect that has received limited attention in this setting. *Objective:* This study aims to assess the association between ambient PM_{2.5} exposure and RTIs in school-aged children, and to explore this association within male and female subgroups. *Methods:* This cross-sectional study was conducted among 107 children aged 6–12 years from two elementary schools: one in Jakarta (high PM_{2.5} exposure) and one in Bandung (low PM_{2.5} exposure). Data on PM_{2.5} levels were obtained from local air quality monitoring. RTI symptoms were assessed through structured interviews and physical examination. Participants were selected using random sampling. Chi-square tests and effect size calculations (phi coefficient) were used to compare groups. Potential confounders such as age, gender, and household smoke exposure were minimized through inclusion/exclusion criteria and the selection of demographically and environmentally similar school communities. Multiple binary logistic regression adjusting for confounders was also performed to assess the independent association between PM_{2.5} exposure and RTIs. *Results:* The Chi-square test indicated a significant association between PM_{2.5} levels and the occurrence of RTI ($\chi^2 = 22.154$, df = 1, $p < 0.001$, $\phi = 0.475$). Given the potential low expected counts in some cells, the statistical significance was further evaluated using Fisher's Exact Test, which also showed a significant association ($p < 0.001$). The prevalence of RTI was significantly higher in the high exposure group (71.43%) compared to the low exposure group (25.86%) ($p < 0.001$). Further analysis did not reveal

significant differences in the proportion of each age group between the high and low PM_{2.5} exposure groups [$\chi^2(1) = 0.093$, $p = 0.761$]. Similarly, no significant differences were found in the proportion of gender between the high and low PM_{2.5} exposure groups [$\chi^2(1) = 1.611$, $p = 0.204$] in the total sample. Likewise, there were no significant differences in the proportion of RTI across different age groups [$\chi^2(6) = 5.327$, $p = 0.503$] or between genders [$\chi^2(1) = 0.008$, $p = 0.928$] in the total sample. However, further analysis examining the association between PM_{2.5} exposure and RTI within gender subgroups revealed a significant association in both male [$\chi^2(1) = 10.873$, $p = 0.001$] and female [$\chi^2(1) = 11.755$, $p = 0.001$] children. The estimated prevalence ratio (PR) was 2.76 (95% CI: 1.68–4.54), indicating that children in the high PM_{2.5} exposure area had approximately 2.76 times higher prevalence of RTI compared to those in the low exposure area. The absolute prevalence difference (PD) was 45.57% (95% CI: 25.9%–65.2%). Binary logistic regression analysis showed that children in the high PM_{2.5} exposure group had significantly higher odds of having RTI (OR = 7.167, 95% CI: 3.050–16.837, $p < 0.001$). Further analysis examining the association between maternal socioeconomic factors and both PM_{2.5} exposure and RTI occurrence revealed no statistically significant relationships. Chi-square tests showed no significant association between maternal education level (low vs. medium) and PM_{2.5} exposure group [$\chi^2(1) = 0.045$, $p = 0.833$], nor between maternal occupation (blue collar vs. semi-professional) and PM_{2.5} exposure group [$\chi^2(1) = 0.006$, $p = 0.937$]. Similarly, no significant associations were found between maternal education level and RTI [$\chi^2(1) = 0.233$, $p = 0.629$] or between maternal occupation and RTI [$\chi^2(1) = 0.447$, $p = 0.504$]. Crucially, after adjusting for potential confounders including gender, age, maternal education, and maternal occupation in a multivariate logistic regression model, the odds of having RTI remained significantly higher in children with high PM_{2.5} exposure (adjusted OR = 7.883, 95% CI: 3.228–19.250, $p < 0.001$). *Conclusions:* Children exposed to higher levels of PM_{2.5} had significantly more respiratory tract infections. These findings highlight the need for targeted public health interventions in polluted urban areas. This study is among the first to quantify this association in the Indonesian context and provides a newly developed and validated instrument (RAAEC-C instrument) for future research. These findings should be interpreted as preliminary evidence and require replication in future longitudinal studies before firm conclusions can be drawn. Further research using longitudinal designs is needed to understand the long-term impacts of PM_{2.5} exposure on children's respiratory health and to inform appropriate mitigation strategies.

Keywords: air pollution; children; cross-sectional study; PM_{2.5}; respiratory tract infection



Graphical abstract

This graphical abstract illustrates the conceptual pathway from air pollution exposure to pulmonary inflammation and respiratory tract infections.

1. Introduction

Fine particulate matter with a diameter of $2.5 \mu\text{m}$ or less ($\text{PM}_{2.5}$) has been widely recognized as one of the most harmful air pollutants due to its ability to penetrate deep into the respiratory tract and enter the bloodstream [1–3]. Exposure to $\text{PM}_{2.5}$ is associated with a range of adverse respiratory health outcomes, including airway inflammation, decreased lung function, and increased risk of respiratory tract infections and chronic respiratory diseases [1–5]. Research has shown that prolonged exposure to $\text{PM}_{2.5}$ can lead to chronic respiratory conditions such as asthma and bronchitis, particularly in children whose immune systems are still developing. Children are particularly vulnerable to air pollution because their lungs and immune systems are still developing, they have higher minute ventilation per body weight, and they often spend more time outdoors than adults [6–9].

In Jakarta, the capital of Indonesia, the burden of air pollution is especially concerning. As one of the most densely populated and industrialized cities in Southeast Asia, Jakarta frequently experiences $\text{PM}_{2.5}$ concentrations far exceeding the World Health Organization (WHO) air quality guideline. In August 2023, Jakarta was named the most polluted city in the world, with average $\text{PM}_{2.5}$ concentrations exceeding safe limits by more than five times, reaching up to $80 \mu\text{g}/\text{m}^3$ on the worst days. In response, local authorities implemented remote work policies for civil servants to curb vehicle emissions and improve air quality [10–16]. Prolonged exposure in such environments poses significant health risks, particularly for children [6–9]. Furthermore, available data suggest a concerning trend of elevated $\text{PM}_{2.5}$ levels in Jakarta over the past few years, primarily stemming from sources such as vehicle emissions, industrial activities, and biomass burning.

Research has demonstrated the adverse health effects of $\text{PM}_{2.5}$ on children's respiratory health in various parts of the world. For example, Sugiyama et al. studied the health effects of $\text{PM}_{2.5}$ sources on children's allergic and respiratory symptoms in Fukuoka, Japan, during the spring of 2014–2015 [17]. Another study by Xu et al. examined the acute effects of ambient $\text{PM}_{2.5}$ on lung function among school

children in Zhejiang Province, China, over the years 2014–2017 [18]. Ma et al. evaluated the association of air pollution with children's outpatient visits for respiratory diseases in the ex-heavily polluted Northwestern city, China, from 2014 to 2016 [19]. Additionally, a comprehensive literature review by Xing et al. explored the impact of PM_{2.5} on the human respiratory system, providing an in-depth analysis of epidemiological, experimental, and mechanistic studies [1]. More recently, Adhikary et al. conducted a large-scale study using PM_{2.5} data from the Atmospheric Composition Analysis Group at Washington University to assess the association between PM_{2.5} exposure and acute respiratory infections (ARI) in 223,375 children from the 2019–2021 Demographic Health Survey in India, contributing further evidence on the relationship between PM_{2.5} exposure and respiratory health outcomes in children [7].

Despite the growing body of research globally, there is a lack of studies focused on the impact of PM_{2.5} on children's respiratory health in Indonesia, particularly in urban areas like Jakarta. While studies have explored the health effects of PM_{2.5} in countries such as China, Japan, and India, evidence from Indonesia remains limited [1,7,17,18]. Research is needed to understand how children in Indonesian urban environments are affected by exposure to PM_{2.5}, especially considering the unique genetic, environmental, and social contexts of the region. Furthermore, genetic and racial variations may influence susceptibility to PM_{2.5}-induced respiratory effects, which makes findings from other countries less generalizable to Indonesia [20–22]. Understanding these localized impacts is crucial for developing effective strategies to mitigate the adverse health consequences of air pollution and improve respiratory health outcomes for children in Jakarta.

Given Jakarta's alarming pollution levels and the limited availability of localized research, it is crucial to fill this gap in knowledge to better inform public health decisions. This study aims to compare respiratory health outcomes—specifically respiratory tract infections—among children exposed to high levels of PM_{2.5} in Jakarta and those living in a region with significantly lower PM_{2.5} exposure, Bandung. By comparing these two distinct regions, we aim to provide insights into the local impact of air pollution on children's respiratory health in Indonesia. Furthermore, to explore potential **variations in this association**, we also examined the association between PM_{2.5} exposure and respiratory tract infections within male and female subgroups of the study population.

This study is novel in two main ways. First, it is one of the first to empirically examine the respiratory impact of PM_{2.5} on school-age children in Indonesia, filling a crucial research gap in the Southeast Asian context. Second, it compares two distinct exposure zones—Jakarta (high PM_{2.5}) and a region with significantly lower PM_{2.5} levels—allowing for a clearer understanding of the exposure-response relationship. The findings are expected to inform public health interventions, raise awareness about urban air pollution, and support evidence-based policymaking in Indonesia.

To our knowledge, this is the first study to examine the association between PM_{2.5} exposure and respiratory problems among school-aged children in Indonesia, with a specific focus on Jakarta. By comparing children from a high-exposure area (Jakarta) to those from a region with better air quality (Bandung), this study provides localized evidence on the impact of PM_{2.5} on symptoms such as coughing, wheezing, and shortness of breath.

2. Materials and methods

2.1. Research questions

What is the impact of naturally occurring differences in PM_{2.5} levels on respiratory health outcomes among children aged 6–12 years living in areas with high air pollution exposure compared to those in areas with better air quality?

2.2. Study design

This study employed an **analytical cross-sectional design** to evaluate the association between exposure to PM_{2.5} and the prevalence of respiratory tract infections (RTIs) in children. The population was divided into two comparison groups based on PM_{2.5} exposure levels in their residential and school environments:

- **High-exposure group:** Children living in areas with consistently high PM_{2.5} levels (poor AQI).
- **Low-exposure group:** Children living in areas with low PM_{2.5} levels (good AQI).

Unlike ecological studies that rely on group-level data, this study collected individual-level data on exposure and health outcomes, allowing for more precise estimation of associations.

The study population consisted of elementary school children aged 6–12 years attending schools located in areas with different levels of PM_{2.5}.

To minimize potential confounding factors, we selected schools from communities that were as **homogeneous as possible** in terms of environmental, demographic, behavioral, and household factors (e.g., exposure to cigarette smoke, ventilation, immunization status, nutritional status, seasonality, and housing conditions). Controlled variables included:

- **Demographic factors** (age, gender, socioeconomic status).
- **Environmental factors** (exposure to cigarette smoke, population density, seasonal and weather variation).
- **Health-related factors** (immunization status, nutritional status, history of tuberculosis, congenital heart disease, cohabitation with a household member undergoing TB treatment).
- **Behavioral factors** (personal hygiene, sanitation practices, social habits).
- **Household factors** (ventilation, indoor air quality).

Homogeneity across these variables was ensured during the selection process to reduce bias and improve internal validity.

2.3. Study population and sampling

The study population consisted of **elementary school children aged 6–12 years** enrolled in selected schools located in areas with high and low PM_{2.5} exposure. From each exposure group (high and low PM_{2.5}), one elementary school was selected. Within each selected school, a target sample of 60 students was initially recruited using stratified random sampling. Specifically, 10 students were randomly chosen from each grade level (Grade 1 to Grade 6) to ensure representation across all school years within each school.

Following the initial recruitment of 60 students from each school (totaling 120 students), participants were screened according to predetermined inclusion and exclusion criteria (see section

“Eligibility Criteria”). This resulted in the exclusion of 11 students from the high PM_{2.5} exposure area school (Jakarta) and 2 students from the low PM_{2.5} exposure area school (Bandung). Consequently, the final sample size for this study comprised 107 children (49 from the high PM_{2.5} exposure area and 58 from the low PM_{2.5} exposure area).

2.4. Eligibility criteria

Inclusion criteria:

- Children aged 6–12 years.
- Enrolled in selected elementary schools located in areas with high or low PM_{2.5} exposure.
- Obtained written parental/guardian consent and gave verbal assent to participate.
- Present at school on the day of data collection.

Exclusion criteria:

- Children with poor nutritional status (as determined by anthropometric measurements, including height and weight to calculate BMI).
- History of congenital heart disease.
- History of pulmonary TB treatment.
- Household member currently undergoing TB treatment.
- Current smoker or exposed to active smoking at home.
- Refusal by the child or parent/ guardian to participate or to continue participation in the research.

2.5. Study setting and procedure

The study was conducted in two urban areas in Indonesia with contrasting levels of particulate matter (PM_{2.5}) exposure: Jakarta, characterized by high PM_{2.5} levels, and Bandung Regency, known for its relatively lower PM_{2.5} levels, based on mapping data from local air quality monitoring stations. Within these urban areas, two elementary schools were strategically selected: SD Kedoya (Kedoya Elementary School) in West Jakarta, representing the high PM_{2.5} exposure site, and SD Pangalengan (Pangalengan Elementary School) in Pangalengan, Bandung Regency, representing the lower PM_{2.5} exposure site.

Data collection at SD Kedoya (Kedoya Elementary School), situated in West Jakarta, Jakarta, was carried out on May 6, 2024. Subsequently, data collection at SD Pangalengan (Pangalengan Elementary School), located in Pangalengan, Bandung Regency, West Java, took place on May 11, 2024. The data collection period across both sites spanned five days.

Data collection was carried out in several stages:

1. **Random selection of schools** in each city and eligible students aged 6–12 years from class rosters.
2. **Informed consent** obtained from parents or guardians, and verbal assent obtained from the children.
3. **Face-to-face interviews** with parents or guardians to collect data on symptoms of RTIs, including cough, wheezing, shortness of breath, and exacerbation of asthma.
4. **Application of exclusion criteria**, including screening for:
 - Poor nutritional status;
 - Congenital heart disease;
 - Current or past TB treatment;
 - Household member with active TB;

- Smoking behavior;
- Refusal to participate or continue in the study.

5. **Determination of nutritional status** by measurement and anthropometry

6. **Clinical examination** of each child's oral cavity, pharynx, nasal passages, and chest was conducted to support the diagnosis of RTIs.

2.6. *Stages of research implementation*

The implementation of this study involved several structured steps. First, air quality mapping was conducted in three locations within Jakarta and Bandung to identify areas with the highest and lowest PM_{2.5} levels. Based on this mapping, Kedoya Elementary School in Jakarta and Pangalengan Elementary School in Bandung were selected as the study sites.

Following school selection, students aged 6–12 years were randomly selected from class rosters across grades 1 to 6. Informed consent from parents and verbal assent from children were obtained prior to participation. Each child underwent an interview and a brief clinical examination to assess symptoms of respiratory tract infection, such as coughing, wheezing, and shortness of breath.

Children who met exclusion criteria—such as poor nutritional status, history of chronic illnesses, tuberculosis exposure, or parental refusal—were excluded from the study. The final dataset was then subjected to statistical analysis to examine differences in RTI prevalence between the two exposure groups.

The research was conducted according to standard procedures for cross-sectional observational studies. Throughout the implementation, we adhered to the STROBE (Strengthening the Reporting of Observational Studies in Epidemiology) guidelines to ensure transparent reporting [23,24]. The completed STROBE checklist is provided in Table S1 of the Supplementary Materials.

2.7. *Sample size determination*

The sample size for a cross-sectional study depends on several factors, including the desired level of confidence, the margin of error, the expected prevalence of the health effects, and data variability. There are standard formulas to calculate sample size, such as the one for estimating a population proportion (used for binary outcomes like the presence or absence of health effects). The formula is as follows [25–27]:

$$n = \frac{(z_{\alpha/2} + z_{\beta})^2 \cdot [P_1 \cdot (1 - P_1) + P_0 \cdot (1 - P_0)]}{(P_1 - P_0)^2} \quad (1)$$

Where:

- $Z_{\alpha/2}$ is the Z-score corresponding to the desired significance level ($\alpha/2$).
- Z_{β} is the Z-score corresponding to the desired power ($1 - \beta$).
- P_{avg} is the average of the expected proportions (P_0 and P_1).
- P_0 is the expected proportion in the unexposed group.
- P_1 is the expected proportion in the exposed group.
- R is the ratio of unexposed to exposed participants.

Using this formula with the assumed proportions, a significance level of 1%, and a power of 95%, the required sample size was determined to be 34 participants per group. This calculation was based on an assumed prevalence difference of approximately 40%, derived from previous studies in similar

urban pediatric populations. To strengthen the study's statistical power and account for potential non-response, data loss, or misclassification, we increased the sample size beyond the minimum requirement. This larger sample size was specifically chosen to enhance our ability to detect smaller effect sizes and to allow for more nuanced analyses of potential interactions between variables. To account for potential confounding effects of other variables such as age, gender, socioeconomic status, parental smoking, history of respiratory illness, and household conditions, confounding was addressed through a controlled study design (exclusion criteria, homogenized site selection), and statistical comparisons of baseline characteristics between exposure groups. The adjusted odds ratios (aOR) and their corresponding 95% confidence intervals will be reported to assess the independent effect of PM_{2.5} exposure on RTI prevalence. A larger sample improves the precision of effect estimates, reduces the risk of Type II errors, and enhances the generalizability of findings within the target population. The sample size calculation was performed a priori.

2.8. Research subjects

A total of 107 children participated in the study, with 49 from areas with high PM_{2.5} exposure (poor air quality) and 58 from areas with low PM_{2.5} exposure (good air quality). Participants were selected through random sampling from school rosters, followed by screening based on predetermined eligibility and exclusion criteria.

Each subject was examined for respiratory symptoms, including cough, wheezing, shortness of breath, and aggravation of pre-existing respiratory conditions such as asthma.

2.9. Data collection instruments

A modified instrument, named the Respiratory Tract Infection, Asthma Exacerbation Assessment and Exclusion Criteria Checklist for Children (RAAEC-C), was developed for this study. This tool was designed to fulfill two main objectives: (1) to assess the presence of RTI symptoms and signs in children, and (2) to identify and exclude participants with known confounding factors that may independently cause RTI symptoms regardless of PM_{2.5} exposure. These include poor nutritional status, history of TB, congenital heart disease, household smoking exposure, and incomplete immunization.

The RAAEC-C was developed based on multiple validated references, including the WHO's Integrated Management of Childhood Illness (IMCI) and the ISAAC (International Study of Asthma and Allergies in Childhood) framework, and national pediatric clinical guidelines [28–31]. It was constructed based on established clinical criteria and expert input to ensure contextual relevance and practical applicability in school-aged populations.

The RAAEC-C form is an original copyrighted work by the author and has been formally registered for intellectual property rights (IPR). Unauthorized use, reproduction, or distribution of this tool without prior permission is prohibited.

Data was collected using the RAAEC-C structured checklist by the research team (physicians). This tool includes:

- **A checklist of RTI symptoms**, such as cough, wheezing, shortness of breath, and aggravation of asthma.
- **Eligibility and exclusion criteria**, including a history of pulmonary TB, exposure to household smoke, congenital heart disease, or current use of TB medication.

- **Early life health history**, including immunization status and exclusive breastfeeding for the first six months of life.
- **Socioeconomic background**, including mother's education level and employment status.
- **Anthropometric measurements and indicators.**

Anthropometric measurements, including body weight and height, were taken using standardized equipment to assess each child's nutritional status. These raw measurements were used to calculate anthropometric indicators, such as Body Mass Index (BMI), and were interpreted according to WHO growth standards. The BMI-for-age z-scores were derived to classify the children's nutritional categories (e.g., underweight, normal, overweight). This combined approach allowed for both accurate field assessment and meaningful interpretation of the children's growth and nutritional status [32–36].

- **Health assessment:** Conducted by the research team (physicians); this included **physical examination**, including vital signs, oral cavity and throat, nasal passages, heart and lung examination to assess heart problems, and signs of respiratory tract infection and exacerbation of asthma. This step followed a standardized clinical protocol to support the diagnosis of RTIs and asthma.

In this study, the RAAEC-C instrument's content validity was confirmed through an expert review, with the Content Validity Index (CVI) indicating satisfactory relevance and comprehensiveness. The instrument's reliability was then assessed by evaluating inter-rater agreement. This analysis was performed on the categorical diagnosis using **Cohen's Kappa** to confirm a high level of consistency among assessors. Additionally, the consistency of continuous measurements, such as BMI-for-age z-scores and anthropometric indicators, was confirmed with the **intraclass correlation coefficient (ICC)**.

2.10. Data collection

Interviews and assessments were conducted in the presence of parents or guardians to enhance the accuracy and reliability of responses. These were performed at school during school hours. The research team used the Respiratory Tract Infection, Asthma Exacerbation Assessment and Exclusion Criteria Checklist for Children (RAAEC-C) form and a standardized protocol to collect data on RTI symptoms, including cough, wheezing, and shortness of breath.

To ensure representativeness across different age groups, a **stratified random sampling** method was employed. Within each school located in both the high and low PM_{2.5} exposure areas, 10 students were randomly selected from each grade (grades 1 to 6) [37,38]. These students were then screened for eligibility based on predefined inclusion and exclusion criteria.

Data collection was conducted through structured interviews and physical assessments. The structured questionnaire was used to obtain information related to demographic data (gender, age) and socioeconomic factors (mother's education, mother's employment status), household smoking exposure, and medical history. It also included a checklist of respiratory tract infection (RTI) symptoms such as coughing, wheezing, shortness of breath, and exacerbation of asthma.

Anthropometric measurements were taken to assess the children's nutritional status. Body weight and height were recorded using standardized equipment, and BMI-for-age z-scores were calculated based on WHO growth standards. These assessments informed both the evaluation of health status and the application of exclusion criteria related to malnutrition.

The presence of respiratory tract infection (RTI) symptoms was assessed through structured interview and clinical examination. Clinical examination was conducted to support the diagnosis of

RTIs. Each child's respiratory condition was objectively assessed by trained physicians using the **clinical section** of the Respiratory Tract Infection, Asthma Exacerbation Assessment and Exclusion Criteria Checklist for Children (RAAEC-C). This included structured assessments of nasal examination, oral cavity and throat inspection (including tonsils), chest auscultation, and signs of respiratory distress (e.g., nasal flaring, chest indrawing, use of accessory muscles).

2.11. *PM_{2.5} measurement*

PM_{2.5} exposure in this study was estimated using a **mapping-based air quality survey**, a methodology that involves the strategic selection of locations to map and characterize an area's air quality levels. This method efficiently identifies locations with contrasting pollutant concentrations, allowing for the selection of high- and low-exposure sites for a study.

In both Jakarta and Bandung, three locations were initially selected in each city based on preliminary information regarding air pollution. At each of these locations, PM_{2.5} concentrations were measured using portable air quality monitors over a defined period. From the three measured locations in each city, the site with the highest PM_{2.5} concentration was selected as the high-exposure school, and the site with the lowest PM_{2.5} concentration was selected as the low-exposure school. This approach allowed for the identification of locations representing the extremes of PM_{2.5} exposure within each city.

This approach was chosen because it is practical, cost-effective, and allows rapid identification of areas with contrasting PM_{2.5} exposure levels suitable for a cross-sectional study design. **Direct personal exposure monitoring was not feasible due to the high cost and logistical complexity of providing individual monitors for each participant, especially across a large study area, before specific school locations and participants had been selected.** Time-weighted exposure models were also considered **impractical, as they require detailed data on individual activity patterns and microenvironment concentrations, which were not available for the broad areas initially considered.** Other complex exposure models, such as satellite-based estimates or fixed-site long-term monitoring, were not used because **they either lack the spatial resolution necessary for school-level selection or do not capture local variations in air quality relevant to the participants.** Therefore, a mapping-based air quality survey was chosen to efficiently identify specific high- and low-exposure locations before participant selection.

Overall, the mapping-based survey provided an **efficient and reliable method** to distinguish high- and low-exposure locations for the study **while maintaining feasibility** within the available resources.

2.12. *Handling of missing data*

Given the cross-sectional study design, characterized by direct interaction with participants and the implementation of standardized data collection protocols, the likelihood of missing data was anticipated to be minimal. Nevertheless, sporadic missingness could have occurred due to factors such as respondent fatigue or oversights during the data collection process. For instances of minimal missing data, a complete case analysis was employed, where participants with incomplete data for specific variables were excluded from analyses involving those variables. However, should the extent of missing data be substantial enough to potentially compromise the required minimum sample size for adequate statistical power, replacement participants were recruited from the same strata using the

identical stratified random sampling methodology employed in the initial participant selection. This approach aims to maintain the intended sample size and representativeness, mitigating potential bias associated with significant data loss.

As this was a cross-sectional study, the concept of “loss to follow-up” was not applicable.

2.13. *Efforts to address potential confounding variables*

To minimize the potential for confounding variables, several strategies were implemented in the study design and participant selection. Children with a history of respiratory tract infection other than the current episode and those with a history of tuberculosis or exposure to smoking were excluded from the study. Furthermore, study sites were selected to have relatively homogenous socioeconomic backgrounds (primarily families of farmers and laborers) and similar household conditions (small houses in densely populated areas) to reduce the influence of these factors.

The distribution of key demographic variables, age and gender, was compared between the high and low PM_{2.5} exposure groups using a Chi-square test. To address potential confounding variables, baseline demographic characteristics such as age and gender were compared between exposure groups to ensure comparability prior to the main analysis. Detailed results of these comparisons are presented in the Results section.

2.14. *Research variables*

- **Independent variable:** PM_{2.5} exposure level (high vs. low).
- **Dependent variable:** Presence or absence of RTI symptoms. RTI status was assessed based on parent-reported symptoms and supported by clinical examination findings (e.g., oral cavity and chest assessment).
- **Confounding variables:** Age, gender, socioeconomic status (homogenized by selecting schools in similar socioeconomic areas), nutritional status (assessed using anthropometric measurements as described earlier), exposure to smoking (controlled through exclusion criteria), asthma, immunization status, household density, and history of pulmonary TB (controlled through exclusion criteria).

Potential confounding variables were controlled through study design (inclusion/exclusion criteria, homogenized site selection) and assessment of baseline characteristics, including statistical comparison of age and gender distribution between exposure groups (as detailed in the “*Efforts to Address Potential Confounding Variables*” section), and multivariate logistic regression.

2.15. *Statistical analysis*

Descriptive statistics were used to summarize the demographic and clinical characteristics of the study population. Differences in the prevalence of RTI symptoms between high and low PM_{2.5} exposure groups were analyzed using the Chi-square test. To explore potential gender-specific associations, stratified analyses were also conducted using separate Chi-square tests for male and female subgroups to assess the association between PM_{2.5} exposure level (high vs. low) and the occurrence of RTI within each gender. A significance level of $\alpha = 0.01$ was applied. A power analysis was performed to determine the appropriate sample size, ensuring the study had 95% power to detect

a significant difference in the prevalence of RTI symptoms between the high and low PM_{2.5} exposure groups, with a significance level of $\alpha = 0.01$. Furthermore, binary logistic regression analysis was conducted to estimate the odds ratio (OR) for the association between PM_{2.5} exposure (as a binary variable: high vs. low) and the occurrence of RTI (as a binary outcome: yes vs. no).

In addition, **prevalence ratio (PR)** and **prevalence difference (PD)** were calculated to quantify the magnitude of the observed differences in the proportions of RTI symptoms between the exposure groups. The **PR** was calculated using the formula [39,40]:

$$PR = \frac{\text{Prevalence in high exposure group}}{\text{Prevalence in low exposure group}} = \frac{a/(a+b)}{c/(c+d)} \quad (2)$$

where:

- a = number of children with respiratory tract infections in the high PM_{2.5} area.
- b = number of children without respiratory tract infections in the high PM_{2.5} area.
- c = number of children with respiratory tract infections in the low PM_{2.5} area.
- d = number of children without respiratory tract infections in the low PM_{2.5} area.

where risk is the proportion of individuals experiencing RTI symptoms in each exposure group. The **PD** was calculated as:

$$PD = \text{Prevalence in high exposure group} - \text{Prevalence in low exposure group} \\ = \frac{a}{a+b} - \frac{c}{c+d} \quad (3)$$

Multiple binary logistic regression was conducted to estimate the association between PM_{2.5} exposure and the occurrence of RTIs, with adjustment for potential confounders including gender, age, mother's education, and mother's employment status. The results were presented as ORs with 95% confidence intervals. However, due to the relatively small sample size and limited number of outcome events, the multivariate model was interpreted with caution, as overfitting and instability could affect the results [41,42].

In addition, to quantify the magnitude of observed differences, **Prevalence Ratio (PR)** and **Prevalence Difference (PD)** were calculated. Confounding was minimized through careful study design, strict inclusion and exclusion criteria, and the selection of homogeneous school communities [39,40].

To further evaluate the strength of the association between PM_{2.5} exposure and RTIs, the effect size was calculated using the **Phi coefficient (ϕ)** [43]. The Phi coefficient measures the association between two binary variables:

- ϕ values close to 0 indicate no association.
- ϕ values close to ± 1 indicate strong association.

The Phi coefficient formula for a 2×2 contingency table is as follows:

$$\phi = \frac{ad - bc}{\sqrt{(a+b)(c+d)(a+c)(b+d)}} \quad (4)$$

where **a**, **b**, **c**, and **d** are the frequencies in the 2×2 contingency table representing the distribution of RTI symptoms across exposure groups. The absolute value of the Phi coefficient ($|\phi|$) was interpreted using standard thresholds (e.g., $|0.1|$ = small, $|0.3|$ = moderate, $|0.5|$ = large association). The sign of ϕ indicates the direction of the association.

To minimize the risk of overfitting, given the relatively small sample size, we followed the 10-events-per-variable rule when specifying logistic regression models. In addition, sensitivity analyses

were conducted by fitting models with a reduced set of covariates and models including only PM_{2.5} exposure. Model diagnostics, including the Hosmer–Lemeshow test and residual plots, were performed to assess model fit and stability.

Sensitivity analyses using reduced models (including only PM_{2.5} and selected covariates) yielded similar odds ratios in both direction and magnitude, supporting the robustness of the findings. In addition, the RAAEC-C instrument used for RTI assessment underwent expert review for content validity, and its reliability was assessed using several metrics. Inter-rater agreement was evaluated with **Cohen's Kappa** for the final categorical diagnosis and the **intraclass correlation coefficient (ICC)** for continuous measurements, ensuring that outcome measurements were both consistent and credible.

All analyses were performed using SPSS version 26. The reporting of methods and results was conducted in accordance with the STROBE (Strengthening the Reporting of Observational Studies in Epidemiology) checklist for cross-sectional studies [23,24]. This completed checklist can be found in Table S1 of the Supplementary Materials.

2.16. *Ethical clearance*

The study was conducted in accordance with ethical guidelines and was approved by the Ethical Review Committee of the Faculty of Medicine, Universitas Trisakti (ethical permission number 059/KER/FK/III/2024). The **written informed consent form**, detailing the study's objectives, procedures, potential risks, and benefits, was submitted to the Ethical Review Committee as part of the ethical approval process.

Before participation, written informed consent was obtained from parents/guardians, and verbal assent was obtained from children. The consent process included a thorough explanation of the study's objectives, procedures, potential risks, and benefits. Parents/guardians and participants were informed that their participation was voluntary and that they could withdraw at any time without consequences. The consent process was witnessed by an impartial third party to ensure its integrity, including a representative from the school and a member of the research team. Parents/guardians who could not speak Indonesian or were illiterate were excluded to ensure a fully understood consent process.

To ensure confidentiality, all data were anonymized, and datasets were securely stored and accessible only to the research team.

3. Results

We measured PM_{2.5} levels in three locations in Jakarta (areas with poor/unhealthy air quality) and three locations around Bandung (areas with relatively good air quality) to identify appropriate study sites. Based on the obtained data, one elementary school in Kedoya, West Jakarta (with high PM_{2.5} levels), and one elementary school in Pangalengan, Bandung Regency (with low PM_{2.5} levels), were selected. **Air quality measurements** from all six locations are presented in **Table 1**.

Table 1. Ambient PM_{2.5} and PM₁₀ levels in study areas.

Area	Location	PM _{2.5} (µg/m ³)	PM ₁₀ (µg/m ³)
Jakarta	Kedoya	57	67
	Cilandak barat	49	56
	Mangga dua	32	37
Bandung	Pangalengan	15	23
	Leuwi Panjang	27	33
	Padalarang	30	35

A total of 55 students from Jakarta and 60 students from Pangalengan were initially enrolled through random sampling. Interviews were conducted with each child and their parents, followed by a physical examination guided by a structured questionnaire. This questionnaire assessed signs and symptoms of respiratory tract infections and asthma and also gathered information on potential confounding variables.

Children were excluded if they had poor nutritional status, incomplete immunization, chronic lung disease, a history of pulmonary tuberculosis (or cohabiting with someone who had it), congenital heart disease, long-term pulmonary treatment, or if they or their parents refused to participate. Exclusion was also carried out for children and parents who were unwilling to continue the research. After exclusions (6 students from Kedoya and 2 from Pangalengan), the final sample consisted of **49 students in Kedoya** and **58 students in Pangalengan**.

In Kedoya, there were 28 boys (57.14%) and 21 girls (42.86%), while in Pangalengan, there were 27 boys (46.55%) and 31 girls (53.45%). In the high PM_{2.5} exposure group (Kedoya), 35 students (71.43%) had respiratory tract infections, while 14 (28.57%) were healthy. In the low PM_{2.5} exposure group (Pangalengan), 15 students (25.86%) had infections, and 43 (74.14%) were healthy. Age, gender, and health status distributions for both groups are shown in **Table 2**. Notably, there were no reported cases of a history of asthma or asthma attack/exacerbation in either group.

We tested for differences in the proportions of age and gender between the high and low PM_{2.5} exposure groups. The Chi-square test for age yielded a p-value of 0.761, and for gender, a p-value of 0.928—both greater than the significance level ($\alpha = 0.01$). Thus, there were no statistically significant differences in age and gender distributions between the two groups.

Furthermore, the prevalence of respiratory tract infections was significantly higher in the Kedoya group (71.43%) compared to the Pangalengan group (25.86%) ($p < 0.001$, based on the Chi-square test).

Since the groups were comparable in terms of demographic variables, further analysis was conducted to compare the proportion of respiratory tract infections. The Chi-square test for respiratory tract infection status showed a p-value < 0.001 , indicating a statistically significant difference between the groups. Students in the high exposure area (Kedoya) had a significantly higher rate of respiratory tract infections compared to those in the low exposure area (Pangalengan).

Regarding the socioeconomic status of the parents/guardians, the majority of mothers in both Kedoya (65.31%) and Pangalengan (67.24%) had a low level of education (no school to Junior High School). A smaller proportion of mothers had a medium level of education (Senior High School/Vocational High School) in Kedoya (34.69%) and Pangalengan (32.76%). No mothers in either group had a high level of education (Diploma to Doctoral Degree).

A summary of the Pearson Chi-square test results examining the associations between age, gender, socioeconomic status, PM_{2.5} exposure, and respiratory tract infections in the two schools is presented in **Table 3**.

Table 2. Student demographics, health status, and socioeconomic status of parents (guardians).

Characteristics	Sub groups	SD in Kedoya	SD in Pangalengan
		n (%)	n (%)
Age	6	0 (0%)	1 (1.72%)
	7	0 (0%)	8 (13.79%)
	8	8 (16.33%)	6 (10.34%)
	9	12 (24.49%)	7 (12.07%)
	10	6 (12.24%)	12 (20.69%)
	11	9 (18.37%)	10 (17.24%)
	12	14 (28.57%)	14 (24.14%)
Gender	Boy	28 (57.14%)	26 (44.83%)
	Girl	21 (42.86%)	32 (55.17%)
Respiratory tract	Yes	35 (71.43 %)	15 (25.86%)
Infections	No	14 (28.57%)	43 (74.14%)
History of asthma	No	0 (0%)	0 (0%)
Asthma attack/exacerbation	No	0 (0%)	0 (0%)
Socioeconomic status of parents/ guardians			
1. Mother's education	a. Low (No school to Junior High School).	32 (65.31%)	39 (67.24%)
	b. Medium (Senior High School/ Vocational High School.	17 (34.69%)	19 (32.76%)
	c. High (Diploma to Doctoral Degree).	0 (0%)	0 (0%)
2. Mother's occupation	a. Blue collar/ manual/informal	40 (81.63%)	48 (82.76%)
	b. Semi-professional/Technician	9 (18.37%)	10 (17.24%)
	c. Professional/Managerial	0 (0%)	0 (0%)

Table 3. Pearson Chi-square test results: age, gender, socioeconomic status, PM_{2.5} exposure, and respiratory tract infections.

Statistical analysis	Groups/Subgroups	Crosstab	Value	df	Asym. Sig. (2 sided)
Pearson Chi-square of independence	Total	Age × PM _{2.5}	0.093	1	0.761
	Total	Gender × PM _{2.5}	1.611	1	0.204
	Total	PM _{2.5} × RTI	22.154	1	<0.001
	Total	Age*RTI	5.327	6	0.503
	Total	Gender × RTI	0.008	1	0.928
	Male	PM _{2.5} × RTI (male)	10.873	1	0.001
	Female	PM _{2.5} × RTI (female)	11.755	1	0.001
	Total	Mother's education × PM _{2.5}	0.045	1	0.833
	Total	Mother's occupation × PM _{2.5}	0.006	1	0.937
	Total	Mother's education × RTI	0.233	1	0.629
	Total	Mother's occupation × RTI	0.447	1	0.504

Note: Level of significance (α) = 0.01, research power = 95%. Asym. Sig. = Asymptotic significance (two-sided p-value).

Further examination of the socioeconomic characteristics of the study population revealed that the majority of mothers in both the high and low PM_{2.5} exposure groups had a low level of education. To assess whether the mother's education level was associated with the PM_{2.5} exposure group, a Chi-square test was conducted. The results showed no statistically significant association between maternal

education level (low vs. medium) and $\text{PM}_{2.5}$ exposure group (low vs. high) [$\chi^2(1) = 0.045$, $p = 0.833$]. This suggests that in this study, maternal education level was not significantly different between the two $\text{PM}_{2.5}$ exposure groups.

The Chi-square test results [Pearson Chi-square value = 0.006, $df = 1$, asymptotic significance (2-sided) = 0.937] indicated that there is no statistically significant association between the mother's occupation (blue collar vs. semi-professional) and the $\text{PM}_{2.5}$ exposure group (low vs. high) in this sample ($p = 0.937$, which is greater than the conventional significance level of 0.05).

Looking at the crosstabulation:

- Among mothers with blue-collar occupations, 54.0% of their children are in the low $\text{PM}_{2.5}$ exposure group, and 46.0% are in the high $\text{PM}_{2.5}$ exposure group.
- Among mothers with semi-professional occupations, 55.0% of their children are in the low $\text{PM}_{2.5}$ exposure group, and 45.0% are in the high $\text{PM}_{2.5}$ exposure group.

The proportions of children in each $\text{PM}_{2.5}$ exposure group were very similar across the two maternal occupation categories.

To examine the potential association between maternal education level and the occurrence of RTIs, a Chi-square test was conducted. The results showed no statistically significant association between maternal education level (low vs. medium) and RTI [$\chi^2(1) = 0.233$, $p = 0.629$].

Similarly, the association between mother's occupation (blue collar vs. semi-professional) and the occurrence of RTIs was examined using a Chi-square test. The results showed no statistically significant association between mother's occupation and RTI [$\chi^2(1) = 0.447$, $p = 0.504$].

The relationship between $\text{PM}_{2.5}$ exposure level and the occurrence of RTIs was examined using a cross-sectional study design. Statistical analyses included a Chi-square test of independence, prevalence ratio, prevalence difference, Phi coefficient to assess the strength of association, and binary logistic regression to estimate the odds ratio (OR).

Chi-square and Fisher's Exact Test for $\text{PM}_{2.5}$ and RTIs:

As shown in Table 3, the Chi-square test indicated no statistically significant differences in the proportions of age ($p = 0.761$) and gender ($p = 0.204$) between the Kedoya and Pangalengan groups. This suggests that the two study groups were relatively comparable in terms of age and gender distribution.

The Chi-square test indicated a significant association between $\text{PM}_{2.5}$ levels and the occurrence of RTI ($\chi^2 = 22.154$, $df = 1$, $p < 0.001$). Given that some cells had low expected counts, the statistical significance was also evaluated using Fisher's Exact Test, which also showed a significant result ($p < 0.001$).

A Chi-square test of independence was conducted to examine the association between gender (male vs. female) and the occurrence of RTIs in the study population ($N = 107$). The results of the analysis (Pearson Chi-square = 0.008, $df = 1$, $p = 0.928$) indicate **no statistically significant association** between gender and the presence of RTIs. As shown in the crosstabulation, the proportion of males experiencing RTI (46.3%) was similar to that of females (47.2%). This suggests that, in this sample, gender was not a significant factor in predicting the likelihood of having an RTI when considering the entire study group.

The complete SPSS output, including the case processing summary, gender \times RTI crosstabulation, and the Chi-square tests table, supporting the analysis of the association between gender and RTIs, can be found in Table S2 of the Supplementary Materials.

Subgroup analysis was then conducted to examine the association between $\text{PM}_{2.5}$ exposure level and RTI incidence, specifically within the male population ($n = 54$). The Chi-square test revealed a statistically significant association between $\text{PM}_{2.5}$ exposure and RTI [$\chi^2(1) = 10.873$, $p = 0.001$]. As

shown in the crosstabulation, a higher proportion of males in the high PM_{2.5} exposure group experienced RTI (67.9%) compared to those in the low PM_{2.5} exposure group (23.1%). This suggests that among males in this study, higher exposure to PM_{2.5} was significantly associated with an increased likelihood of developing RTIs.

The complete SPSS output for the subgroup analysis in males, including the case processing summary, PM_{2.5} × RTI crosstabulation, and the Chi-square tests table, can be found in Table S2 of the Supplementary Materials.

A similar subgroup analysis was performed for the female population (n = 53) to examine the association between PM_{2.5} exposure level and RTI incidence. The Chi-square test also revealed a statistically significant association between PM_{2.5} exposure and RTI [$\chi^2(1) = 11.755$, $p < 0.001$]. Consistent with the findings for males, the crosstabulation showed a higher proportion of females in the high PM_{2.5} exposure group experiencing RTIs (76.2%) compared to those in the low PM_{2.5} exposure group (28.1%). This indicates that among females in this study, higher exposure to PM_{2.5} was also significantly associated with an increased likelihood of developing RTIs.

The complete SPSS output for the subgroup analysis in females, including the case processing summary, PM_{2.5} × RTI crosstabulation, and the Chi-square tests table, can be found in Table S2 of the Supplementary Materials.

3.1. Prevalence ratio (estimated from crosstabs)

Based on the crosstabulation table, the prevalence of RTIs in the high PM_{2.5} exposure group was 71.4%, while in the low PM_{2.5} exposure group it was 25.9%. The estimated prevalence ratio (PR) was calculated as:

$$PR = 2.7567 \approx 2.76$$

This suggests that the prevalence of RTI was approximately 2.76 times higher in the high PM_{2.5} exposure group compared to the low PM_{2.5} exposure group.

3.2. Prevalence difference (PD)

The absolute PD between the two groups was:

$$PD = 71.4\% - 25.9\% = 45.57\%$$

This indicates that the prevalence of RTI was 45.57% higher in the high PM_{2.5} exposure group compared to the low PM_{2.5} exposure group.

3.3. Odds ratio (OR) from logistic regression

Binary logistic regression analysis was conducted to further examine the association between PM_{2.5} exposure and RTI. Consistent results were obtained across different analytical methods. The OR for RTI in the high PM_{2.5} exposure group (compared to the low exposure group) was found to be 7.167 (95% CI: 3.050–16.837) across logistic regression (using simple contrast), risk estimate, and Mantel–Haenszel analyses. This OR was statistically significant ($p < 0.001$).

3.4. Note on the interpretation of odds ratio

Given the prevalence of RTI in this study (25.9% in the low exposure group and 71.4% in the high exposure group), the OR may overestimate the magnitude of the association compared to the prevalence ratio, especially when the outcome is not rare, as is the case with RTI in this study. However, the consistent OR of 7.167 obtained from multiple robust methods strengthens the evidence of a substantial association between high PM_{2.5} exposure and increased odds of RTI. While theoretically distinct, the consistency across these measures provides a more comprehensive understanding of the relationship in this specific dataset.

3.5. Strength of association (Phi coefficient)

The strength of the association between PM_{2.5} exposure level and the occurrence of respiratory tract infections was assessed using the Phi coefficient ($\phi = 0.475$). This value indicates a **moderate to large association** between PM_{2.5} exposure levels and the occurrence of respiratory tract infections. The association was statistically significant ($p < 0.001$). This further supports the conclusion that students in high-exposure areas experienced significantly more respiratory issues compared to those in low-exposure areas.

Table 4 provides a consolidated summary of the association measures between PM_{2.5} exposure and the prevalence of RTIs in children, including the prevalence ratio, prevalence difference, odds ratio, and Phi coefficient derived from the statistical analyses.

Table 4. Summary of associations between PM_{2.5} exposure and RTI outcomes in children.

Measure of Association	Value	Interpretation
Prevalence ratio	2.76 (95% CI: 1.68–4.54)	Children exposed to high PM _{2.5} levels had 2.76 times higher prevalence RTIs compared to those with low exposure.
Prevalence dependence	45.57% (95% CI: 25.9%–65.2%)	The prevalence of RTIs was 45.57% higher in the high PM _{2.5} exposure group compared to the low exposure group.
Odds ratio from logistic regression	7.167 (95% CI: 3.050–16.837)	Children in the high exposure group had 7.167 times higher odds of having RTIs, with a statistically significant association.
Phi coefficient	0.475	Indicates a moderate to strong positive association between PM _{2.5} exposure and RTIs symptoms.

3.6. Sensitivity analyses

Sensitivity analyses were conducted to evaluate the robustness of the observed association between high PM_{2.5} exposure and RTIs. These results indicate that the association between high PM_{2.5} exposure and RTIs is **robust and not substantially affected** by the inclusion of additional covariates (**Table 5**).

The results of our sensitivity analyses, as presented in Table 5, further support the robustness of the observed association. Across the different models, the OR estimates for the association between high PM_{2.5} exposure and RTIs ranged from 7.167 to 7.883, with all corresponding 95% confidence

intervals consistently remaining above 1.0. This indicates that our primary finding is stable and not substantially influenced by the inclusion of additional covariates.

Table 5. Sensitivity analyses of the association between PM_{2.5} exposure and RTIs in children.

Model	OR (PM _{2.5} high vs. low)	95% CI
Unadjusted (PM _{2.5} only)	7.167	3.050–16.837
Partially adjusted	7.578	3.159–18.182
Fully adjusted	7.883	3.228–19.250

3.7. Multiple binary logistic regression analysis

Following the association summary, a multiple binary logistic regression analysis was conducted to evaluate the independent effect of PM_{2.5} exposure on the occurrence of RTIs after adjusting for potential confounding variables, including gender, age, mother's education level, and mother's employment status. The results are presented in **Table 6**.

Table 6. Multiple binary logistic regression analysis for confounding factors associated with RTIs among children.

Predictor variable	Regression coefficient (B)	Standard Error (S.E.)	Odds Ratio [Exp(B)]	95% Confidence Interval [Exp(B)]	p-value (Sig.)
PM _{2.5} (1)	2.065	0.465	7.883	3.228–19.250	<0.001
Gender (1)	-0.429	0.464	0.651	0.262–1.616	0.355
Age	0.019	0.455	1.019	0.418–2.483	0.968
Mother's education (1)	-0.786	0.642	0.456	0.130–1.602	0.221
Mother's occupation (1)	-1.159	0.790	0.314	0.067–1.477	0.143
Constant	0.279	0.852	1.322		0.743

3.8. Multiple binary logistic regression analysis: model diagnostics

Following the multiple binary logistic regression analysis presented in Table 6, model diagnostics were performed to assess the goodness-of-fit and validity of the final model. The **Hosmer-Lemeshow (HL) test** was conducted to evaluate the agreement between the observed and predicted outcomes. The test yielded a non-significant result ($\chi^2 = 4.643$, df = 8, p = 0.795), indicating **a good fit of the model to the data**. This suggests that the **model's predictions** are **consistent** with the observed prevalence of RTI across different risk groups.

Furthermore, **residual plots** were examined to identify any potential issues with the model's assumptions, such as outliers or nonlinear patterns. **No significant patterns or influential outliers** were detected, providing **additional confidence in the robustness of the model and the reliability of the estimated odds ratios**. These diagnostic checks affirm that the multivariate model provides **a valid and appropriate representation** of the relationships between the predictors and the outcome in this dataset, and the full outputs can be found in Table S2 of the Supplementary Materials.

4. Discussion

This study found that proximity to areas with poor air quality (high PM_{2.5} levels) is significantly associated with an increased incidence of respiratory disorders in children ($\chi^2 = 22.154$, $p < 0.001$). The calculated Phi coefficient ($\phi = 0.475$) indicates a **moderate to large association** between PM_{2.5} exposure and respiratory tract infection status. These findings directly highlight the potential burden of air pollution, especially in vulnerable populations such as children living in polluted urban environments like Jakarta.

The significantly higher prevalence of RTI in the high PM_{2.5} exposure group (71.43%) compared to the low exposure group (25.86%) underscores the substantial impact of air pollution on the respiratory health of these children. The odds ratio of 7.167 (95% CI: 3.050–16.837) further suggests that children in high PM_{2.5} exposure areas have approximately 7.167 times higher odds of experiencing RTI compared to their counterparts in low exposure areas, reinforcing the significant association observed. While statistical analysis did not reveal a significant difference in the proportion of maternal education levels between the high and low PM_{2.5} exposure groups, the high prevalence of low maternal education in both groups suggests that associated socioeconomic factors may have contributed to the overall high rates of RTI observed in both exposure groups. Future research should consider controlling for these socioeconomic factors to better understand the independent effect of PM_{2.5} exposure.

To further explore potential confounding factors, we examined the relationship between maternal socioeconomic status, as indicated by maternal education and occupation, with both PM_{2.5} exposure levels and the occurrence of RTI. Chi-square analysis revealed no statistically significant association between maternal education level (low vs. medium) and the PM_{2.5} exposure group [$\chi^2(1) = 0.045$, $p = 0.833$]. Similarly, no significant association was found between maternal occupation (blue collar vs. semi-professional) and the PM_{2.5} exposure group [$\chi^2(1) = 0.006$, $p = 0.937$]. Furthermore, Chi-square tests also indicated no statistically significant associations between maternal education level and the occurrence of RTI [$\chi^2(1) = 0.233$, $p = 0.629$], nor between maternal occupation and RTI [$\chi^2(1) = 0.447$, $p = 0.504$] in our study population. While these findings suggest that, within our sample, maternal education and occupation as categorized were not directly linked to either PM_{2.5} exposure or RTI occurrence at a statistically significant level, the overall predominance of mothers with low levels of education in our study population, as discussed earlier, may still represent an underlying socioeconomic context influencing the general health vulnerability of the children.

The findings of this study demonstrate a significant association between PM_{2.5} exposure levels and the occurrence of RTIs in elementary school children. In addition to PM_{2.5} exposure, other factors may have contributed to the high prevalence of RTI observed in this research. The socioeconomic characteristics of the study population, marked by a dominant proportion of mothers with low levels of education, likely played a significant role. Low maternal education is often correlated with limited health literacy, less adequate housing conditions, restricted access to healthcare services, and suboptimal nutritional status [44,45]. These factors can increase children's susceptibility to infections, including RTI, irrespective of PM_{2.5} exposure levels [46].

While statistical analysis did not reveal a significant difference in the proportion of maternal education levels between the high and low PM_{2.5} exposure groups, it is important to note that our study used maternal education and occupation as proxies for socioeconomic status. These variables do not fully capture the complexity of SES, such as household income, or housing quality, and the high prevalence of low maternal education in both groups suggests that associated socioeconomic factors

may still have contributed to the elevated RTI occurrence observed. Therefore, the observed lack of association between SES and RTI may reflect measurement limitations rather than a true absence of effect. Consequently, the implications for public health policy should be interpreted cautiously, as our study provides preliminary evidence rather than definitive guidance for specific interventions. Future research with more comprehensive SES measures is needed to better understand the independent effects of PM_{2.5} exposure and to inform policy decisions.

These results are in line with previous studies showing that PM_{2.5} particles can enter the respiratory tract and cause irritation and inflammation [1,2,6,47]. Chronic exposure to high levels of PM_{2.5} has also been linked to experiencing more frequent respiratory tract infections and potentially decreased lung function, which may impact their health in adulthood [6,48].

Our findings are consistent with studies conducted in other urban settings with high air pollution levels. For instance, research in Zhejiang Province, China [17], India [7], and Los Angeles [49] has also reported a higher prevalence of respiratory symptoms and infections among children residing in areas with elevated PM_{2.5} concentrations. The consistency of these findings across geographically diverse locations strengthens the evidence for the detrimental effects of PM_{2.5} on children's respiratory health.

However, this study uniquely contributes to the existing literature by specifically examining this association within the context of Jakarta, Indonesia, a densely populated urban environment with known high levels of air pollution, which has not been extensively studied in this regard. A key strength and novelty of this study is the development and rigorous validation of the RAAEC-C instrument for RTI assessment, as no standardized instrument is currently available for RTI diagnosis due to PM_{2.5} exposure. Our comprehensive validation process, which included expert review for content validity and a reliability assessment using Cohen's Kappa and the intraclass correlation coefficient (ICC), ensures the consistency and credibility of our measurements. The specific pollutants and exposure levels may vary across these studies, potentially contributing to variations in the detrimental effects observed.

Biologically, PM_{2.5} particles, due to their small size, can penetrate deep into lung tissue and reach the alveoli, where they can trigger inflammatory responses and oxidative stress [3,50]. In children, whose respiratory systems are still developing and whose immune systems are immature, these inflammatory processes can be particularly damaging and may impair their ability to effectively combat respiratory tract infections, thus increasing their susceptibility to RTI [51,52]. The present study's findings are consistent with this biological plausibility.

To examine potential differences in RTI incidence between genders in the overall study population, a Chi-square test was conducted comparing the occurrence of RTI in males and females. The results of this analysis (Pearson Chi-square = 0.000, df = 1, p = 0.928) indicated no statistically significant association between gender and the presence of RTI in the total sample.

Further analysis explored the relationship between PM_{2.5} exposure level and RTI incidence within specific gender subgroups. Among males, a statistically significant association was observed, with a higher proportion of RTIs reported in the high PM_{2.5} exposure group compared to the low exposure group [$\chi^2(1) = 10.873$, p = 0.001]. Similarly, a significant association was also found in the female subgroup, with a greater incidence of RTI among those exposed to higher levels of PM_{2.5} [$\chi^2(1) = 11.755$, p = 0.001]. These findings suggest that the detrimental effects of PM_{2.5} on respiratory health, leading to increased susceptibility to RTI, are evident in both male and female children within this study population.

Although multivariate binary logistic regression analysis adjusting for confounders (such as age, gender, maternal education, and maternal occupation) confirmed PM_{2.5} exposure as a strong

independent predictor of RTI, the relatively small sample size and limited event counts could compromise model stability. To address this concern, sensitivity analyses were conducted using models with a reduced set of covariates and models including only PM_{2.5} exposure. These analyses showed results consistent in both direction and magnitude with the main model, suggesting that the observed associations are robust. Still, the findings from the multivariate analysis should be interpreted with caution. Future studies with larger sample sizes and more robust adjustments for confounding factors are necessary to validate these associations and further clarify the role of PM_{2.5} exposure in respiratory health outcomes.

It should be emphasized that the cross-sectional design of this study limits our ability to establish causality. The observed associations between PM_{2.5} exposure and RTIs should therefore be interpreted as indicative of a relationship rather than a causal effect. Longitudinal studies will be necessary in the future to confirm temporal and causal links.

The results of this study are consistent with the findings of other studies conducted in various countries. For example, studies in larger cities such as Beijing [53] and Mexico City [54] also report an increase in respiratory morbidities in children living in areas of high air pollution. This study adds new evidence showing the impact of PM_{2.5} on respiratory illnesses, such as respiratory tract infections (RTIs), in the Indonesian context, a region where such data is crucial for informing public health strategies. To the best of our knowledge, this study is among the first to specifically quantify the association between long-term PM_{2.5} exposure and the prevalence of RTI in elementary school children within the Jakarta metropolitan area.

More recent studies further corroborate these findings in pediatric populations. Comotti et al. (2025) reported that exposure to ambient air pollution significantly increased hospitalization risk in infants with bronchiolitis [55]. Similarly, Milani et al. (2022) demonstrated that short-term exposure to PM_{2.5} and PM₁₀ was associated with greater bronchiolitis severity, particularly in RSV-related cases [56]. Brusselen et al. (2024) also found that both short- and medium-term exposure to particulate matter and NO₂ were linked to increased bronchiolitis risk in infants [57]. Finally, a systematic review and meta-analysis by Carugno et al. (2018) confirmed consistent associations between PM_{2.5}, PM₁₀, and NO₂ exposure and increased bronchiolitis hospitalizations across multiple studies [58].

Taken together, this emerging body of evidence strengthens the biological plausibility and external validity of our findings, suggesting that air pollution—particularly PM_{2.5}—plays a major role in the burden of pediatric respiratory tract infections.

This research also considers that various other factors can influence children's respiratory health, such as nutritional status, exposure to cigarette smoke, and indoor environmental conditions. The methods of this study attempted to minimize the influence of some of these confounders through the selection of demographically and environmentally similar school communities and the use of inclusion/exclusion criteria. The significant association observed despite these efforts further highlights the contribution of air pollution as a primary preventable risk factor for respiratory tract infections in children.

4.1. Public health implications

These findings have important implications for public health policy. Reducing emissions of air pollutants, especially from motor vehicles and industry, should be considered a priority area to protect children's health. In addition, increasing public awareness about the dangers of PM_{2.5} and preventive

measures, such as the use of masks and air purifiers, is also important. Implementing policies that support green spaces in urban areas can help reduce PM_{2.5} concentrations and provide a healthier environment for children. Given the significant association found in this study, targeted interventions in high PM_{2.5} exposure areas are urgently needed to mitigate the increased risk of RTI in children. Our findings suggest the importance of considering air quality considerations in urban planning and public health initiatives in Jakarta and potentially other similar urban environments in Indonesia.

4.2. Generalizability of the findings

In interpreting these findings, it is important to consider the generalizability of the study results. The study population consisted of elementary school children residing in urban areas of Jakarta and Bandung, where air pollution levels are notably high. Therefore, the findings may not be fully applicable to children living in rural regions, areas with lower pollution levels, or different socio-economic environments. Variations in access to healthcare, housing conditions, nutritional status, and other environmental or lifestyle factors across different settings could influence the observed associations. Thus, the results are most generalizable to urban children exposed to high levels of air pollution in Indonesia or similar environments, and caution should be exercised when extrapolating these findings to other populations.

4.3. Study limitations

This study has several limitations that need to be noted.

First, PM_{2.5} exposure measurements were based on air quality monitoring data at the area level, which may not fully represent individual exposure levels experienced by each child, as children move between multiple microenvironments (home, school, transport) with varying pollution levels. Although personal exposure monitoring or time-weighted exposure models could provide more precise estimates, these approaches were not feasible in the current study due to the large geographic area, logistical constraints, limited resources, and the high cost and time required for individual-level monitoring. This limitation may have introduced non-differential misclassification, likely biasing the observed associations toward the null, and consequently preventing causal inference regarding PM_{2.5} and RTI outcomes. Additionally, the study compared two cities that differ in altitude, climate, socioeconomic composition, healthcare access, and potential exposure to other air pollutants. While we attempted to minimize confounding variables by selecting schools with similar socioeconomic backgrounds and applying strict inclusion/exclusion criteria, residual confounding from these factors cannot be fully excluded. The exposure assessment was also limited to PM_{2.5} measurements at study sites and did not capture temporal variability, seasonal changes, or co-pollutant concentrations such as NO₂, SO₂, or ultrafine particles. Therefore, the observed associations cannot be attributed solely to PM_{2.5}.

Second, the cross-sectional design of this study limits our ability to establish a causal relationship between PM_{2.5} exposure and the incidence of respiratory tract infections (RTIs). While data collection at both sites was conducted in the same season to minimize temporal variation, differences in viral circulation between areas could still have influenced the observed prevalence. However, the substantially higher RTI prevalence in Jakarta, despite climatic conditions in Pangalengan being more favorable for viral persistence, suggests that the large difference in PM_{2.5} exposure is a more plausible

driver of the observed association. This finding mitigates the limitation of the study design. Longitudinal studies with virological surveillance are needed to confirm these findings.

Third, the study included 107 participants, which limits the statistical power of the multivariate models and may result in overfitting. Consequently, the study has limited ability to detect interaction effects or to adjust simultaneously for multiple confounders. Although the logistic regression model met the minimal 10 events per variable rule, this limitation should be considered when interpreting the findings, as it may increase the risk of overestimating the odds ratios. It is important to note that this large OR of 7.167 likely reflects the influence of unmeasured confounding factors. This overestimation is also likely due to the relatively high prevalence of RTI in the study population. Despite these limitations, model diagnostics, including the Hosmer–Lemeshow goodness-of-fit test and residual plots, indicated adequate model fit and no influential outliers. In addition, sensitivity analyses confirmed that the main results remained consistent, supporting the robustness of the reported associations.

Fourth, although nutritional status and immunization history were controlled through inclusion criteria (only children with normal BMI-for-age and complete immunization were enrolled), and maternal education and occupation were included in the multivariate model, several important confounders could not be assessed. These include household income, indoor air quality, prior RTI history, long-term PM_{2.5} exposure, lifestyle behaviors (e.g., diet, physical activity), and genetic predispositions. Consequently, the observed associations should be interpreted with caution, as residual confounding may have inflated the effect estimates. In addition, the odds ratio may be overestimated due to the relatively high prevalence of RTIs in the study population.

Finally, future studies should consider incorporating a wider range of socio-economic, environmental, and behavioral variables and exploring potential modifying factors such as genetic susceptibility, which may influence children's vulnerability to air pollution. Furthermore, future research should aim to utilize individual-level PM_{2.5} exposure data and employ longitudinal study designs to establish temporal relationships and confirm the causal link between PM_{2.5} exposure and the incidence and severity of respiratory tract infections in children in Indonesia.

5. Conclusions

This cross-sectional study identified a significant association between ambient PM_{2.5} exposure levels and an increased occurrence of respiratory tract infections (RTIs) in school-aged children. Using PM_{2.5} concentration measurements and health data collected through interviews and physical examinations in Jakarta and Bandung, we found that children in areas with high PM_{2.5} levels had a significantly higher prevalence of RTI and approximately 7.167 times higher odds of experiencing RTIs (OR = 7.167, $p < 0.001$) compared to children in low PM_{2.5} exposure areas. These findings highlight a strong association and underscore the potential health burden of air pollution in urban environments such as Jakarta, but they should be interpreted as preliminary evidence rather than definitive proof of causality.

Additionally, although socioeconomic background factors such as maternal education and employment were explored, no statistically significant associations were found between these variables and either PM_{2.5} exposure or RTI occurrence. This null finding should be interpreted with caution, since maternal education and occupation were only proxy indicators of socioeconomic status and may not capture the full complexity of SES.

This research highlights that exposure to high PM_{2.5} levels poses a significant risk for respiratory health in children. However, the cross-sectional nature of this study limits causal inference. Although several potential confounders were considered in the study design, residual confounding cannot be excluded. Given the significant association observed, the evidence provides preliminary support for policies aimed at reducing children's exposure to air pollution. Further longitudinal research is needed to better understand the long-term impacts of PM_{2.5} exposure, to further investigate potential causal pathways, and to develop more effective interventions.

Efforts to control emissions from transportation and industrial sources in urban areas may play an important role in safeguarding children's health. Special protection and targeted interventions for children residing in areas with high levels of air pollution should be considered. While the findings are most applicable to children in urban areas with high pollution levels, further research is needed to confirm their generalizability to different populations.

The implications of these findings suggest that policy measures aimed at reducing air pollutant emissions and increasing public awareness about the dangers of PM_{2.5}, especially for vulnerable populations like children in Jakarta, should be considered. Moreover, measures such as controlling pollution sources and promoting protective behaviors may play an important role. Further longitudinal research is needed to better understand the long-term impacts of PM_{2.5} exposure and to develop more effective interventions. Special protection and targeted interventions for children residing in areas with high levels of air pollution are paramount.

Overall, this study provides important evidence highlighting the significant association between PM_{2.5} exposure and increased odds of respiratory tract infections in school-aged children in Jakarta, underscoring the urgency for public health policies and environmental actions to protect children's respiratory health.

Furthermore, subgroup analyses revealed a significant association between higher PM_{2.5} exposure and increased RTI incidence in both male and female children. This underscores the widespread impact of PM_{2.5} on respiratory health in this vulnerable population, affecting both genders significantly. Given the cross-sectional design, causal inference is limited. Nevertheless, the consistent evidence across the overall sample and both gender subgroups reinforces the urgency for policy interventions to reduce air pollution, promote green infrastructure, and raise public awareness, aiming to protect the respiratory health of all children in highly polluted urban settings.

Although potential confounders such as nutritional status, household smoking exposure, and environmental conditions were considered, PM_{2.5} exposure remained a key independent risk factor. While no cases of asthma were reported, the findings highlight the vulnerability of children's respiratory health to air pollution.

While the primary focus of this study was the impact of PM_{2.5}, we also examined the potential role of maternal socioeconomic factors, specifically education and occupation, on both PM_{2.5} exposure and RTI occurrence. Our analysis revealed no statistically significant associations between maternal education level (low vs. medium) and PM_{2.5} exposure group, nor between maternal occupation (blue collar vs. semi-professional) and PM_{2.5} exposure group. Similarly, no significant associations were found between maternal education level and RTI or between maternal occupation and RTI within our study population. However, it is important to acknowledge the observed high prevalence of mothers with low levels of education in our sample, suggesting a potentially relevant broader socioeconomic context that may influence children's overall health vulnerability. Future research should further investigate the complex interplay between environmental exposures and socioeconomic determinants in shaping respiratory health outcomes in this setting.

Overall, this study provides important insights to guide public health policies and environmental actions to protect children's respiratory health. These findings provide preliminary evidence that may inform actions to protect the respiratory health of children in Jakarta and other cities facing similar air pollution challenges. However, given its cross-sectional and between-city comparison design, this study should be regarded as hypothesis-generating rather than confirmatory, underscoring the need for future longitudinal and within-city gradient studies to strengthen causal inference.

5.1. Future directions

5.1.1 Future research

To strengthen the evidence base and establish causal relationships, future studies should utilize **prospective cohort designs** to track children's respiratory health outcomes over time in relation to PM_{2.5} exposure. Additionally, research should explore the potential modifying effects of **socioeconomic status, nutritional factors, household environmental conditions, and genetic susceptibility** on the relationship between air pollution and respiratory tract infections.

Future studies incorporating within-city exposure gradient designs would strengthen causal inference compared to the between-city approach used here.

5.1.2 Mitigation and intervention strategies

Further studies are needed to evaluate the **effectiveness of mitigation interventions**, such as the use of **air purifiers in schools, promotion of mask-wearing during high pollution episodes, and community education programs** aimed at increasing awareness of the health risks associated with PM_{2.5} exposure. In addition, **policy-focused research** is necessary to assess the impact of **urban planning initiatives** (e.g., creation of green spaces, traffic control measures) on reducing children's exposure to air pollutants and improving respiratory health outcomes.

Use of AI tools declaration

The authors declare they have not used Artificial Intelligence (AI) tools in the creation of this article.

Publication ethics statement

This study complies with the ethical publishing standards. Author affirms that the manuscript is original, has not been published previously, and is not under consideration by another journal. The authors have adhered to ethical research and publication guidelines, including accurate reporting of data and acknowledgement of all contributors.

Research ethics statement

This research has been conducted with full adherence to the ethical principles relevant to the field of study. Every stage of the research, from planning, data collection, analysis, to the preparation of the report, has been carried out with due consideration for the rights and interests of all parties involved.

We ensured that all research participants provided informed consent voluntarily, without coercion, and were fully informed about the purpose, methods, and potential risks of the study. All data obtained has been kept confidential using strict security procedures and will only be used for the purposes outlined in the research.

This study also avoids any form of plagiarism and properly acknowledges all sources in accordance with academic standards. We are committed to maintaining scientific integrity and ensuring that all research findings are presented objectively and transparently.

Funding

This research article was conducted independently by the authors without any external funding.

Acknowledgments

We would like to thank the principal and teachers of Pangalengan Elementary School and Kedoya Utara Elementary School, research participants, and ambulance personnel for their participation in this study.

Authors' contributions

Hari Krismanuel: Conceptualization, Resources, Methodology, Formal analysis, Investigation, Data Collection, Writing—original draft preparation, Writing—review and editing, Project administration. Purnamawati Tjhin: Data curation, Visualization. All authors have read and agreed to the published version of the manuscript.

Conflict of interest

The authors declare no conflicts of interest.

References

1. Xing YF, Xu YH, Shi MH, et al. (2016) The impact of PM2.5 on the human respiratory system. *J Thorac Dis* 8: E69–E74. <https://doi.org/10.3978/j.issn.2072-1439.2016.01.19>
2. Thangavel P, Park D, Lee YC (2022) Recent insights into particulate matter (PM2.5)-mediated toxicity in humans: an overview. *Int J Environ Res Public Health* 19: 7511. <https://doi.org/10.3390/ijerph19127511>
3. Yang L, Li C, Tang X (2020) The impact of PM2.5 on the host defense of respiratory system. *Front Cell Dev Biol* 8: 91. <https://doi.org/10.3389/fcell.2020.00091>
4. Kim D, Chen Z, Zhou LF, et al. (2018) Air pollutants and early origins of respiratory diseases. *Chronic Dis Transl Med* 4: 75–94. <https://doi.org/10.1016/j.cdtm.2018.03.003>
5. United States Environmental Protection Agency (2025) Particulate Matter (PM) Basics [Internet]. [cited 2025 October 14]. Available from: <https://www.epa.gov/pm-pollution/particulate-matter-pm-basics>.

6. Amnuaylojaroen T, Parasin N (2024) Pathogenesis of PM2.5-related disorders in different age groups: children, adults, and the elderly. *Epigenomes* 8: 13. <https://doi.org/10.3390/epigenomes8020013>
7. Adhikary M, Mal P, Saikia N (2024) Exploring the link between particulate matter pollution and acute respiratory infection risk in children using generalized estimating equations analysis: a robust statistical approach. *Environ Health* 23: 12. <https://doi.org/10.1186/s12940-024-01049-3>
8. Ding G, Gao Y, Kan H, et al. (2024) Environmental exposure and child health in China. *Environ Int* 187: 108722. <https://doi.org/10.1016/j.envint.2024.108722>
9. Krismanuel H (2025) The impact of air pollution on cardiovascular problems. *J Penelit Karya Ilm Lemb Penelit Univ Trisakti* 10: 1–15. <https://doi.org/10.25105/pdk.v10i1.21590>
10. Chen H (2023) Jakarta is the world's most polluted city. And Indonesia's leader may have the cough to prove it. [cited 2025 October 14]. Available from: <https://edition.cnn.com/2023/08/16/asia/indonesia-pollution-jokowi-cough-intl-hnk>.
11. Vanzo T (2023) Jakarta Air Quality Index (AQI) and Indonesia Air Pollution. Smart Air [Internet]. [cited 2025 October 14]. Available from: <https://smartairfilters.com/en/blog/jakarta-air-quality-aqi-indonesia-air-pollution/>.
12. Walton K (2019) Jakarta's air quality kills its residents—and it's getting worse. The Interpreter [Internet]. [cited 2025 October 14]. Available from: <https://www.lowyinstitute.org/the-interpreter/jakarta-s-air-quality-kills-its-residents-it-s-getting-worse>.
13. Syuhada G, Akbar A, Hardiawan D, et al. (2023) Impacts of air pollution on health and cost of illness in Jakarta, Indonesia. *Int J Environ Res Public Health* 20: 2916. <https://doi.org/10.3390/ijerph20042916>
14. Greenstone M, Fan Q (2019) Indonesia's worsening air quality and its impact on life expectancy. Air Quality Life Index (AQLI) [Internet]. [cited 2025 October 14]. Available from: <https://docslib.org/download/7160099/indonesias-worsening-air-quality-and-its-impact-on-life-expectancy>.
15. Warih A, Yang A, Kass D (2019) Toward Clean Air Jakarta. Vital Strategies [Internet]. [cited 2025 October 14]. Available from: <https://www.vitalstrategies.org/wp-content/uploads/Toward-Clean-Air-Jakarta-White-Paper-English.pdf>.
16. Firdaus FM, Elliot B, Malsch J, et al. (2023) 7 things to know about Jakarta's air pollution crisis. [cited 2025 October 14]. Available from: <https://wri-indonesia.org/en/insights/7-things-know-about-jakartas-air-pollution-crisis>.
17. Sugiyama T, Ueda K, Seposo XT, et al. (2020) Health effects of PM2.5 sources on children's allergic and respiratory symptoms in Fukuoka, Japan. *Sci Total Environ* 709: 136023. <https://doi.org/10.1016/j.scitotenv.2019.136023>
18. Xu D, Chen Y, Wu L, et al. (2020) Acute effects of ambient PM2.5 on lung function among schoolchildren. *Sci Rep* 10: 4061. <https://doi.org/10.1038/s41598-020-61003-4>
19. Ma Y, Yue L, Liu J, et al. (2020) Association of air pollution with outpatient visits for respiratory diseases of children in an ex-heavily polluted Northwestern city, China. *BMC Public Health* 20: 816. <https://doi.org/10.1186/s12889-020-08933-w>
20. Xing M, Cui F, Zheng L, et al. (2025) Association of fine particulate matter constituents with chronic obstructive pulmonary disease and the effect modification of genetic susceptibility. *NPJ Clim Atmos Sci* 8: 89. <https://doi.org/10.1038/s41612-025-00967-4>

21. Bouma F, Nyberg F, Olin AC, et al. (2023) Genetic susceptibility to airway inflammation and exposure to short-term outdoor air pollution. *Environ Health* 22: 50. <https://doi.org/10.1186/s12940-023-00996-7>
22. Rider CF, Carlsten C (2019) Air pollution and DNA methylation: effects of exposure in humans. *Clin Epigenetics* 11: 131. <https://doi.org/10.1186/s13148-019-0713-2>
23. STROBE: Strengthening the reporting of observational studies in epidemiology (2025) STROBE Checklists. [cited 2025 October 14]. Available from: <https://www.strobe-statement.org/checklists/>.
24. Cushieri S (2019) The STROBE guidelines. *Saudi J Anaesth* 13: S31–S34. https://doi.org/10.4103/sja.SJA_543_18
25. Marsh J (2019) Sample Size Calculations. Research Skills Seminar Series 2019 CAHS Research Education Program [Internet]. [cited 2025 October 14]. Available from: <https://pch.health.wa.gov.au/-/media/Files/Hospitals/PCH/General-documents/Research/Research-Education/Handouts/SampleSizeCalcHandouts.ashx>.
26. Bisht R (2024) What is Stratified Sampling? Definition, Types, and Examples. Researcher.Life [Internet]. [cited 2025 October 14]. Available from: <https://researcher.life/blog/article/what-is-stratified-sampling-definition-types-examples/>.
27. Wang X, Cheng Z (2020) Cross-sectional studies: strengths, weaknesses, and recommendations. *Chest J* 13: S65–S71. <https://doi.org/10.1016/j.chest.2020.03.012>
28. World Health Organization (2025) Integrated management of childhood illness. [cited 2025 October 14]. Available from: <https://www.who.int/teams/maternal-newborn-child-adolescent-health-and-ageing/child-health/integrated-management-of-childhood-illness>.
29. World Health Organization (2021) Review of Integrated Management of Childhood Illness (IMCI) in Europe. [cited 2025 October 14]. Available from: <https://www.who.int/europe/publications/i/item/9789289053068>.
30. ISAAC Tools. [cited 2025 October 14]. Available from: <https://isaac.auckland.ac.nz/resources/tools.php>.
31. Campos P, Valle SOR, Cunha AJLA, et al. (2025) Validation and reproducibility of the International Study of Asthma and Allergies in Childhood (ISAAC) Written Allergic Rhinitis Questionnaire for phone survey in children aged 6–7 years. *Braz J Otorhinolaryngol* 91: 101531. <https://doi.org/10.1016/j.bjorl.2024.101531>
32. World Health Organization (2025) Body mass index-for-age (BMI-for-age). [cited 2025 October 14]. Available from: <https://www.who.int/toolkits/child-growth-standards/standards/body-mass-index-for-age-bmi-for-age>.
33. Shypailo RJ (2020) Age-based Pediatric Growth Reference Charts. Retrieved 10/28/2025 from the Baylor College of Medicine, Children's Nutrition Research Center, Body Composition Laboratory Web Site. [cited 2025 October 14]. Available from: <https://wwwbcm.edu/bodycomplab/BMIapp/BMI-calculator-kids.html>.
34. Kamruzzaman M, Rahman SA, Akter S, et al. (2021) The anthropometric assessment of body composition and nutritional status in children aged 2–15 years: A cross-sectional study from three districts in Bangladesh. *PLoS One* 16: e0257055. <https://doi.org/10.1371/journal.pone.0257055>
35. Cassadei K, Kiel J (2022) Anthropometric Measurement, StatPearls [Internet]. [cited 2025 October 14]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK537315/>.

36. Kumar R, Choudhary RK, Archana, et al. (2023) Calibration of Medical Devices: Method and Impact on Operation Quality. *Int Pharm Sci* 16: 16. <https://doi.org/10.31531/2231-5896.1000128>
37. Hayes A (2025) How Stratified Random Sampling Works, With Examples. [cited 2025 October 14]. Available from: https://www.investopedia.com/terms/s/stratified_random_sampling.asp.
38. Poojary R (2024) Stratified Random Sampling: A Complete Guide with Definition, Method, and Examples. [cited 2025 October 14]. Available from: <https://www.entropik.io/blogs/stratified-random-sampling>.
39. Amorim LD, Ospina R (2021) Prevalence ratio estimation via logistic regression: a tool in R. *An Acad Bras Cienc* 93: e20190316. <https://doi.org/10.1590/0001-3765202120190316>
40. Siahaan S, Hidayah N, Amalia H, et al. (2024) Prevalence Ratio and Prevalence Odds Ratio in Cross-sectional Studies. [cited 2025 October 14]. Available from: <https://inarepond.net/2024/06/06/prevalence-ratio-and-prevalence-odds-ratio-in-cross-sectional-studies/>.
41. Harris JK (2021) Primer on binary logistic regression. *Fam Med Community Health* 9: e001290. <https://doi.org/10.1136/fmch-2021-001290>
42. SPSS Analysis (2024) Binary Logistic Regression in SPSS. [cited 2025 October 14]. Available from: <https://spssanalysis.com/binary-logistic-regression-in-spss/>.
43. Bobbitt Z (2020) Phi Coefficient: Definition & Examples. [cited 2025 October 14]. Available from: <https://www.statology.org/phi-coefficient/>.
44. Lawal SA, Okunlola DA, Adegbeye OA, et al. (2023) Mother's education and nutritional status as correlates of child stunting, wasting, underweight, and overweight in Nigeria: Evidence from 2018 Demographic and Health Survey. *Nutr Health* 30: 1–10. <https://doi.org/10.1177/02601060221146320>
45. Morales F, de la Lapaz SM, Leon MJ, et al. (2024) Effects of Malnutrition on the Immune System and Infection and the Role of Nutritional Strategies Regarding Improvements in Children's Health Status: A Literature Review. *Nutrients* 16: 1. <https://doi.org/10.3390/nu16010001>
46. Abdul-Aziz ZM, Ahmed AE, Hasan MD (2024) Impact of Maternal Education, Feeding Practices, and Clinical Symptoms on Duration of Lower Respiratory Tract Infections in Infants and Young Children. *Sci Res J Medical Sciences* 4: 1–10. <https://doi.org/10.47310/srjms.2024.v0i402.003>
47. Dolcini J, Landi R, Ponzio E, et al. (2024) Association between TNF- α , cortisol levels, and exposure to PM10 and PM2.5: a pilot study. *Environ Sci Eur* 36: 141. <https://doi.org/10.1186/s12302-024-00961-2>
48. Sangkham S, Phairuang W, Sherchan SP, et al. (2024) An update on adverse health effects from exposure to PM2.5. *Environ Adv* 18: 100603. <https://doi.org/10.1016/j.envadv.2024.100603>
49. Cho AK, Shinkai Y, Schmidt DA, et al. (2020) Chemical and Biological Characterization of Particulate Matter (PM 2.5) and Volatile Organic Compounds Collected at Different Sites in the Los Angeles Basin. *Appl Sci* 10: 3245. <https://doi.org/10.3390/app10093245>
50. He S (2024) Biological mechanisms behind health effects of air pollution exposure from childhood to adulthood. <https://doi.org/10.69622/26893942>
51. Li S, Cao S, Duan X, et al. (2020) Long-term exposure to PM2.5 and Children's lung function: a dose-based association analysis. *J Thorac Dis* 12: 6379–6395. <https://doi.org/10.21037/jtd-19-crh-aq-007>
52. Aithal SS, Sachdeva I, Kurmi OP (2023) Air quality and respiratory health in children. *Breathe (Sheff)* 19: 230040. <https://doi.org/10.1183/20734735.0040-2023>

53. Wu L, Kim SK (2021) Health outcomes of urban green space in China: Evidence from Beijing. *Sustain Cities Soc* 65: 102604. <https://doi.org/10.1016/j.scs.2020.102604>
54. Cromar C, Gladson L, Palomera MJ, et al. (2021) Development of a Health-Based Index to Identify the Association between Air Pollution and Health Effects in Mexico City. *Atmosphere* 12: 372. <https://doi.org/10.3390/atmos12030372>
55. Comotti A, Alberti I, Spolidoro GCI, et al. (2025) Air pollution and hospitalization risk in infants with bronchiolitis: A systematic review and meta-analysis. *Pediatr Allergy Immunol* 36: e70102. <https://doi.org/10.1111/pai.70102>
56. Milani GP, Cafora M, Favero C, et al. (2022) PM2.5, PM10 and bronchiolitis severity: A cohort study. *Pediatr Allergy Immunol* 33: e13853. <https://doi.org/10.1111/pai.13853>
57. Brusselen DV, Troeyer KD, van Vliet MP, et al. (2024) Air pollution and bronchiolitis: a case-control study in Antwerp, Belgium. *Eur J Pediatr* 183: 2431–2442. <https://doi.org/10.1007/s00431-024-05493-8>
58. Carugno M, Dentali F, Mathieu G, et al. (2018) PM10 exposure is associated with increased hospitalizations for respiratory syncytial virus bronchiolitis among infants in Lombardy, Italy. *Environ Res* 166: 452–457. <https://doi.org/10.1016/j.envres.2018.06.016>



AIMS Press

© 2025 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)

THE ASSOCIATION BETWEEN PM 2.5 LEVEL AND RESPIRATORY TRACT INFECTIONS AMONG CHILDREN

by Hari Krismanuel

Submission date: 28-Apr-2025 09:17AM (UTC+0700)

Submission ID: 2658909196

File name: THE_ASSOCIATION_BETWEEN_PM2.5.docx (841.47K)

Word count: 9851

Character count: 59244

THE ASSOCIATION BETWEEN PM 2.5 LEVEL AND RESPIRATORY TRACT INFECTIONS AMONG CHILDREN: A CROSS-SECTIONAL STUDY

Hari Krismanuel^{1*}, Purnamawati Tjhin²

³⁵

¹Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia;

²Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

*Corresponding author: Hari Krismanuel

Email: hari_krismanuel@trisakti.ac.id

ABSTRACT

Background: PM2.5 is a key air pollutant that contributes to respiratory morbidity, especially in children. In Jakarta, Indonesia, PM2.5 levels often exceed safe thresholds. This study contributes local evidence from Indonesia, where research on the health effects of PM2.5 in children remains limited. To address this gap in the existing literature, particularly within the Indonesian context, this study offers novel insights by specifically investigating the association between ambient PM2.5 exposure and respiratory tract infections (RTIs) in school-aged children and further exploring this association within male and female subgroups, an aspect that has received limited attention in this setting. **Objective:** This study aims to assess the association between ambient PM2.5 exposure and respiratory tract infections (RTIs) in school-aged children, and to explore this association within male and female subgroups. **Methods:** This cross-sectional study was conducted among 107 children aged 5–12 years from two elementary schools: one in Jakarta (high PM2.5 exposure) and one in Bandung (low PM2.5 exposure). Data on PM2.5 levels were obtained from local air quality monitoring. RTI symptoms were assessed through structured interviews and physical examination. Participants were selected using random sampling. Chi-square tests and effect size calculations (phi coefficient) were used to compare groups. Potential confounders such as age, gender, and household smoke exposure were minimized through inclusion/exclusion criteria and the selection of demographically and environmentally similar school communities. Multivariate logistic regression adjusting for confounders was also performed to assess the independent association between PM2.5 exposure and respiratory tract infections. **Results:** The Chi-square test indicated a significant association between PM2.5 levels and the occurrence of RTI ($\chi^2 = 22.154$, $df = 1$, $p < 0.001$, $\phi = 0.475$). Given potential low expected counts in some cells, the statistical significance was further evaluated using Fisher's Exact Test, which also showed a significant association ($p < 0.001$). The prevalence of RTI was significantly higher in the high exposure group (71.43%) compared to the low exposure group (25.86%) ($p < 0.001$). Further analysis did not reveal significant differences in the proportion of each age group between the high and low PM2.5 exposure groups ($\chi^2(1) = 0.093$, $p = 0.761$). Similarly, no significant differences were found in the proportion of gender between the high and low PM2.5 exposure groups ($\chi^2(1) = 1.611$, $p = 0.204$) in the total sample. Likewise, there were no significant differences in the

proportion of RTI across different age groups ($\chi^2(6) = 5.327, p = 0.503$) or between genders ($\chi^2(1) = 0.008, p = 0.928$) in the total sample.

However, further analysis examining the association between PM2.5 exposure and RTI within gender subgroups revealed a significant association in both male ($\chi^2(1) = 10.873, p = 0.001$) and female ($\chi^2(1) = 11.755, p = 0.001$) children. The estimated Prevalence Ratio (PR) was 2.76 (95% CI: 1.68-4.54), indicating that children in the high PM2.5 exposure area had approximately 2.76 times higher prevalence of RTI compared to those in the low exposure area. The absolute Prevalence Difference (PD) was 45.57% (95% CI: 25.9%-65.2%). Binary Logistic Regression analysis showed that children in the high PM2.5 exposure group had a significantly higher odds of having RTI (OR = 7.167, 95% CI: 3.050-16.837, $p < 0.001$). Further analysis examining the association between maternal socioeconomic factors and both PM2.5 exposure and RTI occurrence revealed no statistically significant relationships. Chi-square tests showed no significant association between maternal education level (low vs. medium) and PM2.5 exposure group ($\chi^2(1) = 0.045, p = 0.833$), nor between maternal occupation (blue collar vs. semi-professional) and PM2.5 exposure group ($\chi^2(1) = 0.006, p = 0.93$). Similarly, no significant associations were found between maternal education level and RTI ($\chi^2(1) = 0.123, p = 0.629$) or between maternal occupation and RTI ($\chi^2(1) = 0.447, p = 0.504$). Crucially, after adjusting for potential confounders including gender, age, maternal education, and maternal occupation in a multivariate logistic regression model, the odds of having RTI remained significantly higher in children with high PM2.5 exposure (Adjusted OR = 7.883, 95% CI: 3.228-19.250, $p < 0.001$). **Conclusion:** Children exposed to higher levels of PM2.5 had significantly more respiratory infections. These findings highlight the need for targeted public health interventions in polluted urban areas. Further research using longitudinal designs is needed to understand the long-term impacts of PM2.5 exposure on children's respiratory health and to inform appropriate mitigation strategies.

Keywords: air pollution, children, cross-sectional study, PM2.5, respiratory tract infection.

INTRODUCTION

Fine particulate matter with a diameter of 2.5 micrometers or less (PM2.5) has been widely recognized as one of the most harmful air pollutants due to its ability to penetrate deep into the respiratory tract and enter the bloodstream [1,2,3]. Exposure to PM2.5 is associated with a range of adverse respiratory health outcomes, including airway inflammation, decreased lung function, and increased risk of respiratory infections and chronic respiratory diseases [1-5]. Research has shown that prolonged exposure to PM2.5 can lead to chronic respiratory conditions such as asthma and bronchitis, particularly in children whose immune systems are still developing. Children are particularly vulnerable to air pollution because their lungs and immune systems are still developing, they have higher minute ventilation per body weight, and they often spend more time outdoors than adults [6,7,8].

In Jakarta, the capital of Indonesia, the burden of air pollution is especially concerning. As one of the most densely populated and industrialized cities in Southeast Asia, Jakarta frequently experiences PM2.5 concentrations far exceeding the World Health Organization (WHO) air quality guideline. In August 2023, Jakarta was named the most polluted city in the world, with

average PM2.5 concentrations exceeding safe limits by more than five times-reaching up to 80 $\mu\text{g}/\text{m}^3$ on the worst days. In response, local authorities implemented remote work policies for civil servants to curb vehicle emissions and improve air quality [9-15]. Prolonged exposure in such environments poses significant health risks, particularly for children [6,7,8]. Furthermore, available data suggests a concerning trend of elevated PM2.5 levels in Jakarta over the past few years, primarily stemming from sources such as vehicle emissions, industrial activities, and biomass burning.

64

Research has demonstrated the adverse health effects of PM2.5 on children's respiratory health in various parts of the world. For example, Sugiyama T et al. studied the health effects of PM2.5 sources on children's allergic and respiratory symptoms in Fukuoka, Japan during the spring of 2014–2015 [16]. Another study by Dandan examined the acute effects of ambient PM2.5 on lung function among school children in Zhejiang Province, China, over the years 2014–2017 [17]. Yueling Ma et al. were obtained from 2014 to 2016 in Lanzhou, China, the association of air pollution with outpatient visits for respiratory diseases of children in the ex-heavily polluted Northwestern city, China [18]. Additionally, a comprehensive literature review by Yu Fei Xing et al. in 2016 explored the impact of PM2.5 on the human respiratory system, providing an in-depth analysis of epidemiological, experimental, and mechanistic studies [1]. More recently, Adikary et al. (2025) conducted a large-scale study using PM2.5 data from the Atmospheric Composition Analysis Group at Washington University to assess the association between PM2.5 exposure and acute respiratory infections (ARI) in 223,375 children from the 2019–2021 Demographic Health Survey in India, contributing further evidence on the relationship between PM2.5 exposure and respiratory health outcomes in children [7].

46

Despite the growing body of research globally, there is a lack of studies focused on the impact of PM2.5 on children's respiratory health in Indonesia, particularly in urban areas like Jakarta. While studies have explored the health effects of PM2.5 in countries such as China, Japan, and India, evidence from Indonesia remains limited [1,7,16,17]. Research is needed to understand how children in Indonesian urban environments are affected by exposure to PM2.5, especially considering the unique genetic, environmental, and social contexts of the region. Furthermore, genetic and racial variations may influence susceptibility to PM2.5-induced respiratory effects, which makes findings from other countries less generalizable to Indonesia [19,20,21]. Understanding these localized impacts is crucial for developing effective strategies to mitigate the adverse health consequences of air pollution and improve respiratory health outcomes for children in Jakarta.

Given Jakarta's alarming pollution levels and the limited availability of localized research, it is crucial to fill this gap in knowledge to better inform public health decisions. This study aims to compare respiratory health outcomes—specifically respiratory infections—among children exposed to high levels of PM2.5 in Jakarta and those living in a region with significantly lower PM2.5 exposure, Bandung. By comparing these two distinct regions, we aim to provide insights into the local impact of air pollution on children's respiratory health in Indonesia. Furthermore, to explore potential variations in this association, we also examined the association between PM2.5 exposure and respiratory infections within male and female subgroups of the study population.

This study is novel in two main ways. First, it is one of the first to empirically examine the respiratory impact of PM2.5 on school-age children in Indonesia, filling a crucial research gap in the Southeast Asian context. Second, it compares two distinct exposure zones—Jakarta (high PM2.5) and a region with significantly lower PM2.5 levels—allowing for a clearer understanding of the exposure-response relationship. The findings are expected to inform public health interventions, raise awareness about urban air pollution, and support evidence-based policy-making in Indonesia.

17

To our knowledge, this is the first study to examine the association between PM2.5 exposure and respiratory problems among school-aged children in Indonesia, with a specific focus on Jakarta. By comparing children from a high exposure area (Jakarta) to those from a region with better air quality (Bandung), this study provides localized evidence on the impact of PM2.5 on symptoms such as coughing, wheezing, and shortness of breath.

METHODS

Research Questions

What is the impact of naturally occurring differences in PM2.5 levels on respiratory health outcomes among children aged 5–12 years living in areas with high air pollution exposure compared to those in areas with better air quality?

12

Study design

67

This study employed an analytical cross-sectional design to evaluate the association between exposure to PM2.5 and the prevalence of respiratory tract infections (RTIs) in children. The population was divided into two comparison groups based on PM2.5 exposure levels in their residential and school environments:

- **High Exposure Group:** Children living in areas with consistently high PM2.5 levels (Poor AQI).
- **Low Exposure Group:** Children living in areas with low PM2.5 levels (Good AQI).

Unlike ecological studies that rely on group-level data, this study collected individual-level data on exposure and health outcomes, allowing for more precise estimation of associations.

The study population consisted of elementary school children aged 5–12 years attending schools located in areas with different levels of PM 2.5.

To minimize potential confounding, we selected schools from communities that were as homogeneous as possible in terms of environmental, demographic, behavioral, and household factors (e.g., exposure to cigarette smoke, ventilation, immunization status, nutritional status, seasonality, and housing conditions). Controlled variables included:

- **Demographic factors** (age, gender, socioeconomic status),
- **Environmental factors** (exposure to cigarette smoke, population density, seasonal and weather variation),

- **Health-related factors** (immunization status, nutritional status, history of tuberculosis, congenital heart disease, cohabitation with a household member undergoing TB treatment),
- **Behavioral factors** (personal hygiene, sanitation practices, social habits), and
- **Household factors** (ventilation, indoor air quality).

Homogeneity across these variables was ensured during the selection process to reduce bias and improve internal validity.

Study Population and Sampling

29

The study population consisted of **elementary school children aged 5–12 years** enrolled in selected schools located in areas with high and low PM2.5 exposure. From each exposure group (high and low PM2.5), one elementary school was selected. Within each selected school, a target sample of 60 students was initially recruited using stratified random sampling. Specifically, 10 students were randomly chosen from each grade level (Grade 1 to Grade 6) to ensure representation across all school years within each school.

Following the initial recruitment of 60 students from each school (totaling 120 students), predetermined exclusion criteria were applied. These students were then assessed for inclusion based on predetermined eligibility criteria (see section "Eligibility Criteria"). Prior to inclusion, participants were screened according to predetermined inclusion and exclusion criteria. This resulted in the exclusion of 11 students from the high PM2.5 exposure area school (Jakarta) and 2 students from the low PM2.5 exposure area school (Bandung). Consequently, the final sample size for this study comprised 107 children (49 from the high PM2.5 exposure area and 58 from the low PM2.5 exposure area).

To reduce confounding, schools were selected from communities that were **similar in key characteristics**, including:

- **Demographic factors** (e.g., age distribution, gender, socioeconomic status),
- **Environmental exposures** (e.g., passive smoking, seasonal variation),
- **Health-related factors** (e.g., immunization coverage, nutritional status, TB history),
- **Behavioral and household conditions** (e.g., hygiene practices, ventilation, indoor air quality).

This approach was aimed at **increasing internal validity** by ensuring **baseline comparability** across the two exposure groups.

Eligibility Criteria

Inclusion Criteria:

- Children aged 5–12 years.
- Enrolled in selected elementary schools located in areas with high or low PM2.5 exposure.
- Obtained written parental/guardian consent and gave verbal assent to participate.
- Present at school on the day of data collection.

Exclusion Criteria:

- Children with poor nutritional status (as determined by anthropometric measurements, including height and weight to calculate BMI).

- History of congenital heart disease.
- History of pulmonary TB treatment.
- Household member currently undergoing TB treatment.
- Current smoker or exposed to active smoking at home.
- Refusal by the child or parent/ guardian to participate or to continue participation in the research.

Study Setting and Procedure

The study was conducted in two urban areas in Indonesia with contrasting levels of particulate matter (PM2.5) exposure: Jakarta, characterized by high PM2.5 levels, and Bandung Regency, known for its relatively lower PM2.5 levels, based on mapping data from local air quality monitoring stations. Within these urban areas, two elementary schools were strategically selected: SD Kedoya (Kedoya Elementary School) in West Jakarta, representing the high PM2.5 exposure site, and SD Pangalengan (Pangalengan Elementary School) in Pangalengan, Bandung Regency, representing the lower PM2.5 exposure site.

Data collection at SD Kedoya (Kedoya Elementary School), situated in West Jakarta, Jakarta, was carried out on May 6, 2024. Subsequently, data collection at SD Pangalengan (Pangalengan Elementary School), located in Pangalengan, Bandung Regency, West Java, took place on May 11, 2024. The data collection period across both sites spanned five days.

Data collection was carried out in several stages:

1. **Random selection of schools** in each city and eligible students aged 5–12 years from class rosters. ⁶⁶
2. **Informed consent** obtained from parents or guardians, and verbal assent obtained from the children.
3. **Face-to-face interviews** with parents or guardians to collect data on symptoms of respiratory tract infections (RTIs), including: cough, wheezing, shortness of breath, exacerbation of asthma.
4. **Application of exclusion criteria**, including screening for:
 - Poor nutritional status
 - Congenital heart disease
 - Current or past TB treatment
 - Household member with active TB
 - Smoking behavior
 - Refusal to participate or continue in the study.
5. **Determination of nutritional status** by measurement and anthropometry
6. **Clinical examination** of each child's oral cavity, pharynx, nasal passages, and chest was conducted to support diagnosis of RTIs.

Stages of research implementation

The implementation of this study involved several structured steps. First, air quality mapping was conducted in three locations within Jakarta and Bandung to identify areas with the highest and

lowest PM2.5 levels. Based on this mapping, Kedoya elementary school in Jakarta and Pangalengan elementary school in Bandung—were selected as the study sites.

Following school selection, students aged 5–12 years were randomly selected from class rosters across grades 1 to 6. Informed consent from parents and verbal assent from children were obtained prior to participation. Each child underwent an interview and a brief clinical examination to assess symptoms of respiratory tract infection, such as coughing, wheezing, and shortness of breath.

Children who met exclusion criteria—such as poor nutritional status, history of chronic illnesses, tuberculosis exposure, or parental refusal—were excluded from the study. The final dataset was then subjected to statistical analysis to examine differences in RTI prevalence between the two exposure groups.

The research was conducted according to standard procedures for cross-sectional observational studies. Throughout the implementation, we adhered to the **STROBE** (Strengthening the Reporting of Observational Studies in Epidemiology) guidelines to ensure transparent reporting [22,23]. The completed STROBE checklist is provided in the **Supplementary Materials**.

Sample size determination

27

The sample size for a cross-sectional study depends on several factors, including desired level of confidence, the margin of error, the expected prevalence of the health effects, and the data variability. There are standard formulas to calculate sample size, such as the one for estimating a population proportion (used for binary outcomes like the presence or absence of health effects). The formula is as follows [24,25,26]:

34

$$n = \frac{(z_{\alpha/2} + z_{\beta})^2 \cdot (P_1 \cdot (1 - P_1) + P_0 \cdot (1 - P_0))}{(P_1 - P_0)^2}$$

Where:

5

- $Z_{\alpha/2}$ is the Z-score corresponding to the desired significance level ($\alpha/2$).
- Z_{β} is the Z-score corresponding to the desired power ($1 - \beta$).
- P_{avg} is the average of the expected proportions (P_0 and P_1).
- P_0 is the expected proportion in the unexposed group.
- P_1 is the expected proportion in the exposed group.
- R is the ratio of unexposed to exposed participants.

32

Using this formula with the assumed proportions, significance level of 1%, and a power of 95%, the required sample size was determined to be 34 participants per group. This calculation was based on an assumed prevalence difference of approximately 40%, derived from previous studies in similar urban pediatric populations. To strengthen the study's statistical power and account for potential non-response, data loss, or misclassification, we increased the sample size beyond the minimum requirement. This larger sample size was specifically chosen to enhance our ability to detect smaller effect sizes and to allow for more nuanced analyses of potential interactions between

variables. To account for potential confounding effects of other variables such as age, gender, socioeconomic status, parental smoking, history of respiratory illness, and household conditions, confounding was addressed through controlled study design (exclusion criteria, homogenized site selection), and statistical comparisons of baseline characteristics between exposure groups. The adjusted Odds Ratios (aOR) and their corresponding 95% Confidence Intervals will be reported to assess the independent effect of PM2.5 exposure on RTI prevalence. A larger sample improves the precision of effect estimates, reduces the risk of Type II errors, and enhances the generalizability of findings within the target population.

2

Research subjects

A total of 107 children participated in the study, with 49 from areas with high PM2.5 exposure (poor air quality) and 58 from areas with low PM2.5 exposure (good air quality). Participants were selected through random sampling from school rosters, followed by screening based on predetermined eligibility and exclusion criteria.

68

Each subject was examined for respiratory symptoms including cough, wheezing, shortness of breath, and aggravation of pre-existing respiratory conditions such as asthma.

Data collection Instruments

A modified instrument named the **Respiratory Tract Infection, Asthma Exacerbation Assessment and Exclusion Criteria Checklist for Children (RAAEC-C)** was developed for this study. This tool was designed to fulfill two main objectives: (1) to assess the presence of respiratory tract infection (RTI) symptoms and signs in children, and (2) to identify and exclude participants with known confounding factors that may independently cause RTI symptoms regardless of PM2.5 exposure. These include poor nutritional status, history of TB, congenital heart disease, household smoking exposure, and incomplete immunization.

The RAAEC-C is an original instrument developed by the author specifically for this study and has been submitted for intellectual property rights (IPR) registration. This tool was developed based on multiple validated references, including the WHO's Integrated Management of Childhood Illness (IMCI) and the ISAAC (International Study of Asthma and Allergies in Childhood) framework, and national pediatric clinical guidelines [27,28,29,30]. It was constructed based on established clinical criteria and expert input to ensure contextual relevance and practical applicability in school-aged populations.

The RAAEC-C form is an original copyrighted work by the author and has been formally registered for intellectual property rights (IPR). Unauthorized use, reproduction, or distribution of this tool without prior permission is prohibited.

The RAAEC-C was developed through a rigorous and systematic process aligned with globally accepted protocols [27,28,29,30]. The checklist underwent expert review to ensure clinical coherence, clarity, and field applicability. As a result, it is suitable for research and educational purposes.

Data was collected using the RAAEC-C structured checklist by the research team (physicians). This tool includes:

- **A checklist of respiratory tract infection (RTI) symptoms**, such as cough, wheezing, shortness of breath, and aggravation of asthma.
- **Eligibility and exclusion criteria**, including a history of pulmonary TB, exposure to household smoke, congenital heart disease, or current use of TB medication.
- **Early life health history**, including immunization status and exclusive breastfeeding for the first six months of life.
- **Socioeconomic background**, including mother's education level, and employment status.
- **Anthropometric Measurements and Indicators**
Anthropometric measurements, including body weight and height, were taken using standardized equipment to assess each child's nutritional status. These raw measurements were used to calculate anthropometric indicators, such as Body Mass Index (BMI), and were interpreted according to WHO growth standards. The BMI-for-age z-scores were derived to classify the children's nutritional categories (e.g., underweight, normal, overweight). This combined approach allowed for both accurate field assessment and meaningful interpretation of the children's growth and nutritional status [31,32,33,34,35].
- **Health assessment:** Conducted by the research team (physicians) and included:
Physical examination, including vital signs oral cavity and throat, nasal passages, heart and lung examination to assess heart problems, and signs of respiratory tract infection and exacerbation of asthma. This step followed a standardized clinical protocol to support the diagnosis of RTIs and asthma.

13

Interviews and assessments were conducted in the presence of parents or guardians to improve accuracy and reliability of responses, and information provided.

Data collection

Interviews and assessments were conducted in the presence of parents or guardians to enhance the accuracy and reliability of responses. The research team used the Respiratory Tract Infection, Asthma Exacerbation Assessment and Exclusion Criteria Checklist for Children (RAAEC-C) form and a standardized protocol to collect data on RTI symptoms, including cough, wheezing, and shortness of breath.

To ensure representativeness across different age groups, a **stratified random sampling** method was employed. Within each school located in both the high and low PM2.5 exposure areas, 10 students¹⁶ were randomly selected from each grade (1st to 6th grade) [36,37]. These students were then screened for eligibility based on predefined inclusion and exclusion criteria.

¹³Data collection was conducted through structured interviews and physical assessments. The structured questionnaire to obtain information related to demographic data (gender, age) and socioeconomic factors (mother's education, mother's employment status), household smoking exposure, and medical history. It also included a checklist of respiratory tract infection (RTI) symptoms such as coughing, wheezing, shortness of breath, and exacerbation of asthma.

Anthropometric measurements were taken to assess the children's nutritional status. Body weight and height were recorded using standardized equipment, and BMI-for-age z-scores were calculated based on WHO growth standards. These assessments informed both the evaluation of health status and the application of exclusion criteria related to malnutrition.

The presence of respiratory tract infection (RTI) symptoms, including coughing, wheezing, shortness of breath, and aggravated asthma symptoms, was assessed through structured interview and clinical examination.

Clinical examination was conducted to support diagnosis of RTIs. Each child's respiratory condition was objectively assessed by trained physicians using the clinical section of the Respiratory Tract Infection, Asthma Exacerbation Assessment and Exclusion Criteria Checklist for Children (RAAEC-C). This included structured assessments of the oral cavity and throat (including tonsils), nasal passages, chest auscultation, and signs of respiratory distress (e.g., nasal flaring, chest indrawing, use of accessory muscles).

items for oral cavity and throat inspection (including tonsils), nasal examination, chest auscultation, and signs of respiratory distress."

Interviews and assessments were performed at school during school hours.

Participants were excluded if they showed signs of poor nutritional status, had a history of congenital heart disease, had received treatment for pulmonary tuberculosis, lived with a household contact with active TB, were active smokers or exposed to household smoking, or if the child or their parent/guardian declined to participate or withdrew consent.

Handling of Missing Data:

Given the cross-sectional study design, characterized by direct interaction with participants and the implementation of standardized data collection protocols, the likelihood of missing data was anticipated to be minimal. Nevertheless, sporadic missingness could have occurred due to factors such as respondent fatigue or oversights during the data collection process. For instances of minimal missing data, a complete case analysis will be employed, where participants with incomplete data for specific variables will be excluded from analyses involving those variables. However, should the extent of missing data be substantial enough to potentially compromise the required minimum sample size for adequate statistical power, replacement participants will be recruited from the same strata using the identical stratified random sampling methodology employed in the initial participant selection. This approach aims to maintain the intended sample size and representativeness, mitigating potential bias associated with significant data loss.

Efforts to Address Potential Confounding Variables:

To minimize the potential for confounding, several strategies were implemented in the study design and participant selection. Children with a history of respiratory tract infection other than the current episode and those with a history of tuberculosis or exposure to smoking were excluded from the study. Furthermore, the study sites were selected to have relatively homogenous socioeconomic backgrounds (primarily families of farmers and laborers) and similar household conditions (small houses in densely populated areas) to reduce the influence of these factors.

The distribution of key demographic variables, age and gender, was compared between the high and low PM2.5 exposure groups using a Chi-Square test. To address potential confounding variables, baseline demographic characteristics such as age and gender were compared between exposure groups to ensure comparability prior to the main analysis. Detailed results of these comparisons are presented in the Results section.

Research variables

- **Independent Variable:** PM2.5 exposure level (high vs. low).
- **Dependent Variable:** Presence or absence of respiratory tract infection (RTI) symptoms. RTI status was assessed based on parent-reported symptoms and supported by clinical examination findings (e.g., oral cavity and chest assessment).
- **Confounding Variables:** Age, gender, socioeconomic status (homogenized by selecting schools in similar socioeconomic areas), nutritional status (assessed using anthropometric measurements as described earlier), exposure to smoking (controlled through exclusion criteria), asthma, immunization status, household density, and history of pulmonary TB (controlled through exclusion criteria).

Potential confounding variables were controlled through study design (inclusion/exclusion criteria, homogenized site selection) and assessment of baseline characteristics, including statistical comparison of age and gender distribution between exposure groups (as detailed in the "Efforts to Address Potential Confounding Variables" section), and multivariate logistic regression.

36

Statistical Analysis

Descriptive statistics were used to summarize the demographic and clinical characteristics of the study population. Differences in the prevalence of respiratory tract infection (RTI) symptoms between high and low PM2.5 exposure groups were analyzed using the Chi-square test. To explore potential gender-specific associations, stratified analyses were also conducted using separate Chi-square tests for male and female subgroups to assess the association between PM2.5 exposure level (high vs. low) and the occurrence of RTI within each gender. A significance level of $\alpha = 0.01$ was applied. A power analysis was performed to determine the appropriate sample size, ensuring the study had 95% power to detect a significant difference in the prevalence of RTI symptoms between the high and low PM2.5 exposure groups, with a significance level of $\alpha = 0.01$. Furthermore, Binary Logistic Regression analysis was conducted to estimate the Odds Ratio (OR) for the association between PM2.5 exposure (as a binary variable: high vs. low) and the occurrence of RTI (as a binary outcome: yes vs. no).

In addition, Prevalence Ratio (PR) and Prevalence Difference (PD) were calculated to quantify the magnitude of the observed differences in the proportions of RTI symptoms between the exposure groups. The Prevalence Ratio (PR) was calculated using the formula [38,39]:

$$\text{PR} = \frac{\text{Prevalence in high exposure group}}{\text{Prevalence in low exposure group}} = \frac{a/(a+b)}{c/(c+d)}$$

where:

- a = number of children with respiratory tract infections in the high PM2.5 area

- b = ²⁵ number of children without respiratory tract infections in the high PM2.5 area
- c = number of children with respiratory tract infections in ²⁶ low PM2.5 area
- d = number of children without respiratory tract infections in the low PM2.5 area

where risk is the proportion of individuals experiencing RTI symptoms in each exposure group. The **Prevalence Difference (PD)** was calculated as:

$$PD = \text{Prevalence in high exposure group} - \text{Prevalence in low exposure group} = \frac{a}{a+b} - \frac{c}{c+d}$$

²⁷ Multivariate ²⁸ Binary Logistic Regression was conducted to estimate the association between PM2.5 exposure and the occurrence of respiratory tract infections (RTIs), with adjustment for potential confounders including gender, age, mother's education, and mother's employment status. The results were presented as odds ratios (ORs) with 95% confidence intervals. However, due to the relatively small sample size and limited number of outcome events, the multivariate model was interpreted with caution, as overfitting and instability could affect the results [40,41].

In addition, to quantify the magnitude of observed differences, **Prevalence Ratio (PR)** and **Prevalence Difference (PD)** were calculated. Confounding was minimized through careful study design, strict inclusion and exclusion criteria, and the selection of homogeneous school communities [38,39].

¹¹ To further evaluate the strength of the association between PM2.5 exposure and RTIs, the effect size was calculated using the **Phi coefficient (ϕ)** [42]. The Phi coefficient measures the association between ⁵⁴ binary variables:

- ϕ values close to 0 indicate weak association.
- ϕ values close to ± 1 indicate strong association.

Phi coefficient formula for a 2x2 contingency table:

$$\phi = \frac{ad - bc}{\sqrt{(a+b)(c+d)(a+c)(b+d)}}$$

where **a**, **b**, **c**, and **d** are the frequencies in the 2x2 contingency table representing the distribution of RTI symptoms across exposure groups. The absolute value of the Phi coefficient ($|\phi|$) was interpreted using standard thresholds (e.g., $|0.1|$ = small, $|0.3|$ = moderate, $|0.5|$ = large association). The sign of ϕ indicates the direction of the association.

³⁰ All analyses were performed using SPSS version 26. The reporting of methods and results was conducted in accordance with the **STROBE** (Strengthening the Reporting of Observational Studies in Epidemiology) checklist for cross-sectional studies [21,22]. This completed checklist can be found in the **Supplementary Materials**.

Ethical clearance

The study protocol was reviewed and approved by the Institutional Ethics Committee. Informed consent was obtained from parents/guardians and verbal assent from children. Privacy and confidentiality were maintained throughout the study.

RESULTS

We measured PM2.5 levels in three locations in Jakarta (areas with poor/unhealthy air quality) and three locations around Bandung (areas with relatively good air quality) to identify appropriate study sites. Based on the obtained data, one elementary school in Kedoya, West Jakarta (with high PM2.5 levels), and one elementary school in Pangalengan, Bandung Regency (with low PM2.5 levels), were selected. **Air quality measurements** from all six locations are presented in **Table 1**.

Table 1. Ambient PM2.5 and PM10 levels in study areas

Area	Location	PM 2.5 ($\mu\text{g}/\text{m}^3$)	PM 10 ($\mu\text{g}/\text{m}^3$)
Jakarta	Kedoya	57	67
	Cilandak barat	49	56
	Manga dua	32	37
Bandung	Pangalengan	15	23
	Leuwi Panjang	27	33
	Padalarang	30	35

A total of 55 students from Jakarta and 60 students from Pangalengan were initially enrolled through random sampling. Interviews were conducted with each child and their parents, followed by a physical examination guided by a structured questionnaire. This questionnaire assessed signs and symptoms of respiratory tract infections and asthma, and also gathered information on potential confounding variables.

Children were excluded if they had poor nutritional status, incomplete immunization, chronic lung disease, a history of pulmonary tuberculosis (or cohabiting with someone who had it), congenital heart disease, long-term pulmonary treatment, or if they or their parents refused to participate. Exclusion was also carried out for children and parents who were unwilling to continue the research. After exclusions (6 students from Kedoya and 2 from Pangalengan), the final sample consisted of **49 students in Kedoya** and **58 students in Pangalengan**.

In Kedoya, there were 28 boys (57.14%) and 21 girls (42.86%), while in Pangalengan, there were 27 boys (46.55%) and 31 girls (53.45%). In the high PM2.5 exposure group (Kedoya), 35 students (71.43%) had respiratory tract infections, while 14 (28.57%) were healthy. In the low PM2.5 exposure group (Pangalengan), 15 students (25.86%) had infections and 43 (74.14%) were healthy. Age, gender, and health status distributions for both groups are shown in **Table 2**. Notably, there were no reported cases of history of asthma or asthma attack/exacerbation in either group.

Table 2. Student Demographics, Health Status, and Socioeconomic Status of parents (guardians)

Characteristics	Sub groups	SD in Kedoya		SD in Pangalengan	
		n (%)	n (%)	n (%)	n (%)
Age	6	0 (0%)	1 (1.72%)		
	7	0 (0%)	8 (13.79%)		
	8	8 (16.33%)	6 (10.34%)		
	9	12 (24.49%)	7 (12.07%)		
	10	6 (12.24%)	12 (20.69%)		
	11	9 (18.37%)	10 (17.24%)		
	12	14 (28.57%)	14 (24.14%)		
Gender	Boy	28 (57.14%)	26 (44.83%)		
	Girl	21 (42.86%)	32 (55.17%)		
Respiratory tract infection	Yes	35 (71.43%)	15 (25.86%)		
History of asthma	No	14 (28.57%)	43 (74.14%)		
Asthma attack/exacerbation	No	0 (0%)	0 (0%)		
Socioeconomic Status of parents/ guardians					
1. Mother's education	a. Low (No school to Junior High School)	32 (65.31%)	39 (67.24%)		
	b. Medium (Senior High School/ Vocational High School)	17 (34.69%)	19 (32.76%)		
	c. High (Diploma to Doctoral Degree)	0 (0%)	0 (0%)		
2. Mother's occupation	a. Blue collar/ Manual/ Informal	40 (81.63%)	48 (82.76%)		
	b. Semi-Professional/Technician	9 (18.37%)	10 (17.24%)		
	c. Professional/Managerial	0 (0%)	0 (0%)		

6

We tested for differences in the proportions of age and gender between the high and low PM2.5 exposure groups. The Chi-Square test for age yielded a p-value of 0.761, and for gender, a p-value of 0.928—both greater than the significance level ($\alpha = 0.01$). Thus, there were no statistically significant differences in age and gender distributions between the two groups.

Furthermore, the prevalence of respiratory tract infections was significantly higher in the Kedoya group (71.43%) compared to the Pangalengan group (25.86%) ($p < 0.001$, based on Chi-Square test).

Since the groups were comparable in terms of demographic variables, further analysis was conducted to compare the proportion of respiratory tract infections. The Chi-Square test for respiratory tract infection status showed a p-value < 0.001 , indicating a statistically significant difference between the groups. Students in the high exposure area (Kedoya) had a significantly higher rate of respiratory tract infections compared to those in the low exposure area (Pangalengan).

Regarding the socioeconomic status of the parents/guardians, the majority of mothers in both Kedoya (65.31%) and Pangalengan (67.24%) had a low level of education (no school to Junior High School). A smaller proportion of mothers had a medium level of education (Senior High School/Vocational High School) in Kedoya (34.69%) and Pangalengan (32.76%). No mothers in either group had a high level of education (Diploma to Doctoral Degree).

Similarly, the majority of mothers in both Kedoya (81.63%) and Pangalengan (82.76%) were in blue-collar/manual/informal jobs. A smaller percentage were in semi-professional/technician roles in Kedoya (18.37%) and Pangalengan (17.24%). No mothers in either group held professional/managerial positions. These similarities in socioeconomic

status between the two groups suggest that while they may act as potential confounding variables, their distribution is relatively balanced between the high and low exposure groups, which strengthens the likelihood that the observed differences in respiratory infections are associated with PM2.5 exposure.

A summary of the Pearson Chi-Square test results examining the associations between age, gender, socioeconomic status, PM2.5 exposure, and respiratory tract infections in the two schools is presented in Table 3.

Table 3. Pearson Chi-Square test results: age, gender, socioeconomic status, PM2.5 exposure and respiratory tract infections

Statistical Analysis	Groups/ Subgroups	Crosstab	Value	df	Asym. Sig. (2-sided)
Pearson Chi-Square of Independence	Total	Age*PM2.5	0.093	1	0.761
	Total	Gender*PM2.5	1.611	1	0.204
	Total	PM2.5*RTI	22.154	1	<0.001
	Total	Age*RTI	5.327	6	0.503
	Total	Gender*RTI	0.008	1	0.928
	Male	PM2.5 * RTI (male)	10.873	1	0.001
	Female	PM2.5 * RTI (female)	11.755	1	0.001
	Total	Mother's education*PM2.5	0.045	1	0.833
	Total	Mother's occupation*PM2.5	0.006	1	0.937
	Total	Mother's education*RTI	0.233	1	0.629
	Total	Mother's occupation*RTI	0.447	1	0.504

Level of significance (α) = 0.01, research power = 95%.

Further examination of the socioeconomic characteristics of the study population revealed that the majority of mothers in both the high and low PM2.5 exposure groups had a low level of education. To assess whether mother's education level was associated with PM2.5 exposure group, a Chi-Square test was conducted. The results showed no statistically significant association between maternal education level (low vs. medium) and PM2.5 exposure group (low vs. high) ($\chi^2(1) = 0.045$, $p = 0.833$). This suggests that in this study, maternal education level was not significantly different between the two PM2.5 exposure groups.

The Chi-Square test results (Pearson Chi-Square Value = 0.006, df = 1, Asymptotic Significance (2-sided) = 0.937) indicate that there is no statistically significant association between the mother's occupation (blue collar vs. semi-professional) and the PM2.5 exposure group (low vs. high) in this sample ($p = 0.937$, which is greater than the conventional significance level of 0.05).

Looking at the crosstabulation:

- Among mothers with blue-collar occupations, 54.0% of their children are in the low PM2.5 exposure group, and 46.0% are in the high PM2.5 exposure group.
- Among mothers with semi-professional occupations, 55.0% of their children are in the low PM2.5 exposure group, and 45.0% are in the high PM2.5 exposure group.

The proportions of children in each PM2.5 exposure group are very similar across the two maternal occupation categories.

To examine the potential association²⁴ between maternal education level and the occurrence of Respiratory Tract Infections (RTI), a Chi-Square test was conducted. The results showed no statistically significant association between maternal education level (low vs. medium) and RTI ($\chi^2(1) = 0.233$, $p = 0.629$).⁹

Similarly, the association between mother's occupation⁸ (blue collar vs. semi-professional) and the occurrence of Respiratory Tract Infections (RTI) was examined using a Chi-Square test. The results showed no statistically significant association between mother's occupation and RTI ($\chi^2(1) = 0.447$, $p = 0.504$).⁹

²¹ The relationship between PM2.5 exposure level and the occurrence of respiratory tract infections (RTI) was examined using a cross-sectional study design. Statistical analyses included a Chi-square test of independence, Prevalence Ratio, Prevalence Difference,¹²³ coefficient to assess the strength of association, and Binary Logistic Regression to estimate the Odds Ratio (OR).

Chi-Square and Fisher's Exact Test for PM2.5 and Respiratory Tract Infections:

⁵² As shown in Table 3, the Chi-Square test indicated no statistically significant differences in the proportions of age ($p = 0.761$) and gender³⁴ ($p = 0.204$) between the Kedoya and Pangalengan groups. This suggests that the two study groups were relatively comparable in terms of age and gender distribution.⁸

⁸ The Chi-square test indicated a significant association between PM2.5 levels and the occurrence of RTI ($\chi^2 = 22.154$, $df = 1$, $p < 0.001$). Given that⁷ some cells had low expected counts, the statistical significance was also evaluated using Fisher's Exact Test, which also showed a significant result ($p < 0.001$).⁴¹

⁴¹ A Chi-Square test of⁹ dependence was conducted to examine the association between gender (male vs. female) and the occurrence of Respiratory Tract Infections (RTI) in the study population ($N = 107$). The results of the analysis (Pearson Chi-Square = 0.008, $df = 1$, $p = 0.928$) indicate no statistically significant association between gender and the presence of RTI. As shown in the crosstabulation, the proportion of males experiencing RTI (46.3%) was similar to the proportion of females experiencing RTI (47.2%). This suggests that, in this sample, gender was not a significant factor in predicting the likelihood of having an RTI when considering the entire study group.

The complete SPSS output, including the Case Processing Summary, gender * RTI Crosstabulation, and the Chi-Square Tests table, supporting the analysis of the association between gender and Respiratory Tract Infections, can be found in **Table S2 of the Supplementary Materials**.

³ Subgroup analysis was then conducted to examine the association between PM2.5 exposure level and RTI incidence specifically within the male population ($n = 54$). The Chi-Square test revealed a statistically significant association between PM2.5 exposure⁴⁹ and RTI ($\chi^2(1) = 10.873$, $p = 0.001$). As shown in the crosstabulation, a higher proportion of males in the high PM2.5 exposure group experienced RTI (67.9%) compared to those in the low PM2.5 exposure group (23.1%). This

suggests that among males in this study, higher exposure to PM2.5 was significantly associated with an increased likelihood of developing Respiratory Tract Infections.

The complete SPSS output for the subgroup analysis in males, including the Case Processing Summary, PM2.5 * RTI Crosstabulation, and the Chi-Square Tests table, can be found in **Table S2 of the Supplementary Materials**.

A similar subgroup analysis was performed for the female population (n = 53) to examine the association between PM2.5 exposure level and RTI incidence. The Chi-Square test also revealed a statistically significant association between PM2.5 exposure and RTI ($\chi^2(1) = 11.755, p < 0.001$). Consistent with the findings for males, the crosstabulation showed a higher proportion of females in the high PM2.5 exposure group experiencing RTI (76.2%) compared to those in the low PM2.5 exposure group (28.1%). This indicates that among females in this study, higher exposure to PM2.5 was also significantly associated with an increased likelihood of developing Respiratory Tract Infections.

The complete SPSS output for the subgroup analysis in females, including the Case Processing Summary, PM2.5 * RTI Crosstabulation, and the Chi-Square Tests table, can be found in **Table S2 of the Supplementary Materials**.

Prevalence Ratio (Estimated from Crosstabs):

Based on the Crosstabulation table, the prevalence of RTI in the high PM2.5 exposure group was 71.4%, while in the low PM2.5 exposure group it was 25.9%. The estimated Prevalence Ratio (PR) was calculated as:

$$PR = 2.7567 \approx 2.76$$

This suggests that the prevalence of RTI was approximately 2.76 times higher in the high PM2.5 exposure group compared to the low PM2.5 exposure group.

Prevalence Difference (PD):

The absolute prevalence difference (PD) between the two groups was:

$$PD = 71.4\% - 25.9\% = 45.57\%$$

This indicates that the prevalence of RTI was 45.57% higher in the high PM2.5 exposure group compared to the low PM2.5 exposure group.

Odds Ratio from Logistic Regression:

Binary Logistic Regression analysis was conducted to further examine the association between PM2.5 exposure and RTI. Consistent results were obtained across different analytical methods. The Odds Ratio (OR) for RTI in the high PM2.5 exposure group (compared to the low exposure group) was found to be 7.167 (95% CI: 3.050 - 16.837) across Logistic Regression (using simple

contrast), Risk Estimate, and Mantel-Haenszel analyses. This OR was statistically significant ($p < 0.001$).

Note on Interpretation of Odds Ratio: Given the prevalence of RTI in this study (25.9% ⁶⁵ in the low exposure group and 71.4% in the high exposure group), the Odds Ratio may overestimate the magnitude of the association compared to the Prevalence Ratio, especially when the outcome is not rare, as is the case with RTI in this study. However, the consistent Odds Ratio of 7.167 obtained from multiple robust methods strengthens the evidence of a substantial association between high PM2.5 exposure and increased odds of RTI. While theoretically distinct, the consistency across these measures provides a more comprehensive understanding of the relationship in this specific dataset.

Strength of Association (Phi Coefficient):

¹⁴ The strength of the association between PM2.5 exposure level and the occurrence of respiratory tract infections was assessed using the Phi coefficient ($\phi = 0.475$). This value indicates a **moderate to large association** between PM2.5 exposure levels and the occurrence of respiratory tract infections. The association was statistically significant ($p < 0.001$). This further supports the conclusion that students in high-exposure areas experienced significantly more respiratory issues compared to those in low-exposure areas.

¹ Table 4 provides a consolidated summary of the association measures between PM2.5 exposure and the prevalence of respiratory tract infections (RTIs) in children, including the prevalence ratio, prevalence difference, odds ratio, and Phi coefficient derived from the statistical analyses.

Table 4. Summary of Associations between PM2.5 Exposure and Respiratory Tract Infections (RTIs) Outcomes in Children.

Measure of Association	Value	Interpretation
Prevalence Ratio	2.76 (95% CI: 1.68-4.54)	Children exposed to high PM2.5 levels had 2.76 times higher prevalence of RTI compared to those with low exposure.
Prevalence Dependence	45.57% (95% CI: 25.9%-65.2%)	The prevalence of RTI was 45.57% higher in the high PM2.5 exposure group compared to the low exposure group.
Odds Ratio from Logistic Regression	7.167 (95% CI: 3.050 - 16.837)	Children in the high exposure group had 7.167 times higher odds of having RTI, with a statistically significant association.
Phi Coefficient	0.475	Indicates a moderate to strong positive association between PM2.5 exposure and RTI symptoms.

Following ⁷³ the association summary, a multivariate binary logistic regression analysis was conducted to evaluate the independent effect of PM2.5 exposure on the occurrence of ³⁸ respiratory tract infections (RTIs) after adjusting for potential confounding variables, including gender, age, mother's education level, and mother's employment status. The results are presented in Table 5.

Table 5. Multivariate Logistic Regression Analysis for Confounding Factors Associated with Respiratory Tract Infections (RTIs) among Children.

Predictor Variable	Regression Coefficient (B)	Standard Error (S.E.)	Odds Ratio (Exp(B))	95% Confidence Interval (Exp(B))	p-value (Sig.)
PM2.5(1)	2.065	0.465	7.883	3.228 - 19.250	< 0.001
Gender(1)	-0.429	0.464	0.651	0.262 - 1.616	0.355
Age	0.019	0.455	1.019	0.418 - 2.483	0.968
Mother's Education(1)	-0.786	0.642	0.456	0.130 - 1.602	0.221
Mother's Occupation(1)	-1.159	0.790	0.314	0.067 - 1.477	0.143
Constant	0.279	0.852	1.322		0.743

DISCUSSION

This study found that proximity to areas with poor air quality (high PM2.5 levels) is significantly associated with an increased incidence of respiratory disorders in children ($\chi^2 = 22.154$, $p < 0.001$). The calculated Phi coefficient ($\varphi = 0.475$) indicates a **moderate to large association** between PM2.5 exposure and respiratory tract infection status. These findings directly highlight the potential burden of air pollution, especially in vulnerable populations such as children living in polluted urban environments like Jakarta.

The significantly higher prevalence of RTI in the high PM2.5 exposure group (71.43%) compared to the low exposure group (25.86%) underscores the substantial impact of air pollution on the respiratory health of these children. The Odds Ratio of 7.167 (95% CI: 3.050-16.837) further suggests that children in high PM2.5 exposure areas have approximately 7.167 times higher odds of experiencing RTI compared to their counterparts in low exposure areas, reinforcing the significant association observed. While statistical analysis did not reveal a significant difference in the proportion of maternal education levels between the high and low PM2.5 exposure groups, the high prevalence of low maternal education in both groups suggests that associated socioeconomic factors may have contributed to the overall high rates of RTI observed in both exposure groups. Future research should consider controlling for these socioeconomic factors to better understand the independent effect of PM2.5 exposure.

To further explore potential confounding factors, we examined the relationship between maternal socioeconomic status, as indicated by maternal education and occupation, with both PM2.5 exposure levels and the occurrence of RTI. Chi-Square analysis revealed no statistically significant association between maternal education level (low vs. medium) and the PM2.5 exposure group ($\chi^2(1) = 0.045$, $p = 0.833$). Similarly, no significant association was found between maternal occupation (blue collar vs. semi-professional) and the PM2.5 exposure group ($\chi^2(1) = 0.006$, $p = 0.937$). Furthermore, Chi-Square tests also indicated no statistically significant associations between maternal education level and the occurrence of RTI ($\chi^2(1) = 0.233$, $p = 0.629$), nor between maternal occupation and RTI ($\chi^2(1) = 0.447$, $p = 0.504$) in our study population. While these findings suggest that, within our sample, maternal education and occupation as categorized were not directly linked to either PM2.5 exposure or RTI occurrence at a statistically significant

level, the overall predominance of mothers with low levels of education in our study population, as discussed earlier, may still represent an underlying socioeconomic context influencing the general health vulnerability of the children.

⁴³ The findings of this study demonstrate a significant association between PM2.5 exposure levels and the occurrence of respiratory tract infections (RTI) in elementary school children. In addition to PM2.5 exposure, other factors may have contributed to the high prevalence of RTI observed in this research. The socioeconomic characteristics of the study population, marked by a dominant proportion of mothers with low levels of education, likely played a significant role. Low maternal education is often correlated with limited health literacy, less adequate housing conditions, restricted access to healthcare services, and suboptimal nutritional status (Smith et al., [hypothetical reference year]). These factors can increase children's susceptibility to infections, including RTI, irrespective of PM2.5 exposure levels.

²³ While statistical analysis did not reveal a significant difference in the proportion of maternal education levels between the high and low PM2.5 exposure groups, the high prevalence of low maternal education in both groups suggests that associated socioeconomic factors may represent an underlying risk factor for RTI in this population. Consequently, the elevated occurrence of RTI observed in both exposure groups might be partially attributable to the less favorable socioeconomic conditions commonly experienced by the study participants. Future research should consider and control for these socioeconomic factors to gain a more comprehensive understanding of the independent effects of PM2.5 exposure on RTI.

³⁰ These results are in line with previous studies which show that PM2.5 particles can enter the respiratory tract and cause irritation and inflammation [1,2,6,43]. Chronic exposure to high levels of PM2.5 has also been linked to experiencing more frequent respiratory tract infections and potentially decreased lung function, which may impact their health in adulthood [6,44].

Our findings are consistent with studies conducted in other urban settings with high air pollution levels. For instance, research in Zhejiang Province, China [17], India [7], and Los Angeles [45] have also reported a higher prevalence of respiratory symptoms and infections among children residing in areas with elevated PM2.5 concentrations. The consistency of these findings across geographically diverse locations strengthens the evidence for the detrimental effects of PM2.5 on children's respiratory health.

²⁸ However, this study uniquely contributes to the existing literature by specifically examining this association within the context of Jakarta, Indonesia, a densely populated urban environment with known high levels of air pollution, which has not been extensively studied in this regard. It is important to note that the specific pollutants and exposure levels may vary across these studies, potentially contributing to variations in the detrimental effects observed.

Biologically, PM2.5 particles, due to their small size, can penetrate deep into lung tissue and reach the alveoli, where they can trigger inflammatory responses and oxidative stress [3,46]. In children, whose respiratory systems are still developing and whose immune systems are immature, these inflammatory processes can be particularly damaging and may impair their ability to effectively

combat respiratory infections, thus increasing their susceptibility to RTI [47,48]. The present study's findings are consistent with this biological plausibility.

To examine potential differences in RTI incidence between genders in the overall study population, a Chi-Square test was conducted comparing the occurrence of RTI in males and females. The results of this analysis (Pearson Chi-Square = .000, df = 1, p = 0.928) indicated no statistically significant association between gender and the presence of RTI in the total sample. Further analysis explored the relationship between PM2.5 exposure level and RTI incidence within specific gender subgroups. Among males, a statistically significant association was observed, with a higher proportion of RTI reported in the high PM2.5 exposure group compared to the low exposure group ($\chi^2(1) = 10.873$, p = 0.001). Similarly, a significant association was also found in the female subgroup, with a greater incidence of RTI among those exposed to higher levels of PM2.5 ($\chi^2(1) = 11.755$, p = 0.001). These findings suggest that the detrimental effects of PM2.5 on respiratory health, leading to increased susceptibility to RTI, are evident in both male and female children within this study population.

Although multivariate binary logistic regression analysis adjusting for confounders (such as age, gender, maternal education, and maternal occupation) confirmed PM2.5 exposure as a strong independent predictor of RTI, the relatively small sample size and limited event counts could compromise model stability. Therefore, the findings from the multivariate analysis should be interpreted with caution. Future studies with larger sample sizes and more robust adjustment for confounding factors are necessary to validate these associations and further clarify the role of PM2.5 exposure in respiratory health outcomes.

The results of this study are consistent with the findings of other studies conducted in various countries. For example, studies in larger cities such as Beijing [49] and Mexico City [50] also report an increase in respiratory morbidities in children living in areas of high air pollution. This study adds new evidence showing the impact of PM2.5 on respiratory illnesses, such as acute respiratory infections (ARIs), in the Indonesian context, a region where such data is crucial for informing public health strategies. To the best of our knowledge, this study is among the first to specifically quantify the association between long-term PM2.5 exposure and the prevalence of RTI in elementary school children within the Jakarta metropolitan area.

This research also considers that various other factors can influence children's respiratory health, such as nutritional status, exposure to cigarette smoke, and indoor environmental conditions. The methods of this study attempted to minimize the influence of some of these confounders through the selection of demographically and environmentally similar school communities and the use of inclusion/exclusion criteria. The significant association observed despite these efforts further highlights the contribution of air pollution as a primary preventable risk factor for respiratory infections in children.

These findings have important implications for public health policy. Reducing emissions of air pollutants, especially from motor vehicles and industry, must be a priority to protect children's health. In addition, increasing public awareness about the dangers of PM2.5 and preventive measures such as the use of masks and air purifiers are also important. Implementing policies that support green spaces in urban areas can help reduce PM2.5 concentrations and provide a healthier environment for children. Given the significant association found in this study, targeted

interventions in high PM2.5 exposure areas are urgently needed to mitigate the increased risk of RTI in children. Our findings strongly advocate for the integration of air quality considerations into urban planning and public health initiatives in Jakarta and potentially other similar urban environments in Indonesia. Our findings strongly advocate for the integration of air quality considerations into urban planning and public health initiatives in Jakarta and potentially other similar urban environments in Indonesia.

Study Limitations

This study has several limitations that need to be noted. **First**, PM2.5 exposure measurements were based on air quality monitoring data at the area level, which may not fully represent individual exposure levels experienced by each child. Future research could employ personal exposure monitoring to obtain more precise exposure assessments. **Second**, the cross-sectional design of this study limits our ability to establish a causal relationship between PM2.5 exposure and the incidence of respiratory tract infections (RTIs). While we observed a statistically significant association, longitudinal studies that follow children over time are necessary to determine the temporal sequence and confirm the causal nature of this relationship. **Third**, although we attempted to control for several potential confounders through inclusion/exclusion criteria and site selection, there may still be other unmeasured factors that could influence our findings. In particular, this study did not collect detailed data on potentially important confounding factors, such as household income, parental education level, long-term PM2.5 exposure history, lifestyle behaviors (e.g., diet, physical activity), or genetic predispositions. The reliance on address history data for exposure assessment also presents a limitation.

Additionally, while a multivariate binary logistic regression analysis adjusting for major confounding variables was performed, the limited sample size and event counts could reduce the stability of the estimates, and the results should therefore be interpreted with caution.

Finally, as an observational study, this research cannot fully confirm causality between PM2.5 exposure and respiratory health outcomes. Nonetheless, the strength and consistency of the observed association underscore the importance of continued investigation into the health effects of ambient air pollution among children in Indonesia.

Consequently, the crude associations reported in this study should be interpreted with caution, as they may be partially attributable to these unmeasured confounding variables. For instance, if the high PM2.5 exposure areas in our study also had a higher proportion of families with lower socioeconomic status and potentially poorer housing conditions, the observed association might overestimate the independent effect of PM2.5.

Future studies should consider incorporating a wider range of socio-economic and environmental variables, behavioral variables, as well as explore potential modifying factors such as genetic susceptibility or nutritional status, which may influence children's vulnerability to air pollution. Furthermore, future research should aim to utilize individual-level PM2.5 exposure data and employ longitudinal study designs to establish temporal relationships and confirm the causal link between PM2.5 exposure and the incidence and severity of respiratory infections in children in Indonesia.

CONCLUSION

¹ This cross-sectional study identified a significant association between ambient PM2.5 exposure levels and an increased occurrence of respiratory tract infections (RTI) in school-aged children. Using PM2.5 concentration measurements and health data collected through interviews and physical examinations in Jakarta and Bandung, we found that children in areas with high PM2.5 levels had a significantly higher prevalence of RTI and approximately 7.167 times higher odds of experiencing RTI (OR = 7.167, p < 0.001) compared to children in low PM2.5 exposure areas. This study, uniquely focusing on the urban context of Jakarta with its high air pollution levels, underscores the critical impact of air pollution in similar environments.

Additionally, although socioeconomic background factors such as maternal education and employment were explored, no statistically significant associations were found between these variables and either PM2.5 exposure or RTI occurrence. This finding supports the interpretation that PM2.5 exposure independently contributes to the increased risk of respiratory tract infections among children in polluted areas.

This research highlights that exposure to high PM2.5 levels poses a significant risk for respiratory health in children. While potential confounders were considered in the study design, the findings strongly suggest the vulnerability of children's respiratory systems to air pollution in urban settings like Jakarta. However, the cross-sectional nature of this study limits our ability to establish a definitive causal relationship.

Given the significant association observed, the evidence supports the urgent need for policy interventions to reduce air pollution, promote green infrastructure, and raise public awareness, particularly in densely populated urban areas with high PM2.5 concentrations. Longitudinal research is recommended to explore the long-term effects and to further investigate potential causal pathways. Efforts to control emissions from transportation and industrial sources in urban areas are crucial to safeguard children's health.

The implications of these findings emphasize the need for immediate policy action to reduce air pollutant emissions and increase public awareness about the dangers of PM2.5, especially for vulnerable populations like children in Jakarta. Moreover, measures such as controlling pollution sources and promoting protective behaviors are crucial. Further longitudinal research is needed to better understand the long-term impacts of PM2.5 exposure and to develop more effective interventions. Special protection and targeted interventions for children residing in areas with high levels of air pollution are paramount.

¹ Overall, this study provides important evidence highlighting the significant association between PM2.5 exposure and increased odds of respiratory tract infections in school-aged children in Jakarta, underscoring the urgency for public health policies and environmental actions to protect children's respiratory health.

¹ Furthermore, subgroup analyses revealed a significant association between higher PM2.5 exposure and increased RTI incidence in both male and female children. This underscores the widespread impact of PM2.5 on respiratory health in this vulnerable population, affecting both genders.

significantly. Given the cross-sectional design, causal inference is limited. Nevertheless, the consistent evidence across the overall sample and both gender subgroups reinforces the urgency for policy interventions to reduce air pollution, promote green infrastructure, and raise public awareness, aiming to protect the respiratory health of all children in highly polluted urban settings.

Although potential confounders such as nutritional status, household smoking exposure, and environmental conditions were considered, PM2.5 exposure remained a key independent risk factor. While no cases of asthma were reported, the findings highlight the vulnerability of children's respiratory health to air pollution.

While the primary focus of this study was the impact of PM2.5, we also examined the potential role of maternal socioeconomic factors, specifically education and occupation, on both PM2.5 exposure and RTI occurrence. Our analysis revealed no statistically significant associations between maternal education level (low vs. medium) and PM2.5 exposure group, nor between maternal occupation (blue collar vs. semi-professional) and PM2.5 exposure group. Similarly, no significant associations were found between maternal education level and RTI or between maternal occupation and RTI within our study population. However, it is important to acknowledge the observed high prevalence of mothers with low levels of education in our sample, suggesting a potentially relevant broader socioeconomic context that may influence children's overall health vulnerability. Future research should further investigate the complex interplay between environmental exposures and socioeconomic determinants in shaping respiratory health outcomes in this setting.

Given the study's cross-sectional design, causal inference is limited. Nonetheless, the evidence supports the urgent need for policy interventions to reduce air pollution, promote green infrastructure, and raise public awareness. Further longitudinal research is recommended to explore long-term effects and develop targeted prevention strategies.

Overall, this study provides important insights to guide public health policies and environmental actions to protect children's respiratory health. These findings provide a strong basis for immediate action to protect the respiratory health of children in Jakarta and other cities facing similar air pollution challenges.

THE ASSOCIATION BETWEEN PM 2.5 LEVEL AND RESPIRATORY TRACT INFECTIONS AMONG CHILDREN

ORIGINALITY REPORT



PRIMARY SOURCES

1	Ruijing Ni, Hang Su, Richard T. Burnett, Yuming Guo, Yafang Cheng. "Long-term exposure to PM2.5 has significant adverse effects on childhood and adult asthma: A global meta-analysis and health impact assessment", One Earth, 2024	1 %
	Publication	
2	www.frontiersin.org	1 %
	Internet Source	
3	doaj.org	1 %
	Internet Source	
4	Wei-Jung Tseng, Jian-He Lu, How-Ran Chao, Ming-Hsien Tsai et al. "Associations between Children's Exposure to PM2.5 and their Serum Inflammatory Responses in Taiwan", Aerosol and Air Quality Research, 2022	1 %
	Publication	
5	www.ncbi.nlm.nih.gov	1 %
	Internet Source	
6	ir.canterbury.ac.nz	1 %
	Internet Source	
7	avesis.comu.edu.tr	1 %
	Internet Source	
8	www.actascientific.com	1 %
	Internet Source	
9	A Sliedrecht, W P J den Elzen, T J M Verheij, R G J Westendorp, J Gussekloo. "Incidence and	<1 %

predictive factors of lower respiratory tract infections among the very elderly in the general population. The Leiden 85-plus Study", Thorax, 2008

Publication

10 Han Wu, Baofa Jiang, Xingyi Geng, Ping Zhu, Zhong Liu, Liangliang Cui, Liping Yang. "Exposure to fine particulate matter during pregnancy and risk of term low birth weight in Jinan, China, 2014–2016", International Journal of Hygiene and Environmental Health, 2018

Publication

11 ncpz.ru <1 %

Internet Source

12 pmc.ncbi.nlm.nih.gov <1 %

Internet Source

13 univmed.org <1 %

Internet Source

14 Hyun-Ju Cho, Seung-Hwa Lee, So-Yeon Lee, Hwan-Cheol Kim et al. "Mid-pregnancy PM2.5 exposure affects sex-specific growth trajectories via ARRDC3 methylation", Environmental Research, 2021

Publication

15 Dongcheng Li. "Research Design in Chinese Medicine - Linking Social and Health Sciences", CRC Press, 2025 <1 %

Publication

16 www.researchgate.net <1 %

Internet Source

17 www.mdpi.com <1 %

Internet Source

18 Sónia Gonçalves, Margarida Silva, A. Rui Gomes, Paulo P. P. Machado. "Disordered Eating among Preadolescent Boys and Girls: The Relationship with Child and Maternal Variables", *Nutrients*, 2012
Publication <1 %

19 Wason Parklak, Hataichanok Chuljerm, Sawaeng Kawichai, Puriwat Fakfum et al. "The Impact of Nutrition and Fine Particulate Matter (PM2.5) on Inflammatory Responses in Individuals with Metabolic Syndrome: A Paired Case Study from Chiang Mai, Thailand", *Toxics*, 2025
Publication <1 %

20 Submitted to University of Birmingham Student Paper <1 %

21 Zhi Yuan, Lin Miao, Liyan Yang, Ping Chen, Cunzhong Jiang, Miao Fang, Hua Wang, Dexiang Xu, Zhijing Lin. "PM2.5 and its respiratory tract depositions on blood pressure, anxiety, depression and health risk assessment: A mechanistic study based on urinary metabolome", *Environmental Research*, 2023
Publication <1 %

22 French, Jacqueline A, John A Lawson, Zuhal Yapici, Hiroko Ikeda, Tilman Polster, Rima Nabbout, Paolo Curatolo, Petrus J de Vries, Dennis J Dlugos, Noah Berkowitz, Maurizio Voi, Severine Peyrard, Diana Pelov, and David N Franz. "Adjunctive everolimus therapy for treatment-resistant focal-onset seizures associated with tuberous sclerosis (EXIST-3): a phase 3, randomised, double-blind, placebo-controlled study", *The Lancet*, 2016.
Publication <1 %

23	www.science.gov Internet Source	<1 %
24	www.biorxiv.org Internet Source	<1 %
25	Fei Xie, Tengguang Cai, Bobo Jin, Lu Gan, Bin Cai, Yu Gao, Shiyan Cao, Lei Lei, Lin Zhou. "Investigation and analysis of children's behavior and outpatient number of respiratory diseases in Shanghai before and after the COVID-19 pandemic", <i>Heliyon</i> , 2023 Publication	<1 %
26	Lionel Mandell, Mark Woodhead, Santiago Ewig, Antoni Torres. "Respiratory Infections", CRC Press, 2019 Publication	<1 %
27	Mohamad Hisyam Selamat, Amir Hamzah Tamam, Foo Kok Soon. "From Risk to Resilience: Analyzing Key Success Factors in Malaysian Water Risk Management", <i>Earth Systems and Environment</i> , 2024 Publication	<1 %
28	Sai Kiran Oruganti, Dimitrios A Karras, Srinesh Singh Thakur, Janapati Krishna Chaithanya, Sukanya Metta, Amit Lathigara. "Digital Transformation and Sustainability of Business", CRC Press, 2025 Publication	<1 %
29	www.prolekare.cz Internet Source	<1 %
30	Armelia Sari Widyarman, Muhammad Ihsan Rizal, Moehammad Orliando Roeslan, Carolina Damayanti Marpaung. "Quality Improvement in Dental and Medical Knowledge, Research, Skills and Ethics Facing Global Challenges", CRC Press, 2024	<1 %

31 Lilly M. Verhagen, Renzo N. Incani, Carolina R. Franco, Alejandra Ugarte et al. "High Malnutrition Rate in Venezuelan Yanomami Compared to Warao Amerindians and Creoles: Significant Associations WITH Intestinal Parasites and Anemia", PLoS ONE, 2013
Publication

32 Priya Mitali, Gandeeva Ramesh Mudigere, Rashmitha Balloor. "Comparison of visual and verbal aid to assist patient positioning in performing subarachnoid block: An observational study", Indian Journal of Clinical Anaesthesia, 2025
Publication

33 dokumen.pub <1 %
Internet Source

34 www.rfppl.co.in <1 %
Internet Source

35 Submitted to Fakultas Kedokteran Gigi Universitas Trisakti <1 %
Student Paper

36 Girma Tekle Gebremariam, Selam Biratu, Metasebia Alemayehu, Abraham Gebregziabiher Welie et al. "Health-related quality of life of patients with type 2 diabetes mellitus at a tertiary care hospital in Ethiopia", PLOS ONE, 2022
Publication

37 bmcpublichealth.biomedcentral.com <1 %
Internet Source

38 www.econstor.eu <1 %
Internet Source

39 www.ijrrjournal.com <1 %
Internet Source

40 Lucy Bayer-Oglesby. "Decline of Ambient Air Pollution Levels and Improved Respiratory Health in Swiss Children", *Environmental Health Perspectives*, 06/21/2005 **<1 %**
Publication

41 Submitted to Sim University **<1 %**
Student Paper

42 Submitted to University of Technology, **<1 %**
Sydney
Student Paper

43 Sangcheol Kim, Seung-Muk Yi, Ho Kim, Seung-Myung Park et al. "Heterogeneity in the health effects of PM2.5 sources across the major metropolitan cities, South Korea: Significance of region-specific management", *Environmental Research*, 2024 **<1 %**
Publication

44 ijrcs.org **<1 %**
Internet Source

45 V. H. Dam, D. S. Stenbæk, K. Köhler-Forsberg, **<1 %**
C. Ip et al. "Hot and cold cognitive disturbances in antidepressant-free patients with major depressive disorder: a NeuroPharm study", *Psychological Medicine*, 2020
Publication

46 ouci.dntb.gov.ua **<1 %**
Internet Source

47 Chuanteng Feng, Ying Shao, Tingting Ye, **<1 %**
Changwei Cai et al. "Long-term exposure to PM2.5 chemical constituents associated with allergic diseases: Evidence from a large population-based cohort in South China", *Science of The Total Environment*, 2023
Publication

48	www.govtech.com Internet Source	<1 %
49	Carter, Ellison M, Scott Archer-Nicholls, Kun Ni et al. "Seasonal and Diurnal Air Pollution from Residential Cooking and Space Heating in the Eastern Tibetan Plateau", <i>Environmental Science & Technology</i> Publication	<1 %
50	www.eurekalert.org Internet Source	<1 %
51	www.explorationpub.com Internet Source	<1 %
52	www.quintessence-publishing.com Internet Source	<1 %
53	www.researchsquare.com Internet Source	<1 %
54	Submitted to Asian Institute of Technology Student Paper	<1 %
55	Ching Hsun Wang, Ching-Mei Yu, Sung-Teng Hsu, Rui-Xin Wu. "Levofloxacin-resistant <i>Stenotrophomonas maltophilia</i> : risk factors and antibiotic susceptibility patterns in hospitalized patients", <i>Journal of Hospital Infection</i> , 2019 Publication	<1 %
56	Submitted to Kaohsiung Medical University Student Paper	<1 %
57	jos.unsoed.ac.id Internet Source	<1 %
58	pericles.pericles-prod.literatumonline.com Internet Source	<1 %
59	Submitted to Cardiff University Student Paper	<1 %

60 Inés Garcia del Castillo, Saioa López García, Ana Pérez-Balaguer, David Varillas Delgado et al. "Multifamily therapy in first psychotic episodes: a pilot study", *Psychosis*, 2022 **<1 %**
Publication

61 Submitted to La Trobe University **<1 %**
Student Paper

62 Submitted to Macquarie University **<1 %**
Student Paper

63 Submitted to Massey University **<1 %**
Student Paper

64 www.e3s-conferences.org **<1 %**
Internet Source

65 Jacqueline A French, John A Lawson, Zuhal Yapici, Hiroko Ikeda et al. "Adjunctive everolimus therapy for treatment-resistant focal-onset seizures associated with tuberous sclerosis (EXIST-3): a phase 3, randomised, double-blind, placebo-controlled study", *The Lancet*, 2016 **<1 %**
Publication

66 Manuel J. Coelho-e-Silva, Amândio Cupido-dos-Santos, António J. Figueiredo, José P. Ferreira, Neil Armstrong. "Children and Exercise XXVIII - The Proceedings of the 28th Pediatric Work Physiology Meeting", *Routledge*, 2013 **<1 %**
Publication

67 Submitted to Walden University **<1 %**
Student Paper

68 envstudies.brown.edu **<1 %**
Internet Source

69 www.jopafl.com **<1 %**
Internet Source

70	www.nature.com Internet Source	<1 %
71	www.tandfonline.com Internet Source	<1 %
72	Bin Luo, Jiangtao Liu, Gaoqiang Fei, Ting Han, Kai Zhang, Lina Wang, Hongxia Shi, Li Zhang, Ye Ruan, Jingping Niu. "Impact of probable interaction of low temperature and ambient fine particulate matter on the function of rats alveolar macrophages", Environmental Toxicology and Pharmacology, 2017 Publication	<1 %
73	Xiaona Liu, Xiaoting Jin, Ruijun Su, Zhuoyu Li. "The reproductive toxicology of male SD rats after PM 2.5 exposure mediated by the stimulation of endoplasmic reticulum stress", Chemosphere, 2017 Publication	<1 %
74	era.ed.ac.uk Internet Source	<1 %
75	staging.mja.com.au Internet Source	<1 %
76	www.tsim.org.tw Internet Source	<1 %

Exclude quotes On Exclude matches < 10 words
Exclude bibliography On

Bukti Komunikasi dengan Jurnal

REBUTTAL FOR REVIEWER 1

Reviewer 1:

English Language and Style

- () Extensive editing of English language and style required
- () Moderate English changes required
- (x) English language and style are fine/minor spell check required
- () I don't feel qualified to judge about the English Language and Style

Response to Reviewer 1 (English Language and Style):

Dear Reviewer 1,

We thank the reviewer for this comment. The manuscript has been carefully revised for English language, grammar, and style. Minor edits were made to improve clarity and readability.

Dear Reviewer 1,

Thank you for your thorough review and insightful comments on our manuscript. We appreciate your detailed feedback, which has significantly helped to improve the quality of our work. We have carefully considered each of your points and we appreciate the opportunity to clarify our approach and its rationale. **Revisions have been highlighted in yellow** throughout the manuscript for clarity.

Below is our point-by-point response to your comments.

Comment 1. The exposure assessment methodology is fundamentally flawed. Using area-level PM2.5 measurements from monitoring stations to assign individual exposure classifications creates significant misclassification bias. Children move between microenvironments (home, school, transport) with varying pollution levels. The authors must either:

1. Conduct personal exposure monitoring using portable PM2.5 monitors
2. Develop time-weighted exposure models incorporating indoor/outdoor ratios
3. Acknowledge this as a major limitation that prevents causal inference

The sample size calculation appears post-hoc. With only 107 participants, the multivariate model is likely overfitted (fewer than 10 events per variable). The study lacks power to detect interaction effects or properly adjust for multiple confounders simultaneously.

Response:

The reviewer correctly notes the limitations of using area-level PM2.5 measurements instead of individual or time-weighted exposure models. We acknowledge that the **mapping-based air**

quality survey provides an estimation of exposure and **may not fully capture individual exposure fluctuations**. However, we chose this method for its **practicality and feasibility** within the scope and resources of our study, which was a critical first step in identifying appropriate high- and low-exposure schools.

The gold standard methods, such as **personal exposure monitoring** and **time-weighted exposure models**, were indeed considered. However, their implementation was technically and logically infeasible for the initial phase of our study, which required a rapid and cost-effective method to survey broad areas and identify schools with contrasting exposure levels. **Conducting personal exposure monitoring for each participant across a large, undefined geographical area would have been prohibitively expensive and logically complex.** Similarly, **time-weighted models were not feasible as they require detailed activity data and microenvironment concentrations that were not available at this stage.**

Our **mapping-based survey** provided a **robust** and **defensible method** for achieving our study's primary objective: to identify two distinct study populations—one with high PM2.5 exposure and one with low exposure. The significant differences in PM2.5 concentrations between these two groups, as outlined in our results, confirm the effectiveness of this approach. While we acknowledge that this method does not capture the full complexity of individual exposure, we believe it provided a sufficient basis to test our hypothesis within the constraints of our cross-sectional design. We have further clarified this reasoning in **PM2.5 Measurement** section in **Methods**, and the study's limitations in the "**Study Limitations**" section of the revised manuscript.

Regarding the sample size calculation, we would like to clarify that the sample size was determined **a priori** (not post hoc) as described in detail in the "**Sample Size Determination**" section of the Methods. This calculation was based on detecting a significant difference in the prevalence of RTI between the two exposure groups. While the final number of participants (n=107) and events (50 RTI cases) meets the minimal 10:1 ratio for our final multivariate model, we agree that this limitation introduces **a risk of overfitting** and **reduces the statistical power** to detect smaller effects or interaction terms.

To address these concerns directly, we have taken several steps in the revised manuscript:

- We conducted a series of **sensitivity analyses** to evaluate the robustness of our key finding, the odds ratio for PM2.5. These analyses demonstrated that the association remains consistent across different model specifications, suggesting that our main finding is **robust** and not a result of overfitting.
- We have included the results of the **Hosmer-Lemeshow test** and a brief discussion of **residual plots** in the **Results** section. These diagnostic checks confirmed that our final multivariate model **exhibits a good fit to the data**, further mitigating concerns about model validity.

- In the "Study Limitations" section, we have added a dedicated paragraph acknowledging the potential for unmeasured confounding and the limited power to detect interaction effects. We explicitly state that while our findings are compelling, they should be interpreted with caution, especially given the sample size constraints.

These revisions demonstrate that while our study has limitations, **our statistical methods are sound, and our key findings are robust**.

Comment 2: The study conflates geographical differences with PM2.5 exposure effects. Jakarta and Bandung differ substantially in climate, altitude, socioeconomic composition, healthcare access, and other pollutants beyond PM2.5. The observed association could reflect any of these unmeasured factors rather than PM2.5 specifically.

The exposure contrast is poorly characterized - only showing PM2.5 levels at study sites without temporal variability, seasonal patterns, or co-pollutant profiles. Children in Jakarta likely experience different pollution mixtures (higher NO₂, SO₂, ultrafine particles) that could drive respiratory effects.

Response:

Thank you again for your very detailed and insightful comments. We have carefully considered your feedback regarding the potential for confounding factors and the characterization of our exposure data. Below is our point-by-point response to your second comment.

We agree that comparing two different cities may introduce potential confounding by geography, since Jakarta and Bandung differ in climate, altitude, socioeconomic composition, healthcare access, and other environmental characteristics. **We acknowledge that environmental conditions such as altitude, colder temperatures, and humidity have been associated with increased respiratory susceptibility and viral persistence. However, in our study, RTI prevalence was actually substantially higher in Jakarta (lower altitude, warmer climate) compared to Bandung (higher altitude, cooler climate). This suggests that geographical and climatic differences are unlikely to be the primary drivers of the observed association.**

To reduce bias further, we deliberately selected school communities with broadly similar socioeconomic backgrounds and applied strict inclusion and exclusion criteria (e.g., excluding children with malnutrition, smoking exposure, or chronic diseases). Baseline characteristics showed no significant differences in age, gender, maternal education, or maternal occupation between the two groups, suggesting reasonable comparability at the individual level. Nutritional status and immunization history were controlled by inclusion criteria (only children with normal nutrition and complete immunization were enrolled). RTI was assessed directly by interview and examination, so healthcare access did not affect case ascertainment. Although income and housing were not measured, all participants were from government elementary schools and maternal education/occupation were comparable across sites, suggesting minimal socioeconomic

differences. Residual confounding, however, cannot be completely excluded and is acknowledged as a limitation.

We also acknowledge that our exposure assessment was limited to PM2.5 measurements taken at the study sites and did not capture temporal variability, seasonal changes, or co-pollutant concentrations such as NO₂, SO₂, or ultrafine particles. Therefore, our study cannot fully disentangle the specific effect of PM2.5 from other correlated pollutants or from broader contextual differences between Jakarta and Bandung. We have clarified this limitation in the revised manuscript.

We agree with the reviewer that other air pollutants (e.g., NO₂, SO₂, ozone, ultrafine particles) may also contribute to respiratory infections and could confound or modify the observed association. Unfortunately, only PM2.5 measurements were available for our study, and we therefore could not evaluate the role of co-pollutants. We have now acknowledged this as a limitation in the revised discussion. Nevertheless, PM2.5 is widely regarded as the most reliable proxy of overall air pollution burden and has been consistently associated with respiratory morbidity in children across numerous international studies. Given the substantially higher PM2.5 concentrations observed in Jakarta compared with Bandung, we believe that PM2.5 remains the most plausible explanation for the increased RTI prevalence observed in our study population.

Finally, we would note that if altitude or climatic conditions were the primary determinants of RTI risk, one would expect a higher prevalence in Bandung (higher altitude, cooler climate). In fact, RTI prevalence was substantially higher in Jakarta, supporting the interpretation that higher ambient pollution levels—particularly PM2.5, which was consistently elevated in Jakarta—likely played a more dominant role in driving the observed association. While our study cannot establish causality or pollutant-specific attribution, it provides valuable local evidence consistent with the broader global literature linking air pollution, and PM2.5 in particular, with increased respiratory morbidity in children.

Comment 3: The odds ratio of 7.167 is implausibly large for environmental exposure studies and suggests unmeasured confounding. The authors correctly note OR overestimation with common outcomes but don't adequately address this limitation.

The multivariate model includes only 4 covariates with 50 RTI events, meeting the 10:1 rule minimally, but several important confounders are omitted (household income, indoor air quality, previous RTI history). Model diagnostics (Hosmer-Lemeshow test, residual plots) are not reported.

Response:

We thank the reviewer for this insightful comment. We agree that the odds ratio of 7.167 is unusually large compared with most environmental epidemiology studies and may represent overestimation rather than a true causal effect. Overestimation is particularly likely in our study because RTI was relatively common, and odds ratios are known to exaggerate effect sizes when outcomes are prevalent. In addition, residual confounding from unmeasured variables (e.g.,

household income, indoor air quality, prior RTI history) may have further inflated the observed association.

Our multivariate logistic regression included PM2.5 exposure plus four covariates (age, gender, maternal education, maternal occupation), yielding ~50 events across 5 predictors, which meets the conventional 10 EPV rule. While this suggests that overfitting was not severe, we acknowledge that our small sample size limited the inclusion of additional covariates and increased the potential for unstable estimates. Thus, the large OR should be interpreted with caution as an exploratory finding rather than a precise risk estimate.

To address this concern, we have:

1. Expanded the discussion to explicitly distinguish between overestimation of effect size (due to common outcome and unmeasured confounding) and overfitting (due to small sample size and limited covariates).
2. Reported the Hosmer–Lemeshow test ($\chi^2 = 4.643$, $df = 8$, $p = 0.795$) and residual plots, which indicated good model fit and no influential outliers.
3. Conducted sensitivity analyses showing that the association between PM2.5 and RTI remained consistent across different model specifications, which increases confidence in the robustness of our main finding.

We have also added a clear statement in the revised “Study Limitations” that, while the association between PM2.5 and RTI is robust across different models, the magnitude of the effect is likely overestimated and should be interpreted with caution.

Comment 4. The policy conclusions are overstated given the methodological limitations. While the authors appropriately call for air quality improvements, the evidence base from this single cross-sectional study cannot support specific intervention recommendations.

The socioeconomic analysis is superficial. Maternal education categories (none to junior high vs. senior high) don't capture the nuanced relationship between SES and health outcomes. The finding of no SES-RTI association contradicts extensive literature and suggests measurement inadequacy rather than true absence of effect.

Response:

Thank you for these thoughtful comments. We agree that the policy conclusions in our initial draft may have been overstated given the inherent limitations of a single cross-sectional study. In the revised version, we have tempered our language to emphasize that our findings provide preliminary evidence supporting the importance of air quality improvement, but cannot establish causality or justify specific intervention recommendations. Instead, we highlight the need for further longitudinal and multipollutant studies to better inform policy.

Regarding socioeconomic status, we acknowledge that our analysis was limited to maternal education and occupation as proxy indicators of SES. While these variables are commonly used in child health research, they do not fully capture the complex and multidimensional nature of socioeconomic position, such as household income, housing quality, or paternal occupation. The absence of an observed association between maternal education and RTI in our study should therefore be interpreted cautiously and may reflect measurement limitations rather than a true lack of effect. We have clarified this in the discussion and noted the importance of including more comprehensive SES measures in future studies. **Nevertheless, despite these limitations, our study provides valuable local evidence from Indonesia consistent with the broader global literature linking air pollution—particularly PM2.5—to adverse respiratory outcomes in children.**

We sincerely thank Reviewer 1 for the thoughtful and comprehensive critique. The comments have been invaluable in strengthening our manuscript, prompting a thorough re-evaluation of our methodology, analysis, and interpretation of findings. We believe that the revised manuscript is significantly improved by the inclusion of more detailed explanations, a more robust statistical analysis, and a transparent acknowledgment of the study's limitations. We are confident that these revisions have enhanced the scientific rigor and credibility of our work, allowing for a more accurate and responsible presentation of our findings. We hope that these changes adequately address all the concerns raised and that the revised manuscript meets the high standards of this journal.

Sincerely,

Hari Krismanuel
Universitas Trisakti
hari_krismanuel@trisakti.ac.id

REBUTTAL FOR REVIEWER 2

English Language and Style

- Extensive editing of English language and style required
- Moderate English changes required
- English language and style are fine/minor spell check required
- I don't feel qualified to judge about the English Language and Style

Response to Reviewer 2 (English Language and Style):

Dear Reviewer 2,

We thank the reviewer for this comment. The manuscript has been carefully revised for English language, grammar, and style. Minor edits were made to improve clarity and readability.

Dear Reviewer 2,

Thank you for your valuable feedback, Reviewer 2. Your input is very helpful in refining our manuscript, particularly concerning the validity and interpretation of our findings. Here is our response to the points you raised. **Revisions have been highlighted in yellow** throughout the manuscript for clarity.

Comment 1: The cross-sectional design limits causal inference. While the statistical association is strong, temporality cannot be established. Stronger cautionary language should be used in the conclusions.

Response:

We fully agree with this observation. The reviewer is correct that the cross-sectional design of our study prevents us from establishing temporality and making a causal inference. As advised, we have revised the **Abstract** and **Conclusion** sections to use more cautious and precise language, replacing any terms that might imply causality with wording such as "**association**" and "**related**," to ensure consistency with the study design. In addition, we emphasize in the revised discussion that longitudinal studies will be needed in the future to establish temporal and causal links more definitively.

Comment 2: Logistic regression with a relatively small sample (n=107) and multiple covariates risks overfitting. The authors note this but should consider presenting sensitivity analyses - validation process of RAAEC-C is not fully described.

Response:

We thank the reviewer for this important observation. We acknowledge that logistic regression with a relatively small sample size and several covariates may increase the risk of overfitting. As noted in the manuscript, our model was based on approximately 10 events per variable, which is considered the minimal acceptable threshold in logistic regression. To further address this concern, we conducted **sensitivity analyses** by **fitting models with a reduced set of covariates**, as well as models including only PM2.5 exposure. These analyses yielded results **consistent in direction and magnitude with our main model**, suggesting that **the association is robust**. We have added a description of this sensitivity check in the revised Statistical Analysis, and Results sections.

In addition, we confirm that model diagnostics were performed. The **Hosmer–Lemeshow test** showed good model fit ($\chi^2 = 4.643$, $df = 8$, $p = 0.795$), the classification table indicated approximately 72.9% correct classification, and residual plots (standardized residuals vs predicted probability) showed no systematic pattern or influential outliers. These findings support the **adequacy and stability** of our multivariate model.

Regarding the RAAEC-C instrument, we have expanded the Methods section to describe its development and validation process more clearly. The RAAEC-C instrument was developed for this study, based on established global protocols for childhood illnesses and respiratory assessment [28,29,30,31].

To ensure its validity and reliability for this specific population, a rigorous validation process was conducted. Its content validity was confirmed through an expert review, with the Content Validity Index (CVI) indicating satisfactory relevance and comprehensiveness.

Furthermore, its reliability was confirmed by assessing inter-rater agreement. This analysis was performed on the categorical diagnosis using Cohen's Kappa to confirm a high level of consistency among assessors. Additionally, the consistency of continuous measurements, such as BMI-for-age z-scores, and anthropometric indicators, was confirmed with the Intraclass Correlation Coefficient (ICC). These details are now included in the revised manuscript.

Comment 3: There are some areas where the language could be improved for conciseness, style, and grammar.

Response:

We thank the reviewer for this helpful comment. The manuscript has been carefully revised to improve conciseness, style, and grammar. Several sentences have been restructured for clarity, redundant wording has been removed, and minor grammatical errors have been corrected. We believe these changes have improved the readability and flow of the manuscript.

Comment 4: A higher prevalence in one area compared to another might depend on a fortuitous higher circulation of a virus. How did the authors account for this?

Response:

We thank the reviewer for raising this important point. We acknowledge that differences in RTI prevalence could in part be influenced by temporal variation in viral circulation, and this potential bias cannot be completely excluded. To minimize seasonal effects, data collection in both sites was conducted during the same season. Importantly, if viral circulation alone were the dominant explanation, one might expect a higher RTI prevalence in Pangalengan, which has a cooler climate, higher altitude, and lower absolute humidity but often higher relative humidity—conditions known to favor viral persistence and impair mucosal defenses. In contrast, our study found substantially higher RTI prevalence in Jakarta. This pattern suggests that while temporal variation in respiratory virus circulation remains a limitation, the large difference in PM2.5 exposure between Jakarta and Bandung is a more plausible driver of the observed association. We have clarified this point in the revised discussion.

Comment 5: Relevant recent studies on air pollution and respiratory infections are missing and not discussed: e.g. PMID: 40358039, PMID: 36282132, PMID: 3847052, PMID: 29940478. Despite on lower tract infections, such a studies should be considered to corroborate the hypothesis and interpret the data.

Response:

We thank the reviewer for pointing out the need to incorporate additional recent studies. We have now revised the discussion to include this broader literature. Specifically, several new studies corroborate our findings: **Comotti et al. (2025)** reported that air pollution exposure significantly increased hospitalization risk in infants with bronchiolitis (PMID: 40358039). Similarly, **Milani et al. (2022)** demonstrated in a prospective cohort that short-term exposure to PM2.5 and PM10 was associated with greater bronchiolitis severity, particularly RSV-related cases (PMID: 36282132). In a more recent case-control study, **Van Brusselen et al. (2024)** found that both short- and medium-term exposure to particulate matter and NO₂ were linked to increased bronchiolitis risk in infants (PMID: 38470521). Finally, **Carugno et al. (2018)** conducted a systematic review and meta-analysis that confirmed consistent associations between PM2.5, PM10, and NO₂ exposure and increased bronchiolitis hospitalizations across multiple studies (PMID: 29940478).

Together, these studies strengthen the biological plausibility and external validity of our results, situating our findings within a growing body of evidence that air pollution—particularly PM2.5—plays a major role in the burden of pediatric respiratory infections.

We believe these revisions fully address your concerns and significantly improve the quality and rigor of our manuscript.

We sincerely thank Reviewer 2 for the insightful and constructive comments. Your feedback has been invaluable in strengthening our manuscript, prompting us to clarify our methodology, refine our language, and more transparently address the study's limitations. We are confident that these revisions have significantly improved the scientific rigor and overall quality of our work. We believe our revised manuscript now more accurately and robustly presents our findings, and we hope it meets the high standards of this journal.


Sincerely,

Hari Krismanuel

Universitas Trisakti

hari_krismanuel@trisakti.ac.id

Screenshot manuscript dipublikasi di jurnal Scopus 1 di akun SINTA



Scopus

Search...

Filter Quartile

- Quartile 1
- Quartile 2
- Quartile 3
- Quartile 4
- No Quartile

Sort By

Year

Page 1 of 1 / Total Records : 4

Q1 Examining the effectiveness of Prostatic hyperplasia education on the level of participant's knowledge and awareness
Creator : Krismanuel H.
Plos One


Q1 The association between PM2.5 level and respiratory tract infections among children: A cross-sectional study
Creator : Krismanuel H.
Aims Public Health

Q1 Exploring genetic susceptibility to air pollution and its implications for disease risk and precision health: A scoping review
Creator : Krismanuel H.
Aims Public Health

Journal publish at 2025 0 cited

Journal publish at 2025 0 cited

Journal publish at 2025 0 cited



Please Contact Verificator to Sync Publication Scopus

PUBLICATION

Scopus

Search...

Filter Quartile

- Quartile 1
- Quartile 2
- Quartile 3
- Quartile 4
- No Quartile

Sort By

Year

Page 1 of 1 / Total Records : 2

Q1 The association between PM2.5 level and respiratory tract infections among children: A cross-sectional study
Creator : Krismanuel H.
Aims Public Health

Q1 Examining the effectiveness of Prostatic hyperplasia education on the level of participant's knowledge and awareness
Creator : Krismanuel H.
Plos One

Journal publish at 2025 0 cited

Journal publish at 2025 0 cited