



LAMPIRAN I : Cover Journal of Drug Delivery & Therapeutics



LAMPIRAN II : Daftar Isi Edisi Terbit Manuskrip Publikasi “*Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years*”



Journal of
DRUG DELIVERY & THERAPEUTICS
ISSN 2256-1177


Home About Current Archives Guidelines Editorial Board Indexing Contact Search

HOME | ARCHIVES | Vol. 13 No. 7 (2023): Volume 13, Issue 7, July 2023

Vol. 13 No. 7 (2023): Volume 13, Issue 7, July 2023

PUBLISHED: 2023-07-15

Register Login



On the ISSN: 2256-1177

MAKE A SUBMISSION

Research

PREVENTIVE EFFECT OF TOTAL ETHANOLIC EXTRACT OF BUTEA MONOSPERMIA ON BONE LOSS IN OSTEOPENIC RATS
Abstract Display: 654 | PDF Downloads: 685 | HTML Downloads: 44
Soni Khanna, Anandap Laxmi, Pooja Gupta, Kishu Sharma, Ramesh Maurya, Nabakya Chatterjee, Divya Singh 6-14
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jddt.v13i7.8887>

FORMULATION AND EVALUATION OF BUCCODHESIVE TABLET OF VERAPAMIL HYDROCHLORIDE FOR THE TREATMENT OF HYPERTENSION
Abstract Display: 512 | PDF Downloads: 608 | HTML Downloads: 104
Suman Sukantia Tripta, Anil V Sawale, Shashtha Dilip Mureshwar, Ghanshi A Gulkare, Sakshi S Gulkar 15-21
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jddt.v13i7.8133>

PHENOLIC PROFILE AND ANTIOXIDANT POTENTIAL OF AQUEOUS EXTRACTS OF SELECTED TRADITIONAL ANTI-COUGH PLANTS
Abstract Display: 723 | PDF Downloads: 670 | HTML Downloads: 174
Okin Lilian Orangau, Chid Uzoma Igwe, Chiamaka Paryatus Nwabude, Oyinlola Chikoko Njoku, Juliet Chiona Ejiro 23-29
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jddt.v13i7.8136>

EMERGING PROBLEM EVALUATION OF HYPOCALCEMIA IN SUPER VASOL HAIR DYE POISONING PATIENTS: CLINICAL STUDY IN BIMS, KADAPA
Abstract Display: 673 | PDF Downloads: 512 | HTML Downloads: 292
K. Leela Prasad Babu, Shakti Kowamulla, M. Ramani, L. Nagarani, B. Abhinav, K. Sravathi Bai 30-33
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jddt.v13i7.8138>

CANI MORINGA OLEIFERA LEAF ETHYL ACETATE EXTRACT INHIBIT CANDIDA ALBICANS PLANKTONIC CELL GROWTH AND BIOFILM FORMATION IN VITRO?
Abstract Display: 629 | PDF Downloads: 767 | HTML Downloads: 100
Harneet Sahayani, Shilpa Sankar, Dewal Rameshwar, Dewal Agastha 34-37
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jddt.v13i7.8139>

EXPLORING DRUG PRESCRIPTION PATTERNS IN THE NEUROLOGY DEPARTMENT OF A TERTIARY CARE HOSPITAL: A COMPREHENSIVE ANALYSIS
Abstract Display: 679 | PDF Downloads: 682 | HTML Downloads: 235
R. Nitya, Seleny, K. Neetha, UDAYAKUMAR Narmatha, R. R. Badi, A. Ranjya, B. Malathi 38-45
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jddt.v13i7.8139>

FORMULATION AND EVALUATION OF POLYHERBAL TABLET BY USING NEEM, TULSI, TURMERIC AND GINGER EXTRACT
Abstract Display: 2373 | PDF Downloads: 1394 | HTML Downloads: 2136
Souvik Paul, Tanmay Dey, Prabir Koleka, Siddhar Tanay, Soumik Bhattacharya, Rajar Das 46-51
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jddt.v13i7.8889>

FACTORS ASSOCIATED WITH PLASMA MALONDIALDEHYDE LEVELS IN PEOPLE OVER 40 YEARS
Abstract Display: 1089 | PDF Downloads: 840 | HTML Downloads: 62
Majani Mejan, Tohana, E'elina Margo, Juli Chuti, Pucahri, Meida Alfa Faradisa 52-56
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jddt.v13i7.8142>

ANALYSIS OF THE FACTORS THAT INFLUENCE THE FAMILY IN THE USE OF TRADITIONAL MEDICINE AS SELF MEDICATION IN SUR TEBUK II DISTRICT
Abstract Display: 603 | PDF Downloads: 617 | HTML Downloads: 57
Habun Niza, Saah-Harrah-Az-Zahra Anwar, Wahyudi Wahyudi 57-61
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jddt.v13i7.8148>

EVALUATION OF THE POTENTIAL FOR MEDICATION ERROR IN OUTPATIENT ELDERLY PATIENTS AT SITI FATMAH AZ-ZAHRA REGIONAL HOSPITAL
Abstract Display: 661 | PDF Downloads: 704 | HTML Downloads: 81
Habun Niza, Saah-Harrah-Az-Zahra Anwar, Sindil Ubani 62-64
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jddt.v13i7.8149>

STRUCTURE-BASED DRUG DESIGN, ADMET AND MOLECULAR DOCKING ANALYSES OF ANTI-VIRAL AGENTS AGAINST SARS-COV-2 VIRUS, ZIKA VIRUS AND HEPATITIS C VIRUS
Abstract Display: 711 | PDF Downloads: 616 | HTML Downloads: 77
Peer Sitis, Georgiade Zeleng, Mulyati Lohi, Arifin Hekika, Alimulrahman Zahir 65-71
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jddt.v13i7.8890>

POTENTIAL ACTIVITY OF STEVIA REBAUDIANA BERT. IN INHIBITING CYCLOOXYGENASE AND LIPOXYGENASE ENZYMES AS ANTI-INFLAMMATORY CANDIDATES: A MOLECULAR DOCKING STUDY AND ADMET PREDICTION
Abstract Display: 511 | PDF Downloads: 651 | HTML Downloads: 67
Maida Adilia, Aif Setiawanayath, Rina SC Sitindon 75-86
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jddt.v13i7.8137>

PHYTOCHEMICAL ANALYSIS OF DIFFERENT EXTRACT OF HYPTIS SUAVEOLENS(L.)POIT
Abstract Display: 703 | PDF Downloads: 840 | HTML Downloads: 482
Saurav Gokhale, Sonali Deokar, Akanksha Adarkar 87-92
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jddt.v13i7.8816>

CATEGORIES

- Aims & Scope
- Editorial Policies
- Publication Ethics
- Plagiarism Policy
- Retraction Policy
- Open Access Policy
- Pricing Charges and Waiver Policy
- Indexing

LATEST PUBLICATIONS

- 100% PDF
- 100% HTML
- 100% PDF

MOLECULAR DOCKING ANALYSIS OF PIPERONAL AND ITS ANALOGUES AS PROMISING CANCER THERAPEUTICS BY MODULATING ANGIOGENESIS

Abstract Display: 432 | PDF Downloads: 472 | HTML Downloads: 351
Anusha Venkatesan, Ashish Prakash Karurkar, Saeed Al Mohamed Yacoub, Sarika Mathanaiselvan 93-98
[VIEW ABSTRACT](#) [PDF](#) [HTML](#)
[DOWNLOAD PDF](#) [DOWNLOAD HTML](#) <https://doi.org/10.22075/jghe.v13i7.2817>

FORMULATION DEVELOPMENT AND EVALUATION OF NEW POLYHERBAL GEL FORMULATIONS FOR THEIR WOUND HEALING ACTIVITY IN RAT

Abstract Display: 1107 | PDF Downloads: 719 | HTML Downloads: 132
Yadavendra Dutt Shukla, Karunakar Shukla, Sakshith Kumar Jaiswal 99-105
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jghe.v13i7.2803>

COMPARISON OF ANTIBIOTIC-RESISTANT PATTERN OF EXTENDED SPECTRUM BETA-LACTAMASE AND CARBAPENEM-RESISTANT ESCHERICHIA COLI ISOLATES FROM CLINICAL AND NON-CLINICAL SOURCES

Abstract Display: 703 | PDF Downloads: 606 | HTML Downloads: 56
Biju Sundaraj Joseph, Jeema Craylyne Okalo, Evangelina Chinyere Ukwuezo, Chinyere Emelita Nwankwo, Ikennedy Udeme 107-118
Peer-Review: Anura Prasad Ojha, Ranvijay Kumar, Gopikrishna Srinivas
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jghe.v13i7.2818>

ISOLATION, CHARACTERIZATION AND ANTIFERTILITY ACTIVITY OF ETHANOLIC EXTRACTS OF TABERNAEMONTANA SYMPLICATA (L.) BR. EX ROEM. & SCHULT

Abstract Display: 1201 | PDF Downloads: 645 | HTML Downloads: 143
Anshu Khatun, Suresh Jain, Rajendra S. Rajan 119-126
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jghe.v13i7.2807>

FORMULATION AND EVALUATION OF MICRO BEADS OF CLONIDINE FOR THE TREATMENT OF ANXIETY AND HYPERTENSIVE DISORDER

Abstract Display: 4611 | PDF Downloads: 615 | HTML Downloads: 235
Aruna Deviand, Nageswara Mohanraj, Mahima Dubey, Anjali Namdeo 127-133
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jghe.v13i7.2820>

Review

A COMPREHENSIVE REVIEW ON EFFERESCENT TABLETS

Abstract Display: 6456 | PDF Downloads: 4171 | HTML Downloads: 1523
Kam Gopinath Varshni, Deepjyoti Varshni, Saradeep Bhattacharya, Sarika Lakshmi Narasimhan 141-150
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jghe.v13i7.2120>

A COMPREHENSIVE OVERVIEW ON PHYTOMEDICINES AS AN UPCOMING/EMERGING CANDIDATE FOR THE MANAGEMENT OF EPILEPSY

Abstract Display: 718 | PDF Downloads: 587 | HTML Downloads: 91
Priya Kumari, Pawan Kumar, Mahendra Singh Anand, Pankaj Sharma 151-160
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jghe.v13i7.2128>

A SYSTEMATIC REVIEW ON COMPARISON OF DEEP BRAIN STIMULATION (DBS) AND TRANSCRANIAL MAGNETIC STIMULATION (TMS) IN THE TREATMENT OF REFRACTORY OBSSIVE COMPULSIVE DISORDER (OCB)

Abstract Display: 616 | PDF Downloads: 599 | HTML Downloads: 221
K Divya, Bibha Samal 161-171
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jghe.v13i7.2126>

AN DETAILED OVERVIEW ON MOUTH DISSOLVING FILM

Abstract Display: 1622 | PDF Downloads: 4089 | HTML Downloads: 148
Priyanka Jain, Ashish Gupta, Gajanan Dawahekar 172-176
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jghe.v13i7.2121>

THERAPEUTIC ASPECTS OF BIOLOGICALLY POTENT VANILLIN DERIVATIVES: A CRITICAL REVIEW

Abstract Display: 1535 | PDF Downloads: 1743 | HTML Downloads: 300
Raju Senthil Kumar, Sundhararajan Naveena, Sekar Praveen, Nagalingam Yogathasabhi 177-189
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jghe.v13i7.2129>


REVIEW ON GREEN CHEMISTRY

Abstract Display: 1062 | PDF Downloads: 1532 | HTML Downloads: 2017
Tapan Wale, Shikhar Surjansh, Vinod Havan, Pallavi Phalke, Manisha Ghoshale 190-193
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jghe.v13i7.2819>

SACCOPETALUM TOMENTOSUM: REVIEW OF ITS BOTANY, MEDICINAL USES, PHARMACOLOGICAL ACTIVITIES AND PHYTOCHEMISTRY

Abstract Display: 572 | PDF Downloads: 538 | HTML Downloads: 98
Poojabharti Kumar, Karunakar Shukla 194-199
[VIEW ABSTRACT](#) [PDF](#) [DOWNLOAD PDF](#) [HTML](#) <https://doi.org/10.22075/jghe.v13i7.2829>

LAMPIRAN III : Editorial Team



HOME Editorial Board

Editorial Board

EDITOR-IN-CHIEF

Dr. Ram C Dhalwar, Hospital Pharmacy, SRG Hospital and Medical College, Jalgaon, India-320001, India. Email: rdc@rcdhalwar.com; Phone: +91-678380207 [Google Scholar](#) [ResearchGate](#) [ORCID](#) [SCOPUS](#) Research fields: Novel Drug Delivery, Nanotechnology, Microspheres, TDDS

ASSOCIATE EDITORS

Dr. Aamir Jalil Al-Mozawil, Advisor And Expert Trainer, Baghdad Medical City And The National Training And Development Center, Iraq Ministry Of Health, Baghdad, Iraq. [ResearchGate](#) [Google Scholar](#) [SCOPUS](#) [ORCID](#) [NCBI-NLM](#) Email: amozawil@yahoo.com Research Fields: Pediatrics, pediatric psychiatry, nephrology, pediatric neurology, genetics

Dr. S K Prajapati, Professor & Head, Institute of Pharmacy, Bundelkhand University, Jhansi (U.P.), India, India. Email: skprajapati@gmail.com [ResearchGate](#) [Website](#) [Google Scholar](#) Research Area: Novel Drug Delivery, Microspheres, Nanoparticles

Dr. Rajesh Ramasamy, Associate Professor, Head of Department, Department of Pathology, Faculty Medicine and Health Science, Universiti Putra Malaysia, Malaysia. Phone: +603 8947 2377 Mobile: +6012 383 6444 Fax: +603 8941 2787 Email: rajesh@upm.edu.my [Google Scholar](#) [ResearchGate](#) [Website](#) [SCOPUS](#) ID: 26024447300 Research Fields: Stem Cells, Immunology, Diabetes, Medicinal Plant Research, Siddha Medicine

Dr. Syed Mohd Danish Rizvi, Associate Professor, Department of Pharmaceutics, College of Pharmacy, University of Hail, Hail, Saudi Arabia. Email: syedmohd19@gmail.com Official Email: sm.danish@uh.edu.sa | [ORCID ID](#) | [Google Scholar](#) | [Scopus](#) | [Research Gate](#) | [Website](#) | Research fields: Microspheres, Nanoparticles, Cancer Therapy

SECTION EDITORS (Scientific Committee Members)

Dr. Narendra Dudhipala, Department of Pharmaceutics and Drug Delivery, School of Pharmacy, University of Mississippi, USA. Research Interests: Formulation development; preclinical evaluation of small molecules for ocular delivery and oral delivery; Tumor delivery; lipid nanoparticles; lyophilization. Mobile: +1 261 818 5548 Email: narendrad@com.olemiss.edu - [View Profile](#) [ResearchGate](#)

Dr. Priyanka Dhari, Department of Pharmaceutical Sciences, College of Pharmacy, University of South Florida, 12801 Bruce B. Downs Blvd., MDC03, Tampa, FL 33613, Email: pdhari@health.usf.edu [Google Scholar](#) [ResearchGate](#) [Website](#)

Dr. Siddhartha Paul, Research Associate, Centre of Excellence, Khulna University, Odisha, India. +91-884577507006251379 psiddhartha@gmail.com [ResearchGate](#) [ORCID](#)

Dr. Iqbal A. Ansari, Assistant Professor, Department of Pharmaceutical Chemistry, Prof. Ravindra Nandan College of Pharmacy, Gandhinagar, Dhule, 424002, Maharashtra, India. Mobile: No-91 9284541386. Email: ansariiqbal@gmail.com [Google Scholar](#) [ResearchGate](#) [Website](#) Research Fields: EGFR Inhibitors, CADD, Docking, MD Simulation

Dr. Praveen Doodipati, Carver College of Medicine, University of Iowa, USA. Research Fields: Molecular Biology - Immunology Cardiovascular Biology [ResearchGate](#) [Website](#) [ORCID ID](#) Email: praveend@iastate.edu

Dr. Ashok K Dasgupta, Assistant Professor, National Institute of Pharmaceutical Education and Research, Rohtak, Uttar Pradesh, 200002, India Email: ashokdasgupta@gmail.com [Google Scholar](#) [ResearchGate](#) [Website](#) Research Interests: Neuroscience, Pharmacology and Toxicology

Dr. Dinesh Nyanandi, Formulation Scientist, Continuous Pharmaceuticals, 25 Olympia Ave, Woburn, Massachusetts, USA. Dhoni - (P) +1(603) 555-6555 (E) ndinesh64@gmail.com; dnyanandi@continuouspharma.com [ResearchGate](#) [Scopus](#): Oral drug delivery, Lipid based drug delivery, Strategies for enhancement of solubility and bioavailability, Gastro-intestine Drug delivery, Ocular Drug Delivery, Topical Drug Delivery [Website](#) [Google Scholar](#) [ORCID](#) [SCOPUS](#)

INTERNATIONAL EDITORIAL MEMBERS

Prof. Dr. nec. Nec. Ibrahim A. El-Dahmy, Director, Institute of Bioproduct Development (IBD), Universiti Teknologi Malaysia (UTM), 81010 UTM, Johor Bahru, Malaysia. Tel: 006-07-5511573 (office), Fax: 006-07-5589706, E-mail: ibrahima@ibd.umy.ac.my [ResearchGate](#) [Google Scholar](#)

Dr. Stanisz Selcher Narda, Department of Chemistry, Myongji University, Yongin, South Korea. Email: staniszselcher@gmail.com [Google Scholar](#) [ResearchGate](#) [Website](#)

Dr. Kishor Wazirunde, Associate Professor at Jashon University of Science and Technology, Chattogram, Bangladesh Email: wazirunde2003@yahoo.com wazirunde@juast.edu.bd [LinkedIn](#) [Google Scholar](#) [ResearchGate](#) [ORCID ID](#)

Dr. Priya Dhawan, Analytical Scientist, Continuous Pharmaceuticals, MA, USA. [Google Scholar](#) [ResearchGate](#) [Website](#) Email: pdhawan@continuouspharma.com Interests fields: Carbohydrates, synthesis, Chemical biology, Supramolecular chemistry, Analytical method development and qualifications of drug molecules, Nanomaterials, and organic chemistry

Dr. Ahmed Zaib Sahib, Professor, Dept of Pharmacology & Toxicology, College of Pharmacy, University of Kerbala, Iraq. Email: ahmedzsaib73@yahoo.com ahmedzsaib@yahoo.com Tel : +961-781-385576 [Google Scholar](#) [ResearchGate](#) [Website](#) Interests fields: Pharmacology, Clinical Pharmacology, Pharmacogenomics

Prof. Fahad I. Al-Salkhan, Department of Clinical Pharmacy, Prince Sultan Bin Abdulaziz University, College of Pharmacy, Kingdom of Saudi Arabia. Email: fahad@psu.edu.sa Telephone: +96615366274 [ResearchGate](#) [Website](#) Interests fields: Primary care/inpatient care, health outcomes, pharmacogenomics

Dr. Najih Al-Din Al-Jamal, Professor, Department of Clinical Pharmacy, Malaysian Academy of Pharmacy, Malaysia. [ResearchGate](#) [ORCID](#) [Website](#) [LinkedIn](#) [Google Scholar](#) [Official website](#) Email: aljamal1977@yahoo.com Research interest: Pharmaceutics (drug delivery; nanotechnology); Pharmaceutical Analysis (method development and validation); Clinical Pharmacy; Pharmacy Education; Pharmacy Practice & Public Health.

Dr. Vijai Kumar Reddy Tangadanchu, Ph. D. Postdoctoral Research Associate, Mallekrodt Institute of Radiology, Washington University School of Medicine, Saint Louis, Missouri-63110, USA. E-mail: vijai.kumar@wustl.edu vijai@imr.wustl.edu Mobile: +1 314-665-8974. [ResearchGate](#) [Publons](#) [SciProfiles](#)

Dr. Soheila Ghafeeli, Department of Pharmaceutics, Faculty of Pharmacy, Islamic Azad University, Tehran, Iran; Email: soheila@iaui.ac.ir [Google Scholar](#) [ResearchGate](#) Tel: +9891252721 Fax: +989124880081

Dr. Mostafa Madrolli, Emergency Medical Technicians, Dezful University of Medical Sciences, Dezful, Iran. Email: mostafamadrolli@yahoo.com Phone: [968370170590] Interests fields: Diabetes, Alzheimer's Disease.

Dr. Vijay Kumar Siripuram, Laboratory of Molecular and Cellular Neuroscience The Rockefeller University, New York, NY, USA. Email: vijay.kumar@rockefeller.edu ; Phone: (+1) 646-238-5620. Interests fields: NMR, FTIR, HPLC, GC/MS/LCMS, column chromatography, crystallization [Google Scholar](#) [Publons](#)

Dr. Sakar Patel, Research Associate at Memorial Sloan Kettering Cancer Center, Department of Radiology, New York, 10005, USA. Interests fields: Organic chemistry, new molecule design, developing new methods for hypoxia reduction via dissolution DNP and real time metabolism (in vivo/in vitro) Phone: +1 929-553-4660 Email: sakarpatel13@gmail.com sakar13@mskcc.org [Google Scholar](#) [Website](#)

Dr. Arun Suresddy, Ph.D., Scientist, Protein Pharmaceutical Development, Regen Inc, Massachusetts, USA. [ResearchGate](#) [Google Scholar](#) [Website](#) Email: suresddyarun@gmail.com Mobile: +1 (647)-360-4558

Dr. Madhukiran Parvathani, Manager - Global Regulatory Affairs, CRC Pharma, New Jersey, USA. Interests Fields: Regulatory Affairs, Clinical Research, Pharmaceuticals and Medical Devices. Ph: +1 (908)662-6496 Email: madhukiran.parvathani@gmail.com [Website](#) [ResearchGate](#) [Google Scholar](#) [LinkedIn](#)

Praveen Kalini, Department of Pharmaceutics and Drug Delivery, University of Mississippi, Faxon Hall, MA, 38077, USA. Mail: praveenkalinipani@gmail.com [Google Scholar](#) [ResearchGate](#) [LinkedIn](#)



Online ISSN: 2256-1177

MAKE A SUBMISSION

CATEGORIES

- [Aims & Scope](#)
- [Editorial Policies](#)
- [Publication Ethics](#)
- [Page/Issue Policy](#)
- [Retraction Policy](#)
- [Open Access Policy](#)
- [Processing Charges and Waiver Policy](#)
- [Indexing](#)

LATEST PUBLICATIONS

[View All](#)

[New Article](#)

[New Review](#)

ASSISTANT EDITORS

Dr. Dito Anarogo, M.D., M.Sc., International PhD Program for Cell Therapy and Regeneration Medicine (IPCTRM), College of Medicine, Taipei Medical University Taiwan and Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Makassar, Makassar, Indonesia. Mail: dito.anarogo@ipctrm.tmu.edu.tw or dito110200@ipctrm.tmu.edu.tw Phone: +62-81224122660. [Google Scholar](#) [Publons](#) [Linkedin](#) [Research Interest](#): drug and delivery, pharmacology, regenerative medicine, oncology, neurosciences, molecular biology & medicine, bioinformatics, nanomedicine.

Dr. Jacopo Forte, Fixed-term Researcher (RTD-A) at the Department of Basic Biotechnological Sciences, Catholic University of the Sacred Heart, Rome. [Research Interest](#): drug delivery like nanosystems for anticancer drug delivery, vesicular systems based on phospholipids and surfactants suitable for antibacterial drugs, and nanocarriers for targeted delivery to the bladder. [ResearchGate](#) [SciProfiles](#) [Website](#) [ORCID](#) [Email](#): jacopo.forte@uniroma2.it

Dr. Alimtar Rizkielena Ahmad, Faculty of Pharmacy, Universitas Muallim Indonesia, Ulp Samudra Km. 5, 80331, Indonesia. [Email](#): alimtarizkielena@uim.ac.id [Phone](#): +6281355184454 [Google Scholar](#) [ResearchGate](#)

Dr. Smita Datta Maurya, Sardar Vallabhbhai Patel College of Pharmacy, Azamgarh, India. [Email](#): smita1993@gmail.com [Google Scholar](#) [ResearchGate](#) [Interested fields](#): Pharmacy (Novel Drug delivery, Liposomes, Ethosomes.)

Dr. Sandeep K. Singh, Postdoctoral Research Associate, Department of Pharmacy Practice and Science, University of Nebraska Medical Center, Omaha, NE, USA. [Google Scholar](#) sasingh10@gmail.com

Dr. Mousumi Kar, Professor and Head, Faculty of Pharmacy Medi-Caps University, Indore, INDIA, India. [Email](#): karmousumi@hotmail.com [Website](#) [Google Scholar](#) [ResearchGate](#) [Linkedin](#) [Website](#) [Interested fields](#): Pharmaceutics, Novel Drug Delivery

Dr. Sachin Goyal, Research Associate, Department of Anesthesiology and Critical Care medicine, School of medicine, University of New Mexico, Albuquerque, NM 87106 USA. [Email](#): sgoyal@salud.unm.edu

Mr. Vaidhar Patel, Director (Quality Assurance), Florida Oncology Inc., 1 Deer Park Drive, Suite E Monmouth Junction, NJ 08852, USA. [Website](#) [Email](#): vpatel0213@gmail.com [Google Scholar](#) [Phone No](#): (571) 208-5652

Mr. Girish Nihalani, Research Fellow of Drug Product Development at Teva Pharmaceuticals Inc. (Active) based at Florida, USA. [email](#): gnihal73@gmail.com gnihal73@gmail.com [Contact](#): +1- 854-647-7763 [LinkedIn](#) [Google Scholar](#) [Interested fields](#): Pharmaceutics, Novel Drug Delivery

Dr. Rajeshwar Patel, Ph.D. (Pharmaceutics), Senior Scientist, Product Development Granules Pharmaceuticals Inc, 3701 Concorda Play, Chantilly, Virginia 20151, USA. [Mobile](#): +1 571 742 9296 [Email](#): rajeshwarpatel@gmail.com [Google Scholar](#) [ResearchGate](#) [ORCID](#)

LAMPIRAN IV : Manuskrip Publikasi “Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years”

← Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years

Available online on 15.07.2023 at <http://jddtonline.info>

Journal of Drug Delivery and Therapeutics

Open Access to Pharmaceutical and Medical Research

Copyright © 2023 The Author(s). This is an open-access article distributed under the terms of the CC BY-NC 4.0 which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited

Open Access Full Text Article

Research Article

Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years

Meiyanti*, Yohana, Eveline Margo, Juni Chudri, Puspardini, Meutia Atika Faradilla

Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Article Info:

Abstract

Article History:

Received 08 May 2023
Reviewed 11 June 2023
Accepted 24 June 2023
Published 15 July 2023

Cite this article as:

Meiyanti, Yohana, Margo E, Chudri J, Puspardini, Faradilla MA. Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years. Journal of Drug Delivery and Therapeutics. 2023; 13(7):52-56

DOI: <http://dx.doi.org/10.22270/jddt.v13i7.6142>

*Address for Correspondence:

Meiyanti, Department of Pharmacology and Medical Pharmacy, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Aging is a progressive process caused by a physiological decline in body functions. It is known that several factors can accelerate aging, namely age, lifestyle, oxidative stress, lipid peroxidation, and mitochondrial damage. Various pollutants such as cigarette smoke, motor vehicle fumes, industrial waste, and exposure to sunlight / ultraviolet light will form free radicals. Malondialdehyde (MDA) level is one of the indicators used to measure oxidative stress. Increased MDA levels are found in various degenerative diseases. This study aims to determine plasma MDA levels in the age group of 40 years and over and factors that can affect plasma MDA levels. This study was a cross-sectional study by including 42 subjects. Data collection was carried out using a questionnaire to obtain subject demographic data and physical activity, as well as physical examinations complemented by laboratory examinations and MDA levels. The mean age of the subjects was 52.7 years, and most of the 32 (76.2%) subjects were categorized as obese and had high activity. Pearson correlation test obtained body mass index statistically significantly correlated with plasma MDA levels with a value of $p = 0.039$, as well as physical activity statistically significantly correlated with a negative correlation, with a value of $r = -0.26$ and $p = 0.043$. Conclusion Body mass index and activity are significantly correlated with plasma MDA levels.

Keywords: malondialdehyde, stress, oxidative, radicals

← Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years

INTRODUCTION

Aging is a progressive process caused by a physiological decline in body function. ¹ A decrease in body homeostasis is accompanied by an increased risk of various chronic diseases including diabetes mellitus, cancer, neurodegenerative diseases, and heart disease. It is known that several factors can accelerate aging, namely age, lifestyle, oxidative stress, lipid peroxidation, and mitochondrial damage. Previous studies have shown that changes in sphingomyelin metabolism in the central nervous system and age are associated with neurodegenerative diseases. ² In the US, aging is one of the most important factors in diabetes mellitus and heart disease. In 2015, an estimated 12% of the population had diabetes mellitus and an undetected 24% had diabetes mellitus. Heart disease is the highest contributor to mortality in the population aged 65-84 years. ³ Recent studies have shown that non-diabetic populations whose fasting glucose levels increase in the blood will increase the risk of coronary heart disease. But on the other hand, the data on fasting blood glucose increase is still debated in the population of cardiovascular disease patients who do not have diabetes mellitus. A recent study in North Sumatra showed a significant relationship between MDA levels in obese and non-obese populations. ⁴

Free radicals are formed through two processes, namely exogenous and endogenous processes. Various pollutants such as cigarette smoke, motor vehicle fumes, industrial waste, and exposure to sunlight/ultraviolet will form free radicals. Illness, stress, and excessive exercise/physical activity are endogenous factors produced by the human body that can form free radicals. The formation of free radicals occurs due to oxidation processes and metabolic processes that take place during exercise or activity, inflammatory processes, consumption of certain foods, and others. Free radicals will begin to activate with molecules in the surrounding cells to obtain electrons to stabilize the condition of one's body. The electrons that have been taken in the cell molecules will trigger the occurrence of free radicals. If this continues for a long time, it can cause oxidative stress which can cause inflammation, and DNA disease can also trigger the disease. Oxidative stress conditions characterized by increased production of Reactive Oxygen Species (ROS) have implications for various diseases such as hypertension, atherosclerosis, diabetes mellitus, stroke, chronic renal failure, heart failure, and other chronic diseases. ⁵⁻⁷

Oxidative stress is an imbalance between oxidants (free radicals) and antioxidants. Free radicals that commonly cause oxidative damage are reactive oxygen species (ROS). ⁸ ROS compounds are formed through aerobic metabolism. These compounds can cause extensive damage through catalytic reactions. They are responsible for damage at the cellular level by oxidizing proteins, lipids, DNA (Deoxyribonucleic acid), and other biomolecules. Various compounds can be used as markers of oxidative stress such as malondialdehyde (MDA), and protein carbonylation. 4-hydroxy-2-nonenal (4HNE), and 8-hydroxy-deoxyguanosine (8OHdG). ⁹

Malondialdehyde (MDA) is a result of lipid peroxidation in the body. The concentration of MDA indicates an oxidation process that occurs in the body's cell membranes due to metabolic processes. One of the factors that cause aging is the production of excess reactive oxygen species (ROS). ROS are generated through the process of cell respiration in the mitochondria and the phagocytosis process of foreign body elimination. To stabilize the amount of ROS, the body can use endogenous and exogenous antioxidants. Disruption of the balance between ROS and antioxidants can cause irreversible damage to cell organelle components up to the DNA level. ^{10,11}

MDA levels are strongly influenced by many factors such as age, illness, stress, various pollutants, food consumption, antioxidant supplements and excessive physical activity or exercise, and others. The other study found that more than 30% of subjects had high MDA levels, and concluded that there was a significant relationship between eating habits, body mass index, glucose levels with MDA levels, as well as smoking habits significantly related to MDA levels. ¹² Recent studies have shown that plasma MDA levels in populations aged 70 years and 60 years are not significantly different but there is an increase in the specific activity of the catalase enzyme that increases with age. ¹⁰ In another study, there was a significant difference in plasma MDA levels in the obesity group without metabolic syndrome compared to the obesity group with metabolic syndrome. ⁴

There are still variations in MDA data in various groups with different ages and different disease groups, so this study aims to determine plasma MDA levels in the age group 40 years and over and factors that can affect plasma MDA levels.

Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years

METHOD

This observational analytic study with a cross-sectional design approach was conducted from November 2022 to February 2023 in Angke village, West Jakarta, Indonesia. The inclusion criteria for this study were: men and women aged ≥ 40 years, willing to participate in the study and sign informed consent. Meanwhile, the exclusion criteria were having a history of malignancy, kidney failure, a history of autoimmune disease, taking drugs related to psychiatric conditions, and having limitations in mobility. This study used the calculation of sample size with the formula $N / 1 + N \cdot \alpha^2$, and obtained the minimum number of samples required was 40 subjects. Demographic data and physical activity were obtained from distributed questionnaires. Blood chemistry examination (complete blood, glucose, total cholesterol, liver enzymes (SGOT and SGPT) using an enzymatic method. Examination of malondialdehyde levels was carried out using the spectrophotometric method. This research has obtained approval from the ethics committee of the Faculty of Medicine, Trisakti University, Number: 178/KER/FK/1X/2022.

RESULTS

A total of 42 subjects participated in this study, most of the subjects were female 27 (64.3%), and the mean age of the subjects was 52.7 years. Anthropometric data of the subjects are as follows body weight 67.53 ± 13.64 and height 156.02 ± 8.73 cm, while for abdominal circumference the average is 93.09 cm. Most of the subjects 59.5% fell into the normotensive category, with an average systolic blood pressure of 141.62 mmHg and an average diastolic blood pressure of 82.45 mmHg. A total of 32 (76.2%) subjects were categorized as obese with BMI ≥ 25.0 , and most subjects had high activity levels. (Table 1)

Table 1: Distribution of Subject Characteristics (n=42)

Characteristics	n (%)	Mean \pm SD
Gender		
Male	15 (35.7)	
Female	27 (64.3)	
Blood pressure (mmHg)		
Normal (< 140/90)	25 (59.5)	
Hypertension ($\geq 140/90$)	17 (40.5)	
Physical Activity		
Low (< 600 Mets)	11 (26.2)	
High (≥ 600 Mets)	31 (73.8)	
Body Mass Index (kg/m²)		
Not obese (< 25.0)	10 (23.8)	27.69 \pm 4.90
Obese (≥ 25.0)	32 (76.2)	
Age (years)		52.7 \pm 10.60
Body weight (kg)		67.53 \pm 13.64
Height (cm)		156.02 \pm 8.73

Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years

Abdominal circumference (cm)	93.09 \pm 10.54
Blood pressure (mmHg)	
Systole	141.62 \pm 20.77
Diastole	82.45 \pm 12.70

The examination of the subject's plasma malondialdehyde levels obtained a mean of 0.77 with the highest value of 1.03 and the lowest of 0.51. Hemoglobin levels averaged 13.16 g/dL. Fasting blood glucose level was 95.06 \pm 41.01 mg/dL and total cholesterol was 179.83 \pm 31.85 mg/dL. Laboratory examinations to determine liver function were carried out by measuring SGPT and SGOT levels, obtaining mean values of 24.21 U/L and 26.31 U/L respectively. (Table 2)

Table 2: Laboratory examination results of subjects (n=42)

Variable	Mean \pm SD
Hemoglobin (g/dL)	13.16 \pm 1.56
Fasting blood glucose (mg/dL)	95.06 \pm 41.01
Total cholesterol (mg/dL)	179.83 \pm 31.85
SGPT (U/L)	24.21 \pm 13.50
SGOT (U/L)	26.31 \pm 11.51
Plasma malondialdehyde	0.77 \pm 0.26

Determining the type of correlation test to be used by first conducting a data normality test. The normality test used was Shapiro-Wilk because the number of subjects was less than 50 subjects. The correlation test used to assess the correlation between blood glucose, total cholesterol, SGPT, SGOT levels, age, body mass index and physical activity with plasma MDA levels was the Pearson correlation test. The test results are presented in Table 3. Blood glucose, total cholesterol, SGPT, SGOT, and age did not have a significant correlation with MDA levels. Body mass index has a significant correlation with plasma MDA ($p = 0.039$), with a positive correlation direction, and the strength of the correlation is statistically weak with a value of $r = 0.396$ (< 0.4). The physical activity variable has a significant correlation with plasma MDA levels ($p = 0.043$) with a negative correlation direction but with a statistically weak correlation strength ($r = -0.26$).

Table 3: Correlation Test of Age, Gender, Glucose, SGOT, SGPT, BMI, Physical Activity with Plasma MDA Levels (n=42)

	Plasma MDA Level	
	r	p
Age	0.098	0.756
Gender	0.124	0.108

Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years

Glucose	0.330	0.835
SGOT	0.104	0.513
SGPT	0.008	0.614
Body Mass Index	0.396	0.039*
Physical activity	-0.260	0.043*

* p < 0.05

DISCUSSION

The level of oxidative stress can be measured using Malondialdehyde (MDA) concentration. MDA is the result of lipid peroxidation, which will cause cell damage. Oxidant compounds such as free radicals or non-radical species attack lipids that contain carbon-carbon double bonds, especially unsaturated fatty acids (PUFA). MDA is one of the biomarkers widely used to assess oxidative stress. The mean serum MDA in this study 0.77 was found to be lower than the previous study¹⁰, this is because the age difference of the subjects in this study was younger than in the previous study. In contrast, a study conducted in Iraq on hypertensive patients concluded that the increase in MDA was associated with an increase in age, gender, and duration of disease.¹³

Free radicals such as ROS are the etiology of most chronic diseases, as it is known that active free radicals are the result of metabolic reactions. Damage to body tissue cells occurs because active free electrons remain looking for other electrons to settle, if no other electrons are available then active free electrons will attack cells resulting in cell damage. Although free radicals naturally have benefits for the body, when the level of free radicals exceeds the normal value, it will cause harm to the body.^{11,12}

Measurement of oxidative stress can be done by measuring plasma malondialdehyde (MDA) levels. MDA is a secondary result of the lipid peroxidation process, producing various active compounds that will damage cells. The process of MDA formation through enzymatic and non-enzymatic processes. Currently, MDA is one of the biomarkers used to assess oxidative stress.¹⁴ Some previous researchers concluded that higher serum MDA was found in patients with hyperlipidemia. The condition of increased fat content in the body is in line with increased oxidative stress. Hyperlipidemia is associated with physical changes in cell membranes that facilitate the release of free radicals from the mitochondrial electron transport chain or activation of NADPH oxidase. In patients with diabetes mellitus (DM) there is also an increase in MDA levels, malondialdehyde interacts irreversibly and reversibly with proteins and phospholipids to have a profound effect, especially on the collagen of the cardiovascular system. MDA levels remain within normal limits due to the antioxidant effect, but in DM, this protective effect is impaired resulting in increased MDA levels.^{1,15}

Increased lipid peroxidation is considered a consequence of oxidative stress that occurs when the dynamic balance between prooxidants and antioxidants is disrupted. The statin class of drugs for hyperlipidemia therapy is proven to be anti-atherogenic, can improve endothelial function, inhibit vascular inflammation, and stabilize atherosclerotic plaques. In addition to statins, non-steroidal anti-inflammatory drugs (NSAIDs) can affect MDA concentration. Previous studies have found that NSAIDs have significant antioxidant properties to scavenge free radicals.^{16,17}

In this study, there was no significant correlation between age and MDA levels. Similar results were also obtained by other researchers, cell damage due to oxidative stress is not related to increasing age but is associated with habits, behavior patterns, and physiological changes due to age. Various habitual patterns such as smoking, fatty food habits, lack of activity, and inflammatory processes can lead to increased oxidative stress. The habit of taking vitamin C (ascorbic acid) has a beneficial impact due to its antioxidant effect so cell damage and DNA damage are lower in this group.¹⁸ MDA levels were significantly higher among all levels of smokers. MDA levels were lower in light smokers compared to heavy or moderate smokers. Higher vitamin C levels were found in the non-smoker group. All body tissues are susceptible to free radical damage, but based on location, the respiratory epithelium is particularly vulnerable to cigarette smoke. Oxygen radicals generated close to the cell membrane oxidize membrane phospholipids which will result in lipid peroxidation with the end products of this reaction being malondialdehyde, ethane, and pentane. This causes an increase in MDA levels in smokers.¹⁹ Another study conducted in Klaten on foundry workers concluded that type of work, body mass index and smoking habits did not affect MDA levels, but length of work and marital status were associated with MDA levels. Long and continuous exposure to metallic materials will result in the accumulation of these materials in the body and will increase the oxidative process of polyunsaturated fats in cell membranes and worsen the level of lipid peroxidase, characterized by an increase in MDA levels

Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years

as a reflection of the intensity and rate of lipid peroxidase reactions of damage to body cells.²⁰

Hardiani's study also concluded that there was no significant difference in plasma MDA levels in the age population, but increasing age along with an increase in catalase-specific activity.¹⁰ This increase in catalase-specific activity is a protective effect especially for elderly women to overcome high levels of oxidative damage. The role of endogenous antioxidant catalase is to suppress oxidative damage.^{18,21} Oxidative stress plays an important role in the incidence of cardiovascular diseases including coronary artery disease. Oxidative stress is considered an important mechanism for the development of cardiovascular diseases especially atherosclerosis. Hypertension, cardiac hypertrophy, and ischemia-reperfusion injury are three other major cardiovascular conditions for which gender is a risk factor. Gender is also associated with differences in oxidative stress. Under physiological conditions, women appear to be less susceptible to oxidative stress. This may be due to the antioxidant properties of estrogen, sex differences in NADPH-oxidase activity or other mechanisms yet to be determined. Sex hormones including estrogen are globally present with their receptors in the heart so sex hormones can exert effects on the cardiovascular system. Estrogen can improve heart function and reduce injury to the heart. In addition, estrogen has antioxidant effects so that it can function as a cardioprotective.^{22,23}

This study found a significant correlation between body mass index and MDA levels, the higher the BMI value, the higher the MDA levels. Similar results were also obtained in a study in India that concluded that in metabolic syndrome patients with one of the signs is an increase in BMI, there is an increase in MDA levels compared to the control group. In obese subjects, the level of MDA is higher than normal IMT. In a healthy body, there is redox homeostasis that occurs in cells, if an imbalance arises it will cause oxidative stress. Obese patients will have an increased metabolic load and mechanical load on the myocardium, increased body mass, and excessive food intake will cause the formation of lipid peroxidation, and free radicals.²⁴ The increase in some of these things will stimulate antioxidant enzymes in the long run, this will cause antioxidant enzyme reserves to be depleted and unable to overcome and balance the increase in oxidative stress.^{25,26}

These types of training activities are exhausting, long exercise, overtraining syndrome, various types of exercise will induce a significant response to oxidative stress. Exercise activities such as light exercise, low-intensity exercise, and prolonged training will have an impact on improving endogenous antioxidant status. Reactive oxygen species play an important role in cell signaling and in regulating antioxidant gene expression. Exertion results in hyperregulation of nuclear factor kappa B and mitogen-activated protein kinases that activate gene expression of a number of enzymes and proteins with important roles in maintaining intracellular oxidative or antioxidant homeostasis.^{1,6,27} The limitation of this study is that some factors that may affect MDA levels such as consumption of vitamin supplements, diet, and length of illness suffered were not examined in this study.

CONCLUSIONS

Body mass index and activity were significantly correlated with MDA levels, while age, gender, blood glucose levels, and liver enzymes were not significantly correlated with MDA levels.

Conflict of Interest:

The investigators declare there is no conflict of interest in this study.

Acknowledgments

The authors thank all the subjects who participated in this study, as well as the laboratory staff who assisted in the implementation of this study.

REFERENCES

1. Sharifi-Rad M, Anil Kumar N V, Zucca P, Varoni EM, Dini L, Panzarini E, et al. Lifestyle, Oxidative Stress, and Antioxidants: Back and Forth in the Pathophysiology of Chronic Diseases. *Front Physiol*. 2020; 11(July):1-21. <https://doi.org/10.3389/fphys.2020.00694>
2. Park MH, Jin HK, Bae J sung. Potential therapeutic target for aging and age-related neurodegenerative diseases: the role of acid sphingomyelinase. *Exp Mol Med [Internet]*. 2020; 52(3):380-9. <https://doi.org/10.1038/s12276-020-0399-8>
3. Całyniuk B, Grochowska-Niedworok E, Walkiewicz K, Kawecka S, Popiolek E, Fatyga E. Malondialdehyde (MDA) - product of lipid peroxidation as marker of homeostasis disorders and aging. *Ann Acad Medicae Silesiensis*. 2016; 70:224-8. <https://doi.org/10.18794/aams/65692>

← Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years

4. Savitra M, Rusdiana A, Widjaja SS, Syahputra M. Comparison Malondialdehyde (MDA) Level between Obesity Non Metabolic Syndrome and Obesity with Metabolic Syndrome Patients. 2020; (5):644-7.
5. Yekti R, Bukhari A, Jafar N, Thaha AR. Measurement of malondialdehyde (MDA) as a good indicator of lipid peroxidation. Int J Allied Med Sci Clin [Internet]. 2018; 6(4):6-8. Available from: www.ijamsc.com
6. Denise Mafra ME. Influence of Physical Exercise on Oxidative Stress and Inflammation in Hemodialysis Patients. J Mol Biomark Diagn. 2015; 06(03). <https://doi.org/10.4172/2155-9929.1000e120>
7. Mao C, Yuan JQ, Lv Y Bin, Gao X, Yin ZX, Kraus VB, et al. Associations between superoxide dismutase, malondialdehyde and all-cause mortality in older adults: A community-based cohort study. BMC Geriatr. 2019; 19(1):1-9. <https://doi.org/10.1186/s12877-019-1109-z>
8. Ayala A, Muñoz MF, Argüelles S. Lipid peroxidation: Production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. Oxid Med Cell Longev. 2014; 2014. <https://doi.org/10.1155/2014/360438>
9. Situmorang N, Zulham Z. Malondialdehyde (MDA) (Zat Oksidan Yang Mempercepat Proses Penuaan). J Keperawatan Dan Fisioter. 2020; 2(2):117-23. <https://doi.org/10.35451/kf.v2i2.338>
10. Hardiany NS, Sucitra S, Paramita R. Profile of malondialdehyde (MDA) and catalase specific activity in plasma of elderly woman. Heal Sci J Indones. 2020; 10(2):132-6. <https://doi.org/10.22435/hj.v1i2i2.2239>
11. Hardiany NS, Sadikin M, Siregar N, Wanandi SI. The suppression of manganese superoxide dismutase decreased the survival of human glioblastoma multiforme T98G cells. Med J Indones. 2017; 26(1):19-25. <https://doi.org/10.13181/mj.v26i1.1511>
12. 439-445_nova.pdf.
13. A. Hassan A, Sayyah SG. Oxidative Stress Marker Malondialdehyde and Glutathione Antioxidant in Hypertensive Patients. Eur J Biomed Res. 2023; 2(1):31-6. <https://doi.org/10.24018/ejbiomed.2023.2.1.47>
14. Khoubnasab Jafari M, Ansarin K, Jouyban A. Comments on "use of malondialdehyde as a biomarker for assessing oxidative stress in different disease pathologies: A review." Iran J Public Health. 2015; 44(5):714-5.
15. Shalash M, Badra M, Imbaby S, ElBanna E. Malondialdehyde in Type 2 Diabetics and Association With Cardiovascular Risk Factors. J Med Res Inst. 2020; 41(2):21-30. <https://doi.org/10.21608/jmalex.2020.147116>
16. Oesterle A, Laufs U, Liao JK. Pleiotropic Effects of Statins on the Cardiovascular System. Circ Res. 2017; 120(1):229-43. <https://doi.org/10.1161/CIRCRESAHA.116.308537>
17. Sekunda KN, Ariosta A, Limantoro C, Setiawan AA. Difference in Profiles of Oxidative Stress Marker (MDA) in STEMI and NSTEMI. Diponegoro Med J (Jurnal Kedok Diponegoro). 2021; 10(2):145-50. <https://doi.org/10.14710/dmj.v10i2.29673>
18. Block G, Dietrich M, Norkus EP, Morrow JD, Hudes M, Caan B, et al. Factors associated with oxidative stress in human populations. Am J Epidemiol. 2002; 156(3):274-85. <https://doi.org/10.1093/aje/kwf029>
19. Kubihal C V, Naik HD. Effect of smoking on vitamin C and MDA: a cross sectional comparative study. Int J Res Med Sci. 2019; 7(3):746. <https://doi.org/10.18203/2320-6012.ijrms20190511>
20. Art Suwondo, Ahmad Mohan, Melyana Nurul Widyawati M. Factors Related to Malondialdehyde (MDA) Levels in Metal Casting Industry Workers. Preprints. 2020; 1(October):1-8. <https://doi.org/10.21203/rs.3.rs-99597/v1>
21. Schöttker B, Brenner H, Jansen EHJM, Gardiner J, Peasey A, Kubinová R, et al. Evidence for the free radical/oxidative stress theory of ageing from the CHANCES consortium: A meta-analysis of individual participant data. BMC Med. 2015; 13(1):1-15. <https://doi.org/10.1186/s12916-015-0537-7>
22. Kander MC, Cui Y, Liu Z. Gender difference in oxidative stress: a new look at the mechanisms for cardiovascular diseases. J Cell Mol Med. 2017; 21(5):1024-32. <https://doi.org/10.1111/jcmm.13038>
23. Bell JR, Bernasocchi GB, Varma U, Raaijmakers AJA, Delbridge LMD. Sex and sex hormones in cardiac stress - Mechanistic insights. J Steroid Biochem Mol Biol. 2013; 137:124-35. <https://doi.org/10.1016/j.jsbmb.2013.05.015>
24. P G Curcio F, Bulli G, Aran L, Della-morte D, Testa G, et al. Oxidative Stress and Diseases. Oxidative Stress Dis. 2012; 757-72.
25. Sabitha K, Venugopal B, Rafi M, V Ramana K. Role of Antioxidant Enzymes in Glucose and Lipid Metabolism in Association with Obesity and Type 2 Diabetes. Am J Med Sci Med. 2014; 2(1):21-4. <https://doi.org/10.12691/ajmsm.2-1-5>
26. Manna P, Jain SK. Obesity, Oxidative Stress, Adipose Tissue Dysfunction, and the Associated Health Risks: Causes and Therapeutic Strategies. Metab Syndr Relat Disord. 2015; 13(10):423-44. <https://doi.org/10.1089/met.2015.0095>
27. Lewis NA, Howatson G, Morton K, Hill J, Pedlar CR. Alterations in Redox Homeostasis in the Elite Endurance Athlete. Sport Med. 2015; 45(3):379-409. <https://doi.org/10.1007/s40279-014-0276-5>

LAMPIRAN V : Bukti Bukti Screenshot pada Sinta

The screenshot shows the SINTA Executive dashboard. On the left is a navigation menu with options: Dashboard, Explore SINTA, Mutation History, List Verificator PT, and My SINTA. The top right corner displays the user profile for JUNI CHUDRI with Sinta ID 6727331 and a notification of 3 records. The main content area lists three publications:

Publication Title	Author	Journal	Year	Cited
Pengukuran Kadar Kolesterol Total Dan Pelayanan Konsultasi Sindrom Metabolik Di Kelurahan Rawamangun, Jakarta Timur	A Handayani, F Chondro, LM Surjadi, J Chudri	AMMA: Jurnal Pengabdian Masyarakat 2 (5), 515-519, 2023	2023	0
Factors associated with plasma malondialdehyde levels in people over 40 years	M Melyanti, E Margo, J Chudri, MA Faradilla	Journal of Drug Delivery and Therapeutics 13 (7), 52-56, 2023	2023	2
Penyuluhan Protokol 3M Dalam Pencegahan COVID-19 Pada Penduduk Kelurahan Pluit, Kecamatan Penjaringan, Jakarta Utara	J Chudri, F Chondro, A Handayani	AMMA: Jurnal Pengabdian Masyarakat 2 (9: Oktober), 1099-1102, 2023	2023	0

The second row, corresponding to the publication 'Factors associated with plasma malondialdehyde levels in people over 40 years', is highlighted with a red rectangular border.

Factors Associated with Plasma
Malondialdehyde Levels in
People over 40 Years

by Meivanti Meivanti

Submission date: 28-Jun-2023 08:09PM (UTC+0700)

Submission ID: 2123934068

File name: aldehyde Levels in People over 40 Years_JDDT_Meivanti_et_al.docx (61.09K)

Word count: 3703

Character count: 20873

Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years

Meiyanti¹, Yohana, Eveline Margo, Juni Chudoi, Pusparini, Meutia Atika Faradilla

Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Corresponding Email: meiyanti@trisakti.ac.id

Address for correspondence: Meiyanti, Department of Pharmacology and Medical Pharmacy, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Acknowledgments

The authors thank all the subjects who participated in this study, as well as the laboratory staff who assisted in the implementation of this study.

Abstract. Aging is a progressive process caused by a physiological decline in body functions. It is known that several factors can accelerate aging, namely age, lifestyle, oxidative stress, lipid peroxidation, and mitochondrial damage. Various pollutants such as cigarette smoke, motor vehicle fumes, industrial waste, and exposure to sunlight / ultraviolet light will form free radicals. Malondialdehyde (MDA) level is one of the indicators used to measure oxidative stress. Increased MDA levels are found in various degenerative diseases. This study aims to determine plasma MDA levels in the age group of 40 years and over and factors that can affect plasma MDA levels. This study was a cross-sectional study by including 42 subjects. Data collection was carried out using a questionnaire to obtain subject demographic data and physical activity, as well as physical examinations complemented by laboratory examinations and MDA levels. The mean age of the subjects was 52.7 years, and most of the 32 (76.2%) subjects were categorized as obese and had high activity. Pearson correlation test obtained body mass index statistically significantly correlated with plasma MDA levels with a value of $p = 0.039$, as well as physical activity statistically significantly correlated with a negative correlation, with a value of $r = -0.26$ and $p = 0.043$. Conclusion Body mass index and activity are significantly correlated with plasma MDA levels.

Keywords: malondialdehyde, stress, oxidative, radicals

INTRODUCTION

Aging is a progressive process caused by a physiological decline in body function.⁽¹⁾ A decrease in body homeostasis is accompanied by an increased risk of various chronic diseases including diabetes mellitus, cancer, neurodegenerative diseases, and heart disease. It is known that several factors can accelerate aging, namely age, lifestyle, oxidative stress, lipid peroxidation, and mitochondrial damage. Previous studies have shown that changes in sphingomyelin metabolism in the central nervous system and age are associated with neurodegenerative diseases.⁽²⁾ In the US, aging is one of the most important factors in diabetes mellitus and heart disease. In 2015, an estimated 12% of the population had diabetes mellitus and an undetected 24% had diabetes mellitus. Heart disease is the highest contributor to mortality in the population aged 65-84 years.⁽³⁾ Recent studies have shown that non-diabetic populations whose fasting glucose levels increase in the blood will increase the risk of coronary heart disease. But on the other hand, the data on fasting blood glucose increase is still debated in the population of cardiovascular disease patients who do not have diabetes mellitus. A recent study in North Sumatra showed a significant relationship between MDA levels in obese and non-obese populations.⁽⁴⁾

Free radicals are formed through two processes, namely exogenous and endogenous processes. Various pollutants such as cigarette smoke, motor vehicle fumes, industrial waste, and exposure to sunlight/ultraviolet will form free radicals. Illness, stress, and excessive exercise/physical activity are endogenous factors produced by the human body that can form free radicals. The formation of free radicals occurs due to oxidation processes and metabolic processes that take place during exercise or activity, inflammatory processes, consumption of certain foods, and others. Free radicals will begin to activate with molecules in the surrounding cells to obtain electrons to stabilize the condition of one's body. The electrons that have been taken in the cell molecules will trigger the

occurrence of free radicals. If this continues for a long time, it can cause oxidative stress which can cause inflammation, and DNA damage can also trigger the disease. Oxidative stress conditions characterized by increased production of Reactive Oxygen Species (ROS) have implications for various diseases such as hypertension, atherosclerosis, diabetes mellitus, stroke, chronic renal failure, heart failure, and other chronic diseases.⁽⁵⁻⁷⁾

Oxidative stress is an imbalance between oxidants (free radicals) and antioxidants. Free radicals that commonly cause oxidative damage are reactive oxygen species (ROS).⁽⁸⁾ ROS compounds are formed through aerobic metabolism. These compounds can cause extensive damage through catalytic reactions. They are responsible for damage at the cellular level by oxidizing proteins, lipids, DNA (Deoxyribonucleic acid), and other biomolecules. Various compounds can be used as markers of oxidative stress such as malondialdehyde (MDA), and protein carbonylation, 4-hydroxy-2-nonenal (4HNE), and 8-hydroxy-deoxyguanosine (8OHdG).⁽⁹⁾

Malondialdehyde (MDA) is a result of lipid peroxidation in the body. The concentration of MDA indicates an oxidation process that occurs in the body's cell membranes due to metabolic processes. One of the factors that cause aging is the production of excess reactive oxygen species (ROS). ROS are generated through the process of cell respiration in the mitochondria and the phagocytosis process of foreign body elimination. To stabilize the amount of ROS, the body can use endogenous and exogenous antioxidants. Disruption of the balance between ROS and antioxidants can cause irreversible damage to cell organelle components up to the DNA level.^(10,11)

MDA levels are strongly influenced by many factors such as age, illness, stress, various pollutants, food consumption, antioxidant supplements and excessive physical activity or exercise, and others. The other study found that more than 30% of subjects had high MDA levels, and concluded that there was a significant relationship between eating habits, body mass index, glucose levels with MDA levels, as well as smoking habits significantly related to MDA levels.⁽¹²⁾ Recent studies have shown that plasma MDA levels in populations aged 70 years and 60 years are not significantly different but there is an increase in the specific activity of the catalase enzyme that increases with age.⁽¹³⁾ In another study, there was a significant difference in plasma MDA levels in the obesity group without metabolic syndrome compared to the obesity group with metabolic syndrome.⁽¹⁴⁾

There are still variations in MDA data in various groups with different ages and different disease groups, so this study aims to determine plasma MDA levels in the age group 40 years and over and factors that can affect plasma MDA levels.

METHOD

This observational analytic study with a cross-sectional design approach was conducted from November 2022 to February 2023 in Angke village, West Jakarta, Indonesia. The inclusion criteria for this study were: men and women aged ≥ 40 years, willing to participate in the study and sign informed consent. Meanwhile, the exclusion criteria were having a history of malignancy, kidney failure, a history of autoimmune disease, taking drugs related to psychiatric conditions, and having limitations in mobility. This study used the calculation of sample size with the formula $N / 1 + N \alpha^2$, and obtained the minimum number of samples required was 40 subjects. Demographic data and physical activity were obtained from distributed questionnaires. Blood chemistry examination (complete blood, glucose, total cholesterol, liver enzymes (SGOT and SGPT) using an enzymatic method. Examination of malondialdehyde levels was carried out using the spectrophotometric method. This research has obtained approval from the ethics committee of the Faculty of Medicine, Trisakti University, Number: 178/KER/FR/IX/2022.

RESULTS

A total of 42 subjects participated in this study, most of the subjects were female 27 (64.3%), and the mean age of the subjects was 52.7 years. Anthropometric data of the subjects are as follows: body weight 67.33 ± 13.64 and height 156.02 ± 8.73 cm, while for abdominal circumference the average is 93.99 cm. Most of the subjects 59.5% fell into the normotensive category, with an average systolic blood pressure of 141.62 mmHg and an average diastolic blood pressure of 82.45 mmHg. A total of 32 (76.2%) subjects were categorized as obese with BMI ≥ 25.0 , and most subjects had high activity levels. (Table 1)

Table 1. Distribution of Subject Characteristics (n=42)

Characteristics	n (%)	Mean ± SD
Gender		
Male	15 (35.7)	
Female	27 (64.3)	
Blood pressure (mmHg)		
Normal (< 140/90)	25 (59.5)	
Hypertension (≥ 140/90)	17 (40.5)	
Physical Activity		
Low (< 600 Mets)	11 (26.2)	
High (≥ 600 Mets)	31 (73.8)	
Body Mass Index (kg/m²)		
Not obese (< 25.0)	10 (23.8)	27.69 ± 4.90
Obese (≥ 25.0)	32 (76.2)	
Age (years)		52.7 ± 10.60
Body weight (kg)		67.53 ± 13.64
Height (cm)		156.02 ± 8.73
Abdominal circumference (cm)		93.09 ± 10.54
Blood pressure (mmHg)		
Systole		141.62 ± 20.77
Diastole		82.45 ± 12.70

The examination of the subject's plasma malondialdehyde levels obtained a mean of 0.77 with the highest value of 1.03 and the lowest of 0.51. Hemoglobin levels averaged 13.16 g/dL. Fasting blood glucose level was 95.06 ± 41.01 mg/dL and total cholesterol was 179.83 ± 31.85 mg/dL. Laboratory examinations to determine liver function were carried out by measuring SGPT and SGOT levels, obtaining mean values of 24.21 U/L and 26.31 U/L, respectively. (Table 2)

Table 2. Laboratory examination results of subjects (n=42)

Variable	Mean ± SD
Hemoglobin (g/dL)	13.16 ± 1.56
Fasting blood glucose (mg/dL)	95.06 ± 41.01
Total cholesterol (mg/dL)	179.83 ± 31.85
SGPT (U/L)	24.21 ± 13.50
SGOT (U/L)	26.31 ± 11.51
Plasma malondialdehyde	0.77 ± 0.26

Determining the type of correlation test to be used by first conducting a data normality test. The normality test used was Shapiro-Wilk because the number of subjects was less than 50 subjects. The correlation test used to assess the correlation between blood glucose, total cholesterol, SGPT, SGOT levels, age, body mass index and physical activity with plasma MDA levels was the Pearson correlation test. The test results are presented in Table 3. Blood glucose, total cholesterol, SGPT, SGOT, and age did not have a significant correlation with MDA levels. Body mass index has a significant correlation with plasma MDA ($p = 0.039$), with a positive correlation direction, and the strength of the correlation is statistically weak with a value of $r = 0.396$ ($r^2 = 0.157$). The physical activity variable has a significant correlation with plasma MDA levels ($p = 0.043$) with a negative correlation direction but with a statistically weak correlation strength ($r = -0.26$).

Table 3. Correlation Test of Age, Gender, Glucose, SGOT, SGPT, BMI, Physical Activity with Plasma MDA Levels (n=42)

	Plasma MDA Level	
	r	p
Age	0.098	0.756
Gender	0.124	0.108
Glucose	0.330	0.835
SGOT	0.104	0.513
SGPT	0.008	0.614
Body Mass Index	0.396	0.039*
Physical activity	0.260	0.043*

* p < 0.05

Discussion

The level of oxidative stress can be measured using Malondialdehyde (MDA) concentration. MDA is the result of lipid peroxidation, which will cause cell damage. Oxidant compounds such as free radicals or non-radical species attack lipids that contain carbon-carbon double bonds, especially unsaturated fatty acids (PUFA). MDA is one of the biomarkers widely used to assess oxidative stress. The mean serum MDA in this study 0.77 was found to be lower than the previous study⁽¹⁰⁾, this is because the age difference of the subjects in this study was younger than in the previous study. In contrast, a study conducted in Iraq on hypertensive patients concluded that the increase in MDA was associated with an increase in age, gender, and duration of disease.⁽¹³⁾

Free radicals such as ROS are the etiology of most chronic diseases, as it is known that active free radicals are the result of metabolic reactions. Damage to body tissue cells occurs because active free electrons remain looking for other electrons to settle, if no other electrons are available then active free electrons will attack cells resulting in cell damage. Although free radicals naturally have benefits for the body, when the level of free radicals exceeds the normal value, it will cause harm to the body.^(11,12)

Measurement of oxidative stress can be done by measuring plasma malondialdehyde (MDA) levels. MDA is a secondary result of the lipid peroxidation process, producing various active compounds that will damage cells. The process of MDA formation through enzymatic and non-enzymatic processes. Currently, MDA is one of the biomarkers used to assess oxidative stress.⁽¹⁴⁾ Some previous researchers concluded that higher serum MDA was found in patients with hyperlipidemia. The condition of increased fat content in the body is in line with increased oxidative stress. Hyperlipidemia is associated with physical changes in cell membranes that facilitate the release of free radicals from the mitochondrial electron transport chain or activation of NADPH oxidase. In patients with diabetes mellitus (DM) there is also an increase in MDA levels, malondialdehyde interacts irreversibly and reversibly with proteins and phospholipids to have a profound effect, especially on the collagen of the cardiovascular system. MDA levels remain within normal limits due to the antioxidant effect, but in DM, this protective effect is impaired resulting in increased MDA levels.^(1,15)

Increased lipid peroxidation is considered a consequence of oxidative stress that occurs when the dynamic balance between prooxidant and antioxidants is disrupted. The statin class of drugs for hyperlipidemia therapy is proven to be anti-atherogenic, can improve endothelial function, inhibit vascular inflammation, and stabilize atherosclerotic plaques. In addition to statins, non-steroidal anti-inflammatory drugs (NSAIDs) can affect MDA concentration. Previous studies have found that NSAIDs have significant antioxidant properties to scavenge free radicals.^(16,17)

In this study, there was no significant correlation between age and MDA levels. Similar results were also obtained by other researchers, cell damage due to oxidative stress is not related to increasing age but is associated with habits, behavior patterns, and physiological changes due to age. Various habitual patterns such as smoking, fatty food habits, lack of activity, and inflammatory processes can lead to increased oxidative stress. The habit of taking vitamin C (ascorbic acid) has a beneficial impact due to its antioxidant effect so cell damage and DNA damage are lower in this group.⁽¹⁸⁾ MDA levels

were significantly higher among all levels of smokers. MDA levels were lower in light smokers compared to heavy or moderate smokers. Higher vitamin C levels were found in the non-smoker group. All body tissues are susceptible to free radical damage, but based on location, the respiratory epithelium is particularly vulnerable to cigarette smoke. Oxygen radicals generated close to the cell membrane oxidize membrane phospholipids which will result in lipid peroxidation with the end products of this reaction being malondialdehyde, ethane, and pentane, this causes an increase in MDA levels in smokers.⁽¹⁸⁾ Another study conducted in Klanten on foundry workers concluded that type of work, body mass index and smoking habits did not affect MDA levels, but length of work and marital status were associated with MDA levels. Long and continuous exposure to metallic materials will result in the accumulation of these materials in the body and will increase the oxidative process of polyunsaturated fats in cell membranes and worsen the level of lipid peroxidase, characterized by an increase in MDA levels as a reflection of the intensity and rate of lipid peroxidase reactions of damage to body cells.⁽²⁰⁾

Hardiandy's study also concluded that there was no significant difference in plasma MDA levels in the age population, but increasing age along with an increase in catalase-specific activity.⁽¹⁹⁾ This increase in catalase-specific activity is a protective effect especially for elderly women to overcome high levels of oxidative damage. The role of endogenous antioxidant catalase is to suppress oxidative damage.^(18,21) Oxidative stress plays an important role in the incidence of cardiovascular diseases including coronary artery disease. Oxidative stress is considered an important mechanism for the development of cardiovascular diseases especially atherosclerosis. Hypertension, cardiac hypertrophy, and ischemia-reperfusion injury are three other major cardiovascular conditions for which gender is a risk factor. Gender is also associated with differences in oxidative stress. Under physiological conditions, women appear to be less susceptible to oxidative stress. This may be due to the antioxidant properties of estrogen, sex differences in NADPH-oxidase activity or other mechanisms yet to be determined. Sex hormones including estrogen are globally present with their receptors in the heart so sex hormones can exert effects on the cardiovascular system. Estrogen can improve heart function and reduce injury to the heart. In addition, estrogen has antioxidant effects so that it can function as a cardioprotective.^(22,23)

This study found a significant correlation between body mass index and MDA levels, the higher the BMI value, the higher the MDA levels. Similar results were also obtained in a study in India that concluded that in metabolic syndrome patients with one of the signs is an increase in BMI, there is an increase in MDA levels compared to the control group. In obese subjects, the level of MDA is higher than normal IMT. In a healthy body, there is redox homeostasis that occurs in cells, if an imbalance arises it will cause oxidative stress. Obese patients will have an increased metabolic load and mechanical load on the myocardium, increased body mass, and excessive food intake will cause the formation of lipid peroxidation, and free radicals.⁽²⁴⁾ The increase in some of these things will stimulate antioxidant enzymes in the long run, this will cause antioxidant enzyme reserves to be depleted and unable to overcome and balance the increase in oxidative stress.^(25,26)

These types of training activities are exhausting, long exercise, overtraining syndrome, various types of exercise will induce a significant response to oxidative stress. Exercise activities such as light exercise, low-intensity exercise, and prolonged training will have an impact on improving endogenous antioxidant status. Reactive oxygen species play an important role in cell signaling and in regulating antioxidant gene expression. Exertion results in hyperregulation of nuclear factor kappa B and mitogen-activated protein kinases that activate gene expression of a number of enzymes and proteins with important roles in maintaining intracellular oxidative or antioxidant homeostasis.^(16,27) The limitation of this study is that some factors that may affect MDA levels such as consumption of vitamin supplements, diet, and length of illness suffered were not examined in this study.

Conclusions

Body mass index and activity were significantly correlated with MDA levels, while age, gender, blood glucose levels, and liver enzymes were not significantly correlated with MDA levels.

Conflict of Interest:

The investigators declare there is no conflict of interest in this study.

References

1. Sharifi-Rad M, Anil Kumar N V, Zucca P, Varoni EM, Dini L, Panzarini E, et al. Lifestyle, Oxidative Stress, and Antioxidants: Back and Forth in the Pathophysiology of Chronic Diseases. *Front Physiol*. 2020;11(July):1–21.
2. Park MH, Jin HK, Bao J sung. Potential therapeutic target for aging and age-related neurodegenerative diseases: the role of acid sphingomyelinase. *Exp Mol Med [Internet]*. 2020;52(3):380–9. Available from: <http://dx.doi.org/10.1038/s12276-020-0399-8>
3. Calyniuk B, Grochowska-Niedworok E, Walkiewicz K, Kawecka S, Popielek E, Fatyga E. Malondialdehyde (MDA) – product of lipid peroxidation as marker of homeostasis disorders and aging. *Ann Acad Medicus Silesiensis*. 2016;70:224–8.
4. Savita M, Rusdiana ., Widjaja SS, Syahputra M. Comparison Malondialdehyde (MDA) Level between Obesity Non Metabolic Syndrome and Obesity with Metabolic Syndrome Patients. 2020;(5):644–7.
5. Yekti R, Bukhari A, Jafar N, Thaha AR. Measurement of malondialdehyde (MDA) as a good indicator of lipid peroxidation. *Int J Allied Med Sci Clin [Internet]*. 2018;6(4):6–8. Available from: www.ijamsc.com
6. Denise Mafrá ME. Influence of Physical Exercise on Oxidative Stress and Inflammation in Hemodialysis Patients. *J Mol Biomark Diagn*. 2015;06(03).
7. Mao C, Yuan JQ, Lv Y Bin, Gao X, Yin ZX, Kraus VB, et al. Associations between superoxide dismutase, malondialdehyde and all-cause mortality in older adults: A community-based cohort study. *BMC Geriatr*. 2019;19(1):1–9.
8. Ayala A, Muñoz MF, Argüelles S. Lipid peroxidation: Production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. *Oxid Med Cell Longev*. 2014;2014.
9. Situmorang N, Zulham Z. Malondialdehyde (Mda) (Zat Oksidan Yang Mempercepat Proses Penuaan). *J Keperawatan Dan Fisioter*. 2020;2(2):117–23.
10. Hardiany NS, Sucitra S, Paramita R. Profile of malondialdehyde (MDA) and catalase specific activity in plasma of elderly woman. *Heal Sci J Indones*. 2020;10(2):132–6.
11. Hardiany NS, Sadikin M, Siregar N, Wanandi SI. The suppression of manganese superoxide dismutase decreased the survival of human glioblastoma multiforme T98G cells. *Med J Indones*. 2017;26(1):19–25. 439-445_nova.pdf.
13. A. Hassan A, Sayyah SG. Oxidative Stress Marker Malondialdehyde and Glutathione Antioxidant in Hypertensive Patients. *Eur J Biomed Res*. 2023;2(1):31–6.
14. Khoubnasab Jafari M, Ansarin K, Jouyban A. Comments on "use of malondialdehyde as a biomarker for assessing oxidative stress in different disease pathologies: A review." *Iran J Public Health*. 2015;44(5):714–5.
15. Shalash M, Badra M, Imbuby S, ElBanna E. Malondialdehyde in Type 2 Diabetics and Association With Cardiovascular Risk Factors. *J Med Res Inst*. 2020;41(2):21–30.
16. Oesterle A, Laufs U, Liao JK. Pleiotropic Effects of Statins on the Cardiovascular System. *Circ Res*. 2017;120(1):229–43.
17. Sekunda KN, Ariosta A, Limantoro C, Setiawan AA. Difference in Profiles of Oxidative Stress Marker (MDA) in STEMI and NSTEMI. *Diponegoro Med J (Jurnal Kedokt Diponegoro)*. 2021;10(2):145–50.
18. Block G, Dietrich M, Norkus EP, Morrow JD, Hudes M, Caan B, et al. Factors associated with oxidative stress in human populations. *Am J Epidemiol*. 2002;156(3):274–85.
19. Kutshah C V., Naik HD. Effect of smoking on vitamin C and MDA: a cross sectional comparative study. *Int J Res Med Sci*. 2019;7(3):746.
20. Ari Sawondo, Ahmad Mohan, Melyana Nural Widyanawati M. Factors Related to Malondialdehyde (MDA) Levels in Metal Casting Industry Workers. Preprints.

2020;1(October):1-8.

21. Schöttker B, Brenner H, Jansen EHJM, Gardiner J, Peasey A, Kubínová R, et al. Evidence for the free radical/oxidative stress theory of ageing from the CHANCES consortium: A meta-analysis of individual participant data. *BMC Med.* 2015;13(1):1-15.
22. Kander MC, Cui Y, Liu Z. Gender difference in oxidative stress: a new look at the mechanisms for cardiovascular diseases. *J Cell Mol Med.* 2017;21(5):1024-32.
23. Bell JR, Bernasocchi GB, Varma U, Raaijmakers AJA, Delbridge LMD. Sex and sex hormones in cardiac stress - Mechanistic insights. *J Steroid Biochem Mol Biol.* 2013;137:124-35.
24. P G, Curcio F, Bulli G, Aran L, Della-morte D, Testa G, et al. Oxidative Stress and Diseases. *Oxidative Stress Dis.* 2012;757-72.
25. Sabitha K, Venugopal B, Rafi M, V Ramana K. Role of Antioxidant Enzymes in Glucose and Lipid Metabolism in Association with Obesity and Type 2 Diabetes. *Am J Med Sci Med.* 2014;2(1):21-4.
26. Manna P, Jain SK. Obesity, Oxidative Stress, Adipose Tissue Dysfunction, and the Associated Health Risks: Causes and Therapeutic Strategies. *Metab Syndr Relat Disord.* 2015;13(10):423-44.
27. Lewis NA, Howatson G, Morton K, Hill J, Pedlar CR. Alterations in Redox Homeostasis in the Elite Endurance Athlete. *Sport Med.* 2015;45(3):379-409.

Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years

ORIGINALITY REPORT

18% SIMILARITY INDEX	17% INTERNET SOURCES	13% PUBLICATIONS	8% STUDENT PAPERS
--------------------------------	--------------------------------	----------------------------	-----------------------------

PRIMARY SOURCES

1	onlinelibrary.wiley.com Internet Source	3%
2	ejournal3.undip.ac.id Internet Source	3%
3	jddtonline.info Internet Source	2%
4	www.ncbi.nlm.nih.gov Internet Source	2%
5	innovpub.org Internet Source	1%
6	www.researchgate.net Internet Source	1%
7	M. Shalash, M. Badra, S. Imbaby E. ElBanna. "MALONDIALDEHYDE IN TYPE 2 DIABETICS AND ASSOCIATION WITH CARDIOVASCULAR RISK FACTORS", Journal of the Medical Research Institute, 2020 Publication	1%

8	Submitted to Universitas Airlangga Student Paper	1%
9	Submitted to Trisakti University Student Paper	1%
10	repository.stikim.ac.id Internet Source	1%
11	mdpi-res.com Internet Source	1%
12	www.pubmedcentral.nih.gov Internet Source	1%
13	Anna Thomson, Ellen Christensen, Maria Wiklund, Monica Christianson. "A safe place – Adolescents' and young adults' perceptions of youth clinics in northern Sweden", Sexual & Reproductive Healthcare, 2022 Publication	1%
14	www.researchsquare.com Internet Source	1%
15	Warisya Miftah Amanda, Chandra Isabella Hostonida Purba, Ristina Mirwanti. "DIFFERENCE OF BOWEL SOUND RETURN TIME AMONG POST-LAPARATOMY SURGERY PATIENTS AFTER CHEWING GUM", Belitung Nursing Journal, 2019 Publication	<1%



Exclude quotes Off
Exclude bibliography On

Exclude matches < 15 words



Factors Associated with Plasma Malondialdehyde Levels in People over 40 Years

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7