

ISSN 2675-1240

Multidisciplinary Science Journal

Advancing Research, Innovation, and Impact Across Disciplines

Journal Metrics

CiteScore (Scopus, 2025)

1.1

CiteScoreTracker (Scopus,
2026)

1.1

SCImago Journal Rank
(2025)

Q3

Acceptance rate (First
Half, 2026)

22%

Days to first decision
(2026)

45

[ABOUT METRICS](#)



Current issue

VOLUME 9 | ISSUE 2 | FEBRUARY 2027

[VIEW CURRENT ISSUE](#)

[BROWSE ALL ISSUES](#)

Editor-in-chief
Dr. Maiko Dantas

Periodicity
Monthly

Journal features

>> **Open Access** Journal

>> **International** Indexing

>> **Agile and Efficient** Peer-Review

>> Indexed on **Scopus!**

Most recent



26.08.2026

26.08.2026

26.08.2026

26.08.2026

Published in 25 January 2024



[The Victims of Terrorism — and Everything the Attack Does Not End](#)

21/08/2026

[Climate Change: What Do Polar Bears Have to Do with Hospital Queues?](#)

07/08/2026

Publish with Malque Publishing and Advance your Academic Career

[Publisher home](#)

[About us](#)

[Register](#)

[Login](#)

Our Journals

[Applied Veterinary Research](#)

[Humanities Journal](#)

[Journal of Animal Behaviour and
Biometeorology](#)

[Multidisciplinary Science Journal](#)

[Multidisciplinary Reviews](#)

[Malque Health](#)

[NexusSDGs](#)

[NexusTech](#)

[Wild Animals](#)

Editorial Office

Malque Publishing, West Home and Business
- Av. João da Escóssia, 3715 - Sala 57 - Bela
Vista, Mossoró - RN, 59607-330, Brazil.

multiscience@malque.pub



[X @malquepub](#)

[f @malquepub](#)

[@malquepub](#)

[Whatsapp](#)

Editorial Team

Editors-in-Chief



Dr. Maiko Dantas

*Applied Animal Biometeorology Research
Center – ThermoBio, Federal University
of Semi-Arid, Brazil*
maiko.dantas@malque.pub



Dr. Fatemeh Mollaamin

Kastamonu University, Turkey
smollaamin@gmail.com



Most Read Articles

**The economic legacy of world expos:
Analyzing long-term benefits for host
nations**

Panwar et al. 676

**Child pornography in India – issues
and challenges**

Deshpande et al. 653

**Lyophilized and fermented beverage of
hibiscus flowers (*Hibiscus sabdariffa*
L.)**

Rogel-Barrezueta et al. 567

**Management accounting
effectiveness in tourism enterprises:
The roles of technology readiness,
information technology infrastructure,
and data quality in an ai-driven
business environment**

Trang et al. 554



Associate Editors



Dr. Thibério Castelo

*Federal Rural University of Semi-Arid,
Brazil*

thiberio.castelo@malque.pub



Dr. Hitesh Panchal

*Government Engineering College Patan,
India*

panchal_hitesh1@gtu.edu.in



Acceptability of sweet potato
(*Ipomoea batatas*) flour and toasted
rice (*Oryza sativa*) powder as base
ingredients in making polvoron

Refardo et al.

541

Editorial Board Members

Dr. Ala Ali Nemer Amourah

Sohar University, Oman

Email: aamourah@su.edu.om



Dr. Ahmed Obaid

University of Kufa, Iraq

Email: ahmedj.aljanaby@uokufa.edu.iq



Dr. Alejandro Regalado

Universidad del Mar, Mexico

Email: alejandro.regalado33@gmail.com



Dr. Aman B. Upaganlawar

Dr. Anam Azam

Dr. Anand Joshi



SNJB's Shriman Suresh Dada Jain College
of Pharmacy, India

Email: amanrxy@gmail.com



Fraunhofer Institute for Systems and
Innovation Research, Germany

Email: anamrafiq95@gmail.com



Parul University, India
Email: anandyjoshi@gmail.com



Dr. Andee Dzulkarnaen Zakaria

Universiti Sains Malaysia, Malaysia

Email: andee778@gmail.com



Dr. Anis Ahmad Chaudhary

Al-Imam Muhammad Ibn Saud Islamic
University, Riyadh, Saudi Arabia

Email: anis.chaudhary@gmail.com



Dr. Anjanna Matta

ICFAI Foundation for Higher Education,
India

Email: anjireddyitm@gmail.com



Dr. Atif Amin Baig

Management & Science University,
Malaysia

Email: atif_amin@msu.edu.my



Dr. Ayo Olasupo

KU School of Engineering, United States

Email: ayoolasupo@ku.edu



Dr. Bahtiar Mohamad

Universiti Utara Malaysia, Malaysia

Email: mbahtiar@uum.edu.my



Dr. Carlos Alberto Acosta-Olivo

Universidad Autónoma de Nuevo León,

Dr. Ágnes Csiszárík-Kocsir

Obuda University, Budapest, Hungary

Email: kocsir.agnes@kgk.uni-obuda.hu

Dr. Denys Miroshnichenko

National Technical University Kharkiv



Mexico
Email: dr.carlosacosta@gmail.com



Polytechnic Institute, Ukraine
Email: dvmir79@gmail.com



Dr. Farooz Ahmad Lone

Sher-e-Kashmir University of Agricultural
Sciences and Technology of Kashmir,
India
Email: dr.farooz462@gmail.com



Dr. Jafar Alasad Alshraideh

The University of Jordan, Jordan
Email: jalasad@ju.edu.jo



Dr. João Queiroz

Federal University of Roraima, Brazil
Email: jp.fernandes@yahoo.com.br



Dr. K. Karthick

GMR Institute of Technology, India
Email: karthick.k@gmr.it.edu.in



Dr. Kehinde Obayomi

Victoria University Melbourne, Australia
Email: obayomikehindeshola@gmail.com



Dr. Khongdet Phasinam

Pibulsongkram Rajabhat University,
Thailand
Email: phasinam@psru.ac.th



Dr. Leonel Avendaño Reyes

Universidad Autónoma de Baja California,

Dr. Lukman Raimi

Universiti Brunei Darussalam, Brunei
Email: luq_man2001@yahoo.com

Dr. Manish Kumar Dubey

Chandigarh University, India
Email: mkmkdubey@gmail.com



Mexico
Email: lar62@uabc.edu.mx



Dr. Md. Fahad Jubayer

Sylhet Agricultural University, Bangladesh

Email: fahadbau21@hotmail.com



Dr. Mohamed Nor Azhari Azman

*Universiti Pendidikan Sultan Idris,
Malaysia*

Email: mnazhari@ftv.upsi.edu.my



Dr. Musa Danladi Ibrahim

Kogi State University, Nigeria

Email: dimusa55@gmail.com



Dr. Nitalo Machado

Federal University of Maranhao, Brazil

Email: nitalo.farias@ufma.br



Dr. Palanivel Velmurugan

*Bharath Institute of Higher Education
and Research, India*

Email: palanivelmurugan2008@gmail.com



Dr. Potula Sree Brahmanandam

*Shri Vishnu Engineering College for
Women, India*

Email: dranandpotula@svecw.edu.in



Dr. Pranav Mehta

Dharmsinh Desai University, India
Email: pranavmehta.mech@ddu.ac.in

Dr. Rakesh Kumar Ameta

*Gandhinagar University, Gandhinagar,
India*
Email: ametarakesh40@gmail.com

Dr. Ram Chandra Chaurasia

Indian Institute of Technology, India
Email: rcc.0761@gmail.com





Dr. Ramona Birau

Craiova University, Romania
Email: ramona.f.birau@gmail.com



Dr. Rohail Hassan

Universiti Utara Malaysia, Malaysia
Email: rohail.hassan@uum.edu.my



Dr. Rohit Sahu

*GL Bajaj Institute of Technology and
Management, India*
Email: rohitsahu87@gmail.com



Dr. Roman Mylostyvyi

*Dnipro State Agrarian and Economic
University, Ukraine*
Email: mylostyvyi.r.v@dsau.dp.ua



Dr. Romanus Osabohien

Universiti Tenaga Nasional, Malaysia
Email: romik247@gmail.com



Dr. Ruchi Juyal

*Himalayan Institute of Medical Sciences,
India*
Email: ruchijuyal@srhu.edu.in



Dr. Stanly Tebug

*International Committee of the Red
Cross, South Sudan*
Email: stanlytebug@yahoo.com

Dr. Vinod Joshi

Sharda University, India
Email: vinodjoshidmrc@gmail.com



Dr. Wunhong Su

Hangzhou Dianzi University, China
Email: whsu@hdu.edu.cn





Publish with Malque Publishing and Advance your Academic Career

[Publisher home](#)

[About us](#)

[Register](#)

[Login](#)

Our Journals

[Applied Veterinary Research](#)

[Humanities Journal](#)

[Journal of Animal Behaviour and
Biometeorology](#)

[Multidisciplinary Science Journal](#)

[Multidisciplinary Reviews](#)

[Malque Health](#)

[NexusSDGs](#)

[NexusTech](#)

[Wild Animals](#)

Editorial Office

Malque Publishing, West Home and
Business – Av. João da Escóssia, 3715 –
Sala 57 – Bela Vista, Mossoró – RN,
59607-330, Brazil.

multiscience@malque.pub



 [_malquepub](#)

 [_malquepub](#)

 [_malquepub](#)

 [_Whatsapp](#)

Exclusive OJS theme plugin developed with ♥ by



Indexing and Archiving

Indexing and Archiving

Get to know the databases in which Multidisciplinary Science Journal is currently indexed:

- [Scopus \(Elsevier\)](#)
- [Google Scholar](#)
- [Latindex](#)
- [Diadorim](#)
- [Scilit](#)
- [ScienceGate](#)
- [Dimensions](#)
- [CiteFactor](#)
- [Directory of Research Journals Indexing](#)
- [Research Biblie](#)



- [WorldCat](#)
- [CrossRef \(DOI\)](#)
- [Cosmos Impact Factor: 3.774](#)

We are already being evaluated for indexing in Web of Science (Clarivate Analytics).

Publish with Malque Publishing and Advance your Academic Career

[Publisher home](#)

[About us](#)

[Register](#)

[Login](#)

Our Journals

[Applied Veterinary Research](#)

[Humanities Journal](#)

[Journal of Animal Behaviour and
Biometeorology](#)

[Multidisciplinary Science Journal](#)

[Multidisciplinary Reviews](#)

[Malque Health](#)

[NexusSDGs](#)

[NexusTech](#)

[Wild Animals](#)

Editorial Office

Malque Publishing, West Home and
Business – Av. João da Escóssia, 3715 –
Sala 57 – Bela Vista, Mossoró – RN,
59607-330, Brazil.

multiscience@malque.pub



 [_malquepub](#)

 [_malquepub](#)


 [_malquepub](#)

 [_Whatsapp](#)

Exclusive OJS theme plugin developed with ♥ by 



Vol. 7 Issue 8 (2025)



Navigating personal guarantor liability in insolvency: Transformative and transactional shifts in the legal framework

ANGEL SHAJI, SHILPA ML...

28.02.2025



Voting behavior in mayoral election: A case study in Sibolga city, Indonesia

ADIL ARIFIN, FAJAR UTAMA RITONGA, INDRA FAUZAN.

27.02.2025



Institutional and economic drivers of GNI growth in the context of income inequality: Evidence from Southeast Europe

QAZIM TMAVA, MUHAMED SPAHIU, ESAT DURGUTI, ARTA KUÇI.

26.02.2025

Share:




August

Editor-in-chief

Dr. Maiko Dantas

Periodicity

Monthly




Innovative approaches to marine environmental protection: Assessing the impact of community engagement, technology, and policy



Public policy administration in phase III relocation of Mount Sinabung disaster victims: A case study of Mardinding Village, Karo Regency, North Sumatra, Indonesia

AGUNG SUHARYANTO, BEBBY MASITHO BATUBARA, REHIA KARENINA ISABELLA BARUS, AUDIA JUNITA.

25.02.2025



Development of solar powered smart telehealth system for conditioning monitoring in IMC



Assessing the mediating role of top management support in the adoption of cloud computing in the public sector: The case of Ghana

ERIC ENTSIE, ISAAC BRIGHT HURSON, ANTHONY VAZ.


24.02.2025



Automated defect detection in perfume bottle packaging using machine vision approach for improved quality control

AHMAD NAJMUDDEN ISMAIL, NUR AIDAWATY RAFAN, NOR AFFENDY NORIZAN.

25.02.2025



Examining the role of normative influences in sustainable clothing disposal behaviors

MAGHFIKA RIZKY MAULANI, INDAH FATMAWATI, NURYAKIN.

24.02.2025





Career-related parent support and career decision-making self-efficacy among Vietnamese university students: The moderating role of



Thought of romanticism and literacy of classicism: An analysis composition techniques of *Rondo Capriccioso Op.14* by Felix Mendelssohn


XINWEN YAN, JIAN ZHANG.

21.02.2025



De *kretek* konning from Kudus: H. M. Nitisemito's business mindset and strategy against foreign business hegemony in the Dutch East Indies

KADHUNG PRAYOGA, JOKO MARIYONO, ANNISA




The emerging role of dynamic contrast enhanced mri in differentiation between various histologic subtypes of oral cavity squamous cell carcinoma

JUBIN JOHN, VADLAMUDI NAGENDRA.

20.02.2025



Legal politics of law number 11 Of 2020 concerning job creation in the labor legal aspects



Odontogenic potential of dental pulp stem cells in response to bioactive root canal sealers combined with ultrasound stimulation

HAMED ALSHAWKANI.

19.02.2025






Anesthetic management and perioperative care in a high-risk patient with bladder cancer, hypertension, and chronic kidney disease undergoing radical cystectomy with ileal conduit urinary diversion



Evaluating pulmonary sequelae in patients recovering from COVID-19: Short- and mid term complications and correlations with initial disease severity

MRINMAYEE MAYEKAR, AMREESH PAUL.

15.02.2025



Technopreneurship, innovation capability and social media marketing as catalysts for competitive advantages: A study of batik MSMEs in Pekalongan, Indonesia




Anesthetic management of a 17-month-old patient with severe dengue fever, anemia, ascites, and secondary acinetobacter baumannii infection: A case report

RENUKA PATOND, NIKHIL BHALRAO, AMREESH PAUL.

15.02.2025



The dynamics of criminal prosecution for Treason in Papua: A critical study on freedom of expression and the principle of national security



Strategic urbanization: Pestel analysis of subterranean spaces in the evolving demographic environment

ZARA MANVELYAN, TAGUHI BARSEGHYAN, HERMINE SAMVELYAN, EMILIA SARGSYAN, HAYK DHERYAN.

14.02.2025






Key factors shaping students' intention to choose green industrial products: A Hanoi, Vietnam case study




Modeling highly dispersed crash data with sichel GAMLSS: An alternative approach to traditional methods

YAJIE ZOU.

12.02.2025



Assessment of village accessibility and community mobility to enhance socioeconomic conditions of rural communities



Applications of numerical analysis in computer simulations


XHEVDET SPAHIU, SENAD ORHANI.

12.02.2025



Seizure onset after head injury due to road traffic accident: A comprehensive case report

ANKIT SHIRODE, AKHILESH SINGH PARIHAR,



Machine learning analysis of financial behavior: A study of Gen Y and Gen Z preferences

KANCHAN TOLANI, JANMEJAY VIJAYKUMAR SHUKLA,
RAHUL MOHARE, TEJASVINI ALOK PARALKAR.


12.02.2025





Socioeconomic effects of *Pestalotiopsis* rubber leaf fall disease on smallholders in South Sumatra, Indonesia

MASAHIKO OTA, LINA FATAYATI SYARIFA, APRIZAL ALAMSYAH, IMAN SATRA NUGRAHA, HAJAR ASYWADI,



Examining natural number bias processes in elementary school students: A case study in Indonesia


SRI RAHAYUNINGSIH, ADE EKA ANGGRANI, MARDHATILLAH, SITI FAIZAH, MOHAMMAD YUSUF RANDY, ET AL.

10.02.2025



'Maritime Big Play Strategy': Solidifying the role of Aukus in the Indo-Pacific region


RESEARCH ARTICLE | VOL. 7 NO. 8 (2025)



Factors influencing the intention to use artificial intelligence for online advertising on social networks

NGUYEN THI PHUONG GIANG, NGUYEN BINH PHUONG DUY, NGUYEN HO THIEN ANH, DOAN NGUYEN TRA GIANG, DANG MINH TRIET, ET AL.

09.02.2025



Contribution of geophysical methods for detecting underground cavities in the Abda-Doukkala region (Morocco)

MUSTAPHA RHAROUSS, KHADIJA BENHADDOU, LATIFA OUADIF, MARIAME BOUCHAOUR, MOHAMMED MENZHI.




Revealing perceptions of corruption behavior in Indonesian higher education institutions

RUDY HARTANTO, DWI RAHAYU, MEY MAEMUNAH, NOPI HERNAWATI.

09.02.2025





Design and implementation of outdoor

RESEARCH ARTICLE | VOL. 7 NO. 8 (2025)





Figure 1 Conceptual framework.

The effect of CG on tax avoidance: The moderating role of company size

CAI YUN CHEN, SAIDATUL NURUL HIDAYAH
JANNATUN NAIM NOR-AHMAD, RAJA HASLINDA BINTI
RAJA MOHD. ALI.


07.02.2025

RESEARCH ARTICLE | VOL. 7 NO. 8 (2025)



Effectiveness of ethnoscience oriented

CASE REPORT/CLINICAL REPORT | VOL. 7 NO. 8 (2025)




Novel ayurveda management of central serous retinopathy (c.s.r.): an explorative case report

NIKHIL MALI, CHITRAVEENA NIBRAD, SHAMLI HIWARE,
RAZA SHEIKH.

07.02.2025

RESEARCH ARTICLE | VOL. 7 NO. 8 (2025)



Harmonizing heritage: A two-decade scoping bibliometric analysis of music in cultural tourism

RESEARCH ARTICLE | VOL. 7 NO. 8 (2025)



Determining factors of lung tuberculosis among children in community health centers: A cross-sectional study


MACHRUMNIZAR MACHRUMNIZAR, YULIANA YULIANA,
ADANG BACHTIAR.

06.02.2025





Artificial intelligence and data privacy in international trade law




A postmenopausal woman with generalized osteoporosis and risk for fracture: A case report

REZARTA STENA, DASANTILA TAHIRAJ, FLORA ZEQAJ LAMÇJA, KLARA HYSENAJ, BLERINA BANI, ET AL.

04.02.2025



Teachers' performance and




Students' critical thinking in numeracy problem-solving through moderate self-Efficacy: A mixed-methods study

VIA YUSTITIA, DIAN KUSMAHARTI, IMAS SRINANA WARDANI.

04.02.2025



Communication of women's empowerment through MSMEs as a CSR program in the cement subsector industry




Social-emotional development in adolescents: The role of peer interaction, social media engagement, and self-esteem

YANE ARDIAN RACHMAN, UJANG SUMARWAN, MELLY LATIFAH, TIN HERAWATI.


03.02.2025





Marital quality of disaster survivor farming families in cianjur district,

RESEARCH ARTICLE | VOL. 7 NO. 8 (2025)



iDDCS: Cervical intraepithelial neoplasia severity detection during vagina birth using artificial intelligence approach

AKANKSHA KAPRUWAN, SACHIN SHARMA, HIMANSHU RAI GOYAL.

02.02.2025



Advancing innovation through university-industry collaboration: Empirical evidence from a developed

RESEARCH ARTICLE | VOL. 7 NO. 8 (2025)



Disinformation's Influence on Public Perception and Defense Policy Implementation in Indonesia

ARIS SARJITO, NORA LELYANA.

01.02.2025



Maternal knowledge of Cushuro (*Nostoc* sp.) and prevalence of iron deficiency anemia: A study at the "Liberación Social" Health Center, Trujillo - 2024

NORIEGA DIEGO CINTHIA SOLANSH, MERCADO VALENCIA YANELI YOSALIN, GARCIA CADILLO LIZ MERCEDES, VEGA PEÑA ELSIBELL NAYL, LEONELA ESMITH JIMENEZ CUEVA, ET AL.

02.02.2025



Publish with Malque Publishing and Advance your Academic Career

[Publisher home](#)

[About us](#)

[Register](#)

[Login](#)

Our Journals

[Applied Veterinary Research](#)

[Humanities Journal](#)

[Journal of Animal Behaviour and
Biometeorology](#)

[Multidisciplinary Science Journal](#)

[Multidisciplinary Reviews](#)

[Malque Health](#)

[NexusSDGs](#)

[NexusTech](#)

[Wild Animals](#)

Editorial Office

Malque Publishing, West Home and
Business – Av. João da Escóssia, 3715 –
Sala 57 – Bela Vista, Mossoró – RN,
59607–330, Brazil.

multiscience@malque.pub



 [_malquepub](#)

 [_malquepub](#)

 [_malquepub](#)

 [_Whatsapp](#)

Exclusive OJS theme plugin developed with ♥ by





Determining factors of lung tuberculosis among children in community health centers: A cross-sectional study

Machrumnizar Machrumnizar^{ab}  | Yuliana Yuliana^b  | Adang Bachtiar^c 

^aDoctoral Program, Faculty of Public Health, Universitas Indonesia, Indonesia.

^bDepartment of Parasitology, Faculty of Medicine, Universitas Trisakti, Indonesia.

^cDepartment of Health Policy and Administration, Faculty of Public Health, Universitas Indonesia, Indonesia.

Abstract Tuberculosis is a serious global health issue and is particularly prevalent among children. Early and correct diagnosis is very important for effective management, but tuberculosis diagnosis in children is often difficult because the clinical symptoms are nonspecific and difficult to collect specimens. This study aims to determine risk factors based on sociodemographic, socioeconomic, nutrition and clinical status associated with tuberculosis. This study is a cross-sectional study. Children aged 0 to 14 years who came to the community health center with one or more clinical symptoms related to TB from January to December 2023 were studied. Research participants underwent two diagnostic methods, clinical and bacteriological. Positive Clinical diagnosis was found to be 72.4%. Meanwhile, bacteriologically confirmed participants were 10.3%, the remainder were unable to produce sputum. Most respondents were men (62.8%), with 66% of respondents aged ≥ 5 years, and those sociodemographic factors have a significant relationship with tuberculosis in children ($p < 0.05$). This study found that 64.7% of respondents had a BMI < 18.5 and significantly related to tuberculosis in children ($p < 0.05$). Furthermore, tuberculosis incidence and malnutrition also have a significant relationship ($p < 0.05$). Contact with active tuberculosis cases shows a significant difference in the proportion of tuberculosis and nontuberculosis cases ($p = 0.009$), while 57.1% of children were exposed to cigarette smoke ($p < 0.05$). Government health insurance (BPJS) was used by 70.5% of respondents who were tested positive for tuberculosis, and it is statistically significant ($p < 0.05$). Other socioeconomic factors were also significant ($p < 0.05$) for tuberculosis cases in children whose parents had low education at 85.9%; have 96.8% of jobs with a household income equal to or more than the provincial minimum wage (UMP) of 60.3%. In the future, these factors can be utilized to construct prediction models for clinical diagnosis in children suspected of tuberculosis, especially machine learning-based models, in addition to tuberculosis-related clinical symptoms.

Keywords: pediatrics, exposure to tuberculosis, tuberculosis, risk factor

1. Introduction

Among the leading sources of morbidity and death in children aged 0–14 years are tuberculosis (TB), which accounts for approximately 12% of the total 10.6 million TB cases in the world in 2023 and contributes to 3–25% of the total TB cases (World Health Organization, 2023). Children might represent up to 20% of the global burden of TB at endemic locations; hence, the real situation can vary from forecasts given adequate reporting systems and appropriate diagnoses (Richens et al., 2020). Nevertheless, there is a fairly modest rate of TB-identified cases and notifications, particularly in developing nations with limited resources, such as Indonesia; the estimated TB incidence of children aged 0–14 in 2022 is 110,881 children, or approximately 15.3% of all TB cases in Indonesia (Ministry of Health of the Republic of Indonesia, 2023). According to a survey by the United Nations International Children's Emergency Fund (UNICEF), 35% of instances of TB among children between the ages of 0 and 14 are unreported, meaning that these 110,881 children constitute only 65% of the total child TB cases in 2022 (UNICEF Indonesia, 2022). These data indicate that the level of community transmission is relatively high.

Bacteriological testing is the standard diagnostic method for adult TB but has low sensitivity and specificity when used in children (Basile et al., 2022). Since the prevalence of TB in children serves as an immediate indicator and surrogate for TB spread in the community, there is a dearth of novel, easily used, and inexpensive diagnostic methods for TB, especially in remote areas with limited laboratory and X-ray facilities, which adds to existing problems (UNICEF Indonesia, 2022). The diagnosis of tuberculosis in children aged 0–14 years without specific symptoms is a major challenge for doctors, academics, and program managers because it can hinder diagnosis and lead to worse treatment outcomes (Siddalingaiah et al., 2023). Current diagnostic methods have limited sensitivity, especially in children aged 0–14 years (Namuganga et al., 2021). The



scoring system, an alternative for diagnosing TB in children, has not yet been fully implemented according to guidelines (Ministry of Health of the Republic of Indonesia, 2023; World Health Organization, 2022).

In addition to clinical and biological variables, several socioeconomic and environmental variables increase the likelihood of TB in children between the ages of 0 and 14. These risk factors are very useful for predicting the development of TB disease. Therefore, identifying the risk factors for TB in children is crucial for making a diagnosis and initiating treatment as soon as possible for the best outcomes. Community health centers in Indonesia are the main place where TB patients seek treatment, so they are the starting point for TB case care, which includes early diagnosis, treatment, and reporting. The purpose of this study was to assess the sociodemographic, socioeconomic, and clinical risk factors for TB in children aged 0 to 14 years who exhibited one or more TB-related clinical symptoms.

2. Materials and methods

2.1. Data collection

This research was conducted at a community health center in the administrative area of South Jakarta, Indonesia. The respondents were children aged 0–14 years who came with a cough to receive treatment at community health centers between January and December 2023.

2.2. Research design

We used a cross-sectional design and studied children with normal nutritional status or malnutrition, including all genders, aged 0–14 years, who presented with a cough to receive treatment at community health centers between January and December 2023. The number of respondents was determined via consecutive sampling, and the sample size was determined via the total sampling method. All respondents who met the inclusion criteria were included in the study. TB diagnosis is categorized on the basis of clinical diagnosis (accompanied by X-rays and/or tuberculin tests) and bacteriology. The clinical diagnosis was determined by the occurrence or absence of symptoms of a persistent cough lasting more than 2 weeks, weight depletion or no weight gain in the previous two months, fever ≥ 2 weeks, and malaise ≥ 2 weeks. Sociodemographic factors (age, sex, age group, weight, height, nutritional status) were recorded during the medical history. Contact with active TB cases, history of BCG immunization, and exposure to cigarette smoke/air pollution were categorized dichotomously. Socioeconomic factors (education, employment, parental income, and health insurance) were also categorized dichotomously. This study also recorded the presence or absence of comorbid factors (HIV and DM). A child's nutritional status is determined by body mass index (BMI), which is calculated as the ratio of body weight in kilograms to height in meters squared. Furthermore, the BMI value of children aged 0–60 months must be compared with the WHO 2005 standard BMI value (WHO, 2006) and BMI values for children and adolescents (5–19 years old) in comparison to the WHO/NCHS 2007 reference (WHO, 2007).

2.3. Data analysis

All the data that had been collected and entered into standard Microsoft Excel were then analyzed via SPSS for Windows (SPSS Inc. version 25.0, Chicago, IL). The variables were initially examined univariately, and then a chi-square statistic was used to evaluate categorical variables to determine risk factors in children with tuberculosis. The McNemar statistic was used to estimate the proportions of risk variables between TB patients and non-TB patients. A statistically significant probability is smaller than 0.05. The odds ratio (OR) and 95% confidence interval (CI) were computed to ascertain the strength of the link.

3. Results and Discussion

The purpose of this cross-sectional study was to identify the predictive variables for both TB and non-TB diagnoses among children aged 0–14 years in Community Health Center facilities in Indonesia. TB risk factors such as HIV and diabetes were not found in the child respondents, so no analysis was carried out.

In this study, 113 out of 156 respondents were diagnosed with pulmonary TB from January to December 2023. Among the TB patients who received treatment, 72.4% of the respondents were clinically diagnosed with pulmonary TB. Bacteriologically confirmed TB cases were found in 10.3% of the respondents, and 89.7% of the respondents were unable to produce phlegm. Most of the respondents (62.8%) were boys, with 34% being less than five years old and 66% being older than five years (Table 1). Negative smear test results reportedly contribute to 13–20% of transmission and are, on average, 20–25% more infectious than positive test results (Asadi et al., 2022). Previous studies reported that most TB cases were diagnosed on the basis of clinical criteria (50–58%) (Chilyabanyama et al., 2024; Zürcher et al., 2019). Similarly, a previous cohort study reported that 16% (58/372), 95% CI, and 12–19% of the cases were smear-negative (Chilyabanyama et al., 2024), indicating that 84% of the TB cases were smear positive. However, another study reported that 65.2% of TB patients had negative BTA results and that 34.8% had positive BTA results (Campos et al., 2016).

Table 1 Characteristics related to sociodemographic factors, TB contact, BCG immunization, nutritional status, and exposure to smoking/air pollution between TB and non-TB respondents.

Characteristics	Frequency		TB Non-TB		p Value	Odd Ratio (OR)	95% CI
	N, mean	%, SD					
Bacteriological diagnosis							
Yes	16	10.3	16	0	0.009*	-	-
No	140	89.7	97	43			
Age (year)							
<5 years	53	34	38	15	0.000¶	0.964	0.594 – 1.564
≥5 years	103	66	75	28		1.019	0.790 – 1.315
Gender							
Boy	98	62.8	72	26	0.000¶	1.054	0.797 – 1.393
Girl	58	37.2	41	17		0.918	0.589 – 1.429
Weight (kg)	19.6218	8.94918			0.000*	-	18.21 – 21.04
Height (cm)	104.3910	26.23235			0.000*	-	100.24 – 108.54
Body Mass Index (BMI)							
<18.5	101	64.7	91	10	0.000*	3.463	1.997 – 6.005
>18.4	55	35.3	22	33		0.254	0.168 – 0.382
Contacts with active TB patients							
Yes	85	54.5	62	23	0.002¶	1.026	0.741 – 1.420
None	71	45.5	51	20		0.970	0.664 – 1.418
BCG immunization							
Yes	114	73.1	85	29	0.328*	1.115	0.883 – 1.408
No	42	26.9	28	14		0.761	0.445 – 1.302
nutritional status							
Malnutrition	100	64.1	90	10	0.000*	3.425	1.974 – 5.941
Normal	56	35.9	23	33		0.265	0.178 – 0.396
Smoke exposure/air pollution							
Yes	89	57.1	65	24	0.006¶	1.031	0.756 – 1.405
None	67	42.9	48	19		0.961	0.645 – 1.432

*Chi-square test, p value <0.05; ¶McNemar test, p value <0.05.

TB disease is generally determined on the basis of clinical diagnosis and is accepted when bacteriological evidence is unavailable (Abdullahi et al., 2021). As is often reported, most respondents were boys. Various studies have reported similar findings, with TB cases occurring more often in boys (Alao et al., 2020; Campos et al., 2016; Chidambaram et al., 2021) and variations in the age range of children who contract TB. Younger children are believed to be more affected than older children are (Hamid et al., 2019; Ogbudebe et al., 2018; Tilahun & Gebre-Selassie, 2016; Vukugah et al., 2022) due to household contact with active TB patients and immature immunity. However, exposure to active TB can also occur in the community. Boys tend toward physical activity and high socialization, so they enjoy playing outside the house. This causes boys to potentially encounter adult active TB and be exposed to cigarette smoke or air pollution (Martínez-Andrés et al., 2017; Miller et al., 2021).

Malnutrition and tuberculosis (TB) are reciprocally associated: malnutrition increases the likelihood of having active TB by six to ten times, and TB contributes to malnutrition in patients (Feleke et al., 2019; Musuenge et al., 2020). Patients with active TB tend to be underweight and have a BMI less than 18.5 kg/m², which is considered an index of malnutrition. Thus, TB incidence is strongly correlated with poor nutrition and low BMI (Musuenge et al., 2020). The possibility of TB increases due to malnutrition, both micro- and macrodeficits, and compromised immunity. Infection with TB can intensify metabolism and require more energy, which can lead to eased appetite and nutritional malabsorption, ultimately increasing the possibility of being underweight (Feleke et al., 2019; Vonasek et al., 2022). A previous study revealed that, compared with the control group, TB patients had a greater likelihood of having a BMI of less than 18.5 kg/m² (Kitonsa et al., 2020). The nutritional status in this study was assessed via BMI; 64.7% of the respondents had a BMI <18.5, which was significantly associated with the incidence of childhood TB (p<0.05).

This study revealed that 54.5% of the respondents had a history of interaction with active TB patients. The proportions of TB and non-TB cases in the child group differed significantly when the children were exposed to cases of active TB (p value=0.002). The strong connection between TB disease in children and a history of interaction with the source of TB cases is crucial since it is well known that the risk of developing TB is directly proportional to the number of germs to which a child is exposed (Koura et al., 2023). Multivariate analysis in previous research revealed a significant relationship between household contacts of children with active TB, with OR = 15 288 and 95% CI: 5,378–43 457; household contacts with smokers, with OR = 7,094 and 95% CI: 2,128–23 648; and contacts > 18 hours with TB sufferers, with OR = 4.681 and 95% CI: 1.198–18.294 (Laghari et al., 2019). Children under the age of five, especially those under 1 year, who meet a source of TB cases usually have a 50% chance of contracting TB, and ninety-five percent of TB cases occur within a year after contact. Home contacts, caregivers, and

medical professionals are considered close contacts of active tuberculosis patients (Patra et al., 2015) because they have a greater chance of contracting *Mycobacterium tuberculosis* and developing primary active tuberculosis (Rea et al., 2022; Reichler et al., 2018).

This study revealed that 73.1% of the respondents had been vaccinated against BCG. The sole licensed and amply used vaccine to prevent TB prompted by *M. tuberculosis* is Bacillus Calmette-Guérin (BCG) (Casey et al., 2016; Katelaris et al., 2019). BCG immunization is highly effective in preventing meningitis and miliary TB in children, with a 70 to 80% success rate (Katelaris et al., 2019; Roy et al., 2019). An 18-year cohort study in Tunisia reported that 36.5% of TB cases occurred in individuals who received BCG vaccination, whereas 63.3% of unvaccinated individuals did (Bennasrallah et al., 2019).

The respondents in this study more often had a history of exposure to cigarette smoke (57.1%), but none of the respondents had a history of HIV or diabetes (Table 1). In numerous countries where the prevalence of tuberculosis is high, most children are exposed to cigarette smoke or secondhand smoke. A comprehensive analysis encompassing eighteen observational studies revealed that exposure to tobacco-related smoke resulted in children being 3.41 times more likely to contract tuberculosis and 1.64 times more likely to develop latent tuberculosis infection (Patra et al., 2015). Research in Spain reported that exposure to cigarette smoke in children (95% CI: 9.63–18.17) increased the prevalence of TB (13.48%) (Altet et al., 2022). Greater vulnerability to pulmonary tuberculosis has been attributed to several biological factors, including poor mucosal secretion clearance, impaired alveolar macrophage phagocytic capacity, diminished immunological reactions, and/or CD4+ lymphopenia caused by cigarette nicotine (Burusie, Enquesilassie, Addissie, Dessalegn, & Lamaro, 2020; Wang et al., 2020). Another study that examined the relationship between TB disease and exposure to cigarette smoke reported that the detrimental impact of smoking on tuberculosis transmission was significantly reduced immediately after smoking cessation, highlighting the value of smoking cessation programs in the fight against TB (Chu et al., 2021). Case-control studies carried out in Brazil and India have shown that firewood or biomass smoke, apart from cigarette smoke, is a separate threat to TB. However, there is currently still a dearth of information about the pathway through which biomass smoke results in chronic pulmonary illness (Báez-Saldaña et al., 2021; Sarkar et al., 2017).

The World Health Organization (WHO) lists fever, inexplicable weight decrease, lethargy or exhaustion, sweating at night, chest pain, and cough lasting longer than two weeks as manifestations and indicators of TB (World Health Organization, 2024). Given the risk of overdue diagnosis in those infected and subsequent spread via close contacts, over time, cough is the most noticed manifestation of tuberculosis, thus summoning inquiry, particularly in resource-constrained states where tuberculosis incidence rises (defined as ≥ 100 in 100 000 inhabitants) (Field et al., 2018). Most respondents in the current study, 79.5%, came to the health center with cough symptoms, and the remaining 20.5% did not experience cough complaints. The change in body weight was not very different; only 49.4% of the respondents experienced weight loss. A total of 55.8% of the respondents had a fever, and 44.2% did not have a fever; malaise was present in 59.6% of the respondents, and 40.4% did not have malaise (Table 1). In the study population conducted in the Philippines, which provided positive TB test results, 84.9% of patients experienced cough ≥ 2 weeks, fever ≥ 2 weeks of unexplained cause (6.9%), and considerable and unintended weight loss (9.5%), and 6.3% of patients experienced exhaustion, sluggishness, and malaise (Lee et al., 2019). The findings of these studies revealed that all participants with cough symptoms, regardless of the length of time for TB examination, presented increased sensitivity from forty-two percent to fifty-one percent (Cheng et al., 2015).

This research revealed that 70.5% of respondents who had national health insurance facilities (BPJS) and 75.5% of respondents tested positive for pulmonary TB. Meanwhile, research in the Philippines reported that 16% of individuals who tested positive for TB were enrolled in a health insurance program (51%) (Lee et al., 2019). Since TB is known to be a disease of poverty, the national health insurance program in Indonesia both improves the capacity of TB patients to seek treatment and contributes significantly to the TB reporting rate. Another study revealed that 60.3% of patients' household income was either per or exceeded the provincial minimum wage (UMP). Almost all the respondents' parents had jobs, 96.8% of

Whom were grouped as nongovernment workers (85.1%) or government workers (10.9%), but only 14.1% had higher education (Table 2). A systematic review revealed a correlation between tuberculosis and poor educational attainment, with a pooled odds ratio of 2.34 (Ricvan Dana et al., 2024). In a South African study including adolescents aged 12 to 18 years, the characteristics that were predictive of TB were older age, ethnic background, low household earnings, and basic schooling or less for both parents (Tchakounte et al., 2022).

With respect to socioeconomic factors (Table 2), TB in childhood occurs more often in families with lower parental education ($p = 0.005$), parents who have jobs ($p < 0.001$), parents with incomes greater than or equal to the UMP ($p < 0.001$), and those with health insurance ($p < 0.001$). The associations between clinical features and the incidence of bacteriological TB in pediatric patients are displayed in Table 3. Symptoms of cough ($p < 0.001$), subfebrile fever ($p < 0.001$), malaise ($p < 0.001$), and weight loss or no weight gain ($p < 0.001$) were significantly different between TB and non-TB events. Owing to the correlation with the increasing prevalence of TB, socioeconomic determinants are linked to the availability of healthcare services, the capacity to fulfill nutritional demands, and the availability of decent and standardized living spaces (Hameed et al., 2019). Working parents are linked to the external environment and to tuberculosis infection, which can spread to children. The exposure that children obtain from a household member who has TB is nearly fourfold inclined to contract the disease (Long et al., 2022; Martinez et al., 2017).

Table 2 Characteristics related to socioeconomic factors, clinical symptoms, and smoke exposure between TB and non-TB respondents.

Characteristics	Frequency		TB	Non-TB	p Value*	Odd Ratio (OR)	95% CI
	N, mean	%, SD					
Parents education							
Low	134	85.9	97	37	0.005¶	0.998	0.66 – 1.150
High	22	14.1	16	6		1.015	0.425 – 2.422
Parents occupation							
Not working	5	3.2	4	1	0.000¶	1.522	0.175 – 13.238
Working	151	96.8	109	42		0.988	0.932 – 1.047
Household income							
< UMP	62	39.7	42	20	0.000¶	0.799	0.536 – 1.192
>= UMP	94	60.3	71	23		1.175	0.859 – 1.606
Government health insurance							
None	46	29.5	30	16	0.000¶	0.713	0.435 – 1.170
Yes	110	70.5	83	27		1.170	0.906 – 1.510
Chronic coughing ≥2 weeks							
Yes	124	79.5	90	34	0.937*	1.007	0.842 – 1.206
None	32	20.5	23	9		0.972	0.490 – 1.931
Weight loss or does not increase in the last 2 months							
Yes	77	49.4	60	17	0.000¶	1.343	0.893 – 2.020
None	79	50.6	53	26		0.776	0.568 – 1.059
Subfebris febris ≥2 weeks							
Yes	87	55.8	63	24	0.003¶	0.999	0.731 – 1.365
None	69	44.2	50	19		1.001	0.675 – 1.486
Malaise ≥ 2 weeks							
Yes	93	59.6	67	26	0.024¶	0.981	0.737 – 1.305
None	63	40.4	46	17		1.030	0.669 – 1.585

*Chi-square test, p value <0.05; ¶McNemar test, p value <0.05. UMP = provincial minimum wage.

Table 3 Characteristics related to clinical factors among TB and non-TB respondents on the basis of bacteriological diagnosis.

Characteristics	Frequency		TB	Non-TB	p Value*	Odd Ratio (OR)	95% CI
	N, mean	%, SD					
Chronic cough ≥2 weeks							
Yes	124	79.5	14	110	0.000¶	1.114	0.908 – 1.366
2. None	32	20.5	2	30		0.583	0.154 – 2.216
Weight drop or did not rise in the previous 2 months							
Yes	77	49.4	7	70	0.000¶	0.875	0.490 – 1.562
2. None	79	50.6	9	70		1.125	0.708 – 1.787
Subfebris fever ≥2 weeks							
Yes	87	55.8	11	76	0.000¶	1.266	0.880 – 1.822
2. None	69	44.2	5	64		0.684	0.323 – 1.446
Malaise ≥ 2 weeks							
Yes	93	59.6	10	83	0.000¶	1.054	0.704 – 1.578
2. None	63	40.4	6	57		0.921	0.474 – 1.788

¶McNemar test, p value <0.05.

4. Conclusions

Despite this modest study, this study underscores the importance of sociodemographic factors, socioeconomic factors, active TB contacts, malnutrition conditions, and exposure to cigarette smoke or air pollution in the diagnosis of TB in children, in addition to TB-related clinical symptoms (cough, fever, malaise, and weight loss).

Ethical considerations

All procedures involving patients in this study were approved by the Ethical Review Committee of the Faculty of Medicine, Trisakti University No.055/KER/FK/I/2024. Informed consent was signed by all patients' legal guardians.

Conflict of interest

The authors have disclosed no conflicts of interest.

Funding

This research received no external financial support.

References

- Abdullahi, O., Moses, N., Sanga, D., & Annie, W. (2021). The effect of empirical and laboratory-confirmed tuberculosis on treatment outcomes. *Scientific Reports*, 11(1), 14854. doi:10.1038/s41598-021-94153-0
- Alao, M. A., Maroushek, S. R., Chan, Y. H., Asinobi, A. O., Slusher, T. M., & Gbadero, D. A. (2020). Treatment outcomes of Nigerian patients with tuberculosis: A retrospective 25-year review in a regional medical center. *PloS one*, 15(10), e0239225. doi:10.1371/journal.pone.0239225
- Altet, N., Latorre, I., Jiménez-Fuentes, M. Á., Soriano-Arandes, A., Villar-Hernández, R., Milà, C., . . . Group, o. b. o. P. S. S. W. (2022). Tobacco Smoking and Second-Hand Smoke Exposure Impact on Tuberculosis in Children. *Journal of Clinical Medicine*, 11(7), 2000. Retrieved from <https://www.mdpi.com/2077-0383/11/7/2000>
- Asadi, L., Croxen, M., Heffernan, C., Dhillon, M., Paulsen, C., Egedahl, M. L., . . . Long, R. (2022). How much do smear-negative patients really contribute to tuberculosis transmissions? Re-examining an old question with new tools. *eClinicalMedicine*, 43. doi:10.1016/j.eclinm.2021.101250
- Báez-Saldaña, R., Canseco-Raymundo, A., Ixcot-Mejía, B., Juárez-Verdugo, I., Escobar-Rojas, A., Rumbo-Nava, U., . . . Arrieta, O. (2021). Case-control study about magnitude of exposure to wood smoke and risk of developing lung cancer. *European Journal of Cancer Prevention*, 30(6), 462-468. doi:10.1097/cej.0000000000000644
- Basile, F. W., Nabeta, P., Ruhwald, M., & Song, R. (2022). Pediatric Tuberculosis Diagnostics: Present and Future. *Journal of the Pediatric Infectious Diseases Society*, 11(Supplement_3), S85-S93. doi:10.1093/jpids/piac082
- Bennasrallah, C., Kacem, M., Dhoub, W., Zemni, I., Ben Fredj, M., Abroug, H., . . . Sriha Belguith, A. (2019). BCG vaccination and tuberculosis prevention: A forty years cohort study, Monastir, Tunisia. *PloS one*, 14(8), e0219991. doi:10.1371/journal.pone.0219991
- Burusie, A., Enquesilassie, F., Addissie, A., Dessalegn, B., & Lamaro, T. (2020). Effect of smoking on tuberculosis treatment outcomes: A systematic review and meta-analysis. *PloS one*, 15(9), e0239333. doi:10.1371/journal.pone.0239333
- Campos, L. C., Rocha, M. V. V., Willers, D. M. C., & Silva, D. R. (2016). Characteristics of Patients with Smear-Negative Pulmonary Tuberculosis (TB) in a Region with High TB and HIV Prevalence. *PloS one*, 11(1), e0147933. doi:10.1371/journal.pone.0147933
- Casey, R. M., Dumolard, L., Danovaro-Holliday, M. C., Gacic-Dobo, M., Diallo, M. S., Hampton, L. M., & Wallace, A. S. (2016). Global Routine Vaccination Coverage, 2015. *MMWR Morb Mortal Wkly Rep*, 65(45), 1270-1273. doi:10.15585/mmwr.mm6545a5
- Cheng, J., Wang, L., Zhang, H., & Xia, Y. (2015). Diagnostic Value of Symptom Screening for Pulmonary Tuberculosis in China. *PloS one*, 10(5), e0127725. doi:10.1371/journal.pone.0127725
- Chidambaram, V., Tun, N. L., Majella, M. G., Ruelas Castillo, J., Ayeh, S. K., Kumar, A., . . . Karakousis, P. C. (2021). Male Sex Is Associated With Worse Microbiological and Clinical Outcomes Following Tuberculosis Treatment: A Retrospective Cohort Study, a Systematic Review of the Literature, and Meta-analysis. *Clinical Infectious Diseases*, 73(9), 1580-1588. doi:10.1093/cid/ciab527
- Chilyabanyama, R., Kamanga, N., & Mwandia, J. N. (2024). Factors associated with tuberculosis treatment outcomes among TB patients aged 15 years and older at chawama level one hospital in Lusaka, Zambia. *Global Public Health*, 19(1), 2307979. doi:10.1080/17441692.2024.2307979
- Chu, A. L., Lecca, L. W., Calderón, R. I., Contreras, C. C., Yataco, R. M., Zhang, Z., . . . Huang, C.-C. (2021). Smoking Cessation in Tuberculosis Patients and the Risk of Tuberculosis Infection in Child Household Contacts. *Clinical Infectious Diseases*, 73(8), 1500-1506. doi:10.1093/cid/ciab504
- Feleke, B. E., Feleke, T. E., & Biadlegne, F. (2019). Nutritional status of tuberculosis patients, a comparative cross-sectional study. *BMC Pulmonary Medicine*, 19(1), 182. doi:10.1186/s12890-019-0953-0
- Field, S. K., Escalante, P., Fisher, D. A., Ireland, B., Irwin, R. S., Adams, T. M., . . . Weir, K. (2018). Cough Due to TB and Other Chronic Infections: CHEST Guideline and Expert Panel Report. *CHEST*, 153(2), 467-497. doi:10.1016/j.chest.2017.11.018
- Hameed, S., Zuberi, F. F., Hussain, S., & Ali, S. K. (2019). Risk factors for mortality among inpatients with smear positive pulmonary tuberculosis: Smear positive pulmonary tuberculosis. *Pakistan Journal of Medical Sciences*, 35(5). doi:10.12669/pjms.35.5.919
- Hamid, M., Brooks, M. B., Madhani, F., Ali, H., Naseer, M. J., The Childhood Tuberculosis Karachi, G., . . . Amanullah, F. (2019). Risk factors for unsuccessful tuberculosis treatment outcomes in children. *PloS one*, 14(9), e0222776. doi:10.1371/journal.pone.0222776
- Katellaris, A. L., Jackson, C., Southern, J., Gupta, R. K., Drobniowski, F., Lalvani, A., . . . Abubakar, I. (2019). Effectiveness of BCG Vaccination Against Mycobacterium tuberculosis Infection in Adults: A Cross-sectional Analysis of a UK-Based Cohort. *The Journal of Infectious Diseases*, 221(1), 146-155. doi:10.1093/infdis/jiz430
- Kitonsa, P. J., Nalutaaya, A., Mukiibi, J., Nakasolya, O., Isooba, D., Kamoga, C., . . . Kendall, E. A. (2020). Evaluation of underweight status may improve identification of the highest-risk patients during outpatient evaluation for pulmonary tuberculosis. *PloS one*, 15(12), e0243542. doi:10.1371/journal.pone.0243542
- Koura, K. G., Mbitikon, O. B., Fiogbé, A. A., Ouédraogo, A. R., Kuete Kuete, A., Magassouba, A. S., . . . Badoum, G. (2023). Programmatic Implementation of Contact Investigation in Eight African Countries. *Tropical Medicine and Infectious Disease*, 8(1), 29. Retrieved from <https://www.mdpi.com/2414-6366/8/1/29>
- Laghari, M., Sulaiman, S. A. S., Khan, A. H., Talpur, B. A., Bhatti, Z., & Memon, N. (2019). Contact screening and risk factors for TB among the household contact of children with active TB: a way to find source case and new TB cases. *BMC Public Health*, 19(1), 1274. doi:10.1186/s12889-019-7597-0
- Lee, S., Lau, L., Lim, K., Ferma, J., Dodd, W., & Cole, D. (2019). The Presence of Cough and Tuberculosis: Active Case Finding Outcomes in the Philippines. *Tuberculosis Research and Treatment*, 2019(1), 4578329. doi:https://doi.org/10.1155/2019/4578329
- Long, R., Divangahi, M., & Schwartzman, K. (2022). Chapter 2: Transmission and pathogenesis of tuberculosis. *Canadian Journal of Respiratory, Critical Care, and Sleep Medicine*, 6(sup1), 22-32. doi:10.1080/24745332.2022.2035540
- Martínez-Andrés, M., Bartolomé-Gutiérrez, R., Rodríguez-Martín, B., Pardo-Guijarro, M. J., & Martínez-Vizcaino, V. (2017). "Football is a boys' game": children's perceptions about barriers for physical activity during recess time. *International Journal of Qualitative Studies on Health and Well-being*, 12(1), 1379338. doi:10.1080/17482631.2017.1379338
- Martinez, L., Shen, Y., Mupere, E., Kizza, A., Hill, P. C., & Whalen, C. C. (2017). Transmission of Mycobacterium Tuberculosis in Households and the Community: A Systematic Review and Meta-Analysis. *American Journal of Epidemiology*, 185(12), 1327-1339. doi:10.1093/aje/kwx025
- Miller, P. B., Zalwango, S., Galiwango, R., Kakaire, R., Sekandi, J., Steinbaum, L., . . . Kiwanuka, N. (2021). Association between tuberculosis in men and social network structure in Kampala, Uganda. *BMC Infectious Diseases*, 21(1), 1023. doi:10.1186/s12879-021-06475-z



- Ministry of Health of the Republic of Indonesia. Tata Laksana Tuberkulosis Anak dan Remaja. Ministry of Health of the Republic of Indonesia Tahun 2023. Accessed on May 2, 2024. <https://tbindonesia.or.id/wp-content/uploads/2024/02/Final-Petunjuk-Teknis-Tata-Laksana-TBC-Anak-Remaja-2023.pdf>.
- Musunge, B. B., Poda, G. G., & Chen, P.-C. (2020). Nutritional Status of Patients with Tuberculosis and Associated Factors in the Health Centre Region of Burkina Faso. *Nutrients*, 12(9), 2540. Retrieved from <https://www.mdpi.com/2072-6643/12/9/2540>
- Namuganga, A. R., Chegou, N. N., & Mayanja-Kizza, H. (2021). Past and Present Approaches to Diagnosis of Active Pulmonary Tuberculosis. *Frontiers in Medicine*, 8. doi:10.3389/fmed.2021.709793
- Ogbudebe, C. L., Adepoju, V., Ekerete-Udofia, C., Abu, E., Egesemba, G., Chukwueme, N., & Gidado, M. (2018). Childhood Tuberculosis in Nigeria: Disease Presentation and Treatment Outcomes. *Health Services Insights*, 11, 1178632918757490. doi:10.1177/1178632918757490
- Patra, J., Bhatia, M., Suraweera, W., Morris, S. K., Patra, C., Gupta, P. C., & Jha, P. (2015). Exposure to Second-Hand Smoke and the Risk of Tuberculosis in Children and Adults: A Systematic Review and Meta-Analysis of 18 Observational Studies. *PLOS Medicine*, 12(6), e1001835. doi:10.1371/journal.pmed.1001835
- Rea, E., Huard, J., & Lee, R. (2022). Chapter 11: Tuberculosis contact investigation and outbreak management. *Canadian Journal of Respiratory, Critical Care, and Sleep Medicine*, 6(sup1), 167-183. doi:10.1080/24745332.2022.2037909
- Reichler, M. R., Khan, A., Sterling, T. R., Zhao, H., Moran, J., McAuley, J., . . . Team, T. E. S. C. T. O. (2018). Risk and Timing of Tuberculosis Among Close Contacts of Persons with Infectious Tuberculosis. *The Journal of Infectious Diseases*, 218(6), 1000-1008. doi:10.1093/infdis/jiy265
- Richens, J. G., Lee, C. M., & Johri, S. (2020). Improving the accuracy of medical diagnosis with causal machine learning. *Nature Communications*, 11(1), 3923. doi:10.1038/s41467-020-17419-7
- Ricvan Dana, N., Rika, S., Pudia M, I., Benny Alexander, M., Muthia, S., Linda, R., . . . Mochammad Fariz, A. (2024). MODIFIABLE AND NON-MODIFIABLE RISK FACTORS FOR TUBERCULOSIS AMONG ADULTS IN INDONESIA: A SYSTEMATIC REVIEW AND META-ANALYSIS. *African Journal of Infectious Diseases (AJID)*, 18(2), 19-28. doi:10.21010/Ajidv18i2.3
- Roy, P., Vekemans, J., Clark, A., Sanderson, C., Harris, R. C., & White, R. G. (2019). Potential effect of age of BCG vaccination on global paediatric tuberculosis mortality: a modelling study. *The Lancet Global Health*, 7(12), e1655-e1663. doi:10.1016/S2214-109X(19)30444-9
- Sarkar, M., Srinivasa, Madabhavi, I., & Kumar, K. (2017). Tuberculosis associated chronic obstructive pulmonary disease. *The Clinical Respiratory Journal*, 11(3), 285-295. doi:<https://doi.org/10.1111/crj.12621>
- Siddalingaiah, N., Chawla, R., Nagaraja, S. B., & Hazra, D. (2023). Risk factors for the development of tuberculosis among the pediatric population: a systematic review and meta-analysis. *European Journal of Pediatrics*, 182(7), 3007-3019. doi:10.1007/s00431-023-04988-0
- Tchakounte Youngui, B., Tchounga, B. K., Graham, S. M., & Bonnet, M. (2022). Tuberculosis Infection in Children and Adolescents. *Pathogens*, 11(12), 1512. Retrieved from <https://www.mdpi.com/2076-0817/11/12/1512>
- Tilahun, G., & Gebre-Selassie, S. (2016). Treatment outcomes of childhood tuberculosis in Addis Ababa: a five-year retrospective analysis. *BMC Public Health*, 16(1), 612. doi:10.1186/s12889-016-3193-8
- UNICEF Indonesia. Desk Review: Pediatric Tuberculosis with a Focus on Indonesia 2022. *UNICEF Indonesia*. Accessed on May 2, 2024. <https://www.unicef.org/indonesia/reports/desk-review-pediatric-tuberculosis-focus-indonesia>.
- Vonasek, B. J., Radtke, K. K., Vaz, P., Buck, W. C., Chabala, C., McCollum, E. D., . . . Garcia-Prats, A. J. (2022). Tuberculosis in children with severe acute malnutrition. *Expert Review of Respiratory Medicine*, 16(3), 273-284. doi:10.1080/17476348.2022.2043747
- Vukugah, T. A., Akoku, D. A., Tchoupa, M. M., & Lambert, E. (2022). Epidemiology of Pediatric Tuberculosis and Factors Associated with Unsuccessful Treatment Outcomes in the Centre Region of Cameroon: A Three-Year Retrospective Cohort Study. *Interdisciplinary Perspectives on Infectious Diseases*, 2022(1), 2236110. doi:<https://doi.org/10.1155/2022/2236110>
- Wang, E. Y., Arrazola, R. A., Mathema, B., Ahluwalia, I. B., & Mase, S. R. (2020). The impact of smoking on tuberculosis treatment outcomes: a meta-analysis. *Int J Tuberc Lung Dis*, 24(2), 170-175. doi:10.5588/ijtld.19.0002
- World Health Organization. World health statistics 2024: monitoring health for the SDGs, Sustainable Development Goals. Geneva: WHO. Accessed on August 21, 2024. <https://www.who.int/data/gho/publications/world-health-statistics>
- World Health Organization. WHO Global Tuberculosis Report 2023. Geneva: WHO. Accessed on May 2, 2024. <https://www.who.int/teams/global-tuberculosis-programme/data>.
- World Health Organization. WHO consolidated guidelines on tuberculosis: Module 5: Management of tuberculosis in children and adolescents 2022. Geneva: WHO. Accessed on May 2, 2024. https://www.ncbi.nlm.nih.gov/books/NBK579387/pdf/Bookshelf_NBK579387.pdf.
- Zürcher, K., Ballif, M., Kiertiburanakul, S., Chenal, H., Yotebieng, M., Grinsztejn, B., . . . consortium, t. I. E. D. t. E. A. (2019). Diagnosis and clinical outcomes of extrapulmonary tuberculosis in antiretroviral therapy programmes in low- and middle-income countries: a multicohort study. *Journal of the International AIDS Society*, 22(9), e25392. doi:<https://doi.org/10.1002/jia2.25392>

Revisi 2

by Machrumnizar FK

Submission date: 08-Jun-2024 08:03PM (UTC+0700)

Submission ID: 2398197390

File name: REVISI_6624__UAS_PIP__Machrumnizar__2306197952.docx (59.05K)

Word count: 5502

Character count: 32168

Determining Factors of Lung Tuberculosis among Indonesian Children in Community Health Centers: A Cross-sectional Study

Machrumnizar Machrumnizar^{1*}, Adang Bachtiar²

¹Doctoral Program, Faculty of Public Health, University of Indonesia, Indonesia

²³Departboyst of Health Analysis and Policy, Faculty of Public Health, University of Indonesia, Indonesia

Corresponding author: *machrumnizar_md@trisakti.ac.id

Abstract

Background: Tuberculosis is a serious global health issue and is particularly prevalent among children. Early and correct diagnosis is very important for effective management, but tuberculosis diagnosis in children is often difficult because the clinical symptoms are non-specific and difficult to collect specimens. This study aims to determine risk factors based on sociodemographic, socioeconomic, nutrition and clinical status associated with tuberculosis. **Methods:** Children aged 0 to 14 years who presented to the community health center with one or more clinical symptoms related to TB from January to December 2023 were studied. Research participants underwent two diagnostic methods, clinical and bacteriological. **Results:** Clinical diagnosis was found to be 72.4% and 27.6% were non-tuberculosis. Meanwhile, bacteriologically confirmed participants were 10.3%, the remainder were unable to produce sputum. Most respondents were boys (62.8%), with 66% of respondents aged ≥ 5 years, and those sociodemographic factors have a significant relationship with tuberculosis in children ($p < 0.05$). This study found that 64.7% of respondents had a BMI < 18.5 and significantly related to tuberculosis in children ($p < 0.05$). Furthermore, tuberculosis incidence and malnutrition also have a significant relationship ($p < 0.05$). Contact with active tuberculosis cases shows a significant difference in the proportion of tuberculosis and non-tuberculosis cases ($p = 0.009$), while 57.1% of children exposed to cigarette smoke ($p < 0.05$). Government health insurance (BPJS) was used by 70.5% of respondents who tested positive for tuberculosis, and it is statistically significant ($p < 0.05$). Other socio-economic factors were also significant ($p < 0.05$) for tuberculosis cases in children whose parents had low education at 85.9%; have 96.8% of jobs with a household income equal to or more than the provincial minimum wage (UMP) of 60.3%. **Conclusion:** In the future, these factors can be utilized to construct prediction models for clinical diagnosis in children suspected of tuberculosis, especially machine learning-based models, in addition to tuberculosis-related clinical symptoms.

Keyword: pediatrics, tuberculosis contact, tuberculosis, risk factor, comorbidity

Introduction

Among the leading source for morbidity and death in children aged 0 – 14 years is tuberculosis (TB), which is around 12% of the total 10.6 million TB sufferers in the world in 2023 and contributes to 3 – 25% of the total TB caseload [1]. Children might represent up to 20% of the global burden of TB in endemic spots, hence the real situation can vary from forecasts given adequate reporting systems and appropriate diagnosis [2]. Still, there is a fairly modest rate of TB

identified cases and notification, particularly in developing nations with limited resources, such as Indonesia, the estimated TB incidence of children aged 0 – 14 in 2022 is 110,881 children or around 15.3% of all TB cases in Indonesia [3]. According to a survey by The United Nations International Children's Emergency Fund (UNICEF), 35 percent of instances of TB among children between the ages of 0 and 14 go unreported, meaning these 110,881 children are only 65% of the total child TB cases in 2022 [4]. Mentioned data indicate the level of community transmission is relatively high.

Bacteriological testing is the standard diagnosis in adult TB but has low sensitivity and specificity when used in children [5]. Since the prevalence of TB in children serves as an immediate clue and surrogate for TB spreads in the community, there is a dearth of novel, easily used, and inexpensive diagnostic methods for TB, especially in remote areas with limited laboratory and x-ray facilities, adds to existing problems [4]. Diagnosis of tuberculosis in children aged 0 – 14 years without specific symptoms is a big challenge for doctors, academics, and program managers because it can hinder diagnosis and lead to worse treatment outcomes [6]. Current diagnostic methods have limited sensitivity, especially in children aged 0 – 14 years [7]. The scoring system, which is an alternative for diagnosing TB in children, is not yet fully implemented according to guidelines [3,8].

Despite clinical and biological variables, there are several socioeconomic and environmental variables that increase the likelihood of TB in children between the ages of 0 and 14. These risk factors are very useful for predicting the development of TB disease. Therefore, knowing the risk factors for TB in children is crucial for making the diagnosis and initiating treatment as soon as possible for the best outcomes. Community health centers in Indonesia are the main place where TB patients seek treatment, so they are the starting point for TB case care which includes early diagnosis, treatment, and reporting. The purpose of this study is to assess the sociodemographic, socioeconomic, and clinical risk factors for TB in children ages 0 to 14 who exhibit one or more TB-related clinical symptoms.

Methods

Research design

This research was conducted at a community health center in the administrative area of South Jakarta, Indonesia. We used a cross-sectional design and studied children with normal nutritional status or malnutrition, including all genders, aged 0 to 14 years, who came with a cough to receive treatment at the community health centers, between January and December 2023. Respondents were determined using consecutive sampling and the sample size was determined using the total sampling method, all respondents who met the inclusion criteria would be included in the study. TB diagnosis is categorized based on clinical diagnosis (accompanied by x-rays and/or tuberculin tests) and bacteriology. The clinical diagnosis was determined by the occurrence or absence of symptoms of a persistent cough lasting more than 2 weeks, weight depletion or no weight gain in the previous two months, fever ≥ 2 weeks, and malaise ≥ 2 weeks. Socio-demographic factors (age, gender, age group, weight, height, nutritional status) were recorded during the history taking. Contact with active TB cases, history of BCG immunization and exposure to cigarette smoke/air pollution were categorized dichotomously. Socio-economic factors (education, employment and parental income, and health insurance) were also categorized dichotomously. This study also recorded the presence or absence of comorbid factors (HIV and DM). A child's nutritional status is determined by the body mass index, or BMI, is calculated using the ratio between body weight in kilograms and height in meters squared. Furthermore, the BMI value of children aged 0-60

months must be compared with the WHO 2005 standard BMI value (WHO, 2006); and BMI values for children and adolescents (5–19 years old) in comparison to the WHO/NCHS 2007 reference (WHO, 2007).

Data analysis

All data that had been collected and entered standard Microsoft Excel was then analyzed using SPSS for Windows (SPSS Inc version 25.0, Chicago, IL). Variables were initially examined univariately, and then a Chi-square statistic was used to evaluate categorical variables in order to determine risk factors in children with tuberculosis. The McNemar statistic was used to estimate the proportion of risk variables between TB and non-TB patients. A statistically significant probability if smaller than 0.05. The odds ratio (OR) and 95% confidence interval (CI) were computed to ascertain the strength of the link.

Results

In this study, 113 out of 156 respondents were diagnosed with pulmonary TB from January to December 2023. Of the TB cases who received treatment, 72.4% of respondents were clinically diagnosed with pulmonary TB. Meanwhile, bacteriologically confirmed TB cases were found in 10.3% of respondents and 89.7% of respondents were unable to produce phlegm. Most of responders (62.8%) were boys, with 34% being less than five years old and 66% being older than five (**Table 1**).

The nutritional status was assessed using the BMI, 64.7% of respondents had a BMI <18.5 and it was significantly associated with the incidence of childhood TB ($p<0.05$). A previous record of interaction with active tuberculosis patients was reported by 54.5% of the respondents. The proportion of TB and non-TB occurrences in the child group differs significantly when exposed to cases of active TB ($p\text{-value}=0.002$). Most respondents received BCG immunization, namely 73.1%. Respondents more often had a history of exposure to cigarette smoke, namely 57.1%, but all respondents did not have a history of HIV and diabetes (**Table 1**).

Most respondents came to the health center with cough symptoms, namely 79.5% and the remaining 20.5% did not experience cough complaints. The change in body weight was not much different, only 49.4% of respondents experienced weight loss. 55.8% of respondents had fever and 44.2% did not have fever, malaise was present in 59.6% of respondents and 40.4% did not have malaise (**Table 1**). All respondents did not use private insurance, but most respondents used BPJS insurance facilities, 70.5%, while 29.5% did not have insurance. Most of the patient's household income corresponds to or exceeds the provincial minimum wage (UMP) of 60.3%. Almost all the respondents' parents had jobs, 96.8% of whom were grouped as non-PNS 85.9% and PNS 10.9%, but only 14.1% had higher education (**Table 2**).

Most respondents were diagnosed with TB based on clinical diagnosis, because it was difficult to obtain children's sputum specimens for bacteriological examination. A bacteriological diagnosis could only be made in 10.3% of respondents and the majority (81.25%) were >10 years old. According to sociodemographic data (**Table 1**) indicates that there are statistically significant variations in proportions of boys and girls ($p<0.001$), body weight ($p<0.001$), height ($p<0.001$). This study showed a notable association between the incidence of TB and nutritional status ($p=0.035$), exposure to cigarette smoke or air pollution ($p=0.006$), but not significantly with BCG immunization ($p=0.328$).

Regarding socioeconomic factors (**Table 2**), TB in childhood occurs more often in families with lower parental education ($p=0.005$), parents who have jobs ($p<0.001$), parental income greater

than or equal to the UMP ($p < 0.001$) and have health insurance ($p < 0.001$). The association among clinical features and the incidence of pediatric TB patients with bacteriological diagnosis is displayed in **Table 3**. Symptoms of cough ($p < 0.001$), subfebrile fever ($p < 0.001$), malaise ($p < 0.001$), and weight loss or no weight gain ($p < 0.001$) showed significant differences in proportion between TB and non-TB events.

Discussion

The purpose of this cross-sectional study was to identify the predictive variables for both TB and non-TB diagnoses among children aged 0–14 years in Community Health Center facilities in Indonesia. TB risk factors such as HIV and diabetes were not found in child respondents, so no analysis was carried out.

Participants in this study underwent two diagnostic methods, namely clinical and bacteriological. Negative smear test results reportedly contribute to 13–20% of transmission and are, on average, 20–25% more infectious than those with positive test results [9]. Previous studies reported that most TB cases were diagnosed based on clinical criteria (50–58%) [10,11]. Likewise, a previous cohort study found 16% (58/372), 95% CI, 12–19% of cases came from smear negative sources [9], meaning 84% of TB cases were smear positive. However, another study reported that 65.2% of TB patients had negative BTA results and 34.8% had positive BTA results [12]. In children, TB disease is generally determined based on clinical diagnosis, which is the accepted method when bacteriological evidence is not available [13]. As is often reported, most respondents were boys. Various studies have similar findings where TB cases occur more often in boys [12,14,15] and variations in the age range of children who contract TB. Younger children are believed to be more affected than older children [16–19] due to household contacts of active TB patients and immature immunity. However, exposure to active TB can also occur in the community. Boys have a tendency for physical activity and high socialization, so they enjoy playing outside the house. This causes boys to potentially encounter adult active TB and be exposed to cigarette smoke or air pollution [20,21].

Malnutrition and tuberculosis (TB) have a reciprocal association in which malnutrition raises the likelihood of getting active TB by six to ten times and TB contributes patient to malnutrition [22,23]. Patients with active TB tend to be underweight, having a BMI below 18.5 kg/m², which is considered an index of malnutrition. Thus, TB incidence is strongly correlated with poor nutrition and lowly BMI [23]. The possibility of TB increases due to malnutrition, both micro and macro deficits, compromised immunity. Being infected with TB can intensify metabolism and require more energy, which can lead to ease appetites and nutrition malabsorption, ultimately raising the possibility of underweight [22,24]. Previous study found that, compared to the control group, TB patients had a higher likelihood of having a BMI of less than 18.5 kg/m² [25]. Numerous countries where tuberculosis prevalence is elevated, most children are exposed to cigarette smoke or secondhand smoke. A comprehensive analysis encompassing eighteen observational studies revealed that exposure to tobacco-related smoke resulted in children being 3.41 times more likely to contract tuberculosis and a 1.64-times more likely to develop latent tuberculosis infection [26]. Research in Spain reported that exposure to cigarette smoke in children (95% CI: 9.63–18.17) and had an increased prevalence of TB (13.48%) [27]. Higher vulnerability to pulmonary tuberculosis illness has been attributed to several biological factors, including poor mucosal secretion clearance, impaired alveolar macrophage phagocytic capacity, diminished immunological reaction, and/or CD4+ lymphopenia brought on by cigarette nicotine [28,29]. Another study that looked at the relationship between TB disease and exposure to cigarette smoke also reported that the detrimental impact of smoking on tuberculosis transmission was significantly

reduced immediately after smoking cessation, hence highlighting the value of smoking cessation programs in the fight against TB [30]. Case-control studies carried out in Brazil and India have shown that firewood or biomass smoke, apart from cigarette smoke, is a separate threat to TB. But there is currently still a dearth of information about the pathway via which biomass smoke results in chronic pulmonary illness [31,32].

Noticing a high connection between TB disease in children and a history of interaction with the source of TB cases is crucial since it is well-known that the risk of developing TB is directly proportionate to the quantity of germs one receives exposure to [33]. Multivariate analysis in previous research showed a significant relationship between household contacts of children with active TB, obtained OR = 15,288 and 95% CI: 5,378–43,457, household contacts with smokers obtained OR = 7,094 and 95% CI: 2,128–23,648, contacts > 18 hours with TB sufferers obtained OR = 4.681 and 95% CI: 1.198–18.294 [34]. Children beneath the age of five, especially under 1 year, who meet a source of TB cases usually have a 50% chance of contracting TB, and ninety-five percent of TB occurs within a year after contact. Home contacts, caregivers, and medical professionals are considered close contacts of active tuberculosis cases [26] because they have a greater chance of contracting *Mycobacterium tuberculosis* and developing primary active tuberculosis [35,36].

The sole licensed and amply used vaccine to prevent TB promoted by *M. tuberculosis* is Bacillus Calmette-Guérin (BCG) [37,38]. The BCG immunization has proven to be highly effective in preventing meningitis and miliary TB in children, with a 70 to 80 percent success rate [37, 39]. An 18-year cohort study in Tunisia found 36.5% of TB cases in individuals who received BCG vaccination compared to 63.3% of unvaccinated individuals [40]. The World Health Organization (WHO) lists fever, inexplicable weight decrease, lethargy or exhaustion, sweating at night, chest pain, and cough lasting longer than two weeks as manifestation and indicators of TB. Given the risk of overdue diagnosis in those infected and subsequent spread via close contacts, overtime cough is the most noticed manifestation in tuberculosis thus summons inquiry, particularly in resource-constrained states where tuberculosis incidence rises (defined as ≥ 100 in 100,000 inhabitants) [41]. In the study population conducted in the Philippines, providing positive TB test results, 84.9% experienced cough ≥ 2 weeks, fever ≥ 2 weeks of unexplained cause (6.9%), considerable and unintended weight loss (9.5%), and 6.3% of patients experienced exhaustion, sluggishness, and malaise [42]. Study finding concluded that all participants with cough symptoms of whichever length for TB examining versed rose sensitivity from forty-two percent to fifty-one percent [43].

This research shows that 70.5% of respondents who have national health insurance facilities (BPJS), 75.5% of respondents tested positive for pulmonary TB. Meanwhile, research in the Philippines reported that 16% of individuals who tested positive for TB were enrolled in a health insurance program (51%) [42]. Since TB is known to be a disease of poverty, national health insurance program in Indonesia's both improves the capacity of TB patients to seek treatment and contributes significantly to TB reporting rate. The study's finding revealed the respondents' parents who were positive for TB had non-PNS jobs with a household income equal to or more than the provincial minimum wage but had low education. A systematic review of studies found a correlation between tuberculosis and a poor educational attainment with a pooled odds ratio = 2.34 [44]. In a South African study including adolescents aged 12 to 18, characteristics that were predictive of TB were older age, ethnic background, low household earning, and basic schooling or lesser for both parents [45]. Due to the correlation with the increasing prevalence of TB, socioeconomic determinants are linked to the availability of healthcare services, the capacity to

fulfill nutrition demands, also the availability of a decent and standardized living space [46]. Working parents are linked to external environment and tuberculosis infection, which can spread to children. Exposure that children obtain from a household member who has TB is nearly fourfold inclined to contract the disease [47,48].

Conclusion

According to the study's findings, children between the ages of 0 and 14 who have a history of cough, fever, malaise, and weight loss, together with malnourished conditions, will increase their risk of developing tuberculosis. Despite the modest study, the results underscore the utility of sociodemographic, socioeconomic, active TB contacts, nutritional status, and exposure to cigarette smoke or air pollution factors in diagnosing TB in children in addition to TB-related clinical symptoms. The use of these simple parameters can help develop digital technology-based diagnostic tools that have the potential to improve early detection of tuberculosis in children in community health center with limited resources and facilities.

Study limitations

This research was conducted at a community health center and there were no patient home visits so it did not analyze environmental factors which are one of the determinants of TB. However, other factors managed to show meaningful results. Additionally, children have difficulty understanding sputum discharge, thereby limiting confirmed cases.

Ethical considerations

All procedures involving patients in this study were approved by the Ethical Review Committee of the Faculty of Medicine, Trisakti University No.055/KER/FK/I/2024. Informed consent was signed by all patients' legal guardians.

Acknowledgments

Not Applicable.

Conflict of Interest

The authors have no conflicts of interest to declare.

Availability of data and materials

The datasets utilized and/or analyzed in the present study are available from the corresponding author on reasonable request.

Authors' contributions

M.M: study concept, writing up, data interpretation, methodology writing, statistical analysis, revise manuscript, and approved the final version.

A.B: study concept, revise manuscript, and approved the final version.

Y.Y: data collection, revise manuscript, and approved the final version.

References

1. WHO Global Tuberculosis Report [Internet]. Geneva: World Health Organization; 2023. Available at: <https://www.who.int/teams/global-tuberculosis-programme/data>. Accessed on Mei 2nd, 2024.

2. Richens, J.G., Lee, C.M. & Johri, S. Improving the accuracy of medical diagnosis with causal machine learning. *Nat Commun* 11, 3923 (2020). Doi: 10.1038/s41467-020-17419-7.
3. Keboysterian Kesehatan RI. Direktorat Jenderal Pencegahan dan Pengendalian Penyakit Petunjuk Teknis Tata Laksana Tuberkulosis Anak dan Remaja [Internet]. Jakarta: Keboysterian Kesehatan RI. 2023. Available at <https://tbindonesia.or.id/wp-content/uploads/2024/02/Final-Petunjuk-Teknis-Tata-Laksana-TBC-Anak-Remaja-2023.pdf>
4. UNICEF Indonesia. Desk Review: Pediatric Tuberculosis with a Focus on Indonesia [Internet]. 2022. Accessed on Mei 2nd, 2024. Available at: <https://www.unicef.org/indonesia/reports/desk-review-pediatric-tuberculosis-focus-indonesia>
5. Basile, F.W., Nabeta, P., Ruhwald, M., Song, R. (2022). Pediatric Tuberculosis Diagnostics: Present and Future. *J Pediatric Infect Dis Soc*, 11(Suppleboyst_3), S85-S93. doi: 10.1093/jpids/piac082
6. Siddalingaiah N, Chawla K, Nagaraja SB, Hazra D. Risk factors for the development of tuberculosis among the pediatric population: a systematic review and meta-analysis. *Eur J Pediatr*. 2023 Jul;182(7):3007-3019. doi: 10.1007/s00431-023-04988-0.
7. Namuganga AR, Chegou NN, Mayanja-Kizza H. Past and Present Approaches to Diagnosis of Active Pulmonary Tuberculosis. *Front Med (Lausanne)*. 2021 Sep 23; 8: 709793. doi: 10.3389/fmed.2021.709793. PMID: 34631731; PMCID: PMC8495065.
8. WHO consolidated guidelines on tuberculosis: Module 5: Management of tuberculosis in children and adolescents [Internet]. Geneva: World Health Organization; 2022. Available at https://www.ncbi.nlm.nih.gov/books/NBK579387/pdf/Bookshelf_NBK579387.pdf. Accessed on Mei 2nd, 2024.
9. Asadi, L., Croxson, M., Heffernan, C., Dhillon, M., Paulsen, C., Egedahl, M. L., Tyrrell, G., Doroshenko, A., & Long, R. (2022). How much do smear-negative patients really contribute to tuberculosis transmissions? Re-examining an old question with new tools. *EClinicalMedicine*, 43, 101250. <https://doi.org/10.1016/j.eclinm.2021.101250>
10. Chilyabanyama, R., Kamanga, N., & Mwandia, J. N. (2024). Factors associated with tuberculosis treatment outcomes among TB patients aged 15 years and older at chawama level one hospital in Lusaka, Zambia. *Global Public Health*, 19(1). <https://doi.org/10.1080/17441692.2024.2307979>
11. Zürcher K, Ballif M, Kiertiburanakul S, Chenal H, Yotebieng M, Grinsztejn B, Michael D, Sterling TR, Ngonyani KM, Mandalakas AM, Egger M, Pettit AC, Fenner L; International Epidemiology Databases to Evaluate AIDS (IeDEA) consortium. Diagnosis and clinical outcomes of extrapulmonary tuberculosis in antiretroviral therapy programmes in low- and middle-income countries: a multicohort study. *J Int AIDS Soc*. 2019 Sep;22(9): e25392. doi: 10.1002/jia2.25392
12. Campos LC, Rocha MVV, Willers DMC, Silva DR (2016) Characteristics of Patients with Smear-Negative Pulmonary Tuberculosis (TB) in a Region with High TB and HIV Prevalence. *PLoS ONE* 11(1): e0147933. <https://doi.org/10.1371/journal.pone.0147933>
13. Abdullahi, O., Moses, N., Sanga, D. *et al*. The effect of empirical and laboratory-confirmed tuberculosis on treatment outcomes. *Sci Rep* 11, 14854 (2021). <https://doi.org/10.1038/s41598-021-94153-0>
14. Chidambaram, V., Tun, N. L., Majella, M. G., Castillo, J. R., Ayeh, S. K., Kumar, A., Neupane, P., Sivakumar, R. K., Win, E. P., Abbey, E. J., Wang, S., Zimmerman, A., Blanck, J., Gupte, A., Wang, J.-Y., & Karakousis, P. C. (2021). Sex differences in TB treatment outcomes:

- Retrospective cohort study and meta-analysis. *medRxiv*, 2021.04.26.21256155. <https://doi.org/10.1101/2021.04.26.21256155>
15. Alao MA, Maroushek SR, Chan YH, Asinobi AO, Slusher TM, Gbadero DA. Treatment outcomes of Nigerian patients with tuberculosis: A retrospective 25-year review in a regional medical center. *PLoS One*. 2020 Oct 29;15(10): e0239225. doi: 10.1371/journal.pone.0239225
 16. Vukugah, T. A., Akoku, D. A., Tchoupa, M. M., & Lambert, E. (2022). Epidemiology of Pediatric Tuberculosis and Factors Associated with Unsuccessful Treatment Outcomes in the Centre Region of Cameroon: A Three-Year Retrospective Cohort Study. *Interdisciplinary perspectives on infectious diseases*, 2022, 2236110. <https://doi.org/10.1155/2022/2236110>
 17. Ogbudebe CL, Adepoju V, Ekerete-Udofia C, Abu E, Egesemba G, Chukwueme N, Gidado M. Childhood Tuberculosis in Nigeria: Disease Presentation and Treatment Outcomes. *Health Serv Insights*. 2018; 11: 1178632918757490. doi: 10.1177/1178632918757490
 18. Tilahun, G., & Gebre-Selassie, S. (2016). Treatment outcomes of childhood tuberculosis in Addis Ababa: a five-year retrospective analysis. *BMC public health*, 16, 612. <https://doi.org/10.1186/s12889-016-3193-8>
 19. Hamid M, Brooks MB, Madhani F, Ali H, Naseer MJ; Childhood Tuberculosis Karachi Group; Becerra M, Amanullah F. Risk factors for unsuccessful tuberculosis treatment outcomes in children. *PLoS One*. 2019 Sep 25;14(9): e0222776. doi: 10.1371/journal.pone.0222776
 20. Martínez-Andrés M, Bartolomé-Gutiérrez R, Rodríguez-Martín B, Pardo-Guijarro MJ, Martínez-Vizcaíno V. (2017). "Football is a boys' game": children's perceptions about barriers for physical activity during recess time. *Int J Qual Stud Health Well-being*, 12(sup2): 1379338. doi: 10.1080/17482631.2017.1379338.
 21. Miller, P.B., Zalwango, S., Galiwango, R. et al. (2021). Association between tuberculosis in men and social network structure in Kampala, Uganda. *BMC Infect Dis*, 21, 1023. <https://doi.org/10.1186/s12879-021-06475-z>
 22. Feleke, B. E., Feleke, T. E., & Biadlegne, F. (2019). Nutritional status of tuberculosis patients, a comparative cross-sectional study. *BMC pulmonary medicine*, 19(1), 182. <https://doi.org/10.1186/s12890-019-0953-0>
 23. Musuenge, B. B., Poda, G. G., & Chen, P. C. (2020). Nutritional Status of Patients with Tuberculosis and Associated Factors in the Health Centre Region of Burkina Faso. *Nutrients*, 12(9), 2540. <https://doi.org/10.3390/nu12092540>
 24. Vonasek, B. J., Radtke, K. K., Vaz, P., Buck, W. C., Chabala, C., McCollum, E. D., Marcy, O., Fitzgerald, E., Kondwani, A., & Garcia-Prats, A. J. (2022). Tuberculosis in children with severe acute malnutrition. *Expert review of respiratory medicine*, 16(3), 273–284. <https://doi.org/10.1080/17476348.2022.2043747>
 25. Kitonsa, P. J., Nalutaaya, A., Mukibi, J., Nakasolya, O., Isooba, D., Kamoga, C., Baik, Y., Robsky, K., Dowdy, D. W., Katamba, A., & Kendall, E. A. (2020). Evaluation of underweight status may improve identification of the highest-risk patients during outpatient evaluation for pulmonary tuberculosis. *PloS one*, 15(12), e0243542. <https://doi.org/10.1371/journal.pone.0243542>
 26. Patra J, Bhatia M, Suraweera W, Morris SK, Patra C, Gupta PC, et al. (2015) Exposure to Second-Hand Smoke and the Risk of Tuberculosis in Children and Adults: A Systematic Review and Meta-Analysis of 18 Observational Studies. *PLoS Med* 12(6): e1001835. <https://doi.org/10.1371/journal.pmed.1001835>
 27. Altet, N., Latorre, I., Jiménez-Fuentes, M. Á., Soriano-Arandes, A., Villar-Hernández, R., Milà, C., Rodríguez-Fernández, P., Muriel-Moreno, B., Comella-Del-Barrio, P., Godoy, P.,

- Millet, J. P., de Souza-Galvão, M. L., Jiménez-Ruiz, C. A., Domínguez, J., & On Behalf Of Pii Smoking Separ Working Group (2022). Tobacco Smoking and Second-Hand Smoke Exposure Impact on Tuberculosis in Children. *Journal of clinical medicine*, 11(7), 2000. <https://doi.org/10.3390/jcm11072000>
28. Burusie, A., Enquesilassie, F., Addissie, A., Dessalegn, B., & Lamaro, T. (2020). Effect of smoking on tuberculosis treatment outcomes: A systematic review and meta-analysis. *PloS one*, 15(9), e0239333. <https://doi.org/10.1371/journal.pone.0239333>
 29. Wang, E. Y., Arrazola, R. A., Mathema, B., Ahluwalia, I. B., & Mase, S. R. (2020). The impact of smoking on tuberculosis treatment outcomes: a meta-analysis. *The international journal of tuberculosis and lung disease: the official journal of the International Union against Tuberculosis and Lung Disease*, 24(2), 170–175. <https://doi.org/10.5588/ijtld.19.0002>
 30. Chu, A. L., Lecca, L. W., Calderón, R. I., Contreras, C. C., Yataco, R. M., Zhang, Z., Becerra, M. C., Murray, M. B., & Huang, C. C. (2021). Smoking Cessation in Tuberculosis Patients and the Risk of Tuberculosis Infection in Child Household Contacts. *Clinical infectious diseases: an official publication of the Infectious Diseases Society of America*, 73(8), 1500–1506. <https://doi.org/10.1093/cid/ciab504>
 31. Báez-Saldaña R, Canseco-Raymundo A, Ixcot-Mejía B, Juárez-Verdugo I, Escobar-Rojas A, Rumbo-Nava U, Castillo-González P, León-Dueñas S, Arrieta O. Case-control study about magnitude of exposure to wood smoke and risk of developing lung cancer. *Eur J Cancer Prev*. 2021 Nov 1;30(6):462-468. doi: 10.1097/CEJ.0000000000000644
 32. Sarkar M, Srinivasa, Madabhavi I, Kumar K. Tuberculosis associated chronic obstructive pulmonary disease. *Clin Respir J*. 2017 May;11(3):285-295. doi: 10.1111/crj.12621
 33. Shaw J.B., Wynn-Williams N. Infectivity of pulmonary tuberculosis in relation to sputum status. *Am Rev Tuberc*. 1954 May;69(5):724-32. doi: 10.1164/art.1954.69.5.724
 34. Laghari, M., Sulaiman, S.A.S., Khan, A.H. *et al*. Contact screening and risk factors for TB among the household contact of children with active TB: a way to find source case and new TB cases. *BMC Public Health* 19, 1274 (2019). <https://doi.org/10.1186/s12889-019-7597-0>
 35. Rea, E., Huard, J., & Lee, R. (2022). Chapter 11: Tuberculosis contact investigation and outbreak management. *Canadian Journal of Respiratory, Critical Care, and Sleep Medicine*, 6(sup1), 167–183. <https://doi.org/10.1080/24745332.2022.2037909>
 36. Reichler, M. R., Khan, A., Sterling, T. R., Zhao, H., Moran, J., McAuley, J., Bessler, P., Mangura, B., & Tuberculosis Epidemiologic Studies Consortium Task Order 2 Team (2018). Risk and Timing of Tuberculosis Among Close Contacts of Persons with Infectious Tuberculosis. *The Journal of infectious diseases*, 218(6), 1000–1008. <https://doi.org/10.1093/infdis/jiy265>
 37. Katelaris AL, Jackson C, Southern J, Gupta RK, Drobniewski F, Lalvani A, Lipman M, Mangtani P, Abubakar I. Effectiveness of BCG Vaccination Against Mycobacterium Tuberculosis Infection in Adults: A Cross-sectional Analysis of a UK-Based Cohort. *J Infect Dis*. 2020 Jan 1;221(1):146-155. doi: 10.1093/infdis/jiz430
 38. Casey RM, Dumolard L, Danovaro-Holliday MC, Gacic-Dobo M, Diallo MS, Hampton LM, Wallace AS. Global Routine Vaccination Coverage, 2015. *MMWR Morb Mortal Wkly Rep*. 2016 Nov 18;65(45):1270-1273. doi: 10.15585/mmwr.mm6545a5
 39. Trunz BB, Fine P, Dye C. Effect of BCG vaccination on childhood tuberculous meningitis and miliary tuberculosis worldwide: a meta-analysis and assessment of cost-effectiveness. *Lancet*. 2006 Apr 8;367(9517):1173-80. doi: 10.1016/S0140-6736(06)68507-3

40. Bennasrallah C, Kacem M, Dhouib W, Zemni I, Ben Fredj M, Abroug H, et al. (2019) BCG vaccination and tuberculosis prevention: A forty-year cohort study, Monastir, Tunisia. *PLoS ONE* 14(8): e0219991. <https://doi.org/10.1371/journal.pone.0219991>
41. Field, S. K., Escalante, P., Fisher, D. A., Ireland, B., Irwin, R. S., & CHEST Expert Cough Panel (2018). Cough Due to TB and Other Chronic Infections: CHEST Guideline and Expert Panel Report. *Chest*, 153(2), 467–497. <https://doi.org/10.1016/j.chest.2017.11.018>
42. Lee, S., Lau, L., Lim, K., Ferma, J., Dodd, W., & Cole, D. (2019). The Presence of Cough and Tuberculosis: Active Case Finding Outcomes in the Philippines. *Tuberculosis research and treatment*, 2019, 4578329. <https://doi.org/10.1155/2019/4578329>
43. Cheng J, Wang L, Zhang H, Xia Y. Diagnostic value of symptom screening for pulmonary tuberculosis in China. *PLoS One*. 2015 May 22;10(5): e0127725. doi: 10.1371/journal.pone.0127725.
44. Dana, N. R., Rika, S., M, I. P., Alexander, M. B., Muthia, S., Linda, R., Astri, W., Zuhrah, T., Rahman, A. D., Rahmi, F., Nomira, P., Setia, N. D. A. W., Arif, L. B. L., Ainil, M., Octarini, E. M., Nova, L., Tamia, M. Y., Sri, R. A., Permata, S. A., Mimin, O., ... Fariz, A. M. (2024). Modifiable and non-modifiable risk factors for tuberculosis among adults in Indonesia: A Systematic review and Meta-analysis. *African journal of infectious diseases*, 18(2), 19–28. <https://doi.org/10.21010/Ajidv18i2.3>
45. Tchakounte Youngui B, Tchounga BK, Graham SM, Bonnet M. Tuberculosis Infection in Children and Adolescents. *Pathogens*. 2022 Dec 9;11(12):1512. doi: 10.3390/pathogens11121512.
46. Khaliq, A., Khan, I.H., Akhtar, M.W., & Chaudhry, M.N. (2015). Environmental Risk Factors and Social Determinants of Pulmonary Tuberculosis in Pakistan. *Epidemiology (Sunnyvale)* 2015;2015(5):3. doi: <https://doi.org/10.4172/2161-1165.1000201>
47. Long R., Diwangahi M., Schwartzman K. Chapter 2: Transmission and Pathogenesis of Tuberculosis. *Can. J. Breathing. Critics. Sleep Care Med.* 2022; 6 :22–32. doi: 10.1080/24745332.2022.2035540.
48. Martinez L., Shen Y., Mupere E., Kizza A., Hill PC, Whalen CC Transmission of Mycobacterium tuberculosis in Households and Communities: A Systematic Review and Meta-Analysis. *I. J. Epidemiol.* 2017; 185 :1327–1339. doi: 10.1093/aje/kwx025.



Table 1. Karakteristik terkait faktor sosiodemografi, kontak TB, imunisasi BCG, dan status nutrisi diantara responden TB dan non-TB.

Karakteristik	Frekuensi		TB	Non-TB	p-Value	Odd Ratio (OR)	95% CI
	N, mean	%, SD					
Diagnosis bakteriologis							
1. Ya	16	10.3	16	0	0.009*	-	-
2. Tidak	140	89.7	97	43			
Usia (tahun)	6.1795	3.29317					
1. <5 tahun	53	34	38	15	0.000[§]	0.964	0.594 – 1.564
2. ≥5 tahun	103	66	75	28		1.019	0.790 – 1.315
Jenis kelamin							
1. Laki-laki	98	62.8	72	26	0.000[§]	1.054	0.797 – 1.393
2. Perempuan	58	37.2	41	17		0.918	0.589 – 1.429
Berat Badan (kg)	19.6218	8.94918			0.000*	-	18.21 – 21.04
Tinggi Badan (cm)	104.3910	26.23235			0.000*	-	100.24 – 108.54
Indeks Massa Tubuh (IMT)	17.5742	2.87826					
1. <18.5	101	64.7	91	10	0.000*	3.463	1.997 – 6.005
2. >18.4	55	35.3	22	33		0.254	0.168 – 0.382
Kontak pasien TB aktif							
1. Ada	85	54.5	62	23	0.002[§]	1.026	0.741 – 1.420
2. Tidak ada							

	71	45.5	51	20		0.970	0.664 – 1.418
Imunisasi BCG							
1. Ya	114	73.1	85	29	0.328*	1.115	0.883 – 1.408
2. Tidak	42	26.9	28	14		0.761	0.445 – 1.302
Status gizi							
1. Gizi kurang	100	64.1	90	10	0.000*	3.425	1.974 – 5.941
2. Normal	56	35.9	23	33		0.265	0.178 – 0.396
Paparan rokok/polusi udara							
1. Ada	89	57.1	65	24	0.006 [§]	1.031	0.756 – 1.405
2. Tidak ada	67	42.9	48	19		0.961	0.645 – 1.432

*Chi-square test, p-value <0.05; [§]McNemar test, p-value <0.05

Table 2. Karakteristik terkait faktor sosioekonomi, gejala klinis, dan paparan asap diantara responden TB dan non-TB.

Karakteristik	Frekuensi		TB	Non-TB	p-Value*	Odd Ratio (OR)	95 % CI
	N, mean	%, SD					
Pendidikan orangtua							
1. Rendah	134	85.9	97	37	0.005 [§]	0.998	0.66 – 1.150
2. Tinggi	22	14.1	16	6		1.015	0.425 – 2.422
Pekerjaan orangtua							
• Tidak bekerja	5	3.2	4	1	0.000 [§]	1.522	0.175 – 13.238
• Bekerja	151	96.8	109	42		0.988	0.932 – 1.047
Pendapatan rumah tangga/orangtua							
• < UMP	62	39.7	42	20	0.000 [§]	0.799	0.536 – 1.192
• ≥ UMP	94	60.3	71	23		1.175	0.859 – 1.606
Asuransi kesehatan							
1. Tidak ada	46	29.5	30	16	0.000 [§]	0.713	0.435 – 1.170
2. BPJS	110	70.5	83	27		1.170	0.906 – 1.510
Batuk kronis ≥2 minggu							
1. Ada	124	79.5	90	34	0.937*	1.007	0.842 – 1.206
2. Tidak ada	32	20.5	23	9		0.972	0.490 – 1.931
BB turun atau tidak naik dalam 2 bulan sebelumnya							

1. Ada 2. Tidak ada	77 79	49.4 50.6	60 53	17 26	0.000[§]	1.343 0.776	0.893 – 2.020 0.568 – 1.059
Demam subfebris ≥2 minggu							
1. Ada 2. Tidak ada	87 69	55.8 44.2	63 50	24 19	0.003[§]	0.999 1.001	0.731 – 1.365 0.675 – 1.486
Malaise ≥ 2 minggu							
1. Ada 2. Tidak ada	93 63	59.6 40.4	67 46	26 17	0.024[§]	0.981 1.030	0.737 – 1.305 0.669 – 1.585

*Chi-square test, *p-value* <0.05; [§]McNemar test, *p-value* <0.05.

Table 3. Karakteristik terkait faktor klinis diantara responden TB dan non-TB berdasarkan diagnosis bakteriologi.

§McNemar test, *p*-value <0.05

Karakteristik	Frekuensi		TB	Non-TB	p-Value*	Odd Ratio (OR)	95 % CI
	N, mean	%, SD					
Batuk kronis ≥2 minggu							
1. Ada	124	79.5	14	110	0.000§	1.114	0.908 – 1.366
2. Tidak ada	32	20.5	2	30		0.583	0.154 – 2.216
BB turun atau tidak naik dalam 2 bulan sebelumnya							
1. Ada	77	49.4	7	70	0.000§	0.875	0.490 – 1.562
2. Tidak ada	79	50.6	9	70		1.125	0.708 – 1.787
Demam subfebris ≥2 minggu							
1. Ada	87	55.8	11	76	0.000§	1.266	0.880 – 1.822
2. Tidak ada	69	44.2	5	64		0.684	0.323 – 1.446
Malaise ≥ 2 minggu							
1. Ada	93	59.6	10	83	0.000§	1.054	0.704 – 1.578
2. Tidak ada	63	40.4	6	57		0.921	0.474 – 1.788



Revisi 2

ORIGINALITY REPORT

5%

SIMILARITY INDEX

5%

INTERNET SOURCES

4%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

1

repo.stikesicme-jbg.ac.id

Internet Source

1%

2

www.clin-lab-publications.com

Internet Source

1%

3

www.mdpi.com

Internet Source

1%

4

Kedir Usmael, Tsegahun Manyazewal, Hussen Mohammed, Getnet Yimer et al. "Patterns of childhood tuberculosis diagnosis in Ethiopia: A multicenter cross-sectional study", Research Square Platform LLC, 2023

Publication

1%

5

topsecretapiaccess.dovepress.com

Internet Source

1%

6

www.balimedicaljournal.org

Internet Source

1%