

Relationship between Serum 25-Hydroxyvitamin D Levels and Cardiovascular Disease Risk among Road Sanitations Workers

Magdalena Wartono,^{1,2,3} Tjam Diana Samara,^{1,2,3} Adrianus Kosasih,⁴ Dian Mediana^{1,3}

¹Department of Anatomy, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

²Department of Occupational Health, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

³Study Center of Occupational Medicine, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

⁴Department of Internal Disease, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

Abstract

Background: Road sanitation workers are frequently exposed to sunlight during outdoor work, which may influence vitamin D status. Although several studies suggest that outdoor workers have a lower risk of vitamin D deficiency than indoor workers, evidence regarding the association between 25-hydroxy vitamin D [25-(OH)D] levels and cardiovascular disease (CVD) risk among sanitation workers remains limited. This study aimed to examine the relationship between serum 25(OH)D levels and CVD risk among road sanitation workers.

Methods: This analytical cross-sectional study included 105 road sanitation workers in West Cengkareng District, Jakarta, Indonesia selected using consecutive random. The 10-year cardiovascular risk was calculated using the World Health Organization/International Society of Hypertension (WHO/ISH) risk charts. Serum 25-(OH)D levels were measured from peripheral venous blood samples. Data were analyzed using the chi-square test and Fisher's exact test, with statistical significance defined as $p < 0.05$.

Results: Vitamin D deficiency was observed in 59.05% of participants, whereas 85.7% had low-to-moderate CVD risk. No significant association was found between 25(OH)D levels and CVD risk ($p = 0.582$). However, significant associations were identified between age ($p = 0.001$), body mass index ($p = 0.003$), blood pressure ($p = 0.037$), smoking status ($p = 0.037$) and cardiovascular disease risk.

Conclusions: Vitamin D deficiency is not associated with increased CVD risk among road sanitation workers. However, regular monitoring of vitamin D status may support occupational health and overall well-being.

Keywords: Cardiovascular disease risk, occupational health, road sanitation workers, vitamin D

Althea Medical Journal.
2026;13(1):28-34

Received: November 6, 2025
Accepted: February 18, 2026
Published: February 28, 2026

Correspondence:

Magdalena Wartono,
Department of Anatomy
Faculty of Medicine,
Universitas Trisakti,
Jl. Kyai Tapa Bo. 260, Kampus B,
Tomang, Jakarta, Indonesia

E-mail:
magdalena_w@trisakti.ac.id

Introduction

The mortality rate due to cardiovascular disease (CVD) has been increasing in many developing countries.¹ In Indonesia, about 37% of deaths are caused by cardiovascular disease, especially stroke, followed by coronary heart disease.² The number of CVD-related deaths has more than doubled, increasing from 292,000 cases in 1990 to 659,000 cases in 2019. During the same period, the mortality rate increased from 346.4 to 387.2 deaths per 100,000 population, while the prevalence of

CVD increased by 6.1%.³

Cardiovascular disease also contributes substantially to the global burden of disease. In 2012, CVD accounted for approximately 18,000 disability-adjusted life years (DALYs), of which around 17,500 DALYs were attributed to premature mortality, while the remainder resulted from years of healthy life lost due to disability.² Among ASEAN countries, Indonesia has the second-highest DALYs burden from CVD after Laos. At the provincial level, the highest DALYs rates have been reported in Bangka Belitung, followed by South Kalimantan and

Yogyakarta.³

A study conducted in Malang among adults aged 40 years and older reported that the 10-year CVD risk was lowest in rural areas (26.2%) and highest in urban areas (31.6%). By occupation, the highest risk was observed among unemployed individuals, while the lowest was among private-sector employees.⁴ In addition, sanitation workers were reported to have a higher risk of CVD than hotel housekeepers across several cardiovascular risk prediction models.⁵

Major risk factors for cardiovascular disease include hypertension, smoking, obesity, and diabetes mellitus (DM).^{6,7} The World Health Organization (WHO) Global Burden of Disease (GBD) report in 2013 noted that the prevalence of hypertension, hypercholesterolemia, and smoking has increased in low- and middle-income countries, while declining in high-income countries.⁸ According to the Indonesian National Health Survey, approximately one-quarter of the population suffers from hypertension, around 30% were overweight with hypercholesterolemia, 7% had diabetes, most men (65%) were smokers, and 23% of the population had low physical activity levels.²

Another condition that may raise the risk of CVD is deficiency of vitamin D. Several studies have reported associations between serum 25-hydroxyvitamin D [25(OH)D] and various CVD risk factors, including hypertension, dyslipidemia, and diabetes mellitus.⁹ A case-control study conducted in Kansas City revealed that individuals with vitamin D deficiency and no previous myocardial infarction experienced a notably reduced risk of mortality when their (25-OH)D levels increased above 20 ng/mL and 30 ng/mL compared to those who did not receive such supplementation. Individuals who maintained (25-OH)D levels ≥ 30 ng/mL also had a lower risk of myocardial infarction.¹⁰ However, a study conducted in New Zealand found no significant difference in CVD incidence between groups receiving vitamin D supplementation and those receiving placebo.¹¹

Governor Regulation of the Special Capital Region of Jakarta No. 196/2015 outlines the responsibilities of road sanitation workers, who maintain road infrastructure and sidewalks across two shifts (07:00–15:00 and 15:00–23:00 WIB).¹² Given these outdoor working hours, higher levels of ultraviolet B (UVB) exposure and subsequent vitamin D synthesis might be expected. However, research among Korean fishermen, another

outdoor-working cohort, revealed that 78% had suboptimal vitamin D levels, highlighting a discrepancy between outdoor exposure and actual serum levels.¹³

Since data on serum 25(OH)D levels and cardiovascular risks among road sanitation workers remains limited, further investigation is required. Therefore, this study aimed to assess the relationship between serum 25(OH)D levels and CVD risk among sanitation workers in West Cengkareng district, Jakarta, Indonesia.

Methods

This cross-sectional analytical study was performed in West Jakarta, Indonesia in August 2025. The population of the study consisted of Public Facilities and Infrastructure Management officers, commonly known as road sanitation workers, in the West Cengkareng district, Jakarta. Eligible participants were male sanitation workers aged 40–64 years who had worked for at least one year and provided written informed consent to participate in the study. Respondents with a history of consuming cholesterol-lowering medications, anti-diabetic drugs, heart disease medication, or vitamin D supplements were excluded. Individuals with a history of stroke, coronary heart disease, or kidney failure were also excluded. Participants were recruited using a consecutive non-random sampling technique.

Sociodemographic and clinical data, including age, employment status, length of employment, history of comorbidities, and medication use, were collected through structured interviews using a standardized questionnaire.

Blood pressure was measured twice using an automatic sphygmomanometer (OMRON M3, Omron Healthcare Co., Ltd., Kyoto, Japan) after the participants had rested for at least 10 minutes. The final recorded value was used for analysis. Hypertension was defined as systolic blood pressure ≥ 140 mmHg or diastolic blood pressure ≥ 90 mmHg. Participants receiving antihypertensive therapy were also classified as having hypertension.

Body weight and height were measured using a SECA model 700 scale and a SECA 220 stadiometer (SECA GmbH, Hamburg, Germany). Body mass index (BMI) was calculated as weight in kilograms (kg) divided by the square of height in meters (kg/m^2).

The 10-year CVD risk was stratified based on the WHO non-laboratory-based cardiovascular risk charts for the South East

Asia region. Risk stratification was evaluated by determining age, gender, smoking status, systolic blood pressure, and BMI. The predicted 10-year cardiovascular risk was categorized into five groups: low risk (<5%), moderate risk (5–<10%), high risk (10–<20%), very high risk (20–<30%), and extremely high risk (≥30%). For analytical purposes, participants were further grouped into low-to-moderate risk (<10%) and high risk (≥10%).¹⁴

Serum 25-(OH)D levels were measured from peripheral venous blood samples. A serum level ≥30 ng/mL was considered normal, while levels <30 ng/mL was considered vitamin D deficiency.

Statistical analysis was performed using the chi-square statistical test, Fisher's exact test, and logistic regression analysis. Data processing was carried out using IBM SPSS Statistics version 26.0 (IBM Corp., Armonk, NY, USA). A p-value <0.05 was considered statistically significant. Ethical approval for this study was obtained from the Ethics Review Committee of the Faculty of Medicine, Trisakti University (No. 047/KER/FK/08/2024).

Results

Table 1 summarizes the sociodemographic

and clinical characteristics of the respondents (n=105). Most participants were aged 40–49 years (67.6%) and had a normal body mass index (53.4%). A high proportion of respondents presented with elevated blood pressure (79.0%) and a history of smoking (81.0%). The majority had worked for less than 10 years (93.3%). More than half of the respondents had low 25-(OH)D levels (59.1%). Despite these clinical risk factors, most participants were classified as having a low-to-moderate risk of cardiovascular disease (85.7%), while a smaller proportion were categorized as high risk (14.3%).

The analysis revealed that age (p=0.001), smoking habits (p=0.037), BMI (p=0.003) and blood pressure (p=0.037) were significantly associated with CVD risk. Individuals aged 50–59 years had significantly higher odds of high CVD risk compared with those aged 40–49 years (OR=9.10; 95% CI: 2.32–35.65). Overweight individuals were also more likely to have high CVD risk compared with those with normal or underweight BMI (OR=5.50; 95% CI: 1.62–18.73). Smoking habits and blood pressure were significantly associated with cardiovascular disease risk; however, the distribution showed a higher proportion of high-risk cases among non-smokers compared

Table 1 Sociodemographic and Clinical Characteristics of Respondents (n=105)

Characteristic	n	%
Age (years)		
40–49	71	67.6
50–59	34	32.4
Body mass index (BMI)		
Underweight (≤18.49 kg/m ²)	8	7.6
Normal (18.5–24.9 kg/m ²)	56	53.4
Overweight (≥25 kg/m ²)	41	39.0
Blood pressure		
Normal (<120/<80 mmHg)	22	21.0
High (≥120/≥80 mmHg)	83	79.0
Smoking history		
No	20	19.0
Yes	85	81.0
Length of employment		
1–9 years	98	93.3
10–24 years	7	6.7
25-Hydroxyvitamin D		
Normal	43	40.9
Low	62	59.1
Cardiovascular disease risk		
Low-moderate risk	90	85.7
High risk	15	14.3

Table 2 Association Between Respondent Characteristics, 25-Hydroxyvitamin D Levels, and Risk of Cardiovascular Disease

Variable	Low-Moderate Risk n (%)	High Risk n (%)	p-value	OR	95% CI
Age (years)					
40–49	67 (94.4)	4 (5.6)	0.001^	9.10	2.32–35.65
50–59	23 (67.6)	11 (32.4)			
Smoking habits					
No	14 (70.0)	6 (30.0)	0.037^	0.28	0.09–0.90
Yes	76 (89.4)	9 (10.6)			
Length of employment (years)					
1–9	83 (84.7)	15 (15.3)	0.590^	N/A	N/A
10–24	7 (100.0)	0 (0.0)			
Body mass index					
Underweight–normal	60 (93.8)	4 (6.2)	0.003*	5.50	1.62–18.73
Overweight	30 (73.2)	11 (26.8)			
Blood pressure					
Normal	22 (100.0)	0 (0.0)	0.037^	N/A	N/A
High	68 (81.9)	15 (18.1)			
25-hydroxyvitamin D (ng/mL)					
Normal (30–100)	38 (88.4)	5 (11.6)	0.582*	–	–
Low (<30)	52 (83.9)	10 (16.1)			

Notes: CVD = cardiovascular disease; OR = odds ratio; CI = confidence interval; N/A = not applicable. ^ Fisher’s exact test; * Chi-square test

with smokers. In contrast, no significant association was observed between length of employment or 25-hydroxyvitamin D levels and cardiovascular disease risk (Table 2).

Discussion

In this study, 67.6% of the 105 respondents were aged between 40 and 49 years old, placing the majority in middle adulthood, a stage associated with increasing vulnerability to CVD. The incidence of CVD events, including ischemic heart disease (IHD) and stroke, generally remains low before the age of 40 but increases steeply thereafter, particularly among male.¹⁵

Aging is widely recognized as a major risk factor for CVD such as atherosclerosis, stroke, and myocardial infarction in both male and female.¹⁶ Numerous studies have shown that increasing age is associated with a higher risk of CVD. Research conducted in New Zealand found that the heart age gap decreases with advancing age; individuals aged 65–74 years have approximately half the heart age gap observed in individuals aged 35–44 years.¹⁷ Although the heart age gap decreases with age, the absolute risk of short-term heart disease events continues to increase.¹⁷ Thus,

age remains an unmodified and independent risk factor for atherogenesis and also further CVD.¹⁷

This study also demonstrates an association between BMI and cardiovascular disease risk. A higher proportion of overweight individuals are classified as having high CVD risk compared with respondents with underweight or normal BMI, with an odds ratio greater than five. These findings are consistent with previous study, which reported a concomitant rise in atherosclerotic CVD prevalence with increasing BMI, along with other CVD risk factors such as insulin resistance and lipoprotein(a).¹⁸ Similarly, data from an American cohort study indicated that a one-standard deviation increase in BMI variability was correlated with an 8% higher risk of adverse events of CVD.¹⁹

Smoking habits are also significantly associated with CVD risk. In this study, the majority of the non-smoking group are classified as having low-to-moderate CVD risk. The analysis demonstrated that non-smokers had lower odds of experiencing adverse cardiovascular outcomes compared to smokers. Previous evidence supports the strong association between smoking and cardiovascular disease. The Cross-Cohort Collaboration study found that smoking 2–5

cigarettes per day increases the risk of CVD by 26–57% than non-smokers, whereas smoking cessation for more than 20 years reduces the risk by over 80%.²⁰

Blood pressure is also associated with cardiovascular disease risk in this study. All respondents with normal blood pressure are classified as having low-to-moderate CVD risk. Findings from a randomized controlled trial conducted in China indicate that targeting a systolic target of <120 mm Hg reduces major vascular events in high-risk hypertensive patients (independent of diabetes or stroke status) compared with a target of <140 mmHg, although with a slightly increased risk of adverse effects.²¹ These findings highlight the importance of blood pressure control in reducing cardiovascular risk.

Sunlight exposure is the primary natural source of vitamin D, as ultraviolet radiation stimulates vitamin D synthesis in the skin.²² However, previous study among road sanitation workers have shown that many individuals still have vitamin D levels below normal despite frequent outdoor exposure. In this study, 11.6% of respondents with normal 25-hydroxyvitamin D levels had a high risk of CVD compared with 16.1% among those with low vitamin D levels. However, no significant association was found between vitamin D levels and cardiovascular disease risk in this population. These findings are consistent with the study conducted in New Zealand, which reported no significant difference between monthly high-dose vitamin D supplementation and placebo in preventing CVD.¹¹ Similar results were reported in a meta-analysis evaluating vitamin D supplementation and major adverse cardiovascular events.²³

However, another study has reported conflicting findings. An Australian study found that participants receiving vitamin D supplementation had a lower risk of myocardial infarction and coronary revascularization compared with those receiving placebo.²⁴ Another study also suggested that vitamin D deficiency may increase the likelihood of coronary artery disease.²⁵

Vitamin D plays a vital role in maintaining calcium and phosphorus homeostasis. More recently, vitamin D receptors have been identified in various extra-skeletal tissues, suggesting broader physiological effects beyond bone metabolism.^{9,26} Vitamin D may influence cardiovascular health by regulating the renin-angiotensin-aldosterone system (RAAS), endothelial function, and vascular smooth muscle cell activity. Vitamin D

deficiency has also been associated with increased mortality and a higher incidence of cardiovascular disease.²⁶ Although no universal consensus exists regarding optimal vitamin D levels for preventing CVD, serum concentrations above 30 ng/mL are generally considered adequate.²⁷

Vitamin D deficiency may contribute to vascular inflammation through activation of the NF-κB pathway, promoting inflammatory processes in blood vessel walls and epicardial adipose tissue that contribute to atherosclerosis.^{26,28}

Furthermore, vitamin D deficiency has been linked to increased arterial stiffness, endothelial dysfunction, and vascular inflammation, all of which may accelerate atherosclerosis.²⁹ Several studies have suggested that vitamin D may support cardiovascular health by influencing blood pressure regulation through its effects on blood vessel cells. Vitamin D deficiency has also been associated with increased incidence of myocardial infarction, stroke, and peripheral artery disease.⁹

In this study, other risk factors such as length of employment were not significantly related to CVD risk. This may be due to the relatively small number of respondents who had worked for more than nine years. Further studies with larger sample sizes are needed to provide more accurate estimates.

One limitation of this study is the lack of homogeneity among respondents, particularly in terms of age and cumulative exposure to ultraviolet-B radiation, which may influence 25-hydroxyvitamin D levels. These factors could potentially affect the interpretation of the relationship between vitamin D status and cardiovascular risk.

In conclusion, this study demonstrates that vitamin D deficiency is not a factor that increases cardiovascular risk in road sanitation workers. However, age, BMI, blood pressure and smoking habit are associated with increased cardiovascular risk. Public health education remains important to encourage individuals to maintain adequate vitamin D levels and adopt healthy lifestyles to reduce the risk of cardiovascular and other extra-skeletal diseases.

Acknowledgements

The authors would like to express their sincere gratitude to the Cengkareng Subdistrict Administration for their support in facilitating the successful conduct of this study.

Authors' Contributions

MW conceived the research idea, conducted the study, contributed to manuscript drafting, and performed revisions and proofreading. TDS participated in conducting the study, analyzed the data, drafted the manuscript, and assisted with revisions. AK contributed to the research concept and assisted with data collection and data analysis. DM participated in data collection and provided education to respondents.

Conflict of Interest

The authors declare no conflict of interest.

Funding

This study was supported by the Faculty of Medicine, Universitas Trisakti.

Generative AI Disclosure Statement

During the preparation of this work, the authors used Perplexity AI to identify and verify relevant references and Google Gemini to improve English translation. The authors reviewed and edited all generated content as needed and take full responsibility for the final manuscript.

References

1. Harmadha WSP, Muharram FR, Gaspar RS, Azimuth Z, Sulistya HA, Firmansyah F, et al. Explaining the increase of incidence and mortality from cardiovascular disease in Indonesia: A global burden of disease study analysis (2000–2019). *PLoS One*. 2023;18(12):e0294128. doi: 10.1371/journal.pone.0294128.
2. Chow CK, Atkins E, Islam SMS, Lung T. Reducing the burden of cardiovascular disease in Indonesia: evidence review. Newtown: The George Institute for Global Health; 2017. doi: 10.13140/RG.2.2.20937.57445
3. Muharram FR, Multazam CECZ, Mustofa A, Socha W, Andrianto, Martini S, et al. The 30 years of shifting in the Indonesian cardiovascular burden—analysis of the global burden of disease study. *J Epidemiol Glob Health*. 2024;14(1):193–212. doi: 10.1007/s44197-024-00187-8.
4. Maharani A, Sujarwoto, Praveen D, Oceandy D, Tampubolon G, Patel A. Cardiovascular disease risk factor prevalence and estimated 10-year cardiovascular risk scores in Indonesia: The SMARThealth Extend study. *PLoS One*. 2019;14(4):e0215219. doi: 10.1371/journal.pone.0215219.
5. López-González AA, Manzanero Z, González San Miguel HM, Arroyo Bote S, Riutord Sbert P, Rigo Vives M del M, et al. Differences in cardiovascular risk levels between cleaning staff and hotel housekeepers. *J Occup Health*. 2022;64(1):e12320. doi: 10.1002/1348-9585.12320.
6. Chen X, Zhou L, Zhang Y, Yi D, Liu L, Rao W, et al. Risk factors of stroke in Western and Asian countries: a systematic review and meta-analysis of prospective cohort studies. *BMC Public Health*. 2014;14(1):776. doi: 10.1186/1471-2458-14-776.
7. Li Z, Yu S, Han X, Liu J, Yao H. Changes to cardiovascular risk factors over 7 years: a prospective cohort study of in situ urbanised residents in the Chaoyang District of Beijing. *BMJ Open*. 2020;10(3):e033548. doi: 10.1136/bmjopen-2019-033548.
8. Yeates K, Lohfeld L, Sleeth J, Morales F, Rajkotia Y, Ogedegbe O. A global perspective on cardiovascular disease in vulnerable populations. *Can J Cardiol*. 2015(9):1081–93. doi: 10.1016/j.cjca.2015.06.035.
9. Norman PE, Powell JT. Vitamin D and cardiovascular disease. *Circ Res*. 2014;114(2):379–93. doi: 10.1161/CIRCRESAHA.113.301241.
10. Acharya P, Dalia T, Ranka S, Sethi P, Oni OA, Safarova MS et al. The effects of vitamin D supplementation and 25-hydroxyvitamin D levels on the risk of myocardial infarction and mortality. *J Endocr Soc*. 2022;6(1). doi: 10.1210/jendso/bvab164.
11. Scragg R, Stewart AW, Waayer D, Lawes CMM, Toop L, Sluyter J, et al. Effect of monthly high-dose vitamin D supplementation on cardiovascular disease in the vitamin D assessment study. *JAMA Cardiol*. 2017;2(6):608. doi: 10.1001/jamacardio.2017.0175.
12. DKI Jakarta Provincial Government. Governor Regulation No. 169 of 2015 on the Management of Public Infrastructure and Facilities at the Subdistrict Level [Internet]. Jakarta: DKI Jakarta Provincial Government; 2015 [cited 2026 Jan 21]. Available from: <https://peraturan.bpk.go.id/Details/229959/pergub-prov-dki-jakarta-no-169-tahun-2015>.
13. Lee DH, Park KS, Cho MC. Laboratory confirmation of the effect of occupational sun exposure on serum 25-hydroxyvitamin D concentration. *Medicine (Baltimore)*.

- 2018;97:e11419. doi:10.1097/MD.00000000000011419.
14. World Health Organization. Cardiovascular disease risk non-laboratory-based chart [Internet]. WHO; 2019 [cited 2026 Jan 21]. Available from: https://www.who.int/docs/default-source/cardiovascular-diseases/southeast-asia.pdf?sfvrsn=cbee748_2.
15. Ramroth J, Shakir R, Darby SC, Cutter DJ, Kuan V. Cardiovascular disease incidence rates: a study using routinely collected health data. *Cardiooncology*. 2023;9(1):41. doi: 10.1186/s40959-023-00189-8.
16. Brown JC, Gerhardt TE, Kwon E. Treasure Island (FL): StatPearls Publishing. 2024 [cited 2024 Jul 24]. Risk factors for coronary artery disease. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK554410/>.
17. Stjernholm K, Kerr A, Poppe KK, Jensen AE, Mehta S, Nielsen JB, Jackson R, Wells S. Age-stratified comparison of heart age and predicted cardiovascular risk in 370 000 primary care patients. *Heart*. 2025;111(4):166–71. doi: 10.1136/heartjnl-2024-324156.
18. Asztalos BF, Russo G, He L, Diffenderfer MR. Body mass index and cardiovascular risk markers: a large population analysis. *Nutrients*. 2025;17(5):740. doi:10.3390/nu17050740
19. Almuwaqqat Z, Hui Q, Liu C, Al Rifai M, Lima BB, Vaccarino V, et al. Long-term body mass index variability and adverse cardiovascular outcomes. *JAMA Netw Open*. 2024;7(3):e243062. doi:10.1001/jamanetworkopen.2024.3062.
20. Tasdighi E, Yao ZQ, Dardari ZE, Jha KK, Osuji N, Rajan T, et al. Association between cigarette smoking status, intensity, and cessation duration with long-term incidence of nine cardiovascular and mortality outcomes: the cross-cohort collaboration (CCC). *PLoS Med*. 2025;22(11):e1004561. doi:10.1371/journal.pmed.1004561.
21. Liu JM, Li Y, Ge JZ, Yan XF, Zhang HB, Zheng X, et al. Lowering systolic blood pressure to less than 120 mm Hg versus less than 140 mm Hg in patients with high cardiovascular risk with and without diabetes or previous stroke: an open-label, blinded-outcome, randomised trial. *Lancet*. 2024;404:245–
55. doi: 10.1016/S0140-6736(24)01028-6.
22. Kheiri B, Abdalla A, Osman M, Ahmed S, Hassan M, Bachuwa G. Vitamin D deficiency and risk of cardiovascular diseases: a narrative review. *Clin Hypertens*. 2018;24(1):9. doi: 10.1186/s40885-018-0094-4.
23. Barbarawi M, Kheiri B, Zayed Y, Barbarawi O, Dhillon H, Swaid B, et al. Vitamin D supplementation and cardiovascular disease risks in more than 83,000 individuals in 21 randomized clinical trials: a meta-analysis. *JAMA Cardiol*. 2019;4(8):765–76. doi:10.1001/jamacardio.2019.1870.
24. Thompson B, Waterhouse M, English D R, McLeod D S, Armstrong B K, Baxter C et al. Vitamin D supplementation and major cardiovascular events: D-Health randomised controlled trial. *BMJ*. 2023;381:e075230. doi:10.1136/bmj-2023-075230.
25. Xu R, Li YY, Ma LL, Yang HN. Association of vitamin D status with coronary artery disease in postmenopausal women. *Medicine (Baltimore)*. 2020;99(11):e19544. doi:10.1097/MD.00000000000019544.
26. Maretzke F, Bechthold A, Egert S, Ernst JB, Melo van Lent D, Pilz S, et al. Role of vitamin D in preventing and treating selected extraskeletal diseases—an umbrella review. *Nutrients*. 2020;12(4):969. doi: 10.3390/nu12040969.
27. Bikle DD. Vitamin D: Production, Metabolism, and Mechanism of Action. In: Feingold KR, Adler RA, Ahmed SF, et al., editors. *Endotext* [Internet]. South Dartmouth (MA): MDText.com, Inc.; 2000 [Cited 2025 Nov 21] Available from: <https://www.ncbi.nlm.nih.gov/books/NBK278935/>.
28. de la Guía-Galipienso F, Martínez-Ferran M, Vallecillo N, Lavie CJ, Sanchis-Gomar F, Pareja-Galeano H. Vitamin D and cardiovascular health. *Clin Nutr*. 2021;40(5):2946–57. doi: 10.1016/j.clnu.2020.12.025.
29. Haider F, Ghafoor H, Hassan OF, Farooqui K, Bel Khair AOM, Shoaib F. Vitamin D and cardiovascular diseases: an update. *Cureus*. 2023. doi: 10.7759/cureus.49734.

About the Journal



Althea Medical Journal is an online journal published since February 2014 (e-ISSN [2337-4330](https://doi.org/10.24127/amj.v10i4.2337-4330)) by the Faculty of Medicine Universitas Padjadjaran, Bandung, Indonesia.

Althea Medical Journal is indexed in major databases, including Science and Technology Index (SINTA) Rank 2, Directory of Open Access Journals (DOAJ), and [SCOPUS](https://scopus.com).

The journal is published quarterly (February, May, August, and November). Althea Medical Journal, which previously publishes articles related to research in the fields of biomedical science, clinical medicine, family-community medicine, and public health, since March 2024 has only accepted articles related to **aging-**

related diseases, wellness, and healthy lifestyle which includes neurodegenerative diseases, cardiovascular diseases, cancer, immune system disorders, and musculoskeletal disorders.

Update to Publication Schedule

📅 2026-02-03

The Editorial Board of **Althea Medical Journal** announces an update to the journal's **publication schedule**.

Previously, **Althea Medical Journal** was published quarterly in **March, June, September, and December**. Starting from the current publication cycle, the journal will now be published in **February, May, August, and November**.

This adjustment is intended to improve the editorial workflow and to ensure the timely publication of high-quality scholarly articles.

We kindly encourage **authors, reviewers, and readers** to take note of this updated publication schedule. Thank you for your continued support of **Althea Medical Journal**.

[Read More](#) ➔

AMJ TEMPLATE




Manuscript Template

Download the official AMJ template



ACCREDITATION & METRICS

Scopus & SCImago



SINTA Accreditation

Research Articles

Physical Activity Levels Based on Indonesian Population Characteristics: Evidence from the Indonesian Family Life Survey-5

Givana Felicia, Felicia Kumiawan, Yunisa Astiarani, Kevin Kristian, Bryany Titi Santi

Early Metabolic Alterations and Predictors of Obesity Among Young Adults in Indonesia: Focus on Lipid Abnormalities and Cardiometabolic Risk

Mirasari Putri, Eva Rianti Indrasari, Yafma Ferni Helta Heriady, Rizky Suganda Prawiradilaga

Insomnia as a Risk Factor for Cardiovascular Disease Among Adults in Eastern Indonesia

Helty Helty, Firman Firman, Elmukhsinur Elmukhsinur, Wa Ode Aisya Zoharia, Noviati Noviati

Relationship between Serum 25-Hydroxyvitamin D Levels and Cardiovascular Disease Risk among Road Sanitations Workers

Magdalena Wartono, Tjam Diana Samara, Adrianus Kosasih, Dian Mediana

Correlation between Nicotine Dependence and Cardiorespiratory Fitness within the Wellness Matrix Framework among University Students

Muhammad Ikhsan Ghalfazzari Ghafar, Ronny Lesmana, Setiawan Setiawan

Effects of Moderate-Intensity Treadmill Training on Cardiac Mitochondrial Mitophagy and Dynamics in Young and Aged Wistar Rats

Julia Windi Gunadi, Difa Hidayat Ramadhan, Bernike Tirsa Safira, Andra Alifya Ramadhan, Roro Wahyudianingsih, Yenni Limyati, Hanna Goenawan, Ronny Lesmana

Reduced Left Ventricular Global Longitudinal Strain in Obese Adolescents

Rilda Dwi Febrianda, Tisnasari Hafsa, Sri Endah Rahayuningsih, Dzulfikar Djalil Lukmanul Hakim, Rahmat Budi Kuswiyanto, Putria Rayani Apandi, Dida Akhmad Gurnida

Evaluation of Urinary hsa-miR-21-5p Expression as a Non-Invasive Biomarker for Prostate Cancer

Resa Paksi Mandariska, Adzka Fahma Rodliya

Effects of Cucumber (*Cucumis sativus*) Extract Cream on Aquaporin-3 and Hyaluronic Acid Levels in Wistar Rats with Xerosis Cutis

Ajeng Destian Suparwi, Pasid Harlisa, Siti Thomas Zulaikhah

Effects of Aloe Vera Extract on Basal Cell Thickness and Lymphocyte Infiltration at the Gastroesophageal Junction in a Rat Model of Gastroesophageal Reflux Disease

Shellia Violita, Tena Djuartina, Vetinly Vetinly, Iskandar Rahardjo

[Home](#) / [Archives](#) / Vol. 13 No. 1 (2026)

Vol. 13 No. 1 (2026)



DOI: <https://doi.org/10.15850/amj.v13n1>

Published: 2026-02-28

Articles

Physical Activity Levels Based on Indonesian Population Characteristics: Evidence from the Indonesian Family Life Survey-5

Givana Felicia, Felicia Kurniawan, Yunisa Astiarani, Kevin Kristian, Bryany Titi Santi

1-9



PDF

Early Metabolic Alterations and Predictors of Obesity Among Young Adults in Indonesia: Focus on Lipid Abnormalities and Cardiometabolic Risk

Mirasari Putri, Eva Rianti Indrasari, Yafma Femin Helta Heriady, Rizky Suganda Prawiradilaga

10-19



PDF

Insomnia as a Risk Factor for Cardiovascular Disease Among Adults in Eastern Indonesia

Helitty Helitty, Firman Firman, Elmukhsinur Elmukhsinur, Wa Ode Aisya Zoharia, Noviati Noviati
20-27



Relationship between Serum 25-Hydroxyvitamin D Levels and Cardiovascular Disease Risk among Road Sanitations Workers

Magdalena Wartono, Tjam Diana Samara, Adrianus Kosasih, Dian Mediana

28-34



Correlation between Nicotine Dependence and Cardiorespiratory Fitness within the Wellness Matrix Framework among University Students

Muhammad Ikhsan Ghaifazzari Ghafar, Ronny Lesmana, Setiawan Setiawan

35-42



Effects of Moderate-Intensity Treadmill Training on Cardiac Mitochondrial Mitophagy and Dynamics in Young and Aged Wistar Rats

Julia Windi Gunadi, Difa Hidayat Ramadhan, Bernike Tirsa Safira, Andra Alifya Ramadhan, Roro Wahyudianingsih, Yenni Limyati, Hanna Goenawan, Ronny Lesmana

43-51



Reduced Left Ventricular Global Longitudinal Strain in Obese Adolescents

Rilda Dwi Febrianda, Tisnasari Hafsah, Sri Endah Rahayuningsih, Dzulfikar Djalil Lukamnul Hakim, Rahmat Budi Kuswiyanto, Putria Rayani Apandi, Dida Akhmad Gurnida

52-58



Evaluation of Urinary hsa-miR-21-5p Expression as a Non-Invasive Biomarker for Prostate Cancer

Resa Paksi Mandariska, Adzka Fahma Rodliya

59-64



Effect of Cucumber (*Cucumis sativus*) Extract Cream on Aquaporin-3 and Hyaluronic Acid Levels in Wistar Rats with Xerosis Cutis

Ajeng Destian Suparwi, Pasid Harlisa, Siti Thomas Zulaikhah

65-73



Effects of Aloe Vera Extract on Basal Cell Thickness and Lymphocyte Infiltration at the Gastroesophageal Junction in a Rat Model of Gastroesophageal Reflux Disease

Shella Violita, Tena Djuartina, Vetinly Vetinly, Iskandar Rahardjo Budianto

74-80



ALTHEA MEDICAL JOURNAL

Journal Resources

Editorial policies and information for authors

Online Submission

Submit your manuscript online

Focus and Scope

Author Guidelines

Editorial Board

Peer Reviewers

EDITORIAL POLICIES

Publication Ethics

Peer Review Process

Journal Policies

VISIBILITY AND FEES

Abstracting & Indexing

Scopus Citedness

Article Processing Charges

Editorial Board

Editor in Chief

Edhyana Sahiratmadja

Scholar

Scopus

ORCID

Department of Biomedical Sciences, Faculty of Medicine Universitas Padjadjaran, Indonesia

Managing Editor

Mas Rizky A. A. Syamsunarno

Scholar

Scopus

ORCID

Department of Biomedical Sciences, Faculty of Medicine Universitas Padjadjaran, Indonesia

Editorial Board Members

Gerard Pals

Scholar

Scopus

VU University Medical Center Amsterdam, Netherlands

Alexander Kwarteng

Scholar

Scopus

ORCID

Department of Biochemistry and Biotechnology, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana

Mohd Hanafi Bin Ahmad Damanhuri

Scholar

Department of Biochemistry, Faculty of Medicine, Universiti Kebangsaan Malaysia (UKM), Kuala Lumpur, Malaysia

Kittiphong Paiboonsukwong

Scholar

Scopus

ORCID

Thalassemia Research Center, Institute of Molecular Biosciences, Mahidol University, Thailand

Winda Ariyani

Scholar

Scopus

ORCID

Gunma University Graduate School of Medicine, Gunma, Japan

Zulkarnain Jaafar

Scholar

Scopus

ORCID

Universiti Malaya, Kuala Lumpur, Malaysia

R. Tedjo Sasmono

Scholar

Scopus

ORCID

National Agency for Research and Innovation, Jakarta, Indonesia

Jarir At Thobari

Scopus

ORCID

Faculty of Medicine, Public Health, and Nursing Universitas Gadjah Mada, Yogyakarta, Indonesia

Soegianto Ali

Scholar

Scopus

ORCID

School of Medicine and Health Sciences, Atma Jaya Catholic University of Indonesia, Indonesia

Trevino Aristarkus Pakasi

Scholar

Scopus

Faculty of Medicine Universitas Indonesia, Indonesia

Andani Eka Putra

Faculty of Medicine Universitas Andalas, Indonesia

Pudji Lestari

ORCID

Faculty of Medicine Universitas Airlangga Surabaya, Indonesia

Ronny Lesmana

Scholar

Scopus

ORCID

Division of Physiology, Department of Biomedical Sciences, Faculty of Medicine Universitas Padjadjaran, Indonesia

Yunia Sribudiani

Scholar

Scopus

ORCID

Study Center of Clinical Genetics, Faculty of Medicine Universitas Padjadjaran, Indonesia

Setiawan Setiawan

ORCID

Physiology Division, Department of Biomedical Sciences, Faculty of Medicine Universitas Padjadjaran, Indonesia

Hanna Goenawan

Scholar

Scopus

ORCID

Division of Physiology, Department of Biomedical Sciences, Faculty of Medicine Universitas Padjadjaran, Indonesia

Nur Atik

Scholar

Scopus

ORCID

Division of Anatomy and Cell Biology, Department of Biomedical Sciences, Faculty of Medicine Universitas Padjadjaran, Indonesia

Arifudin Achmad

Scholar

Scopus

ORCID

Department of Nuclear Medicine and Molecular Theranostics, Faculty of Medicine Universitas Padjadjaran/Dr. Hasan Sadikin General Hospital Bandung, Indonesia

Mohammad Ghozali

Scholar

Scopus

ORCID

Division of Biochemistry and Molecular Biology, Department of Biomedical Sciences, Faculty of Medicine Universitas Padjadjaran, Indonesia

Astrid Feinisa Khairani

Division of Anatomy and Cell Biology, Department of Biomedical Sciences, Faculty of Medicine Universitas Padjadjaran, Indonesia

Irma Ruslina Defi

Scholar

Scopus

ORCID

Department of Physical and Rehabilitation Medicine, Faculty of Medicine Universitas Padjadjaran/Dr. Hasan Sadikin General Hospital Bandung, Indonesia

Sri Endah Rahayuningsih

Scopus

ORCID

Department of Child Health, Faculty of Medicine Universitas Padjadjaran/Dr. Hasan Sadikin General Hospital Bandung, Indonesia

Eko Fuji Ariyanto

Scholar

Scopus

Division of Biochemistry and Molecular Biology, Department of Biomedical Sciences, Faculty of Medicine Universitas Padjadjaran, Indonesia

Julia Ramadhanti

Division of Pharmacology and Therapy, Department of Biomedical Sciences, Faculty of Medicine Universitas Padjadjaran, Indonesia

Copyeditors

Poppy Siti C. Djen Amar

Scientific Journal Management Unit, Faculty of Medicine Universitas Padjadjaran, Indonesia

Editorial Assistants

Ayu Aeniah

Scientific Journal Management Unit, Faculty of Medicine Universitas Padjadjaran, Indonesia

Layout Editor

Devi Fabiola

Scientific Journal Management Unit, Faculty of Medicine Universitas Padjadjaran, Indonesia

ALTHEA MEDICAL JOURNAL

Journal Resources

Editorial policies and information for authors

Online Submission

Submit your manuscript online

99+

Mail

1

Chat

Meet

Compose

Inbox6,080

Starred

Snoozed

Sent

Drafts2

Categories

More

Labels

Contacts

Emailed Contacts

althea

Active

3 of 45

Letter of Acceptance and Proofreading Approval

ExternalInbox x

Althea Medical Journal<chiefeditor.amj@gmail.com>

to me, Tjam, Adrianus, Dian

Mar 15, 2026, 12:01 PM

Dear Magdalena Wartono, Tjam Diana Samara, Adrianus Kosasih, Dian Mediana,

We are pleased to inform you that your manuscript entitled "Relationship between Serum 25-Hydroxyvitamin D Levels and Cardiovascular Disease Risk among Road Sanitations Workers" is now ready for publication in the Althea Medical Journal (AMJ Volume 13 No. 1/February Year 2026) and will be available on <https://journal.fk.unpad.ac.id/index.php/amj/issue/archive>.

However, because we have made some editorial changes to the manuscript, we need your assistance to check if there are any mistakes in our layout and please state your agreement and fill in the form attached. If you find any mistakes in the design, please do not hesitate to let us know. In addition, due to our updated editorial policy, please kindly add the following sections where applicable: **Acknowledgments (if any), Authors' Contributions, Conflict of Interest, Funding, and Generative AI Disclosure Statement** (Please read the guidelines ([Editorial Policies](#))).

Please find the attachment to view the draft of the manuscript and the proofreading form. When you have completed all your corrections/comments, please send a copy of your comment and send it along with your filled "Proofreading Approval by Author" to our email address at chiefeditor.amj@gmail.com.

Thank you very much for submitting your article to the Althea Medical Journal.

ReplyReply allForwardShare in chat

Compose

Inbox6,080

Starred

Snoozed

Sent

Drafts2

Categories

More

Labels

Contacts

Emailed Contacts

althea

7 of 45

[AMJ] Editor Decision

ExternalInbox x

Edhyana Sahiratmadja<amj.fkunpad@gmail.com> via noreply.unpad.ac.id

to me, Tjam, Adrianus, Dian

Jan 29, 2026, 8:53 PM

Dear Dr. Magdalena Wartono,

We have reached a decision regarding your submission to Althea Medical Journal, "Relationship between 25-Hydroxy Vitamin D and the Risk of Cardiovascular Disease in Sanitations Workers".

One of our reviewers has raised a concern about your statistical result:
- when p value is not significant, is it needed to show OR and CI95%?
Please revise accordingly and send your manuscript in a single file.
Therefore, our decision is: Revisions Required

We appreciate your contribution to the Althea Medical Journal and look forward to receiving your revised manuscript, due 31 January 2026.

Thank you for your cooperation and for choosing Althea Medical Journal for your research. Should you have any questions or require further assistance, please do not hesitate to contact us.

ReplyForwardShare in chat

Compose

Inbox6,080

Starred

Snoozed

Sent

Drafts2

Categories

More

Labels

Contacts

Emailed Contacts

althea

10 of 45

Edhyana Sahiratmadja

amj.fk.unpad@gmail.com via noreply.unpad.ac.id

to me, Tjam, Adrianus, Dian

Tue, Dec 30, 2025, 1:07 AM

☆

😊

↶

⋮

Dear Magdalena Wartono,

We have reached a decision regarding your submission to Althea Medical Journal, "Relationship between 25-Hydroxy Vitamin D and the Risk of Cardiovascular Disease in Sanitations Workers".

Our decision is: Revisions Required

Please download all files from the reviewers. Revisions can be made to one of the reviewers' articles.

When submitting a revised manuscript, please include a COVER LETTER outlining, point by point, the revisions made in response to the reviewers' comments and guidelines. Besides, please use the TRACK CHANGES feature in MS Word or highlight the text to indicate which sections have been revised.

Once revised, please upload the manuscript in MS Word format to our website (Login > Authors > Submissions > Review > Editor's Decision > Upload Author Version).

We appreciate your contribution to the Althea Medical Journal and look

↶ Reply

↶ Reply all

↷ Forward

😊

🗨 Share in chat

Compose

Inbox6,080

Starred

Snoozed

Sent

Drafts2

Categories

More

Labels

Contacts

Emailed Contacts

althea

11 of 45

[AMJ] Submission Confirmation Letter

External

Inbox x

Althea Medical Journal

chiefeditor.amj@gmail.com via noreply.unpad.ac.id

to me, Tjam, Adrianus, Dian

Thu, Nov 6, 2025, 3:36 PM

★

😊

↶

⋮

Dear Dr. Magdalena Wartono, Dr. Tjam Diana Samara, Dr. Adrianus Kosasih, Dr. Dian Mediana

Thank you very much for submitting your manuscript entitled:

Relationship between 25-Hydroxy Vitamin D and the Risk of Cardiovascular Disease in Sanitations Workers

to Althea Medical Journal. Your manuscript has been assigned the ID of 4570.

The manuscript has undergone an initial checking process, including a plagiarism check using Turnitin. The result shows a similarity index of 20%, which is an acceptable similarity.

We appreciate your submission to Althea Medical Journal. Your manuscript will now proceed to the peer review process.

↶ Reply

↶ Reply all

↷ Forward

😊

 SINTA Executive

Dashboard

Explore SINTA

Mutation History

List Verificator PT

My SINTA

MAGDALENA WARTONO
Sinta ID : 5990641

Search...

Sort By

Year

Page 1 of 6 | Total Records : 59

Correlation between COHb, HIF-1? levels, and smoking habits with physical fatigue in online motorcycle taxi drivers

Author : M Wartono, TD Samara, AD Lestari, A Kosasih, N Novendy
Indonesian Journal of Biomedicine and Clinical Sciences 58 (26), 1-9, 2026

publish at
2026

0 cited

Relationship between Serum 25-Hydroxyvitamin D Levels and Cardiovascular Disease Risk among Road Sanitations Workers

Author : M Wartono, TD Samara, A Kosasih, D Mediana
Althea Medical Journal 13 (1), 28-34, 2026

publish at
2026

0 cited

41 The Historical Evolution of Occupational Exposure to Asbestos, Silica, and Inorganic Dusts in Indonesia: Regulation, Industry Influence, and Shifting Recognition of ...

Author : N Halrunisa, A Dwi Lestari, H Nuriah, A Mohamad Ridwan, L T. Merijanti, ...
Journal of Occupational Medicine and Toxicology 21 (Suppl1), 41, 2026

publish at
2026

0 cited

HUBUNGAN POLA ASUH ORANG TUA DENGAN PERILAKU MELUKAI DIRI PADA DEWASA MUDA

Author : NA Fadlyah, M Wartono

publish at
2025

0 cited

PENYULUHAN PENTINGNYA VITAMIN D DALAM PENCEGAHAN PENYAKIT JANTUNG KEPADA PETUGAS

publish at
2025

0 cited

 SINTA Executive

 Dashboard

 Explore SINTA

 Mutation History

 List Verificator PT

 My SINTA

MAGDALENA WARTONO
Sinta ID : 5990641



DASHBOARD

Your Overview

SINTA SCORE OVERALL

443

SINTA SCORE 3YR

153

AFFILIATION SCORE OVERALL

443

AFFILIATION SCORE 3YR

153

SCOPUS H-INDEX

1

WOS H-INDEX

2

GOOGLE SCHOLAR H-INDEX

7