



ISSN (Online): 2589-8655  
(Print): 2589-8647

# Journal of Medical Case Reports and Reviews



**Volume 07  
Year 2024**

 <http://jmcrr.info/index.php/jmcrr>



Current Archives Indexing About

**Journal Issue**

ISSN (Online): 2589-8655  
ISSN (Print): 2589-8647

**Journal of Medical Case Reports and Reviews**

Volume 07  
Year 2024

Published by  
**Medical Editor & Educational Research Publisher**  
View Current Issue

### Most Popular Articles

- Validity and reliability of visual analogue Scale (vas) for pain measurement
- Return to Article DetailsThe Contents of Calcium, Chlorine, Iodine, Potassium, Magnesium, Manganese, and Sodium in Thyroid Malignant Nodules and Thyroid Tissue Adjacent to Nodules
- A case report Tongue injury after car accident in 1.5 years old boy :A case report
- COVID-19 Vaccine Acquired Haemophilia:A case study and literature review
- Incidentally Morcellated High grade LeiomyosarcomaTreated by Cytoreductive Surgery and simultaneous Hyperthermic Intraperitoneal Chemotherapy

### Keywords

extracted... life time risk... agent

### Published Volumes

- 2024
- 2023
- 2022
- 2021
- 2020
- 2019
- 2018

### EDITORIAL BOARD MEMBER

**Andrii Puzyrenko, M.D., Ph.D.**  
(312) 610-2845 | Email: apuzyrenko@mcw.edu  
Pathology and Laboratory Medicine, Medical College of Wisconsin, Milwaukee, WI, USA

**Ozgur Karcioğlu**  
Email: Okarcioğlu@gmail.com  
Istanbul Education and Research Hospital,  
Dept. of Emergency Medicine, Fatih, Istanbul.

**Anna Pantouvaki**  
Email: anna.pantouvaki@gmail.com  
Head of Physiotherapy Department of "Venizeleio"  
General Hospital, Heraklion-Crete Greek

**Xiangxia Liu MD, PhD**  
Email: liuxsha@mail.sysu.edu.cn  
Division of Plastic Surgery  
First Affiliated Hospital, Sun Yat-sen University  
58# Zhongshan Road II  
Guangzhou, China 510080

**Maryam Sadat Hosseini Zare**  
Email: maryamhoseinizare@gmail.com  
Paper review for Journal of Drug Research and Development  
in USA .9111 lakes at 610 DR. Houston, TX, USA

**Sorush Nik Namian**  
Email: niknamian@mail.co.uk  
ACP Chapter member in United States Army and Air Force  
Board Member of United Breast Cancer Foundation

**Dr. Musa Basheer Musa Mnasour**  
Email: musabasheer97@yahoo.com  
Consultant family medicine, CME/CPD  
Instructor. TA in MLTA, Brazil.

**Dr. Palani ELUMALAI**  
Email: palani.elumalai@tamu.edu  
Department of Chemistry, Science Program,  
Texas A&M University, Texas, USA

**Dr. Abbas Jedariforoughi MD**  
Email: abbas.jedariforoughi@gmail.com  
ORCID: 0000-0001-8154-3792  
ResearcherID: AAQ-3025-2021  
MD Medical Doctor, Copenhagen university, Copenhagen, Denmark

**Dr. Nikolaos A. Chrysanthakopoulos**  
Email: nikolaos\_c@hotmail.com  
ORCID 0000-0002-8295-2819  
Dental and Oral Surgeon (DDSc) Greek

**Filippo Manelli MD**  
Email: filippo.manelli@asst-valcamonica.it  
ORCID: 0000-0003-0797-0360  
Director of Emergency Unit at ASST Bergamo EST (Esine, Italy)

**Professor Dr. Seyed Saeid, Zamanieh Shahri, MD**  
Email: saeid.zamanieh@cnsu.edu  
Faculty Member in California Northstate University, CNSU, USA  
University Professor in Losrios Community College District, USA

**Dr. Sonia Sayyedalhosseini, MD**  
Email: sonia.sayyedalhosseini@cnsu.edu  
Faculty Member in California Northstate University, CNSU, USA  
University Professor in Losrios Community College District, USA

**Maria Sofia Cotelli**  
Email: cotellim@gmail.com  
ORCID: 0000-0002-7010-2809  
SCOPUS: 55198442600  
Neurological Unit, ASST Valcamonica, Esine, Brescia,, Italy

**Dr. Grigorios Kastanis**  
Email: kastanis@gmail.com  
Senior Consultant of department of Reconstruction  
Hand Surgery of General Hospital of Heraklion- Venizeleio,  
Crete, Greece

**Hazim Abdul Rahman Alhiti**  
Email: hazim4436@gmail.com  
ORCID 0000-0003-0000-8267  
Scopus ID: 191002-007776  
General Surgeon Specialist M.D  
Al-Ramadi Teaching Hospital, general surgeon specialist

### Meet Our Editorial Team

**Dr. Andrii Puzyrenko**  
*Editor-in-chief*  
Milwaukee, WI, USA  
apuzyrenko@mcw.edu

**Dr. Ozgur Karcioğlu**  
*Lead Editor*  
Fatih, Istanbul  
Okarcioğlu@gmail.com


**Dr. Anna Pantouvaki**  
*Senior Editor*  
Heraklion-Crete Greek  
anna.pantouvaki@gmail.com

**Dr. Xiangxia Liu**  
*Associate Editor*  
Guangzhou, China  
liuxsha@mail.sysu.edu.cn

Read More

### Citation

	All	Since 2019
Citations	5	5
h-index	1	1
i10-index	0	0



### Language

- English
- Nederlands

### Information

- For Readers
- For Authors
- For Librarians

[Current](#) [Archives](#) [Indexing](#) [About](#) ▾

## Journal Issue



Published by  
**Medical Editor & Educational Research  
Publisher**  
[View Current Issue](#)

## Most Popular Articles

Validity and reliability of visual analogue Scale (vas) for pain measurement

Return to Article Details  
The Contents of Calcium, Chlorine, Iodine, Potassium, Magnesium, Manganese, and Sodium in Thyroid Malignant Nodules and Thyroid Tissue Adjacent to Nodules

A case report Tongue injury after car accident in 1.5 years old boy :A case report

COVID-19 Vaccine Acquired Haemophilia:A case study and literature review

Incidentally Morcellated High grade Leiomyosarcoma Treated by Cytoreductive Surgery and simultaneous Hyperthermic Intraperitoneal Chemotherapy

## Keywords




## Published Volumes

2018-2024

- 2024 ▾
- 2023 ▾
- 2022 ▾
- 2021 ▾
- 2020 ▾
- 2019 ▾
- 2018 ▾

[Home](#) / [Archives](#) / Vol. 7 No. 02 (2024)

## Articles



### The Stigma of COVID-19 And Its Influence on The Management of COVID-19

Stigma of COVID-19

Nany Hairunisa , Husnun Amalia

Citations ?

Online First: February 8, 2024 | DOI : 10.52845/JMCRR/2024/7-2-1 | Google Scholar | Abstract : 164

**Cite this:** *Journal of Medical Case Reports and Reviews*, Vol. 7 No. 02 (2024), **February 8, 2024**

[PDF](#)



### Book Review ' School Service Guidelines' (Malay Version)

Uqbah Iqbal

Citations ?


Online First: February 23, 2024 | DOI : 10.52845/JMCRR/2024/7-2-2 | Google Scholar | Abstract : 37

**Cite this:** *Journal of Medical Case Reports and Reviews*, Vol. 7 No. 02 (2024), **February 23, 2024**

[PDF](#)




## Meet Our Editorial Team



**Dr. Andrii Puzyrenko**  
*Editor-in-chief*  
Milwaukee, WI, USA  
apuzzyrenko@mcw.edu



**Dr. Ozgur Karcioğlu**  
*Lead Editor*  
Fatih, Istanbul  
Okarcioğlu@gmail.com



**Dr. Anna Pantouvaki**  
*Senior Editor*  
Heraklion-Crete Greek  
anna.pantouvaki@gmail.com




**Dr. Xiangxia Liu**  
*Associate Editor*  
Guangzhou, China  
liuxsha@mail.sysu.edu.cn

[Read More](#)

## Citation

Updated weekly

	All	Since 2019
Citations	5	5
h-index	1	1
i10-index	0	0



## Language

- English
- Nederlands

## Information

- For Readers
- For Authors
- For Librarians



## The Stigma of COVID-19 and Its Influence on the Management of COVID-19

Nany Hairunisa<sup>†</sup> <sup>1\*</sup>, Husnun Amalia<sup>2</sup>, Isra Sabrina<sup>3</sup>, Nashita Amira Zaina<sup>4</sup>, Laila Musfirah<sup>5</sup>, Yasmine Mashabi<sup>6</sup>, Emad Yousif<sup>7</sup>

1. Department of Occupational Medicine, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia
2. Department of Ophthalmology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia.
3. Department of Cardiology, Siloam Hospital, Banten, Indonesia
4. Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia
5. The National Narcotics Board Pratama Clinic, Banda Aceh, Indonesia
6. Department of Clinical Pathology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia
7. Department of Chemistry, College of Science, Al-Nahrain University, Baghdad, Iraq

**Abstract:-** Background: Negative stigma related to COVID-19, which has a social impact, among others, can cause an individual to hide their illness to avoid discrimination, prevent people from seeking immediate health care when experiencing symptoms, prevent individuals from developing healthy behaviors, and contribute to more severe health problems, ongoing transmission, and difficulties in controlling the spread of the COVID-19 virus. Objectives: The purpose and aim of conducting this research are to determine the effect of the stigma of COVID-19 on health care for individuals infected with COVID-19. Materials and methods: This research is descriptive research with quantitative and qualitative methods as a complement. Respondents involved in the quantitative method were 291 people, and qualitative (in-depth interviews) were 21 key informants. The research was carried out in 4 cities in Indonesia from December 2022 to June 2023.

Results: Most were women (69.8%), aged 26-35 years (37.1%), Diploma/Undergraduate (66.7%), and worked as government employees (28.9%). This research founds that 59.1% of respondents have moderate knowledge about COVID-19, and 83.8% answered that they really need information related to COVID-19. Regarding trust in information sources, most respondents (83.8%) responded that they trust it. We also found 79.4% of the respondent has moderate behavior towards the pandemic, and 74.6% has average emotional well-being. Stigma was evaluated through questionnaires distributed in a guided manner with the result that 6.19% of the 291 respondents experienced stigma, while 93.8% had no stigma related to COVID-19. Multiple linear regression test found that the significance value for the influence of knowledge, the need for information, trust in information sources, behavior towards pandemics and emotional well-being simultaneously for stigma is  $0.000 < 0.05$ . The calculated F value is  $4.716 > F$  table 2.25, so it can be concluded that the above factors significantly influence simultaneous Stigma.

Conclusion: This study found that knowledge, need for information, trust in information sources, attitudes towards pandemics and emotional well-being simultaneously have a significant effect on Stigma.

**Keyword:-** COVID-19, Knowledge, Behavior, Stigma.

**Copyright :** © 2024 The Authors. Published by Publisher. This is an open access article under the CC BY-NC-ND license

(<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Supplementary information** The online version of this article (<https://doi.org/xx.xxx/xxx.xx>) contains supplementary material, which is available to autho-rized users.

**Corresponding Author:** Nany Hairunisa<sup>†</sup>, Department of Occupational Medicine, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

\*ORCID ID: <https://orcid.org/0000-0003-2414-9224>

## Introduction

The COVID-19 pandemic caused by infection with the SARS-CoV-2 virus was first discovered in Wuhan, China, at the end of 2019 and was later declared a pandemic by WHO in March 2020. Since it first appeared until March 2022, 440,807,756 cases have been found worldwide, and precisely as many as 5,639,702 cases in Indonesia. [1], [2] Globally, the COVID-19 pandemic is causing health problems that impact a person's physical condition and emotional well-being.[3] Individuals infected with SARS-CoV-2 may have psychiatric comorbidities as well as problems related to depression, anxiety, stress, as well as negative stigma, which refers to disapproval or negative attitudes towards people infected with COVID-19.[4] Stigma is closely related to mental health problems, especially depression, so this social stigma then becomes an important part influencing public discourse and social phenomena towards the COVID-19 pandemic.[5]

During the COVID-19 pandemic, stigma has become a public health challenge. Stigma related to COVID-19 refers to disapproving or negative self-attitudes stemming from individuals infected with, or who have close contact with, COVID-19.[6], [7] Recent findings indicate that some survivors of COVID-19 and their family members are rejected by neighbors, landlords, and even their employers.[8] Certain subpopulations, such as people suspected of having a high risk of contracting COVID-19, such as foreign travelers and individuals returning home from quarantine, also experience various forms of stigmatization, including social exclusion, stereotyping, and even humiliation.[9]

The social impacts of negative stigma related to COVID-19 include causing an individual to hide their illness to avoid discrimination, preventing people from seeking immediate health care when experiencing symptoms, preventing individuals from developing healthy behaviors, and contributing to more serious health problems. Severe, ongoing transmission and difficulties in controlling the spread of the COVID-19 virus.[9]

Social stigma in health is a negative association between a person or group of people with certain characteristics and certain diseases. In this case, labels, stereotypes, or discrimination against people infected with the COVID-19 virus exist.[10] Stigma can encourage someone to tend to hide their illness to avoid discrimination, prevent people from seeking immediate health care, and prevent them from adopting healthy behaviors. The magnitude of a discriminatory social stigma will hinder the discovery and treatment of COVID-19 sufferers.[9]

This then also has a negative impact on the treatment and care of patients. The current COVID-19 pandemic has triggered social stigma and discriminatory behavior towards individuals from certain ethnic backgrounds as well as individuals who are considered to have had contact with the virus.[9] A prejudiced and discriminatory public view of COVID-19 can lead to reduced access to treatment, inability to make decisions regarding treatment, and treatment being enforced. In addition, stigma also has a negative impact on coping policies and programs aimed at overcoming the COVID-19 pandemic.[11], [12]

The COVID-19 pandemic has caused serious threats to human physical health and life. This triggers various psychological problems; panic disorder, anxiety, and depression.[13]

## Materials and Methods

The research design was cross-sectional and conducted from September 2022 to July 2023. The research locations were in 4 cities/districts in 4 regions of Indonesia (Jakarta, Aceh, Bali, and Maluku). The selection of samples was consecutively non-random sampling. The questionnaire-based survey (quantitative) and qualitative methodologies were nested into it.

Data were collected from December 2022 to March 2023 in 4 cities/districts in Indonesia. Quantitative data comprised 291 respondents, with the sample population from each province calculated using the proportion of qualitative respondents. In addition, a total of 21 respondents were conducted in in-depth interviews (in-depth interviews), which will still be carried out with several respondents in Greater Jakarta.

The questionnaire was built with survey tools and COVID-19 guidelines from WHO (WHO, 2020).[14] The questionnaire includes six significant dimensions, knowledge, need for information, trust in information sources, behavior towards the pandemic, emotional well-being, and stigma. In addition to primary demographic details,

respondents were also asked to determine the distance to a health facility, positive Covid-19 status, prevention and treatment, and treatment-seeking behavior for COVID-19.

## Statistical Analyses

Univariate and multivariate analyzes were performed with IBM SPSS Statistics 25.0 (Statistical Package for Social Sciences, New York, USA) program. In multivariate analysis, multiple linear regression was used to determine the effect of several factors on stigma. The t-test was used to determine the partial effect of several factors on stigma, and the F test to determine the simultaneous effect of several factors on stigma. The coefficient of termination is also used to determine the percentage of influence that several factors simultaneously have on stigma.

## Results

Data collection was carried out from December 2022 to March 2023 in 4 cities/regencies in Indonesia for quantitative data consisting of 291 respondents. In addition, 21 respondents conducted in-depth interviews, which complemented this study.

**Table 1. The Demographics of Respondent**

Characteristic	N	%
Age		
18–25-year-old	104	35.7
26–35-year-old	108	37.1
36–45-year-old	50	17.2
>45-year-old	29	10
Gender		
Male	88	30.2
Female	203	69.8
Education		
High school	91	31.3
Diploma/Undergraduate	194	66.7
Postgraduate	6	2.1
Marital Status		
Married	153	52.6
Not married/Divorced	138	47.4
Occupation		
Government employee	84	28.9
Health care worker	66	22.7
Student	69	23.7
Private employee	49	16.8
Businessman	15	5.2
Others	8	2.7
Religion		
Islam	222	76.3
Christian/Catholic	18	6.2
Hindu	47	16.2
Buddhist	4	1.4
Residence		
Aceh	67	23.0
Bali	73	25.1
JABODETABEK	78	26.8
Maluku	73	25.1

Most of the respondents in this study were women, as much as 69.8% and aged 22-35 years, as much as 37.1%. The age of respondents ranged from 18–58 years, with a mean age of 30.47. At the same time, the median and mode were 29 and 21 years, respectively. The highest age group was 26-35 years (37.1%). Around two-thirds of the

participants were female. And 52.6% of individuals were married. Most respondents' educational level is Diploma/Undergraduate 66.7% and as much as 28.9% are government employees. While the religion that is followed by the majority is Islam, 76.3%.

### Univariate analysis

In our questionnaire, we assessed several factors related to COVID-19, such as knowledge, need for information, trust in information sources, behavior toward the pandemic, emotional well-being and the stigma of COVID-19 with the results we can see in Table 2.

A total of 29 questions in the questionnaire related to knowledge about COVID-19, ten questions related to the need for information about COVID-19 and seven questions about trust in sources of information were asked of the respondents. The majority of respondents have moderate knowledge about COVID-19, as much as 59.1%. When asked whether they felt they needed more detailed information about COVID-19, 83.8% of respondents answered that they needed it. Regarding trust in sources of information, respondents were asked how much they trust doctors, the mass media, hospitals, the Ministry of Health and the role of community leaders or religious leaders in properly handling Covid-19, and 83.8% of respondents answered they trust. This was reinforced by the results of our in-depth interviews with key informants in collecting qualitative data, almost all of whom responded that the role of doctors down to the community level, in this case, religious leaders, plays a significant role in efforts to reduce transmission and in efforts to minimize the stigma that exists in society against COVID-19 patients so that management can be carried out properly. The following is one of the key informant comments from Bali. *“They work hand in hand. They coordinate well in Bali. Suppose there is this case, for example. We have officers at the Puskesmas doing surveillance; we communicate directly with Klian. In Bali, Klian is the head of the village. They have officials in their respective villages. In their village, there is an officer handling covid; they immediately went down to survey the place.”* Said Mrs D, 54 years old from Bali.

**Table 2. Univariate analysis**

Variable	N	%
Knowledge about COVID-19		
Good	24	8.2
Moderate	172	59.1
Lack	95	32.6
The Need for Information on COVID-19		
Really need	244	83.8
Simply need	23	7.9
Did not need	24	8.2
Trust in information sources		
Trust	244	83.8
Did not trust	47	16.2
Behavior towards the Pandemic		
Good	42	14.4
Average	231	79.4
Poor	18	6.2
Emotional Well-being		
Good	69	23.7
Moderate	217	74.6
Poor	5	1.7
Stigma on COVID-19		
Yes	18	6.2
No	273	93.8

The questionnaire also asked 13 questions about behavior towards a pandemic and ten questions about emotional well-being. The majority of respondents have relatively good behavior, namely 79.4%. The behavior here includes asking families not to visit during a pandemic and buying personal protective equipment (gloves, masks) etc.; almost

all respondents have mediocre behavior, namely 79.4%. As for the emotional well-being aspect, 74.6% of respondents have average emotional well-being.

Regarding stigma, we asked 31 questions related to stigma, and the result was 93.8% of respondents did not experience stigma associated with COVID-19, but 6.2% did experience stigma.

### Multivariate analysis

The analysis was carried out using Multiple Linear Regression with the results as we can see in Table 3.

**Table 3. Multiple linear regression for stigma**

#### Coefficients<sup>a</sup>

Model	Unstandardized Coefficients		Standardized Coefficients	t	Sig.
	B	Std. Error	Beta		
Knowledge about COVID-19	0.062	0.070	0.059	0.889	0.375
The Need for Information on COVID-19	0.045	0.041	0.078	1.105	0.270
Trust in information sources	0.155	0.054	0.192	2.885	0.004
Behavior towards the Pandemic	-0.029	0.072	-0.026	-0.399	0.690
Emotional Well-being	0.063	0.071	0.054	0.879	0.380

a. Dependent Variable: Stigma

From the calculations that have been done, the t-table value is 1,968. Analysis using multiple linear regression shows that the effect of knowledge on stigma is  $0.375 > 0.05$ , and the t-value is  $0.889 < 1.968$ , so it can be concluded that there is no significant effect between knowledge and stigma. The effect of the Need for Information on COVID-19 on stigma is  $0.270 > 0.05$ , and the t count is  $1.105 < 1.968$ , so it can be concluded that there is no significant effect of the need for information on stigma. The effect of behavior on stigma is  $0.690 > 0.05$ , and the t value is  $-0.399 < 1.968$ , so it can be concluded that behavior towards the pandemic has no effect on stigma. Likewise, the effect of emotional well-being on stigma is  $0.380 > 0.005$ , and the t-value is  $0.879 < 1.968$ , so it can be concluded that emotional well-being does not affect stigma. Meanwhile, the effect of Trust in information sources is  $0.004 < 0.05$ , and the t-value is  $2.885 < 1.968$ , so it can be concluded that Trust in information sources significantly affects stigma.

**Table 4. ANOVA<sup>a</sup>**

Model	Sum of Squares	df	Mean Square	F	Sig.
Regression	4127.995	5	825.599	4.716	0.000 <sup>b</sup>
Residual	49897.448	285	175.079		
Total	54025.443	290			

a. Dependent Variable: Stigma

b. Predictors: (Constant), Emotional Well-being, Information Needs, Behavior, Knowledge, Trust in Information Sources

Calculations have been made for the value of the F table, and the result is 2.25. As seen in Table 4, the ANOVA test obtained the value of Sig.  $0.000 < 0.05$  and calculated F value of  $4.716 > F$  table 2.25 so that it can be concluded that there is an effect of variable knowledge about COVID-19; the need for information on COVID-19; trust in information sources; behavior towards the pandemic and emotional well-being simultaneously towards stigma.

**Table 5. The coefficient of termination**

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	0.276 <sup>a</sup>	0.076	0.060	13.232

a. Predictors: (Constant), Emotional Well-being, Information Needs, Behavior, Knowledge, Trust in Information Sources

In the results of the subsequent analysis, we can see in Table 5, which shows the R square value of 0.076, which states that the effect of the variable, emotional well-being, information needs, behavior, knowledge, and trust in information sources simultaneously on stigma is 7.6%.



## Discussion

The spread of the coronavirus and its global reach has become a critical concern and calls for action to prevent further spread of the virus collectively. Even though most people are worried about staying safe during this pandemic, minimizing the stigma associated with the coronavirus because it can exacerbate this quite tense situation is very important.[15]

The World Health Organization (WHO) emphasizes the effectiveness of interventions to reduce stigma, pointing to activities related to building trust in health care, providing timely and proven information, showing empathy to people affected by illness and creating an environment that facilitates an open discussion between people and health professionals. Promoting an understanding of the disease and taking adequate and practical precautions to ensure the safety of oneself and loved ones is crucial. How we communicate about COVID-19 is critical to helping us take effective action to fight the disease and avoid the fear and stigma it triggers. An environment must be created in which diseases and related problems can be discussed and dealt with in an open, honest and compelling manner. CIFRC, UNICEF, WHO. Social Stigma Associated with COVID-19.[16], [17]

In a multivariate analysis, it was found that knowledge about COVID-19 did not significantly affect the stigma of COVID-19. However, the univariate results found that most respondents from the four cities studied had moderate knowledge about COVID-19. This is in line with the results of a study conducted by Singh et al. on the Indian population, where it was found that most respondents had a moderate level of covid related knowledge.[18] In addition, the level of knowledge about COVID-19 shown in this study is higher than the Bangladeshis.[19]

In this study, it was found that 83.8% of respondents stated that they really needed information related to COVID-19. However, in the multivariate analysis, it was found that the need for information did not have a significant effect on stigma. In a study conducted by Bhanot et al., it was found stigma produces long-term emotional and social consequences that stigma produces beyond the infection itself. The main reason for the stigma attached to the disease is ignorance. When the COVID-19 pandemic emerged, information about the causative agent, the nature of the virus, the severity of the infection, and the mode of transmission and management was lacking because the virus was still new, so there was a lot of misinformation. Today, in the era of social media, many statements are spread that are not verified, and then they go "viral". The need for accurate information is urgently needed by the community so that stigma can be suppressed. All parties must be committed to providing valid and correct details on COVID-19.[20]

The role of community leaders or traditional leaders in certain areas significantly reduces the stigma in the community related to COVID-19, especially the stigma against health workers who treat COVID-19 patients or provide health services during a pandemic. This has more or less affected the incidence of COVID-19 in Indonesia. Information from sources that are unreliable, either through word of mouth, mass media and social media, still dominates among the public, causing distrust of the dangers of COVID-19 which then affects compliance with health protocols.[21] This is in line with the results of our research, the sources of information are very important. In a univariate analysis, it was found that 83.8% of respondents stated that they trusted sources of information such as doctors, health services, traditional leaders, and religious leaders up to the level of the Ministry of Health. In the multivariate analysis results, it was found that trust in information sources has a significant effect on stigma.

In this study, statistical testing showed behavior has no significant effect on stigma. This aligns with research conducted by Devantoro et al. in a cross-sectional epidemiological study involving 2,240 Indonesian population. Statistical tests showed that there was no significant relationship between behavior and stigma.[22] The pandemic and how it can affect people's behavior towards health workers are still of little concern to researchers. Stigma is a predictable problem, but of course, it can lead to more direct severe consequences for the performance of health workers.[23] When health workers experience stigma, which results in increased stress, this can result in patients not getting proper care.[24]

In our study, most of the respondents had fairly good emotional well-being. In the multivariate analysis, it was found that emotional well-being did not have a significant effect on stigma. This aligns with research conducted by Lades et al. that researched differences in emotional well-being and how people spend their time during the pandemic. They found that most time spent at home and outside the home was associated with increased positive affect and

decreased negative emotions. These findings highlight activities that may play a protective role concerning well-being during a pandemic and the importance of limiting exposure to media coverage related to COVID-19.[25]

We also conducted a multivariate analysis to assess the effect of knowledge about COVID-19, need for information, trust in information sources, behavior towards COVID-19 and emotional well-being simultaneously on stigma. Our research found a significant effect of all these variables on COVID-19. In a study conducted by Danborn et al., the same results were found that there was a significant effect between the knowledge, attitudes, practices (KAP), fear and stigma of the population towards COVID-19 from state to state.[26]

### **Limitation of the study**

In answering the questionnaire, the respondents answered themselves, so the results depended on the respondents' honesty and their ability to remember it might create a memory bias. The mixed research design (quantitative and qualitative) makes it less possible for us to conduct statistical validity and reliability analysis. Furthermore, the research was conducted in 4 cities with only 291 respondents, so it does not represent the diverse Indonesian population.

### **Conclusion**

This study found that knowledge, need for information, trust in information sources, attitudes towards pandemics and emotional well-being simultaneously have a significant effect on Stigma. The role of community leaders or traditional leaders in certain areas significantly reduces the stigma in the community related to COVID-19, especially the stigma against health workers who treat COVID-19 patients or provide health services during a pandemic. This has more or less affected the incidence of COVID-19 in Indonesia.

### **Ethical Declarations**

#### **Acknowledgements**

We thank all those who have helped carry out this research, especially the Abiansema 1 Public Health Center (Bali), the Puskesmas Soasio (Ternate, North Maluku), the Lembah Seulawah Public Health Center (Aceh) and the Greater Jakarta Health Facility.

#### **Ethics Approval and Consent to Participate**

Ethical clearance from the Faculty of Medicine Universitas Trisakti Research Ethics Committee No. 180/KER/FK/X/2022 and research permits from the Ministry of Home Affairs No.400.5/7800/Polpum were obtained. All respondents filled out informed consent and had the right to resign whenever the respondent wanted. The interviews' results were kept confidential and strictly protected by removing all personal identifiers from the form to protect the respondents' privacy.

#### **Consent for Publication**

Not applicable (no individual personal data included).

#### **Availability of Data and Material**

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

#### **Competing Interests**

The authors declare that there is no conflict of interest.

#### **Funding**

This research was funded by an internal grant from Universitas Trisakti

#### **Authors' Contributions**

All authors listed have made a substantial, direct, and intellectual contribution to the work, and approved it for publication.

## References

- [1] W. H. O. WHO, “WHO Coronavirus (COVID-19) Dashboard | WHO Coronavirus (COVID-19) Dashboard With Vaccination Data,” *World Health Organization*, 2022. <https://covid19.who.int> (accessed Jun. 29, 2023).
- [2] DKI JAKARTA Government, “COVID-19 Response Team,” *DKI Jakarta Government*, 2020. <https://corona.jakarta.go.id/en/data> (accessed Jun. 29, 2023).
- [3] I. Sabrina, N. Hairunisa, and E. Yousif, “Long covid: A review on cardiovascular disease (CVD) in post-acute sequelae of SARS-CoV-2 infection (PASC),” *Biointerface Research in Applied Chemistry*, vol. 12, no. 3. AMG Transcend Association, pp. 3989–3995, 2022. doi: 10.33263/BRIAC123.39893995.
- [4] W. Duan, H. Bu, and Z. Chen, “COVID-19-related stigma profiles and risk factors among people who are at high risk of contagion,” *Soc Sci Med*, vol. 266, pp. 1–10, Dec. 2020, doi: 10.1016/j.socscimed.2020.113425.
- [5] Y. Yuan *et al.*, “COVID-19-related stigma and its sociodemographic correlates: a comparative study,” *Global Health*, vol. 17, no. 1, Dec. 2021, doi: 10.1186/s12992-021-00705-4.
- [6] E. Alison Holman, R. R. Thompson, D. R. Garfin, and R. C. Silver, “The unfolding COVID-19 pandemic: A probability-based, nationally representative study of mental health in the United States,” *Sci Adv*, vol. 6, no. 42, Oct. 2020, doi: 10.1126/sciadv.abd5390.
- [7] C. for D. C. and P. CDC, “Stigma Reduction,” *Centers for Disease Control and Prevention*, 2020. <https://www.cdc.gov/stopoverdose/stigma/index.html> (accessed Jun. 29, 2023).
- [8] D. Adom, J. A. Mensah, and M. Osei, “The psychological distress and mental health disorders from COVID-19 stigmatization in Ghana,” *Social Sciences & Humanities Open*, vol. 4, no. 1, p. 100186, 2021, doi: 10.1016/j.ssaho.2021.100186.
- [9] W. H. O. WHO, “Situation Report-35 SITUATION IN NUMBERS total and new cases in last 24 hours,” Geneva, Feb. 2020. Accessed: Jun. 29, 2023. [Online]. Available: <https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200224-sitrep-35-covid-19.pdf>
- [10] IFRC, UNICEF, and W. H. O. WHO, “Panduan-untuk-mencegah-dan-mengatasi-stigma-sosial,” *World Health Organization*, Feb. 24, 2020. [https://www.who.int/docs/default-source/searo/indonesia/covid19/panduan-untuk-mencegah-dan-mengatasi-stigma-sosial.pdf?sfvrsn=4f8bc734\\_2](https://www.who.int/docs/default-source/searo/indonesia/covid19/panduan-untuk-mencegah-dan-mengatasi-stigma-sosial.pdf?sfvrsn=4f8bc734_2) (accessed Jun. 29, 2023).
- [11] A. Aprianti, D. Puspitaningtyas Laksana, and F. Dewi Puspita Anggraini, “STIGMA MASYARAKAT INDONESIA PADA PASIEN DAN TENAGA KESEHATAN COVID-19 BERDASARKAN TEORI HEALTH BELIEF MODEL,” *Jurnal Kesehatan Masyarakat Andalas*, vol. 15, no. 2, pp. 15–21, Jun. 2021, [Online]. Available: <http://jurnal.fkm.unand.ac.id/index.php/jkma/>
- [12] L. Dannatt *et al.*, “The Impact of Stigma on Treatment Services for People With Substance Use Disorders During the COVID-19 Pandemic—Perspectives of NECPAM Members,” *Front Psychiatry*, vol. 12, pp. 1–4, Mar. 2021, doi: 10.3389/fpsy.2021.634515.
- [13] J. Qiu, B. Shen, M. Zhao, Z. Wang, B. Xie, and Y. Xu, “A nationwide survey of psychological distress among Chinese people in the COVID-19 epidemic: Implications and policy recommendations,” *Gen Psychiatr*, vol. 33, no. 2, Mar. 2020, doi: 10.1136/gpsych-2020-100213.
- [14] World Health Organization (WHO), “SURVEY TOOL AND GUIDANCE Rapid, simple, flexible behavioural insights on COVID-19,” Copenhagen, 2020. [Online]. Available: <http://www.euro.who.int/pubrequest>
- [15] S. UNICEF, “COVID-19 & stigma: How to prevent and address social stigma in your community | UNICEF Sudan,” *UNICEF*, 2020. <https://www.unicef.org/sudan/covid-19-stigma-how-prevent-and-address-social-stigma-your->

