

VOL. 5

NO. 4 | APRIL 2024

JISS

Jurnal of Indonesian Social Science

E-ISSN: 2723-6692 | P-ISSN: 2723-6595

jiss.publikasiindonesia.id



[HOME](#) / Editorial Team

Editorial Team

Editor-in-Chief

Ikhsan Nendi

Politeknik Siber Cerdika Internasional

Journal Manager

Abdurokhim

Politeknik Siber Cerdika Internasional, Indonesia [[Scopus](#)] [[Scholar](#)]

Editors

Muhammad Iqbal

National Cheng Kung University, Taiwan [[Scopus](#)]

Al-Harath Abdulaziz Mohammed Ateik

I-Madinah International University, Malaysia [[Scopus](#)] [[Scholar](#)]

Md. Isahaque Ali

Universiti Sains Malaysia, Malaysia [[Scopus](#)] [[Scholar](#)]

Juliana

Universitas Pelita Harapan, Indonesia, [[Scopus](#)] [[Scholar](#)]

Mohd Norazmi Bin Nordin

Universiti Tun Hussein Onn, Malaysia, [[Scopus](#)] [[Scholar](#)]

Johanuddin Lahap

Universitas Teknologi Mara, Malaysia, [[Scopus](#)] [[Scholar](#)]

Leni Pebriantika

Universitas Baturaja, Indonesia, [[Scopus](#)] [[Scholar](#)]

Wresni Pujiyati

Universitas Wiralodra, Indonesia, [[Scopus](#)] [[Scholar](#)]

Zul Anwar

Institut Keguruan dan Ilmu Pendidikan Mataram, Indonesia, [[Scopus](#)] [[Scholar](#)]

Muhammad Ali Equatora

Politeknik Ilmu Pemasaran, Indonesia, [[Scopus](#)] [[Scholar](#)]

Ady Muzwardi

Universitas Maritim Raja Ali Haji, Indonesia, [[Scopus](#)] [[Scholar](#)]

Andiyan

Universitas Falatehan Bandung, Indonesia, [[Scopus](#)] [[Sholar](#)]

Fidya Arie Pratama

Institut Agama Islam Bunga Bangsa Cirebon, Indonesia, [[Scopus](#)] [[Scholar](#)]

Riesti Triyanti

Research Center for Marine and Fisheries Socio Economics, Jakarta, Indonesia, [[Scopus](#)] [[Scholar](#)]

Siti Mahmudah

UIN Raden Intan Lampung, Indonesia [[Scopus](#)] [[Scholar](#)]



PEOPLE

Editorial Team

Reviewers

ABOUT JOURNAL

Focus and Scope

Peer Review Process

Publication Ethics and Misconduct

Artificial Intelligence Policy

Article Processing Charges

Copyright & License Statement

Abstract & Indexing

Plagiarism Policy

Publication Frequency

Open Access Policy

Digital Preservation

Withdrawal of Manuscript

Correction & Retraction

Revenue Source

SUBMISSION

Author Guidelines

Reviewer Guidelines

TEMPLATE ENGLISH



**TEMPLATE OF
MANUSCRIPT**

TEMPLATE INDONESIA

TEMPLAT NASKAH

SUPPORTED & TOOLS



INFORMATION

[For Readers](#)

[For Authors](#)

[For Librarians](#)

FLAG COUNTER



statcounter



Jurnal Indonesia Sosial Sains

Alamat

Greenland Sendang Residence Blok H1, Sendang,
Kec. Sumber, Cirebon, Jawa Barat 45611

Principal Contact

Publikasi Indonesia

Phone: 085659237884


Email: jiss.jsi@gmail.com

This journal is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-sa/4.0/).

© 2026 **jiss**

[HOME](#) / [ARCHIVES](#) / Vol. 5 No. 04 (2024): Jurnal Indonesia Sosial Sains

Vol. 5 No. 04 (2024): Jurnal Indonesia Sosial Sains



DOI: <https://doi.org/10.59141/jiss.v5i04>

PUBLISHED: 2024-04-25

ARTICLES

Visual Design of Pop-Up Book Publication Binturong the Rare Cat Bear

Dermawan Syahputra, Universitas Potensi Utama, Indonesia

Muhammad Rusdi Tanjung, Universitas Potensi Utama, Indonesia



Dermawan Syahputra ,



Muhammad Rusdi Tanjung



DOI: [10.59141/jiss.v5i04.1071](https://doi.org/10.59141/jiss.v5i04.1071)

788-798

PDF

HTML

Criminal Law Reform of The Existence of Article 378 of The Criminal Code in Land Cases

Case Study of Judgment No. 1154/Pid.B/2021/PN. JKT.SE

Fransiska Khatrine, Universitas Kristen Indonesia, Indonesia



Fransiska Khatrine

DOI: [10.59141/jiss.v5i04.1082](https://doi.org/10.59141/jiss.v5i04.1082)

859-875

PDF

HTML

Legal Protection for Shareholders of Dissolved Insurance Companies

Oktaviandi Bangun Tri Anugrah Sitorus, Universitas Indonesia, Indonesia

Oktaviandi Bangun Tri Anugrah Sitorus

DOI: [10.59141/jiss.v5i04.1088](https://doi.org/10.59141/jiss.v5i04.1088)

876-881

PDF

HTML

Determinants in Used Car Purchase Results

El Ashar Hakim, PPM School of Management, Indonesia**Yusuf A. Zaki**, PPM School of Management, Indonesia**M. Fauzan Labib**, PPM School of Management, Indonesia**M. Fikri Arafa**, PPM School of Management, Indonesia**Andhika Hardiman P.**, PPM School of Management, Indonesia

El Ashar Hakim ,



Yusuf A. Zaki ,



M. Fauzan Labib ,



M. Fikri Arafa ,



Andhika Hardiman P,

DOI: [10.59141/jiss.v5i04.1093](https://doi.org/10.59141/jiss.v5i04.1093)

940-948

PDF

HTML

The Evaluation of the Implementation of the Independent Curriculum E-Report at Ngasinan Elementary School, Semarang Regency

Murtiari Setyaningtyas, Universitas Kristen Satya Wacana, Indonesia**Yari Dwikurnaningsih**, Universitas Kristen Satya Wacana, Indonesia**Bambang Ismanto**, Universitas Kristen Satya Wacana, Indonesia

Murtiari Setyaningtyas ,



Yari Dwikurnaningsih ,



Bambang Ismanto

DOI: [10.59141/jiss.v5i04.1055](https://doi.org/10.59141/jiss.v5i04.1055)

644-657

PDF

HTML

The Importance of Anchor Chain Maintenance at MT. Griya Flores

Andarum Septianto, Politeknik Pelayaran Sulawesi Utara, Indonesia**Iksan Saifudin**, Politeknik Pelayaran Sulawesi Utara, Indonesia

Andarum Septianto ,



Iksan Saifudin

DOI: [10.59141/jiss.v5i04.1065](https://doi.org/10.59141/jiss.v5i04.1065)

730-738

PDF

HTML

Comparative Strategy Analysis: Battle of Dien Bien Phu 1954 Vs General Attack 1 March 1949 Reflections on The History of Decolonization

Tri Yudha Ismanto, Universitas Pertahanan Republik Indonesia, Indonesia**Eko Lutvie**, Universitas Pertahanan Republik Indonesia, Indonesia**Irdham Ahmad**, Universitas Pertahanan Republik Indonesia, IndonesiaTri Yudha Ismanto ,  Eko Lutvie ,  Irdham AhmadDOI: [10.59141/jiss.v5i04.1080](https://doi.org/10.59141/jiss.v5i04.1080)

805-810

PDF

HTML

Risk Factors Affecting Internet Use in Adolescents: A Literature Review

Fika Maulid, Universitas Bina Bangsa Getsempena, Indonesia**Linur Steffi Harkensia**, Universitas Bumi Persada, IndonesiaFika Maulid ,  Linur Steffi HarkensiaDOI: [10.59141/jiss.v5i04.1085](https://doi.org/10.59141/jiss.v5i04.1085)

901-907

PDF

HTML

Dynamics of Advocacy and Community Empowerment Stage through the Resettlement Action Plan Program for the Kampung Bayam Community in the Construction of the Jakarta International Stadium Project by PT Jakarta Propertindo (Perseroda)

Hifdzi Mujtahid, Universitas Indonesia, Indonesia**Sari Viciawati Machdum**, Universitas Indonesia, IndonesiaHifdzi Mujtahid ,  Sari Viciawati MachdumDOI: [10.59141/jiss.v5i04.1091](https://doi.org/10.59141/jiss.v5i04.1091)

924-934

PDF

HTML

Exploring the Relationship Between Sod1, 2 And 3 Gene Polymorphisms With Post-Covid19 Symptoms

Jihan Samira Thabit, Universitas Trisakti, Indonesia**Sisca Sisca**, Universitas Trisakti, Indonesia**Monica Dwi Hartanti**, Universitas Trisakti, Indonesia

Noviani Prasetyaningsih, Universitas Trisakti, Indonesia

Alvionita Kogoya, Universitas Trisakti, Indonesia

Arleen Devita, Universitas Trisakti, Indonesia

Isa Bela, Universitas Trisakti, Indonesia

Ida Effendi, Universitas Trisakti, Indonesia

 Jihan Samira Thabit ,  Sisca Sisca ,  Monica Dwi Hartanti ,  Noviani Prasetyaningsih ,  Alvionita Kogoya , 
Arleen Devita ,  Isa Bela ,  Ida Effendi



DOI: [10.59141/jiss.v5i04.1097](https://doi.org/10.59141/jiss.v5i04.1097)

949-959

PDF

HTML

The Effect of Viral Marketing and E-Service Quality on Consumer Buying Interest in Disney+ Hotstar Services

Rama Raihan Syihab, Universitas Islam Indonesia, Indonesia

Erlita Ridanasti, Universitas Islam Indonesia, Indonesia

 Rama Raihan Syihab ,  Erlita Ridanasti



DOI: [10.59141/jiss.v5i04.1029](https://doi.org/10.59141/jiss.v5i04.1029)

842-858

PDF

HTML

Representation of Environmental Issues in Marjan's "Baruna the Guardian of the Ocean" Ad 2023 Episode 1 and 2 (Analysis Semiotika Charles Sanders Peirce)

Fatimah Nur Aliya Mahruf, Universitas Mulawarman, Indonesia

Kheyene Molekandella Boer, Universitas Mulawarman, Indonesia

Johantan Alfando Wikandana sucipta, Universitas Mulawarman, Indonesia

Jaka Farih Agustian, Universitas Mulawarman, Indonesia

 Fatimah Nur Aliya Mahruf ,  Kheyene Molekandella Boer ,  Johantan Alfando Wikandana sucipta ,  Jaka Farih Agustian



DOI: [10.59141/jiss.v5i04.1058](https://doi.org/10.59141/jiss.v5i04.1058)

667-686

PDF

HTML

A Comprehensive Analysis of Public Choices in Mass Transport and Assessing Development Challenges in The Transportation Sector

Ipoeng Martha Marsikun, Universitas Jenderal Soedirman, Indonesia

Paulus Israwan Setyoko, Universitas Jenderal Soedirman, Indonesia

Hikmah Nuraini, Universitas Jenderal Soedirman, Indonesia

Muslih Faozanudin, Universitas Jenderal Soedirman, Indonesia

Denok Kurniasih, Universitas Jenderal Soedirman, Indonesia

 Ipoeng Martha Marsikun ,  Paulus Israwan Setyoko ,  Hikmah Nuraini ,  Muslih Faozanudin ,  Denok Kurniasih



DOI: [10.59141/jiss.v5i04.1072](https://doi.org/10.59141/jiss.v5i04.1072)

705-718

[PDF](#)[HTML](#)

Women and The Journalist Profession (Phenomenological Study of The Professionalism of Women Journalists Within the Samarinda City Government)

Ellysa Fitri, Universitas Mulawarman, Indonesia

Jaka Farih Agustian, Universitas Mulawarman, Indonesia

Silviana Purwanti, Universitas Mulawarman, Indonesia

Hairunnisa Hairunnisa, Universitas Mulawarman, Indonesia

 Ellysa Fitri ,  Jaka Farih Agustian ,  Silviana Purwanti ,  Hairunnisa Hairunnisa



DOI: [10.59141/jiss.v5i04.1083](https://doi.org/10.59141/jiss.v5i04.1083)

[PDF](#)[HTML](#)

The Effect of Work Posture on Work Fatigue in Furniture Workers in the East Jakarta Furniture Industry Center

Iwan Jaya Azis, Universitas Mohammad Husni Thamrin, Indonesia

Nur Asniati Djaali, Universitas Mohammad Husni Thamrin, Indonesia

Brian Sri Prahastuti, Universitas Mohammad Husni Thamrin, Indonesia

 Iwan Jaya Azis ,  Nur Asniati Djaali ,  Brian Sri Prahastuti



DOI: [10.59141/jiss.v5i04.1089](https://doi.org/10.59141/jiss.v5i04.1089)

882-886

[PDF](#)[HTML](#)

The Influence of Motivation, Organizational Culture, and Transformational Leadership Style on Performance at Bhaktiasih General Hospital

Mohammad Abubakar, Universitas Esa Unggul, Indonesia

Sandra Dewi, Universitas Esa Unggul, Indonesia

Arrozi MF., Universitas Esa Unggul, Indonesia

 Mohammad Abubakar ,  Sandra Dewi ,  Arrozi MF.



DOI: [10.59141/jiss.v5i04.1095](https://doi.org/10.59141/jiss.v5i04.1095)

887-900

[PDF](#)[HTML](#)

The Effect of The Application of Good Corporate Governance (GCG) Principles on Job Satisfaction and Employee Performance

Rio Mubaraq Nazam, Universitas Esa Unggul, Indonesia

Rojuaniah Rojuaniah, Universitas Esa Unggul, Indonesia

 Rio Mubaraq Nazam ,  Rojuaniah Rojuaniah



DOI: [10.59141/jiss.v5i04.1056](https://doi.org/10.59141/jiss.v5i04.1056)

719-729

[PDF](#)[HTML](#)

Design and Build a Public Complaint Feature Via WhatsApp on the Adu.in Website with the Scrum Method

Case Study: West Java DPRD

Aprilita Firsty Hazdia, Universitas Telkom, Indonesia

Nur Ichsan Utama, Universitas Telkom, Indonesia

Sinung Suakanto, Universitas Telkom, Indonesia



Aprilita Firsty Hazdia ,



Nur Ichsan Utama ,



Sinung Suakanto



DOI: [10.59141/jiss.v5i04.1070](https://doi.org/10.59141/jiss.v5i04.1070)

760-772

[PDF](#)[HTML](#)

Silica Filter Membrane by Utilizing Palm Oil Empty Bunches Waste

Reco Rambang Putra Neksagis, PLN Sumatera Selatan, Indonesia

Hari Firmansah, PLN Sumatera Selatan, Indonesia

Ultra Marsedi, PLN Sumatera Selatan, Indonesia



Reco Rambang Putra Neksagis ,



Hari Firmansah ,



Ultra Marsedi



DOI: [10.59141/jiss.v5i04.1081](https://doi.org/10.59141/jiss.v5i04.1081)

773-781

[PDF](#)[HTML](#)

Governance of The Corporate Social Responsibility (CSR) Program of Karang Mukti Village in The Development of Embung Batu Beureum Tourism

Samsudin Samsudin, Program Pascasarjana Magister Ilmu Komunikasi STIAM, Indonesia

Muhammad Rifqi Hamidi, Program Pascasarjana Magister Ilmu Komunikasi STIAM, Indonesia

Wulan Furrie, Program Pascasarjana Magister Ilmu Komunikasi STIAM, Indonesia



Samsudin Samsudin ,



Muhammad Rifqi Hamidi ,



Wulan Furrie



DOI: [10.59141/jiss.v5i04.1086](https://doi.org/10.59141/jiss.v5i04.1086)

782-787

[PDF](#)[HTML](#)

The Application of Finger Painting Games to Children's Fine Motor Development in Kb Nurul Barokah, Wangkal Village, Gading District, Probolinggo District

Fatimatus Zahro, Universitas Islam Zainul Hasan Genggong, Indonesia

Endah Tri Wisudaningsih, Universitas Islam Zainul Hasan Genggong, Indonesia

Debby Adelita Febrianti Purnamasari, Universitas Islam Zainul Hasan Genggong, Indonesia

 Fatimatus Zahro ,  Endah Tri Wisudaningsih ,  Debby Adelita Febrianti Purnamasari



DOI: [10.59141/jiss.v5i04.1092](https://doi.org/10.59141/jiss.v5i04.1092)

811-824

PDF

HTML

The Importance of The Role of The Community in Amdal For The Goal of Environmentally Sound Sustainable Development

Case Study of Constitutional Court Decision No. 91/PUU-XVIII/2020 Concerning Judicial Review Law NO. 11 OF 2020 Concerning Job Creation

Sri Subekti, Kantor Notaris & PPAT Dr. Sri Subekti Kota Semarang, Indonesia

 Sri Subekti



DOI: [10.59141/jiss.v5i04.1098](https://doi.org/10.59141/jiss.v5i04.1098)

960-968

PDF

HTML

Increased Initial Reading Through Picture Word Cards

Latipah Syaripah, Universitas Muhammadiyah Sukabumi, Indonesia

 Latipah Syaripah



DOI: [10.59141/jiss.v5i04.1047](https://doi.org/10.59141/jiss.v5i04.1047)

739-745

PDF

HTML

The Effect of Motivation and Job Satisfaction on the Performance of MSME Credit Account Officer Employees at Bank BJB

Muhammad Adhitya Mulyadi, Sekolah Tinggi Ilmu Ekonomi (STIE) Ekuitas, Indonesia

Neneng Hayati, Sekolah Tinggi Ilmu Ekonomi (STIE) Ekuitas, Indonesia

 Muhammad Adhitya Mulyadi ,  Neneng Hayati



DOI: [10.59141/jiss.v5i04.1060](https://doi.org/10.59141/jiss.v5i04.1060)

687-704

PDF

HTML

The Role of Mass Media in The Emergence of Deviant Behavior in Society

Lianda Rachmadany Sinaga, Universitas Esa Unggul, Indonesia

Ayu Larasati, Universitas Esa Unggul, Indonesia

 Lianda Rachmadany Sinaga ,  Ayu Larasati

DOI: [10.59141/jiss.v5i04.1079](https://doi.org/10.59141/jiss.v5i04.1079)

799-804

PDF

HTML

Implementation of Vector-Based Melody Extraction for Plagiarism Detection Using Szymkiewicz-Simpson Coefficient

Nindyo Artha Dewantara Wardhana, Universitas Esa Unggul, Indonesia**Agung Mulyo Widodo**, Universitas Esa Unggul, Indonesia**Gerry Firmansyah**, Universitas Esa Unggul, Indonesia**Budi Tjahjono**, Universitas Esa Unggul, Indonesia

Nindyo Artha Dewantara Wardhana , Agung Mulyo Widodo , Gerry Firmansyah , Budi Tjahjono

DOI: [10.59141/jiss.v5i04.1084](https://doi.org/10.59141/jiss.v5i04.1084)

746-759

PDF

HTML

The Relationship of Hypertension with Death in Covid-19 Patients Treated at Tugurejo Hospital Semarang

Nanindya Ailsa Pramudhita, Universitas Muhammadiyah Semarang, Indonesia**Setyoko Setyoko**, Universitas Muhammadiyah Semarang, Indonesia**Susilo Budi Pratama**, Universitas Muhammadiyah Semarang, Indonesia

Nanindya Ailsa Pramudhita , Setyoko Setyoko , Susilo Budi Pratama

DOI: [10.59141/jiss.v5i04.1090](https://doi.org/10.59141/jiss.v5i04.1090)

935-939

PDF

HTML

Blue Economy from Perspective of Al-Qur'an

Ahmad Hery Sultoni, Universitas PTIQ, Indonesia**Hamdani Anwar**, Universitas Islam Negeri Syarif Hidayatullah, Indonesia**Nur Arfiyah Febriani**, Universitas Islam Negeri Syarif Hidayatullah, Indonesia

Ahmad Hery Sultoni , Hamdani Anwar , Nur Arfiyah Febriani

DOI: [10.59141/jiss.v5i04.1096](https://doi.org/10.59141/jiss.v5i04.1096)

908-923

PDF

HTML

The Influence of Using Documentary Films on Students' Writing Ability of Explanation Text

Murniasih Murniasih, Sekolah Tinggi Ilmu Ekonomi PPI, Indonesia

Melia Najihatul Ummah, Sekolah Tinggi Ilmu Ekonomi PPI, Indonesia



Murniasih Murniasih,  Melia Najihatul Ummah



DOI: [10.59141/jiss.v5i04.1057](https://doi.org/10.59141/jiss.v5i04.1057)

658-666

[HTML](#)[PDF](#)

sinta **S4**
Science and Technology Index

PEOPLE

Editorial Team

Reviewers

ABOUT JOURNAL

Focus and Scope

Peer Review Process

Publication Ethics and Misconduct

Artificial Intelligence Policy

Article Processing Charges

Copyright & License Statement

Abstract & Indexing

Plagiarism Policy

Publication Frequency

Open Access Policy

Digital Preservation

Withdrawal of Manuscript

Correction & Retraction

Revenue Source

SUBMISSION

Author Guidelines

Reviewer Guidelines

TEMPLATE ENGLISH



**TEMPLATE OF
MANUSCRIPT**

TEMPLATE INDONESIA

TEMPLAT NASKAH

SUPPORTED & TOOLS



INFORMATION

[For Readers](#)

[For Authors](#)

[For Librarians](#)

FLAG COUNTER



statcounter



Jurnal Indonesia Sosial Sains

Alamat

Greenland Sendang Residence Blok H1, Sendang,
Kec. Sumber, Cirebon, Jawa Barat 45611

Principal Contact

Publikasi Indonesia

Phone: 085659237884

Email: jiss.jsi@gmail.com

This journal is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-sa/4.0/) .

© 2026 **jiss**

Exploring the Relationship Between Sod1, 2 And 3 Gene Polymorphisms With Post-Covid19 Symptoms

Jihan Samira Thabit, Sisca, Monica Dwi Hartanti, Noviani Prasetyaningsih, Alvionita Kogoya, Arleen Devita, Isa Bella, Ida Effendi

Universitas Trisakti, Jakarta, Indonesia

Email: airajn08@gmail.com, sisca@trisakti.ac.id, mdhartanti@trisakti.ac.id,
novianip@trisakti.ac.id, alvionita030001900165@std.trisakti.ac.id,
arleen.devita@trisakti.ac.id, isabella@trisakti.ac.id, idaeffendi@trisakti.ac.id

Correspondence: airajn08@gmail.com*

KEYWORDS

Covid19; Long Covid19; SOD;
Genomic variation

ABSTRACT

The coronavirus disease 19 (COVID-19) has become a challenge for the media world. Even though they have been declared cured, some Covid-19 survivors still have health complaints. Abnormal symptoms, signs, or clinical parameters that persist two weeks or more after the onset of COVID-19 and do not return to their initial healthy state are potentially considered long-term effects of the disease. Although such changes are primarily reported in people with severe and critical illness, lasting effects also occur in individuals with mild infections that do not require hospitalization. This study aims to explore the role of antioxidants on the pathogenesis of Covid19 and its relationship with SOD1, 2 and 3 genomic variations. Getting alternative biomarkers for long covid19 detection. Research Method: The research sample is blood and questionnaires that will be taken from respondents affected by Covid19 a maximum of 6 months before data collection is held. Furthermore, DNA isolation, DNA amplification, cutting with restriction enzymes, and DNA band documentation with gel electrophoresis will be carried out. Symptoms related to systemic are the most common symptoms found in respondents with Long Covid19. A total of 8 respondents had systemic related symptoms, namely weakness, lethargy and sweating, while skin-related complaints were found in three respondents and one respondent had lung-related complaints. The average SOD activity value of respondents was 2.19 U/ml. In this study, more symptoms of long COVID-19 were found associated with systemic and cardiovascular symptoms. The complaints were not differentiated by the sex of the respondents. The output draft will be processed immediately after getting other results.

Attribution-ShareAlike 4.0 International (CC BY-SA 4.0)



1. Introduction

Coronavirus disease 19 (COVID-19) has become a challenge for the media world. Although related to the respiratory system, the pathogenesis of this disease is still not fully understood. COVID-19 was designated a pandemic by the World Health Organization (WHO) in March 2020 because it has spread worldwide and is considered a disease with a high transmission ability from person to person. Symptoms of Covid19 include early symptoms of low-grade fever and dry cough, followed by high fever, fatigue and difficulty breathing with dyspnea, and in more severe cases, hypoxia with oxygen saturation of less than 85%, multi-organ damage and even death (Xu et al., 2020). Even though they have been declared cured, some COVID-19 survivors still have health complaints. Abnormal symptoms, signs, or clinical parameters that persist two weeks or more after the onset of COVID-19 and do not return to their initial healthy condition are potentially considered long-term effects of the disease (Tenforde et al., 2020). Although such changes are primarily reported in people with severe and critical illnesses, lasting effects also occur in individuals with mild infections that do not require hospitalization (Townsend et al., 2021).

Once the virus enters the respiratory tract, the viral replication process begins, and the natural immune response begins to activate macrophages and dendritic cells via toll-like receptors and NODs against the production of inflammatory cytokines and reactive oxygen species (ROS) (Uehara et al., 2015). The spread of the virus in the blood will cause damage to erythrocytes by ROS and other inflammatory mechanisms (Li et al., 2021), leading to the formation of free hemoglobin and Fe. In addition, activated macrophages and neutrophils will produce respiratory bursts that will form superoxide radicals and H₂O₂, which lead to oxidative stress. Oxidative stress, along with free Fe, will convert soluble plasma fibrinogen into abnormal fibrin clumps in the form of dense matted deposits (DMD) that are resistant to degradation by enzymes. This will cause microthrombosis in the vascular system and pulmonary microcirculation. Cytokine storms arise through the upregulation of cytokine expression via NF- κ B. Furthermore, cytokine storms will induce oxidative stress through macrophages and neutrophil respiratory bursts and vice versa. This cycle will provoke serious tissue damage that is not related to viruses. In addition, mitochondria produce ROS, which increases iNOS expression through NF- κ B so that NO is formed which will induce mitochondrial dysfunction and cause cytopathic hypoxia. The virus will also inhibit Nrf2, which is responsible for increased antioxidant enzymes, causing oxidative stress. In conclusion, low haemoglobin levels and high lung exudate protein will cause pulmonary hypoxia, cytopathic hypoxia and endothelial damage and disseminated coagulation, which will cause multi-organ collapse (Cecchini & Cecchini, 2020).

Reactive Oxygen Species (ROS) can be characterized by several markers, including the enzyme Superoxide Dismutase (SOD). SOD functions to accelerate the reaction of the superoxide anion (O₂⁻) with itself to form hydrogen peroxide (H₂O₂) and oxygen (2O₂⁻ + 2H⁺ → H₂O₂ + O₂) (Wang et al., 2018). This reaction is important for maintaining redox equilibrium (Sies, 2015). However, excessive ROS production can exceed the capacity of the antioxidant defence system, causing oxidative stress and causing cellular damage by oxidized proteins, lipids, DNA/RNA and other macromolecules. There are three types of SOD; two of them are often found in mitochondria, namely SOD1 and SOD2. SOD 1 can be found in the intermembrane space, while SOD2 can be found in the mitochondrial matrix (Kim et al., 2017). SOD3 is often found extracellularly (Kim et al., 2017).

A relationship between the variation of the third SOD genotype and the risk of occurrence of several diseases has been found, for example, the CC rs4880 genotype is associated with increased hepatotoxicity after asparaginase therapy, the CC genotype rs2234694 increases the risk of T2DM, and the rs1041740 and rs17880487 variants in the SOD1 gene are associated with cardiovascular mortality and rs7655372 the SOD3 gene is associated with ischemic stroke (Alachkar et al., 2017; Ghattas & Abo-Elmatty, 2012; Otaki et al., 2016; Yang et al., 2021). However, the relationship between the third genotypic variant of SOD and post-Covid19 symptoms is still unknown.

The aim of this study was to explore the role of antioxidants in the pathogenesis of COVID-19. The other objective is to Analyze the SOD1, SOD2 and SOD3 mRNA expression levels of COVID-19 survivors. Analyzing SOD1, SOD2 and SOD3 levels in the plasma of Covid19 survivors, Analyzing the genotype variation of SOD1, SOD2 and SOD3 of Covid19 survivors, Analyzing the demographic picture and clinical symptoms of Covid19 survivors, Analyzing the relationship between expression levels and plasma SOD1, SOD2 and SOD3 levels with the duration of recovery and clinical symptoms of long covid19, Analyzing the relationship between SOD1, SOD2 and SOD3 genotype variations with the length of recovery and clinical symptoms of long covid19.

2. Materials and Methods

Research Design

The study will be designed as a case-control study that will test two groups of COVID-19 survivors with and without long-term COVID-19 symptoms.

Research Subjects

This study's research population is composed of COVID-19 survivors in Jakarta. Research subjects who meet the inclusion and exclusion criteria will be asked to join the study and sign an informed consent form.

Research Sample

Research samples are divided into 2 types, namely blood and tears. Blood is taken from a 5cc cubital vein and inserted into an EDTA vacutainer. Cheek mucosal smears will be performed using sterile cotton swabs that are daubed into the cheeks of research subjects and stored in buffers to prevent DNA and RNA damage. All blood samples and cheek mucosal smears were transferred to the laboratory in a styrofoam box filled with ice and stored in a refrigerator at - 80C for further analysis. Inclusion and exclusion criteria. The inclusion criteria are:

- Willing to sign informed consent.
- Willing to follow research procedures.

The exclusion criteria are:

- Individuals confirmed with COVID-19.

Sample Size

The five groups' gene expression differences were determined using G*Power 3.1.9.4 software. Based on the calculation of G*Power, which takes into account the effect size of 0.5, the total number of samples needed is 128 (Figure 1), with a total sample of 64 for each group.




Figure 1 Calculation of total sample size on G*Power

Research materials and methods

The materials needed in this study are:

- 1) RNA isolation kit
- 2) Kit synthesis cDNA
- 3) The DNA is the.
- 4) Enzyme restrictions

- 5) Conventional PCR mix
- 6) SYBR Green PCR Master Mix
- 7) Primer
- 8) Agarose Powder
- 9) Buffer TAE 1x
- 10) DNA Fluorosafe

Research Methods

1) RNA Isolation Research Methods

RNA will be extracted from blood and cervical smear samples in tubes containing RNA later. RNA extraction and DNase cleaning using RNAqueous-Micro® Kit (Cat# AM1931, Thermo Fisher Scientific, Waltham, MA, USA) in accordance with the manufacturer's protocol. The RNA concentration was determined using a nanodrop spectrophotometer based on a wavelength of 260. All samples with an absorption ratio of 2.0 to 260:280 will be used for further analysis.

2) Synthesis Cdna

Complement DNA (cDNA) will be synthesized from 200 ng of DNase-treated RNA. This method uses 250 ng/ul random hexamers (Sigma, Adelaide, SA, Australia) and 200 U Superscript Reverse Transcriptase III (Thermo Fisher Scientific, Waltham, MA, USA). To ensure the absence of genomic contamination, negative controls will be analyzed by adding water that DEPC has added in place of Superscript Reverse Transcriptase III.

3) Quantitative real-time PCR

For Quantitative real-time PCR (qPCR) analysis, primers will be designed based on publicly published RNA sequences using Primer3 plus (Rozen and Skaletsky, 2000) and primary Net software (PREMIERE Biosof). Quantitative real-time PCR analysis will be performed using the Rotor-Gene 6000 series 1.7 thermal cyclers (Qiagen GmbH) by duplication at 95C for 15 seconds, then 60C for 60 seconds for 40 cycles. CDNA amplification will be prepared in a 10 ml reaction containing 2 ml 1:20 cDNA, 5 ml Power SYBR Green PCR Master Mix (Applied Biosystems), 0.2 ml forward and reserve primers for target genes and 2.6 ml water mixed with DEPC. The Ct value is determined using Rotor-Gene 6000 software (Q series; Qiagen GmbH) at a threshold of 0.05 normalized with fluorescence units. Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) is used as a "housekeeping" gene due to its stable expression in human tissues.

4) DNA Isolation

According to the manufacturer's protocol, DNA will be isolated from venous blood and cheek mucosal smears using Quick-DNA™ Miniprep Plus Kit (Zymo Research, Irvine, CA, USA).

The concentration of DNA was determined using nanophotometers based on 260 wavelengths. All samples with a ratio of 1.8 to absorbance of 260:280 will be used for further analysis.

5) Target gene amplification

The target gene will be amplified using conventional PCR machines. According to the manufacturer's protocol, DNA will be mixed with MyTag™ HS Red Mix (Bioline-Meridian Bioscience Memphis, Tennessee, USA).

6) Incubation with restriction enzymes

The amplification results are incubated at a certain time and temperature with restriction enzymes (Thermo Fisher Scientific, Waltham, MA, USA) according to the manufacturer's protocol.

7) Gel Elektroforesis

Electrophoresis using 2% agarose gel to see the size of the incubation restriction fragment band

Analysis Methods

Data In this study, the data obtained on quantitative PCR will be entered into the following formula:

$$2^{-ave\Delta Ct}$$

Where:

Average ΔCt = mean of the results of reducing the cycle threshold of the GAPDH gene and the target gene studied.

The formula compares the expression of the target gene with the expression of the GAPDH (housekeeping gene) gene, which is constantly expressed in each type of body cell. The final result will be numerical data describing the target expression. The higher number describes the higher gene expression, while the lower number also describes the lower gene expression.

All statistical analysis will be performed using Microsoft Office Excel 2010 and GraphPad Prism Version 6.00 (GraphPad Software Inc.). Results will be presented in mean $\pm$ SEM if the data distribution is normal, and results will be presented in median if the data distribution is abnormal. All bivariate data will be analyzed using the t-test in the normal distribution; for normal distribution, data will be analyzed in non-Mann-Whitney statistical tests. The result will be considered significant at $p < 0.05$ with a 95% confidence interval.

Table 1 Research Achievement Indicators

Aspects	CHECKLIST
Featured Scale	International Scale
	National Scale ✓
	Local Scale
Research Topics/Themes	Top Down
	Semi Top Down
	Bottom Up ✓
Funding Scheme	Block Grant
	Competitive
RESEARCH IMPLEMENTER	Research Center
	Individual ✓
	Riset Group
Source Of Funds	Decentralized Fund

3. Result and Discussion

This study analyzes the demographic characteristics of patients and their relationship with symptoms of long-term COVID-19 that occur a maximum of 6 months after being declared recovered from COVID-19. Symptoms will be described based on the organ systems of the human body. There have been 55 respondents with the same portion for gender. Most respondents had never had Covid19 before but 10 respondents had had Covid19 before twice. Generally, respondents are self-isolating at home, and most respondents have no comorbidities for COVID-19 disease and no previous history of smoking.

Table 2 Characteristics of research respondents (n = 55)

No.	Information	Frequency (n)	Percentage (%)
1.	Gender		
	Man	27	49.09
	Woman	28	50.91
2.	History of Covid19		
	Never	24	43.64
	Once	21	38.18
	Twice	10	18.18
3.	Hospital history		
	Self-isolation	54	98.18
	Treated	1	1.82
4.	Comorbid		
	None	49	89.09
	Diabetes Mellitus	2	3.64
	Hypertensive	3	5.45
	Heart Disease	0	0
	Lung Disease	1	1.82
	Other	1	1.82
5.	Habit history		
	Smoke	16	29.09
	Drinking alcohol	0	0
	Drugs	0	0
	Other	39	70.91
6.	Symptoms of Long COVID-19		
	Yes	28	51
	No	27	49

Of the 55 respondents, there were 27 respondents without long-term covid symptoms and 28 respondents with long-term covid symptoms

Based on the data, it is stated that most patients who experience symptoms of long-term COVID-19 are women, while male patients have a tendency not to experience symptoms of long-term COVID-19. Based on the analysis, it was found that around 60% of patients suffering from long covid were women. This is in line with a publication from ISOS related to Long Covid which states that people who have weak immunity to infection, women, and people with severe disease are more likely to experience Long COVID.

As for the history of COVID-19, patients who experience symptoms of long-term COVID-19 are mostly patients who have previously experienced positive COVID-19 once. All patients who experience symptoms are patients undergoing self-isolation. Some patients do not have comorbidities. As many as 3 patients suffered from comorbidity, namely diabetes mellitus, as many as 1 person and 2 people with hypertension. Most patients do not have the habit of smoking, drinking alcohol or using drugs (Proskurnina et al., 2020; Uehara et al., 2015). Only 5 patients had a smoking habit. As for

patients who smoke, both regular cigarettes and vapes. Their smoking habits are intense sometimes, and they spend as much as 6 cigarettes to 1 pack of cigarettes in one day.

The virus that causes Covid19 is SARS-CoV-2 which has mutated several times. To date, there have been thousands of variants WHO has recorded in three categories; the dangerous one is the Variant of Concern (VOC). At the beginning of the COVID-19 pandemic, the alpha and beta variants had spread throughout the world, but it was indeed the delta variant that had a high morbidity and mortality rate compared to other variants.

The virus must be able to infect host cells and evade the immune system in order to survive. Therefore, the virus will continue to mutate. Similar to SARS-CoV2, this virus also continues to mutate, resulting in many variants, including the omicron variant currently detected in many countries worldwide.

Symptoms related to systemic are the most common symptoms found in respondents with Long Covid-19. A total of 8 respondents had systemic-related symptoms, namely weakness, lethargy and sweating, while skin-related complaints were found in three respondents, and one respondent had lung-related complaints.

Table 3 Long COVID-19 Complaints Related to Systemic Symptoms

No.	Information	Frequency (n)
1	Systemic Symptoms	
	Weak and lethargic	6
	Perspire	2
2	Lungs	
	Shortness of breath Skin	1
3	Hair loss	1
	Dry scaly skin	1
	Red stain	1
4	Cardiovascular	
	Low blood pressure	2
	Heart palpitations	1
	Chest pain	1

Low blood pressure is the most common cardiovascular-related symptom, followed by heart palpitations and chest pain.

From the results of SOD activity, it was found that the average SOD activity was 2.19 U/ml, with the highest SOD activity value being 3,164 U/ml while the lowest was 1,062 U/ml.

Table 4 The Value of SOD Activity in Respondents

No. Respond	SOD activity (U/ml)
1	2.570
2	2.088
3	2.317
4	2.852
5	3.164
6	2.852
7	2.570
8	2.341
9	1.786
10	2.570
11	2.088
12	2.317
13	1.882
14	2.088
15	2.317
16	2.317
17	2.707
18	2.852
19	2.440
20	2.852
21	1.529
22	2.199
23	1.451
24	2.199
25	2.088
26	2.570
27	1.982
28	1.882
29	1.882
30	2.852
31	2.088
32	1.529
33	2.088
34	1.882
35	1.378

36	2.570
37	1.786
38	1.882
39	1.062
40	1.696

Lopez-Leon et al. (2022) showed that there are five most common manifestations: fatigue, headache, attention disorder, hair loss, and shortness of breath. Other symptoms are related to breathing, cardiovascular, and neurological organs. Some studies report that complaints are more common in women, especially fatigue and panting (Townsend et al., 2021; Xiong et al., 2021).

4. Conclusion

In this study, more symptoms of long COVID-19 were found associated with systemic and cardiovascular symptoms. The complaints were not differentiated by the sex of the respondent.

5. References

- Alachkar, H., Fulton, N., Sanford, B., Malnassy, G., Mutonga, M., Larson, R. A., Bloomfield, C. D., Marcucci, G., Nakamura, Y., & Stock, W. (2017). Expression and polymorphism (rs4880) of mitochondrial superoxide dismutase (SOD2) and asparaginase induced hepatotoxicity in adult patients with acute lymphoblastic leukemia. *The Pharmacogenomics Journal*, 17(3), 274–279. <https://doi.org/10.1038/tpj.2016.7>
- Cecchini, R., & Cecchini, A. L. (2020). SARS-CoV-2 infection pathogenesis is related to oxidative stress as a response to aggression. *Medical Hypotheses*, 143, 110102. <https://doi.org/10.1016/j.mehy.2020.110102>
- Ghattas, M. H., & Abo-Elmatty, D. M. (2012). Association of Polymorphic Markers of the Catalase and Superoxide Dismutase Genes with Type 2 Diabetes Mellitus. *DNA and Cell Biology*, 31(11), 1598–1603. <https://doi.org/10.1089/dna.2012.1739>
- Kim, Y., Gupta Vallur, P., Phaëton, R., Mythreye, K., & Hempel, N. (2017). Insights into the Dichotomous Regulation of SOD2 in Cancer. *Antioxidants*, 6(4), 86. <https://doi.org/10.3390/antiox6040086>
- Li, S., Li, S., Disoma, C., Zheng, R., Zhou, M., Razzaq, A., Liu, P., Zhou, Y., Dong, Z., Du, A., Peng, J., Hu, L., Huang, J., Feng, P., Jiang, T., & Xia, Z. (2021). SARS-CoV-2: Mechanism of infection and emerging technologies for future prospects. *Reviews in Medical Virology*, 31(2). <https://doi.org/10.1002/rmv.2168>
- Otaki, Y., Watanabe, T., Nishiyama, S., Takahashi, H., Arimoto, T., Shishido, T., Miyamoto, T., Konta, T., Shibata, Y., Sato, H., Kawasaki, R., Daimon, M., Ueno, Y., Kato, T., Kayama, T., & Kubota, I. (2016). The Impact of Superoxide Dismutase-1 Genetic Variation on Cardiovascular and All-Cause Mortality in a Prospective Cohort Study: The Yamagata (Takahata) Study. *PLOS ONE*, 11(10), e0164732. <https://doi.org/10.1371/journal.pone.0164732>
- Proskurnina, E. V., Izmailov, D. Yu., Sozarukova, M. M., Zhuravleva, T. A., Leneva, I. A., & Poromov, A. A. (2020). Antioxidant Potential of Antiviral Drug Umifenovir. *Molecules*, 25(7), 1577. <https://doi.org/10.3390/molecules25071577>

- Sies, H. (2015). Oxidative stress: a concept in redox biology and medicine. *Redox Biology*, 4, 180–183. <https://doi.org/10.1016/j.redox.2015.01.002>
- Tenforde, M. W., Kim, S. S., Lindsell, C. J., Billig Rose, E., Shapiro, N. I., Files, D. C., Gibbs, K. W., Erickson, H. L., Steingrub, J. S., Smithline, H. A., Gong, M. N., Aboodi, M. S., Exline, M. C., Henning, D. J., Wilson, J. G., Khan, A., Qadir, N., Brown, S. M., Peltan, I. D., ... Wu, M. J. (2020). Symptom Duration and Risk Factors for Delayed Return to Usual Health Among Outpatients with COVID-19 in a Multistate Health Care Systems Network — United States, March–June 2020. *MMWR. Morbidity and Mortality Weekly Report*, 69(30), 993–998. <https://doi.org/10.15585/mmwr.mm6930e1>
- Townsend, L., Dowds, J., O'Brien, K., Sheill, G., Dyer, A. H., O'Kelly, B., Hynes, J. P., Mooney, A., Dunne, J., Ni Cheallaigh, C., O'Farrelly, C., Bourke, N. M., Conlon, N., Martin-Loeches, I., Bergin, C., Nadarajan, P., & Bannan, C. (2021). Persistent Poor Health after COVID-19 Is Not Associated with Respiratory Complications or Initial Disease Severity. *Annals of the American Thoracic Society*, 18(6), 997–1003. <https://doi.org/10.1513/AnnalsATS.202009-1175OC>
- Uehara, E. U., Shida, B. de S., & de Brito, C. A. (2015). Role of nitric oxide in immune responses against viruses: beyond microbicidal activity. *Inflammation Research*, 64(11), 845–852. <https://doi.org/10.1007/s00011-015-0857-2>
- Wang, Y., Branicky, R., Noë, A., & Hekimi, S. (2018). Superoxide dismutases: Dual roles in controlling ROS damage and regulating ROS signaling. *Journal of Cell Biology*, 217(6), 1915–1928. <https://doi.org/10.1083/jcb.201708007>
- Xiong, Q., Xu, M., Li, J., Liu, Y., Zhang, J., Xu, Y., & Dong, W. (2021). Clinical sequelae of COVID-19 survivors in Wuhan, China: a single-centre longitudinal study. *Clinical Microbiology and Infection*, 27(1), 89–95. <https://doi.org/10.1016/j.cmi.2020.09.023>
- Xu, Z., Shi, L., Wang, Y., Zhang, J., Huang, L., Zhang, C., Liu, S., Zhao, P., Liu, H., Zhu, L., Tai, Y., Bai, C., Gao, T., Song, J., Xia, P., Dong, J., Zhao, J., & Wang, F.-S. (2020). Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *The Lancet Respiratory Medicine*, 8(4), 420–422. [https://doi.org/10.1016/S2213-2600\(20\)30076-X](https://doi.org/10.1016/S2213-2600(20)30076-X)
- Yang, X., Yang, S., Xu, H., Liu, D., Zhang, Y., & Wang, G. (2021). Superoxide Dismutase Gene Polymorphism is Associated With Ischemic Stroke Risk in the China Dali Region Han Population. *The Neurologist*, 26(2), 27–31. <https://doi.org/10.1097/NRL.0000000000000301>

Exploring the Relationship Between Sod1, 2 And 3 Gene Polymorphisms With Post-Covid19 Symptoms

by Turnitin.library FK

Exploring the Relationship Between Sod1, 2 And 3 Gene Polymorphisms With Post-Covid19 Symptoms

Jihan Samira Thabit, Sisca, Monica Dwi Hartanti, Noviani Prasetyaningsih, Alvionita Kogoya, Arleen Devita, Isa Bella, Ida Effendi

Universitas Trisakti, Jakarta, Indonesia

Email: airajn08@gmail.com, sisca@trisakti.ac.id, mdhartanti@trisakti.ac.id,
novianip@trisakti.ac.id, alvionita030001900165@std.trisakti.ac.id,
arleen.devita@trisakti.ac.id, isabella@trisakti.ac.id, idaeffendi@trisakti.ac.id

Correspondence: airajn08@gmail.com*

KEYWORDS

Covid19; Long Covid19; SOD;
Genomic variation

ABSTRACT

The coronavirus disease 19 (COVID-19) has become a challenge for the media world. Even though they have been declared cured, some Covid-19 survivors still have health complaints. Abnormal symptoms, signs, or clinical parameters that persist two weeks or more after the onset of COVID-19 and do not return to their initial healthy state are potentially considered long-term effects of the disease. Although such changes are primarily reported in people with severe and critical illness, lasting effects also occur in individuals with mild infections that do not require hospitalization. This study aims to explore the role of antioxidants on the pathogenesis of Covid19 and its relationship with SOD1, 2 and 3 genomic variations. Getting alternative biomarkers for long covid19 detection. Research Method: The research sample is blood and questionnaires that will be taken from respondents affected by Covid19 a maximum of 6 months before data collection is held. Furthermore, DNA isolation, DNA amplification, cutting with restriction enzymes, and DNA band documentation with gel electrophoresis will be carried out. Symptoms related to systemic are the most common symptoms found in respondents with Long Covid19. A total of 8 respondents had systemic related symptoms, namely weakness, lethargy and sweating, while skin-related complaints were found in three respondents and one respondent had lung-related complaints. The average SOD activity value of respondents was 2.19 U/ml. In this study, more symptoms of long COVID-19 were found associated with systemic and cardiovascular symptoms. The complaints were not differentiated by the sex of the respondents. The output draft will be processed immediately after getting other results.

20

Attribution-ShareAlike 4.0 International (CC BY-SA 4.0)



1. Introduction

Coronavirus disease 19 (COVID-19) has become a challenge for the media world. Although related to the respiratory system, the pathogenesis of this disease is still not fully understood. COVID-19 was designated a pandemic by the World Health Organization (WHO) in March 2020 because it has spread worldwide and is considered a disease with a high transmission ability from person to person. Symptoms of Covid19 include early symptoms of low-grade fever and dry cough, followed by high fever, fatigue and difficulty breathing with dyspnea, and in more severe cases, hypoxia with oxygen saturation of less than 85%, multi-organ damage and even death (Xu et al., 2020). Even though they have been declared cured, some COVID-19 survivors still have health complaints. Abnormal symptoms, signs, or clinical parameters that persist two weeks or more after the onset of COVID-19 and do not return to their initial healthy condition are potentially considered long-term effects of the disease (Tenforde et al., 2020). Although such changes are primarily reported in people with severe and critical illnesses, lasting effects also occur in individuals with mild infections that do not require hospitalization (Townsend et al., 2021).

Once the virus enters the respiratory tract, the viral replication process begins, and the natural immune response begins to activate macrophages and dendritic cells via toll-like receptors and NODs against the production of inflammatory cytokines and reactive oxygen species (ROS) (Uehara et al., 2015). The spread of the virus in the blood will cause damage to erythrocytes by ROS and other inflammatory mechanisms (Li et al., 2021), leading to the formation of free hemoglobin and Fe. In addition, activated macrophages and neutrophils will produce respiratory bursts that will form superoxide radicals and H₂O₂, which lead to oxidative stress. Oxidative stress, along with free Fe, will convert soluble plasma fibrinogen into abnormal fibrin clumps in the form of dense matted deposits (DMD) that are resistant to degradation by enzymes. This will cause microthrombosis in the vascular system and pulmonary microcirculation. Cytokine storms arise through the upregulation of cytokine expression via NF- κ B. Furthermore, cytokine storms will induce oxidative stress through macrophages and neutrophil respiratory bursts and vice versa. This cycle will provoke serious tissue damage that is not related to viruses. In addition, mitochondria produce ROS, which increases iNOS expression through NF- κ B so that NO is formed which will induce mitochondrial dysfunction and cause cytopathic hypoxia. The virus will also inhibit Nrf2, which is responsible for increased antioxidant enzymes, causing oxidative stress. In conclusion, low haemoglobin levels and high lung exudate protein will cause pulmonary hypoxia, cytopathic hypoxia and endothelial damage and disseminated coagulation, which will cause multi-organ collapse (Cecchini & Cecchini, 2020).

Reactive Oxygen Species (ROS) can be characterized by several markers, including the enzyme Superoxide Dismutase (SOD). SOD functions to accelerate the reaction of the superoxide anion (O₂⁻) with itself to form hydrogen peroxide (H₂O₂) and oxygen ($O_2 + 2H \rightarrow H_2O_2 + O_2$) (Wang et al., 2018). This reaction is important for maintaining redox equilibrium (Sies, 2015). However, excessive ROS production can exceed the capacity of the antioxidant defence system, causing oxidative stress and causing cellular damage by oxidized proteins, lipids, DNA/RNA and other macromolecules. There are three types of SOD; two of them are often found in mitochondria, namely SOD1 and SOD2. SOD 1 can be found in the intermembrane space, while SOD2 can be found in the mitochondrial matrix (Kim et al., 2017). SOD3 is often found extracellularly (Kim et al., 2017).

A relationship between the variation of the third SOD genotype and the risk of occurrence of several diseases has been found, for example, the CC rs4880 genotype is associated with increased hepatotoxicity after asparaginase therapy, the CC genotype rs2234694 increases the risk of T2DM, and the rs1041740 and rs17880487 variants in the SOD1 gene are associated with cardiovascular mortality and rs7655372 the SOD3 gene is associated with ischemic stroke (Alachkar et al., 2017; Ghattas & Abo-Elmatty, 2012; Otaki et al., 2016; Yang et al., 2021). However, the relationship between the third genotypic variant of SOD and post-Covid19 symptoms is still unknown.

The aim of this study was to explore the role of antioxidants in the pathogenesis of COVID-19. The other objective is to Analyze the SOD1, SOD2 and SOD3 mRNA expression levels of COVID-19 survivors. Analyzing SOD1, SOD2 and SOD3 levels in the plasma of Covid19 survivors, Analyzing the genotype variation of SOD1, SOD2 and SOD3 of Covid19 survivors, Analyzing the demographic picture and clinical symptoms of Covid19 survivors, Analyzing the relationship between expression levels and plasma SOD1, SOD2 and SOD3 levels with the duration of recovery and clinical symptoms of long covid19, Analyzing the relationship between SOD1, SOD2 and SOD3 genotype variations with the length of recovery and clinical symptoms of long covid19.

2. Materials and Methods

Research Design

The study will be designed as a case-control study that will test two groups of COVID-19 survivors with and without long-term COVID-19 symptoms.

Research Subjects

This study's research population is composed of COVID-19 survivors in Jakarta. Research subjects who meet the inclusion and exclusion criteria will be asked to join the study and sign an informed consent form.

Research Sample

Research samples are divided into 2 types, namely blood and tears. Blood is taken from a 5cc cubital vein and inserted into an EDTA vacutainer. Cheek mucosal smears will be performed using sterile cotton swabs that are daubed into the cheeks of research subjects and stored in buffers to prevent DNA and RNA damage. All blood samples and cheek mucosal smears were transferred to the laboratory in a styrofoam box filled with ice and stored in a refrigerator at - 80C for further analysis. Inclusion and exclusion criteria. The inclusion criteria are:

- Willing to sign informed consent.
- Willing to follow research procedures.

The exclusion criteria are:

- Individuals confirmed with COVID-19.

Sample Size

The five groups' gene expression differences were determined using ⁸G*Power 3.1.9.4 software. Based on the calculation of G*Power, which takes into account the effect size of 0.5, the total number of samples needed is 128 (Figure 1), with a total sample of 64 for each group.




Figure 1 Calculation of total sample size on G*Power

Research materials and methods

The materials needed in this study are:

- 1) RNA isolation kit
- 2) Kit synthesis cDNA
- 3) The DNA is the.
- 4) Enzyme restrictions

- 5) Conventional PCR mix
- 6) SYBR Green PCR Master Mix
- 7) Primer
- 8) Agarose Powder
- 9) Buffer TAE 1x
- 10) DNA Fluorosafe

Research Methods

1) RNA Isolation Research Methods

RNA will be extracted from blood and cervical smear samples in tubes containing RNA later. RNA extraction and DNase cleaning using RNAqueous-Micro® Kit (Cat# AM1931, Thermo Fisher Scientific, Waltham, MA, USA) in accordance with the manufacturer's protocol. The RNA concentration was determined using a nanodrop spectrophotometer based on a wavelength of 260. All samples with an absorption ratio of 2.0 to 260:280 will be used for further analysis.

2) Synthesis Cdna

Complement DNA (cDNA) will be synthesized from 200 ng of DNase-treated RNA. This method uses 250 ng/ul random hexamers (Sigma, Adelaide, SA, Australia) and 200 U Superscript Reverse Transcriptase III (Thermo Fisher Scientific, Waltham, MA, USA). To ensure the absence of genomic contamination, negative controls will be analyzed by adding water that DEPC has added in place of Superscript Reverse Transcriptase III.

3) Quantitative real-time PCR

For Quantitative real-time PCR (qPCR) analysis, primers will be designed based on publicly published RNA sequences using Primer3 plus (Rozen and Skaletsky, 2000) and primary Net software (PREMIERE Biosof). Quantitative real-time PCR analysis will be performed using the Rotor-Gene 6000 series 1.7 thermal cyclers (Qiagen GmbH) by duplication at 95C for 15 seconds, then 60C for 60 seconds for 40 cycles. CDNA amplification will be prepared in a 10 ml reaction containing 2 ml 1:20 cDNA, 5 ml Power SYBR Green PCR Master Mix (Applied Biosystems), 0.2 ml forward and reserve primers for target genes and 2.6 ml water mixed with DEPC. The Ct value is determined using Rotor-Gene 6000 software (Q series; Qiagen GmbH) at a threshold of 0.05 normalized with fluorescence units. Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) is used as a "housekeeping" gene due to its stable expression in human tissues.

4) DNA Isolation

According to the manufacturer's protocol, DNA will be isolated from venous blood and cheek mucosal smears using Quick-DNA™ Miniprep Plus Kit (Zymo Research, Irvine, CA, USA). The concentration of DNA was determined using nanophotometers based on 260 wavelengths. All samples with a ratio of 1.8 to absorbance of 260:280 will be used for further analysis.

5) Target gene amplification

The target gene will be amplified using conventional PCR machines. According to the manufacturer's protocol, DNA will be mixed with MyTag™ HS Red Mix (Bioline-Meridian Bioscience Memphis, Tennessee, USA).

6) Incubation with restriction enzymes

The amplification results are incubated at a certain time and temperature with restriction enzymes (Thermo Fisher Scientific, Waltham, MA, USA) according to the manufacturer's protocol.

7) Gel Elektroforesis

Electrophoresis using 2% agarose gel to see the size of the incubation restriction fragment band

Analysis Methods

Data In this study, the data obtained on quantitative PCR will be entered into the following formula:

$$2^{-ave\Delta Ct}$$

Where:

Average ΔCt = mean of the results of reducing the cycle threshold of the GAPDH gene and the target gene studied.

The formula compares the expression of the target gene with the expression of the GAPDH (housekeeping gene) gene, which is constantly expressed in each type of body cell. The final result will be numerical data describing the target expression. The higher number describes the higher gene expression, while the lower number also describes the lower gene expression.

All statistical analysis will be performed using Microsoft Office Excel 2010 and GraphPad Prism Version 6.00 (GraphPad Software Inc.). Results will be presented in mean $\pm$ SEM if the data distribution is normal, and results will be presented in median if the data distribution is abnormal. All bivariate data will be analyzed using the t-test in the normal distribution; for normal distribution, data will be analyzed in non-Mann-Whitney statistical tests. The result will be considered significant at $p < 0.05$ with a 95% confidence interval.

Table 1 Research Achievement Indicators

Aspects	CHECKLIST
Featured Scale	International Scale
	National Scale ✓
	Local Scale
Research Topics/Themes	Top Down
	Semi Top Down
	Bottom Up ✓
Funding Scheme	Block Grant
	Competitive
RESEARCH IMPLEMENTER	Research Center
	Individual ✓
	Riset Group
Source Of Funds	Decentralized Fund

3. Result and Discussion

This study analyzes the demographic characteristics of patients and their relationship with symptoms of long-term COVID-19 that occur a maximum of 6 months after being declared recovered from COVID-19. Symptoms will be described based on the organ systems of the human body. There have been 55 respondents with the same portion for gender. Most respondents had never had Covid19 before but 10 respondents had had Covid19 before twice. Generally, respondents are self-isolating at home, and most respondents have no comorbidities for COVID-19 disease and no previous history of smoking.

Table 2 Characteristics of research respondents (n = 55)

No.	Information	Frequency (n)	Percentage (%)
1.	Gender		
	Man	27	49.09
	Woman	28	50.91
2.	History of Covid19		
	Never	24	43.64
	Once	21	38.18
	Twice	10	18.18
3.	Hospital history		
	Self-isolation	54	98.18
	Treated	1	1.82
4.	Comorbid		
	None	49	89.09
	Diabetes Mellitus	2	3.64
	Hypertensive	3	5.45
	Heart Disease	0	0
	Lung Disease	1	1.82
	Other	1	1.82
5.	Habit history		
	Smoke	16	29.09
	Drinking alcohol	0	0
	Drugs	0	0
	Other	39	70.91
6.	Symptoms of Long COVID-19		
	Yes	28	51
	No	27	49

Of the 55 respondents, there were 27 respondents without long-term covid symptoms and 28 respondents with long-term covid symptoms

Based on the data, it is stated that most patients who experience symptoms of long-term COVID-19 are women, while male patients have a tendency not to experience symptoms of long-term COVID-19. Based on the analysis, it was found that around 60% of patients suffering from long covid were women. This is in line with a publication from ISOS related to Long Covid which states that people who have weak immunity to infection, women, and people with severe disease are more likely to experience Long COVID.

As for the history of COVID-19, patients who experience symptoms of long-term COVID-19 are mostly patients who have previously experienced positive COVID-19 once. All patients who experience symptoms are patients undergoing self-isolation. Some patients do not have comorbidities. As many as 3 patients suffered from comorbidity, namely diabetes mellitus, as many as 1 person and 2 people with hypertension. Most patients do not have the habit of smoking, drinking alcohol or using drugs (Proskurnina et al., 2020; Uehara et al., 2015). Only 5 patients had a smoking habit. As for

patients who smoke, both regular cigarettes and vapes. Their smoking habits are intense sometimes, and they spend as much as 6 cigarettes to 1 pack of cigarettes in one day.

The virus that causes Covid19 is SARS-CoV-2 which has mutated several times. To date, there have been thousands of variants WHO has recorded in three categories; the dangerous one is the Variant of Concern (VOC). At the beginning of the COVID-19 pandemic, the alpha and beta variants had spread throughout the world, but it was indeed the delta variant that had a high morbidity and mortality rate compared to other variants.

The virus must be able to infect host cells and evade the immune system in order to survive. Therefore, the virus will continue to mutate. Similar to SARS-CoV2, this virus also continues to mutate, resulting in many variants, including the omicron variant currently detected in many countries worldwide.

Symptoms related to systemic are the most common symptoms found in respondents with Long Covid-19. A total of 8 respondents had systemic-related symptoms, namely weakness, lethargy and sweating, while skin-related complaints were found in three respondents, and one respondent had lung-related complaints.

Table 3 Long COVID-19 Complaints Related to Systemic Symptoms

No.	Information	Frequency (n)
1	Systemic Symptoms	
	Weak and lethargic	6
	Perspire	2
2	Lungs	
	Shortness of breath Skin	1
3	Hair loss	1
	Dry scaly skin	1
	Red stain	1
4	Cardiovascular	
	Low blood pressure	2
	Heart palpitations	1
	Chest pain	1

Low blood pressure is the most common cardiovascular-related symptom, followed by heart palpitations and chest pain.

From the results of SOD activity, it was found that the average SOD activity was 2.19 U/ml, with the highest SOD activity value being 3,164 U/ml while the lowest was 1,062 U/ml.

Table 4 The Value of SOD Activity in Respondents

No. Respond	SOD activity (U/ml)
1	2.570
2	2.088
3	2.317
4	2.852
5	3.164
6	2.852
7	2.570
8	2.341
9	1.786
10	2.570
11	2.088
12	2.317
13	1.882
14	2.088
15	2.317
16	2.317
17	2.707
18	2.852
19	2.440
20	2.852
21	1.529
22	2.199
23	1.451
24	2.199
25	2.088
26	2.570
27	1.982
28	1.882
29	1.882
30	2.852
31	2.088
32	1.529
33	2.088
34	1.882
35	1.378

36	2.570
37	1.786
38	1.882
39	1.062
40	1.696

16

Lopez-Leon et al. (2022) showed that there are five most common manifestations: fatigue, headache, attention disorder, hair loss, and shortness of breath. Other symptoms are related to breathing, cardiovascular, and neurological organs. Some studies report that complaints are more common in women, especially fatigue and panting (Townsend et al., 2021; Xiong et al., 2021).

4. Conclusion

13

In this study, more symptoms of long COVID-19 were found associated with systemic and cardiovascular symptoms. The complaints were not differentiated by the sex of the respondent.

5. References

- Alachkar, H., Fulton, N., Sanford, B., Malnassy, G., Mutonga, M., Larson, R. A., Bloomfield, C. D., Marcucci, G., Nakamura, Y., & Stock, W. (2017). Expression and polymorphism (rs4880) of mitochondrial superoxide dismutase (SOD2) and asparaginase induced hepatotoxicity in adult patients with acute lymphoblastic leukemia. *The Pharmacogenomics Journal*, 17(3), 274–279. <https://doi.org/10.1038/tpj.2016.7>
- Cecchini, R., & Cecchini, A. L. (2020). SARS-CoV-2 infection pathogenesis is related to oxidative stress as a response to aggression. *Medical Hypotheses*, 143, 110102. <https://doi.org/10.1016/j.mehy.2020.110102>
- Ghattas, M. H., & Abo-Elmatty, D. M. (2012). Association of Polymorphic Markers of the Catalase and Superoxide Dismutase Genes with Type 2 Diabetes Mellitus. *DNA and Cell Biology*, 31(11), 1598–1603. <https://doi.org/10.1089/dna.2012.1739>
- Kim, Y., Gupta Vallur, P., Phaëton, R., Mythreye, K., & Hempel, N. (2017). Insights into the Dichotomous Regulation of SOD2 in Cancer. *Antioxidants*, 6(4), 86. <https://doi.org/10.3390/antiox6040086>
- Li, S., Li, S., Disoma, C., Zheng, R., Zhou, M., Razzaq, A., Liu, P., Zhou, Y., Dong, Z., Du, A., Peng, J., Hu, L., Huang, J., Feng, P., Jiang, T., & Xia, Z. (2021). SARS-CoV-2: Mechanism of infection and emerging technologies for future prospects. *Reviews in Medical Virology*, 31(2). <https://doi.org/10.1002/rmv.2168>
- Otaki, Y., Watanabe, T., Nishiyama, S., Takahashi, H., Arimoto, T., Shishido, T., Miyamoto, T., Konta, T., Shibata, Y., Sato, H., Kawasaki, R., Daimon, M., Ueno, Y., Kato, T., Kayama, T., & Kubota, I. (2016). The Impact of Superoxide Dismutase-1 Genetic Variation on Cardiovascular and All-Cause Mortality in a Prospective Cohort Study: The Yamagata (Takahata) Study. *PLOS ONE*, 11(10), e0164732. <https://doi.org/10.1371/journal.pone.0164732>
- Proskurnina, E. V., Izmailov, D. Yu., Sozarukova, M. M., Zhuravleva, T. A., Leneva, I. A., & Poromov, A. A. (2020). Antioxidant Potential of Antiviral Drug Umifenovir. *Molecules*, 25(7), 1577. <https://doi.org/10.3390/molecules25071577>

- Sies, H. (2015). Oxidative stress: a concept in redox biology and medicine. *Redox Biology*, 4, 180–183. <https://doi.org/10.1016/j.redox.2015.01.002>
- Tenforde, M. W., Kim, S. S., Lindsell, C. J., Billig Rose, E., Shapiro, N. I., Files, D. C., Gibbs, K. W., Erickson, H. L., Steingrub, J. S., Smithline, H. A., Gong, M. N., Aboodi, M. S., Exline, M. C., Henning, D. J., Wilson, J. G., Khan, A., Qadir, N., Brown, S. M., Peltan, I. D., ... Wu, M. J. (2020). Symptom Duration and Risk Factors for Delayed Return to Usual Health Among Outpatients with COVID-19 in a Multistate Health Care Systems Network — United States, March–June 2020. *MMWR. Morbidity and Mortality Weekly Report*, 69(30), 993–998. <https://doi.org/10.15585/mmwr.mm6930e1>
- Townsend, L., Dowds, J., O'Brien, K., Sheill, G., Dyer, A. H., O'Kelly, B., Hynes, J. P., Mooney, A., Dunne, J., Ni Cheallaigh, C., O'Farrelly, C., Bourke, N. M., Conlon, N., Martin-Loeches, I., Bergin, C., Nadarajan, P., & Bannan, C. (2021). Persistent Poor Health after COVID-19 Is Not Associated with Respiratory Complications or Initial Disease Severity. *Annals of the American Thoracic Society*, 18(6), 997–1003. <https://doi.org/10.1513/AnnalsATS.202009-1175OC>
- Uehara, E. U., Shida, B. de S., & de Brito, C. A. (2015). Role of nitric oxide in immune responses against viruses: beyond microbicidal activity. *Inflammation Research*, 64(11), 845–852. <https://doi.org/10.1007/s00011-015-0857-2>
- Wang, Y., Branicky, R., Noë, A., & Hekimi, S. (2018). Superoxide dismutases: Dual roles in controlling ROS damage and regulating ROS signaling. *Journal of Cell Biology*, 217(6), 1915–1928. <https://doi.org/10.1083/jcb.201708007>
- Xiong, Q., Xu, M., Li, J., Liu, Y., Zhang, J., Xu, Y., & Dong, W. (2021). Clinical sequelae of COVID-19 survivors in Wuhan, China: a single-centre longitudinal study. *Clinical Microbiology and Infection*, 27(1), 89–95. <https://doi.org/10.1016/j.cmi.2020.09.023>
- Xu, Z., Shi, L., Wang, Y., Zhang, J., Huang, L., Zhang, C., Liu, S., Zhao, P., Liu, H., Zhu, L., Tai, Y., Bai, C., Gao, T., Song, J., Xia, P., Dong, J., Zhao, J., & Wang, F.-S. (2020). Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *The Lancet Respiratory Medicine*, 8(4), 420–422. [https://doi.org/10.1016/S2213-2600\(20\)30076-X](https://doi.org/10.1016/S2213-2600(20)30076-X)
- Yang, X., Yang, S., Xu, H., Liu, D., Zhang, Y., & Wang, G. (2021). Superoxide Dismutase Gene Polymorphism is Associated With Ischemic Stroke Risk in the China Dali Region Han Population. *The Neurologist*, 26(2), 27–31. <https://doi.org/10.1097/NRL.0000000000000301>

Exploring the Relationship Between Sod1, 2 And 3 Gene Polymorphisms With Post-Covid19 Symptoms

ORIGINALITY REPORT

16%

SIMILARITY INDEX

12%

INTERNET SOURCES

12%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1	files01.core.ac.uk Internet Source	2%
2	Kanisha Prita Addina, Matthew Hendra, Jeffrey Tedjasulaksana, Muhammad Fadlan Hidayat. "Profit Clues in Environmental, Social, and Governance Trails: Evaluation of Machine and Deep Learning Algorithms", 2025 8th International Seminar on Research of Information Technology and Intelligent Systems (ISRITI), 2025 Publication	2%
3	wwjmr.com Internet Source	2%
4	www.nature.com Internet Source	2%
5	Motoc Nicoleta Stefania, Claudia Toma, Cosmina Ioana Bondor, Ruta Victoriua Maria, Petrariu Florin, Man Milena Adina. "Long COVID and Lung Involvement: A One-Year Longitudinal, Real-Life Study", Medicina, 2025 Publication	1%
6	www.spandidos-publications.com Internet Source	1%
7	research-repository.griffith.edu.au Internet Source	1%
8	mdpi-res.com Internet Source	1%
9	www.karyailmiah.trisakti.ac.id Internet Source	1%
10	rupress.org Internet Source	1%
11	Patrick Manu, Clara Cheung, Akilu Yunusa-Kaltungo, Fidelis Emuze, Tarcisio Abreu Saurin, Bonaventura Hadikusumo. "Construction Safety, Health and Well-being in the COVID-19 Era", Routledge, 2023 Publication	<1%

12	wyndhamresidence.com Internet Source	<1 %
13	Krishna Mohan Surapaneni, Manmohan Singhal, Sofia Rani Saggu, Ashruti Bhatt, Priya Shunmathy, Ashish Joshi. "A Scoping Review on Long COVID-19: Physiological and Psychological Symptoms Post-Acute, Long-Post and Persistent Post COVID-19", Healthcare, 2022 Publication	<1 %
14	Monica D. Hartanti, Roseanne Rosario, Katja Hummitzsch, Nicole A. Bastian et al. "Could perturbed fetal development of the ovary contribute to the development of polycystic ovary syndrome in later life?", PLOS ONE, 2020 Publication	<1 %
15	core.ac.uk Internet Source	<1 %
16	nc.inverse.com Internet Source	<1 %
17	www.publish.csiro.au Internet Source	<1 %
18	Nara Aline Costa, Natália Baraldi Cunha, Ana Lucia Gut, Paula Schmidt Azevedo et al. "Erythrocyte SOD1 activity, but not SOD1 polymorphisms, is associated with ICU mortality in patients with septic shock", Free Radical Biology and Medicine, 2018 Publication	<1 %
19	erj.ersjournals.com Internet Source	<1 %
20	cnrs.hal.science Internet Source	<1 %

Exclude quotes

Off

Exclude matches

< 10 words

Exclude bibliography

On

Exploring the Relationship Between Sod1, 2 And 3 Gene Polymorphisms With Post-Covid19 Symptoms

GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/0

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11