




AccScience Publishing / EJMO



[Journal Flyer](#)

Eurasian Journal of Medicine and Oncology

[Gold Open Access](#)

[Charge APCs](#)

[Double-Blind Peer Review](#)

[Archived in Portfolio](#)

Electronic ISSN: 2587-196X

Print ISSN: 2587-2400

Date of Foundation: 2017-08-03

Indexing: ESCI, Scopus, EBSCO, CNKI, Google Scholar, ProQuest, Gale

Journal Rank: Impact Factor 2025: 1.3 - Q2(Medicine, General& Internal), Q4(Oncology); CiteScore 2025: 1.1 - Q3(Medicine(Internal Medicine), Q4(Oncology))

Editors-in-Chief: Bernd Kaina, Hilmi Kodaz, Mingzhu Yin [View Editorial Board](#)



The *Eurasian Journal of Medicine and Oncology (EJMO)* is an international peer-reviewed journal that publishes high-quality research across the full spectrum of oncology and general medicine. *EJMO* accepts submissions in English only and aims to disseminate impactful research and clinical insights to an international readership.

Eurasian Journal of Medicine and Oncology

Singapore | Universities and research institutions | Media Ranking

Country

Singapore



Subject Area and Category

Medicine

Internal Medicine
Oncology

Publisher

AccScience Publishing

SJR 2025

0.238

Q3

H-Index

15

Publication type
Journals

ISSN
2587196X, 25872400

Coverage
2019-2025

Editor https://accscience.com/journal/EJMO/about/editorial_board



Submit to EJMO

Join in Editorial Board

Journal Browser

Volume | Year

Issue

Search

> Forthcoming Issue

> Current Issue

Volume 10	Volume 9
Volume 8	Volume 7
Volume 6	Volume 5
Volume 4	Volume 3
Volume 2	Volume 1

[View All](#)

News and

Announcements

The EJMO 2025 impact factor has been released! It's 1.3, and the quartile is Q2!
7 June 2025

The XXII SIS World Congress in Breast Cancer and Breast Health
1 April 2025

Call for Papers: New Developments in Bladder Cancer Treatment and Management
6 February 2025

Call for Papers: Nursing Care in Cancer
9 February 2025

Call for Papers: Natural Products in Disease Prevention and Treatment
9 January 2025

Eurasian Journal of Medicine and Oncology (EJMO) is

Editorial Board

Filter editorial board members

Editors-in-Chief



Prof. Dr. Bernd Kaina   

Institute of Toxicology, University Medical Center of the Johannes Gutenberg University Mainz, Mainz, Germany

Interests: Oncology; Glioblastoma; DNA repair; Damage response; Anticancer drugs; Mutagenesis; Carcinogenesis

Profile: Professor of Toxicology, Ph.D. habil., Senior researcher, Director (em) of the Institute of Toxicology and group leader, University Mainz...

[Show More](#)



PhD. Hilmi Kodaz   

Department of Medical Oncology, Acibadem Eskisehir Hospital, Eskisehir, Turkey

Interests: Oncology; Prognostic factors of tumors; Pathological diagnosis of tumors

Profile: Member of the Board of Directors of the Turkish Society of Medical Oncology



Prof. Mingzhu Yin   

1. Clinical Research Center (CRC), Medical Pathology Center (MPC), Cancer Early Detection and Treatment Center (CEDTC) and Translational Medicine Research Center (TMRC), Chongqing University Three Gorges Hospital, Chongqing University, Chongqing, China
2. Chongqing Technical Innovation Center for Quality Evaluation and Identification of Authentic Medicinal Herbs, Chongqing, China
3. School of Medicine, Chongqing University, Chongqing, China
4. Department of Dermatology, Hunan Engineering Research Center of Skin Health and Disease, Hunan Key Laboratory of Skin Cancer and Psoriasis, Xiangya Hospital, Central South University, Changsha, China

Interests: Tumor immune regulation mechanism; Synergistic mechanism of tumor epigenetics and immunity

Profile: Chief Physician, Professor and Doctoral Supervisor of Three Gorges Hospital, Chongqing University
Director of the Medical Pathology Center, Clinical Research Center, Early Diagnosis and Treatment Center for Tumors and...

[Show More](#)

Executive Editor-in-Chief



Prof. Tian Li  

Tianjin Medical University, Tianjin, China

Interests: Quality check; Evidence-based medicine; Clinical study; Oncology

Profile: Member of Cochrane Collaboration; World's Top 2% Scientists (2023-2025), Elsevier & Stanford University; Wiley China
Highly Contribution Author 2023-2025; Chinese Top Ten Advances in Traditional Chinese Medicine 2023, MedSci.

Associate Editors



Hu Zhou   

Affiliated Cancer Hospital of Zhengzhou University, Zhengzhou, China

Interests: Hematology; Immunology

Profile: Chief Physician of the Department of Hematology, Henan Cancer Hospital
National Committee Member of the Hemostasis and Thrombosis Group of the Chinese Medical Association...

[Show More](#)



Back to top

Volume 10 Issue 1 2026

Order by Published Date ▼

Full Issue: [Download](#)

REVIEW ARTICLE

Advances in understanding the mechanism and treatment strategies of radiation myelopathy

Yuhang Yu, Yukai Tang, Shengyi Liu, Limin Liu

Eurasian Journal of Medicine and Oncology **2026**, 10(1), 1–14; <https://doi.org/10.36922/EJMO025100046>

[Abstract](#) [Download PDF\(598.25KB\)](#) [XML](#)

120 Download 4239 Views

REVIEW ARTICLE

Targeting PI3K/AKT/mTOR signaling: A promising therapeutic approach for colorectal cancer

Zhaoxia Qu, Xiaozhen Pen, Xingjun Lu, Shunli Luo

Eurasian Journal of Medicine and Oncology **2026**, 10(1), 15–31; <https://doi.org/10.36922/EJMO025190175>

[Abstract](#) [Download PDF\(897.06KB\)](#) [XML](#)

79 Download 6768 Views

REVIEW ARTICLE

Understanding the molecular basis of radioresistance in lung cancer: A comprehensive review

Fatima Zohra Attouahri, Boutaina Addoum, Leila Benbacer, Samira Mimount, ... Mohammed El Mzibri

Eurasian Journal of Medicine and Oncology **2026**, 10(1), 32–59; <https://doi.org/10.36922/EJMO025190168>

[Abstract](#) [Download PDF\(3.64MB\)](#) [XML](#)

96 Download 3472 Views



ORIGINAL RESEARCH ARTICLE

Risk factors and human papillomavirus genotypes: Investigating their interplay in c

Raditya Wratsangka, Monica D. Hartanti, Gatot Purwoto, Lily M. Surjadi, ... Alvionita Kogoya

Eurasian Journal of Medicine and Oncology **2026**, 10(1), 318–327; <https://doi.org/10.36922/EJMO025490509>

[Abstract](#) [Download PDF\(505.93KB\)](#) [XML](#)

SHORT COMMUNICATION

Effects of inhaled methyl methacrylate monomers from acrylic prostheses on mous

Dhona Afriza, Utmi Arma, Hana Azzahra, Alimuddin Tofrizal

Eurasian Journal of Medicine and Oncology **2026**, 10(1), 328–335; <https://doi.org/10.36922/EJMO025190174>

[Abstract](#) [Download PDF\(1.87MB\)](#) [XML](#)

SHORT COMMUNICATION

Impact of residential proximity to hospital on diagnostic inequality: A retrospective Central Poland

ORIGINAL RESEARCH ARTICLE

Risk factors and human papillomavirus genotypes: Investigating their interplay in cervical cancer

Raditya Wratsangka^{1†}, **Monica D. Hartanti^{2,3†}**, **Gatot Purwoto⁴**, **Lily M. Surjadi¹**, **Miranti P. Aulia⁵**, and **Alvionita Kogoya⁶**

¹Department of Obstetric and Gynecology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

²Department of Biology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

³Center for Biomedical Research, Research Organization for Health, National Research and Innovation Agency, Bogor, West Java, Indonesia

⁴Department of Obstetrics and Gynecology, Cipto Mangunkusumo National Hospital, Faculty of Medicine, Universitas Indonesia, Jakarta, Indonesia

⁵Independent Researcher, Jakarta, Indonesia

⁶Biomolecular Laboratory, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

†These authors contributed equally to this work.

*Corresponding author:

Monica D. Hartanti
(mdhartanti@trisakti.ac.id)

Citation: Wratsangka R, Hartanti MD, Purwoto G, Surjadi LM, Aulia MP, Kogoya A. Risk factors and human papillomavirus genotypes: Investigating their interplay in cervical cancer. *Eurasian J Med Oncol.* 2026;10(1):318-327.
doi: 10.36922/EJMO025490509

Received: December 6, 2025

Revised: January 14, 2026

Accepted: January 20, 2026

Published online: February 27, 2026

Copyright: © 2026 Author(s). This is an Open-Access article distributed under the terms of the Creative Commons Attribution License, permitting distribution, and reproduction in any medium, provided the original work is properly cited.

Publisher's Note: AccScience Publishing remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Abstract

Introduction: Cervical cancer continues to pose a substantial worldwide health challenge, primarily linked to human papillomavirus (HPV) infection. In regions like Indonesia, understanding local genotype distribution is critical for refining screening and vaccination efforts.

Objective: This research explores the complex relationship between various risk factors and the types of HPV detected in cervical cancer patients at a national referral center in Indonesia.

Methods: Ninety-one women diagnosed with cervical cancer were recruited for this cross-sectional study at Cipto Mangunkusumo National Hospital, Jakarta—the top-tier referral facility in Indonesia. This setting provides a diverse patient demographic representing various geographical regions across the archipelago. Cervical swabs were collected for HPV genotyping using dot blot hybridization. Various clinical risk factors, including age, age at marriage, age at first pregnancy, parity, smoking habits, hormonal contraceptive history, and the stage of cervical cancer diagnosis, were assessed. Statistical analysis was conducted using Fisher's exact test to accommodate the distribution of the data.

Results: High-risk (HR) HPV was detected in 62.6% of participants. HPV type 18 was the most prevalent (15.4%), followed by types 52 (14.3%) and 16 (12.1%). Statistically significant associations were observed between HR-HPV presence and both early age at marriage ($p = 0.004$) and the use of hormonal contraception ($p = 0.002$). No significant correlations were observed between some patient characteristics (age, parity, and smoking habits) and HPV genotypes ($p > 0.05$). A total of 35.2% of samples were HPV-negative, potentially reflecting technical challenges or viral integration in advanced-stage cases.

Conclusion: These findings suggest that early marriage and hormonal contraceptive use are significant clinical markers for HR-HPV in this Indonesian population. The dominance of HPV 18 and 52 highlights the need for locally relevant diagnostic tools and the potential transition to nonavalent vaccines. These data are essential for developing locally relevant diagnostic tools and optimizing cervical cancer prevention strategies tailored to the Indonesian demographic.

Keywords: Cervical cancer; HPV genotypes; Hormonal contraception; Age at marriage; Indonesia

1. Introduction

Cervical cancer stands as a significant global health issue, ranking as the fourth most prevalent malignancy among women worldwide.¹ This malignancy represents a profound burden on healthcare systems, particularly in developing nations where screening infrastructure is often fragmented. According to data from the World Health Organization (WHO) in 2022, approximately 660,000 new cases and 350,000 deaths from cervical cancer were reported, with approximately 94% occurring in low- and middle-income countries.² The disparity in mortality rates between high-income and low-income regions underscores the critical need for localized epidemiological data to inform public health strategies. In many developing regions, the lack of centralized registries and limited access to pathology services further complicates the accurate mapping of disease prevalence.

The global incidence of cervical cancer ranges from 10 to 75 cases per 100,000 women.³ Indonesia, as of 2022 data, recorded 36,000 cases of cervical cancer and 21,000 associated deaths.⁴ The high mortality-to-incidence ratio in Indonesia suggests that many cases are detected at advanced stages, limiting treatment efficacy. This clinical reality is exacerbated by the geographical complexity of the Indonesian archipelago, where the distribution of healthcare resources is often uneven, leaving rural populations with limited access to life-saving interventions. In 2021, the WHO recommended the DNA-HPV test as a primary screening method.⁵ This shift from cytology-based screening to molecular-based testing represents a paradigm shift in early detection, aiming to identify high-risk viral presence before cellular abnormalities manifest.

Cervical cancer is preventable, with human papillomavirus (HPV) infection being a crucial etiological factor. HPV is present in over 95% of cervical cancer lesions.^{6,7} The virus targets the basal layer of the stratified squamous epithelium, utilizing the host cell's machinery to replicate its double-stranded DNA genome. While there is no direct treatment for the virus, some topical

microbicides have shown intrinsic antiviral activity.⁸ Despite these developments, the primary defense against cervical malignancy remains a combination of prophylactic vaccination and secondary prevention through rigorous screening protocols. Although most HPV infections are asymptomatic and can be resolved on their own, persistent HPV infection poses a risk of developing into precancerous lesions, ultimately leading to cervical cancer.⁹ High-risk HPV types, notably HPV 16 and 18, are responsible for low- and high-grade squamous intraepithelial lesions (LSIL and HSIL) in the cervix. Timely identification and treatment of precancerous lesions can prevent progression.¹⁰ Vaccination, protecting individuals from HPV infection, has demonstrated success in reducing cervical cancer incidence in developed countries alongside regular screening for adult women. However, the efficacy of these vaccines is largely dependent on alignment between the vaccine's targeted genotypes and the circulating genotypes within a specific geographic population.

Low- and middle-income countries often face limitations in preventive facilities and access, leading to advanced-stage diagnoses, as observed in Indonesia, where approximately 66.4% of patients are diagnosed at stages IIB-IVB.¹⁰ In Indonesia, logistical barriers, socio-economic factors, and a lack of widespread awareness contribute to the high prevalence of late-stage presentations.^{11,12} Furthermore, cultural taboos regarding pelvic examinations and reproductive health can deter women from seeking early screening, even when services are available.^{12,13}

As a member of the Papillomaviridae family, HPV encompasses a diverse group of over 200 distinct genotypes. These viruses specifically target the cutaneous and mucosal squamous epithelium, leading to their primary classification by tissue type. Regarding oncogenesis, these strains are further divided into low-risk types, which typically result in benign growths, and high-risk types, which are the fundamental drivers of malignant transformations.¹⁴ The pathogenic mechanism of high-risk HPV centers on the activity of the E6 and E7 oncoproteins; these viral products

facilitate cancer progression by inhibiting the essential tumor-suppressing functions of the p53 and pRb proteins.

Globally, HPV represents a pervasive sexually transmitted infection that impacts all genders. Epidemiological data indicate that genital HPV prevalence varies significantly, ranging from 3.5% to 45% in male populations and 2% to 44% in female cohorts, with comparable rates of transmission observed between them.¹⁵ In men, the majority of infections remain subclinical and resolve without medical intervention. When symptoms do manifest in males, they usually present as benign genital warts. Similarly, while a high percentage of women contract cervical HPV during their lifetime, the vast majority of these cases do not escalate into invasive malignancy.¹⁶

A critical milestone in the transition to cervical cancer is the integration of the HPV genome into the host cell's DNA. This molecular event frequently disrupts the viral *E2* gene, causing an unchecked surge in *E6* and *E7* expression that drives rapid cellular transformation. Because symptoms often remain dormant for months or even years following the initial infection, clinical detection is frequently delayed until the appearance of physical warts or advanced lesions.¹⁷⁻²⁰ To combat this, the WHO has established the 90-70-90 targets for 2030: ensuring 90% vaccination coverage for girls by age 15, 70% screening participation for women at ages 35 and 45, and 90% access to treatment for those diagnosed with cervical disease.²¹ Indonesia has aligned with these goals by integrating HPV vaccination into its national immunization framework for adolescents. While this strategy is vital for long-term control, its success depends on logistical stability and public trust. Currently, the program does not yet extend to the adult female demographic.

To align with the government's goals of developing domestic diagnostic tools and expanding screening reach, it is crucial to develop cost-effective, sensitive, and specific diagnostic reagents tailored to the high-risk HPV genotype prevalence in the Indonesian population.²² Current diagnostic kits often use primers designed for Western populations, which may not achieve optimal sensitivity due to the unique genotype distribution in Southeast Asia. This preliminary study aims to correlate risk factors with HPV genotypes among women in Indonesia. Given the cohort size, this research serves as an exploratory analysis to identify patterns that warrant larger, multicenter investigations into the complex interplay between patients' reproductive histories and viral characteristics. By establishing a clearer picture of the local molecular landscape, we can better inform the selection of vaccines and the development of localized screening protocols that are both economically viable and clinically effective.

2. Materials and methods

2.1. Ethical clearance

This study was approved by the ethical clearance committee of the Faculty of Medicine, Universitas Indonesia, with ethical clearance number KET-1262/UN2.F1/ETIK/PPM.00.02/2022. Written informed consent was obtained from subjects before enrolment, ensuring that all participants were fully apprised of the study's objectives and the nature of the clinical specimens to be collected. The informed consent process was conducted in the local language to ensure a comprehensive understanding of the voluntary nature of participation and the confidentiality of genomic data.

2.2. Study design and population

This was a cross-sectional study, observing the relationship between risk factors and HPV genotyping among women in Indonesia. It was conducted in 2022 at Cipto Mangunkusumo National Hospital, Jakarta, with targeted participants being women aged 18–65 years old. The hospital serves as a national referral center, providing a diverse patient demographic for oncological research. Only women who had signed the informed consent were included in this study. Women who have had a prior HPV vaccination or a treatment related to cervical cancer were excluded from this study to prevent the confounding effects of medical intervention on the natural distribution of HPV genotypes. These exclusion criteria were strictly maintained to ensure that the observed viral prevalence reflects the natural history of the infection within the local community rather than the impact of clinical prophylaxis.

2.3. Data collection

Ninety-one participants were recruited consecutively according to inclusion and exclusion criteria. While the cohort size reflects the preliminary and exploratory nature of this research, rigorous data collection protocols were maintained. Questionnaires were distributed to participants, containing detailed questions related to marriage status, smoking, parity status, and hormonal contraceptive use. Information regarding age at marriage and first pregnancy was recorded to assess early exposure to potential risk factors. For HPV genotyping, cervical swabs were collected in ThinPrep containers and stored at 4 °C for analysis. Maintaining a strict cold chain was essential to prevent viral DNA degradation by endogenous nucleases in clinical samples.

2.4. HPV genotyping

DNA extraction was performed from cervical swabs using the QIAamp® DNA Blood Mini Kit (cat. 51104, Qiagen,

Netherlands) according to the manufacturer's protocol. Briefly, the sample in pellet resuspension was added to a mixture of Qiagen Protease (proteinase K) and Buffer AL. The addition of proteinase K is a critical step to ensure the complete lysis of cervical epithelial cells and the digestion of proteins that might otherwise sequester viral DNA. After a 10-min incubation at 56 °C, 96–100% ethanol was added. The mixture was then transferred to the QIAamp Mini Spin Column and centrifuged at $6000 \times g$ for 1 min. The use of high-fidelity spin columns ensures the removal of potential polymerase chain reaction (PCR) inhibitors, though the final DNA yield remains dependent on the initial cellularity of the swab. These inhibitors, often present in clinical mucus and blood, can lead to false-negative results if not thoroughly washed away during purification. Buffers AW2 and AE were added separately after serial centrifugation at different speeds and times.

The extracted DNA was then amplified, and HPV was genotyped using a reverse dot blot “flow-through” hybridization method with GeneFlow Human Papillomavirus (HPV) array test kit (cat. 92007; DiagCor Life Science Ltd., Hong Kong). The flow-through hybridization technology uses a patented active-pumping mechanism to direct target DNA to the probes, significantly increasing reaction speed and sensitivity compared to traditional passive hybridization. This hybridization-based approach enables simultaneous identification of multiple genotypes in a single sample, a critical step for detecting coinfections. This method can detect different high-risk HPV types, including 16, 18, 31, 33, 35, 39, 45, 51, 52, 53, 56, 58, 59, 66/68, 73, 81 and 82; and low-risk HPV types, including 6, 11, 26, 40, 42, 43, 44, 54, 55, 57, 61, 70, 71, 72, and 84. The broad coverage of this array is particularly advantageous for exploratory studies in regions where non-standard genotypes may be prevalent.

2.5. Statistical analysis

The data were collected and stored in Microsoft Excel 2025 (version 2511, Microsoft, USA). Categorical variables were summarized as absolute counts and percentages and analyzed using Fisher's exact test. This test is preferred to avoid the inaccuracies of the Chi-square approximation in sparse contingency tables, as it provides an exact probability value even when several cells have expected values of less than five. A p -value < 0.05 was considered significant. All statistical analyses were carried out using GraphPad version 9.0.0 (Dotmatics, United States). Due to the relatively small sample size ($n = 91$), this study lacked sufficient power to perform multivariable logistic regression to control confounding factors such as age and cancer stage. Consequently, the findings presented here are descriptive and exploratory. The high rate of HPV-negative results was analyzed separately to ensure no systematic bias

related to patient age or diagnostic stage was introduced.

3. Results

This study highlighted the characteristics of women with cervical cancer and correlated those characteristics with HPV genotyping. These findings illuminate the multifaceted nature of patient characteristics in the study, underscoring the endeavor to explore the correlations between risk factors and HPV genotyping in cervical cancer among Indonesian patients. While the sample size was 91, the findings remained primarily exploratory; however, the cohort revealed statistically significant associations that had previously been obscured. Notably, the data demonstrated that age at marriage and hormonal contraceptive use were key clinical markers for the presence of high-risk HPV genotypes.

3.1. Characteristics of participants

The study emphasized the significance of patient age and reproductive history, as detailed in [Table 1](#). The age distribution varied, with a minor proportion in the 25–34 age group (12.1%) and the majority (36.3%) aged 50 or older. Age at marriage emerged as a noteworthy factor. Roughly 29.7% of patients were married before age 20, while approximately 69.2% married after age 20. There was substantial variation in the age at which patients experienced their first pregnancy, with 72.5% experiencing it between the ages of 20 and 35. Differences in parity status were observed: 7.7% were nulliparous (never having given birth) and 91.2% were multiparous (having given birth more than once). Smoking habits exhibited significant variability among patients, with 6.6% identified as active smokers and 93.4% as non-smokers. Diagnostic data reflected the severity of cervical cancer in the patients. A minority (12.1%) received a pre-cancer diagnosis, while 25.3% were at stage 2, 7.7% at stage 3, and 1.1% at stage 4. This distribution confirms that the majority of participants had advanced-stage disease. The patterns of contraceptive use were diverse, with 30.8% of patients not using contraception and 69.2% using contraceptive methods.

3.2. HPV genotyping distribution

The study unveiled a diversity of HPV infections. Notably, 35.2% of the cohort ($n = 32$) had no detectable high-risk or low-risk HPV types—a finding that warrants further technical and clinical investigation. High-risk HPV types were detected in 62.6% of participants, while 2.2% had low-risk HPV types. Among the high-risk HPV types identified, HPV 18 was the most prevalent (15.4%), followed by HPV 52 (14.3%) and HPV 16 (12.1%). Other identified high-risk HPV types included HPV 45 (5.5%) and HPV 59 (3.3%). Coinfections were found in three women, possessing HPV 16 and 52; HPV 16 and 31; and

HPV 18, 66, and 68.

Table 1. Characteristics of participants

Characteristics	Count (n)	Percentage (%)
Age (years)		
25–34	11	12.1
35–44	25	27.5
45–50	22	24.2
>50	33	36.3
Age at marriage		
<20	27	29.7
>20	63	69.2
Age at first pregnancy		
<20	18	19.8
20–35	66	72.5
Parity number		
Nulliparous	7	7.7
Multiparous	83	91.2
Smoking		
Yes	6	6.6
No	85	93.4
Cancer stage		
Pre-Cancer	11	12.1
Stadium 1	9	9.9
Stadium 2	23	25.3
Stadium 3	7	7.7
Stadium 4	1	1.1
Hormonal Contraception		
Yes	28	30.8
No	63	69.2
HPV genotyping		
No HPV	32	35.2
High-risk HPV	57	62.6
Low-risk HPV	2	2.2
HPV type		
16	11	12.1
18	14	15.4
26	1	1.1
35	1	1.1
45	5	5.5
52	13	14.3
58	1	1.1
59	3	3.3

3.3. Relationship between risk factors and HPV genotyping

The interplay between clinical risk factors and the specific HPV genotypes was analyzed using Fisher's exact test, with results presented in Table 2. Significant correlations were observed between specific reproductive behaviors and the presence of high-risk HPV genotypes. Specifically, a significant association was observed regarding the age at marriage ($p = 0.004$), where patients who married before the age of 20 were significantly more likely to test positive for high-risk HPV (85.2%) compared to those who married at a later age (53.1%). Furthermore, the use of hormonal contraceptives demonstrated a strong statistical association with high-risk HPV detection ($p = 0.002$), with 85.7% of patients utilizing hormonal methods carrying high-risk HPV genotypes, whereas only 52.4% of non-users tested positive. No significant correlations were observed between HPV risk status and other variables, including age ($p = 0.514$), parity ($p = 1.000$), age at pregnancy ($p = 0.178$), and smoking habits ($p = 0.08$). While certain factors did not reach statistical significance, the findings regarding early marriage and hormonal contraception provide critical insights. These results indicate that in this specific Indonesian referral hospital population, HPV genotype distribution is significantly linked to early sexual exposure and hormonal history, suggesting that preventative strategies and screening in this demographic should prioritize individuals with these specific clinical markers.

4. Discussion

The findings of this study provide a nuanced look into the epidemiological landscape of HPV genotypes among cervical cancer patients in Indonesia. The primary observation of this research is the significant correlation between clinical markers—specifically age at marriage and hormonal contraceptive use—and the presence of high-risk HPV. While established clinical risk factors—such as parity and smoking—are well-documented drivers of cervical carcinogenesis, the lack of statistical significance in our cohort ($p > 0.05$) may be attributed to the overwhelming prevalence of these factors across the entire sample, thereby reducing their discriminatory power in a bivariate analysis. However, the identification of high-risk HPV in 62.6% of participants provides a more robust framework for identifying viral patterns that deviate from Western-centric models.

One of the most striking results is the high prevalence of HPV 18 (15.4%) and HPV 52 (14.3%), which, along with HPV 16, constitute the primary high-risk HPV types in this cohort. This distribution aligns with emerging data from Southeast Asia, suggesting that the regional

virome may differ from that of Western populations, in which HPV 16 and 18 are more dominant.^{20,23} The high frequency of HPV 18 and 52 reflects a specific molecular epidemiology in the Indonesian archipelago. Previous studies in neighboring regions have suggested that HPV 52 and 58 exhibit a higher fitness or transmission efficiency in certain Asian demographics, possibly due to host genetic factors or environmental selection pressures.^{24,25} The high frequency of HPV 52 has profound implications for local public health, particularly regarding the efficacy of current vaccines. Given that HPV 18 and 52 are major drivers in this population, transitioning to nonavalent vaccines or developing locally tailored diagnostic reagents is a clinical priority.²⁶

Table 2. Association of risk factors with HPV genotyping in women with cervical cancer

Risk Factors	No HPV/low-risk HPV	High-risk HPV	<i>p</i> value
Age			
<45	15	21	0.514
≥45	19	36	
Age at marriage			
<20	4	23	0.004*
>20	30	34	
Age at first pregnancy			
<20	4	14	0.178
20–35	30	43	
Parity number			
Nulliparous	3	4	1.000
Multiparous	31	53	
Smoking			
Yes	0	6	0.08*
No	34	51	
Hormonal contraception			
Yes	4	24	0.002*
No	30	33	

Notes: Fisher's test analysis was used to determine the relationships of each variable. **p* < 0.05.

A noteworthy finding in this study is the significant association between early age at marriage and high-risk HPV (*p* = 0.004). Specifically, 85.2% of women who married before age 20 tested positive for high-risk HPV genotypes. Similarly, the use of hormonal contraception (reported by 30.8% participants) was linked to the development of cervical cancer. However, this has become

a controversial topic due to inconsistent results. We did not find any significant association between the use of hormonal contraception and specific HPV genotypes (*p* > 0.05). Long-term use of oral contraceptives has been hypothesized to increase cervical cancer risk by making the cervix more susceptible to HPV DNA integration through hormonal modulation of the viral upstream regulatory region.^{27,28} Specifically, estrogen and progesterone may influence the expression of the viral *E6* and *E7* genes through hormone-responsive elements in the viral long control region.^{29,30} Our findings suggest that in the Indonesian context, the choice of contraception does not appear to influence the type of high-risk HPV that leads to malignancy. This reinforces the conclusion that once a high-risk HPV infection is established, the progression to cancer is likely driven by the oncogenic potential of the virus itself rather than the specific method of birth control used by the patient.

A significant point of discussion is the high proportion of HPV-negative results (35.2%) observed in this study. While cervical cancer is almost universally linked to HPV, literature does report HPV-negative cases, though usually at a lower rate than seen here. Several factors may explain this discrepancy. First, technical limitations in preserving DNA in a tropical climate could lead to sample degradation.³¹ Second, the sensitivity of the reverse dot blot hybridization method, while high, may have a detection threshold that misses low-copy-number viral DNA.³² Furthermore, the molecular architecture of the virus changes as the disease progresses. In advanced-stage cancer (which characterized 8.8% of our participants), the HPV genome often integrates into the host cell DNA, sometimes resulting in the loss of the *L1* gene—the target most PCR-based assays use for detection.³³ This false-negative phenomenon in advanced malignancies is a documented challenge in molecular oncology.^{34,35} Future studies should use multiple primer sets targeting different viral regions (e.g., *E6/E7*) to improve detection rates.

Regarding other reproductive factors, our study found that 91.2% of participants were multiparous. High parity is thought to increase cervical cancer risk through sustained hormonal changes and physical trauma to the transformation zone during childbirth.^{36–38} The repeated expansion and remodeling of the cervical transformation zone during multiple pregnancies may create a microenvironment that is more susceptible to persistent viral infection and subsequent cellular transformation. However, our analysis did not demonstrate an association between parity and specific HPV genotype (*p* = 1.000). This suggests that while parity helps the cancer progress, it may not be a selective pressure on HPV genotype. Similarly,

early age at first pregnancy (often proxies for early sexual debut) was common in our cohort but did not significantly correlate with HPV genotype. These findings reinforce the idea that once a high-risk HPV infection is established, the viral genotype's oncogenic potential is likely independent of the initial transmission timing.

The role of lifestyle and behavioral factors, specifically smoking, also warrants closer examination. Biologically, tobacco metabolites are known to concentrate in the cervical mucus, causing DNA damage and local immunosuppression that facilitates persistent HPV infection.^{39,40} The nicotine and cotinine metabolites act as co-carcinogens, potentially accelerating the transition from a transient infection to a malignant lesion. In our study, 93.4% of participants were non-smokers, and no significant correlation was observed between smoking status and HPV types ($p = 0.08$), potentially due to the very low prevalence of smoking (2.2%) among women in our cohort, hindering the ability to detect any genotype-specific interactions. This suggests that while smoking is a potent co-factor for cervical carcinogenesis in general, it may not exert a selective pressure that favors one HPV genotype over another in this specific demographic.

Finally, the advanced stage of diagnosis for many patients—with 25.3% at Stage 2—highlights a critical clinical reality: late-stage presentation is common in Indonesian referral hospitals. The lack of correlation between cancer stage and HPV genotype suggests that virulence is not necessarily determined by the specific viral strain but rather by delays in secondary prevention and access to screening. Despite the limitations of a cross-sectional design, the identification of HPV 18 and 52 as major genotypes, alongside the significant roles of early marriage and hormonal contraception, provides critical data for the development of domestic diagnostic tools and vaccine policy in Indonesia. Tailoring public health interventions to the local virome is essential for achieving the WHO's goal of eliminating cervical cancer as a public health threat by the end of the decade.

5. Conclusion

This study provides a descriptive overview of HPV genotype distribution among cervical cancer patients in Indonesia, identifying early age at marriage ($p = 0.004$) and hormonal contraceptive use ($p = 0.002$) as significant predictors for high-risk HPV genotype. The findings highlight a unique regional virome dominated by HPV 18, 52, and 16, suggesting that regional viral epidemiology may differ significantly from global patterns, which are dominated by HPV 16 and 18. This prevalence of non-HPV 16 genotypes emphasizes the need for nonavalent vaccines and locally

tailored diagnostic tools.

While other factors, such as high parity and smoking, did not show genotype-specific correlations, they remain common clinical markers in this cohort. Furthermore, the 35.2% HPV-negative rate underscores the technical challenges of detection in advanced malignancies. These results provide a critical foundation for optimizing cervical cancer screening and prevention strategies specifically for the Indonesian population.

Acknowledgments

The authors would like to thank the Center for Biomedical Research, Organization for Health, National Research and Innovation (BRIN), and the Biomolecular Laboratory, Faculty of Medicine, Universitas Trisakti, for supporting this research.

Funding

This research was supported by the RIIM LPDP Grant and BRIN (grant number 51/IV/KS/06/2022).

Conflict of interest

The authors declare they have no competing interests.

Author contributions

Conceptualization: Raditya Wratsangka, Monica D. Hartanti

Formal analysis: Raditya Wratsangka, Monica D. Hartanti, Lily M. Surjadi

Investigation: Monica D. Hartanti, Gatot Purwoto, Miranti P. Aulia, Alvionita Kogoya

Methodology: Monica D. Hartanti, Gatot Purwoto, Miranti P. Aulia

Writing—original draft: Raditya Wratsangka, Monica D. Hartanti, Miranti P. Aulia

Writing—review & editing: All authors

Ethics approval and consent to participate

This study was approved by the ethical clearance committee of the Faculty of Medicine, Universitas Indonesia, with ethical clearance number KET-1262/UN2.F1/ETIK/PPM.00.02/2022. Written informed consent was obtained from participants before enrolment.

Consent for publication

This study involves fully anonymized, cross-sectional data and does not contain any images or information that could identify a subject. Written informed consent was obtained from all participants for the publication of this study. All data have been anonymized to ensure participant

confidentiality.

Availability of data

All data generated or analyzed during this study are included in this published article.

References

1. ASCO. Challenges and Opportunities for Global Cervical Cancer Elimination: How Can We Build a Model for Other Cancers? *Am Soc Clin Oncol Educ Book*. 2025;45(3):e473702. doi: 10.1200/EDBK-25-473702.
2. World Health Organization, 2008. Cervical cancer, Fact Sheet. Updated December 2, 2025. Available from: <https://www.who.int/news-room/fact-sheets/detail/cervical-cancer> (Last accessed on 2025 Feb 23).
3. Wu J, Jin Q, Zhang Y, *et al*. Global burden of cervical cancer: current estimates, temporal trend and future projections based on the GLOBOCAN 2022. *J Natl Cancer Cent*. 2025;5(3):322-329. doi: 10.1016/j.jncc.2024.11.006
4. World Health Organization, 2024. WHO, UNFPA commend Indonesia's efforts to eliminate cervical cancer, urge streamlined vaccine strategy and enhanced screening. Posted November 15, 2024. Available from: <https://www.who.int/indonesia/news/detail/15-11-2024-who--unfpa-commend-indonesia-s-efforts-to-eliminate-cervical-cancer--urge-streamlined-vaccine-strategy-and-enhanced-screening#:~:text=The%20UN%20agencies%2C%20which%20supported,invasive%20cancer%20to%20be%20managed> (Last accessed on 2025 Dec 28).
5. World Health Organization, 2021. WHO recommends DNA testing as a first-choice screening method for cervical cancer prevention. Posted September 11, 2011. Available from: <https://www.who.int/europe/news/item/11-09-2021-who-recommends-dna-testing-as-a-first-choice-screening-method-for-cervical-cancer-prevention> (Last accessed on 2025 Jan 31).
6. Kusakabe M, Taguchi A, Sone K, Mori M, Osuga Y. Carcinogenesis and management of human papillomavirus-associated cervical cancer. *Int J Clin Oncol*. 2023;28(8):965-974. doi: 10.1007/s10147-023-02337-7
7. United Nations Children's Fund (UNICEF). 2023. Fast facts about HPV and cervical cancer. Available from: <https://www.unicef.org/stories/fast-facts-hpv-cervical-cancer> (Last accessed on 2025 Feb 15).
8. Calagna G, Maranto M, Paola C, *et al*. "Secondary prevention" against female HPV infection: literature review of the role of carrageenan. *Expert Rev Anti Infect Ther*. 2020;18(9):865-874. doi: 10.1080/14787210.2020.1770082
9. Capra G, Giovannelli L, Matranga D, *et al*. Potential impact of a nonavalent HPV vaccine on HPV related low- and high-grade cervical intraepithelial lesions: A referral hospital-based study in Sicily. *Hum Vaccin Immunother*. 2017;13(8):1839-1843. doi: 10.1080/21645515.2017.1319026
10. Ministry of Health, Republic of Indonesia. *Pedoman Nasional Pelayanan Kedokteran Tata Laksana Kanker Serviks* [National guidelines for treatment of cervical cancer]. 2018. HK.01.07/MENKES/349/2018. Available from: https://yankes.kemkes.go.id/unduh/fileunduh/1610415049_651524.pdf (Last accessed on 2025 Mar 15).
11. Ministry of Health, Republic of Indonesia, 2023. National Cervical Cancer Elimination (NCCE) Plan for Indonesia 2023-2030. Updated Dec 22, 2023. Available from: <https://kemkes.go.id/id/national-cervical-cancer-elimination-plan-for-indonesia-2023-2030> (Last accessed on 2025 Dec 28).
12. Robbers GML, Bennett LR, Spagnoletti BRM, Wilopo SA. Facilitators and barriers for the delivery and uptake of cervical cancer screening in Indonesia: A scoping review. *Glob Health Action*. 2021;14(1):1979280. doi: 10.1080/16549716.2021.1979280
13. Hutajulu SH, Prabandari YS, Bintoro BS, *et al*. Delays in the presentation and diagnosis of women with breast cancer in Yogyakarta, Indonesia: A retrospective observational study. *PLoS one*. 2022;17(1):e0262468. doi: 10.1371/journal.pone.0262468
14. Burd EM. Human papillomavirus and cervical cancer. *Clin Microbiol Rev*. 2023;16(1):1-17. doi: 10.1128/CMR.16.1.1-17.2003
15. Kombe Kombe AJ, Li B, Zahid A, *et al*. Epidemiology and Burden of Human Papillomavirus and Related Diseases, Molecular Pathogenesis, and Vaccine Evaluation. *Front Public Health*. 2020;8:552028. doi: 10.3389/fpubh.2020.552028
16. Bosco L, Serra N, Fasciana T, *et al*. Potential impact of a nonavalent anti HPV vaccine in Italian men with and without clinical manifestations. *Sci Rep*. 2021;11(1):4096. doi: 10.1038/s41598-021-83639-6
17. Hu Z, Zhu D, Wang W, *et al*. Genome-wide profiling of HPV integration in cervical cancer identifies clustered genomic hot spots and a potential microhomology-mediated integration mechanism. *Nat Genet*. 2015;47(2):158-163. doi: 10.1038/ng.3178
18. Hartanti MD, Rachmadhany EDA, Mustopa ITI, *et al*. High mobility group AT-Hook 2 expression is elevated in cervical cancer. *Int J Pharm Res*. 2020;12:2332-2338. doi: 10.31838/ijpr/2020.12.04.322
19. Doorbar J. The human Papillomavirus twilight zone -

- Latency, immune control and subclinical infection. *Tumour Virus Res.* 2023;16:200268.
doi: 10.1016/j.tvr.2023.200268
20. Kogoya A, Hartanti MD. Correlations of Expression of KLF5, FHIT and DLG2 with Cervical Cancer. In: *Advances in Social Science, Education and Humanities Research*. Paris, France: Atlantis Press SARI; 2024:1302-1309.
doi: 10.2991/978-2-38476-273-6_132
21. World Health Organization. Global strategy to accelerate the elimination of cervical cancer as a public health problem. Updated Nov 17, 2020. Available from: <https://www.who.int/publications/i/item/9789240014107> (Last accessed on 2025 Apr 18).
22. Hartanti MD, Amtha R, Wratsangka R, *et al.* Operational Feasibility and Performance of mRPA Versus PCR for the Detection of HPV Types 16, 18, and 52 in Clinical Samples: An Exploratory Study. *Int J Womens Health.* 2025;17:955-964.
doi: 10.2147/IJWH.S496621
23. Burger EA, Kim JJ, Sy S, *et al.* Age of Acquiring Causal Human Papillomavirus (HPV) Infections: Leveraging Simulation Models to Explore the Natural History of HPV-Induced Cervical Cancer. *Clin Infect Dis.* 2017;65(6):893-899.
doi: 10.1093/cid/cix475
24. Liu F, Chang L, Bai T, Liu X, Hu J. Association of human papillomavirus genotype distribution and cervical cytology: a cross-sectional study. *Epidemiol Infect.* 2021;149:e95.
doi: 10.1017/S0950268821000741
25. Tenjimbayashi Y, Onuki M, Hirose Y, *et al.* Whole-genome analysis of human papillomavirus genotypes 52 and 58 isolated from Japanese women with cervical intraepithelial neoplasia and invasive cervical cancer. *Infect Agent Cancer.* 2017;12:44.
doi: 10.1186/s13027-017-0155-4
26. Lin Y, Lin WY, Lin TW, *et al.* Trend of HPV Molecular Epidemiology in the Post-Vaccine Era: A 10-Year Study. *Viruses.* 2023;15(10):2015.
doi: 10.3390/v15102015
27. Gadducci A, Cosio S, Fruzzetti F. Estro-progestin Contraceptives and Risk of Cervical Cancer: A Debated Issue. *Anticancer Res.* 2020;40(11):5995-6002.
doi: 10.21873/anticancer.14620
28. Bergqvist L, Kalliala I, Aro K, *et al.* Distribution of HPV Genotypes Differs Depending on Behavioural Factors among Young Women. *Microorganisms.* 2021;9(4):750.
doi: 10.3390/microorganisms9040750
29. Lee SA, Baik S, Chung SH. Functional roles of female sex hormones and their nuclear receptors in cervical cancer. *Essays Biochem.* 2021;65(6):941-950.
doi: 10.1042/EBC20200175
30. James CD, Morgan IM, Bristol ML. The Relationship between Estrogen-Related Signaling and Human Papillomavirus Positive Cancers. *Pathogens (Basel, Switzerland).* 2020;9(5):403.
doi: 10.3390/pathogens9050403
31. Qi M, Naranjo AR, Duque AJ, *et al.* Evaluation of Pre-Analytical Variables for Human Papillomavirus Primary Screening from Self-Collected Vaginal Swabs. *J Mol Diagn.* 2024;26(6):487-497.
doi: 10.1016/j.jmoldx.2024.02.006
32. Dai W, Zeng X, Pantanowitz L, Zhao C. Human papillomavirus negative cervical cancers and precancerous lesions: prevalence, pathological and molecular features, and clinical implications. *Gynecol Obstet Clin Med.* 2025;5:e000160.
doi: 10.1136/gocm-2024-000160
33. Martinelli C, Ercoli A, Parisi S, *et al.* Molecular Mechanisms and Clinical Divergences in HPV-Positive Cervical vs. Oropharyngeal Cancers: A Critical Narrative Review. *BMC Medicine.* 2025;23(1):405.
doi: 10.1186/s12916-025-04247-z
34. Patel A, Galaal K, Burnley C, *et al.* Cervical cancer incidence in young women: a historical and geographic controlled UK regional population study. *Br J Cancer.* 2021;106(11):1753-1759.
doi: 10.1038/bjc.2012.148
35. Lau HY, Juang CM, Chen YJ, *et al.* Aggressive characteristics of cervical cancer in young women in Taiwan. *Int J Gynaecol Obstet.* 2009;107(3):220-223
doi: 10.1016/j.ijgo.2009.07.029
36. Li S, Hu T, Lv W, *et al.* Changes in prevalence and clinical characteristics of cervical cancer in the People's Republic of China: A study of 10,012 cases from a nationwide working group. *Oncologist.* 2013;18(10):1101-1107.
doi: 10.1634/theoncologist.2013-0123
37. Itarat Y, Kietpeerakool C, Jampathong N, *et al.* Sexual behavior and infection with cervical human papillomavirus types 16 and 18. *Int J Women's Health.* 2019;11:489-494.
doi: 10.2147/IJWH.S218441
38. Luo D, Peng M, Wei X, *et al.* Prevalence of Human Papillomavirus and Genotype Distribution in Pregnant and Non-Pregnant Women in China. *Risk Manag Healthc Policy.* 2021;14:3147-3157.
doi: 10.2147/RMHP.S288838
39. Aguayo F, Munoz JP, Perez-Dominguez F, *et al.* High-Risk

Human Papillomavirus and Tobacco Smoke Interactions in Epithelial Carcinogenesis. *Cancers*. 2020;12(8):2201.

doi: 10.3390/cancers12082201

40. Ma K, Li S, Wu S, *et al.* Impact of smoking exposure on

human papillomavirus clearance among Chinese women: A follow-up propensity score matching study. *Tobacco Induced Diseases*. 2023;21:42.

doi: 10.18332/tid/161026

Risk factors and human papillomavirus genotypes: Investigating their interplay in cervical cancer

By Raditya Wratsangka

ORIGINAL RESEARCH ARTICLE

Risk factors and human papillomavirus genotypes: Investigating their interplay in cervical cancer

Raditya Wratsangka^{1†}, Monica D. Hartanti^{2,3†}, Gatot Purwoto⁴, Lily M. Surjadi¹, Miranti P. Aulia⁵, and Alvionita Kogoya⁶

¹Department of Obstetric and Gynecology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

²Department of Biology, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

³Center for Biomedical Research, Research Organization for Health, National Research and Innovation Agency, Bogor, West Java, Indonesia

⁴Department of Obstetrics and Gynecology, Cipto Mangunkusumo National Hospital, Faculty of Medicine, Universitas Indonesia, Jakarta, Indonesia

⁵Independent Researcher, Jakarta, Indonesia

⁶Biomolecular Laboratory, Faculty of Medicine, Universitas Trisakti, Jakarta, Indonesia

[†]These authors contributed equally to this work.

***Corresponding author:**
Monica D. Hartanti
(mdhartanti@trisakti.ac.id)

Citation: Wratsangka R, Hartanti MD, Purwoto G, Surjadi LM, Aulia MP, Kogoya A. Risk factors and human papillomavirus genotypes: Investigating their interplay in cervical cancer. *Eurasian J Med Oncol.* 2026;10(1):318-327. doi: 10.36922/EJMO025490509

Received: December 6, 2025

Revised: January 14, 2026

Accepted: January 20, 2026

Published online: February 27, 2026

Copyright: © 2026 Author(s). This is an Open-Access article distributed under the terms of the Creative Commons Attribution License, permitting distribution, and reproduction in any medium, provided the original work is properly cited.

Publisher's Note: AccScience Publishing remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Abstract

Introduction: Cervical cancer continues to pose a substantial worldwide health challenge, primarily linked to human papillomavirus (HPV) infection. In regions like Indonesia, understanding local genotype distribution is critical for refining screening and vaccination efforts.

Objective: This research explores the complex relationship between various risk factors and the types of HPV detected in cervical cancer patients at a national referral center in Indonesia.

Methods: Ninety-one women diagnosed with cervical cancer were recruited for this cross-sectional study at Cipto Mangunkusumo National Hospital, Jakarta—the top-tier referral facility in Indonesia. This setting provides a diverse patient demographic representing various geographical regions across the archipelago. Cervical swabs were collected for HPV genotyping using dot blot hybridization. Various clinical risk factors, including age, age at marriage, age at first pregnancy, parity, smoking habits, hormonal contraceptive history, and the stage of cervical cancer diagnosis, were assessed. Statistical analysis was conducted using Fisher's exact test to accommodate the distribution of the data.

Results: High-risk (HR) HPV was detected in 62.6% of participants. HPV type 18 was most prevalent (15.4%), followed by types 52 (14.3%) and 16 (12.1%). Statistically significant associations were observed between HR-HPV presence and both early age at marriage ($p = 0.004$) and the use of hormonal contraception ($p = 0.002$). No significant correlations were observed between some patient characteristics (age, parity, and smoking habits) and HPV genotypes ($p > 0.05$). A total of 35.2% of samples were HPV-negative, potentially reflecting technical challenges or viral integration in advanced-stage cases.

Conclusion: These findings suggest that early marriage and hormonal contraceptive use are significant clinical markers for HR-HPV in this Indonesian population. The dominance of HPV 18 and 52 highlights the need for locally relevant diagnostic tools and the potential transition to nonavalent vaccines. These data are essential for developing locally relevant diagnostic tools and optimizing cervical cancer prevention strategies tailored to the Indonesian demographic.

Keywords: Cervical cancer; HPV genotypes; Hormonal contraception; Age at marriage; Indonesia

1. Introduction

Cervical cancer stands as a significant global health issue, ranking as the fourth most prevalent malignancy among women worldwide.¹ This malignancy represents a profound burden on healthcare systems, particularly in developing nations where screening infrastructure is often fragmented. According to data from the World Health Organization (WHO) in 2022, approximately 660,000 new cases and 350,000 deaths from cervical cancer were reported, with approximately 94% occurring in low- and middle-income countries.² The disparity in mortality rates between high-income and low-income regions underscores the critical need for localized epidemiological data to inform public health strategies. In many developing regions, the lack of centralized registries and limited access to pathology services further complicates the accurate mapping of disease prevalence.

The global incidence of cervical cancer ranges from 10 to 75 cases per 100,000 women.³ Indonesia, as of 2022 data, recorded 36,000 cases of cervical cancer and 21,000 associated deaths.⁴ The high mortality-to-incidence ratio in Indonesia suggests that many cases are detected at advanced stages, limiting treatment efficacy. This clinical reality is exacerbated by the geographical complexity of the Indonesian archipelago, where the distribution of healthcare resources is often uneven, leaving rural populations with limited access to life-saving interventions. In 2021, the WHO recommended the DNA-HPV test as a primary screening method.⁵ This shift from cytology-based screening to molecular-based testing represents a paradigm shift in early detection, aiming to identify high-risk viral presence before cellular abnormalities manifest.

Cervical cancer is preventable, with human papillomavirus (HPV) infection being a crucial etiological factor. HPV is present in over 95% of cervical cancer lesions.^{6,7} The virus targets the basal layer of the stratified squamous epithelium, utilizing the host cell's machinery to replicate its double-stranded DNA genome. While there is no direct treatment for the virus, some topical

microbicides have shown intrinsic antiviral activity.⁸ Despite these developments, the primary defense against cervical malignancy remains a combination of prophylactic vaccination and secondary prevention through rigorous screening protocols. Although most HPV infections are asymptomatic and can be resolved on their own, persistent HPV infection poses a risk of developing into precancerous lesions, ultimately leading to cervical cancer.⁹ High-risk HPV types, notably HPV 16 and 18, are responsible for low- and high-grade squamous intraepithelial lesions (LSIL and HSIL) in the cervix. Timely identification and treatment of precancerous lesions can prevent progression.¹⁰ Vaccination, protecting individuals from HPV infection, has demonstrated success in reducing cervical cancer incidence in developed countries alongside regular screening for adult women. However, the efficacy of these vaccines is largely dependent on alignment between the vaccine's targeted genotypes and the circulating genotypes within a specific geographic population.

Low- and middle-income countries often face limitations in preventive facilities and access, leading to advanced-stage diagnoses, as observed in Indonesia, where approximately 66.4% of patients are diagnosed at stages IIB-IVB.¹⁰ In Indonesia, logistical barriers, socio-economic factors, and a lack of widespread awareness contribute to the high prevalence of late-stage presentations.^{11,12} Furthermore, cultural taboos regarding pelvic examinations and reproductive health can deter women from seeking early screening, even when services are available.^{12,13}

As a member of the Papillomaviridae family, HPV encompasses a diverse group of over 200 distinct genotypes. These viruses specifically target the cutaneous and mucosal squamous epithelium, leading to their primary classification by tissue type. Regarding oncogenesis, these strains are further divided into low-risk types, which typically result in benign growths, and high-risk types, which are the fundamental drivers of malignant transformations.¹⁴ The pathogenic mechanism of high-risk HPV centers on the activity of the E6 and E7 oncoproteins; these viral products

facilitate cancer progression by inhibiting the essential tumor-suppressing functions of the p53 and pRb proteins.

Globally, HPV represents a pervasive sexually transmitted infection that impacts all genders. Epidemiological data indicate that genital HPV prevalence varies significantly, ranging from 3.5% to 45% in male populations and 2% to 44% in female cohorts, with comparable rates of transmission observed between them.¹⁵ In men, the majority of infections remain subclinical and resolve without medical intervention. When symptoms do manifest in males, they usually present as benign genital warts. Similarly, while a high percentage of women contract cervical HPV during their lifetime, the vast majority of these cases do not escalate into invasive malignancy.¹⁶

A critical milestone in the transition to cervical cancer is the integration of the HPV genome into the host cell's DNA. This molecular event frequently disrupts the viral E2 gene, causing an unchecked surge in E6 and E7 expression that drives rapid cellular transformation. Because symptoms often remain dormant for months or even years following the initial infection, clinical detection is frequently delayed until the appearance of physical warts or advanced lesions.¹⁷⁻²⁰ To combat this, the WHO has published the 90-70-90 targets for 2030: ensuring 90% vaccination coverage for girls by age 15, 70% screening participation for women at ages 35 and 45, and 90% access to treatment for those diagnosed with cervical disease.²¹ Indonesia has aligned with these goals by integrating HPV vaccination into its national immunization framework for adolescents. While this strategy is vital for long-term control, its success depends on logistical stability and public trust. Currently, the program does not yet extend to the adult female demographic.

To align with the government's goals of developing domestic diagnostic tools and expanding screening reach, it is crucial to develop cost-effective, sensitive, and specific diagnostic reagents tailored to the high-risk HPV genotype prevalence in the Indonesian population.²² Current diagnostic kits often use primers designed for Western populations, which may not achieve optimal sensitivity due to the unique genotype distribution in Southeast Asia. This preliminary study aims to correlate risk factors with HPV genotypes among women in Indonesia. Given the cohort size, this research serves as an exploratory analysis to identify patterns that warrant larger, multicenter investigations into the complex interplay between patients' reproductive histories and viral characteristics. By establishing a clearer picture of the local molecular landscape, we can better inform the selection of vaccines and the development of localized screening protocols that are both economically viable and clinically effective.

56

2. Materials and methods

2.1. Ethical clearance

4

This study was approved by the ethical clearance committee of the Faculty of Medicine, Universitas Indonesia, with ethical clearance number KET-1262/UN2.F1/ETIK/PPM.00.02/2022. Written informed consent was obtained from subjects before enrolment, ensuring that all participants were fully apprised of the study's objectives and the nature of the clinical specimens to be collected. The informed consent process was conducted in the local language to ensure a comprehensive understanding of the voluntary nature of participation and the confidentiality of genomic data.

21

2.2. Study design and population

This was a cross-sectional study, observing the relationship between risk factors and HPV genotyping among women in Indonesia. It was conducted in 2022 at Cipto Mangunkusumo National Hospital, Jakarta, with targeted participants being women aged 18–65 years old. The hospital serves as a national referral center, providing a diverse patient demographic for epidemiological research. Only women who had signed the informed consent were included in this study. Women who have had a prior HPV vaccination or a treatment related to cervical cancer were excluded from this study to prevent the confounding effects of medical intervention on the natural distribution of HPV genotypes. These exclusion criteria were strictly maintained to ensure that the observed viral prevalence reflects the natural history of the infection within the local community rather than the impact of clinical prophylaxis.

2.3. Data collection

Ninety-one participants were recruited consecutively according to inclusion and exclusion criteria. While the cohort size reflects the preliminary and exploratory nature of this research, rigorous data collection protocols were maintained. Questionnaires were distributed to participants, containing detailed questions related to marriage status, smoking, parity status, and hormonal contraceptive use. Information regarding age at marriage and first pregnancy was recorded to assess early exposure to potential risk factors. For HPV genotyping, cervical swabs were collected in ThinPrep containers and stored at 4 °C for analysis. Maintaining a strict cold chain was essential to prevent viral DNA degradation by endogenous nucleases in clinical samples.

41

2.4. HPV genotyping

16

DNA extraction was performed from cervical swabs using the QIAamp® DNA Blood Mini Kit (cat. 51104, Qiagen,

Netherlands) according to the manufacturer's protocol. Briefly, the sample in pellet resuspension was added to a mixture of Qiagen Protease (proteinase K) and Buffer AL. The addition of proteinase K is a critical step to ensure the complete lysis of cervical epithelial cells and the digestion of proteins that might otherwise sequester viral DNA. After a 10-min incubation at 56 °C, 96–100% ethanol was added. The mixture was then transferred to the QIAamp Mini Spin Column and centrifuged at $6000 \times g$ for 1 min. The use of high-fidelity spin columns ensures the removal of potential polymerase chain reaction (PCR) inhibitors, though the final DNA yield remains dependent on the initial cellularity of the swab. These inhibitors, often present in clinical mucus and blood, can lead to false-negative results if not thoroughly washed away during purification. Buffers AW2 and AE were added separately after serial centrifugation at different speeds and times.

The extracted DNA was then amplified, and HPV was genotyped using a reverse dot blot “flow-through” hybridization method with GeneFlow Human Papillomavirus (HPV) array test kit (cat. 92007; DiagCor Life Science Ltd., Hong Kong). The flow-through hybridization technology uses a patented active-pumping mechanism to direct target DNA to the probes, significantly increasing reaction speed and sensitivity compared to traditional passive hybridization. This hybridization-based approach enables simultaneous identification of multiple genotypes in a single sample, a critical step for detecting coinfections. This method can detect different high-risk HPV types, including 16, 18, 31, 33, 35, 39, 45, 51, 52, 53, 56, 58, 59, 66/68, 73, 81 and 82; and low-risk HPV types, including 6, 11, 26, 40, 42, 43, 44, 54, 55, 57, 61, 70, 71, 72, and 84. The broad coverage of this array is particularly advantageous for exploratory studies in regions where non-standard genotypes may be prevalent.

25 2.5. Statistical analysis

The data were collected and stored in Microsoft Excel 2025 (version 2511, Microsoft, USA). Categorical variables were summarized as absolute counts and percentages and analyzed using Fisher's exact test. This test is preferred to avoid the inaccuracies of the Chi-square approximation in sparse contingency tables, as it provides an exact probability value even when several cells have expected values of less than five. A p -value < 0.05 was considered significant. All statistical analyses were carried out using GraphPad version 9.0.0 (Dotmatics, United States). Due to the relatively small sample size ($n = 91$), this study lacked sufficient power to perform multivariable logistic regression to control confounding factors such as age and cancer stage. Consequently, the findings presented here are descriptive and exploratory. The high rate of HPV-negative results was analyzed separately to ensure no systematic bias

related to patient age or diagnostic stage was introduced.

3. Results

This study highlighted the characteristics of women with cervical cancer and correlated those characteristics with HPV genotyping. These findings illuminate the multifaceted nature of patient characteristics in the study, underscoring the endeavor to explore the correlations between risk factors and HPV genotyping in cervical cancer among Indonesian patients. While the sample size was 91, the findings remained primarily exploratory; however, the cohort revealed statistically significant associations that had previously been obscured. Notably, the data demonstrated that age at marriage and hormonal contraceptive use were key clinical markers for the presence of high-risk HPV genotypes.

3.1. Characteristics of participants

The study emphasized the significance of patient age and reproductive history, as detailed in Table 1. The age distribution varied, with a minor proportion in the 25–34 age group (12.1%) and the majority (36.3%) aged 50 or older. Age at marriage emerged as a noteworthy factor. Roughly 29.7% of patients were married before age 20, while approximately 69.2% married after age 20. There was substantial variation in the age at which patients experienced their first pregnancy, with 72.5% experiencing it between the ages of 20 and 35. Differences in parity status were observed: 7.7% were nulliparous (never having given birth) and 91.2% were multiparous (having given birth more than once). Smoking habits exhibited significant variability among patients, with 6.6% identified as active smokers and 93.4% as non-smokers. Diagnostic data reflected the severity of cervical cancer in the patients. A minority (12.1%) received a pre-cancer diagnosis, while 25.3% were at stage 2, 7.7% at stage 3, and 1.1% at stage 4. This distribution confirms that the majority of participants had advanced-stage disease. The patterns of contraceptive use were diverse, with 30.8% of patients not using contraception and 69.2% using contraceptive methods.

3.2. HPV genotyping distribution

The study unveiled a diversity of HPV infections. Notably, 35.2% of the cohort ($n = 32$) had no detectable high-risk or low-risk HPV types—a finding that warrants further technical and clinical investigation. High-risk HPV types were detected in 62.6% of participants, while 2.2% had low-risk HPV types. Among the high-risk HPV types identified, HPV 18 was the most prevalent (15.4%), followed by HPV 52 (14.3%) and HPV 16 (12.1%). Other identified high-risk HPV types included HPV 45 (5.5%) and HPV 59 (3.3%). Coinfections were found in three women, possessing HPV 16 and 52; HPV 16 and 31; and

HPV 18, 66, and 68.

36
Table 1. Characteristics of participants

Characteristics	Count (n)	Percentage (%)
Age (years)		
25–34	11	12.1
35–44	25	27.5
45–50	22	24.2
>50	33	36.3
Age at marriage		
<20	27	29.7
>20	63	69.2
Age at first pregnancy		
<20	18	19.8
20–35	66	72.5
Parity number		
Nulliparous	7	7.7
Multiparous	83	91.2
Smoking		
Yes	6	6.6
No	85	93.4
Cancer stage		
Pre-Cancer	11	12.1
Stadium 1	9	9.9
Stadium 2	23	25.3
Stadium 3	7	7.7
Stadium 4	1	1.1
Hormonal Contraception		
Yes	28	30.8
No	63	69.2
HPV genotyping		
No HPV	32	35.2
High-risk HPV	57	62.6
Low-risk HPV	2	2.2
HPV type		
16	11	12.1
18	14	15.4
26	1	1.1
35	1	1.1
45	5	5.5
52	13	14.3
58	1	1.1
59	3	3.3

3.3. Relationship between risk factors and HPV genotyping

The interplay between clinical risk factors and the specific HPV genotypes was analyzed using Fisher's exact test, with results presented in Table 2. Significant correlations were observed between specific reproductive behaviors and the presence of high-risk HPV genotypes. Specifically, a significant association was observed regarding the age at marriage ($p = 0.004$), where patients who married before the age of 20 were significantly more likely to test positive for high-risk HPV (85.2%) compared to those who married at a later age (53.1%). Furthermore, the use of hormonal contraceptives demonstrated a strong statistical association with high-risk HPV detection ($p = 0.002$), with 85.7% of patients utilizing hormonal methods carrying high-risk HPV genotypes, whereas only 52.4% of non-users tested positive. No significant correlations were observed between HPV risk status and other variables, including age ($p = 0.514$), parity ($p = 1.000$), age at pregnancy ($p = 0.178$), and smoking habits ($p = 0.08$). While certain factors did not reach statistical significance, the findings regarding early marriage and hormonal contraception provide critical insights. These results indicate that in this specific Indonesian referral hospital population, HPV genotype distribution is significantly linked to early sexual exposure and hormonal history, suggesting that preventative strategies and screening in this demographic should prioritize individuals with these specific clinical markers.

4. Discussion

35
The findings of this study provide a nuanced look into the epidemiological landscape of HPV genotypes among cervical cancer patients in Indonesia. The primary observation of this research is the significant correlation between clinical markers—specifically at marriage and hormonal contraceptive use—and the presence of high-risk HPV. While established clinical risk factors—such as parity and smoking—are well-documented drivers of cervical carcinogenesis, the lack of statistical significance in our cohort ($p > 0.05$) may be attributed to the overwhelming prevalence of these factors across the entire sample, thereby reducing their discriminatory power in a bivariate analysis. However, the identification of high-risk HPV in 62.6% of participants provides a more robust framework for identifying viral patterns that deviate from Western-centric models.

One of the most striking results is the high prevalence of HPV 18 (15.4%) and HPV 52 (14.3%), which, along with HPV 16, constitute the primary high-risk HPV types in this cohort. This distribution aligns with emerging data from Southeast Asia, suggesting that the regional

virome may differ from that of Western populations, in which HPV 16 and 18 are more dominant.^{20,23} The high frequency of HPV 18 and 52 reflects a specific molecular epidemiology in the Indonesian archipelago. Previous studies in neighboring regions have suggested that HPV 52 and 58 exhibit a higher fitness or transmission efficiency in certain Asian demographics, possibly due to host genetic factors or environmental selection pressures.^{24,25} The high frequency of HPV 52 has profound implications for local public health, particularly regarding the efficacy of current vaccines. Given that HPV 18 and 52 are major drivers in this population, transitioning to nonavalent vaccines or developing locally tailored diagnostic reagents is a clinical priority.²⁶

Table 2. Association of risk factors with HPV genotyping in women with cervical cancer

Risk Factors	No HPV/low-risk HPV	High-risk HPV	p value
Age			
<45	15	21	0.514
≥45	19	36	
Age at marriage			
<20	4	23	0.004*
>20	30	34	
Age at first pregnancy			
<20	4	14	0.178
20–35	30	43	
Parity number			
Nulliparous	3	4	1.000
Multiparous	31	53	
Smoking			
Yes	0	6	0.08*
No	34	51	
Hormonal contraception			
Yes	4	24	0.002*
No	30	33	

Notes: Fisher's test analysis was used to determine the relationships of each variable. *p < 0.05.

A noteworthy finding in this study is the significant association between early age at marriage and high-risk HPV ($p = 0.004$). Specifically, 85.2% of women who married before age 20 tested positive for high-risk HPV genotypes. Similarly, the use of hormonal contraceptives (reported by 30.8% participants) was linked to the development of cervical cancer. However, this has become

a controversial topic due to inconsistent results. We did not find any significant association between the use of hormonal contraceptives and specific HPV genotypes ($p > 0.05$). Long-term use of oral contraceptives has been hypothesized to increase cervical cancer risk by making the cervix more susceptible to HPV DNA integration through hormonal modulation of the viral upstream regulatory region.^{27,28} Specifically, estrogen and progesterone may influence the expression of the viral E6 and E7 genes through hormone-responsive elements in the viral long control region.^{29,30} Our findings suggest that in the Indonesian context, the choice of contraception does not appear to influence the type of high-risk HPV that leads to malignancy. This reinforces the conclusion that once a high-risk HPV infection is established, the progression to cancer is likely driven by the oncogenic potential of the virus itself rather than the specific method of birth control used by the patient.

A significant point of discussion is the high proportion of HPV-negative results (35.2%) observed in this study. While cervical cancer is almost universally linked to HPV, literature does report HPV-negative cases, though usually at a lower rate than seen here. Several factors may explain this discrepancy. First, technical limitations in preserving DNA in a tropical climate could lead to sample degradation.³¹ Second, the sensitivity of the reverse dot blot hybridization method, while high, may have a detection threshold that misses low-copy-number viral DNA.³² Furthermore, the molecular architecture of the virus changes as the disease progresses. In advanced-stage cancer (which characterized 8.8% of our participants), the HPV genome often integrates into the host cell DNA, sometimes resulting in the loss of the L1 gene—the target most PCR-based assays use for detection.³³ This false-negative phenomenon in advanced malignancies is a documented challenge in molecular oncology.^{34,35} Future studies should use multiple primer sets targeting different viral regions (e.g., E6/E7) to improve detection rates.

Regarding other reproductive factors, our study found that 91.2% of participants were multiparous. High parity is thought to increase cervical cancer risk through sustained hormonal changes and physical trauma to the transformation zone during childbirth.^{36–38} The repeated expansion and remodeling of the cervical transformation zone during multiple pregnancies may create a microenvironment that is more susceptible to persistent viral infection and subsequent cellular transformation. However, our analysis did not demonstrate an association between parity and specific HPV genotype ($p = 1.000$). This suggests that while parity helps the cancer progress, it may not be a selective pressure on HPV genotype. Similarly,

early age at first pregnancy (often proxies for early sexual debut) was common in our cohort but did not significantly correlate with HPV genotype. These findings reinforce the idea that once a high-risk HPV infection is established, the viral genotype's oncogenic potential is likely independent of the initial transmission timing.

The role of lifestyle and behavioral factors, specifically smoking, also warrants closer examination. Biologically, tobacco metabolites are known to concentrate in the cervical mucus, causing DNA damage and local immunosuppression that facilitates persistent HPV infection.^{39,40} The nicotine and cotinine metabolites act as co-carcinogens, potentially accelerating the transition from a transient infection to a malignant lesion. In a ³⁴ study, 93.4% of participants were non-smokers, and ³⁴ no significant correlation was observed between smoking status and HPV types ($p = 0.08$), potentially due to the very low prevalence of smoking (2.2%) among women in our cohort, hindering the ability to detect any genotype-specific interactions. This suggests that while smoking is a potent co-factor for cervical carcinogenesis in general, it may not exert a selective pressure that favors one HPV genotype over another in this specific demographic.

Finally, the advanced stage of diagnosis for many patients—with 25.3% at Stage 2—highlights a critical clinical reality: late-stage presentation is common in Indonesian referral hospitals. The lack of correlation between cancer stage and HPV genotype suggests that virulence is not necessarily determined by the specific viral strain but rather by delays in secondary prevention and access to screening. Despite the limitations of a cross-sectional design, the identification of HPV 18 and 52 as major genotypes, alongside the significant roles of early marriage and hormonal contraception, provides critical data for the development of domestic diagnostic tools and vaccine policy in Indonesia. Tailoring public ¹⁸ health interventions to the local virome is essential for achieving the WHO's goal of eliminating cervical cancer as a public health threat by the end of the decade.

5. Conclusion

This study provides a descriptive overview of HPV genotype distribution among cervical cancer ⁴² patients in Indonesia, identifying early age at marriage ($p = 0.004$) and hormonal contraceptive use ($p = 0.002$) as significant predictors for high-risk HPV genotype. The findings highlight a unique regional virome dominated by HPV 18, 52, and 16, suggesting that regional viral epidemiology may differ significantly from global patterns, which are dominated by HPV 16 and 18. This prevalence of non-HPV 16 genotypes emphasizes the need for nonavalent vaccines and locally

tailored diagnostic tools.

While other factors, such as high parity and smoking, did not show genotype-specific correlations, they remain common clinical markers in this cohort. Furthermore, the 35.2% HPV-negative rate underscores the technical challenge ⁴⁹ of detection in advanced malignancies. These results provide a critical foundation for optimizing cervical cancer screening and prevention strategies specifically for the Indonesian population.

Acknowledgments

²⁷ The authors would like to thank the Center for Biomedical Research, Organization for Health, National Research and Innovation (BRIN), and the Biomolecular Laboratory, Faculty of Medicine, Universitas Trisakti, for supporting this research.

¹⁴ Funding

This research was supported by the RIIM LPDP Grant and BRIN (grant number 51/IV/KS/06/2022).

Conflict of interest

¹ The authors declare they have no competing interests.

Author contributions

Conceptualization: Raditya Wratsangka, Monica D. Hartanti

Formal analysis: Raditya Wratsangka, Monica D. Hartanti, Lily M. Surjadi

Investigation: Monica D. Hartanti, Gatot Purwoto, Miranti P. Aulia, Alvionita Kogoya

Methodology: Monica D. Hartanti, Gatot Purwoto, Miranti P. Aulia

Writing—original draft: Raditya Wratsangka, Monica D. Hartanti, Miranti P. Aulia

Writing—review & editing: All authors

³ Ethics approval and consent to participate

This study was approved by the ethical clearance committee of the Faculty of Medicine, Universitas Indonesia, with ethical clearance number KET-1262/UN2.F1/ETIK/PPM.00.02/2022. Written informed consent was obtained from participants before enrolment.

Consent for publication

This study involves fully anonymized, cross-sectional data and does not contain ⁸ any images or information that could identify a subject. Written informed consent was obtained from all participants for the publication of this study. All data have been anonymized to ensure participant

confidentiality.

15

Availability of data

All data generated or analyzed during this study are included in this published article.

References

1. ASCO. Challenges and Opportunities for Global Cervical Cancer Elimination: How Can We Build a Model for Other Cancers? *Am Soc Clin Oncol Educ Book*. 2025;45(3):e473702. doi: 10.1200/EDBK-25-473702.
2. World Health Organization, 2008. Cervical cancer, Fact Sheet. Updated December 2, 2025. Available from: <https://www.who.int/news-room/fact-sheets/detail/cervical-cancer> (Last accessed on 2025 Feb 23).
3. Wu J, Jin Q, Zhang Y, *et al*. Global burden of cervical cancer: current estimates, temporal trend and future projections based on the GLOBOCAN 2022. *J Natl Cancer Cent*. 2025;5(3):322-329. doi: 10.1016/j.jncc.2024.11.006
4. World Health Organization, 2024. WHO, UNFPA commend Indonesia's efforts to eliminate cervical cancer, urge streamlined vaccine strategy and enhanced screening. Posted November 15, 2024. Available from: <https://www.who.int/indonesia/news/detail/15-11-2024-who--unfpa-commend-indonesia-s-efforts-to-eliminate-cervical-cancer--urge-streamlined-vaccine-strategy-and-enhanced-screening#:~:text=The%20UN%20agencies%2C%20which%20supported,invasive%20cancer%20to%20be%20managed> (Last accessed on 2025 Dec 28).
5. World Health Organization, 2021. WHO recommends DNA testing as a first-choice screening method for cervical cancer prevention. Posted September 11, 2021. Available from: <https://www.who.int/europe/news/item/11-09-2021-who-recommends-dna-testing-as-a-first-choice-screening-method-for-cervical-cancer-prevention> (Last accessed on 2025 Jan 31).
6. Kusakabe M, Taguchi A, Sone K, Mori M, Osuga Y. Carcinogenesis and management of human papillomavirus-associated cervical cancer. *Int J Clin Oncol*. 2023;28(8):965-974. doi: 10.1007/s10147-023-02337-7
7. United Nations Children's Fund (UNICEF). 2023. Fast facts about HPV and cervical cancer. Available from: <https://www.unicef.org/stories/fast-facts-hpv-cervical-cancer> (Last accessed on 2025 Feb 15).
8. Calagna G, Maranto M, Paola C, *et al*. "Secondary prevention" against female HPV infection: literature review of the role of carrageenan. *Expert Rev Anti Infect Ther*. 2020;18(9):865-874. doi: 10.1080/14787210.2020.1770082
9. Capra G, Giovannelli L, Matranga D, *et al*. Potential impact of a nonavalent HPV vaccine on HPV related low- and high-grade cervical intraepithelial lesions: A referral hospital-based study in Sicily. *Hum Vaccin Immunother*. 2017;13(8):1839-1843. doi: 10.1080/21645515.2017.1319026
10. Ministry of Health, Republic of Indonesia. *Pedoman Nasional Pelayanan Kedokteran Tata Laksana Kanker Serviks* [National guidelines for treatment of cervical cancer]. 2018. HK.01.07/MENKES/349/2018. Available from: https://yankes.kemkes.go.id/unduh/fileunduh/1610415049_651524.pdf (Last accessed on 2025 Mar 15).
11. Ministry of Health, Republic of Indonesia, 2023. National Cervical Cancer Elimination (NCCE) Plan for Indonesia 2023-2030. Updated Dec 22, 2023. Available from: <https://kemkes.go.id/id/national-cervical-cancer-elimination-plan-for-indonesia-2023-2030> (Last accessed on 2025 Dec 28).
12. Robbers GML, Bennett LR, Spagnoletti BRM, Wilopo SA. Facilitators and barriers for the delivery and uptake of cervical cancer screening in Indonesia: A scoping review. *Glob Health Action*. 2021;14(1):1979280. doi: 10.1080/16549716.2021.1979280
13. Hutajulu SH, Prabandari YS, Bintoro BS, *et al*. Delays in the presentation and diagnosis of women with breast cancer in Yogyakarta, Indonesia: A retrospective observational study. *PLoS one*. 2022;17(1):e0262468. doi: 10.1371/journal.pone.0262468
14. Burd EM. Human papillomavirus and cervical cancer. *Clin Microbiol Rev*. 2023;16(1):1-17. doi: 10.1128/CMR.16.1.1-17.2003
15. Kombe Kombe AJ, Li B, Zahid A, *et al*. Epidemiology and Burden of Human Papillomavirus and Related Diseases, Molecular Pathogenesis, and Vaccine Evaluation. *Front Public Health*. 2020;8:552028. doi: 10.3389/fpubh.2020.552028
16. Bosco L, Serra N, Fasciana T, *et al*. Potential impact of a nonavalent anti HPV vaccine in Italian men with and without clinical manifestations. *Sci Rep*. 2021;11(1):4096. doi: 10.1038/s41598-021-83639-6
17. Hu Z, Zhu D, Wang W, *et al*. Genome-wide profiling of HPV integration in cervical cancer identifies clustered genomic hot spots and a potential microhomology-mediated integration mechanism. *Nat Genet*. 2015;47(2):158-163. doi: 10.1038/ng.3178
18. Hartanti MD, Rachmadhany EDA, Mustopa ITI, *et al*. High mobility group AT-hook 2 expression is elevated in cervical cancer. *Int J Pharm Res*. 2020;12:2332-2338. doi: 10.31838/ijpr/2020.12.04.322
19. Doorbar J. The human Papillomavirus twilight zone -

- Latency, immune control and subclinical infection. *Tumour Virus Res.* 2023;16:200268.
doi: 10.1016/j.tvr.2023.200268
20. Kogoya A, Hartanti MD. Correlations of Expression of KLF5, FHIT and DLG2 with Cervical Cancer. In: *Advances in Social Science, Education and Humanities Research*. Paris, France: Atlantis Press SARL; 2024:1302-1309.
doi: 10.2991/978-2-38476-273-6_132
21. World Health Organization. Global strategy to accelerate the elimination of cervical cancer as a public health problem. Updated Nov 17, 2020. Available from: <https://www.who.int/publications/i/item/9789240014107> (Last accessed on 2025 Apr 18).
22. Hartanti MD, Amtha R, Wratsangka R, et al. Operational Feasibility and Performance of mRPA Versus PCR for the Detection of HPV Types 16, 18, and 52 in Clinical Samples: An Exploratory Study. *Int J Womens Health.* 2025;17:955-964.
doi: 10.2147/IJWH.S496621
23. Burger EA, Kim JJ, Sy S, et al. Age of Acquiring Causal Human Papillomavirus (HPV) Infections: Leveraging Simulation Models to Explore the Natural History of HPV-induced Cervical Cancer. *Clin Infect Dis.* 2017;65(6):893-899.
doi: 10.1093/cid/cix475
24. Liu F, Chang L, Bai T, Liu X, Hu J. Association of human papillomavirus genotype distribution and cervical cytology: a cross-sectional study. *Epidemiol Infect.* 2021;149:e95.
doi: 10.1017/S0950268821000741
25. Tenjimbayashi Y, Onuki M, Hirose Y, et al. Whole-genome analysis of human papillomavirus genotypes 52 and 58 isolated from Japanese women with cervical intraepithelial neoplasia and invasive cervical cancer. *Infect Agent Cancer.* 2017;12:44.
doi: 10.1186/s13027-017-0155-4
26. Lin Y, Lin WY, Lin TW, et al. Trend of HPV Molecular Epidemiology in the Post-Vaccine Era: A 10-Year Study. *Viruses.* 2023;15(10):2015.
doi: 10.3390/v15102015
27. Gadducci A, Cosio S, Fruzzetti F. Estro-progestin Contraceptives and Risk of Cervical Cancer: A Debated Issue. *Anticancer Res.* 2020;40(11):5995-6002.
doi: 10.21873/anticancer.14620
28. Bergqvist L, Kalliala I, Aro K, et al. Distribution of HPV Genotypes Differs Depending on Behavioural Factors among Young Women. *Microorganisms.* 2021;9(4):750.
doi: 10.3390/microorganisms9040750
29. Lee SA, Baik S, Chung SH. Functional roles of female sex hormones and their nuclear receptors in cervical cancer. *Essays Biochem.* 2021;65(6):941-950.
doi: 10.1042/EBC20200175
30. James CD, Morgan IM, Bristol ML. The Relationship between Estrogen-Related Signaling and Human Papillomavirus Positive Cancers. *Pathogens (Basel, Switzerland).* 2020;9(5):403.
doi: 10.3390/pathogens9050403
31. Qi M, Naranjo AR, Duque AJ, et al. Evaluation of Pre-Analytical Variables for Human Papillomavirus Primary Screening from Self-Collected Vaginal Swabs. *J Mol Diagn.* 2024;26(6):487-497.
doi: 10.1016/j.jmoldx.2024.02.006
32. Dai W, Zeng X, Pantanowitz L, Zhao C. Human papillomavirus negative cervical cancers and precancerous lesions: prevalence, pathological and molecular features, and clinical implications. *Gynecol Obstet Clin Med.* 2025;5:e000160.
doi: 10.1136/gocm-2024-000160
33. Martinelli C, Ercoli A, Parisi S, et al. Molecular Mechanisms and Clinical Divergences in HPV-Positive Cervical vs. Oropharyngeal Cancers: A Critical Narrative Review. *BMC Medicine.* 2025;23(1):405.
doi: 10.1186/s12916-025-04247-z
34. Patel A, Galaal K, Burnley C, et al. Cervical cancer incidence in young women: a historical and geographic controlled UK regional population study. *Br J Cancer.* 2021;106(11):1753-1759.
doi: 10.1038/bjc.2012.148
35. Lau HY, Juang CM, Chen YJ, et al. Aggressive characteristics of cervical cancer in young women in Taiwan. *Int J Gynaecol Obstet.* 2009;107(3):220-223.
doi: 10.1016/j.ijgo.2009.07.029
36. Li S, Hu T, Lv W, et al. Changes in prevalence and clinical characteristics of cervical cancer in the People's Republic of China: A study of 10,012 cases from a nationwide working group. *Oncologist.* 2013;18(10):1101-1107.
doi: 10.1634/theoncologist.2013-0123
37. Itarat Y, Kietpeerakool C, Jampathong N, et al. Sexual behavior and infection with cervical human papillomavirus types 16 and 18. *Int J Women's Health.* 2019;11:489-494.
doi: 10.2147/IJWH.S218441
38. Luo D, Peng M, Wei X, et al. Prevalence of Human Papillomavirus and Genotype Distribution in Pregnant and Non-Pregnant Women in China. *Risk Manag Healthc Policy.* 2021;14:3147-3157.
doi: 10.2147/RMHP.S288838
39. Aguayo F, Munoz JP, Perez-Dominguez F, et al. High-Risk

Human Papillomavirus and Tobacco Smoke Interactions in Epithelial Carcinogenesis. *Cancers*. 2020;12(8):2201.

doi: 10.3390/cancers12082201

40. Ma K, Li S, Wu S, *et al.* Impact of smoking exposure on

human papillomavirus clearance among Chinese women: A follow-up propensity score matching study. *Tobacco Induced Diseases*. 2023;21:42.

doi: 10.18332/tid/161026

Risk factors and human papillomavirus genotypes: Investigating their interplay in cervical cancer

ORIGINALITY REPORT

16%
SIMILARITY INDEX

PRIMARY SOURCES

1	api-journal.accscience.com Internet	60 words — 1%
2	Guilherme S. Accorsi, Jeferson R. Zanon, Marcelo H. dos Santos, Ana Carla F. Ubinha et al. "Cervical cancer in young women: Does age impact survival in cervical cancer?", European Journal of Obstetrics & Gynecology and Reproductive Biology, 2025 Crossref	36 words — 1%
3	www.pagepress.org Internet	32 words — 1%
4	f1000research-files.f1000.com Internet	30 words — 1%
5	2020.ipvconference.org Internet	25 words — 1%
6	Kun Lin, Qiyang Hong, Ya Fu, Haijian Tu, Hua Lin, Jiexiang Huang, Yajing Hu, Minjun Huang, Mingqiao Chen. "Cervical HPV infection and related diseases among 149,559 women in Fujian: an epidemiological study from 2018 to 2023", Frontiers in Microbiology, 2024 Crossref	25 words — 1%
7	Joana Kyei Dompim, Abigail Kusi Amponsah, Edward Appiah Boateng, Alberta Dotse Makafui et al. "Pregnant women's beliefs and perceptions of cervical cancer screening: A descriptive phenomenological study", International Journal of Africa Nursing Sciences, 2025 Crossref	24 words — < 1%
8	Yu Wang, Chao Qi. "Comparative Diagnostic Performance of DNA Ploidy Analysis and p16/Ki67 Dual-Staining with Cytology in hrHPV-Defined Subgroups: A Single-Center Evaluation", Cancer Management and Research, 2026 Crossref	23 words — < 1%

9	etheses.whiterose.ac.uk Internet	20 words — < 1%
10	gupea.ub.gu.se Internet	19 words — < 1%
11	Ahmed A.Amer, Hashem Altabbaa, Mohamed Sh.Ramadan, Hamza A. Abdul-Hafez et al. "A multinational cross-sectional study on human papillomavirus and cervical cancer knowledge, vaccination attitudes, and risk factors in the Middle East", Scientific Reports, 2025 Crossref	18 words — < 1%
12	Bijaya Chandra Acharya, Hemnath Subedi, Sarita Rana Gurung, Greta Pandey, Asmita Rana, Hari Prasad Upadhyay. "Prevalence and Associated Factors of HPV DNA Test Status of Women Attending a Tertiary Cancer Hospital of Central Nepal", Journal of College of Medical Sciences-Nepal, 2025 Crossref	18 words — < 1%
13	www.journalcls.org Internet	18 words — < 1%
14	Ari Hardianto, Zahra Silmi Muscifa, Wahyu Widayat, Muhammad Yusuf, Toto Subroto. "The Effect of Ethanol on Lipid Nanoparticle Stabilization from a Molecular Dynamics Simulation Perspective", Molecules, 2023 Crossref	17 words — < 1%
15	ir.ymlib.yonsei.ac.kr Internet	17 words — < 1%
16	www.spandidos-publications.com Internet	16 words — < 1%
17	coek.info Internet	15 words — < 1%
18	www.researchandmarkets.com Internet	14 words — < 1%
19	Canio Martinelli, Alfredo Ercoli, Silvana Parisi, Giuseppe Iatì et al. "Molecular Mechanisms and Clinical Divergences in HPV-Positive Cervical vs. Oropharyngeal Cancers: A Critical Narrative Review", BMC Medicine, 2025 Crossref	13 words — < 1%

20	Human Papillomavirus (HPV)-Associated Oropharyngeal Cancer, 2015. Crossref	13 words — < 1%
21	Théodora Mahoukèdè Zohoncon, Shoukrat Ohuwa Toyin Bello, Prosper Bado, Rogomenoma Alice Ouédraogo et al. "Detection of Human Papillomavirus DNA and E6/E7 mRNA from HPV Genotypes 16, 18, 31 and 33 in Histologically Confirmed Cases of Cervical Cancer and Precancerous Lesions in Burkina Faso", American Journal of Molecular Biology, 2023 Crossref	13 words — < 1%
22	Tuti Pahria, Hartiah Haroen, Hana Rizmadewi Agustina, Citra Windani Mambang Sari et al. "Women's experiences toward cervical cancer care at the initiation of care: a qualitative study closing the gap policy and implementation in Indonesia", BMC Women's Health, 2026 Crossref	13 words — < 1%
23	Ebubechukwu H. Nkadi, Danielle L. Schaal, Rona S. Scott. "Epstein-Barr Virus and Human Papillomavirus Co-Infections in Epithelial Carcinomas", Journal of Medical Virology, 2026 Crossref	12 words — < 1%
24	health.infolabmed.com Internet	12 words — < 1%
25	open.uct.ac.za Internet	11 words — < 1%
26	www2.mdpi.com Internet	11 words — < 1%
27	Marianna Martinelli, Sadaf Sakina Hassan, Emel Yilmaz, Camilla Lagheden et al. "Assessing sample adequacy and clinical performance of self-collected and clinician-collected HPV specimens using internal control Ct values", Journal of Clinical Virology, 2025 Crossref	10 words — < 1%
28	www.researchsquare.com Internet	10 words — < 1%
29	www.sansureglobal.com Internet	10 words — < 1%

-
- 30 Jie Wu, Qianyun Jin, Yunmeng Zhang, Yuting Ji et al. "Global burden of cervical cancer: current estimates, temporal trend and future projections based on the GLOBOCAN 2022", Journal of the National Cancer Center, 2025
Crossref 9 words — < 1%
-
- 31 Karin Butz. "Activation of human papillomavirus type 18 E6-E7 oncogene expression by transcription factor Sp1", Nucleic Acids Research, 1992
Crossref 9 words — < 1%
-
- 32 Lac Thi Kim Ngan, Huynh Van Ba, Dao Hoang Thien Kim, Dang Tran Dang Khoa et al. "Prevalence and distribution of human papillomavirus in genital warts in Can Tho City, Vietnam: a cross-sectional study", Infectious Agents and Cancer, 2025
Crossref 9 words — < 1%
-
- 33 Mebrahtu Teweldemedhin Shfare, Hayelom Kahsay, Gebrecherkos Teame Gebrehiwot, Getachew Kahsu Abay et al. "High-risk human papillomavirus positivity and associated factors among women living with HIV in public health facilities of Mekelle, Northern Ethiopia", Infectious Agents and Cancer, 2026
Crossref 9 words — < 1%
-
- 34 Shanshan Yang, Tianbo Liu, Haiyan Cheng, Zhao Wang, Yue Feng, Jiazhao Yan, Sijia Liu, Yunyan Zhang. "Decreased Expression of Retinoblastoma Protein-Interacting Zinc-Finger Gene 1 Is Correlated With Poor Survival and Aggressiveness of Cervical Cancer Patients", Frontiers in Oncology, 2019
Crossref 9 words — < 1%
-
- 35 bmjopen.bmj.com
Internet 9 words — < 1%
-
- 36 dspace.library.uu.nl
Internet 9 words — < 1%
-
- 37 synapse.koreamed.org
Internet 9 words — < 1%
-
- 38 www.researchprotocols.org
Internet 9 words — < 1%

-
- 39 Amitabha Ray. "Human Papillomavirus and Other Relevant Issues in Cervical Cancer Pathogenesis", International Journal of Molecular Sciences, 2025 8 words — < 1%
[Crossref](#)
-
- 40 Daiwen Liang, Qi Zhang, Wenyu Li, Youkun Lin. "Analysis of urinary tobacco-specific nitrosamine 4- (methylnitrosamino)1-(3-pyridyl)-1-butanol (NNAL) and HPV infection in American women: National health and nutrition examination survey", PLOS ONE, 2024 8 words — < 1%
[Crossref](#)
-
- 41 Di Gao, Xueyin Wang, Juan Juan, Hui Bi, Jiayue Li, Yupeng Shi, Taoye Xu, Yue Wang, Gengli Zhao, Xiaosong Zhang. "Prevalence and genotype distribution of cervical human papillomaviruses infection across cytology: a nationwide, cross-sectional study of over 1 million Chinese women from 2020 to 2024", Journal of Infection, 2026 8 words — < 1%
[Crossref](#)
-
- 42 Giovanni Cangelosi, Francesco Sacchini, Stefano Mancin, Fabio Petrelli et al. "Papillomavirus Vaccination Programs and Knowledge Gaps as Barriers to Implementation: A Systematic Review", Vaccines, 2025 8 words — < 1%
[Crossref](#)
-
- 43 Giuseppina Palena, Irene Stilo, Michele Sorrentino, Claudio Fiorilla, Raffaele Palladino. "Organizational barriers in HPV vaccination uptake: A cross-sectional study among health sciences students", PLOS One, 2025 8 words — < 1%
[Crossref](#)
-
- 44 I Ketut Agus Somia, Made Lady Adelaida Purwanta, Ni Wayan Winarti, Ida Bagus Nyoman Putra Dwija et al. "Prevalence and risk factors of high-risk HPV and cervical abnormalities in HIV-positive women in Bali, Indonesia", BMC Infectious Diseases, 2025 8 words — < 1%
[Crossref](#)
-
- 45 Karol Magiera, Jakub Rybak, Barbara Magiera, Alicja Grabarczyk, Martyna Grabowska-Szczurek. "Cervical cancer – risk factors and new opportunities for prevention in Poland", Environmental Medicine, 2024 8 words — < 1%
[Crossref](#)
-
- 46 Mondraya F. Howard, Ria C. Fyffe-Freil, Christina C. Pierre, Henrietta O. Maku et al. "Self-collection for primary HPV screening using dry swabs: a review of clinical performance, laboratory considerations, and patient preferences", Journal of Clinical Microbiology, 2025 8 words — < 1%
[Crossref](#)

47	Nanqiu Peng, Yiyi Zhang, Li He, Li Xie. "Vaginal microecological diversity and Ureaplasma urealyticum co-infection in high-risk HPV infection: a cross-sectional study", Frontiers in Cellular and Infection Microbiology, 2026 Crossref	8 words — < 1%
48	Valeria Escobar, Alejandro Catalán, Verónica Callejas, Roylester Araya et al. "Human papillomavirus (HPV) genotypes extended prevalence in the female population from a city in Northern Chile", BMC Women's Health, 2025 Crossref	8 words — < 1%
49	Yana Merdzhanova-Gargova, Magdalena Ivanova, Angelina Mollova-Kysebekirova, Anna Mihaylova et al. "Advancements in Cervical Cancer Screening: Enhancing HPV Testing and Triage Strategies for Improved Risk Assessment", Biomedicines, 2025 Crossref	8 words — < 1%
50	belitungraya.org Internet	8 words — < 1%
51	cmr.asm.org Internet	8 words — < 1%
52	nigerianmedjournal.org Internet	8 words — < 1%
53	portalcris.lsmuni.lt Internet	8 words — < 1%
54	www.mp.pl Internet	8 words — < 1%
55	www.preprints.org Internet	8 words — < 1%
56	www.scielo.br Internet	8 words — < 1%
57	Sondos Al Sad, Labiqah Iftikhar, Masa Chamout. "Revisiting HPV vaccination post-COVID: geopolitical, sociocultural, and ethical disparities in global health", International Journal for Equity in Health, 2025 Crossref	7 words — < 1%

58 Medha Karnik, Preethi G. Anantharaju, Arati Sharma, Olga Sukocheva, Edmund Tse, SubbaRao V. Madhunapantula. "Aurora kinase B as a therapeutic target in HPV-induced cervical cancer: mechanisms and future perspectives", Frontiers in Oncology, 2026 6 words — < 1%
[Crossref](#)

59 Rehana V. Hewavisenti, Joshua Arena, Chantelle L. Ahlenstiel, Sarah C. Sasson. "Human papillomavirus in the setting of immunodeficiency: Pathogenesis and the emergence of next-generation therapies to reduce the high associated cancer risk", Frontiers in Immunology, 2023 6 words — < 1%
[Crossref](#)

60 Yujia Kong, Liju Zong, Junjun Yang, Ming Wu, Yang Xiang. "<p>Cervical cancer in women aged 25 years or younger: a retrospective study</p>", Cancer Management and Research, 2019 6 words — < 1%
[Crossref](#)

61 Kim, K.H.. "Development and evaluation of a highly sensitive human papillomavirus genotyping DNA chip", Gynecologic Oncology, 200601 5 words — < 1%
[Crossref](#)

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES OFF
EXCLUDE MATCHES OFF