

Journal of Water and Land Development

COUNTRY

Poland


-  Universities and research institutions in Poland
-  Media Ranking in Poland

SUBJECT AREA AND CATEGORY

- Agricultural and Biological Sciences
 - Agricultural and Biological Sciences (miscellaneous)
- Environmental Science
 - Environmental Engineering
 - Water Science and Technology
- Social Sciences
 - Development
 - Geography, Planning and Development

PUBLISHER

Polish Academy of Sciences

-  Polish Academy of Sciences in Scimago Institutions Rankings

- Seen this ad multiple times
- Ad was inappropriate
- Not interested in this ad

H-INDEX

21

PUBLICATION TYPE

Journals

ISSN




14297426, 20834535

COVERAGE

2006-2022

INFORMATION

- [Homepage](#)
- [How to publish in this journal](#)
- journal@itp.edu.pl



Journal of Water and Land Development

Q2 Agricultural and Biological Sciences (miscellaneous) best quartile

SJR 2022
0.28

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com" data-bbox="700 865 860 885">
```

Cover



Daftar Isi

CONTENT

Journal of Water and Land Development | 2021 | No 49

1 **Monitoring of horizontal displacements and changes of the riverine area of the Dniester River**

[Volodymyr Shevchuk](#) , [Khrystyna Burshtynska](#) , [Iryna Korolik](#) , [Maksym Halochkin](#)

[Journal of Water and Land Development | 2021 | No 49 | 1-15](#)

Keywords: [channel processes](#) [Dniester River](#) [monitoring](#) [riverbed displacements](#) [riverine areas](#) [satellite images](#) [super-vised classification](#) [topographic maps](#)

[Abstract](#) [Download PDF](#) [Download RIS](#) [Download Bibtex](#)
[Authors and Affiliations](#)

2 **Waterlogging tolerance: A review on regulative morpho-physiological homeostasis of crop plants**

[Shubhangani Sharma](#) , [Jyotshana Sharma](#) , [Vineet Soni](#) , [Hazem M. Kalaji](#) , [Nabil I. Elsheery](#)

[Journal of Water and Land Development | 2021 | No 49 | 16-28](#)

Keywords: [antioxidants](#) [cellular metabolites](#) [climate change](#) [photosynthesis](#) [waterlogging](#)

[Abstract](#) [Download PDF](#) [Download RIS](#) [Download Bibtex](#)

3 **Comparative assessment of Al₂O₃-modified biomasses from agricultural residues for nickel and cadmium removal**

[Adriana Herrera-Barros](#) , [Candelaria Tejada-Tovar](#) , [Angel D. Gonzalez-Delgado](#)

[Journal of Water and Land Development | 2021 | No 49 | 29-34](#)

Keywords: [adsorption](#) [agricultural residues](#) [alumina nanoparticles](#) [cellulose](#) [heavy metal ions](#)

[Abstract](#) [Download PDF](#) [Download RIS](#) [Download Bibtex](#)
[Authors and Affiliations](#)

4 **Assessment of groundwater vulnerability using the DRASTIC model: A case study of quaternary catchment A21C, Limpopo River Basin, South Africa**

[Simeneh S. Moges](#) , [Megersa O. Dinka](#)

[Journal of Water and Land Development | 2021 | No 49 | 35-46](#)

Keywords: [A21C Quaternary catchment](#) [DRASTIC model](#) [groundwater vulnerability](#) [sensitivity analysis](#) [South Africa](#)

[Abstract](#) [Download PDF](#) [Download RIS](#) [Download Bibtex](#)
[Authors and Affiliations](#)

5 **Soil erosion control in immature oil palm plantation**

[Halus Satriawan](#) , [Zahrul Fuady](#) , [Rini Fitri](#)

[Journal of Water and Land Development | 2021 | No 49 | 47-54](#)

Keywords: [cover crops](#) [erosion](#) [nutrient leaching](#) [oil palm plantation](#) [sediment trap](#) [surface runoff](#) [terraces](#)

[Abstract](#) [Download PDF](#) [Download RIS](#) [Download Bibtex](#)
[Authors and Affiliations](#)

6 **Effect of magnetic fields and fertilizers on grass and onion growth on technogenic soils**

[Maria Vasilyeva](#) , [Stanislav Kovshov](#) , [Johnny Zambrano](#) , [Maxim Zhemchuzhnikov](#)

[Journal of Water and Land Development | 2021 | No 49 | 55-62](#)

Keywords: [alternating magnetic field](#) [biochar](#) [bioefficiency](#) [biohumus](#) [constant magnetic field](#) [fertilization](#) [physiological parameters of plants](#) [technogenic soil](#)

[Abstract](#) [Download PDF](#) [Download RIS](#) [Download Bibtex](#)
[Authors and Affiliations](#)

7 **An experimental sedimentation tank for enhancing the settling of solid particles**

[Dina A. Yaseen](#) , [Saad Abu-Alhail](#) , [Rusul N. Mohammed](#)

[Journal of Water and Land Development | 2021 | No 49 | 63-73](#)

Keywords: [baffle](#) [clay](#) [secondary settling basin](#) [solid particles](#) [treatment plant](#) [turbid water](#)

[Abstract](#) [Download PDF](#) [Download RIS](#) [Download Bibtex](#)
[Authors and Affiliations](#)

8 **Prospects for the use of microwave energy in grain crop seeding**

[Fedor A. Kipriyanov](#) , [Petr A. Savinykh](#) , [Alexey Yu. Isupov](#) , [Yulia A. Plotnikova](#) , [Natalia A. Medvedeva](#) , [Svetlana V. Belozerova](#)

[Journal of Water and Land Development | 2021 | No 49 | 74-78](#)

Keywords: [disinfection](#) [microwave drying](#) [microwave energy](#) [microwave use on plants](#) [pre-sowing treatment](#)

[Abstract](#) [Download PDF](#) [Download RIS](#) [Download Bibtex](#)
[Authors and Affiliations](#)

9 **The minimization of impact of oil pollution on soils in the area of railways using glauconite**

[Oksana Chayka](#) , [Igor Petrushka](#) , [Maria Ruda](#) , [Nadiya Paranyak](#) , [Olena Matskiv](#)

[Journal of Water and Land Development | 2021 | No 49 | 79-84](#)

Keywords: [ecological safety](#) [glauconite](#) [oil products](#) [pollution](#) [railway transport](#) [soil contamination](#)

[Abstract](#) [Download PDF](#) [Download RIS](#) [Download Bibtex](#)
[Authors and Affiliations](#)

10 **A novel method and device for plastic mulch retriever**

[Kanat M. Khazimov](#) , [Adilkhan K. Niyazbayev](#) , [Zhanbota S. Shekerbekova](#) , [Aigul A. Urymbayeva](#) , [Gulzhanat A. Mukanova](#) , [Tursunkul A. Bazarbayeva](#) , [Vladimir F. Nekrashevich](#) , [Marat Zh. Khazimov](#)

[Journal of Water and Land Development | 2021 | No 49 | 85-94](#)

Keywords: [compaction](#) [microclimatic conditions](#) [mulch retriever](#) [plastic mulch](#) [polymer degradation](#)

[Abstract](#) [Download PDF](#) [Download RIS](#) [Download Bibtex](#)
[Authors and Affiliations](#)

11 **Use of fly larvae *Hermetia illucens* in poultry feeding: A review paper**

[Svetlana V. Sverguzova](#) , [Ildar H. Shaikhiev](#) , [Zhanna A. Saponova](#) , [Ekaterina V. Fomina](#) , [Yulia L. Makridina](#)

[Journal of Water and Land Development | 2021 | No 49 | 95-103](#)

Keywords: [feed](#) [Hermetia illucens](#) [insect feed](#) [poultry](#) [protein flour](#)





[Abstract](#) [Download PDF](#) [Download RIS](#) [Download Bibtex](#)
[Authors and Affiliations](#)


12 Features of the surface water oxygen regime in the Ukrainian Polesie Region

Ella O. Aristarkhova , Tetiana P. Fedoniuk , Ludmila D. Romanchuk , Sergii V. Latushynskyi , Iryna V. Kot

Journal of Water and Land Development | 2021 | No 49 | 104-110

Keywords: correlation analysis dissolved oxygen fish oxygen regime pH seasonal fluctuations surface water temperature toxicity

 Abstract  Download PDF  Download RIS  Download Bibtex





 Authors and Affiliations


13 Spatial and temporal assessment of surface water quality using water quality index. The Saguling Reservoir, Indonesia

Mariana Marselina , Anwar Sabar , Nurul Fahimah

Journal of Water and Land Development | 2021 | No 49 | 111-120

Keywords: inverse distance weight spatial and temporal assessment surface water water quality index

 Abstract  Download PDF  Download RIS  Download Bibtex





 Authors and Affiliations


14 Impact of raw and purified wastewater reuse from the Dar El-Gueddari treatment plant on the soil

Smail Njimat , Fouad Elfettahi , Hajar Griou , Mohammed Y. El Brouzi , Mohammed Aboulouafa , Said Ibn Ahmed

Journal of Water and Land Development | 2021 | No 49 | 121-128

Keywords: dilutions electrical conductivity infiltration organic matter pH treatment plant wastewater irrigation

 Abstract  Download PDF  Download RIS  Download Bibtex





 Authors and Affiliations


15 Field and modelling study for deficit irrigation strategy on roots volume and water productivity of wheat

Ramadan E. Abdelraouf , Mohamed A. El-Shawadfy , Osama M. Dewedar , Mahmoud Hozayn

Journal of Water and Land Development | 2021 | No 49 | 129-138

Keywords: irrigation roots volume SALTMED model soil moisture water productivity wheat

 Abstract  Download PDF  Download RIS  Download Bibtex





 Authors and Affiliations


16 Response of potato biomass and tuber yield under future climate change scenarios in Egypt

Osama Dewedar , Finn Plauborg , Ahmed El-Shafie , Abdelbaset Marwa

Journal of Water and Land Development | 2021 | No 49 | 139-150

Keywords: AquaCrop model biomass climate change CMIP5 scenarios potato yield

 Abstract  Download PDF  Download RIS  Download Bibtex





 Authors and Affiliations


17 Phosphoric regime of light chestnut soil and sugar beet yield with long-term use of phosphorous fertilizers

Balnur Alimbekova , Rakhimzhan Yeleshev , Zhenisgul Bakenova , Aigerim Shibikeyeva , Marzhan Balkozha

Journal of Water and Land Development | 2021 | No 49 | 151-155

Keywords: ammonia nitrogen chestnut soil fertilization gross phosphorus mineral nitrogen mineral phosphorus mobile phosphorus organic phosphorus permanent sugar beet crop

 Abstract  Download PDF  Download RIS  Download Bibtex

 Authors and Affiliations

18 Development of biogas and biorafinery systems in Polish rural communities

Grzegorz Wałowski

Journal of Water and Land Development | 2021 | No 49 | 156-168

Keywords: agricultural biogas biorafination fermentation process pig slurry

[Abstract](#) [Download PDF](#) [Download RIS](#) [Download Bibtex](#)
[Authors and Affiliations](#)

19 Grape production assessment using surface and subsurface drip irrigation methods

Sharad J. Kadbhane , Vivek L. Manekar

Journal of Water and Land Development | 2021 | No 49 | 169-178

Keywords: crop water requirement irrigation with plastic bottles organic mulch plastic mulch stone column sustainable irrigation water productivity

[Abstract](#) [Download PDF](#) [Download RIS](#) [Download Bibtex](#)
[Authors and Affiliations](#)

20 Hydrochemical characteristics and water quality assessment of Lake Dayet Erroumi - Khemisset, Morocco

Mohamed El Qryefy , Jamila Ouardi , Mohamed Najy , Driss Belghyti , Khadija El Kharrim

Journal of Water and Land Development | 2021 | No 49 | 179-187

Keywords: Dayet Erroumi hydrochemical characteristics Morocco salinity water quality

[Abstract](#) [Download PDF](#) [Download RIS](#) [Download Bibtex](#)
[Authors and Affiliations](#)

21 The mechanism of phosphate bacteria in increasing the solubility of phosphorus in Indonesian Andisols

Tamad , Azwar Maas , Eko Hanudin , Jaka Widada

Journal of Water and Land Development | 2021 | No 49 | 188-194

Keywords: adsorption Andisols mineralization phosphate bacteria solubilisation

[Abstract](#) [Download PDF](#) [Download RIS](#) [Download Bibtex](#)
[Authors and Affiliations](#)

22 Estimation and management of the future evolution of groundwater water demand in the Taoura region, North-East Algeria

Farid Rouaibia , Lamine Sayad , Badra Attoui

Journal of Water and Land Development | 2021 | No 49 | 195-203

Keywords: Algeria annual average precipitation groundwater socioeconomic development sustainable resource man-agement Taoura WEAP model

[Abstract](#) [Download PDF](#) [Download RIS](#) [Download Bibtex](#)
[Authors and Affiliations](#)

23 Community enhancement of the environmental quality of riverbank settlements: A case study of Tridi Kampong, Indonesia

Ida A.M. Wahyuni , I M. Weni , Tommy Hariyanto , Laksni Sedyowati

Journal of Water and Land Development | 2021 | No 49 | 204-212








24 The influence of the mixture of vermicompost and sulphur-perlite-containing waste on the yield and the quality of crops

Moldir O. Baikhamurova , Gaukhar A. Sainova , Amankul D. Akbasova , Gulshat D. Anarbekova , Mehmet A. Ozler

Journal of Water and Land Development | 2021 | No 49 | 213-218

Keywords: crops quality root vegetables sierozem soil soil fertility substrate sulphur perlite-containing waste vermicom-post yielding

 Abstract  Download PDF  Download RIS  Download Bibtex





 Authors and Affiliations


25 Assessing domestic factors determining water consumption in a semi-arid area (Sedrata City) using artificial neural networks and principal component analysis

Menal Zeroual , Azzedine Hani , Amir Boustila

Journal of Water and Land Development | 2021 | No 49 | 219-228

Keywords: artificial neural networks domestic water use determinants household water consumption principal component analysis semi-arid area

 Abstract  Download PDF  Download RIS  Download Bibtex

 Authors and Affiliations

26 Management of expenses for fertilizers and chemical plant protection products in the accounting and control system of agrarian formations

Zhanna V. Degaltseva , Victor V. Govdya , Konstantin A. Velichko

Journal of Water and Land Development | 2021 | No 49 | 229-234





Keywords: accounting and control system agrarian formation cost of fertilizers efficiency information technology


27 Characterization of Algerian Sahara groundwater for irrigation and water supply: Adrar region study case

Ali Bendida , Mohammed Amin Kendouci , Abdellatif El-Bari Tidjani

Journal of Water and Land Development | 2021 | No 49 | 235-243

Keywords: Adrar area Algerian Sahara drinking water groundwater hydrochemical irrigation water Piper diagram

 Abstract  Download PDF  Download RIS  Download Bibtex





 Authors and Affiliations


28 Flow characteristics after installation of floating bridge in open channel

Mohamed M. Ibrahim , Mahmoud A.R. Eltoukhy , Adnan D. Ghanim

Journal of Water and Land Development | 2021 | No 49 | 244-251

Keywords: floating bridges flow characteristics flume head loss velocity

 Abstract  Download PDF  Download RIS  Download Bibtex

 Authors and Affiliations

29 Kinetic study of oil-containing wastewater treatment with the use of natural sorbents

Saltanat T. Tleuova , Banu A. Userbayeva , Alibek S. Tleuov , Marina M. Yeskendirova , Raissa R. Yakubova , Kulyash Z. Kerimbayeva

Journal of Water and Land Development | 2021 | No 49 | 252-258

30 Floodway design affected by land use changes in an urbanized area

[Agus Suharyanto](#) , [Yatnanta P. Devia](#) , [Indradi Wijatmiko](#)

[Journal of Water and Land Development](#) | 2021 | No 49 | 259-266

Keywords: channel dimensions channel normalization floodway land cover changes land use runoff coefficient

[Abstract](#) [Download PDF](#) [Download RIS](#) [Download Bibtex](#)
[Authors and Affiliations](#)

31 Hydraulic gradient prediction for solid-water mixture flow through horizontal pipelines of different diameters and roughness and solids concentrations

[Faisal A. Odra](#)

[Journal of Water and Land Development](#) | 2021 | No 49 | 267-272

Keywords: deposit limit velocity hydraulic gradient hydraulic losses slurry flow solid concentrations suspended solids wastewater

[Abstract](#) [Download PDF](#) [Download RIS](#) [Download Bibtex](#)
[Authors and Affiliations](#)

32 Geospatial potential and environmental changes: Case study of complex nature of the Altai Geopark

[Nina A. Kocheeva](#) , [Natalia A. Iurkova](#) , [Mariia G. Sukhova](#) , [Aleksndr Germanovich Redkin](#) , [Olga Valerevna Zhuravleva](#)

[Journal of Water and Land Development](#) | 2021 | No 49 | 273-281

Keywords: Altai environmental changes geological objects geopark geotourism rural development

[Abstract](#) [Download PDF](#) [Download RIS](#) [Download Bibtex](#)
[Authors and Affiliations](#)

Dewan Redaksi

Editorial Board:

Jan Franklin ADAMOWSKI – McGill University, Quebec, Canada

Tariq AFTAB – Aligarh Muslim University, Aligarh, India [id 0000-0002-5927-719X](#)

Suleyman I. ALLAKHVERDIEV – Azerbaijan State University, Baku, Azerbaijan

Ozan ARTUN – Çukurova University, Adana, Turkey [id 0000-0002-6122-2729](#)

Habib R. ATHAR – Institute of Pure and Applied Biology, Zakariya University, Multan, Pakistan [id 0000-0002-8733-3865](#)

Atilgan ATILGAN – Alanya Alaaddin Keykubat University, Alanya, Turkey

Okke BATELAAN – Flinders University, Adelaide, Australia [id 0000-0003-1443-6385](#)

Marian BRESTIC – Slovak University of Agriculture, Nitra, Slovakia

Filippo BUSSOTTI – University of Firenze, Firenze, Italy

Shiguo CHEN – Nanjing Agricultural University, Nanjing, China

Ralf DANNOWSKI – Leibniz-Zentrum für Agrarlandschaftsforschung, Brandenburg, Germany [id 0000-0002-9331-672X](#)

Nabil ELSHERY – Tanta University, Tanta, Egypt [id 0000-0001-9542-1913](#)

Domenica FARCI – Università degli studi di Cagliari, Cagliari, Italy

GOVINDJEE – University of Illinois at Urbana-Champaign, Urbana, USA [id 0000-0003-3774-0638](#)

Francisco GUERRERO – Universidad de Jaén, Jaén, Spain

Salim HEDDAM – Université 20 août 1955-Skikda, Skikda, Algeria [id 0000-0002-8055-8463](#)

Tomasz HORACZEK – Institute of Technology and Life Sciences, Falenty, Poland [id 0000-0001-5534-7544](#)

Miroslava KAČÁNIOVÁ – Slovak University of Agriculture, Nitra, Slovakia

Jan KAZAK – Wrocław University of Environmental and Life Sciences, Poland

Sungwon KIM – Dongyang University, Yeongju, Republic of Korea [id 0000-0002-9371-8884](#)

Ozgur KISI – University of Applied Sciences, Lübeck, Germany [id 0000-0001-7847-5872](#)

Peter KOVALENKO – Institute of Water Problems and Melioration of the National Academy of Agrarian Sciences, Ukraine

Nour-Eddine LAFTOUHI – Cadi Ayyad University, Marrakesh, Morocco

Ramin LOTFI – Dryland Agricultural Research Institute, AREEO, Maragheh, Iran

Grażyna MASTALERCZUK – Warsaw University of Life Sciences – SGGW, Warsaw, Poland

Maria MRÓWCZYŃSKA – University of Zielona Góra, Poland [id 0000-0002-4762-3999](#)

Samar OMAR – Tanta University, Tanta, Egypt

Dario PIANO – Università degli studi di Cagliari, Cagliari, Italy

Karol PLESIŃSKI – Agricultural University in Cracow, Cracow, Poland

Majeti Narasimha Vara PRASAD – University of Hyderabad, Hyderabad, India

Mahendra RAI – SGB Amravati University, Maharashtra, India

Marcin RAPACZ – University of Agriculture, Cracow, Poland

Manzer H. SIDDIQUI – King Saud University, Riyadh, Saudi Arabia

Edyta SIERKA – University of Silesia, Katowice, Poland

Ewa SOBIESZCZUK-NOWICKA – Adam Mickiewicz University in Poznań, Poznań, Poland [id 0000-0002-6603-0582](#)

Kazimierz STRZAŁKA – Jagiellonian University, Cracow, Poland

Wayan SUPARTA – Institute Technology National Yogyakarta, Yogyakarta, Indonesia

Sawsan TAWKAZ – Consultative Group on International Agricultural Research, Montpellier, France

Adam TAŃSKI – West Pomeranian University of Technology, Szczecin, Poland


Renata TOBIASZ-SALACH – Rzeszow University, Rzeszow, Poland

Katarzyna TURNAU – Jagiellonian University, Cracow, Poland

Martin J. WASSEN – Utrecht University, Utrecht, Netherlands

Marek ZIVCAK – Slovak University of Agriculture, Nitra, Slovak Republic

Journal of Water and Land Development



About the Journal

Aims and scope

Editorial Board

Editors

Contact

Open Access Policy

Abstracting & Indexing

Editor-in-Chief

Professor Dr. Hab. Hazem M. Kalaji  0000-0001-2345-6789



Department of Plant Physiology, Faculty of Agriculture and Biology, Warsaw University of Life Sciences SGGW, Warsaw, Poland
Institute of Technology and Life Sciences - National Research Institute, Falenty, Al. Hraboska 3, 05-090 Raszyn, Poland
Interests: fluorescence sensors; chlorophyll fluorescence analysis; photochemistry of photosynthesis; plant stress; physiology of plants and algae; plant talk and machine learning

Managing Editor

Dr Adam Brysiewicz

English Language Editor

Charlotte Aldred (English Native Speaker)

Statistical Editors

dr hab. Dariusz Gozdowski

dr inż. Leszek Sieczko

Subscribe

 **RSS**

Associate Editors

Suzana Gotovac Atlagić (*environmental chemistry, nanotechnology*)

Barbara Borawska-Jarmutowicz (*grassland, agronomy*)

Klaudia Borowiak (*air pollution, landscape evaluation*)

Piotr Bugajski (*wastewater, wetlands*)

Jarosław Chormański (*water management, environmental science, water science*)

Przemysław Czerniejewski (*hydrobiology, ecology*)

Piotr Dąbrowski (*environmental science, green infrastructure*)

Tomasz Gnatowski (*soil water management, irrigation*)

Tomasz Kowalczyk (*water resources, GIS, irrigation, water management*)

Mateusz Malinowski (*environmental engineering, GIS, land degradation*)

Federico Marrone (*hydrobiology, phylogeography*)

Gabriel Minea (*hydrology, water ecosystems*)

Amarena Narayan Misra (*biotechnology, molecular biology*)

Agnieszka Operacz (*water science, soil science, hydrology, hydrogeology*)

Jos Puthur (*agricultural plant science, molecular biology*)

Piotr Stępień (*plant physiology, biochemistry*)

Osama H. Sayed (*ecology, water stress*)

Avinash Chandra Pandey (*agricultural plant science, genetics*)

Mariusz Sojka (*water monitoring, modelling and management*)

Waldemar Spychalski (*soil science, soil protection*)

Irena Suwara (*agronomy, crop production*)

Szymon Szewrański (*landscape architecture, spatial economy*)

Hayat Touchane (*water productivity, climate change*)

Grzegorz Wałowski (*agricultural biogas*)

Krytyna Żuk – Gołaszewska (*agriculture, plant physiology*)



Received 10.04.2020
Reviewed 26.05.2020
Accepted 29.09.2020

Soil erosion control in immature oil palm plantation

Halus SATRIAWAN¹  , Zahrul FUADY¹ , Rini FITRI² 

¹ Almuslim University, Faculty of Agriculture, Department of Agrotechnology, Bireuen, Aceh, 24261, Indonesia

² University of Trisakti, Faculty of Landscape Architecture and Environmental Technology, Department of Landscape Architecture, Jakarta, Indonesia

For citation: Satriawan H., Fuady Z., Fitri R. 2021. Soil erosion control in immature oil palm plantation. *Journal of Water and Land Development*. No. 49 (IV–VI) p. 47–54. DOI 10.24425/jwld.2021.137095.

Abstract

The objectives of developing oil palm plantations should be feasible economically and without causing massive erosion. This research proposes soil and water conservation strategies that are ideal and optimal for oil palm cultivation depending on land capability class. The conservation test for plants was performed according to land capability classes on a plot measuring 22 m × 4 m. Runoff and erosion rates were measured using Multislot Divisor Method. Nutrient leaching was analysed based on the content of C-organic (C_{tot}) (Walkley–Black method), total nitrogen (N_{tot}) (Kjeldahl method), P-available (Bray-1 method) and K_2O (extraction with 1N NH_4OAc at pH 7.0). From the results, land capability class III, cover crops (soybean) + manure (P3) treatment effectively reduced runoff and soil erosion ($22.63 \text{ m}^3 \cdot \text{ha}^{-1} \cdot \text{y}^{-1}$ and $13.04 \text{ Mg} \cdot \text{ha}^{-1} \cdot \text{y}^{-1}$), as well as nutrient leaching, compared to other treatments. Furthermore, sediment trap + cover crop + manure (P3) controlled runoff, erosion and nutrient leaching on land capability class IV, producing the lowest runoff ($129.40 \text{ m}^3 \cdot \text{ha}^{-1} \cdot \text{y}^{-1}$), soil erosion ($11.39 \text{ Mg} \cdot \text{ha}^{-1} \cdot \text{y}^{-1}$), C-organic (1.3%), and P (1.95 mg kg^{-1}). Soil conservation treatment significantly reduced erosion and runoff ($p < 0.05$) on land capability class VI. The bench terrace + cover plants + manure treatment-controlled runoff, erosion, and soil nutrient leaching.

Key words: cover crops, erosion, nutrient leaching, oil palm plantation, sediment trap, surface runoff, terraces

INTRODUCTION

Oil palm is widely grown in the fast-weathered tropical areas, including Indonesia, which has 18.2 mln ha land potential for crop cultivation. In 2010, the country planted 9 mln ha of oil palm [PUTRA *et al.* 2012], rising to 10 mln ha in 2016 [PIRKER *et al.* 2016]. In 2015, the oil palm plantation area in Aceh Province was 208,124 ha, down from 396,644 in 2013 [BPS 2014]. In Bireuen Regency, however, oil palm plantations have developed rapidly in recent years. For example, by 2012, the potential oil palm plantation in 14 regions was 27,434 ha [BPS Bireuen 2014]. Out of this, approximately 4,372 ha of land had been planted, of which 3,109 ha were community plantations. Soil and water conservation in oil palm cultivation is strengthened because this plant is globally associated with environmental damage.

Aceh's oil palm is mostly cultivated on newly cleared or converted land with a slope of 15–60%. Generally, this land falls under capability class III–VII [SATRIAWAN *et al.*

2017]. Lands in capability classes III and IV are agriculturally viable under adequate conservation. In contrast, lands in capability classes V–VII are agriculturally unsuitable owing to their damage vulnerability [SAIDA *et al.* 2013].

Intensive agricultural activities on unsuitable land triggers massive erosion exceeding $21.26 \text{ Mg} \cdot \text{ha}^{-1} \cdot \text{y}^{-1}$ [SATRIAWAN *et al.* 2015b]. This becomes a serious global threat to food security and sustainable development [SUI *et al.* 2016]. The massive soil erosion leads to loss of essential nutrients for growth and production in oil palm plantation. Numerous studies have been conducted on soil erosion in oil palm plantations in Indonesia. However, there are no studies on soil losses based on land capability classes. Soil erosion in oil palm plantations in Sumatera varies with age and conservation techniques ranging within $3.3\text{--}56.4 \text{ Mg} \cdot \text{ha}^{-1} \cdot \text{y}^{-1}$ [YENI *et al.* 2016], $12.20\text{--}18.66 \text{ Mg} \cdot \text{ha}^{-1} \cdot \text{y}^{-1}$ [SATRIAWAN *et al.* 2015a], and $23\text{--}793 \text{ Mg} \cdot \text{ha}^{-1} \cdot \text{y}^{-1}$ [DASTUR SYAH *et al.* 2013]. Therefore, this problem could be overcome through technological innovation. Mechanical soil conservation

technology has been applied in oil palm plantations using individual terraces and sedimentation [HARIBOWO *et al.* 2019; SATRIAWAN *et al.* 2017].

Soil erosion reduction in immature oil palm cultivation requires several conservation practices, including vegetative and mechanical methods. SATRIAWAN *et al.* [2016] and HARIBOWO *et al.* [2019] established that vegetation stabilizes soil against strong erosion flows, especially when using a combination of several plants such as Leguminosae. The research proposed the most effective soil and water conservation techniques according to capability classes.

MATERIALS AND METHODS

SITE DESCRIPTION AND EXPERIMENTAL DESIGN

The research took place in Peusangan Siblah Krueng Subdistrict, Bireuen Regency of Aceh Province (5°4'30" N and 96°45'18" E with 116 m elevation), with the previously determined land capability classes (III, IV and VI). Land capability class III was located on slopes (gradient 8–15%), having a soil solum depth of 85–125 cm, with a mild-moderate erosion. In this area, 25% of topsoil had been eroded, exposing the surface rocks. Land capability class IV was on slopes (15–30%), but in a moderate soil solum depth (80–90 cm). Also, a 50% of the upper layer was eroded. Land capability class VI was on the sloping area (>45%), where 75% of the upper layer had been lost.

This experiment lasted two years, from August 2014 until March 2016. The object of the research was immature plantations, 1–2 years old, with uniform land class groups. Oil palm trees are grown 8 × 8 m apart, based on the slope direction. The soil in the study region is classified as Typic Paleudults [USDA 2010], dominated by andesite rocks and sandy clay. The subsoil solum had a 0.5 m depth of argillic horizon, with clay content between 19 and 37%.

The research used the experimental method (standard of plot erosion) [SATRIAWAN *et al.* 2015a; 2016]. The study involved a standard plot erosion test. Soil conservation practices for oil palm were applied to the land capability classes previously determined. Different technologies were applied based on the recommendation of the individual classes [ARSYAD 2010].

In land capability class III, 4 treatments with 3 replications were analysed, including:

- P0 = no soil conservation,
- P1 = individual terrace (IT),
- P2 = individual terrace + cover crop (soybeans),
- P3 = individual terrace + cover crop (soybeans) + manure 15 Mg·ha⁻¹.

In land capability class IV, 4 treatments with 3 replications were examined, including:

- P0 = no soil conservation,
- P1 = sediment trap (ST),
- P2 = sediment trap + vertical mulch (VM),
- P3 = sediment trap + cover crop (soybeans) + manure 15 Mg·ha⁻¹.

In land capability class VI, 4 treatments with 3 replications were tested, including:

- P0 = no soil conservation,
- P1 = bench terrace (BT),
- P2 = bench terrace + cover crop (*Mucuna bracteata*),
- P3 = bench terrace + cover crop (*Mucuna bracteata*) + manure 15 Mg·ha⁻¹.

OBSERVATION AND DATA COLLECTION

Data was collected on runoff, suspended sediment (soil erosion), the concentration of C-organic, N, P, and K in the sediment.

The experiment was conducted on a plot of 22 × 4 m facing the slope. Surface runoff and erosion were measured using the multi-slot divisor method. The experimental plot boundary was set using an embedded plastic trap +20 cm into the ground and +20 cm above of soil surface. The runoff and erosion sediment collector was at the depth = 2 m, a width = 0.5 m, and a length = 0.5 m with 7 holes, each with a diameter of 5 cm. The middle hole was connected with a PVC pipe, 5 cm in diameter, to channel the overflow into a small 0.5 m × 0.5 m × 0.5 m container (Fig. 1c). In the course of the research, rainfall was recorded using an ombrometer installed near the experimental plot. Erosive rainfall was calculated after each rainfall by directly checking the runoff collector. According to EL KATEB *et al.* [2013], rainfall is non-erosive when precipitation cannot generate measurable runoff, producing no runoff and sediment. Erosive rainfall was characterized by runoff and sediment inside the collector.

Individual terraces have a length and width 1.5 × 1.5 m and a depth of 20 cm (Photo 1, Fig. 1). Sediment traps are long and wide soil trenches, measuring 1.5 m long, 1 m wide, and 0.5 m depth, made in the midst of planting rows perpendicular to slope direction (Photos 2a, b). The community designed the bench terraces, perpendicular to slope direction, with 3 m width and 1.5–2 m height (Photo 3). Bench terraces were constructed when preparing land for oil palm planting. Leguminous cover crops were grown with spacing of 20 cm × 30 cm.



Photo 1. Individual terrace in plot of land capability class III (phot. Fauzan)




Fig. 1. Sediment collector; source: own elaboration



Photo 2. Soil conservation technologies applied in the experiment of land capability class IV: a) sediment trap, b) sediment trap + cover crop + manure (phot. Fauzan)

Soybeans are planted between rows of oil palm trees as cover crops. They are planted throughout the year, planting 3 times a year by burying. During the period of growth, soybeans are treated from pests and diseases and harvesting is



Photo 3. Bench terrace land capability class VI (phot. Fauzan)

done. The interval between planting time and harvest is 3 weeks, allowing preparation time for the new crop. Application of manure was conducted 1 week before the soybeans were planted. The dose given is $15 \text{ Mg}\cdot\text{ha}^{-1}$ by sprinkling on the surface of the soil and mixing with the soil using a hoe. This step is done so that cattle manure is distributed evenly, making it easier to plant the soybeans.

MEASUREMENT OF SURFACE RUNOFF AND SOIL EROSION

Soil erosion is analysed through the filtration of water samples from the reservoir. The residue is dried in an oven at a constant 60°C . The measured sediment weight represents soil erosion, calculated as [SATRIAWAN *et al.* 2015a]:

$$E = \frac{C_{ap} \cdot V_{ap} \cdot 10^{-3}}{A} \quad (1)$$

where: E = eroded soil ($\text{Mg}\cdot\text{ha}^{-1}$), C_{ap} = concentration of sediment load ($\text{kg}\cdot\text{m}^{-3}$), V_{ap} = runoff volume (m^3), A = eroded area (ha), 10^{-3} = conversion from kg to Mg.

Sediment nutrients were analysed to determine C-organic (C_{tot}) (Walkley–Black method), total nitrogen (N_{tot}) (Kjeldahl method), P-available (Bray-1 method) and K_2O (extraction with $1\text{N NH}_4\text{OAc}$ at pH 7.0).

Data on surface runoff volume, erosion, and nutrient loss from erosion sediment were analysed by ANOVA. The mean separation was tested using the least significant difference (*LSD*) test at 5% significance level.

RESULTS

RAINFALL DURING STUDY

Within the experiment period, the total amount of rainfall recorded was 3,053 mm, with the highest values being recorded from October to December 2014 and September to December 2015. There were 24 erosive rainfall events during the observation period, in which a total of 1,895 mm was recorded (Fig. 2).




Fig. 2. Rainfall events over the rainy season in experiment duration (Aug 2014–March 2016); source: own elaboration acc. to rainfall and erosive rainfall data

RUNOFF, SOIL EROSION, AND NUTRIENT LEACHING

The soil conservation technologies applied in this study included single individual terraces (creating dishes around the oil palm canopy), individual terraces combined with crop strips, and cover crop (soybeans) together with manures. The treatment applied to oil palm plantations on land capability class III significantly influenced surface runoff, soil erosion, and C-organic and nitrogen leaching. However, there was no effect on the leaching of phosphate and potassium (Tab. 1). Soil conservation treatment significantly affected surface runoff volume and soil erosion. The P3 treatment was the most effective on surface runoff and soil erosion ($22.63 \text{ m}^3\cdot\text{ha}^{-1}\cdot\text{y}^{-1}$ and $13.04 \text{ Mg}\cdot\text{ha}^{-1}\cdot\text{y}^{-1}$) compared to other treatments ($p < 0.05$). The volumes of surface runoff were in the order, $\text{P3} < \text{P2} < \text{P1} < \text{P0}$. Compared to control treatments, surface runoff in P1, P2, and P3 decreased by 30.25, 47.78, and 69.66%, respectively, with significantly different treatments. Soil erosion had a similar trend order as surface runoff, with a reduction of 19.02%, 28.86%, and 47.65%. The P3 was effective for runoff intercept, due to the high density and large ground soybean coverage, resulting in the lowering of raindrop splash erosion compared to other treatments. Soil conservation treatment influenced nutrient leaching, runoff, and erosion (Tab. 1), in which the percentage of C-organic and total nitrogen were signifi-

cantly affected by the P3 treatment. Phosphate and potassium leaching were not affected by soil conservation treatment.

The treatment on land capability class IV involved the use of sediment trap (P1), sediment trap combined with vertical mulch (P2), as well as a sediment trap, cover crops, and manure (P3). Results showed that treatment with sediment traps cover crops, and manure (P3) significantly reduced surface runoff, soil erosion, and nutrient leaching (Tab. 2).

The treatment by sediment trap, cover crops, and manure (P3) on class IV land was appropriate in controlling surface runoff and soil erosion ($129.40 \text{ m}^3\cdot\text{ha}^{-1}\cdot\text{y}^{-1}$ and $11.39 \text{ Mg}\cdot\text{ha}^{-1}\cdot\text{y}^{-1}$) compared to others ($p < 0.05$). Surface runoff volumes were in the order of $\text{P3} < \text{P2} < \text{P1} < \text{P0}$. Compared with the control treatment, the runoff of P1, P2, and P3 reduced by 9.03, 23.19, and 37.95%, respectively, with significantly different treatments, except for P0 and P1. The soil conservation treatment reduced soil erosion by 50.80, 52.52, and 63.97%, respectively, and the P1, P2, and P3 treatments significantly differed with P0. Soil nutrient leaching, runoff, and erosion were affected by the treatment (Tab. 2), in which the percentage of C-organic and phosphorus were significantly affected. However, the treatment did not affect nitrogen and potassium nutrient removal. In this group, the sediment trap + cover crop + manure (P3) treatment did not prevent nutrient leaching.

Table 1. Runoff, soil erosion and nutrient leaching in sediment on land capability class III

Soil conservation technology	Runoff ($\text{m}^3\cdot\text{ha}^{-1}\cdot\text{y}^{-1}$)	Erosion ($\text{Mg}\cdot\text{ha}^{-1}\cdot\text{y}^{-1}$)	C-organic (%)	N (%)	P (mg kg^{-1})	K (cmol kg^{-1})
Control (P0)	74.66 ^d	24.91 ^d	4.39 ^b	0.43 ^{ab}	2.47	0.44
Individual terrace (P1)	52.07 ^c	20.17 ^c	3.70 ^{ab}	0.42 ^{ab}	2.04	0.42
Individual terrace + cover crop (P2)	38.98 ^b	17.72 ^b	3.40 ^{ab}	0.29 ^{ab}	1.37	0.28
Individual terrace + cover crop + manure (P3)	22.65 ^a	13.04 ^a	2.70 ^a	0.22 ^a	1.20	0.23
<i>LSD</i> _{0.05}	2.01	1.64	1.20	0.20		

Explanations: *LSD* = least significant difference; different letter notations in the same column show significant differences on the 0.05 *LSD* test. Source: own study.

Table 2. Runoff, soil erosion and nutrient leaching in sediment on land capability class IV

Soil conservation technology	Runoff ($\text{m}^3 \cdot \text{ha}^{-1} \cdot \text{y}^{-1}$)	Erosion ($\text{Mg} \cdot \text{ha}^{-1} \cdot \text{y}^{-1}$)	C organic (%)	N (%)	P (mg kg^{-1})	K (cmol kg^{-1})
Control (P0)	208.55 ^c	31.61 ^b	2.72 ^b	0.24	4.07 ^b	1.16
Sediment trap (P1)	189.70 ^c	15.55 ^a	1.65 ^a	0.15	3.93 ^b	1.10
Sediment trap + vertical mulch (P2)	160.18 ^b	14.06 ^a	1.47 ^a	0.14	2.39 ^a	0.91
Sediment trap + cover crop + manure (P3)	129.40 ^a	11.39 ^a	1.30 ^a	0.17	1.95 ^a	0.90
<i>LSD</i> _{0.05}	21.18	11.02	0.9		1.08	

Explanations: *LSD* = least significant difference; different letter notations in the same column show significant differences on the 0.05 *LSD* test. Source: own study.

Table 3. Runoff, soil erosion and nutrient leaching in sediment on land capability class VI

Soil conservation technology	Runoff ($\text{m}^3 \cdot \text{ha}^{-1} \cdot \text{y}^{-1}$)	Erosion ($\text{Mg} \cdot \text{ha}^{-1} \cdot \text{y}^{-1}$)	C organic (%)	N (%)	P (mg kg^{-1})	K (cmol kg^{-1})
Control	318.59 ^b	60.38 ^b	2.83 ^b	0.56 ^b	2.49	0.71
Bench terrace	174.37 ^a	29.58 ^a	1.67 ^a	0.33 ^a	1.70	0.54
Bench terrace + cover crop	139.66 ^a	22.50 ^a	1.41 ^a	0.27 ^a	2.33	0.56
Bench terrace + cover crop + manure	115.98 ^a	21.29 ^a	1.32 ^a	0.24 ^a	1.67	0.67
<i>LSD</i> _{0.05}	63.2	5.2	0.85	0.17		

Explanations: *LSD* = least significant difference; different letter notations in the same column show significant differences on the 0.05 *LSD* test. Source: own study.



Photo 4. Individual terrace + cover crop + manure in plot of land capability class III (phot. Fauzan)



Photo 5. Bench terrace + cover crop in plot of land capability class VI (phot. Fauzan)

Soil conservation treatment on land capability class VI significantly reduced erosion and runoff more than the control treatment ($p < 0.05$) – Table 3. The bench terrace + cover crop + manure treatment significantly reduced surface runoff by 63.59%, compared to the control treatment, followed by bench terrace + cover crop (56.16%) and bench terrace (45.26%), respectively (Photo 4). The bench terrace + cover crop + manure (P3) reduced soil erosion more than other treatments, 64.73% compared to other erosion controls. Soil nutrient leaching, runoff, and erosion were greatly influenced by the conservation treatment applied (Tab. 3 and Photo 5), in which the percentage of C-organic and total nitrogen were significantly affected by the P3 treatment. However, phosphate and potassium leaching was not affected by the treatment. In this group, bench terrace + cover crop + manure (P3) was the most effective in preventing nutrient leaching.

DISCUSSION

Erosion is caused by forced movement of the ground-mass by water, wind, or gravity. The humid and tropical climate in Indonesia increases the risk of water erosion [ARSYAD 2010]. Erosion is accelerated by the intense rainfall, combined with the relatively steep-sloped topography. Intense agricultural activities in these areas also increase soil erosion [SATRIAWAN, FUADY 2014].

Mitigation of surface runoff and soil erosion in agricultural land use, especially on a new oil palm plantation, is the key to soil protection [ZUAZO, PLEGUEZUELO 2008]. In this study, the use of soybeans as cover crops between rows proved to be efficient in managing surface runoff and soil erosion (Photo 4). The dense soybean canopy reduces the erosivity of rainfall, therefore limiting soil loss. Cover crops, therefore, play an essential role in soil conservation. They positively affect physical soil characteristics, including infiltration, moisture, and bulk density [HULUGALLE *et*

al. 1986]. Ground leguminous cover plants increase the content of organic matter, naturally provide nitrogen (N) by using N_2 binding legumes, providing critical nutritional contributions to oil palm plants, which influences their growth [IBEWIRO *et al.* 2000; ILE *et al.* 1996; OBIAGWU 1995; SALAKO, TIAN 2003].

Related to the improvement of soil physical properties, manure helped form stable aggregates in this study. Maintenance of soil particles from the detachment of rainfall erosivity is best achieved using the stable aggregates. Organic matter binds soil particles, yielding microaggregates further bound by plant roots and fungal hyphae. As a result, the primary and secondary soil grains are compacted together [OADES 1984].

Manure stabilizes and increases the size of soil aggregates, raising hydrophobicity and inter-particle cohesion [CHENU *et al.* 2000]. Stable aggregates increase soil porosity, facilitate water absorption and improve the capacity for water retention. Furthermore, soil aggregation and aggregate stability are critical aspects contributing to limiting erosion. JUARSAH *et al.* [2008] established the importance of organic matters to the physical and chemical soil characteristics, including increasing aggregation, preventing aggregate destruction by water, making the soil more tillable, improving porosity and aeration, as well as increasing infiltration and percolation capacities.

Soil conservation systems significantly lowered C-organic, nitrogen, and phosphate loss due to limited surface runoff ($p < 0.05$), although the impact depended on the treatment (Tab. 1). The P3 treatments were appropriate, leading to minimum C-organic, nitrogen, and phosphate, with the lowest concentrations (2.7%, 0.22%, and 1.22 mg kg⁻¹, respectively). According to (Tab. 1), runoff and nutrient concentrations influenced nutrient loss on oil palm cultivated land, since Nitrogen was correlated with C-organic. In general, soil conservation on modified land surface roughness through individual terraces and maximized land cover, therefore preventing nutrient loss through erosion.

Based on land capability class IV in this project, sediment traps collected water and controlled the sediment moved by surface runoff. Apart from cover plants, the soil helped in controlling runoff, with improved aggregation and soil physical properties. This is consistent with previous results that the application effectively suppressed erosion by 71%, based on soil structure and land cover condition. The shorter the distance between the sediment trap on the same slope, the more it reduced erosion and runoff and increased groundwater [BRATA 1998; MONDE 2010; MURILAKSONO *et al.* 2008].

Sediment trap + cover crop (soybean) and manure treatment reduced organic C, N, P, and K loss. Based on Table 2, the lowest level of C-organic, nitrogen, there were phosphate and potassium in the P3 treatment. The lower loss of nutrients in sediment trap + cover crop + was attributed to cover crops' role, which appropriately utilized nutrients. Furthermore, the sediment trapping availed nutritious water to the plant roots. However, the P2 treatment was not significantly different from P0. This is because the P elements,

which are mobile and highly soluble in water, were mainly evident in the control and sediment trap treatment.

In sloping land, the conservation measures considerably influenced erosion by changing the soil surface [MAETENS *et al.* 2012]. Cover crops and strip plants benefit plantation trees because they minimize rainfall erosivity and surface runoff, as well as adding to the soil's organic matter through the stems, twigs, and fallen dead leaves. They transpire, reducing soil water content. Cover plants reduce the strength of the rainwater dispersion and the quantity and speed of runoff, enhancing infiltration. As a result, they lessen erosion and nutrient leaching. According to BAH *et al.* [2014], nutrient loss through runoff and sediments relates to oil palm age, surface feature, land clearing, and adequate soil conservation measures.

Soil and water conservation influenced nutrient loss of land capability class VI, especially to C-organic, nitrogen, and phosphate. The greatest loss was in the control treatment, without soil and water conservation. Of all the treatments, soybean as cover crops and strip plant highly suppressed soil nutrient loss than the other treatments.

Based on the socio-economic benefits of oil palm plantations, land cover crops, and simple conservation buildings effectively reduced surface runoff and erosion, improving farming sustainability in the area. However, in general, soil erosion largely depends on the management system.

CONCLUSIONS

The study conclusions are based on empirical analysis and field discussions. The results show that the soil conservation techniques applied in every land capability class-controlled surface runoff and erosion. In terms of land capability, class III cover crops + manure (P3) treatment effectively reduced runoff and soil erosion (22.63 m³·ha⁻¹·y⁻¹ and 13.04 Mg·ha⁻¹·y⁻¹) and nutrient leaching, compared to other treatments. In land capability class IV, sediment trap + cover crop + manure (P3) reduced runoff, erosion and nutrient leaching, producing the lowest runoff (129.40 m³·ha⁻¹·y⁻¹), soil erosion (11.39 Mg·ha⁻¹·y⁻¹), C organic (1.3%), and P (1.95 mg kg⁻¹). Finally, on land capability class VI, located on the sloping area (>45%), with heavily eroded soil, the conservation treatment significantly reduced erosion and runoff. Bench terrace + cover crop + manure minimized runoff, erosion, and soil nutrient leaching, reducing surface runoff by 63.59%.

Based on the findings of this study, in the cultivation of oil palm on sloping land in Aceh and Indonesia in general, the application of soil conservation techniques is absolutely necessary, given the potential for soil erosion in the tropics is very large. It is important to communicate this policy as negative issues on oil palm becoming global.

ACKNOWLEDGEMENTS

The authors acknowledge the Directorate of Research and Community Service (DRPM) of the Ministry of Research, Technology, and Higher Education for financial support on the Competitive Grant Research Scheme.

REFERENCES

- ARSYAD S. 2010. Konservasi tanah dan air [Soil and water conservation]. 10th ed. Bogor. IPB Press. ISBN 978-493-003-2. pp. 496 [In Indonesian].
- BAH A., HUSNI M.H.A., TEH C.B.S., RAFII M.Y., SYED OMAR S.R., AHMED O.H. 2014. Reducing runoff loss of applied nutrients in oil palm cultivation using controlled-release fertilizers. *Advances in Agriculture*. Vol. 2014. Art. ID 285387. DOI [10.1155/2014/285387](https://doi.org/10.1155/2014/285387).
- BPS 2014. Statistik kelapa sawit Indonesia 2014 [Indonesian oil palm statistics]. Badan Pusat Statistik. Jakarta. pp. 76. [Access 4.8.2020]. Available at: <https://media.neliti.com/media/publications/48713-ID-statistik-kelapa-sawit-indonesia-2014.pdf>
- BPS Bireuen 2014. Komposisi dan produktivitas keadaan perkebunan rakyat tanaman kelapa sawit di Kabupaten Bireuen Tahun 2013. Di: Bireuen dalam angka 2014 [Composition and productivity of oil palm smallholders estates of Bireuen District, 2013. In: Bireuen in figures 2014] [online]. Badan Pusat Statistik kabupaten Bireuen pp. 265. [Access 4.08.2020]. Available at: http://bappeda.bireuenkab.go.id/wp-content/uploads/2015/04/BIREUEN-Dalm-Angka-2014_revisi.com-pressed.pdf
- BRATA K.R. 1998. Pemanfaatan jerami padi sebagai mulsa vertikal untuk pengendalian aliran permukaan, erosi dan kehilangan unsur hara dari pertanian lahan kering [Utilization of rice straw as a vertical mulch for surface flow control, erosion and loss of nutrients from dryland agriculture]. *Jurnal Ilmu Tanah dan Lingkungan*. Vol. 1(1) p. 21–27.
- CHENU C., BISSONNAIS Y.L., ARROUAYS D. 2000. Organic matter influence on clay wettability and soil aggregate stability. *Soil Science Society of America Journal*. Vol. 64 p. 1479–1486. DOI [10.2136/sssaj2000.6441479x](https://doi.org/10.2136/sssaj2000.6441479x).
- DASTUR SYAH M.R., ALIBASYAH, SYAMAUN A.A. 2013. Prediksi erosi pada beberapa tingkat umur kelapa sawit (*Elaeis guineensis* Jacq) [The prediction of soil erosion on various age of oil palm]. *Jurnal Manajemen Sumberdaya Lahan*. Vol. 2. No. 4 p. 304–316.
- EL KATEB H., ZHANG H.F., ZHANG P.C., MOSANDL R. 2013. Soil erosion and surface runoff on different vegetation covers and slope gradients: A field experiment in Southern Shaanxi Province, China. *Catena*. Vol. 105 p. 1–10. DOI [10.1016/j.catena.2012.12.012](https://doi.org/10.1016/j.catena.2012.12.012).
- HARIBOWO R., ANDAWAYANTI U., LUFIRA R.D. 2019. Effectivity test of an eco-friendly sediment trap model as a strategy to control erosion on agricultural land. *Journal of Water and Land Development*. No. 42 (VII–IX) p. 76–82. DOI [10.2478/jwld-2019-0047](https://doi.org/10.2478/jwld-2019-0047).
- HULUGALLE N.R., LAL R., TER KUILE C.H.H. 1986. Amelioration of soil physical properties by Mucuna after mechanized land clearing of a tropical rain forest. *Soil Science*. Vol. 141(1) p. 219–224.
- IBEWIRO B., SANGINGA N., VANLAUWE B., MERCKX R. 2000. Nitrogen contributions from decomposing cover crop residues to maize in a tropical derived savanna. *Nutrient Cycling in Agroecosystems*. Vol. 57 p. 131–140.
- ILE E., HAMADINA M.K., ZUFA K., HENROT J. 1996. Note on effects of a *Mucuna pruriens* var. utilis crop on the growth of maize (*Zea mays*) on an acid Ultisol in southeastern Nigeria. *Journal of Field Crops Research*. Vol. 48(2–3) p. 135–140. DOI [10.1016/S0378-4290\(96\)01028-3](https://doi.org/10.1016/S0378-4290(96)01028-3).
- ISMAIL M., YACOB R.K. 2012. Digital soil map using the capability of new technology in sugar beet area, Nubariya, Egypt. *The Egyptian Journal of Remote Sensing and Space Science*. Vol. 15. Iss. 2 p. 113–124. DOI [10.1016/j.ejrs.2012.08.001](https://doi.org/10.1016/j.ejrs.2012.08.001).
- JUARSAH I., YUSTIKA R.D., ABDURRAHMAN A. 2008. Pengendalian erosi dan kahat bahan organik tanah pada lahan kering berlereng mendukung produksi pangan nasional [The soil erosion control and organic matters deficiencies on dry land to support of national food production]. *Prosiding Seminar Nasional Multifungsi Dan Konversi Lahan Pertanian*. Bogor. Indonesian Soil Research Institute p. 255–273.
- MAETENS W., VANMAERCKE M., POESEN J., JANKAUSKAS B., JANKAUSKIENE G., IONITA I. 2012. Effects of land use on annual runoff and soil loss in Europe and the Mediterranean. A meta-analysis of plot data. *Progress in Physical Geography*. Vol. 36 p. 599–653. DOI [10.1177/0309133312451303](https://doi.org/10.1177/0309133312451303).
- MANSOORI H., KOHANSAL M.R. 2009. Introducing a lexicographic goal programming for environmental conservation program in farm activities: A case study in Iran. *China Agricultural Economic Review*. Vol. 1. No. 4 p. 478–484. DOI [10.1108/17561370910989284](https://doi.org/10.1108/17561370910989284).
- MONDE A. 2010. Pengendalian aliran permukaan dan erosi pada lahan berbasis kakao di DAS Gumbasa Sulawesi Tengah [The control of runoff and erosion on cocoa-based land in the Gumbasa watershed of Central Sulawesi]. *Media Litbang Sulteng* III (2) p. 131–136.
- MURTIKSONO K., SUTARTA E.S., SIREGAR H.H., DARMOSAROKORO W., HIDAYAT Y. 2008. Penarapan teknik konservasi tanah dan air dalam upaya penekanan aliran permukaan dan erosi di kebun kelapa sawit [Application of soil and water conservation techniques in efforts to suppress of runoff and erosion in oil palm plantations] [online]. [Access 12.10.2017]. Available at: http://repository.ipb.ac.id/bitstream/handle/123456789/37076/Seminar%201_Kmurtiaksono.pdf?sequence=1&isAllowed=y
- OADES J.M. 1984. Soil organic matter and structural stability: Mechanisms and implications for management. *Plant Soil*. Vol. 76 p. 319–337.
- OBIAGWU C.J. 1995. Estimated yield and nutrient contributions of legume cover crops intercropped with yam, cassava, and maize in the Benue River basins of Nigeria. *Journal of Plant Nutrition and Soil Science*. Vol. 18(12) p. 2775–2782.
- PIRKER J., MOSNIER A., KRAXNER F., HAVLIK P., OBERSTEINER M. 2016. What are the limits to oil palm expansion?. *Global Environmental Change*. Vol. 40 p. 73–81. DOI [10.1016/j.gloenvcha.2016.06.007](https://doi.org/10.1016/j.gloenvcha.2016.06.007).
- PUTRA E.T.S., SIMATUPANG A.F., SUPRIYATNA, WALUYO S., INDRADAWA D. 2012. The growth of one year-old oil palms intercropped with soybean and groundnut. *Journal of Agricultural Science*. Vol. 4. No. 5 p. 169–180.
- SAIDA, ABDULLAH, JUSOFF K., ILSAN M., HARIS A., NURAENI 2013. Evaluation of land capability for agriculture in the upstream of Jeneberang watershed, South Sulawesi. *American-Eurasian Journal Agricultural & Environmental Sciences*. Vol. 13(8) p. 1027–1033. DOI [10.5829/idosi.ajeaes.2013.13.08.11014](https://doi.org/10.5829/idosi.ajeaes.2013.13.08.11014).
- SALAKO F.K., TIAN G. 2003. Soil water depletion under various leguminous cover crops in the derived savanna of West Africa. *Agriculture, Ecosystems and Environment*. Vol. 100(2–3) p. 173–180.
- SATRIAWAN H., FUADY Z., MAYANI N. 2016. Soil conservation by vegetative systems in oil palm plantation. *Polish Journal of Soil Science*. Vol. 49. No. 2 p. 223–235. DOI [10.17951/pjss/2016.49.2.223](https://doi.org/10.17951/pjss/2016.49.2.223).
- SATRIAWAN H., FUADY Z., AGUSNI A. 2017. Soil conservation techniques in oil palm cultivation for sustainable agriculture. *Jurnal Pengelolaan Sumberdaya Alam dan Lingkungan*. Vol. 7. No. 2 p. 178–183.
- SATRIAWAN H., FUADY Z. 2014. Teknologi konservasi tanah dan air [Soil and water conservation technology]. 1st ed. Yogyakarta. Deepublish. ISBN 6022807832 pp. 142.
- SATRIAWAN H., HARAHAP E.M., RAHMAWATY, KARIM A. 2015a. Effectiveness of soil conservation to erosion control on several

- land use types. *Agriculture (Poľnohospodárstvo)*. Vol. 61(2) p. 61–68. DOI [10.1515/agri-2015-0011](https://doi.org/10.1515/agri-2015-0011).
- SATRIAWAN H., HARAHAP E.M., RAHMAWATY, KARIM A. 2015b. Analysis of soil erosion in agricultural land use in Krueng Sieumpo watershed Aceh Province. *Academic Research International*. Vol. 6(2) p. 18–27.
- SUI Y., QU Y., YAN B., XU X., ROUSSEAU A.N., ZHANG Y. 2016. Assessment of micro-basin tillage as a soil and water conservation practice in the black soil region of northeast China. *Plos One*. Vol. 11(3): e0152313. DOI [10.1371/journal.pone.0152313](https://doi.org/10.1371/journal.pone.0152313).
- USDA 2010. *Keys to soil taxonomy*. 11th ed. Washington, D.C. United States Department of Agriculture Natural Resources Conservation Service pp. 337.
- ZUAZO V.H.D., PLEGUEZUELO C.R.R. 2008. Soil erosion and runoff prevention by plant covers. A review. *Agronomy Sustainable Development*. Vol. 28 p. 65–86. DOI [10.1051/agro:2007062](https://doi.org/10.1051/agro:2007062).
-