



PubMed

ISSN: 0976-237X

# CCD

CONTEMPORARY CLINICAL DENTISTRY



Wolters Kluwer  
Health

Medknow

[www.contempclindent.org](http://www.contempclindent.org)

[Previous Issue](#)



## Apr-Jun 2023 - Volume 14 - Issue 2

- [Table of Contents Outline](#)
- [eTOC Alerts](#)
- [Contributor Index](#)

## About the Journal

The journal **Contemporary Clinical Dentistry** (CCD) (Print ISSN: 0976-237X, E-ISSN:0976- 2361) is peer-reviewed journal published on behalf of Maharishi Markandeshwar University and issues are published quarterly in the last week of March, June, September and December. The Journal publishes Original research papers, clinical studies, case series strictly of clinical interest. Manuscripts are invited from all specialties of Dentistry i.e. Conservative dentistry and Endodontics, Dentofacial orthopedics and Orthodontics, Oral medicine and Radiology, Oral pathology, Oral surgery, Orofacial diseases, Pediatric Dentistry, Periodontics, Clinical aspects of Public Health dentistry and Prosthodontics. Review articles are not accepted. Review, if published, will only be by invitation from eminent scholars and academicians of National and International repute in the field of Medical/Dental education.

## Abstracting and Indexing Information

The journal is registered with the following abstracting partners:

Baidu Scholar, CNKI (China National Knowledge Infrastructure), EBSCO Publishing's Electronic Databases, Ex Libris – Primo Central, Google Scholar, Hinari, Infotrieve, National Science Library, ProQuest, TDNet, Wanfang Data

The journal is indexed with, or included in, the following:

DOAJ, Emerging Sources Citation Index, Indian Science Abstracts, IndMed, PubMed Central, SCOPUS, Web of Science

Impact Factor<sup>®</sup> as reported in the 2022 Journal Citation Reports<sup>®</sup> (Clarivate Analytics, 2023): **1.2**

## Journal Ethics

Wolters Kluwer and Journal/Association are committed to meeting and upholding standards of ethical behavior at all stages of the publication process. We follow closely the industry associations, such as the Committee on Publication Ethics (COPE), International Committee of Medical Journal Editors (ICMJE) and World Association of Medical Editors (WAME), that set standards and provide guidelines for best practices in order to meet these requirements. For a summary of our specific policies regarding duplicate



publication, conflicts of interest, patient consent, etc., please visit [www.Medknow.com/EthicalGuidelines.asp](http://www.Medknow.com/EthicalGuidelines.asp)

### Open Access Publication and Creative Commons Licensing

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

### Digital Archiving

Wolters Kluwer Medknow provides for long-term digital preservation through two primary partnerships, Portico and CLOCKSS.

Portico is a leading digital preservation service worldwide. The content is preserved as an archival version and is not publically accessible via Portico, but is provided when required under specific conditions, such as discontinuation of the collection or catastrophic failure of the website.

CLOCKSS will enable any library to maintain their own archive of content from Wolters Kluwer Medknow and other publishers, with minimal technical effort and using cheaply available hardware.



### Ahead of Print policy

Articles published online under the Ahead of Print model are considered published and can be cited and quoted using the DOI as the reference source. Wolters Kluwer Medknow has a policy that changes will not be made after publication of an article without following accepted procedures for making corrections to the scientific record.

### Advertisements

While advertisements are crucial to this journal to be able to keep all content free for everyone, ethical considerations are in place to ensure the integrity of the journal and its content:

- "Pop-up" and "banner" ads appear on a random, rotating basis. The advertiser has no control or input over the pages where their ads appear.
- The Editorial Board has full and final approval over the content of all advertisements.
- Advertisers will never be shown any manuscripts or other content prior to publication.

### Scope of the Journal

The journal will cover technical and clinical studies related to diagnosis and therapy during the pre-natal stage in humans including ethical and social issues. Articles with clinical interest and implications will be given preference.



### Medknow Publications

Medknow, part of Wolters Kluwer Health, is one of the largest open access publishers worldwide with more than 450+ medical journals in its portfolio. Students, researchers, clinicians and other healthcare professionals worldwide access Medknow journals to help them fuel new discoveries and improve patient care.

Founded in 1997, Medknow is a well-respected medical publisher that fully supports many of the popular open access models through its peer-review management system. Medknow's open access policy has resulted in more than a half a million article downloads monthly for all of its journals.

Wolters Kluwer Health (Philadelphia, PA) is a leading global provider of information, business intelligence and point-of-care solutions for the healthcare industry. Serving more than 150 countries and territories worldwide, Wolters Kluwer Health's customers include professionals, institutions and students in medicine, nursing, allied health and pharmacy. Major brands include Lippincott Williams & Wilkins, Ovid®, UpToDate®, Medi-Span®, Facts & Comparisons®, Pharmacy OneSource®, Lexicomp® and ProVation® Medical. Wolters Kluwer Health is part of Wolters Kluwer, a market-leading global information services company with annual revenues (2019) of €3.4 billion (\$4.7 billion), approximately 19,000 employees worldwide and operations in over 40 countries across Europe, North America, Asia Pacific, and Latin America.

**Medknow Specialties**

Online and Print Scholarly Publishing, Peer-Review System, Specialized in Medical Research, Largest Open Access Publisher

[Back to Top](#)**Never Miss an Issue**

Get new journal Tables of Contents sent right to your email inbox

[Get New Issue Alerts](#)**Browse Journal Content**[Register on the website](#)[Get eTOC Alerts](#)**Customer Service**[Submit a Service Request](#)

800-638-3030 (within the USA)

301-223-2300 (outside of the USA)

[Manage Cookie Preferences](#)

[Privacy Policy \(Updated June 29, 2023\)](#) - [Legal Disclaimer](#) - [Terms of Use](#) - [Open Access Policy](#)  
Copyright © 2023 Contemporary Clinical Dentistry | Published by Wolters Kluwer - Medknow

## Editorial Board

**Editor-in Chief**

Prof. G .M. Sogi  
Maharishi Markandeshwar (Deemed to be University),  
Mullana, Ambala (HR)  
INDIA-133207  
Email- [chiefeditor.ccdjournal@mmumullana.org](mailto:chiefeditor.ccdjournal@mmumullana.org)

**Associate Editors**

Prof. Sivakumar Nuvvula  
Narayana Dental College and Hospital  
Nellore, Andhra Pradesh  
INDIA -524003

Dr. Monika Gupta  
Maharishi Markandeshwar (Deemed to be University),  
Mullana, Ambala (HR)  
INDIA-133207

Dr Deepak Gupta  
Maharishi Markandeshwar (Deemed to be University),  
Mullana, Ambala (HR)  
INDIA-133207

**Assistant Editors**

Dr. Prachi Goyal  
Maharishi Markandeshwar (Deemed to be University),  
Mullana, Ambala (HR)  
INDIA-133207

Dr Jasneet Sudan  
Maharishi Markandeshwar (Deemed to be University),  
Mullana, Ambala (HR)  
INDIA-133207

### **International Advisory Editorial Board**

Dr. Raman Bedi, Head of Centre for International Child Oral Health, King's College, London, United Kingdom

Dr. Prathip Phantumvanit, Dean Emeritus, Faculty of Dentistry, Thammasat University, Pathum Thailand

Dr. Monty Duggal, Professor of Paediatric Dentistry, National University of Singapore, Singapore

Dr. Simrit Malhi, Head, Department of Paediatric Dentistry and Orthodontics, Westmead Hospital, Sydney, Australia

Dr. Paul Mullasseril, Chair of Restorative dentistry at Oklahoma University. USA

Dr. Richard Welbury, Professor / Professor / Honorary Consultant in Paediatric Dentistry., University of Central Lancashire, UK

Dr. Lars Andersson, Chairman of Department of Surgical Sciences, Faculty of Dentistry, Kuwait University, Kuwait

Dr. Arthur M Kemoli, Chairman, Paediatric Dentistry and Orthodontics, University of Nairobi, Kenya

Prof Wim van Palenstein Helderma WHO Collaborating Centre, Nijmegen, The Netherlands

Dr. Richard Widmar, Head of Paediatric Dentistry, Sydney Children's Hospital Network, Sydney, Australia

Dr. Alessandro Cavalcanti, Chair of the Post Graduate Program in Public Health at the State University of Paraiba, Brazil

Dr. Duygu Ilhan, member Education committee of Istanbul Dental Chamber, Turkey

Dr. David M. Okuji, Associate Director-Extramural Residency Training, NYU Lutheran Medical Center of Dental Medicine, New York USA

Dr. Chad Gehani, trustee from the Second District (New York State) American Dental Association USA

Dr. Stephen J Moss, President at Health Education Enterprises, Greater New York City Area

Dr. Denis Bourgeois, Claude Bernard University Lyon, France

Prof. Dr. Drg.Tri Erri Astoeti, MKes, Dean Faculty of Dentistry Trisakti University, Jakarta, Indonesia

Prof. Dr. Armelia Sari Widyarman, Head Department of Oral Microbiology, Faculty of Dentistry, Trisakti University, Jakarta, Indonesia

Dr. Wen Sheng Rong, Professor, Department of Preventive Dentistry, Peking University, School & Hospital of Stomatology, China

Dr. Vineet K Dhar, Associate professor and interim-chair of Orthodontics and Paediatric Dentistry University of Maryland, USA

Dr. How Kim Chuan, Managing Director and Chief Consultant at Imperial Dental Specialist Centre, Malaysia

Dr Ali Farahani, Craniofacial and Special Care Orthodontics (Children's Hospital Los Angeles, USC)

Dr. Maryam Khoroushi, Isfahan University of Medical Sciences Isfahan, Iran

Dr. Soni Stephen, Director and Principal Specialist Southside Paediatric Dentistry, Miranda, Sydney, Australia

Dr. Archana Viswanath, Department of Oral and Maxillofacial Surgery, Tufts University, USA

Dr. Vajihesadat Mortazavi, Professor of Restorative Dentistry, Isfahan University of Medical Sciences, Iran

Dr. Prasad Amaratunga, Professor, Faculty of Dental Sciences and Hospital, University of Peradeniya, Sri Lanka.

---

Dr. Bhumiya Gupta, Assistant Professor of Clinical Dentistry, Eastman Institute for Oral Health University of Rochester, New York, USA

**Section Editors Board**  
**Public Health Dentistry**

Dr. R. K. Bali, Former President, Dental Council of India, New Delhi

Dr. S. S. Hiremath, Senior Professor and Head, Department of Public Health Dentistry, The Oxford Dental College, Hospital and Research Centre, Bengaluru, Karnataka, India

Dr. Ganesh Shenoy, Professor and Head, Department of Public Health Dentistry, Yenepoya Dental College, Mangaluru, India

Dr. M. B. Aswath Narayanan, Professor and Head, Department of Public Health Dentistry, Tamil Nadu Govt. Dental College, Chennai, TN, India

**Conservative Dentistry**

Dr. Dibyendu Mazumder, President, Dental Council of India

Dr. Sanjay Tewari, Principal, Dental College (PGIMER), Rohtak, Haryana, India

Dr. Naseem Shah, Former Head, AIIMS, New Delhi

Dr. T. Rambabu, Professor and Head, Drs. Sudha & Nageswara Rao Siddhartha Institute of Dental Sciences, Chinoutapally, Gannavaram (M), Krishna Dist., Andhra Pradesh

Dr. Rajiv Chugh, Private Practice, New Delhi

Dr. Abhay Kamra, Former Professor & Head, Dept. Conservative Dentistry & Endodontics - C.S.M.S.S. Dental College & Hospital, Aurangabad.

Dr. Devender Chaudhary, Principal, Head, Maharaja Ganga Singh Dental College & Research Centre

Dr. Mithra Hegde, Senior Professor & Head of the Department, A.B Shetty Memorial Institute of Dental Sciences, Mangalore

Dr. Vinod Babu Mathew, King Khalid University, Saudi Arabia

Dr. Karthick, K, KSR Institute of Dental Science and Research, Tiruchengode, TN.

#### **Esthetic Dentistry**

Dr. Sandesh Mayekar, Esthetic Dentist, Private practice, Mumbai

Dr. Poras Turner, Professor of Postgraduate studies in the Dept. of Prosthodontics at A. B. Shetty Institute of Dental Studies, Mangalore, India

Dr. Deepak Mucchala, Private practice, Mumbai

#### **General Dentistry**

Dr. R. Bellie, Private practice, Coimbatore, Tamil Nadu

Dr. George Thomas, Private practice, Kumbakonam, Tamilnadu

Dr. Bhagwant Singh, Private practice, Ludhiana

#### **HIV & Oral Health**

Dr. Amar Pazare, Prof. & HOD Medicine at Seth G. S. Medical College & KEM hospital, Parel, Mumbai

#### **Pedodontics**

Dr. A.K. Munshi, Former Director, Principal, K. D. Dental College & Hospital, Mathura

Dr. Krishan Gauba, Head, Oral Health Sciences Centre, PGI, Chandigarh

Dr. I. K. Pandit, Principal, D. A. V. Dental College & Hospital, Yamuna Nagar, Haryana

Dr. Navneet Grewal, Prof. and Head, Govt. Dental College, Amritsar

Dr. Narendra Chandranee, Former Professor & Head, VSPM Dental College & Research, Nagpur, Maharashtra

Dr. Ashima Goyal, Professor, OHSC, PGI, Chandigarh

Dr. Attiguppe R Prabhakar, Vice Principal, Professor and Head, Bapuji Dental College & Hospital, Davangere

---



Dr. TN Tilak Raj, Professor and Head, Shymla Reddy Dental College, Bangalore

Dr. J. Baby John, Principal, VMS Dental College, Salem

Dr. Shalini Garg, Professor and Head, Sudha Rahstogi Dental College, Faridabad

Dr. B. Nandlal, Professor, J S S, Dental College, Mysore

Dr. Nikhil Srivastava, Principal and Dean, Subharti Dental College and Hospital, Meerut

Dr. Sham Bhat, Vice Principal and Head, Yenepoya Dental College, Mangalore

Dr. Abi Thomas, Principal, Christian Dental College, Ludhiana

Dr. Mousumi Singh, Professor and Head, ITS Dental College, Greater Noida (U.P.)

Dr. Ramakrishna Yeluri, Professor and Head, Teerthankar Mahavidyalaya, Moradabad

Dr. Adesh Kakde, Nair Hospital Dental College, Mumbai

Dr. Amita Tiku, Prof and Head, Bharti Vidhyapeeth Dental College, Navi Mumbai, Maharashtra

Dr. N. Venugopal Reddy, Prof and Head, Mamata Dental College Khammam Andhra Pradesh

Dr. Vijay P Mathur, Professor and Head, AIIMS, New Delhi

Dr. Tejashree Gupte, Professor, Nair Hospital Dental College, Mumbai

### **Periodontics**

Dr. A. Kumarswamy, Periodontist, Private Practice, Chembur East, Mumbai

Dr. D.S Mehta, Professor & Head, Bapuji Dental College

Dr. C. S. Saimbi, Former Professor and Head, Dental College, Lucknow, Uttar Pradesh

Dr. Gururaja Rao, Ex-Principal, AME Dental College, Raichur, Karnataka

Dr. Tamal Kanti Pal, Professor & Head, Guru Nanak Institute of Dental Science & Research and Hospital, Kolkata

Dr. Vijay Kumar Chava, Prof and Head, Narayana Dental College, Nellore, AP.

Dr. Dilip G Naik, Former Dean, Professor, Manipal College of Dental Sciences, Mangalore

Dr. Shalu Bathla, Professor, M.M.College of Dental Sciences, Mullana, Ambala, Haryana.

Dr. Sreenivas Nagarakanti, Professor, Narayana Dental College, Nellore, AP.

Dr. Neeraj Deshpande, Professor, K.M.Shah Dental College, Vadodara ,Gujrat

### **Prosthodontics**

Dr. Hari Prakash- Director General at I.T.S Centre for Dental Studies & Research, Muradnagar,

Dr. Mahesh Verma, Director Principal, Maulana Azad Institute of Dental Sciences, New Delhi

Dr. Manu Rathi, Prof & Head, Post Graduate Institute Of Dental Sciences,Rohatak ,Haryana

Dr. Narendra Gupta, Professor, Baba Banarasidas Dental College, Lucknow

Dr. Himanshu Aeran, Director Principal, Seema Dental College & Hospital, Rishikesh

Dr. Dileepnag Vinnakota, Professor, Narayana Dental College, Nellore, AP.

Dr. Sushant Garg, Yamuna Institute of Dental Sciences, Yamuna Nagar, Haryana

Dr. Sanjay Bansal, Eklavya Dental College, Kotputli, Rajasthan

Dr. Umesh Pai, Manipal College of Dental Sciences, Mangalore, Karnataka

Dr. Meena Aras, Govt.Dental College, Goa.

**Oral Medicine**

Dr. Freny Karjodkar Prof & Head, Nair Hospital Dental College Mumbai

Dr. Jayachandran, Prof and Head, Govt Dental College, Chennai

Dr. Kannan Natarajan, Professor and Head, Narayana Dental College, Nellore, AP.

Dr. Rajendra Patil, Professor and Head, Kothiwal Dental College, Moradabad, UP

**Oral Pathology**

Dr. Dinesh Daftary, Former Scientist. Tata Institute of Fundamental Research, Mumbai

Dr. C.R. Ramchandran, Former Dean, RM Dental College, Annamalai Nagar,Chidambaram,TN.

Dr R.R Paul, Vice Principal, Senior Professor, Guru Nanak Institute of Dental Sciences & Research, Kolkata

Dr. V. K. Hazarey, Former Dean, Government Dental College, Nagpur.

Dr. Ranganathan, Professor and HOD, Ragas Dental College

Dr. B. Ajay Reginald, Principal, Narayana Dental College, Nellore, AP.

**Oral Surgery**

Dr. Neelima Malik, Vice Chancellor, Krishna University Karad

Dr. Mohan Baliga, Associate Dean & Professor of MCODS, Mangalore

Dr. Sripathi Rao BH, Dean, Yenepoya Dental College Manglore

Dr. Rajiv Borle, Vice Chancellor, Datta Meghe Deemeed University Wardha

Dr. Vidya Rattan, Professor PGI Chandigarh

Dr. L. Krishna Prasad, Principal and Dean, Sibar Institute of Dental Sciences, Guntur

Dr. Venkata Kishore Kumar Rayaduragam, Narayana Dental College, Nellore A P

Dr. Rajshekhar Gali, Professor, Narayana Dental College Nellore AP

**Orthodontics**

Dr. Ashok Utreja, Former Head Oral Health PGI , Chandigarh

Dr. Sadashiva Shetty, Principal Prof. and Head, Bapuji Dental College

Dr. O.P. Kharbanda, Prof. and Head, All India Institute and Medical Sciences, New Delhi.

Dr. Ashok Dhoble, Hon. Secretary General at Indian Dental Association (H.O.)

Dr. T. Samraj, Director, VMS Dental College, Lakshmipuram,, Gandhi Road, Salem

Dr. S. P. Singh, Prof of Orthodontics PGI Chandigarh

Dr. U. S. Krishna Nayak, Principal, A.B. Shetty Dental College Manglore

Dr. Vinay Dua, Director, Principal, Professor & Head, National Dental College & Hospital, Dera Bassi (Punjab)

**Biotechnology**

Prof. (Dr). Anil Sharma

Maharishi Markandeshwar (Deemed to be University),  
Mullana, Ambala (HR)

## The Bond Strength of Root Canal Filling after Calcium Hydroxide Removal with a Simple Apical Negative Pressure Kit in Oval-Shaped Root Canal

Leelapornpisid, Warat; Sastraruji, Thanapat; Louwakul, Phumisak; More

*Contemporary Clinical Dentistry*. 14(2):98-103, Apr-Jun 2023.

[Abstract](#) [Favorite](#) [PDF](#) [Permissions](#)

OPEN

## Comparative Evaluation of the Bonding Efficacy of Multimode Adhesive, Two-Step Self-Etch Adhesive, and a Total-Etch System to Pulpal Floor Dentin – An *In vitro* Study

Valsan, Dhanya; Bhaskaran, Sajeev; Mathew, Joy; More

*Contemporary Clinical Dentistry*. 14(2):104-108, Apr-Jun 2023.

[Abstract](#) [Favorite](#) [PDF](#) [Permissions](#)

OPEN

## Antimicrobial and Cytotoxic Activity of *Ocimum tenuiflorum* and *Stevia rebaudiana*-Mediated Silver Nanoparticles – An *In vitro* Study

Pandiyan, Indumathy; Arumugham, Meignana Indiran; Doraikannan, Sri Sakthi; More

*Contemporary Clinical Dentistry*. 14(2):109-114, Apr-Jun 2023.

[Abstract](#) [Favorite](#) [PDF](#) [Permissions](#)

OPEN

## Antifungal Efficacy of *Ocimum Basilicum* Essential Oil in Tissue Conditioner Against *Candida Albicans*: An *In vitro* Study

Rajali, Aiemeeza; Zain, Nurhayati Mohamad; Amran, Nurafiqah Aina; More

*Contemporary Clinical Dentistry*. 14(2):115-122, Apr-Jun 2023.

[Abstract](#) [Favorite](#) [PDF](#) [Permissions](#)

OPEN

## Sequelae of Unreplaced Tooth Loss in Orthodontic Patients: A Cross-Sectional Study in Najran, Kingdom of Saudi Arabia

Alyami, Bandar

*Contemporary Clinical Dentistry*. 14(2):123-127, Apr-Jun 2023.

[⊕ Abstract](#) [☆ Favorite](#) [📄 PDF](#) [© Permissions](#)

OPEN

---

## Comparative Evaluation of Smear Layer Removal Efficacy of Neem Leaf Extract, Propolis, and Orange Oil when used as Endodontic Irrigants: An *in vitro* Scanning Electron Microscopic Study

Setia, Ria; Bajaj, Nitika; Bholra, Meenu; More

*Contemporary Clinical Dentistry*. 14(2):128-134, Apr-Jun 2023.

[⊕ Abstract](#) [☆ Favorite](#) [📄 PDF](#) [© Permissions](#)

OPEN

## Clinical and Radiographic Evaluation of Locally Delivered Plant Stem Cells for Treatment of Periodontitis: Randomized Clinical Trial

Elboraey, Mohamed Omar; Sabra, Reda Saber; Gamal, Sherouk Mohamed Mohamed

*Contemporary Clinical Dentistry*. 14(2):135-140, Apr-Jun 2023.

[⊕ Abstract](#)   [☆ Favorite](#)   [📄 PDF](#)   [© Permissions](#)

OPEN

---

## Comparative Evaluation of Antimicrobial Efficacy of Fluoride-Based and Self-Assembling Peptide P<sub>11</sub>-4-based Tooth Remineralization Agents on *Streptococcus mutans*: A Microbiological Study

Gayas, Zaina; Azher, Umme; Paul, Santhosh T.; More

*Contemporary Clinical Dentistry*. 14(2):141-144, Apr-Jun 2023.

[⊕ Abstract](#)   [☆ Favorite](#)   [📄 PDF](#)   [© Permissions](#)

OPEN

## The Antibacterial and Antibiofilm Effect of Amoxicillin and *Mangifera indica* L. Leaves Extract on Oral Pathogens

Soesanto, Sheila; Hepziba, Evangelista Rachel; Yasnill, ; More

*Contemporary Clinical Dentistry*. 14(2):145-151, Apr-Jun 2023.

[⊕ Abstract](#)   [☆ Favorite](#)   [📄 PDF](#)   [© Permissions](#)

OPEN

---

## Association of Serum Interleukin-10 Level with Glycemic Status to Predict Glycemic Alteration with Periodontitis: A Cross-Sectional, Observational Study

Ikbal, S. K. Aziz; Gupta, Sanjay; Tiwari, Vandana; More

*Contemporary Clinical Dentistry*. 14(2):152-156, Apr-Jun 2023.


[⊕ Abstract](#)   [☆ Favorite](#)   [📄 PDF](#)   [© Permissions](#)

OPEN




## Contemporary Clinical Dentistry

<b>COUNTRY</b> India Universities and research institutions in India Media Ranking in India	<b>SUBJECT AREA AND CATEGORY</b> Dentistry - Oral Surgery - Orthodontics - Periodontics	<b>PUBLISHER</b> Wolters Kluwer   Madknow Publications	<b>H-INDEX</b> <h1>34</h1>
<b>PUBLICATION TYPE</b> Journals	<b>ISSN</b> 09762361, 0976237X	<b>COVERAGE</b> 2012-2022	<b>INFORMATION</b> <a href="#">Homepage</a> <a href="#">How to publish in this journal</a> <a href="mailto:journacod@gmail.com">journacod@gmail.com</a>



**FIND SIMILAR JOURNALS**

<b>1</b> Journal of Contemporary Dental Practice USA <b>85%</b> similarity	<b>2</b> Saudi Dental Journal SAU <b>85%</b> similarity	<b>3</b> World Journal of Dentistry IND <b>85%</b> similarity	<b>4</b> Dental Research Journal IRN <b>84%</b> similarity
---	--	--	---



# The Antibacterial and Antibiofilm Effect of Amoxicillin and *Mangifera indica* L. Leaves Extract on Oral Pathogens

## Abstract

**Objective:** This study aimed to determine the antibacterial and antibiofilm effects of amoxicillin combined with extract of *Mangifera indica* L. leaves against *Staphylococcus aureus* and *Porphyromonas gingivalis*. **Materials and Methods:** This was an experimental laboratory *in vitro* study with a posttest-only control group design. An antibacterial test using the plate count method and an antibiofilm test using the microtiter plate biofilm assay method were conducted. The research samples comprised extract of *M. indica* L. leaves with a concentration of 100%; amoxicillin and extract concentrations of 3.125%, 6.25%, 12.5%, 25%, 50%, and 100%; and amoxicillin. Dimethyl sulfoxide served as a negative control and co-amoxiclav served as a positive control. **Results:** The combination of amoxicillin and the extract exhibited an antibacterial effect against *S. aureus* at a concentration of 12.5% and higher and more effective than co-amoxiclav *P. gingivalis* at a concentration of 3.125% and higher. In the antibiofilm test, the combination of amoxicillin and the extract at a concentration of 25% after 1 h of incubation and a concentration of 6.25% after 3 h of incubation inhibited *S. aureus*. The inhibition of *S. aureus* biofilms at a concentration of 100% after 24 h of incubation was as effective as that of co-amoxiclav. The extract at a concentration of 25% over the entire incubation period showed more potent inhibition against the *P. gingivalis* biofilm than co-amoxiclav. **Conclusions:** The ethanolic extract of *M. indica* L. leaves and the combination of amoxicillin and the extract have the potential to inhibit the growth and formation of *S. aureus* and *P. gingivalis* biofilms.

**Keywords:** Amoxicillin, antibiofilm, arumanis mango, ethanol extract of *Mangifera indica* L. leaves

## Introduction

Based on data from Basic Health Research (Riskesdas) in 2018, Indonesia's dental and oral health problems reached 57.6% of overall health-care problems.<sup>[1]</sup> The latter is influenced by poor oral hygiene, which triggers various diseases, such as dentoalveolar abscesses and periodontitis.<sup>[2,3]</sup> A dentoalveolar abscess is a pathological cavity in the oral cavity that contains pus due to secondary infection caused by caries, trauma, failure of root canal treatment, and poor oral hygiene.<sup>[3]</sup> Periodontitis is a chronic inflammation of the periodontal tissue structure, including gingiva, bone, and the periodontal ligament, which can cause pocket formation, recession, tooth mobility, or tooth loss.<sup>[2]</sup> Dentoalveolar abscesses and periodontitis are associated with bacterial pathogens in biofilms.<sup>[3,4]</sup>

A biofilm is a collection of microbial cells, especially bacteria, attached to the tooth

surface and coated by an extracellular polymeric substance. This coating protects cells and enables accelerated growth rates, along with additional horizontal gene transfer between cells within the coating, which promotes additional problems, such as antibiotic resistance.<sup>[5]</sup> The formation of biofilms in the oral cavity involves complex competition between microflora for initial attachment.<sup>[6]</sup> *Staphylococcus aureus*, a Gram-positive bacterium, initiates adherence in biofilm formation and produces multiple layers of biofilm embedded in a glycocalyx layer.<sup>[7,8]</sup> *S. aureus* has several virulence factors (capsules, adhesins, coagulase, hyaluronidase, staphylokinase, enterotoxin, and leucocidin) that support biofilm formation in a dentoalveolar abscess.<sup>[6]</sup> *Porphyromonas gingivalis*, which belongs to anaerobic Gram-negative bacteria, is the second most common colonizing bacteria in biofilms. *P. gingivalis* virulence factors, such as lipopolysaccharides, capsules, fimbriae, outer membrane proteins, proteases, and enzymes, induce the destruction of periodontal tissue, causing periodontitis.<sup>[9]</sup>

**How to cite this article:** Soesanto S, Hepziba ER, Yasnill, Widyarman AS. The antibacterial and antibiofilm effect of amoxicillin and *Mangifera indica* L. leaves extract on oral pathogens. *Contemp Clin Dent* 2023;14:145-51.

Sheila Soesanto<sup>1</sup>,  
Evangelista Rachel  
Hepziba<sup>2</sup>,  
Yasnill<sup>2</sup>,  
Armelia Sari  
Widyarman<sup>3</sup>

<sup>1</sup>Department of Pharmacology,  
Faculty of Dentistry, Universitas  
Trisakti, Jakarta, Indonesia,  
<sup>2</sup>Undergraduate Student,  
Faculty of Dentistry, Universitas  
Trisakti, Jakarta, Indonesia,  
<sup>3</sup>Department of Microbiology,  
Faculty of Dentistry, Universitas  
Trisakti, Jakarta, Indonesia

Submitted : 26-Jul-2022  
Revised : 06-Jan-2023  
Accepted : 21-Feb-2023  
Published : 30-Jun-2023

**Address for correspondence:**  
Dr. Armelia Sari Widyarman,  
Department of Microbiology,  
Faculty of Dentistry, Universitas  
Trisakti, Kyai Tapa 260, Grogol,  
Jakarta 11440, Indonesia.  
E-mail: armeliasari@trisakti.  
ac.id

Access this article online

Website:  
<https://journals.lww.com/cccd>

DOI: 10.4103/ccd.ccd\_399\_22

Quick Response Code:



This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow\_reprints@wolterskluwer.com

Amoxicillin from the penicillin group is often used to treat dentoalveolar abscesses and periodontitis. Amoxicillin is a broad-spectrum antibiotic that works by binding to penicillin-binding proteins in Gram-positive and Gram-negative bacteria and inhibiting the transpeptidation process in bacteria.<sup>[3,10]</sup> Overuse of amoxicillin can lead to side effects, such as hypersensitivity reactions, nausea, vomiting, diarrhea, thrombocytopenia, dermatological disorders, and resistance.<sup>[11]</sup> Resistance to amoxicillin can also occur due to the destruction of the  $\beta$ -lactam ring by  $\beta$ -lactamase enzymes produced by *S. aureus* and *P. gingivalis*.<sup>[2,12]</sup> Therefore, amoxicillin is often combined with clavulanic acid, known as co-amoxiclav, to reduce resistance.<sup>[13]</sup> Clavulanic acid is a  $\beta$ -lactamase inhibitor that works by inactivating the pathogen's  $\beta$ -lactamase, thereby increasing the antibacterial activity of amoxicillin.<sup>[11]</sup> However, the use of co-amoxiclav can cause side effects, such as itching, redness around the mouth, diarrhea, nausea, and idiosyncratic drug-induced liver injury.<sup>[11,13]</sup> Alternative medicines using herbal plants have the potential to minimize these side effects.<sup>[14]</sup>

Arumanis mango (*Mangifera indica* L) is a herbal plant from India cultivated in various tropical and subtropical regions, including Indonesia.<sup>[15,16]</sup> Indonesians cultivate Arumanis mangos for their sweetness, freshness, and fragrance. The fruit contains Vitamins A, B, and C, which are beneficial for health.<sup>[17]</sup> In addition, the seeds, skin, roots, and leaves of *M. indica* L. have various properties in traditional medicine. *M. indica* L. leaves are generally discarded and considered waste, even though these leaves contain secondary metabolites, including phenolics (mangiferin, tannins, and flavonoids), alkaloids, saponins, terpenoids, glycosides, and steroids that have potential antibacterial, antifungal, antiviral, antiparasitic, antioxidant, anti-inflammatory, antitumor, anticancer, and analgesic effects.<sup>[18]</sup>

Previous studies showed that *M. indica* L. leaf extracts reduced the number of *Streptococcus mutans* and improved the antibacterial effect of clindamycin against *S. aureus*.<sup>[19,20]</sup> However, no research has investigated the potential of combining amoxicillin with *M. indica* L. leaf extracts in combating *S. mutans* and *P. gingivalis*. Thus, to address this research gap, this study aimed to determine the effectiveness of amoxicillin and ethanolic extract of Arumanis mango (*M. indica* L.) in inhibiting the growth and formation of *S. aureus* and *P. gingivalis* biofilms.

## Materials and Methods

This experimental *in vitro* study was performed at the Microbiology Center of Research and Education (MiCORE) Laboratory, Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia. An ethanolic extract of Arumanis mango (*M. indica* L.) leaves was prepared at the Research Institute for Spices and Medicinal Plants (BALITTRO)

in Bogor, West Java, Indonesia. The test solutions used were 10% dimethyl sulfoxide (DMSO) (negative control), co-amoxiclav (positive control), ethanolic extract of *M. indica* L. leaves with a concentration of 100%, and a combination of amoxicillin with extract concentrations of 3.125%, 6.25%, 12.5%, 25%, 50%, and 100%.

The sample size was calculated using Federer's formula, with n equals number of repetition, and t equals as total test group (8 groups comprised 6 treatment groups, 1 positive control, and 1 negative control). Based on this formula, each treatment group was conducted with four-time repetition for all assays.

### Ethanolic extract of *Mangifera indica* L. leaves

The ethanolic extract of *M. indica* L. leaves was prepared using the maceration method. The mango leaves were washed, dried, and mashed, and the simplicial was then soaked in 70% ethanol at a ratio of 1:5. The maceration process was performed for 2–3 h, and the macerate was then allowed to stand for 24 h and filtered. The extract was diluted with 10% DMSO solution to obtain extract concentrations of 50%, 25%, 12.5%, 6.25%, and 3.125%.

### Preparation of antibiotic solution

Amoxicillin and co-amoxiclav solution were prepared by crushing 500 mg of amoxicillin and 625 mg of co-amoxiclav until smooth, using a mortar and pestle. In total, 1.2 mg of amoxicillin and 1.5 mg of co-amoxiclav were each mixed with 6 ml of sterile distilled water until homogeneous to obtain 200 g/ $\mu$ l of amoxicillin and 250 g/ $\mu$ l of co-amoxiclav.

### Phytochemical tests

Qualitative phytochemical tests were performed at BALITTRO to identify secondary metabolites, such as alkaloids, saponins, tannins, phenolics, flavonoids, triterpenoids, steroids, and glycosides, in the ethanolic extract of *M. indica* L. leaves.

### Bacterial culture

*S. aureus* ATCC 25923 and *P. gingivalis* ATCC 33277 were obtained from the MiCORE Laboratory, Faculty of Dentistry, Trisakti University. *S. aureus* was cultured on brain heart infusion broth medium (Sigma Aldrich, St. Louis, Missouri). *P. gingivalis* was cultured on Tryptone Soya Broth (Sigma Aldrich, St. Louis, Missouri) medium enriched with hemin (5 mg/l), Vitamin K (10 mg/l), 0.5% yeast extract, and L-cystine (400 mg/l).<sup>[21]</sup> The medium was incubated in an anaerobic jar (Oxoid, Basingstoke, and Hampshire) at 37° for 24 h under anaerobic conditions. Following the incubation, the bacterial absorbance value was standardized into McFarland standard of 0.5, approximately  $1.5 \times 10^8$  colony forming unit (CFU)/mL (optical density OD<sub>600</sub>  $\pm$  0.132), before the following assays.

### Microdilution and total plate count

To evaluate the antibacterial properties of the extract combined with amoxicillin, an antibacterial test was performed on the plate count by the microdilution method. In total, 100 µL of cultured *S. aureus* ATCC 25923 and *P. gingivalis* ATCC 33277 were distributed into 96-well plates (Nest Biotech, Jiangsu, China). A test solution of 100 µl was added to each well and incubated at 37°C for 24 h under anaerobic conditions. After incubation, each mixture containing treated bacteria was diluted 10<sup>5</sup> times. Five microliters of diluted mixture were then spread on a petri dish containing sterile brain heart infusion agar media. The growth of bacterial colonies was calculated after incubation at 37°C for 24 h.

### Microtiter plate biofilm assay

The bacterial cultures were inserted into each well of the 96-well plates and then incubated at 37°C for 24 h under anaerobic conditions. The supernatant was discarded, leaving a thin layer on the surface of the well. The wells were rinsed using phosphate-buffered saline (PBS) (Biomatrix, Ontario, Canada). Each test solution (200 µl) was added to each well and then incubated for 1, 3, and 24 h at 37°C. The wells were rinsed twice with PBS and fixated over burning spirit lamp. Crystal violet (Merck, Darmstadt, Germany) (200 µl) was then added to each well and allowed to stand for 15 min, followed by rinsing twice and standing for 15 min. In the past step, 200 µl of 96% ethanol was added. The OD was measured using a microplate reader (Safas, Monaco) with a wavelength of 490 nm.

### Statistical analysis

The research data were processed using the Statistical Package for the Social Sciences (SPSS) computer program, version 26 (IBM, Armonk, NY, USA). The Shapiro–Wilk method was used to test the normality of the data. If the data were normally distributed ( $P > 0.05$ ), a one-way analysis of variance test was conducted, followed by Turkey's honestly significant difference test (significance level of  $P < 0.05$ ) to verify the significance between the groups.

## Results

### Phytochemical screening

The results of the phytochemical screening qualitatively proved that the ethanolic extract of *M. indica* L. leaves contains alkaloids, saponins, tannins, phenolics, flavonoids, and steroids [Table 1].

### Antibacterial test using microdilution and plate count methods

The antibacterial test results showed that the ethanolic extract of *M. indica* L. leaves at 100% concentration and the combination of amoxicillin with extracts of various

**Table 1: The results of phytochemical screening of the ethanol extract of *Mangifera indica* L. leaves**

Secondary metabolites	Screening result
Alkaloids	+
Saponins	+
Tannins	+
Phenolics	+
Flavonoids	+
Triterpenoids	–
Steroids	+
Glycoside	–

concentrations exhibited antibacterial and antibiofilm effects against *S. aureus* and *P. gingivalis* [Figure 1]. In terms of inhibition of *S. aureus*, the results obtained by the combination of amoxicillin and the extract were not significantly different from those obtained using the positive control ( $P > 0.05$ ). The extract at a concentration of 12.5% and higher was effective against *S. aureus*, and the extract at a concentration of 3.125% and higher was effective against *P. gingivalis* [Figures 2 and 3].

### Antibiofilm test using the microtiter plate biofilm assay method

The results of the antibiofilm test showed that the addition of extracts to amoxicillin in inhibiting *S. aureus* biofilms had a significantly lower OD value compared to the positive control ( $P < 0.05$ ), starting at a concentration of 25% after 1 h of incubation [Figure 4] and at a concentration of 6.25% after 3 h of incubation [Figure 5]. After 24 h of incubation, following OD measurement, the group with the combination of extract concentration of 100% and amoxicillin proved not to be significantly different from OD of the positive control ( $P > 0.05$ ) [Figure 6]. In the *P. gingivalis* biofilm, the OD values of the addition of amoxicillin and the extract group, starting at a concentration of 25% after 1, 3, and 24 h of incubation, were smaller than the OD values of the positive control amoxiclav. The OD values of the treatment group were significantly different from those of the positive control amoxiclav [Figures 7-9].

## Discussion

As shown by our results, the ethanol extract of *M. indica* L. leaves contains secondary metabolites, including alkaloids, saponins, tannins, phenolics, flavonoids, and steroids. Different mechanisms of action of each compound account for the antibacterial and antibiofilm properties of the extract. Alkaloids inhibit the formation of peptidoglycan in bacterial cells. The alkaloids can disrupt the amino acid structure of bacterial DNA, leading to bacterial lysis.<sup>[22]</sup> Saponins damage the cell membrane and cell wall permeability in the diffusion process, resulting in the release of enzymes, amino acids, nutrients, and water, leading to cell destabilization and cell death.<sup>[23]</sup> Tannins form a complex with protein in the cell wall, namely,




Figure 1: The results of the inhibition test of *Staphylococcus aureus* and *Porphyromonas gingivalis* using the plate count method




Figure 2: Graph of the average total colony of *Staphylococcus aureus* (\* $P < 0.05$ , \*\* $P < 0.01$ )




Figure 3: Graph of the average total colony of *Porphyromonas gingivalis* (\* $P < 0.05$ , \*\* $P < 0.01$ )

proline, which can damage the cell wall.<sup>[22]</sup> The most abundant phenolic compound in *M. indica* L leaves is mangiferin. Mangiferin belongs to the xanthone C-glucosyl group, which can damage cell structure and cell membranes and inhibit bacterial protein synthesis.<sup>[24]</sup> Previous research reported that mangiferin compounds interfere with the mechanism of drug resistance; thus, restoring

bactericidal and bacteriostatic effects of nalidixic acid, ampicillin, tetracycline, and sulfamethoxazole/trimethoprim.<sup>[25]</sup> However, the exact mechanism of how mangiferin interferes with the mechanism of drug resistance is yet to be elucidated. Flavonoids, as antibacterial, play a role in disrupting the activity of cell wall formation by suppressing cytoplasm function, interrupting the nutrient exchange process, and thereby inhibiting the energy supply of bacteria.<sup>[26]</sup> In addition, flavonoids inhibit enzymes from producing quorum-sensing signals, thereby disrupting the communication process between cells during biofilm formation.<sup>[27]</sup> Steroids can cause leakage of lysosomes and membrane phospholipids that reduce the integrity of cell membranes and lead to cell lysis.<sup>[28]</sup>

Thus far, only a few studies have focused on antibacterial and antibiofilm properties of ethanolic extract of *M. indica* L. leaves.<sup>[19,20]</sup> The concentration of the extract used in this study ranged from 3.125% to 100%. As a positive control, co-amoxiclav, a combination of amoxicillin and clavulanic acid, was used. Amoxicillin, a  $\beta$ -lactam antibiotic, works by inhibiting the synthesis of bacterial cell walls.<sup>[11]</sup> The release of  $\beta$ -lactamase enzymes by *S. aureus* and *P. gingivalis* decrease the antibacterial effect of amoxicillin.<sup>[2,12]</sup> The addition of clavulanic acid binds to  $\beta$ -lactamase enzymes from bacteria, inhibiting the enzyme thereby unable to cleave  $\beta$ -lactam ring in amoxicillin, so the amoxicillin can still exhibit its antibacterial activity.<sup>[11]</sup>

In the microdilution and plate count tests, the addition of the ethanolic extract of *M. indica* L. leaves to amoxicillin inhibited the growth of *S. aureus* and *P. gingivalis*, and the combination of the ethanolic extract and amoxicillin was as effective as that of co-amoxiclav. In terms of antibacterial activity, a combination of ethanolic extract at a concentration of 12.5% and amoxicillin was as effective as co-amoxiclav against *S. aureus*, and a concentration as low as 3.125% was as effective as co-amoxiclav against *P. gingivalis*. Therefore, ethanolic extract of *M. indica* L. leaves may be a potential  $\beta$ -lactamase inhibitor equivalent to clavulanic acid. The results of this study are in line with the research of Hartanto *et al.*, who showed that adding




Figure 4: Graph of the average OD value of *Staphylococcus aureus* biofilm after 1 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). OD: Optical density




Figure 5: Graph of the average OD value of *Staphylococcus aureus* biofilm after 3 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). OD: Optical density




Figure 6: Graph of the average OD value of *Staphylococcus aureus* biofilm after 24 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). OD: Optical density




Figure 7: Graph of the average OD value of *Porphyromonas gingivalis* biofilm after 1 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). OD: Optical density




Figure 8: Graph of the average OD value of *Porphyromonas gingivalis* biofilm after 3 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). OD: Optical density




Figure 9: Graph of the average OD value of *Porphyromonas gingivalis* biofilm after 24 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). OD: Optical density

methanolic extract of *M. indica* L. leaves to clindamycin has antibacterial effects against *S. aureus*, especially at a concentration of 100%.<sup>[19]</sup> The bioactive component of *M. indica* L. leaves, namely, mangiferin, is known to have a synergistic effect on tetracycline, ampicillin, nalidixic acid, and trimethoprim in inhibiting the growth of *S. aureus*.<sup>[25]</sup>

The biofilm formation phase begins with pellicle formation, which occurs in the first few seconds to the first min from

the initial contact, with the initial adhesion phase occurring 2–4 h later. After 24 h, the biofilm enters the maturation phase, becoming 1.000 - 1.500 times more resistant than planktonic bacteria.<sup>[29]</sup> The incubation time used in the antibiofilm test in this study was adjusted to the stage of biofilm formation, namely 1, 3, and 24 h. This timing aimed to determine at which stage of biofilm formation amoxicillin and the ethanolic extract of *M. indica* L. leaves

most effectively inhibited *S. aureus* and *P. gingivalis*. In the antibiofilm test, the OD from the combination of amoxicillin with ethanolic extract of *M. indica* L. leaves starting at a concentration of 25% after 1 h of incubation and 6.25% after 3 h of incubation had a significantly lower OD value compared to co-amoxiclav. This result proved that the ethanolic extract of *M. indica* L. leaves at a concentration of 25% after 1 h of incubation and 6.25% after 3 h of incubation was more effective at inhibiting *S. aureus* biofilms than co-amoxiclav. The OD values for the combination of amoxicillin and the extract at a concentration of 100% after 24 h of incubation were lower than those obtained for co-amoxiclav, but the difference was not statistically significant. Therefore, the combination of amoxicillin and an extract concentration of 100% has antibiofilm properties equivalent to those of co-amoxiclav after 24 h of incubation.

In terms of inhibiting *P. gingivalis* biofilm, the extract at a concentration of 25% and higher during the entire incubation period showed a more potent antibiofilm effect than that of co-amoxiclav. The combination of amoxicillin and the ethanolic extract of *M. indica* L. leaves inhibited the formation of *S. aureus* biofilms at a concentration of 6.25% after 3 h of incubation and *P. gingivalis* biofilms at a concentration of 25% after 1 h of incubation. This study supports the findings of previous research, which reported that ethanolic extract of *M. indica* L. leaves reduced *S. aureus* attachment and the number of *S. aureus* biofilms.<sup>[30]</sup>

This study has several limitations. First, the ethanolic extract used in this study was a crude extract. The use of a more refined extract, such as an extract exposed to extraction chromatography, would have resulted in an extract with fewer impurities. Second, only two of the many known oral pathogens were used in this study. Other oral pathogens can also be tested to expand the antibacterial activity of amoxicillin combined with *M. indica* L. leaves ethanolic extract. Further research is needed to determine the toxicity of this combination. Preclinical and clinical tests should also be conducted before the combination can be used as an alternative treatment for dentoalveolar abscesses and periodontitis.

## Conclusions

Ethanolic extract of *M. indica* L. leaves contains alkaloids, saponins, tannins, phenolics, flavonoids, and steroids that have the potential to inhibit the growth and formation of *S. aureus* and *P. gingivalis* biofilms *in vitro*. Within the limitations of this preliminary study, we conclude that the addition of ethanolic extract of *M. indica* L. leaves to amoxicillin could potentially increase the antibacterial and antibiofilm properties of amoxicillin against *S. aureus* and *P. gingivalis*.

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## References

1. Ministry of Health of the Republic of Indonesia. Report on Result of National Basic Health Research 2018. Jakarta: Ministry of Health; 2018. p. 197-207.
2. Japoni A, Vasin A, Noushadi S, Kiany F, Japoni S, Alborzi A. Antibacterial susceptibility patterns of *Porphyromonas gingivalis* isolated from chronic periodontitis patients. *Med Oral Patol Oral Cir Bucal* 2011;16:e1031-5.
3. Shweta N, Prakash SK. Dental abscess: A microbiological review. *Dent Res J (Isfahan)* 2013;10:585-91.
4. Mehrotra N, Singh S. Periodontitis. Treasure Island (FL): StatPearls Publishing; 2011.
5. Archer NK, Mazaitis MJ, Costerton JW, Leid JG, Powers ME, Shirtliff ME. *Staphylococcus aureus* biofilms: Properties, regulation, and roles in human disease. *Virulence* 2011;2:445-59.
6. Huang R, Li M, Gregory RL. Bacterial interactions in dental biofilm. *Virulence* 2011;2:435-44.
7. Periasamy S, Joo HS, Duong AC, Bach TH, Tan VY, Chatterjee SS, *et al.* How *Staphylococcus aureus* biofilms develop their characteristic structure. *Proc Natl Acad Sci U S A* 2012;109:1281-6.
8. Taylor TA, Unakal CG. *Staphylococcus aureus*. Treasure Island (FL): StatPearls Publishing; 2021.
9. How KY, Song KP, Chan KG. *Porphyromonas gingivalis*: An overview of periodontopathic pathogen below the gum line. *Front Microbiol* 2016;7:53.
10. Ciancio SG, Mariotti AJ. Systemic Anti-infective therapy for periodontal diseases. In: Carranza F, Newman M, Takei H, Klokkevold P, editors. Newman and Carranza's Clinical Periodontology. 13<sup>th</sup> ed. Philadelphia: Elsevier; 2019. p. 555-7.
11. Akhavan B, Khanna N, Vijhani P. Amoxicillin. Treasure Island (FL): StatPearls Publishing; 2020.
12. Foster TJ, Geoghegan JA. *Staphylococcus aureus*. In: Molecular Medical Microbiology. Massachusetts: Elsevier; 2015. p. 655-74.
13. Samaranayake L. Essential Microbiology for Dentistry. 5<sup>th</sup> ed. Poland: Elsevier Ltd; 2018. p. 75, 158, 293, 297.
14. Joshua M, Takudzwa M. Antibacterial properties of *Mangifera indica* on *Staphylococcus aureus*. *Afr J Clin Exp Microbiol* 2013;14:62-74.
15. Kalita P. An overview on *Mangifera indica*: Importance and its various pharmacological action. *Pharma Tutoring* 2014;2:72-6.
16. Sivakumar D, Jiang Y, Yahia EM. Maintaining mango (*Mangifera indica* L.) fruit quality during the export chain. *Food Res Int* 2011;44:1254-63.
17. Chiumarelli M, Ferrari CC, Sarantopoulos CI, Hubinger MD. Fresh cut *Tommy Atkins* mango pre-treated with citric acid and coated with cassava (*Manihot esculenta* Crantz) starch or sodium alginate. *Innov Food Sci Emerg Technol* 2011;12:381-7.
18. Gu C, Yang M, Zhou Z, Khan A, Cao J, Cheng G. Purification and characterization of four benzophenone derivatives from *Mangifera indica* L. leaves and their antioxidant, immunosuppressive and  $\alpha$ -glucosidase inhibitory activities. *J Funct Foods* 2019;52:709-14.
19. Hartanto R, Tran V, Khang G, Pham T, Trinh T, Denhara C. Effect of the addition of arumanis mango leaf extract (*Mangifera indica* L.) on clindamycin antibiotics in inhibiting the growth of *Staphylococcus aureus*. *J Prima Med Sains* 2020;2:14-7.

20. Kurniasih R. Effect of Concentrations of Young Mango Arumanis Ethanolic Extract (*Mangifera indica* L.) leaves on growth inhibition of *Streptococcus mutans* *in vitro*. Surakarta: Fakultas Kedokteran Gigi Universitas Muhammadiyah (Faculty of Dentistry, Muhammadiyah University); 2016.
21. Yamanaka T, Furukawa T, Matsumoto-Mashimo C, Yamane K, Sugimori C, Nambu T, *et al.* Gene expression profile and pathogenicity of biofilm-forming *Prevotella intermedia* strain 17. *BMC Microbiol* 2009;9:11.
22. Sylvana D, Amir M, Purnamasari CB, Iskandar A, Asfirizal V. Antibacterial activity of ethanol extract of beluntas leaves on *Streptococcus mutans*, *Porphyromonas gingivalis*, and *Enterococcus faecalis*. *Padjadjaran J Dent* 2021;33:191-8.
23. Sebastian J, Widyarman AS. Roselle flower petals extract inhibits periodontal pathogenic biofilms. *J Dentomaxillofacial Sci* 2021;6:102-5.
24. Tirado-Kulieva V, Atoche-Dioses S, Hernandez-Martinez E. Phenolic compounds of mango (*Mangifera indica*) by-products: Antioxidant and antimicrobial potential, use in disease prevention and food industry, methods of extraction, and microencapsulation. *Sci Agropecu* 2021;12:283-93.
25. Mazlan NA, Azman S, Ghazali NF, Zarith P, Yusri S. Synergistic antibacterial activity of mangiferin with antibiotics against *Staphylococcus aureus*. *Drug Invent Today* 2019;12:14-7.
26. Ma Y, Ding S, Fei Y, Liu G, Jang H, Fang J. Antimicrobial activity of anthocyanins and catechins against foodborne pathogens *Escherichia coli* and *Salmonella*. *Food Control* 2019;106:78-80.
27. Federika AS, Rukmo M, Setyabudi S. Antibiofilm activity of flavonoid mangosteen pericarp extract against *Porphyromonas gingivalis* bacteria. *Conserv Dent J* 2020;10:27-30.
28. Madduluri S, Rao KB, Sitaram B. *In vitro* evaluation of antibacterial activity of five indigenous plants extracts against five bacteria pathogens of humans. *Int J Pharm Pharm Sci* 2013;5:679-84.
29. Bjarnsholt T. The role of bacterial biofilms in chronic infections. *APMIS Suppl* 2013;121:1-51.
30. Manzur AG, Sm Junior V, Morais-Costa F, Mariano EG, Careli RT, da Silva LM, *et al.* Extract of *Mangifera indica* L. leaves may reduce biofilms of *Staphylococcus* spp. in stainless steel and teatcup rubbers. *Food Sci Technol Int* 2020;26:11-20.



# The Antibacterial and Antibiofilm Effect of Amoxicillin and *Mangifera indica* L. Leaves Extract on Oral Pathogens

*by* Sheila Soesanto FKG

---

**Submission date:** 22-Feb-2024 01:24PM (UTC+0700)

**Submission ID:** 2301393628

**File name:** CCD-14-145.pdf (2.12M)

**Word count:** 4635

**Character count:** 25635

# The Antibacterial and Antibiofilm Effect of Amoxicillin and *Mangifera indica* L. Leaves Extract on Oral Pathogens

## Abstract

**Objective:** This study aimed to determine the antibacterial and antibiofilm effects of amoxicillin combined with extract of *Mangifera indica* L. leaves against *Staphylococcus aureus* and *Porphyromonas gingivalis*. **Materials and Methods:** This was an experimental laboratory *in vitro* study with a posttest-only control group design. An antibacterial test using the plate count method and an antibiofilm test using the microtiter plate biofilm assay method were conducted. The research samples comprised extract of *M. indica* L. leaves with a concentration of 100%; amoxicillin and extract concentrations of 3.125%, 6.25%, 12.5%, 25%, 50%, and 100%; and amoxicillin. Dimethyl sulfoxide served as a negative control and co-amoxiclav served as a positive control. **Results:** The combination of amoxicillin and the extract exhibited an antibacterial effect against *S. aureus* at a concentration of 12.5% and higher and more effective than co-amoxiclav *P. gingivalis* at a concentration of 3.125% and higher. In the antibiofilm test, the combination of amoxicillin and the extract at a concentration of 25% after 1 h of incubation and a concentration of 6.25% after 3 h of incubation inhibited *S. aureus*. The inhibition of *S. aureus* biofilms at a concentration of 100% after 24 h of incubation was as effective as that of co-amoxiclav. The extract at a concentration of 25% over the entire incubation period showed more potent inhibition against the *P. gingivalis* biofilm than co-amoxiclav. **Conclusions:** The ethanolic extract of *M. indica* L. leaves and the combination of amoxicillin and the extract have the potential to inhibit the growth and formation of *S. aureus* and *P. gingivalis* biofilms.

**Keywords:** Amoxicillin, antibiofilm, arumanis mango, ethanol extract of *Mangifera indica* L. leaves

## Introduction

Based on data from Basic Health Research (Riskesdas) in 2018, Indonesia's dental and oral health problems reached 57.6% of overall health-care problems.<sup>[1]</sup> The latter is influenced by poor oral hygiene, which triggers various diseases, such as dentoalveolar abscesses and periodontitis.<sup>[2,3]</sup> A dentoalveolar abscess is a pathological cavity in the oral cavity that contains pus due to secondary infection caused by caries, trauma, failure of root canal treatment, and poor oral hygiene.<sup>[3]</sup> Periodontitis is a chronic inflammation of the periodontal tissue structure, including gingiva, bone, and the periodontal ligament, which can cause pocket formation, recession, tooth mobility, or tooth loss.<sup>[2]</sup> Dentoalveolar abscesses and periodontitis are associated with bacterial pathogens in biofilms.<sup>[3,4]</sup>

A biofilm is a collection of microbial cells, especially bacteria, attached to the tooth

surface and coated by an extracellular polymeric substance. This coating protects cells and enables accelerated growth rates, along with additional horizontal gene transfer between cells within the coating, which promotes additional problems, such as antibiotic resistance.<sup>[5]</sup> The formation of biofilms in the oral cavity involves complex competition between microflora for initial attachment.<sup>[6]</sup> *Staphylococcus aureus*, a Gram-positive bacterium, initiates adherence in biofilm formation and produces multiple layers of biofilm embedded in a glycocalyx layer.<sup>[7,8]</sup> *S. aureus* has several virulence factors (capsules, adhesins, coagulase, hyaluronidase, staphylokinase, enterotoxin, and leucocidin) that support biofilm formation in a dentoalveolar abscess.<sup>[6]</sup> *Porphyromonas gingivalis*, which belongs to anaerobic Gram-negative bacteria, is the second most common colonizing bacteria in biofilms. *P. gingivalis* virulence factors, such as lipopolysaccharides, capsules, fimbriae, outer membrane proteins, proteases, and enzymes, induce the destruction of periodontal tissue, causing periodontitis.<sup>[9]</sup>

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow\_reprints@wolterskluwer.com

**How to cite this article:** Soesanto S, Hepziba ER, Yasnill, Widyarman AS. The antibacterial and antibiofilm effect of amoxicillin and *Mangifera indica* L. leaves extract on oral pathogens. *Contemp Clin Dent* 2023;14:145-51.

Sheila Soesanto<sup>1</sup>,  
Evangelista Rachel  
Hepziba<sup>2</sup>,  
Yasnill<sup>2</sup>,  
Armelia Sari  
Widyarman<sup>3</sup>

<sup>1</sup>Department of Pharmacology,  
Faculty of Dentistry, Universitas  
Trisakti, Jakarta, Indonesia,  
<sup>2</sup>Undergraduate Student,  
Faculty of Dentistry, Universitas  
Trisakti, Jakarta, Indonesia,  
<sup>3</sup>Department of Microbiology,  
Faculty of Dentistry, Universitas  
Trisakti, Jakarta, Indonesia

Submitted : 26-Jul-2022  
Revised : 06-Jan-2023  
Accepted : 21-Feb-2023  
Published : 30-Jun-2023

**Address for correspondence:**  
Dr. Armelia Sari Widyarman,  
Department of Microbiology,  
Faculty of Dentistry, Universitas  
Trisakti, Kyai Tapa 260, Grogol,  
Jakarta 11440, Indonesia.  
E-mail: armeliasari@trisakti.  
ac.id

Access this article online

**Website:**  
<https://journals.lww.com/ccod>

**DOI:** 10.4103/ccod.ccd\_399\_22

**Quick Response Code:**



Amoxicillin from the penicillin group is often used to treat dentoalveolar abscesses and periodontitis. Amoxicillin is a broad-spectrum antibiotic that works by binding to penicillin-binding proteins in Gram-positive and Gram-negative bacteria and inhibiting the transpeptidation process in bacteria.<sup>[3,10]</sup> Overuse of amoxicillin can lead to side effects, such as hypersensitivity reactions, nausea, vomiting, diarrhea, thrombocytopenia, dermatological disorders, and resistance.<sup>[11]</sup> Resistance to amoxicillin can also occur due to the destruction of the  $\beta$ -lactam ring by  $\beta$ -lactamase enzymes produced by *S. aureus* and *P. gingivalis*.<sup>[2,12]</sup> Therefore, amoxicillin is often combined with clavulanic acid, known as co-amoxiclav, to reduce resistance.<sup>[13]</sup> Clavulanic acid is a  $\beta$ -lactamase inhibitor that works by inactivating the pathogen's  $\beta$ -lactamase, thereby increasing the antibacterial activity of amoxicillin.<sup>[11]</sup> However, the use of co-amoxiclav can cause side effects, such as itching, redness around the mouth, diarrhea, nausea, and idiosyncratic drug-induced liver injury.<sup>[11,13]</sup> Alternative medicines using herbal plants have the potential to minimize these side effects.<sup>[14]</sup>

Arumanis mango (*Mangifera indica* L) is a herbal plant from India cultivated in various tropical and subtropical regions, including Indonesia.<sup>[15,16]</sup> Indonesians cultivate Arumanis mangos for their sweetness, freshness, and fragrance. The fruit contains Vitamins A, B, and C, which are beneficial for health.<sup>[17]</sup> In addition, the seeds, skin, roots, and leaves of *M. indica* L. have various properties in traditional medicine. *M. indica* L. leaves are generally discarded and considered waste, even though these leaves contain secondary metabolites, including phenolics (mangiferin, tannins, and flavonoids), alkaloids, saponins, terpenoids, glycosides, and steroids that have potential antibacterial, antifungal, antiviral, antiparasitic, antioxidant, anti-inflammatory, antitumor, anticancer, and analgesic effects.<sup>[18]</sup>

Previous studies showed that *M. indica* L. leaf extracts reduced the number of *Streptococcus mutans* and improved the antibacterial effect of clindamycin against *S. aureus*.<sup>[19,20]</sup> However, no research has investigated the potential of combining amoxicillin with *M. indica* L. leaf extracts in combating *S. mutans* and *P. gingivalis*. Thus, to address this research gap, this study aimed to determine the effectiveness of amoxicillin and ethanolic extract of Arumanis mango (*M. indica* L.) in inhibiting the growth and formation of *S. aureus* and *P. gingivalis* biofilms.

## Materials and Methods

This experimental *in vitro* study was performed at the Microbiology Center of Research and Education (MiCORE) Laboratory, Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia. An ethanolic extract of Arumanis mango (*M. indica* L.) leaves was prepared at the Research Institute for Spices and Medicinal Plants (BALITTRO)

in Bogor, West Java, Indonesia. The test solutions used were 10% dimethyl sulfoxide (DMSO) (negative control), co-amoxiclav (positive control), ethanolic extract of *M. indica* L. leaves with a concentration of 100%, and a combination of amoxicillin with extract concentrations of 3.125%, 6.25%, 12.5%, 25%, 50%, and 100%.

The sample size was calculated using Federer's formula, with *n* equals number of repetition, and *t* equals as total test group (8 groups comprised 6 treatment groups, 1 positive control, and 1 negative control). Based on this formula, each treatment group was conducted with four-time repetition for all assays.

### Ethanolic extract of *Mangifera indica* L. leaves

The ethanolic extract of *M. indica* L. leaves was prepared using the maceration method. The mango leaves were washed, dried, and mashed, and the simplicial was then soaked in 70% ethanol at a ratio of 1:5. The maceration process was performed for 2–3 h, and the macerate was then allowed to stand for 24 h and filtered. The extract was diluted with 10% DMSO solution to obtain extract concentrations of 50%, 25%, 12.5%, 6.25%, and 3.125%.

### Preparation of antibiotic solution

Amoxicillin and co-amoxiclav solution were prepared by crushing 500 mg of amoxicillin and 625 mg of co-amoxiclav until smooth, using a mortar and pestle. In total, 1.2 mg of amoxicillin and 1.5 mg of co-amoxiclav were each mixed with 6 ml of sterile distilled water until homogeneous to obtain 200 g/ $\mu$ l of amoxicillin and 250 g/ $\mu$ l of co-amoxiclav.

### Phytochemical tests

Qualitative phytochemical tests were performed at BALITTRO to identify secondary metabolites, such as alkaloids, saponins, tannins, phenolics, flavonoids, triterpenoids, steroids, and glycosides, in the ethanolic extract of *M. indica* L. leaves.

### Bacterial culture

*S. aureus* ATCC 25923 and *P. gingivalis* ATCC 33277 were obtained from the MiCORE Laboratory, Faculty of Dentistry, Trisakti University. *S. aureus* was cultured on brain heart infusion broth medium (Sigma Aldrich, St. Louis, Missouri). *P. gingivalis* was cultured on Tryptone Soya Broth (Sigma Aldrich, St. Louis, Missouri) medium enriched with hemin (5 mg/l), Vitamin K (10 mg/l), 0.5% yeast extract, and L-cystine (400 mg/l).<sup>[21]</sup> The medium was incubated in an anaerobic jar (Oxoid, Basingstoke, and Hampshire) at 37° for 24 h under anaerobic conditions. Following the incubation, the bacterial absorbance value was standardized into McFarland standard of 0.5, approximately  $1.5 \times 10^8$  colony forming unit (CFU)/mL (optical density OD<sub>600</sub>  $\pm$  0.132), before the following assays.

### Microdilution and total plate count

To evaluate the antibacterial properties of the extract combined with amoxicillin, an antibacterial test was performed on the plate count by the microdilution method. In total, 100 µL of cultured *S. aureus* ATCC 25923 and *P. gingivalis* ATCC 33277 were distributed into 96-well plates (Nest Biotech, Jiangsu, China). A test solution of 100 µl was added to each well and incubated at 37°C for 24 h under anaerobic conditions. After incubation, each mixture containing treated bacteria was diluted 10<sup>5</sup> times. Five microliters of diluted mixture were then spread on a petri dish containing sterile brain heart infusion agar media. The growth of bacterial colonies was calculated after incubation at 37°C for 24 h.

### Microtiter plate biofilm assay

The bacterial cultures were inserted into each well of the 96-well plates and then incubated at 37°C for 24 h under anaerobic conditions. The supernatant was discarded, leaving a thin layer on the surface of the well. The wells were rinsed using phosphate-buffered saline (PBS) (Biomatics, Ontario, Canada). Each test solution (200 µl) was added to each well and then incubated for 1, 3, and 24 h at 37°C. The wells were rinsed twice with PBS and fixated over burning spirit lamp. Crystal violet (Merck, Darmstadt, Germany) (200 µl) was then added to each well and allowed to stand for 15 min, followed by rinsing twice and standing for 15 min. In the past step, 200 µl of 96% ethanol was added. The OD was measured using a microplate reader (Safas, Monaco) with a wavelength of 490 nm.

### Statistical analysis

The research data were processed using the Statistical Package for the Social Sciences (SPSS) computer program, version 26 (IBM, Armonk, NY, USA). The Shapiro-Wilk method was used to test the normality of the data. If the data were normally distributed ( $P > 0.05$ ), a one-way analysis of variance test was conducted, followed by Turkey's honestly significant difference test (significance level of  $P < 0.05$ ) to verify the significance between the groups.

## Results

### Phytochemical screening

The results of the phytochemical screening qualitatively proved that the ethanolic extract of *M. indica* L. leaves contains alkaloids, saponins, tannins, phenolics, flavonoids, and steroids [Table 1].

### Antibacterial test using microdilution and plate count methods

The antibacterial test results showed that the ethanolic extract of *M. indica* L. leaves at 100% concentration and the combination of amoxicillin with extracts of various

**Table 1: The results of phytochemical screening of the ethanol extract of *Mangifera indica* L. leaves**

Secondary metabolites	Screening result
Alkaloids	+
Saponins	+
Tannins	+
Phenolics	+
Flavonoids	+
Triterpenoids	-
Steroids	+
Glycoside	-

concentrations exhibited antibacterial and antibiofilm effects against *S. aureus* and *P. gingivalis* [Figure 1]. In terms of inhibition of *S. aureus*, the results obtained by the combination of amoxicillin and the extract were not significantly different from those obtained using the positive control ( $P > 0.05$ ). The extract at a concentration of 12.5% and higher was effective against *S. aureus*, and the extract at a concentration of 3.125% and higher was effective against *P. gingivalis* [Figures 2 and 3].

### Antibiofilm test using the microtiter plate biofilm assay method

The results of the antibiofilm test showed that the addition of extracts to amoxicillin in inhibiting *S. aureus* biofilms had a significantly lower OD value compared to the positive control ( $P < 0.05$ ), starting at a concentration of 25% after 1 h of incubation [Figure 4] and at a concentration of 6.25% after 3 h of incubation [Figure 5]. After 24 h of incubation, following OD measurement, the group with the combination of extract concentration of 100% and amoxicillin proved not to be significantly different from OD of the positive control ( $P > 0.05$ ) [Figure 6]. In the *P. gingivalis* biofilm, the OD values of the addition of amoxicillin and the extract group, starting at a concentration of 25% after 1, 3, and 24 h of incubation, were smaller than the OD values of the positive control amoxiclav. The OD values of the treatment group were significantly different from those of the positive control amoxiclav [Figures 7-9].

## Discussion

As shown by our results, the ethanol extract of *M. indica* L. leaves contains secondary metabolites, including alkaloids, saponins, tannins, phenolics, flavonoids, and steroids. Different mechanisms of action of each compound account for the antibacterial and antibiofilm properties of the extract. Alkaloids inhibit the formation of peptidoglycan in bacterial cells. The alkaloids can disrupt the amino acid structure of bacterial DNA, leading to bacterial lysis.<sup>[22]</sup> Saponins damage the cell membrane and cell wall permeability in the diffusion process, resulting in the release of enzymes, amino acids, nutrients, and water, leading to cell destabilization and cell death.<sup>[23]</sup> Tannins form a complex with protein in the cell wall, namely,




Figure 1: The results of the inhibition test of *Staphylococcus aureus* and *Porphyromonas gingivalis* using the plate count method




Figure 2: Graph of the average total colony of *Staphylococcus aureus* (\* $P < 0.05$ , \*\* $P < 0.01$ )




Figure 3: Graph of the average total colony of *Porphyromonas gingivalis* (\* $P < 0.05$ , \*\* $P < 0.01$ )

proline, which can damage the cell wall.<sup>[22]</sup> The most abundant phenolic compound in *M. indica* L leaves is mangiferin. Mangiferin belongs to the xanthone C-glucosyl group, which can damage cell structure and cell membranes and inhibit bacterial protein synthesis.<sup>[24]</sup> Previous research reported that mangiferin compounds interfere with the mechanism of drug resistance; thus, restoring

bactericidal and bacteriostatic effects of nalidixic acid, ampicillin, tetracycline, and sulfamethoxazole/trimethoprim.<sup>[25]</sup> However, the exact mechanism of how mangiferin interferes with the mechanism of drug resistance is yet to be elucidated. Flavonoids, as antibacterial, play a role in disrupting the activity of cell wall formation by suppressing cytoplasm function, interrupting the nutrient exchange process, and thereby inhibiting the energy supply of bacteria.<sup>[26]</sup> In addition, flavonoids inhibit enzymes from producing quorum-sensing signals, thereby disrupting the communication process between cells during biofilm formation.<sup>[27]</sup> Steroids can cause leakage of lysosomes and membrane phospholipids that reduce the integrity of cell membranes and lead to cell lysis.<sup>[28]</sup>

Thus far, only a few studies have focused on antibacterial and antibiofilm properties of ethanolic extract of *M. indica* L. leaves.<sup>[19,20]</sup> The concentration of the extract used in this study ranged from 3.125% to 100%. As a positive control, co-amoxiclav, a combination of amoxicillin and clavulanic acid, was used. Amoxicillin, a  $\beta$ -lactam antibiotic, works by inhibiting the synthesis of bacterial cell walls.<sup>[11]</sup> The release of  $\beta$ -lactamase enzymes by *S. aureus* and *P. gingivalis* decrease the antibacterial effect of amoxicillin.<sup>[2,12]</sup> The addition of clavulanic acid binds to  $\beta$ -lactamase enzymes from bacteria, inhibiting the enzyme thereby unable to cleave  $\beta$ -lactam ring in amoxicillin, so the amoxicillin can still exhibit its antibacterial activity.<sup>[11]</sup>

In the microdilution and plate count tests, the addition of the ethanolic extract of *M. indica* L. leaves to amoxicillin inhibited the growth of *S. aureus* and *P. gingivalis*, and the combination of the ethanolic extract and amoxicillin was as effective as that of co-amoxiclav. In terms of antibacterial activity, a combination of ethanolic extract at a concentration of 12.5% and amoxicillin was as effective as co-amoxiclav against *S. aureus*, and a concentration as low as 3.125% was as effective as co-amoxiclav against *P. gingivalis*. Therefore, ethanolic extract of *M. indica* L. leaves may be a potential  $\beta$ -lactamase inhibitor equivalent to clavulanic acid. The results of this study are in line with the research of Hartanto *et al.*, who showed that adding




Figure 4: Graph of the average OD value of *Staphylococcus aureus* biofilm after 1 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). OD: Optical density




Figure 5: Graph of the average OD value of *Staphylococcus aureus* biofilm after 3 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). OD: Optical density




Figure 6: Graph of the average OD value of *Staphylococcus aureus* biofilm after 24 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). OD: Optical density




Figure 7: Graph of the average OD value of *Porphyromonas gingivalis* biofilm after 1 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). OD: Optical density




Figure 8: Graph of the average OD value of *Porphyromonas gingivalis* biofilm after 3 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). OD: Optical density




Figure 9: Graph of the average OD value of *Porphyromonas gingivalis* biofilm after 24 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). OD: Optical density

methanolic extract of *M. indica* L. leaves to clindamycin has antibacterial effects against *S. aureus*, especially at a concentration of 100%.<sup>[19]</sup> The bioactive component of *M. indica* L. leaves, namely, mangiferin, is known to have a synergistic effect on tetracycline, ampicillin, nalidixic acid, and trimethoprim in inhibiting the growth of *S. aureus*.<sup>[25]</sup>

The biofilm formation phase begins with pellicle formation, which occurs in the first few seconds to the first min from

the initial contact, with the initial adhesion phase occurring 2–4 h later. After 24 h, the biofilm enters the maturation phase, becoming 1.000 - 1.500 times more resistant than planktonic bacteria.<sup>[29]</sup> The incubation time used in the antibiofilm test in this study was adjusted to the stage of biofilm formation, namely 1, 3, and 24 h. This timing aimed to determine at which stage of biofilm formation amoxicillin and the ethanolic extract of *M. indica* L. leaves

most effectively inhibited *S. aureus* and *P. gingivalis*. In the antibiofilm test, the OD from the combination of amoxicillin with ethanolic extract of *M. indica* L. leaves starting at a concentration of 25% after 1 h of incubation and 6.25% after 3 h of incubation had a significantly lower OD value compared to co-amoxiclav. This result proved that the ethanolic extract of *M. indica* L. leaves at a concentration of 25% after 1 h of incubation and 6.25% after 3 h of incubation was more effective at inhibiting *S. aureus* biofilms than co-amoxiclav. The OD values for the combination of amoxicillin and the extract at a concentration of 100% after 24 h of incubation were lower than those obtained for co-amoxiclav, but the difference was not statistically significant. Therefore, the combination of amoxicillin and an extract concentration of 100% has antibiofilm properties equivalent to those of co-amoxiclav after 24 h of incubation.

In terms of inhibiting *P. gingivalis* biofilm, the extract at a concentration of 25% and higher during the entire incubation period showed a more potent antibiofilm effect than that of co-amoxiclav. The combination of amoxicillin and the ethanolic extract of *M. indica* L. leaves inhibited the formation of *S. aureus* biofilms at a concentration of 6.25% after 3 h of incubation and *P. gingivalis* biofilms at a concentration of 25% after 1 h of incubation. This study supports the findings of previous research, which reported that ethanolic extract of *M. indica* L. leaves reduced *S. aureus* attachment and the number of *S. aureus* biofilms.<sup>[30]</sup>

This study has several limitations. First, the ethanolic extract used in this study was a crude extract. The use of a more refined extract, such as an extract exposed to extraction chromatography, would have resulted in an extract with fewer impurities. Second, only two of the many known oral pathogens were used in this study. Other oral pathogens can also be tested to expand the antibacterial activity of amoxicillin combined with *M. indica* L. leaves ethanolic extract. Further research is needed to determine the toxicity of this combination. Preclinical and clinical tests should also be conducted before the combination can be used as an alternative treatment for dentoalveolar abscesses and periodontitis.

## Conclusions

Ethanolic extract of *M. indica* L. leaves contains alkaloids, saponins, tannins, phenolics, flavonoids, and steroids that have the potential to inhibit the growth and formation of *S. aureus* and *P. gingivalis* biofilms *in vitro*. Within the limitations of this preliminary study, we conclude that the addition of ethanolic extract of *M. indica* L. leaves to amoxicillin could potentially increase the antibacterial and antibiofilm properties of amoxicillin against *S. aureus* and *P. gingivalis*.

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## References

1. Ministry of Health of the Republic of Indonesia. Report on Result of National Basic Health Research 2018. Jakarta: Ministry of Health; 2018. p. 197-207.
2. Japoni A, Vasin A, Noushadi S, Kiany F, Japoni S, Alborzi A. Antibacterial susceptibility patterns of *Porphyromonas gingivalis* isolated from chronic periodontitis patients. *Med Oral Patol Oral Cir Bucal* 2011;16:e1031-5.
3. Shweta N, Prakash SK. Dental abscess: A microbiological review. *Dent Res J (Isfahan)* 2013;10:585-91.
4. Mehrotra N, Singh S. Periodontitis. Treasure Island (FL): StatPearls Publishing; 2011.
5. Archer NK, Mazaitis MJ, Costerton JW, Leid JG, Powers ME, Shirtliff ME. *Staphylococcus aureus* biofilms: Properties, regulation, and roles in human disease. *Virulence* 2011;2:445-59.
6. Huang R, Li M, Gregory RL. Bacterial interactions in dental biofilm. *Virulence* 2011;2:435-44.
7. Periasamy S, Joo HS, Duong AC, Bach TH, Tan VY, Chatterjee SS, et al. How *Staphylococcus aureus* biofilms develop their characteristic structure. *Proc Natl Acad Sci U S A* 2012;109:1281-6.
8. Taylor TA, Unakal CG. *Staphylococcus aureus*. Treasure Island (FL): StatPearls Publishing; 2021.
9. How KY, Song KP, Chan KG. *Porphyromonas gingivalis*: An overview of periodontopathic pathogen below the gum line. *Front Microbiol* 2016;7:53.
10. Ciancio SG, Mariotti AJ. Systemic Anti-infective therapy for periodontal diseases. In: Carranza F, Newman M, Takei H, Klokkevold P, editors. *Newman and Carranza's Clinical Periodontology*. 13<sup>th</sup> ed. Philadelphia: Elsevier; 2019. p. 555-7.
11. Akhavan B, Khanna N, Vjihani P. Amoxicillin. Treasure Island (FL): StatPearls Publishing; 2020.
12. Foster TJ, Geoghegan JA. *Staphylococcus aureus*. In: *Molecular Medical Microbiology*. Massachusetts: Elsevier; 2015. p. 655-74.
13. Samaranyake L. *Essential Microbiology for Dentistry*. 5<sup>th</sup> ed. Poland: Elsevier Ltd; 2018. p. 75, 158, 293, 297.
14. Joshua M, Takudzwa M. Antibacterial properties of *Mangifera indica* on *Staphylococcus aureus*. *Afr J Clin Exp Microbiol* 2013;14:62-74.
15. Kalita P. An overview on *Mangifera indica*: Importance and its various pharmacological action. *Pharma Tutoring* 2014;2:72-6.
16. Sivakumar D, Jiang Y, Yahia EM. Maintaining mango (*Mangifera indica* L.) fruit quality during the export chain. *Food Res Int* 2011;44:1254-63.
17. Chiumarelli M, Ferrari CC, Sarantopoulos CI, Hubinger MD. Fresh cut *Tommy Atkins* mango pre-treated with citric acid and coated with cassava (*Manihot esculenta* Crantz) starch or sodium alginate. *Innov Food Sci Emerg Technol* 2011;12:381-7.
18. Gu C, Yang M, Zhou Z, Khan A, Cao J, Cheng G. Purification and characterization of four benzophenone derivatives from *Mangifera indica* L. leaves and their antioxidant, immunosuppressive and  $\alpha$ -glucosidase inhibitory activities. *J Funct Foods* 2019;52:709-14.
19. Hartanto R, Tran V, Khang G, Pham T, Trinh T, Denhara C. Effect of the addition of arumanis mango leaf extract (*Mangifera indica* L.) on clindamycin antibiotics in inhibiting the growth of *Staphylococcus aureus*. *J Prima Med Sains* 2020;2:14-7.

20. Kurniasih R. Effect of Concentrations of Young Mango Arumanis Ethanolic Extract (*Mangifera indica* L.) leaves on growth inhibition of *Streptococcus mutans* *in vitro*. Surakarta: Fakultas Kedokteran Gigi Universitas Muhammadiyah (Faculty of Dentistry, Muhammadiyah University); 2016.
21. Yamanaka T, Furukawa T, Matsumoto-Mashimo C, Yamane K, Sugimori C, Nambu T, *et al.* Gene expression profile and pathogenicity of biofilm-forming *Prevotella intermedia* strain 17. *BMC Microbiol* 2009;9:11.
22. Sylvana D, Amir M, Purnamasari CB, Iskandar A, Asfirizal V. Antibacterial activity of ethanol extract of beluntas leaves on *Streptococcus mutans*, *Porphyromonas gingivalis*, and *Enterococcus faecalis*. *Padjadjaran J Dent* 2021;33:191-8.
23. Sebastian J, Widyarman AS. Roselle flower petals extract inhibits periodontal pathogenic biofilms. *J Dentomaxillofacial Sci* 2021;6:102-5.
24. Tirado-Kulieva V, Atoche-Dioses S, Hernandez-Martinez E. Phenolic compounds of mango (*Mangifera indica*) by-products: Antioxidant and antimicrobial potential, use in disease prevention and food industry, methods of extraction, and microencapsulation. *Sci Agropecu* 2021;12:283-93.
25. Mazlan NA, Azman S, Ghazali NF, Zarith P, Yusri S. Synergistic antibacterial activity of mangiferin with antibiotics against *Staphylococcus aureus*. *Drug Invent Today* 2019;12:14-7.
26. Ma Y, Ding S, Fei Y, Liu G, Jang H, Fang J. Antimicrobial activity of anthocyanins and catechins against foodborne pathogens *Escherichia coli* and *Salmonella*. *Food Control* 2019;106:78-80.
27. Federika AS, Rukmo M, Setyabudi S. Antibiofilm activity of flavonoid mangosteen pericarp extract against *Porphyromonas gingivalis* bacteria. *Conserv Dent J* 2020;10:27-30.
28. Madduluri S, Rao KB, Sitaram B. *In vitro* evaluation of antibacterial activity of five indigenous plants extracts against five bacteria pathogens of humans. *Int J Pharm Pharm Sci* 2013;5:679-84.
29. Bjarnsholt T. The role of bacterial biofilms in chronic infections. *APMIS Suppl* 2013;121:1-51.
30. Manzur AG, Sm Junior V, Morais-Costa F, Mariano EG, Careli RT, da Silva LM, *et al.* Extract of *Mangifera indica* L. leaves may reduce biofilms of *Staphylococcus* spp. in stainless steel and teacup rubbers. *Food Sci Technol Int* 2020;26:11-20.



# The Antibacterial and Antibiofilm Effect of Amoxicillin and Mangifera indica L. Leaves Extract on Oral Pathogens

## ORIGINALITY REPORT

20%

SIMILARITY INDEX

18%

INTERNET SOURCES

14%

PUBLICATIONS

9%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://bmc.altmetric.com">bmc.altmetric.com</a> Internet Source	3%
2	<a href="http://innovareacademics.in">innovareacademics.in</a> Internet Source	2%
3	Submitted to Fakultas Kedokteran Gigi Universitas Trisakti Student Paper	2%
4	<a href="http://ijnmr.mui.ac.ir">ijnmr.mui.ac.ir</a> Internet Source	2%
5	<a href="http://www2.mdpi.com">www2.mdpi.com</a> Internet Source	1%
6	<a href="http://rcastoragev2.blob.core.windows.net">rcastoragev2.blob.core.windows.net</a> Internet Source	1%
7	<a href="http://www.thieme-connect.com">www.thieme-connect.com</a> Internet Source	1%
8	Submitted to Universitas Airlangga Student Paper	1%

9	Internet Source	1 %
10	<a href="http://www.researchpublish.com">www.researchpublish.com</a> Internet Source	1 %
11	<a href="http://journal.unpad.ac.id">journal.unpad.ac.id</a> Internet Source	1 %
12	<a href="http://link.springer.com">link.springer.com</a> Internet Source	1 %
13	Andressa GB Manzur, Valdo SM Junior, Franciellen Morais-Costa, Emanuely GA Mariano et al. " Extract of L. leaves may reduce biofilms of spp. in stainless steel and teatcup rubbers ", Food Science and Technology International, 2019 Publication	1 %
14	<a href="http://biomedpharmajournal.org">biomedpharmajournal.org</a> Internet Source	1 %
15	Armelia Sari Widyarman, Louise Anastasya Halim, Jesslyn, Heidi Amanda Irma, Mario Richi, Muhammad Ihsan Rizal. "The Potential of Reuterin Derived from Indonesian Strain of Lactobacilus reuteri against Endodontic Pathogen Biofilms in vitro and ex vivo", The Saudi Dental Journal, 2023 Publication	1 %
16	Submitted to Universitas Jember Student Paper	

1 %

17

Armelia Sari Widyarman, Thalia Venessa Maukar, Rosalina Tjandrawinata, Citra Fragrantia Theodora. "Antibiofilm Activity of Parabiotic Reuterin on Acrylic Resin Plates", 2021 IEEE International Conference on Health, Instrumentation & Measurement, and Natural Sciences (InHeNce), 2021

Publication

<1 %

18

[journals.iium.edu.my](http://journals.iium.edu.my)

Internet Source

<1 %

19

[jurnal.pdgi.or.id](http://jurnal.pdgi.or.id)

Internet Source

<1 %

20

[www.scielo.br](http://www.scielo.br)

Internet Source

<1 %

Exclude quotes  On

Exclude matches  < 15 words

Exclude bibliography  On

# The Antibacterial and Antibiofilm Effect of Amoxicillin and Mangifera indica L. Leaves Extract on Oral Pathogens

---

GRADEMARK REPORT

---

FINAL GRADE

GENERAL COMMENTS

**/0**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

Copyright agreement & co-author account activation

Contemporary Clinical Dentistry <editor@contempclinidnt.org>  
to: ms

Tue, Jul 26, 2022, 11:42AM

Dear Mrs Sheila Soesanto,

You have been added as a co-author for article entitled THE ANTIBACTERIAL AND ANTI-BIOFILM EFFECT OF AMOXICILLIN AND Mangifera indica L. LEAVES EXTRACT ON ORAL PATHOGENS submitted to Contemporary Clinical Dentistry.

Please review the Copyright Form and agree on Terms & Conditions for the article by clicking on the below link:

<https://review.iqwi.medknow.com/outside/user-accept-copyright?journal=2836&author=18977271>

To activate your account as co-author, click on the link given below or copy and paste the entire link to browser address bar:

<https://review.iqwi.medknow.com/outside/user-verification?token=9878812f28ab-4fed-9c11-f5bc1a7a7938&email=sheilasoesanto@trusmi.ac.id&journalid=2836&journalCode=ccd>

Your Username is [sheilasoesanto@trusmi.ac.id](mailto:sheilasoesanto@trusmi.ac.id)

This will confirm your registration into our system.

Please note that the manuscript is visible only under the account of submitting author. Please co-ordinate with the submitting author for the article status. You may use your author account to submit new manuscript after the activation.

You have been provided with the following Journal Uri for login:

1 <https://review.iqwi.medknow.com/ccd>

Thank you.

Best Regards  
Editor in chief

Fwd: ccd Manuscript for revision: ccd\_399\_22

mail.google.com/mail/u/3/#label/Jurnal%2FCCD/WhctKkZPGVDLzrCtTRRbPrZQBDxRjhntDJsPrKzrNSLwcvGCHCnK...

USAKTI - SIS SIMPPM SS

Compose

Inbox 6

Starred

Snoozed

Sent

Drafts 2

More

Labels +

- Bintang SISTER SUALI
- HKI
- Jurnal
  - AKAL
  - CCD**
  - ETC
  - Fori
  - Inhence
  - JIDA

label:jurnal-ccd

Fwd: ccd Manuscript for revision: ccd\_399\_22 (External) Jurnal/CCD

Armelia Sari <armeliasari@trisakti.ac.id>  
to me, Armelia

Fri, Dec 22, 2022, 6:19 PM

Sent from my iPhone

Begin forwarded message:

From: Contemporary Clinical Dentistry <ccdit@contemccindent.org>  
Date: 25 December 2022, 16:42:16 GMT+3  
To: armeliasari@trisakti.ac.id  
Subject: ccd Manuscript for revision: ccd\_399\_22

Dear Dr. Widjarmas,

With reference to your manuscript ccd\_399\_22 entitled THE ANTIBACTERIAL AND ANTIBIOFILM EFFECT OF AMOXICILLIN AND Mangifera indica L. LEAVES EXTRACT ON ORAL PATHOGENS, please review the comments of the referees from our site <https://review.jow.medknow.com/ccd>. The manuscript would be reconsidered after requisite modifications as per the comments and instructions provided by the journal.

If you wish to continue with the publication process, kindly make the changes according to the comments and upload the revised manuscript along with clarifications for all the comments clearly indicating the areas where the changes have been made.

Fwd: ccd Manuscript for revision: ccd\_399\_22

mail.google.com/mail/u/3/#label/Jurnal%2FCCD/WhctKkZPGVDLzrCtTRRbPrZQBDxRjhntDJsPrKzrNSLwcvGCHCnK...

USAKTI - SIS SIMPPM SS

Compose

Inbox 6

Starred

Snoozed

Sent

Drafts 2

More

Labels +

- Bintang SISTER SUALI
- HKI
- Jurnal
  - AKAL
  - CCD**
  - ETC
  - Fori
  - Inhence
  - JIDA

label:jurnal-ccd


Do check the FAQ regarding replying to the comments and uploading a file. The template of point-by-point comments file for the reviewers, is available in your dashboard under the 'Downloads' menu option.

The journal allows two weeks for the revision of the manuscript. If we do not hear from you within this period, we will consider it as your decision to withdraw your article from publication. Please also note that the submission of the revised article does not guarantee its final acceptance by the journal.

We thank you for submitting your valuable research work to Contemporary Clinical Dentistry.

With warm personal regards,

Editorial Team  
Contemporary Clinical Dentistry



**UNIVERSITAS TRISAKTI**  
"Is a one stop learning for sustainable development"  
Kampus A, Jl. Kyau Tapa No. 1, Grogol  
Jakarta Barat 11440 - INDONESIA  
[www.trisakti.ac.id](http://www.trisakti.ac.id)  
(t) +62-21-266 3232, (f) +62-21-567 3001

Fwd: Author Resubmission Acknowledgement letter: ccd\_399\_22

Armelia Sari <armeliasari@tricolit.ac.id>  
to me, Armelia

Sent from my iPhone

Begin forwarded message:

From: Contemporary Clinical Dentistry <editor@contemporaryclinical.org>  
Date: 21 November 2022 10:54:19 GMT+3  
To: [armeliasari@tricolit.ac.id](mailto:armeliasari@tricolit.ac.id)  
Subject: Author Resubmission Acknowledgement letter: ccd\_399\_22

Dear Dr. Widjerman,

Contemporary Clinical Dentistry has received your revised manuscript "ccd\_399\_22" entitled "THE ANTIBACTERIAL AND ANTIBIOFILM EFFECT OF AMOXICILLIN AND *Mangifera indica* L. LEAVES EXTRACT ON ORAL PATHOGENS? (in consideration for publication, which was sent for technical modification.

We thank you for submitting your valuable research work to the Contemporary Clinical Dentistry.

Yours sincerely,  
Editors' Team  
Contemporary Clinical Dentistry

Fwd: Author-side fee of your manuscript: ccd\_399\_22

Armelia Sari <armeliasari@tricolit.ac.id>  
to me, Armelia

Sent from my iPhone

Begin forwarded message:

From: Contemporary Clinical Dentistry <editor@contemporaryclinical.org>  
Date: 20 February 2023 10:22:07 GMT+3  
To: [armeliasari@tricolit.ac.id](mailto:armeliasari@tricolit.ac.id)  
Subject: Author-side fee of your manuscript: ccd\_399\_22

Dear Dr. Widjerman,

We are pleased to inform that your manuscript "Original Article: Antibacterial and Antibiofilm Effects of Amoxicillin and *Mangifera indica* L. Leaf Extract on Oral Pathogens" is now acceptable after clearing the dues for publication of the manuscript. The details of the same can be found on the journal website under "Instructions to the Authors" page.

The payment can be done using the following link:  
<https://prepayment.medknow.com/online/index?apicaller=UNIPRR>

Browser window showing a Gmail email. The address bar contains: `mail.google.com/mail/u/3/#label/Jurnal%2FCOD/WhctKKZPGVDLZzOzDFLxMTsPvbnPb5VjdCm5K8qSFxtwmdCJISMjktg...`

**Gmail** Search: `label:jurnal-ccd`

Compose

Inbox 6  
Starred  
Snoozed  
Sent  
Drafts 2  
More

Labels +  
Bintang SISTER SIJALI  
HKI  
Jurnal  
AKAL  
CCD

The following options are available for payment:


- Pay online
- Cheque payment
- Wire transfer

Once the payment is received at our end, the manuscript would be processed further and you would receive an edited version of article in about 2-3 weeks from now for a final check and correction.

We thank you for submitting your valuable research work to Contemporary Clinical Dentistry.

- With warm personal regards,

Yours sincerely,  
Editorial Team  
Contemporary Clinical Dentistry





# AQ2 Antibacterial and Antibiofilm Effects of Amoxicillin and *Mangifera indica* L. Leaf Extract on Oral Pathogens

## Abstract

**Objective:** This study aimed to determine the antibacterial and antibiofilm effects of amoxicillin combined with extract of *Mangifera indica* L. leaves against *Staphylococcus aureus*, and *Porphyromonas gingivalis*. **Materials and Methods:** This was an experimental laboratory *in vitro* study with a posttest-only control group design. An antibacterial test using the plate count method and an antibiofilm test using the microtiter plate biofilm assay method were conducted. The research samples comprised extract of *M. indica* L. leaves with a concentration of 100%; amoxicillin and extract concentrations of 3.125%, 6.25%, 12.5%, 25%, 50%, and 100%; and amoxicillin. Dimethyl sulfoxide served as a negative control and co-amoxiclav served as a positive control. **Results:** The combination of amoxicillin and the extract exhibited an antibacterial effect against *S. aureus* at a concentration of 12.5% and higher and more effective against *P. gingivalis* at a concentration of 3.125% and higher. In the antibiofilm test, the combination of amoxicillin and the extract at a concentration of 25% after 1 h of incubation and a concentration of 6.25% after 3 h of incubation inhibited *S. aureus*. The inhibition of *S. aureus* biofilms at a concentration of 100% after 24 h of incubation was as effective as that of co-amoxiclav. The extract at a concentration of 25% over the entire incubation period showed inhibition against the *P. gingivalis* biofilm. **Conclusions:** The ethanolic extract of *M. indica* L. leaves and the combination of amoxicillin and the extract have the potential to inhibit the growth and formation of *S. aureus* and *P. gingivalis* biofilms.

**Keywords:** Amoxicillin, antibiofilm, arumanis mango, ethanol extract of *Mangifera indica* L. leaves

## Introduction

Based on data from Basic Health Research (Riskesdas) in 2018, Indonesia's dental and oral health problems reached 57.6% of overall health-care problems.<sup>[1]</sup> The latter is influenced by poor oral hygiene, which triggers various diseases, such as dentoalveolar abscesses and periodontitis.<sup>[2,3]</sup> A dentoalveolar abscess is a pathological cavity in the oral cavity that contains pus due to secondary infection caused by caries, trauma, failure of root canal treatment, and poor oral hygiene.<sup>[3]</sup> Periodontitis is a chronic inflammation of the periodontal tissue structure, including gingiva, bone, and the periodontal ligament, which can cause pocket formation, recession, tooth mobility, or tooth loss.<sup>[2]</sup> Dentoalveolar abscesses and periodontitis are associated with bacterial pathogens in biofilms.<sup>[3,4]</sup>

A biofilm is a collection of microbial cells, especially bacteria, attached to the tooth

surface and coated by an extracellular polymeric substance. This coating protects cells and enables accelerated growth rates, along with additional horizontal gene transfer between cells within the coating, which promotes additional problems, such as antibiotic resistance.<sup>[5]</sup> The formation of biofilms in the oral cavity involves complex competition between microflora for initial attachment.<sup>[6]</sup> *Staphylococcus aureus*, a Gram-positive bacterium, initiates adherence in biofilm formation and produces multiple layers of biofilm embedded in a glycocalyx layer.<sup>[7,8]</sup> *S. aureus* has several virulence factors (capsules, adhesins, coagulase, hyaluronidase, staphylokinase, enterotoxin, and leucocidin) that support biofilm formation in a dentoalveolar abscess.<sup>[6]</sup> *Porphyromonas gingivalis*, which belongs to anaerobic Gram-negative bacteria, is the second most common colonizing bacteria in biofilms. *P. gingivalis* virulence factors, such as lipopolysaccharides, capsules, fimbriae, outer membrane

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow\_reprints@wolterskluwer.com

**How to cite this article:** Soesanto S, Hepziba ER, Yasnill, Widyarman AS. Antibacterial and antibiofilm effects of amoxicillin and *Mangifera indica* L. Leaf extract on oral pathogens. Contemp Clin Dent 2023;XX:XX-XX.

Sheila Soesanto<sup>1</sup>,  
Evangelista Rachel  
Hepziba<sup>2</sup>,  
Yasnill<sup>2</sup>,  
Armelia Sari  
Widyarman<sup>3</sup>

<sup>1</sup>Department of Pharmacology,  
Faculty of Dentistry, Universitas  
Trisakti, Jakarta, Indonesia,  
<sup>2</sup>???, Faculty of Dentistry,  
Universitas Trisakti, Jakarta,  
Indonesia, <sup>3</sup>Department of  
Microbiology, Faculty of  
Dentistry, Universitas Trisakti,  
Jakarta, Indonesia

Submitted : 26-Jul-2022  
Revised : 06-Jan-2023  
Accepted : 21-Feb-2023  
Published : \*\*\*

**Address for correspondence:**  
Dr. Armelia Sari Widyarman,  
Department of Microbiology,  
Faculty of Dentistry, Universitas  
Trisakti, Kyai Tapa 260, Grogol,  
Jakarta 11440, Indonesia.  
E-mail: armeliasari@trisakti.  
ac.id

Access this article online

Website:  
www.contempclindent.org

DOI: 10.4103/ccd.ccd\_399\_22

Quick Response Code:



proteins, proteases, and enzymes, induce the destruction of periodontal tissue, causing periodontitis.<sup>[9]</sup>

Amoxicillin from the penicillin group is often used to treat dentoalveolar abscesses and periodontitis. Amoxicillin is a broad-spectrum antibiotic that works by binding to penicillin-binding proteins in Gram-positive and Gram-negative bacteria and inhibiting the transpeptidation process in bacteria.<sup>[3,10]</sup> Overuse of amoxicillin can lead to side effects, such as hypersensitivity reactions, nausea, vomiting, diarrhea, thrombocytopenia, dermatological disorders, and resistance.<sup>[11]</sup> Resistance to amoxicillin can also occur due to the destruction of the  $\beta$ -lactam ring by  $\beta$ -lactamase enzymes produced by *S. aureus* and *P. gingivalis*.<sup>[2,12]</sup> Therefore, amoxicillin is often combined with clavulanic acid, known as co-amoxiclav, to reduce resistance.<sup>[13]</sup> Clavulanic acid is a  $\beta$ -lactamase inhibitor that works by inactivating the pathogen's  $\beta$ -lactamase, thereby increasing the antibacterial activity of amoxicillin.<sup>[11]</sup> However, the use of co-amoxiclav can cause side effects, such as itching, redness around the mouth, diarrhea, nausea, and idiosyncratic drug-induced liver injury.<sup>[11,13]</sup> Alternative medicines using herbal plants have the potential to minimize these side effects.<sup>[14]</sup>

Arumanis mango (*Mangifera indica* L) is a herbal plant from India cultivated in various tropical and subtropical regions, including Indonesia.<sup>[15,16]</sup> Indonesians cultivate Arumanis mangos for their sweetness, freshness, and fragrance. The fruit contains Vitamins A, B, and C, which are beneficial for health.<sup>[17]</sup> In addition, the seeds, skin, roots, and leaves of *M. indica* L. have various properties in traditional medicine. *M. indica* L. leaves are generally discarded and considered waste, even though these leaves contain secondary metabolites, including phenolics (mangiferin, tannins, and flavonoids), alkaloids, saponins, terpenoids, glycosides, and steroids that have potential antibacterial, antifungal, antiviral, antiparasitic, antioxidant, anti-inflammatory, antitumor, anticancer, and analgesic effects.<sup>[18]</sup>

Previous studies showed that *M. indica* L. leaf extracts reduced the number of *Streptococcus mutans* and improved the antibacterial effect of clindamycin against *S. aureus*.<sup>[19,20]</sup> However, no research has investigated the potential of combining amoxicillin with *M. indica* L. leaf extracts in combating *S. mutans* and *P. gingivalis*. Thus, to address this research gap, this study aimed to determine the effectiveness of amoxicillin and ethanolic extract of Arumanis mango (*M. indica* L.) in inhibiting the growth and formation of *S. aureus* and *P. gingivalis* biofilms.

## Materials and Methods

This experimental *in vitro* study was performed at the Microbiology Center of Research and Education (MiCORE) Laboratory, Faculty of Dentistry,

Trisakti University, Jakarta, Indonesia. An ethanolic extract of Arumanis mango (*M. indica* L.) leaves was prepared at the Research Institute for Spices and Medicinal Plants (BALITTRO) in Bogor, West Java, Indonesia. The test solutions used were 10% dimethyl sulfoxide (DMSO) (negative control), co-amoxiclav (positive control), ethanolic extract of *M. indica* L. leaves with a concentration of 100%, and a combination of amoxicillin with extract concentrations of 3.125%, 6.25%, 12.5%, 25%, 50%, and 100%.

The sample size was calculated using Federer's formula, with n equals number of repetition, and t equals as total test group (8 groups comprised 6 treatment groups, 1 positive control, and 1 negative control). Based on this formula, each treatment group was conducted with four-time repetition for all assays.

### Ethanolic extract of *Mangifera indica* L. leaves

The ethanolic extract of *M. indica* L. leaves was prepared using the maceration method. The mango leaves were washed, dried, and mashed, and the simplicial was then soaked in 70% ethanol at a ratio of 1:5. The maceration process was performed for 2–3 h, and the macerate was then allowed to stand for 24 h and filtered. The extract was diluted with 10% DMSO solution to obtain extract concentrations of 50%, 25%, 12.5%, 6.25%, and 3.125%.

### Preparation of antibiotic solution

Amoxicillin and co-amoxiclav solution were prepared by crushing 500 mg of amoxicillin and 625 mg of co-amoxiclav until smooth, using a mortar and pestle. In total, 1.2 mg of amoxicillin and 1.5 mg of co-amoxiclav were each mixed with 6 ml of sterile distilled water until homogeneous to obtain 200 g/ $\mu$ l of amoxicillin and 250 g/ $\mu$ l of co-amoxiclav.

### Phytochemical tests

Qualitative phytochemical tests were performed at BALITTRO to identify secondary metabolites, such as alkaloids, saponins, tannins, phenolics, flavonoids, triterpenoids, steroids, and glycosides, in the ethanolic extract of *M. indica* L. leaves.

### Bacterial culture

*S. aureus* ATCC 25923 and *P. gingivalis* ATCC 33277 were obtained from the MiCORE Laboratory, Faculty of Dentistry, Trisakti University. *S. aureus* was cultured on brain heart infusion broth medium (Sigma Aldrich, St. Louis, Missouri). *P. gingivalis* was cultured on Tryptone Soya Broth (Sigma Aldrich, St. Louis, Missouri) medium enriched with hemin (5 mg/l), Vitamin K (10 mg/l), 0.5% yeast extract, and L-cystine (400 mg/l).<sup>[21]</sup> The medium was incubated in an anaerobic jar (Oxoid, Basingstoke, Hampshire) at 37° for 24 h under anaerobic conditions.

Following the incubation, the bacterial absorbance value was standardized into McFarland standard of 0.5, approximately  $1.5 \times 10^8$  colony forming unit (CFU)/mL (optical density [OD]<sub>600</sub> ± 0.132), before the following assays.

### Microdilution and total plate count

To evaluate the antibacterial properties of the extract combined with amoxicillin, an antibacterial test was performed on the plate count by the microdilution method. In total, 100 µL of cultured *S. aureus* ATCC 25923 and *P. gingivalis* ATCC 33277 were distributed into 96-well plates (Nest Biotech, Jiangsu, China). A test solution of 100 µl was added to each well and incubated at 37°C for 24 h under anaerobic conditions. After incubation, each mixture containing treated bacteria was diluted 10,000 times. Five microliters of diluted mixture were then spread on a petri dish containing sterile brain heart infusion agar media. The growth of bacterial colonies was calculated after incubation at 37°C for 24 h.

### Microtiter plate biofilm assay

The bacterial cultures were inserted into each well of the 96-well plates and then incubated at 37°C for 24 h under anaerobic conditions. The supernatant was discarded, leaving a thin layer on the surface of the well. The wells were rinsed using phosphate-buffered saline (PBS) (Biomatrix, Ontario, Canada). Each test solution (200 µl) was added to each well and then incubated for 1, 3, and 24 h at 37°C. The wells were rinsed twice with PBS and fixated over burning spirit lamp. Crystal violet (Merck, Darmstadt, Germany) (200 µl) was then added to each well and allowed to stand for 15 min, followed by rinsing twice and standing for 15 min. In the past step, 200 µl of 96% ethanol was added. The OD was measured using a microplate reader (Safas, Monaco) with a wavelength of 490 nm.

### Statistical analysis

The research data were processed using the Statistical Package for the Social Sciences (SPSS) computer program, version 26 (IBM, Armonk, NY, USA). The Shapiro–Wilk method was used to test the normality of the data. If the data were normally distributed ( $P > 0.05$ ), a one-way analysis of variance test was conducted, followed by Turkey's honestly significant difference test (significance level of  $P < 0.05$ ) to verify the significance between the groups.

## Results

### Phytochemical screening

The results of the phytochemical screening qualitatively proved that the ethanolic extract of *M. indica* L. leaves contains alkaloids, saponins, tannins, phenolics, flavonoids, and steroids [Table 1].

**Table 1: The results of phytochemical screening of the ethanol extract of *Mangifera indica* L. leaves**

Secondary metabolites	Screening result
Alkaloids	+
Saponins	+
Tannins	+
Phenolics	+
Flavonoids	+
Triterpenoids	–
Steroids	+
Glycoside	–

### Antibacterial test using microdilution and plate count methods

The antibacterial test results showed that the ethanolic extract of *M. indica* L. leaves at 100% concentration and the combination of amoxicillin with extracts of various concentrations exhibited antibacterial and antibiofilm effects against *S. aureus* and *P. gingivalis* [Figure 1] in terms of inhibition of *S. aureus*, the results obtained by the combination of amoxicillin and the extract were not significantly different from those obtained using the positive control ( $P > 0.05$ ). The extract at a concentration of 12.5% and higher was effective against *S. aureus*, and the extract at a concentration of 3.125% and higher was effective against *P. gingivalis* [Figure 3].

### Antibiofilm test using the microtiter plate biofilm assay method

The results of the antibiofilm test showed that the addition of extracts to amoxicillin in inhibiting *S. aureus* biofilms had a significantly lower OD value compared to the positive control ( $P < 0.05$ ), starting at a concentration of 25% after 1 h of incubation [Figure 4] and at a concentration of 6.25% after 3 h of incubation [Figure 5]. After 24 h of incubation, following OD measurement, the group with the combination of extract concentration of 100% and amoxicillin proved not to be significantly different from OD of the positive control ( $P > 0.05$ ) [Figure 6]. In the *P. gingivalis* biofilm, the OD values of the addition of amoxicillin and the extract group, starting at a concentration of 25% after 1, 3, and 24 h of incubation, were smaller than the OD values of the positive control amoxiclav. The OD values of the treatment group were significantly different from those of the positive control amoxiclav [Figures 7-9].

## Discussion

As shown by our results, the ethanol extract of *M. indica* L. leaves contains secondary metabolites, including alkaloids, saponins, tannins, phenolics, flavonoids, and steroids. Different mechanisms of action of each compound account for the antibacterial and antibiofilm properties of the extract. Alkaloids inhibit the formation of peptidoglycan in bacterial cells. The alkaloids can disrupt




Figure 1: The results of the inhibition test of *Staphylococcus aureus* and *Porphyromonas gingivalis* using the plate count method




Figure 2: Graph of the average total colony of *Staphylococcus aureus* (\* $P < 0.05$ , \*\* $P < 0.01$ )




Figure 3: Graph of the average total colony of *Porphyromonas gingivalis* (\* $P < 0.05$ , \*\* $P < 0.01$ )

the amino acid structure of bacterial DNA, leading to bacterial lysis.<sup>[22]</sup> Saponins damage the cell membrane and cell wall permeability in the diffusion process, resulting in the release of enzymes, amino acids, nutrients, and water, leading to cell destabilization and cell death.<sup>[23]</sup> Tannins form a complex with protein in the cell wall, namely, proline, which can damage the cell wall.<sup>[22]</sup> The most

abundant phenolic compound in *M. indica L* leaves is mangiferin. Mangiferin belongs to the xanthone C-glucosyl group, which can damage cell structure and cell membranes and inhibit bacterial protein synthesis.<sup>[24]</sup> Previous research reported that mangiferin compounds interfere with the mechanism of drug resistance; thus, restoring bactericidal and bacteriostatic effects of nalidixic acid, ampicillin, tetracycline, and sulfamethoxazole/trimethoprim.<sup>[25]</sup> However, the exact mechanism of how mangiferin interferes with the mechanism of drug resistance is yet to be elucidated. Flavonoids, as antibacterial, play a role in disrupting the activity of cell wall formation by suppressing cytoplasm function, interrupting the nutrient exchange process, and thereby inhibiting the energy supply of bacteria.<sup>[26]</sup> In addition, flavonoids inhibit enzymes from producing quorum-sensing signals, thereby disrupting the communication process between cells during biofilm formation.<sup>[27]</sup> Steroids can cause leakage of lysosomes and membrane phospholipids that reduce the integrity of cell membranes and lead to cell lysis.<sup>[28]</sup>

Thus far, only a few studies have focused on antibacterial and antibiofilm properties of ethanolic extract of *M. indica L* leaves.<sup>[19,20]</sup> The concentration of the extract used in this study ranged from 3.125% to 100%. As a positive control, co-amoxiclav, a combination of amoxicillin and clavulanic acid, was used. Amoxicillin, a  $\beta$ -lactam antibiotic, works by inhibiting the synthesis of bacterial cell walls.<sup>[11]</sup> The release of  $\beta$ -lactamase enzymes by *S. aureus* and *P. gingivalis* decrease the antibacterial effect of amoxicillin.<sup>[2,12]</sup> The addition of clavulanic acid binds to  $\beta$ -lactamase enzymes from bacteria, inhibiting the enzyme thereby unable to cleave  $\beta$ -lactam ring in amoxicillin, so the amoxicillin can still exhibit its antibacterial activity.<sup>[11]</sup>

In the microdilution and plate count tests, the addition of the ethanolic extract of *M. indica L* leaves to amoxicillin inhibited the growth of *S. aureus* and *P. gingivalis*, and the combination of the ethanolic extract and amoxicillin was as effective as that of co-amoxiclav. In terms of antibacterial activity, a combination of ethanolic extract at




Figure 4: Graph of the average OD value of *Staphylococcus aureus* biofilm after 1 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). OD: Optical density




Figure 5: Graph of the average OD value of *Staphylococcus aureus* biofilm after 3 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). OD: Optical density




Figure 6: Graph of the average OD value of *Staphylococcus aureus* biofilm after 24 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). OD: Optical density




Figure 7: Graph of the average OD value of *Porphyromonas gingivalis* biofilm after 1 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). OD: Optical density




Figure 8: Graph of the average OD value of *Porphyromonas gingivalis* biofilm after 3 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). OD: Optical density




Figure 9: Graph of the average OD value of *Porphyromonas gingivalis* biofilm after 24 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). OD: Optical density

a concentration of 12.5% and amoxicillin was as effective as co-amoxiclav against *S. aureus*, and a concentration as low as 3.125% was as effective as co-amoxiclav against *P. gingivalis*. Therefore, ethanolic extract of *M. indica* L. leaves may be a potential  $\beta$ -lactamase inhibitor equivalent to clavulanic acid. The results of this study are in line with the research of Hartanto *et al.*, who showed that adding methanolic extract of *M. indica* L. leaves to clindamycin

has antibacterial effects against *S. aureus*, especially at a concentration of 100%.<sup>[19]</sup> The bioactive component of *M. indica* L. leaves, namely, mangiferin, is known to have a synergistic effect on tetracycline, ampicillin, nalidixic acid, and trimethoprim in inhibiting the growth of *S. aureus*.<sup>[25]</sup>

The biofilm formation phase begins with pellicle formation, which occurs in the first few seconds to the 1<sup>st</sup> min from

the initial contact, with the initial adhesion phase occurring 2–4 h later. After 24 h, the biofilm enters the maturation phase, becoming 1000-1500 times more resistant than planktonic bacteria.<sup>[29]</sup> The incubation time used in the antibiofilm test in this study was adjusted to the stage of biofilm formation, namely 1, 3, and 24 h. This timing aimed to determine at which stage of biofilm formation amoxicillin and the ethanolic extract of *M. indica* L. leaves most effectively inhibited *S. aureus* and *P. gingivalis*. In the antibiofilm test, the OD from the combination of amoxicillin with ethanolic extract of *M. indica* L. leaves starting at a concentration of 25% after 1 h of incubation and 6.25% after 3 h of incubation had a significantly lower OD value compared to co-amoxiclav. This result proved that the ethanolic extract of *M. indica* L. leaves at a concentration of 25% after 1 h of incubation and 6.25% after 3 h of incubation was more effective at inhibiting *S. aureus* biofilms than co-amoxiclav. The OD values for the combination of amoxicillin and the extract at a concentration of 100% after 24 h of incubation were lower than those obtained for co-amoxiclav, but the difference was not statistically significant. Therefore, the combination of amoxicillin and an extract concentration of 100% has antibiofilm properties equivalent to those of co-amoxiclav after 24 h of incubation.

In terms of inhibiting *P. gingivalis* biofilm, the extract at a concentration of 25% and higher during the entire incubation period showed a more potent antibiofilm effect than that of co-amoxiclav. The combination of amoxicillin and the ethanolic extract of *M. indica* L. leaves inhibited the formation of *S. aureus* biofilms at a concentration of 6.25% after 3 h of incubation and *P. gingivalis* biofilms at a concentration of 25% after 1 h of incubation. This study supports the findings of previous research, which reported that ethanolic extract of *M. indica* L. leaves reduced *S. aureus* attachment and the number of *S. aureus* biofilms.<sup>[30]</sup>

This study has several limitations. First, the ethanolic extract used in this study was a crude extract. The use of a more refined extract, such as an extract exposed to extraction chromatography, would have resulted in an extract with fewer impurities. Second, only two of the many known oral pathogens were used in this study. Other oral pathogens can also be tested to expand the antibacterial activity of amoxicillin combined with *M. indica* L. leaf ethanolic extract. Further research is needed to determine the toxicity of this combination. Preclinical and clinical tests should also be conducted before the combination can be used as an alternative treatment for dentoalveolar abscesses and periodontitis.

## Conclusions

Ethanolic extract of *M. indica* L. leaves contains alkaloids, saponins, tannins, phenolics, flavonoids, and steroids that have the potential to inhibit the growth and formation of *S. aureus* and *P. gingivalis* biofilms *in vitro*. Within the

limitations of this preliminary study, we conclude that the addition of ethanolic extract of *M. indica* L. leaves to amoxicillin could potentially increase the antibacterial and antibiofilm properties of amoxicillin against *S. aureus* and *P. gingivalis*.

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## References

- Ministry of Health of the Republic of Indonesia. Report on Result of National Basic Health Research 2018. Jakarta: Ministry of Health; 2018. p. 197-207.
- Japoni A, Vasin A, Noushadi S, Kiany F, Japoni S, Alborzi A. Antibacterial susceptibility patterns of *Porphyromonas gingivalis* isolated from chronic periodontitis patients. *Med Oral Patol Oral Cir Bucal* 2011;16:e1031-5.
- Shweta N, Prakash SK. Dental abscess: A microbiological review. *Dent Res J (Isfahan)* 2013;10:585-91.
- Mehrotra N, Singh S. Periodontitis. Treasure Island (FL): StatPearls Publishing; 2011.
- Archer NK, Mazaitis MJ, Costerton JW, Leid JG, Powers ME, Shirtliff ME. *Staphylococcus aureus* biofilms: Properties, regulation, and roles in human disease. *Virulence* 2011;2:445-59.
- Huang R, Li M, Gregory RL. Bacterial interactions in dental biofilm. *Virulence* 2011;2:435-44.
- Periasamy S, Joo HS, Duong AC, Bach TH, Tan VY, Chatterjee SS, et al. How *Staphylococcus aureus* biofilms develop their characteristic structure. *Proc Natl Acad Sci U S A* 2012;109:1281-6.
- Taylor TA, Unakal CG. *Staphylococcus aureus*. Treasure Island (FL): StatPearls Publishing; 2021.
- How KY, Song KP, Chan KG. *Porphyromonas gingivalis*: An overview of periodontopathic pathogen below the gum line. *Front Microbiol* 2016;7:53.
- Ciancio SG, Mariotti AJ. Systemic Anti-infective therapy for periodontal diseases. In: Carranza F, Newman M, Takei H, Klokkevold P, editors. *Newman and Carranza's Clinical Periodontology*. 13<sup>th</sup> ed. Philadelphia: Elsevier; 2019. p. 555-7.
- Akhavan B, Khanna N, Vijhani P. Amoxicillin. Treasure Island (FL): StatPearls Publishing; 2020.
- Foster TJ, Geoghegan JA. *Staphylococcus aureus*. In: *Molecular Medical Microbiology*. ???; Elsevier; 2015. p. 655-74.
- Samaranayake L. *Essential Microbiology for Dentistry*. 5<sup>th</sup> ed. Poland: Elsevier Ltd; 2018. p. 75, 158, 293, 297.
- Joshua M, Takudzwa M. Antibacterial properties of *Mangifera indica* on *Staphylococcus aureus*. *Afr J Clin Exp Microbiol* 2013;14:62-74.
- Kalita P. An overview on *Mangifera indica*: Importance and its various pharmacological action. *Pharma Tutoring* 2014;2:72-6.
- Sivakumar D, Jiang Y, Yahia EM. Maintaining mango (*Mangifera indica* L.) fruit quality during the export chain. *Food Res Int* 2011;44:1254-63.
- Chiumarelli M, Ferrari CC, Sarantopoulos CI, Hubinger MD. Fresh cut *Tommy Atkins* mango pre-treated with citric acid and coated with cassava (*Manihot esculenta* Crantz) starch or sodium alginate. *Innov Food Sci Emerg Technol* 2011;12:381-7.
- Gu C, Yang M, Zhou Z, Khan A, Cao J, Cheng G. Purification

and characterization of four benzophenone derivatives from *Mangifera indica* L. leaves and their antioxidant, immunosuppressive and  $\alpha$ -glucosidase inhibitory activities. *J Funct Foods* 2019;52:709-14.

- AQ8 19. Hartanto R, Tran V, Khang G, Pham T, Trinh T, Denhara C. Efek penambahan ekstrak daun mangga arumanis (*Mangifera indica* L.) pada antibiotik klindamisin dalam menghambat pertumbuhan bakteri *Stapylococcus aureus* (Effect of the addition of arumanis mango leaf extract (*Mangifera indica* L.) on clindamycin antibiotics in inhibiting the growth of *Stapylococcus aureus*). *J Prima Med Sains* 2020;2:14-7.
- AQ8 20. Kurniasih R. Pengaruh Konsentrasi Ekstrak Etanol Daun Mangga Arumanis Muda (*Mangifera Indica* L.) Terhadap Hambatan Pertumbuhan Bakteri *Streptococcus Mutans In Vitro* (Effect of Concentrations of Young Mango Arumanis Ethanolic Extract (*Mangifera indica* L.) Leaves on Growth Inhibition of *Streptococcus Mutans in vitro*). Surakarta: Fakultas Kedokteran Gigi Universitas Muhammadiyah (Faculty of Dentistry, Muhammadiyah University); 2016.
21. Yamanaka T, Furukawa T, Matsumoto-Mashimo C, Yamane K, Sugimori C, Nambu T, et al. Gene expression profile and pathogenicity of biofilm-forming *Prevotella intermedia* strain 17. *BMC Microbiol* 2009;9:11.
22. Sylvana D, Amir M, Purnamasari CB, Iskandar A, Asfirizal V. Antibacterial activity of ethanol extract of beluntas leaves on *Streptococcus mutans*, *Porphyromonas gingivalis*, and *Enterococcus faecalis*. *Padjadjaran J Dent* 2021;33:191-8.

23. Sebastian J, Widyarman AS. Roselle flower petals extract inhibits periodontal pathogenic biofilms. *J Dentomaxillofacial Sci* 2021;6:102-5.
24. Tirado-Kulieva V, Atoche-Dioses S, Hernandez-Martinez E. Phenolic compounds of mango (*Mangifera indica*) by-products: Antioxidant and antimicrobial potential, use in disease prevention and food industry, methods of extraction, and microencapsulation. *Sci Agropecu* 2021;12:283-93.
25. Mazlan NA, Azman S, Ghazali NF, Zarith P, Yusri S. Synergistic antibacterial activity of mangiferin with antibiotics against *Staphylococcus aureus*. *Drug Invent Today* 2019;12:14-7.
26. Ma Y, Ding S, Fei Y, Liu G, Jang H, Fang J. Antimicrobial activity of anthocyanins and catechins against foodborne pathogens *Escherichia coli* and *Salmonella*. *Food Control* 2019;106:78-80.
27. Federika AS, Rukmo M, Setyabudi S. Antibiofilm activity of flavonoid mangosteen pericarp extract against *Porphyromonas gingivalis* bacteria. *Conserv Dent J* 2020;10:27-30.
28. Madduluri S, Rao KB, Sitaram B. *In vitro* evaluation of antibacterial activity of five indigenous plants extracts against five bacteria pathogens of humans. *Int J Pharm Pharm Sci* 2013;5:679-84.
29. Bjarnsholt T. The role of bacterial biofilms in chronic infections. *APMIS Suppl* 2013;???:1-51.
30. Manzur AG, Sm Junior V, Morais-Costa F, Mariano EG, Careli RT, da Silva LM, et al. Extract of *Mangifera indica* L. leaves may reduce biofilms of *Staphylococcus* spp. in stainless steel and teatcup rubbers. *Food Sci Technol Int* 2020;26:11-20.

#### Author Queries???

AQ1: Please provide running title

AQ2: Please check article title provided in front page “The Antibacterial and Antibiofilm Effect of Amoxicillin and *Mangifera Indica* L. Leaves Extract on Oral Pathogens” please confirm any one article title.

AQ3: Kindly provide author full name.

AQ4: The author “Yasnill” has not agreed to the copyright terms and conditions which was sent on the authors email address.

AQ5: Kindly provide the department.

AQ6: Kindly mention the significant \* in the image.

AQ7: Please provide publisher location.

AQ8: Kindly provide English Language.

AQ9: Please provide complete reference details such as volume.

AQ10: Please provide citation for figure 2 in the text part

# The Antibacterial and Antibiofilm Effect of Amoxicillin and *Mangifera indica* L. Leaves Extract on Oral Pathogens

## Abstract

**Objective:** This study aimed to determine the antibacterial and antibiofilm effects of amoxicillin combined with extract of *Mangifera indica* L. leaves against *Staphylococcus aureus*, and *Porphyromonas gingivalis*. **Materials and Methods:** This was an experimental laboratory *in vitro* study with a posttest-only control group design. An antibacterial test using the plate count method and an antibiofilm test using the microtiter plate biofilm assay method were conducted. The research samples comprised extract of *M. indica* L. leaves with a concentration of 100%; amoxicillin and extract concentrations of 3.125%, 6.25%, 12.5%, 25%, 50%, and 100%; and amoxicillin. Dimethyl sulfoxide served as a negative control and co-amoxiclav served as a positive control. **Results:** The combination of amoxicillin and the extract exhibited an antibacterial effect against *S. aureus* at a concentration of 12.5% and higher and more effective against *P. gingivalis* at a concentration of 3.125% and higher. In the antibiofilm test, the combination of amoxicillin and the extract at a concentration of 25% after 1 h of incubation and a concentration of 6.25% after 3 h of incubation inhibited *S. aureus*. The inhibition of *S. aureus* biofilms at a concentration of 100% after 24 h of incubation was as effective as that of co-amoxiclav. The extract at a concentration of 25% over the entire incubation period showed inhibition against the *P. gingivalis* biofilm. **Conclusions:** The ethanolic extract of *M. indica* L. leaves and the combination of amoxicillin and the extract have the potential to inhibit the growth and formation of *S. aureus* and *P. gingivalis* biofilms.

**Keywords:** Amoxicillin, antibiofilm, arumanis mango, ethanol extract of *Mangifera indica* L. leaves

## Introduction

Based on data from Basic Health Research (Riskesdas) in 2018, Indonesia's dental and oral health problems reached 57.6% of overall health-care problems.<sup>[1]</sup> The latter is influenced by poor oral hygiene, which triggers various diseases, such as dentoalveolar abscesses and periodontitis.<sup>[2,3]</sup> A dentoalveolar abscess is a pathological cavity in the oral cavity that contains pus due to secondary infection caused by caries, trauma, failure of root canal treatment, and poor oral hygiene.<sup>[3]</sup> Periodontitis is a chronic inflammation of the periodontal tissue structure, including gingiva, bone, and the periodontal ligament, which can cause pocket formation, recession, tooth mobility, or tooth loss.<sup>[2]</sup> Dentoalveolar abscesses and periodontitis are associated with bacterial pathogens in biofilms.<sup>[3,4]</sup>

A biofilm is a collection of microbial cells, especially bacteria, attached to the tooth

surface and coated by an extracellular polymeric substance. This coating protects cells and enables accelerated growth rates, along with additional horizontal gene transfer between cells within the coating, which promotes additional problems, such as antibiotic resistance.<sup>[5]</sup> The formation of biofilms in the oral cavity involves complex competition between microflora for initial attachment.<sup>[6]</sup> *Staphylococcus aureus*, a Gram-positive bacterium, initiates adherence in biofilm formation and produces multiple layers of biofilm embedded in a glycocalyx layer.<sup>[7,8]</sup> *S. aureus* has several virulence factors (capsules, adhesins, coagulase, hyaluronidase, staphylokinase, enterotoxin, and leucocidin) that support biofilm formation in a dentoalveolar abscess.<sup>[6]</sup> *Porphyromonas gingivalis*, which belongs to anaerobic Gram-negative bacteria, is the second most common colonizing bacteria in biofilms. *P. gingivalis* virulence factors, such as lipopolysaccharides, capsules, fimbriae, outer membrane proteins, proteases, and enzymes, induce the destruction of periodontal tissue, causing periodontitis.<sup>[9]</sup>

**How to cite this article:** Soesanto S, Hepziba ER, Yasnill, Widyarman AS. The antibacterial and antibiofilm effect of amoxicillin and *Mangifera indica* L. leaves extract on oral pathogens. *Contemp Clin Dent* 2023;XX:XX-XX.

Sheila Soesanto<sup>1</sup>,  
Evangelista Rachel  
Hepziba<sup>2</sup>,  
Yasnill<sup>2</sup>,  
Armelia Sari  
Widyarman<sup>3</sup>

<sup>1</sup>Department of Pharmacology,  
Faculty of Dentistry, Universitas  
Trisakti, Jakarta, Indonesia,

<sup>2</sup>Undergraduate Student,  
Faculty of Dentistry, Universitas  
Trisakti, Jakarta, Indonesia,

<sup>3</sup>Department of Microbiology,  
Faculty of Dentistry, Universitas  
Trisakti, Jakarta, Indonesia

Submitted : 26-Jul-2022  
Revised : 06-Jan-2023  
Accepted : 21-Feb-2023  
Published : \*\*\*

**Address for correspondence:**  
Dr. Armelia Sari Widyarman,  
Department of Microbiology,  
Faculty of Dentistry, Universitas  
Trisakti, Kyai Tapa 260, Grogol,  
Jakarta 11440, Indonesia.  
E-mail: armeliasari@trisakti.  
ac.id

Access this article online

Website:  
www.contempclindent.org

DOI: 10.4103/ccd.ccd\_399\_22

Quick Response Code:





Amoxicillin from the penicillin group is often used to treat dentoalveolar abscesses and periodontitis. Amoxicillin is a broad-spectrum antibiotic that works by binding to penicillin-binding proteins in Gram-positive and Gram-negative bacteria and inhibiting the transpeptidation process in bacteria.<sup>[3,10]</sup> Overuse of amoxicillin can lead to side effects, such as hypersensitivity reactions, nausea, vomiting, diarrhea, thrombocytopenia, dermatological disorders, and resistance.<sup>[11]</sup> Resistance to amoxicillin can also occur due to the destruction of the  $\beta$ -lactam ring by  $\beta$ -lactamase enzymes produced by *S. aureus* and *P. gingivalis*.<sup>[2,12]</sup> Therefore, amoxicillin is often combined with clavulanic acid, known as co-amoxiclav, to reduce resistance.<sup>[13]</sup> Clavulanic acid is a  $\beta$ -lactamase inhibitor that works by inactivating the pathogen's  $\beta$ -lactamase, thereby increasing the antibacterial activity of amoxicillin.<sup>[11]</sup> However, the use of co-amoxiclav can cause side effects, such as itching, redness around the mouth, diarrhea, nausea, and idiosyncratic drug-induced liver injury.<sup>[11,13]</sup> Alternative medicines using herbal plants have the potential to minimize these side effects.<sup>[14]</sup>

Arumanis mango (*Mangifera indica* L) is a herbal plant from India cultivated in various tropical and subtropical regions, including Indonesia.<sup>[15,16]</sup> Indonesians cultivate Arumanis mangos for their sweetness, freshness, and fragrance. The fruit contains Vitamins A, B, and C, which are beneficial for health.<sup>[17]</sup> In addition, the seeds, skin, roots, and leaves of *M. indica* L. have various properties in traditional medicine. *M. indica* L. leaves are generally discarded and considered waste, even though these leaves contain secondary metabolites, including phenolics (mangiferin, tannins, and flavonoids), alkaloids, saponins, terpenoids, glycosides, and steroids that have potential antibacterial, antifungal, antiviral, antiparasitic, antioxidant, anti-inflammatory, antitumor, anticancer, and analgesic effects.<sup>[18]</sup>

Previous studies showed that *M. indica* L. leaf extracts reduced the number of *Streptococcus mutans* and improved the antibacterial effect of clindamycin against *S. aureus*.<sup>[19,20]</sup> However, no research has investigated the potential of combining amoxicillin with *M. indica* L. leaf extracts in combating *S. mutans* and *P. gingivalis*. Thus, to address this research gap, this study aimed to determine the effectiveness of amoxicillin and ethanolic extract of Arumanis mango (*M. indica* L.) in inhibiting the growth and formation of *S. aureus* and *P. gingivalis* biofilms.

## Materials and Methods

This experimental *in vitro* study was performed at the Microbiology Center of Research and Education (MiCORE) Laboratory, Faculty of Dentistry, Trisakti University, Jakarta, Indonesia. An ethanolic extract of Arumanis mango (*M. indica* L.) leaves was prepared at the Research Institute for Spices and Medicinal Plants (BALITTRO)

in Bogor, West Java, Indonesia. The test solutions used were 10% dimethyl sulfoxide (DMSO) (negative control), co-amoxiclav (positive control), ethanolic extract of *M. indica* L. leaves with a concentration of 100%, and a combination of amoxicillin with extract concentrations of 3.125%, 6.25%, 12.5%, 25%, 50%, and 100%.

The sample size was calculated using Federer's formula, with n equals number of repetition, and t equals as total test group (8 groups comprised 6 treatment groups, 1 positive control, and 1 negative control). Based on this formula, each treatment group was conducted with four-time repetition for all assays.

### Ethanolic extract of *Mangifera indica* L. leaves

The ethanolic extract of *M. indica* L. leaves was prepared using the maceration method. The mango leaves were washed, dried, and mashed, and the simplicial was then soaked in 70% ethanol at a ratio of 1:5. The maceration process was performed for 2–3 h, and the macerate was then allowed to stand for 24 h and filtered. The extract was diluted with 10% DMSO solution to obtain extract concentrations of 50%, 25%, 12.5%, 6.25%, and 3.125%.

### Preparation of antibiotic solution

Amoxicillin and co-amoxiclav solution were prepared by crushing 500 mg of amoxicillin and 625 mg of co-amoxiclav until smooth, using a mortar and pestle. In total, 1.2 mg of amoxicillin and 1.5 mg of co-amoxiclav were each mixed with 6 ml of sterile distilled water until homogeneous to obtain 200 g/ $\mu$ l of amoxicillin and 250 g/ $\mu$ l of co-amoxiclav.

### Phytochemical tests

Qualitative phytochemical tests were performed at BALITTRO to identify secondary metabolites, such as alkaloids, saponins, tannins, phenolics, flavonoids, triterpenoids, steroids, and glycosides, in the ethanolic extract of *M. indica* L. leaves.

### Bacterial culture

*S. aureus* ATCC 25923 and *P. gingivalis* ATCC 33277 were obtained from the MiCORE Laboratory, Faculty of Dentistry, Trisakti University. *S. aureus* was cultured on brain heart infusion broth medium (Sigma Aldrich, St. Louis, Missouri). *P. gingivalis* was cultured on Tryptone Soya Broth (Sigma Aldrich, St. Louis, Missouri) medium enriched with hemin (5 mg/l), Vitamin K (10 mg/l), 0.5% yeast extract, and L-cystine (400 mg/l).<sup>[21]</sup> The medium was incubated in an anaerobic jar (Oxoid, Basingstoke, and Hampshire) at 37° for 24 h under anaerobic conditions. Following the incubation, the bacterial absorbance value was standardized into McFarland standard of 0.5, approximately  $1.5 \times 10^8$  colony forming unit (CFU)/mL (optical density [OD]<sub>600</sub>  $\pm$  0.132), before the following assays.

## Microdilution and total plate count

To evaluate the antibacterial properties of the extract combined with amoxicillin, an antibacterial test was performed on the plate count by the microdilution method. In total, 100 µL of cultured *S. aureus* ATCC 25923 and *P. gingivalis* ATCC 33277 were distributed into 96-well plates (Nest Biotech, Jiangsu, China). A test solution of 100 µl was added to each well and incubated at 37°C for 24 h under anaerobic conditions. After incubation, each mixture containing treated bacteria was diluted 10,000 times. Five microliters of diluted mixture were then spread on a petri dish containing sterile brain heart infusion agar media. The growth of bacterial colonies was calculated after incubation at 37°C for 24 h.

## Microtiter plate biofilm assay

The bacterial cultures were inserted into each well of the 96-well plates and then incubated at 37°C for 24 h under anaerobic conditions. The supernatant was discarded, leaving a thin layer on the surface of the well. The wells were rinsed using phosphate-buffered saline (PBS) (Biomatics, Ontario, Canada). Each test solution (200 µl) was added to each well and then incubated for 1, 3, and 24 h at 37°C. The wells were rinsed twice with PBS and fixated over burning spirit lamp. Crystal violet (Merck, Darmstadt, Germany) (200 µl) was then added to each well and allowed to stand for 15 min, followed by rinsing twice and standing for 15 min. In the past step, 200 µl of 96% ethanol was added. The OD was measured using a microplate reader (Safas, Monaco) with a wavelength of 490 nm.

## Statistical analysis

The research data were processed using the Statistical Package for the Social Sciences (SPSS) computer program, version 26 (IBM, Armonk, NY, USA). The Shapiro–Wilk method was used to test the normality of the data. If the data were normally distributed ( $P > 0.05$ ), a one-way analysis of variance test was conducted, followed by Turkey's honestly significant difference test (significance level of  $P < 0.05$ ) to verify the significance between the groups.

## Results

### Phytochemical screening

The results of the phytochemical screening qualitatively proved that the ethanolic extract of *M. indica* L. leaves contains alkaloids, saponins, tannins, phenolics, flavonoids, and steroids [Table 1].

### Antibacterial test using microdilution and plate count methods

The antibacterial test results showed that the ethanolic extract of *M. indica* L. leaves at 100% concentration and the combination of amoxicillin with extracts of various

**Table 1: The results of phytochemical screening of the ethanol extract of *Mangifera indica* L. leaves**

Secondary metabolites	Screening result
Alkaloids	+
Saponins	+
Tannins	+
Phenolics	+
Flavonoids	+
Triterpenoids	–
Steroids	+
Glycoside	–

concentrations exhibited antibacterial and antibiofilm effects against *S. aureus* and *P. gingivalis* [Figure 1]. In terms of inhibition of *S. aureus*, the results obtained by the combination of amoxicillin and the extract were not significantly different from those obtained using the positive control ( $P > 0.05$ ). The extract at a concentration of 12.5% and higher was effective against *S. aureus*, and the extract at a concentration of 3.125% and higher was effective against *P. gingivalis* [Figures 2 and 3].

### Antibiofilm test using the microtiter plate biofilm assay method

The results of the antibiofilm test showed that the addition of extracts to amoxicillin in inhibiting *S. aureus* biofilms had a significantly lower OD value compared to the positive control ( $P < 0.05$ ), starting at a concentration of 25% after 1 h of incubation [Figure 4] and at a concentration of 6.25% after 3 h of incubation [Figure 5]. After 24 h of incubation, following OD measurement, the group with the combination of extract concentration of 100% and amoxicillin proved not to be significantly different from OD of the positive control ( $P > 0.05$ ) [Figure 6]. In the *P. gingivalis* biofilm, the OD values of the addition of amoxicillin and the extract group, starting at a concentration of 25% after 1, 3, and 24 h of incubation, were smaller than the OD values of the positive control amoxiclav. The OD values of the treatment group were significantly different from those of the positive control amoxiclav [Figures 7-9].

## Discussion

As shown by our results, the ethanol extract of *M. indica* L. leaves contains secondary metabolites, including alkaloids, saponins, tannins, phenolics, flavonoids, and steroids. Different mechanisms of action of each compound account for the antibacterial and antibiofilm properties of the extract. Alkaloids inhibit the formation of peptidoglycan in bacterial cells. The alkaloids can disrupt the amino acid structure of bacterial DNA, leading to bacterial lysis.<sup>[22]</sup> Saponins damage the cell membrane and cell wall permeability in the diffusion process, resulting in the release of enzymes, amino acids, nutrients, and water, leading to cell destabilization and cell death.<sup>[23]</sup> Tannins form a complex with protein in the cell wall, namely,




Figure 1: The results of the inhibition test of *Staphylococcus aureus* and *Porphyromonas gingivalis* using the plate count method




Figure 2: Graph of the average total colony of *Staphylococcus aureus* (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant




Figure 3: Graph of the average total colony of *Porphyromonas gingivalis* (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant

proline, which can damage the cell wall.<sup>[22]</sup> The most abundant phenolic compound in *M. indica* L leaves is mangiferin. Mangiferin belongs to the xanthone C-glucosyl group, which can damage cell structure and cell membranes and inhibit bacterial protein synthesis.<sup>[24]</sup> Previous research reported that mangiferin compounds interfere with the mechanism of drug resistance; thus, restoring

bactericidal and bacteriostatic effects of nalidixic acid, ampicillin, tetracycline, and sulfamethoxazole/trimethoprim.<sup>[25]</sup> However, the exact mechanism of how mangiferin interferes with the mechanism of drug resistance is yet to be elucidated. Flavonoids, as antibacterial, play a role in disrupting the activity of cell wall formation by suppressing cytoplasm function, interrupting the nutrient exchange process, and thereby inhibiting the energy supply of bacteria.<sup>[26]</sup> In addition, flavonoids inhibit enzymes from producing quorum-sensing signals, thereby disrupting the communication process between cells during biofilm formation.<sup>[27]</sup> Steroids can cause leakage of lysosomes and membrane phospholipids that reduce the integrity of cell membranes and lead to cell lysis.<sup>[28]</sup>

Thus far, only a few studies have focused on antibacterial and antibiofilm properties of ethanolic extract of *M. indica* L. leaves.<sup>[19,20]</sup> The concentration of the extract used in this study ranged from 3.125% to 100%. As a positive control, co-amoxiclav, a combination of amoxicillin and clavulanic acid, was used. Amoxicillin, a  $\beta$ -lactam antibiotic, works by inhibiting the synthesis of bacterial cell walls.<sup>[11]</sup> The release of  $\beta$ -lactamase enzymes by *S. aureus* and *P. gingivalis* decrease the antibacterial effect of amoxicillin.<sup>[2,12]</sup> The addition of clavulanic acid binds to  $\beta$ -lactamase enzymes from bacteria, inhibiting the enzyme thereby unable to cleave  $\beta$ -lactam ring in amoxicillin, so the amoxicillin can still exhibit its antibacterial activity.<sup>[11]</sup>

In the microdilution and plate count tests, the addition of the ethanolic extract of *M. indica* L. leaves to amoxicillin inhibited the growth of *S. aureus* and *P. gingivalis*, and the combination of the ethanolic extract and amoxicillin was as effective as that of co-amoxiclav. In terms of antibacterial activity, a combination of ethanolic extract at a concentration of 12.5% and amoxicillin was as effective as co-amoxiclav against *S. aureus*, and a concentration as low as 3.125% was as effective as co-amoxiclav against *P. gingivalis*. Therefore, ethanolic extract of *M. indica* L. leaves may be a potential  $\beta$ -lactamase inhibitor equivalent to clavulanic acid. The results of this study are in line with the research of Hartanto *et al.*, who showed that adding




Figure 4: Graph of the average OD value of *Staphylococcus aureus* biofilm after 1 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant. OD: Optical density




Figure 5: Graph of the average OD value of *Staphylococcus aureus* biofilm after 3 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant. OD: Optical density




Figure 6: Graph of the average OD value of *Staphylococcus aureus* biofilm after 24 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant. OD: Optical density




Figure 7: Graph of the average OD value of *Porphyromonas gingivalis* biofilm after 1 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant. OD: Optical density




Figure 8: Graph of the average OD value of *Porphyromonas gingivalis* biofilm after 3 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant. OD: Optical density




Figure 9: Graph of the average OD value of *Porphyromonas gingivalis* biofilm after 24 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant. OD: Optical density

methanolic extract of *M. indica* L. leaves to clindamycin has antibacterial effects against *S. aureus*, especially at a concentration of 100%.<sup>[19]</sup> The bioactive component of *M. indica* L. leaves, namely, mangiferin, is known to have a synergistic effect on tetracycline, ampicillin, nalidixic acid, and trimethoprim in inhibiting the growth of *S. aureus*.<sup>[25]</sup>

The biofilm formation phase begins with pellicle formation, which occurs in the first few seconds to the 1<sup>st</sup> min from the initial contact, with the initial adhesion phase occurring 2–4 h later. After 24 h, the biofilm enters the maturation phase, becoming 1000-1500 times more resistant than

planktonic bacteria.<sup>[29]</sup> The incubation time used in the antibiofilm test in this study was adjusted to the stage of biofilm formation, namely 1, 3, and 24 h. This timing aimed to determine at which stage of biofilm formation amoxicillin and the ethanolic extract of *M. indica* L. leaves most effectively inhibited *S. aureus* and *P. gingivalis*. In the antibiofilm test, the OD from the combination of amoxicillin with ethanolic extract of *M. indica* L. leaves starting at a concentration of 25% after 1 h of incubation and 6.25% after 3 h of incubation had a significantly lower OD value compared to co-amoxiclav. This result proved that the ethanolic extract of *M. indica* L. leaves at a concentration of 25% after 1 h of incubation and 6.25% after 3 h of incubation was more effective at inhibiting *S. aureus* biofilms than co-amoxiclav. The OD values for the combination of amoxicillin and the extract at a concentration of 100% after 24 h of incubation were lower than those obtained for co-amoxiclav, but the difference was not statistically significant. Therefore, the combination of amoxicillin and an extract concentration of 100% has antibiofilm properties equivalent to those of co-amoxiclav after 24 h of incubation.

In terms of inhibiting *P. gingivalis* biofilm, the extract at a concentration of 25% and higher during the entire incubation period showed a more potent antibiofilm effect than that of co-amoxiclav. The combination of amoxicillin and the ethanolic extract of *M. indica* L. leaves inhibited the formation of *S. aureus* biofilms at a concentration of 6.25% after 3 h of incubation and *P. gingivalis* biofilms at a concentration of 25% after 1 h of incubation. This study supports the findings of previous research, which reported that ethanolic extract of *M. indica* L. leaves reduced *S. aureus* attachment and the number of *S. aureus* biofilms.<sup>[30]</sup>

This study has several limitations. First, the ethanolic extract used in this study was a crude extract. The use of a more refined extract, such as an extract exposed to extraction chromatography, would have resulted in an extract with fewer impurities. Second, only two of the many known oral pathogens were used in this study. Other oral pathogens can also be tested to expand the antibacterial activity of amoxicillin combined with *M. indica* L. leaf ethanolic extract. Further research is needed to determine the toxicity of this combination. Preclinical and clinical tests should also be conducted before the combination can be used as an alternative treatment for dentoalveolar abscesses and periodontitis.

## Conclusions

Ethanolic extract of *M. indica* L. leaves contains alkaloids, saponins, tannins, phenolics, flavonoids, and steroids that have the potential to inhibit the growth and formation of *S. aureus* and *P. gingivalis* biofilms *in vitro*. Within the limitations of this preliminary study, we conclude that the addition of ethanolic extract of *M. indica* L. leaves to amoxicillin could potentially increase the antibacterial and

antibiofilm properties of amoxicillin against *S. aureus* and *P. gingivalis*.

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## References

1. Ministry of Health of the Republic of Indonesia. Report on Result of National Basic Health Research 2018. Jakarta: Ministry of Health; 2018. p. 197-207.
2. Japoni A, Vasin A, Noushadi S, Kiany F, Japoni S, Alborzi A. Antibacterial susceptibility patterns of *Porphyromonas gingivalis* isolated from chronic periodontitis patients. *Med Oral Patol Oral Cir Bucal* 2011;16:e1031-5.
3. Shweta N, Prakash SK. Dental abscess: A microbiological review. *Dent Res J (Isfahan)* 2013;10:585-91.
4. Mehrotra N, Singh S. Periodontitis. Treasure Island (FL): StatPearls Publishing; 2011.
5. Archer NK, Mazaitis MJ, Costerton JW, Leid JG, Powers ME, Shirtliff ME. *Staphylococcus aureus* biofilms: Properties, regulation, and roles in human disease. *Virulence* 2011;2:445-59.
6. Huang R, Li M, Gregory RL. Bacterial interactions in dental biofilm. *Virulence* 2011;2:435-44.
7. Periasamy S, Joo HS, Duong AC, Bach TH, Tan VY, Chatterjee SS, et al. How *Staphylococcus aureus* biofilms develop their characteristic structure. *Proc Natl Acad Sci U S A* 2012;109:1281-6.
8. Taylor TA, Unakal CG. *Staphylococcus aureus*. Treasure Island (FL): StatPearls Publishing; 2021.
9. How KY, Song KP, Chan KG. *Porphyromonas gingivalis*: An overview of periodontopathic pathogen below the gum line. *Front Microbiol* 2016;7:53.
10. Ciancio SG, Mariotti AJ. Systemic Anti-infective therapy for periodontal diseases. In: Carranza F, Newman M, Takei H, Klokkevold P, editors. Newman and Carranza's Clinical Periodontology. 13<sup>th</sup> ed. Philadelphia: Elsevier; 2019. p. 555-7.
11. Akhavan B, Khanna N, Vijhani P. Amoxicillin. Treasure Island (FL): StatPearls Publishing; 2020.
12. Foster TJ, Geoghegan JA. *Staphylococcus aureus*. In: Molecular Medical Microbiology. Massachusetts: Elsevier; 2015. p. 655-74.
13. Samaranayake L. Essential Microbiology for Dentistry. 5<sup>th</sup> ed. Poland: Elsevier Ltd; 2018. p. 75, 158, 293, 297.
14. Joshua M, Takudzwa M. Antibacterial properties of *Mangifera indica* on *Staphylococcus aureus*. *Afr J Clin Exp Microbiol* 2013;14:62-74.
15. Kalita P. An overview on *Mangifera indica*: Importance and its various pharmacological action. *Pharma Tutoring* 2014;2:72-6.
16. Sivakumar D, Jiang Y, Yahia EM. Maintaining mango (*Mangifera indica* L.) fruit quality during the export chain. *Food Res Int* 2011;44:1254-63.
17. Chiumarelli M, Ferrari CC, Sarantopoulos CI, Hubinger MD. Fresh cut *Tommy Atkins* mango pre-treated with citric acid and coated with cassava (*Manihot esculenta* Crantz) starch or sodium alginate. *Innov Food Sci Emerg Technol* 2011;12:381-7.
18. Gu C, Yang M, Zhou Z, Khan A, Cao J, Cheng G. Purification and characterization of four benzophenone derivatives from *Mangifera indica* L. leaves and their antioxidant,

- immunosuppressive and  $\alpha$ -glucosidase inhibitory activities. *J Funct Foods* 2019;52:709-14.
19. Hartanto R, Tran V, Khang G, Pham T, Trinh T, Denhara C. Effect of the addition of arumanis mango leaf extract (*Mangifera indica* L.) on clindamycin antibiotics in inhibiting the growth of *Staphylococcus aureus*. *J Prima Med Sains* 2020;2:14-7.
  20. Kurniasih R. Effect of Concentrations of Young Mango Arumanis Ethanolic Extract (*Mangifera indica* L.) leaves on growth inhibition of *Streptococcus mutans in vitro*. Surakarta: Fakultas Kedokteran Gigi Universitas Muhammadiyah (Faculty of Dentistry, Muhammadiyah University); 2016.
  21. Yamanaka T, Furukawa T, Matsumoto-Mashimo C, Yamane K, Sugimori C, Nambu T, et al. Gene expression profile and pathogenicity of biofilm-forming *Prevotella intermedia* strain 17. *BMC Microbiol* 2009;9:11.
  22. Sylvana D, Amir M, Purnamasari CB, Iskandar A, Asfirizal V. Antibacterial activity of ethanol extract of beluntas leaves on *Streptococcus mutans*, *Porphyromonas gingivalis*, and *Enterococcus faecalis*. *Padjadjaran J Dent* 2021;33:191-8.
  23. Sebastian J, Widyarman AS. Roselle flower petals extract inhibits periodontal pathogenic biofilms. *J Dentomaxillofacial Sci* 2021;6:102-5.
  24. Tirado-Kulieva V, Atoche-Dioses S, Hernandez-Martinez E. Phenolic compounds of mango (*Mangifera indica*) by-products: Antioxidant and antimicrobial potential, use in disease prevention and food industry, methods of extraction, and microencapsulation. *Sci Agropecu* 2021;12:283-93.
  25. Mazlan NA, Azman S, Ghazali NF, Zarith P, Yusri S. Synergistic antibacterial activity of mangiferin with antibiotics against *Staphylococcus aureus*. *Drug Invent Today* 2019;12:14-7.
  26. Ma Y, Ding S, Fei Y, Liu G, Jang H, Fang J. Antimicrobial activity of anthocyanins and catechins against foodborne pathogens *Escherichia coli* and *Salmonella*. *Food Control* 2019;106:78-80.
  27. Federika AS, Rukmo M, Setyabudi S. Antibiofilm activity of flavonoid mangosteen pericarp extract against *Porphyromonas gingivalis* bacteria. *Conserv Dent J* 2020;10:27-30.
  28. Madduluri S, Rao KB, Sitaram B. *In vitro* evaluation of antibacterial activity of five indigenous plants extracts against five bacteria pathogens of humans. *Int J Pharm Pharm Sci* 2013;5:679-84.
  29. Bjarnsholt T. The role of bacterial biofilms in chronic infections. *APMIS Suppl* 2013;121:1-51.
  30. Manzur AG, Sm Junior V, Morais-Costa F, Mariano EG, Careli RT, da Silva LM, et al. Extract of *Mangifera indica* L. leaves may reduce biofilms of *Staphylococcus* spp. in stainless steel and teatcup rubbers. *Food Sci Technol Int* 2020;26:11-20.

Author Queries???

AQ6: Kindly check the significant \* in the image.

# The Antibacterial and Antibiofilm Effect of Amoxicillin and *Mangifera indica* L. Leaves Extract on Oral Pathogens

## Abstract

**Objective:** This study aimed to determine the antibacterial and antibiofilm effects of amoxicillin combined with extract of *Mangifera indica* L. leaves against *Staphylococcus aureus*, *Porphyromonas gingivalis*. **Materials and Methods:** This was an experimental laboratory *in vitro* study with a posttest-only control group design. An antibacterial test using the plate count method and an antibiofilm test using the microtiter plate biofilm assay method were conducted. The research samples comprised extract of *M. indica* L. leaves with a concentration of 100%; amoxicillin and extract concentrations of 3.125%, 6.25%, 12.5%, 25%, 50%, and 100%; and amoxicillin. Dimethyl sulfoxide served as a negative control and co-amoxiclav served as a positive control. **Results:** The combination of amoxicillin and the extract exhibited an antibacterial effect against *S. aureus* at a concentration of 12.5% and higher and **more effective against** *P. gingivalis* at a concentration of 3.125% and higher. In the antibiofilm test, the combination of amoxicillin and the extract at a concentration of 25% after 1 h of incubation and a concentration of 6.25% after 3 h of incubation inhibited *S. aureus*. The inhibition of *S. aureus* biofilms at a concentration of 100% after 24 h of incubation was as effective as that of co-amoxiclav. The extract at a concentration of 25% over the entire incubation period showed inhibition against the *P. gingivalis* biofilm. **Conclusions:** The ethanolic extract of *M. indica* L. leaves and the combination of amoxicillin and the extract have the potential to inhibit the growth and formation of *S. aureus* and *P. gingivalis* biofilms.

**Keywords:** Amoxicillin, antibiofilm, arumanis mango, ethanol extract of *Mangifera indica* L. leaves

## Introduction

Based on data from Basic Health Research (Riskesdas) in 2018, Indonesia's dental and oral health problems reached 57.6% of overall health-care problems.<sup>[1]</sup> The latter is influenced by poor oral hygiene, which triggers various diseases, such as dentoalveolar abscesses and periodontitis.<sup>[2,3]</sup> A dentoalveolar abscess is a pathological cavity in the oral cavity that contains pus due to secondary infection caused by caries, trauma, failure of root canal treatment, and poor oral hygiene.<sup>[3]</sup> Periodontitis is a chronic inflammation of the periodontal tissue structure, including gingiva, bone, and the periodontal ligament, which can cause pocket formation, recession, tooth mobility, or tooth loss.<sup>[2]</sup> Dentoalveolar abscesses and periodontitis are associated with bacterial pathogens in biofilms.<sup>[3,4]</sup>

A biofilm is a collection of microbial cells, especially bacteria, attached to the tooth

surface and coated by an extracellular polymeric substance. This coating protects cells and enables accelerated growth rates, along with additional horizontal gene transfer between cells within the coating, which promotes additional problems, such as antibiotic resistance.<sup>[5]</sup> The formation of biofilms in the oral cavity involves complex competition between microflora for initial attachment.<sup>[6]</sup> *Staphylococcus aureus*, a Gram-positive bacterium, initiates adherence in biofilm formation and produces multiple layers of biofilm embedded in a glycocalyx layer.<sup>[7,8]</sup> *S. aureus* has several virulence factors (capsules, adhesins, coagulase, hyaluronidase, staphylokinase, enterotoxin, and leucocidin) that support biofilm formation in a dentoalveolar abscess.<sup>[6]</sup> *Porphyromonas gingivalis*, which belongs to anaerobic Gram-negative bacteria, is the second most common colonizing bacteria in biofilms. *P. gingivalis* virulence factors, such as lipopolysaccharides, capsules, fimbriae, outer membrane proteins, proteases, and enzymes, induce the destruction of periodontal tissue, causing periodontitis.<sup>[9]</sup>

**How to cite this article:** Soesanto S, Hepziba ER, Yasnill, Widyarman AS. The antibacterial and antibiofilm effect of amoxicillin and *Mangifera indica* L. leaves extract on oral pathogens. *Contemp Clin Dent* 2023;XX:XX-XX.

Sheila Soesanto<sup>1</sup>,  
Evangelista Rachel  
Hepziba<sup>2</sup>,  
Yasnill<sup>2</sup>,  
Armelia Sari  
Widyarman<sup>3</sup>

<sup>1</sup>Department of Pharmacology,  
Faculty of Dentistry, Universitas  
Trisakti, Jakarta, Indonesia,

<sup>2</sup>Undergraduate Student,  
Faculty of Dentistry, Universitas  
Trisakti, Jakarta, Indonesia,

<sup>3</sup>Department of Microbiology,  
Faculty of Dentistry, Universitas  
Trisakti, Jakarta, Indonesia

Submitted : 26-Jul-2022  
Revised : 06-Jan-2023  
Accepted : 21-Feb-2023  
Published : \*\*\*

**Address for correspondence:**  
Dr. Armelia Sari Widyarman,  
Department of Microbiology,  
Faculty of Dentistry, Universitas  
Trisakti, Kyai Tapa 260, Grogol,  
Jakarta 11440, Indonesia.  
E-mail: armeliasari@trisakti.  
ac.id

Access this article online

Website:  
www.contempclindent.org

DOI: 10.4103/ccd.ccd\_399\_22

Quick Response Code:



Amoxicillin from the penicillin group is often used to treat dentoalveolar abscesses and periodontitis. Amoxicillin is a broad-spectrum antibiotic that works by binding to penicillin-binding proteins in Gram-positive and Gram-negative bacteria and inhibiting the transpeptidation process in bacteria.<sup>[3,10]</sup> Overuse of amoxicillin can lead to side effects, such as hypersensitivity reactions, nausea, vomiting, diarrhea, thrombocytopenia, dermatological disorders, and resistance.<sup>[11]</sup> Resistance to amoxicillin can also occur due to the destruction of the  $\beta$ -lactam ring by  $\beta$ -lactamase enzymes produced by *S. aureus* and *P. gingivalis*.<sup>[2,12]</sup> Therefore, amoxicillin is often combined with clavulanic acid, known as co-amoxiclav, to reduce resistance.<sup>[13]</sup> Clavulanic acid is a  $\beta$ -lactamase inhibitor that works by inactivating the pathogen's  $\beta$ -lactamase, thereby increasing the antibacterial activity of amoxicillin.<sup>[11]</sup> However, the use of co-amoxiclav can cause side effects, such as itching, redness around the mouth, diarrhea, nausea, and idiosyncratic drug-induced liver injury.<sup>[11,13]</sup> Alternative medicines using herbal plants have the potential to minimize these side effects.<sup>[14]</sup>

Arumanis mango (*Mangifera indica* L) is a herbal plant from India cultivated in various tropical and subtropical regions, including Indonesia.<sup>[15,16]</sup> Indonesians cultivate Arumanis mangos for their sweetness, freshness, and fragrance. The fruit contains Vitamins A, B, and C, which are beneficial for health.<sup>[17]</sup> In addition, the seeds, skin, roots, and leaves of *M. indica* L. have various properties in traditional medicine. *M. indica* L. leaves are generally discarded and considered waste, even though these leaves contain secondary metabolites, including phenolics (mangiferin, tannins, and flavonoids), alkaloids, saponins, terpenoids, glycosides, and steroids that have potential antibacterial, antifungal, antiviral, antiparasitic, antioxidant, anti-inflammatory, antitumor, anticancer, and analgesic effects.<sup>[18]</sup>

Previous studies showed that *M. indica* L. leaf extracts reduced the number of *Streptococcus mutans* and improved the antibacterial effect of clindamycin against *S. aureus*.<sup>[19,20]</sup> However, no research has investigated the potential of combining amoxicillin with *M. indica* L. leaf extracts in combating *S. mutans* and *P. gingivalis*. Thus, to address this research gap, this study aimed to determine the effectiveness of amoxicillin and ethanolic extract of Arumanis mango (*M. indica* L.) in inhibiting the growth and formation of *S. aureus* and *P. gingivalis* biofilms.

## Materials and Methods

This experimental *in vitro* study was performed at the Microbiology Center of Research and Education (MiCORE) Laboratory, Faculty of Dentistry, Trisakti University, Jakarta, Indonesia. An ethanolic extract of Arumanis mango (*M. indica* L.) leaves was prepared at the Research Institute for Spices and Medicinal Plants (BALITTRO)

in Bogor, West Java, Indonesia. The test solutions used were 10% dimethyl sulfoxide (DMSO) (negative control), co-amoxiclav (positive control), ethanolic extract of *M. indica* L. leaves with a concentration of 100%, and a combination of amoxicillin with extract concentrations of 3.125%, 6.25%, 12.5%, 25%, 50%, and 100%.

The sample size was calculated using Federer's formula, with n equals number of repetition, and t equals as total test group (8 groups comprised 6 treatment groups, 1 positive control, and 1 negative control). Based on this formula, each treatment group was conducted with four-time repetition for all assays.

### Ethanolic extract of *Mangifera indica* L. leaves

The ethanolic extract of *M. indica* L. leaves was prepared using the maceration method. The mango leaves were washed, dried, and mashed, and the simplicial was then soaked in 70% ethanol at a ratio of 1:5. The maceration process was performed for 2–3 h, and the macerate was then allowed to stand for 24 h and filtered. The extract was diluted with 10% DMSO solution to obtain extract concentrations of 50%, 25%, 12.5%, 6.25%, and 3.125%.

### Preparation of antibiotic solution

Amoxicillin and co-amoxiclav solution were prepared by crushing 500 mg of amoxicillin and 625 mg of co-amoxiclav until smooth, using a mortar and pestle. In total, 1.2 mg of amoxicillin and 1.5 mg of co-amoxiclav were each mixed with 6 ml of sterile distilled water until homogeneous to obtain 200 g/ $\mu$ l of amoxicillin and 250 g/ $\mu$ l of co-amoxiclav.

### Phytochemical tests

Qualitative phytochemical tests were performed at BALITTRO to identify secondary metabolites, such as alkaloids, saponins, tannins, phenolics, flavonoids, triterpenoids, steroids, and glycosides, in the ethanolic extract of *M. indica* L. leaves.

### Bacterial culture

*S. aureus* ATCC 25923 and *P. gingivalis* ATCC 33277 were obtained from the MiCORE Laboratory, Faculty of Dentistry, Trisakti University. *S. aureus* was cultured on brain heart infusion broth medium (Sigma Aldrich, St. Louis, Missouri). *P. gingivalis* was cultured on Tryptone Soya Broth (Sigma Aldrich, St. Louis, Missouri) medium enriched with hemin (5 mg/l), Vitamin K (10 mg/l), 0.5% yeast extract, and L-cystine (400 mg/l).<sup>[21]</sup> The medium was incubated in an anaerobic jar (Oxoid, Basingstoke, and Hampshire) at 37° for 24 h under anaerobic conditions. Following the incubation, the bacterial absorbance value was standardized into McFarland standard of 0.5, approximately  $1.5 \times 10^8$  colony forming unit (CFU)/mL (optical density [OD]<sub>600</sub>  $\pm$  0.132), before the following assays.



## Microdilution and total plate count

To evaluate the antibacterial properties of the extract combined with amoxicillin, an antibacterial test was performed on the plate count by the microdilution method. In total, 100  $\mu$ L of cultured *S. aureus* ATCC 25923 and *P. gingivalis* ATCC 33277 were distributed into 96-well plates (Nest Biotech, Jiangsu, China). A test solution of 100  $\mu$ l was added to each well and incubated at 37°C for 24 h under anaerobic conditions. After incubation, each mixture containing treated bacteria was diluted 10,000 times. Five microliters of diluted mixture were then spread on a petri dish containing sterile brain heart infusion agar media. The growth of bacterial colonies was calculated after incubation at 37°C for 24 h.

## Microtiter plate biofilm assay

The bacterial cultures were inserted into each well of the 96-well plates and then incubated at 37°C for 24 h under anaerobic conditions. The supernatant was discarded, leaving a thin layer on the surface of the well. The wells were rinsed using phosphate-buffered saline (PBS) (Biomatics, Ontario, Canada). Each test solution (200  $\mu$ l) was added to each well and then incubated for 1, 3, and 24 h at 37°C. The wells were rinsed twice with PBS and fixated over burning spirit lamp. Crystal violet (Merck, Darmstadt, Germany) (200  $\mu$ l) was then added to each well and allowed to stand for 15 min, followed by rinsing twice and standing for 15 min. In the past step, 200  $\mu$ l of 96% ethanol was added. The OD was measured using a microplate reader (Safas, Monaco) with a wavelength of 490 nm.

## Statistical analysis

The research data were processed using the Statistical Package for the Social Sciences (SPSS) computer program, version 26 (IBM, Armonk, NY, USA). The Shapiro–Wilk method was used to test the normality of the data. If the data were normally distributed ( $P > 0.05$ ), a one-way analysis of variance test was conducted, followed by Turkey's honestly significant difference test (significance level of  $P < 0.05$ ) to verify the significance between the groups.

## Results

### Phytochemical screening

The results of the phytochemical screening qualitatively proved that the ethanolic extract of *M. indica* L. leaves contains alkaloids, saponins, tannins, phenolics, flavonoids, and steroids [Table 1].

### Antibacterial test using microdilution and plate count methods

The antibacterial test results showed that the ethanolic extract of *M. indica* L. leaves at 100% concentration and the combination of amoxicillin with extracts of various

**Table 1: The results of phytochemical screening of the ethanol extract of *Mangifera indica* L. leaves**

Secondary metabolites	Screening result
Alkaloids	+
Saponins	+
Tannins	+
Phenolics	+
Flavonoids	+
Triterpenoids	–
Steroids	+
Glycoside	–

concentrations exhibited antibacterial and antibiofilm effects against *S. aureus* and *P. gingivalis* [Figure 1]. In terms of inhibition of *S. aureus*, the results obtained by the combination of amoxicillin and the extract were not significantly different from those obtained using the positive control ( $P > 0.05$ ). The extract at a concentration of 12.5% and higher was effective against *S. aureus*, and the extract at a concentration of 3.125% and higher was effective against *P. gingivalis* [Figures 2 and 3].

### Antibiofilm test using the microtiter plate biofilm assay method

The results of the antibiofilm test showed that the addition of extracts to amoxicillin in inhibiting *S. aureus* biofilms had a significantly lower OD value compared to the positive control ( $P < 0.05$ ), starting at a concentration of 25% after 1 h of incubation [Figure 4] and at a concentration of 6.25% after 3 h of incubation [Figure 5]. After 24 h of incubation, following OD measurement, the group with the combination of extract concentration of 100% and amoxicillin proved not to be significantly different from OD of the positive control ( $P > 0.05$ ) [Figure 6]. In the *P. gingivalis* biofilm, the OD values of the addition of amoxicillin and the extract group, starting at a concentration of 25% after 1, 3, and 24 h of incubation, were smaller than the OD values of the positive control amoxiclav. The OD values of the treatment group were significantly different from those of the positive control amoxiclav [Figures 7-9].

## Discussion

As shown by our results, the ethanol extract of *M. indica* L. leaves contains secondary metabolites, including alkaloids, saponins, tannins, phenolics, flavonoids, and steroids. Different mechanisms of action of each compound account for the antibacterial and antibiofilm properties of the extract. Alkaloids inhibit the formation of peptidoglycan in bacterial cells. The alkaloids can disrupt the amino acid structure of bacterial DNA, leading to bacterial lysis.<sup>[22]</sup> Saponins damage the cell membrane and cell wall permeability in the diffusion process, resulting in the release of enzymes, amino acids, nutrients, and water, leading to cell destabilization and cell death.<sup>[23]</sup> Tannins form a complex with protein in the cell wall, namely,




Figure 1: The results of the inhibition test of *Staphylococcus aureus* and *Porphyromonas gingivalis* using the plate count method




Figure 2: Graph of the average total colony of *Staphylococcus aureus* (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant




Figure 3: Graph of the average total colony of *Porphyromonas gingivalis* (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant

proline, which can damage the cell wall.<sup>[22]</sup> The most abundant phenolic compound in *M. indica* L leaves is mangiferin. Mangiferin belongs to the xanthone C-glucosyl group, which can damage cell structure and cell membranes and inhibit bacterial protein synthesis.<sup>[24]</sup> Previous research reported that mangiferin compounds interfere with the mechanism of drug resistance; thus, restoring

bactericidal and bacteriostatic effects of nalidixic acid, ampicillin, tetracycline, and sulfamethoxazole/trimethoprim.<sup>[25]</sup> However, the exact mechanism of how mangiferin interferes with the mechanism of drug resistance is yet to be elucidated. Flavonoids, as antibacterial, play a role in disrupting the activity of cell wall formation by suppressing cytoplasm function, interrupting the nutrient exchange process, and thereby inhibiting the energy supply of bacteria.<sup>[26]</sup> In addition, flavonoids inhibit enzymes from producing quorum-sensing signals, thereby disrupting the communication process between cells during biofilm formation.<sup>[27]</sup> Steroids can cause leakage of lysosomes and membrane phospholipids that reduce the integrity of cell membranes and lead to cell lysis.<sup>[28]</sup>

Thus far, only a few studies have focused on antibacterial and antibiofilm properties of ethanolic extract of *M. indica* L. leaves.<sup>[19,20]</sup> The concentration of the extract used in this study ranged from 3.125% to 100%. As a positive control, co-amoxiclav, a combination of amoxicillin and clavulanic acid, was used. Amoxicillin, a  $\beta$ -lactam antibiotic, works by inhibiting the synthesis of bacterial cell walls.<sup>[11]</sup> The release of  $\beta$ -lactamase enzymes by *S. aureus* and *P. gingivalis* decrease the antibacterial effect of amoxicillin.<sup>[2,12]</sup> The addition of clavulanic acid binds to  $\beta$ -lactamase enzymes from bacteria, inhibiting the enzyme thereby unable to cleave  $\beta$ -lactam ring in amoxicillin, so the amoxicillin can still exhibit its antibacterial activity.<sup>[11]</sup>

In the microdilution and plate count tests, the addition of the ethanolic extract of *M. indica* L. leaves to amoxicillin inhibited the growth of *S. aureus* and *P. gingivalis*, and the combination of the ethanolic extract and amoxicillin was as effective as that of co-amoxiclav. In terms of antibacterial activity, a combination of ethanolic extract at a concentration of 12.5% and amoxicillin was as effective as co-amoxiclav against *S. aureus*, and a concentration as low as 3.125% was as effective as co-amoxiclav against *P. gingivalis*. Therefore, ethanolic extract of *M. indica* L. leaves may be a potential  $\beta$ -lactamase inhibitor equivalent to clavulanic acid. The results of this study are in line with the research of Hartanto *et al.*, who showed that adding




Figure 4: Graph of the average OD value of *Staphylococcus aureus* biofilm after 1 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant. OD: Optical density




Figure 5: Graph of the average OD value of *Staphylococcus aureus* biofilm after 3 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant. OD: Optical density




Figure 6: Graph of the average OD value of *Staphylococcus aureus* biofilm after 24 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant. OD: Optical density




Figure 7: Graph of the average OD value of *Porphyromonas gingivalis* biofilm after 1 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant. OD: Optical density




Figure 8: Graph of the average OD value of *Porphyromonas gingivalis* biofilm after 3 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant. OD: Optical density




Figure 9: Graph of the average OD value of *Porphyromonas gingivalis* biofilm after 24 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant. OD: Optical density

methanolic extract of *M. indica* L. leaves to clindamycin has antibacterial effects against *S. aureus*, especially at a concentration of 100%.<sup>[19]</sup> The bioactive component of *M. indica* L. leaves, namely, mangiferin, is known to have a synergistic effect on tetracycline, ampicillin, nalidixic acid, and trimethoprim in inhibiting the growth of *S. aureus*.<sup>[25]</sup>

The biofilm formation phase begins with pellicle formation, which occurs in the first few seconds to the 1<sup>st</sup> min from the initial contact, with the initial adhesion phase occurring 2–4 h later. After 24 h, the biofilm enters the maturation phase, becoming 1000-1500 times more resistant than

planktonic bacteria.<sup>[29]</sup> The incubation time used in the antibiofilm test in this study was adjusted to the stage of biofilm formation, namely 1, 3, and 24 h. This timing aimed to determine at which stage of biofilm formation amoxicillin and the ethanolic extract of *M. indica* L. leaves most effectively inhibited *S. aureus* and *P. gingivalis*. In the antibiofilm test, the OD from the combination of amoxicillin with ethanolic extract of *M. indica* L. leaves starting at a concentration of 25% after 1 h of incubation and 6.25% after 3 h of incubation had a significantly lower OD value compared to co-amoxiclav. This result proved that the ethanolic extract of *M. indica* L. leaves at a concentration of 25% after 1 h of incubation and 6.25% after 3 h of incubation was more effective at inhibiting *S. aureus* biofilms than co-amoxiclav. The OD values for the combination of amoxicillin and the extract at a concentration of 100% after 24 h of incubation were lower than those obtained for co-amoxiclav, but the difference was not statistically significant. Therefore, the combination of amoxicillin and an extract concentration of 100% has antibiofilm properties equivalent to those of co-amoxiclav after 24 h of incubation.

In terms of inhibiting *P. gingivalis* biofilm, the extract at a concentration of 25% and higher during the entire incubation period showed a more potent antibiofilm effect than that of co-amoxiclav. The combination of amoxicillin and the ethanolic extract of *M. indica* L. leaves inhibited the formation of *S. aureus* biofilms at a concentration of 6.25% after 3 h of incubation and *P. gingivalis* biofilms at a concentration of 25% after 1 h of incubation. This study supports the findings of previous research, which reported that ethanolic extract of *M. indica* L. leaves reduced *S. aureus* attachment and the number of *S. aureus* biofilms.<sup>[30]</sup>

This study has several limitations. First, the ethanolic extract used in this study was a crude extract. The use of a more refined extract, such as an extract exposed to extraction chromatography, would have resulted in an extract with fewer impurities. Second, only two of the many known oral pathogens were used in this study. Other oral pathogens can also be tested to expand the antibacterial activity of amoxicillin combined with *M. indica* L. leaf ethanolic extract. Further research is needed to determine the toxicity of this combination. Preclinical and clinical tests should also be conducted before the combination can be used as an alternative treatment for dentoalveolar abscesses and periodontitis.

## Conclusions

Ethanolic extract of *M. indica* L. leaves contains alkaloids, saponins, tannins, phenolics, flavonoids, and steroids that have the potential to inhibit the growth and formation of *S. aureus* and *P. gingivalis* biofilms *in vitro*. Within the limitations of this preliminary study, we conclude that the addition of ethanolic extract of *M. indica* L. leaves to amoxicillin could potentially increase the antibacterial and

antibiofilm properties of amoxicillin against *S. aureus* and *P. gingivalis*.

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## References

- Ministry of Health of the Republic of Indonesia. Report on Result of National Basic Health Research 2018. Jakarta: Ministry of Health; 2018. p. 197-207.
- Japoni A, Vasin A, Noushadi S, Kiany F, Japoni S, Alborzi A. Antibacterial susceptibility patterns of *Porphyromonas gingivalis* isolated from chronic periodontitis patients. *Med Oral Patol Oral Cir Bucal* 2011;16:e1031-5.
- Shweta N, Prakash SK. Dental abscess: A microbiological review. *Dent Res J (Isfahan)* 2013;10:585-91.
- Mehrotra N, Singh S. Periodontitis. Treasure Island (FL): StatPearls Publishing; 2011.
- Archer NK, Mazaitis MJ, Costerton JW, Leid JG, Powers ME, Shirtliff ME. *Staphylococcus aureus* biofilms: Properties, regulation, and roles in human disease. *Virulence* 2011;2:445-59.
- Huang R, Li M, Gregory RL. Bacterial interactions in dental biofilm. *Virulence* 2011;2:435-44.
- Periasamy S, Joo HS, Duong AC, Bach TH, Tan VY, Chatterjee SS, et al. How *Staphylococcus aureus* biofilms develop their characteristic structure. *Proc Natl Acad Sci U S A* 2012;109:1281-6.
- Taylor TA, Unakal CG. *Staphylococcus aureus*. Treasure Island (FL): StatPearls Publishing; 2021.
- How KY, Song KP, Chan KG. *Porphyromonas gingivalis*: An overview of periodontopathic pathogen below the gum line. *Front Microbiol* 2016;7:53.
- Ciancio SG, Mariotti AJ. Systemic Anti-infective therapy for periodontal diseases. In: Carranza F, Newman M, Takei H, Klokkevold P, editors. Newman and Carranza's Clinical Periodontology. 13<sup>th</sup> ed. Philadelphia: Elsevier; 2019. p. 555-7.
- Akhavan B, Khanna N, Vijhani P. Amoxicillin. Treasure Island (FL): StatPearls Publishing; 2020.
- Foster TJ, Geoghegan JA. *Staphylococcus aureus*. In: Molecular Medical Microbiology. Massachusetts: Elsevier; 2015. p. 655-74.
- Samaranayake L. Essential Microbiology for Dentistry. 5<sup>th</sup> ed. Poland: Elsevier Ltd; 2018. p. 75, 158, 293, 297.
- Joshua M, Takudzwa M. Antibacterial properties of *Mangifera indica* on *Staphylococcus aureus*. *Afr J Clin Exp Microbiol* 2013;14:62-74.
- Kalita P. An overview on *Mangifera indica*: Importance and its various pharmacological action. *Pharma Tutoring* 2014;2:72-6.
- Sivakumar D, Jiang Y, Yahia EM. Maintaining mango (*Mangifera indica* L.) fruit quality during the export chain. *Food Res Int* 2011;44:1254-63.
- Chiumarelli M, Ferrari CC, Sarantopoulos CI, Hubinger MD. Fresh cut *Tommy Atkins* mango pre-treated with citric acid and coated with cassava (*Manihot esculenta* Crantz) starch or sodium alginate. *Innov Food Sci Emerg Technol* 2011;12:381-7.
- Gu C, Yang M, Zhou Z, Khan A, Cao J, Cheng G. Purification and characterization of four benzophenone derivatives from *Mangifera indica* L. leaves and their antioxidant,

- immunosuppressive and  $\alpha$ -glucosidase inhibitory activities. *J Funct Foods* 2019;52:709-14.
19. Hartanto R, Tran V, Khang G, Pham T, Trinh T, Denhara C. Effect of the addition of arumanis mango leaf extract (*Mangifera indica* L.) on clindamycin antibiotics in inhibiting the growth of *Staphylococcus aureus*. *J Prima Med Sains* 2020;2:14-7.
  20. Kurniasih R. Effect of Concentrations of Young Mango Arumanis Ethanolic Extract (*Mangifera indica* L.) leaves on growth inhibition of *Streptococcus mutans in vitro*. Surakarta: Fakultas Kedokteran Gigi Universitas Muhammadiyah (Faculty of Dentistry, Muhammadiyah University); 2016.
  21. Yamanaka T, Furukawa T, Matsumoto-Mashimo C, Yamane K, Sugimori C, Nambu T, *et al.* Gene expression profile and pathogenicity of biofilm-forming *Prevotella intermedia* strain 17. *BMC Microbiol* 2009;9:11.
  22. Sylvana D, Amir M, Purnamasari CB, Iskandar A, Asfirizal V. Antibacterial activity of ethanol extract of beluntas leaves on *Streptococcus mutans*, *Porphyromonas gingivalis*, and *Enterococcus faecalis*. *Padjadjaran J Dent* 2021;33:191-8.
  23. Sebastian J, Widyarman AS. Roselle flower petals extract inhibits periodontal pathogenic biofilms. *J Dentomaxillofacial Sci* 2021;6:102-5.
  24. Tirado-Kulieva V, Atoche-Dioses S, Hernandez-Martinez E. Phenolic compounds of mango (*Mangifera indica*) by-products: Antioxidant and antimicrobial potential, use in disease prevention and food industry, methods of extraction, and microencapsulation. *Sci Agropecu* 2021;12:283-93.
  25. Mazlan NA, Azman S, Ghazali NF, Zarith P, Yusri S. Synergistic antibacterial activity of mangiferin with antibiotics against *Staphylococcus aureus*. *Drug Invent Today* 2019;12:14-7.
  26. Ma Y, Ding S, Fei Y, Liu G, Jang H, Fang J. Antimicrobial activity of anthocyanins and catechins against foodborne pathogens *Escherichia coli* and *Salmonella*. *Food Control* 2019;106:78-80.
  27. Federika AS, Rukmo M, Setyabudi S. Antibiofilm activity of flavonoid mangosteen pericarp extract against *Porphyromonas gingivalis* bacteria. *Conserv Dent J* 2020;10:27-30.
  28. Madduluri S, Rao KB, Sitaram B. *In vitro* evaluation of antibacterial activity of five indigenous plants extracts against five bacteria pathogens of humans. *Int J Pharm Pharm Sci* 2013;5:679-84.
  29. Bjarnsholt T. The role of bacterial biofilms in chronic infections. *APMIS Suppl* 2013;121:1-51.
  30. Manzur AG, Sm Junior V, Morais-Costa F, Mariano EG, Careli RT, da Silva LM, *et al.* Extract of *Mangifera indica* L. leaves may reduce biofilms of *Staphylococcus* spp. in stainless steel and teatcup rubbers. *Food Sci Technol Int* 2020;26:11-20.

Author Queries???

AQ6: Kindly check the significant \* in the image.

# The Antibacterial and Antibiofilm Effect of Amoxicillin and *Mangifera indica* L. Leaves Extract on Oral Pathogens

## Abstract

**Objective:** This study aimed to determine the antibacterial and antibiofilm effects of amoxicillin combined with extract of *Mangifera indica* L. leaves against *Staphylococcus aureus*, *Porphyromonas gingivalis*. **Materials and Methods:** This was an experimental laboratory *in vitro* study with a posttest-only control group design. An antibacterial test using the plate count method and an antibiofilm test using the microtiter plate biofilm assay method were conducted. The research samples comprised extract of *M. indica* L. leaves with a concentration of 100%; amoxicillin and extract concentrations of 3.125%, 6.25%, 12.5%, 25%, 50%, and 100%; and amoxicillin. Dimethyl sulfoxide served as a negative control and co-amoxiclav served as a positive control. **Results:** The combination of amoxicillin and the extract exhibited an antibacterial effect against *S. aureus* at a concentration of 12.5% and higher and more effective against *P. gingivalis* at a concentration of 3.125% and higher. In the antibiofilm test, the combination of amoxicillin and the extract at a concentration of 25% after 1 h of incubation and a concentration of 6.25% after 3 h of incubation inhibited *S. aureus* inhibition of *S. aureus* biofilms at a concentration of 100% after 24 h of incubation was as effective as that of co-amoxiclav. The extract at a concentration of 25% over the entire incubation period showed inhibition against the *P. gingivalis* biofilm. **Conclusions:** The ethanolic extract of *M. indica* L. leaves and the combination of amoxicillin and the extract have the potential to inhibit the growth and formation of *S. aureus* and *P. gingivalis* biofilms.

**Keywords:** Amoxicillin, antibiofilm, arumanis mango, ethanol extract of *Mangifera indica* L. leaves

## Introduction

Based on data from Basic Health Research (Riskesdas) in 2018, Indonesia's dental and oral health problems reached 57.6% of overall health-care problems.<sup>[1]</sup> The latter is influenced by poor oral hygiene, which triggers various diseases, such as dentoalveolar abscesses and periodontitis.<sup>[2,3]</sup> A dentoalveolar abscess is a pathological cavity in the oral cavity that contains pus due to secondary infection caused by caries, trauma, failure of root canal treatment, and poor oral hygiene.<sup>[3]</sup> Periodontitis is a chronic inflammation of the periodontal tissue structure, including gingiva, bone, and the periodontal ligament, which can cause pocket formation, recession, tooth mobility, or tooth loss.<sup>[2]</sup> Dentoalveolar abscesses and periodontitis are associated with bacterial pathogens in biofilms.<sup>[3,4]</sup>

A biofilm is a collection of microbial cells, especially bacteria, attached to the tooth

surface and coated by an extracellular polymeric substance. This coating protects cells and enables accelerated growth rates, along with additional horizontal gene transfer between cells within the coating, which promotes additional problems, such as antibiotic resistance.<sup>[5]</sup> The formation of biofilms in the oral cavity involves complex competition between microflora for initial attachment.<sup>[6]</sup> *Staphylococcus aureus*, a Gram-positive bacterium, initiates adherence in biofilm formation and produces multiple layers of biofilm embedded in a glycocalyx layer.<sup>[7,8]</sup> *S. aureus* has several virulence factors (capsules, adhesins, coagulase, hyaluronidase, staphylokinase, enterotoxin, and leucocidin) that support biofilm formation in a dentoalveolar abscess.<sup>[6]</sup> *Porphyromonas gingivalis*, which belongs to anaerobic Gram-negative bacteria, is the second most common colonizing bacteria in biofilms. *P. gingivalis* virulence factors, such as lipopolysaccharides, capsules, fimbriae, outer membrane proteins, proteases, and enzymes, induce the destruction of periodontal tissue, causing periodontitis.<sup>[9]</sup>

**How to cite this article:** Soesanto S, Hepziba ER, Yasnill, Widyarman AS. The antibacterial and antibiofilm effect of amoxicillin and *Mangifera indica* L. leaves extract on oral pathogens. *Contemp Clin Dent* 2023;XX:XX-XX.

Sheila Soesanto<sup>1</sup>,  
Evangelista Rachel  
Hepziba<sup>2</sup>,  
Yasnill<sup>2</sup>,  
Armelia Sari  
Widyarman<sup>3</sup>

<sup>1</sup>Department of Pharmacology,  
Faculty of Dentistry, Universitas  
Trisakti, Jakarta, Indonesia,

<sup>2</sup>Undergraduate Student,  
Faculty of Dentistry, Universitas  
Trisakti, Jakarta, Indonesia,

<sup>3</sup>Department of Microbiology,  
Faculty of Dentistry, Universitas  
Trisakti, Jakarta, Indonesia

Submitted : 26-Jul-2022  
Revised : 06-Jan-2023  
Accepted : 21-Feb-2023  
Published : \*\*\*

**Address for correspondence:**  
Dr. Armelia Sari Widyarman,  
Department of Microbiology,  
Faculty of Dentistry, Universitas  
Trisakti, Kyai Tapa 260, Grogol,  
Jakarta 11440, Indonesia.  
E-mail: armeliasari@trisakti.  
ac.id

Access this article online

Website:  
www.contempclindent.org

DOI: 10.4103/ccd.ccd\_399\_22

Quick Response Code:



Amoxicillin from the penicillin group is often used to treat dentoalveolar abscesses and periodontitis. Amoxicillin is a broad-spectrum antibiotic that works by binding to penicillin-binding proteins in Gram-positive and Gram-negative bacteria and inhibiting the transpeptidation process in bacteria.<sup>[3,10]</sup> Overuse of amoxicillin can lead to side effects, such as hypersensitivity reactions, nausea, vomiting, diarrhea, thrombocytopenia, dermatological disorders, and resistance.<sup>[11]</sup> Resistance to amoxicillin can also occur due to the destruction of the  $\beta$ -lactam ring by  $\beta$ -lactamase enzymes produced by *S. aureus* and *P. gingivalis*.<sup>[2,12]</sup> Therefore, amoxicillin is often combined with clavulanic acid, known as co-amoxiclav, to reduce resistance.<sup>[13]</sup> Clavulanic acid is a  $\beta$ -lactamase inhibitor that works by inactivating the pathogen's  $\beta$ -lactamase, thereby increasing the antibacterial activity of amoxicillin.<sup>[11]</sup> However, the use of co-amoxiclav can cause side effects, such as itching, redness around the mouth, diarrhea, nausea, and idiosyncratic drug-induced liver injury.<sup>[11,13]</sup> Alternative medicines using herbal plants have the potential to minimize these side effects.<sup>[14]</sup>

Arumanis mango (*Mangifera indica* L) is a herbal plant from India cultivated in various tropical and subtropical regions, including Indonesia.<sup>[15,16]</sup> Indonesians cultivate Arumanis mangos for their sweetness, freshness, and fragrance. The fruit contains Vitamins A, B, and C, which are beneficial for health.<sup>[17]</sup> In addition, the seeds, skin, roots, and leaves of *M. indica* L. have various properties in traditional medicine. *M. indica* L. leaves are generally discarded and considered waste, even though these leaves contain secondary metabolites, including phenolics (mangiferin, tannins, and flavonoids), alkaloids, saponins, terpenoids, glycosides, and steroids that have potential antibacterial, antifungal, antiviral, antiparasitic, antioxidant, anti-inflammatory, antitumor, anticancer, and analgesic effects.<sup>[18]</sup>

Previous studies showed that *M. indica* L. leaf extracts reduced the number of *Streptococcus mutans* and improved the antibacterial effect of clindamycin against *S. aureus*.<sup>[19,20]</sup> However, no research has investigated the potential of combining amoxicillin with *M. indica* L. leaf extracts in combating *S. mutans* and *P. gingivalis*. Thus, to address this research gap, this study aimed to determine the effectiveness of amoxicillin and ethanolic extract of Arumanis mango (*M. indica* L.) in inhibiting the growth and formation of *S. aureus* and *P. gingivalis* biofilms.

## Materials and Methods

This experimental *in vitro* study was performed at the Microbiology Center of Research and Education (MiCORE) Laboratory, Faculty of Dentistry, Trisakti University, Jakarta, Indonesia. An ethanolic extract of Arumanis mango (*M. indica* L.) leaves was prepared at the Research Institute for Spices and Medicinal Plants (BALITTRO)

in Bogor, West Java, Indonesia. The test solutions used were 10% dimethyl sulfoxide (DMSO) (negative control), co-amoxiclav (positive control), ethanolic extract of *M. indica* L. leaves with a concentration of 100%, and a combination of amoxicillin with extract concentrations of 3.125%, 6.25%, 12.5%, 25%, 50%, and 100%.

The sample size was calculated using Federer's formula, with n equals number of repetition, and t equals as total test group (8 groups comprised 6 treatment groups, 1 positive control, and 1 negative control). Based on this formula, each treatment group was conducted with four-time repetition for all assays.

### Ethanolic extract of *Mangifera indica* L. leaves

The ethanolic extract of *M. indica* L. leaves was prepared using the maceration method. The mango leaves were washed, dried, and mashed, and the simplicial was then soaked in 70% ethanol at a ratio of 1:5. The maceration process was performed for 2–3 h, and the macerate was then allowed to stand for 24 h and filtered. The extract was diluted with 10% DMSO solution to obtain extract concentrations of 50%, 25%, 12.5%, 6.25%, and 3.125%.

### Preparation of antibiotic solution

Amoxicillin and co-amoxiclav solution were prepared by crushing 500 mg of amoxicillin and 625 mg of co-amoxiclav until smooth, using a mortar and pestle. In total, 1.2 mg of amoxicillin and 1.5 mg of co-amoxiclav were each mixed with 6 ml of sterile distilled water until homogeneous to obtain 200 g/ $\mu$ l of amoxicillin and 250 g/ $\mu$ l of co-amoxiclav.

### Phytochemical tests

Qualitative phytochemical tests were performed at BALITTRO to identify secondary metabolites, such as alkaloids, saponins, tannins, phenolics, flavonoids, triterpenoids, steroids, and glycosides, in the ethanolic extract of *M. indica* L. leaves.

### Bacterial culture

*S. aureus* ATCC 25923 and *P. gingivalis* ATCC 33277 were obtained from the MiCORE Laboratory, Faculty of Dentistry, Trisakti University. *S. aureus* was cultured on brain heart infusion broth medium (Sigma Aldrich, St. Louis, Missouri). *P. gingivalis* was cultured on Tryptone Soya Broth (Sigma Aldrich, St. Louis, Missouri) medium enriched with hemin (5 mg/l), Vitamin K (10 mg/l), 0.5% yeast extract, and L-cystine (400 mg/l).<sup>[21]</sup> The medium was incubated in an anaerobic jar (Oxoid, Basingstoke, and Hampshire) at 37° for 24 h under anaerobic conditions. Following the incubation, the bacterial absorbance value was standardized into McFarland standard of 0.5, approximately  $1.5 \times 10^8$  colony forming unit (CFU)/mL (optical density [OD]<sub>600</sub>  $\pm$  0.132), before the following assays.

## Microdilution and total plate count

To evaluate the antibacterial properties of the extract combined with amoxicillin, an antibacterial test was performed on the plate count by the microdilution method. In total, 100 µL of cultured *S. aureus* ATCC 25923 and *P. gingivalis* ATCC 33277 were distributed into 96-well plates (Nest Biotech, Jiangsu, China). A test solution of 100 µl was added to each well and incubated at 37°C for 24 h under anaerobic conditions. After incubation, each mixture containing treated bacteria was diluted 10,000 times. Five microliters of diluted mixture were then spread on a petri dish containing sterile brain heart infusion agar media. The growth of bacterial colonies was calculated after incubation at 37°C for 24 h.

## Microtiter plate biofilm assay

The bacterial cultures were inserted into each well of the 96-well plates and then incubated at 37°C for 24 h under anaerobic conditions. The supernatant was discarded, leaving a thin layer on the surface of the well. The wells were rinsed using phosphate-buffered saline (PBS) (Biomatics, Ontario, Canada). Each test solution (200 µl) was added to each well and then incubated for 1, 3, and 24 h at 37°C. The wells were rinsed twice with PBS and fixated over burning spirit lamp. Crystal violet (Merck, Darmstadt, Germany) (200 µl) was then added to each well and allowed to stand for 15 min, followed by rinsing twice and standing for 15 min. In the past step, 200 µl of 96% ethanol was added. The OD was measured using a microplate reader (Safas, Monaco) with a wavelength of 490 nm.

## Statistical analysis

The research data were processed using the Statistical Package for the Social Sciences (SPSS) computer program, version 26 (IBM, Armonk, NY, USA). The Shapiro–Wilk method was used to test the normality of the data. If the data were normally distributed ( $P > 0.05$ ), a one-way analysis of variance test was conducted, followed by Turkey's honestly significant difference test (significance level of  $P < 0.05$ ) to verify the significance between the groups.

## Results

### Phytochemical screening

The results of the phytochemical screening qualitatively proved that the ethanolic extract of *M. indica* L. leaves contains alkaloids, saponins, tannins, phenolics, flavonoids, and steroids [Table 1].

### Antibacterial test using microdilution and plate count methods

The antibacterial test results showed that the ethanolic extract of *M. indica* L. leaves at 100% concentration and the combination of amoxicillin with extracts of various

**Table 1: The results of phytochemical screening of the ethanol extract of *Mangifera indica* L. leaves**

Secondary metabolites	Screening result
Alkaloids	+
Saponins	+
Tannins	+
Phenolics	+
Flavonoids	+
Triterpenoids	–
Steroids	+
Glycoside	–

concentrations exhibited antibacterial and antibiofilm effects against *S. aureus* and *P. gingivalis* [Figure 1]. In terms of inhibition of *S. aureus*, the results obtained by the combination of amoxicillin and the extract were not significantly different from those obtained using the positive control ( $P > 0.05$ ). The extract at a concentration of 12.5% and higher was effective against *S. aureus*, and the extract at a concentration of 3.125% and higher was effective against *P. gingivalis* [Figures 2 and 3].

### Antibiofilm test using the microtiter plate biofilm assay method

The results of the antibiofilm test showed that the addition of extracts to amoxicillin in inhibiting *S. aureus* biofilms had a significantly lower OD value compared to the positive control ( $P < 0.05$ ), starting at a concentration of 25% after 1 h of incubation [Figure 4] and at a concentration of 6.25% after 3 h of incubation [Figure 5]. After 24 h of incubation, following OD measurement, the group with the combination of extract concentration of 100% and amoxicillin proved not to be significantly different from OD of the positive control ( $P > 0.05$ ) [Figure 6]. In the *P. gingivalis* biofilm, the OD values of the addition of amoxicillin and the extract group, starting at a concentration of 25% after 1, 3, and 24 h of incubation, were smaller than the OD values of the positive control amoxiclav. The OD values of the treatment group were significantly different from those of the positive control amoxiclav [Figures 7-9].

## Discussion

As shown by our results, the ethanol extract of *M. indica* L. leaves contains secondary metabolites, including alkaloids, saponins, tannins, phenolics, flavonoids, and steroids. Different mechanisms of action of each compound account for the antibacterial and antibiofilm properties of the extract. Alkaloids inhibit the formation of peptidoglycan in bacterial cells. The alkaloids can disrupt the amino acid structure of bacterial DNA, leading to bacterial lysis.<sup>[22]</sup> Saponins damage the cell membrane and cell wall permeability in the diffusion process, resulting in the release of enzymes, amino acids, nutrients, and water, leading to cell destabilization and cell death.<sup>[23]</sup> Tannins form a complex with protein in the cell wall, namely,






Figure 1: The results of the inhibition test of *Staphylococcus aureus* and *Porphyromonas gingivalis* using the plate count method




Figure 2: Graph of the average total colony of *Staphylococcus aureus* (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant




Figure 3: Graph of the average total colony of *Porphyromonas gingivalis* (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant

proline, which can damage the cell wall.<sup>[22]</sup> The most abundant phenolic compound in *M. indica* L leaves is mangiferin. Mangiferin belongs to the xanthone C-glucosyl group, which can damage cell structure and cell membranes and inhibit bacterial protein synthesis.<sup>[24]</sup> Previous research reported that mangiferin compounds interfere with the mechanism of drug resistance; thus, restoring

bactericidal and bacteriostatic effects of nalidixic acid, ampicillin, tetracycline, and sulfamethoxazole/trimethoprim.<sup>[25]</sup> However, the exact mechanism of how mangiferin interferes with the mechanism of drug resistance is yet to be elucidated. Flavonoids, as antibacterial, play a role in disrupting the activity of cell wall formation by suppressing cytoplasm function, interrupting the nutrient exchange process, and thereby inhibiting the energy supply of bacteria.<sup>[26]</sup> In addition, flavonoids inhibit enzymes from producing quorum-sensing signals, thereby disrupting the communication process between cells during biofilm formation.<sup>[27]</sup> Steroids can cause leakage of lysosomes and membrane phospholipids that reduce the integrity of cell membranes and lead to cell lysis.<sup>[28]</sup>

Thus far, only a few studies have focused on antibacterial and antibiofilm properties of ethanolic extract of *M. indica* L. leaves.<sup>[19,20]</sup> The concentration of the extract used in this study ranged from 3.125% to 100%. As a positive control, co-amoxiclav, a combination of amoxicillin and clavulanic acid, was used. Amoxicillin, a  $\beta$ -lactam antibiotic, works by inhibiting the synthesis of bacterial cell walls.<sup>[11]</sup> The release of  $\beta$ -lactamase enzymes by *S. aureus* and *P. gingivalis* decrease the antibacterial effect of amoxicillin.<sup>[2,12]</sup> The addition of clavulanic acid binds to  $\beta$ -lactamase enzymes from bacteria, inhibiting the enzyme thereby unable to cleave  $\beta$ -lactam ring in amoxicillin, so the amoxicillin can still exhibit its antibacterial activity.<sup>[11]</sup>

In the microdilution and plate count tests, the addition of the ethanolic extract of *M. indica* L. leaves to amoxicillin inhibited the growth of *S. aureus* and *P. gingivalis*, and the combination of the ethanolic extract and amoxicillin was as effective as that of co-amoxiclav. In terms of antibacterial activity, a combination of ethanolic extract at a concentration of 12.5% and amoxicillin was as effective as co-amoxiclav against *S. aureus*, and a concentration as low as 3.125% was as effective as co-amoxiclav against *P. gingivalis*. Therefore, ethanolic extract of *M. indica* L. leaves may be a potential  $\beta$ -lactamase inhibitor equivalent to clavulanic acid. The results of this study are in line with the research of Hartanto *et al.*, who showed that adding




Figure 4: Graph of the average OD value of *Staphylococcus aureus* biofilm after 1 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant. OD: Optical density




Figure 5: Graph of the average OD value of *Staphylococcus aureus* biofilm after 3 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant. OD: Optical density




Figure 6: Graph of the average OD value of *Staphylococcus aureus* biofilm after 24 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant. OD: Optical density




Figure 7: Graph of the average OD value of *Porphyromonas gingivalis* biofilm after 1 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant. OD: Optical density




Figure 8: Graph of the average OD value of *Porphyromonas gingivalis* biofilm after 3 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant. OD: Optical density




Figure 9: Graph of the average OD value of *Porphyromonas gingivalis* biofilm after 24 h of incubation (\* $P < 0.05$ , \*\* $P < 0.01$ ). \*: Significant, \*\*: Very significant. OD: Optical density

methanolic extract of *M. indica* L. leaves to clindamycin has antibacterial effects against *S. aureus*, especially at a concentration of 100%.<sup>[19]</sup> The bioactive component of *M. indica* L. leaves, namely, mangiferin, is known to have a synergistic effect on tetracycline, ampicillin, nalidixic acid, and trimethoprim in inhibiting the growth of *S. aureus*.<sup>[25]</sup>

The biofilm formation phase begins with pellicle formation, which occurs in the first few seconds to the first mm from the initial contact, with the initial adhesion phase occurring 2–4 h later. After 24 h, the biofilm enters the maturation phase, becoming 1000–1500 times more resistant than

planktonic bacteria.<sup>[29]</sup> The incubation time used in the antibiofilm test in this study was adjusted to the stage of biofilm formation, namely 1, 3, and 24 h. This timing aimed to determine at which stage of biofilm formation amoxicillin and the ethanolic extract of *M. indica* L. leaves most effectively inhibited *S. aureus* and *P. gingivalis*. In the antibiofilm test, the OD from the combination of amoxicillin with ethanolic extract of *M. indica* L. leaves starting at a concentration of 25% after 1 h of incubation and 6.25% after 3 h of incubation had a significantly lower OD value compared to co-amoxiclav. This result proved that the ethanolic extract of *M. indica* L. leaves at a concentration of 25% after 1 h of incubation and 6.25% after 3 h of incubation was more effective at inhibiting *S. aureus* biofilms than co-amoxiclav. The OD values for the combination of amoxicillin and the extract at a concentration of 100% after 24 h of incubation were lower than those obtained for co-amoxiclav, but the difference was not statistically significant. Therefore, the combination of amoxicillin and an extract concentration of 100% has antibiofilm properties equivalent to those of co-amoxiclav after 24 h of incubation.

In terms of inhibiting *P. gingivalis* biofilm, the extract at a concentration of 25% and higher during the entire incubation period showed a more potent antibiofilm effect than that of co-amoxiclav. The combination of amoxicillin and the ethanolic extract of *M. indica* L. leaves inhibited the formation of *S. aureus* biofilms at a concentration of 6.25% after 3 h of incubation and *P. gingivalis* biofilms at a concentration of 25% after 1 h of incubation. This study supports the findings of previous research, which reported that ethanolic extract of *M. indica* L. leaves reduced *S. aureus* attachment and the number of *S. aureus* biofilms.<sup>[30]</sup>

This study has several limitations. First, the ethanolic extract used in this study was a crude extract. The use of a more refined extract, such as an extract exposed to extraction chromatography, would have resulted in an extract with fewer impurities. Second, only two of the many known oral pathogens were used in this study. Other oral pathogens can also be tested to expand the antibacterial activity of amoxicillin combined with *M. indica* L. leaf ethanolic extract. Further research is needed to determine the toxicity of this combination. Preclinical and clinical tests should also be conducted before the combination can be used as an alternative treatment for dentoalveolar abscesses and periodontitis.

## Conclusions

Ethanolic extract of *M. indica* L. leaves contains alkaloids, saponins, tannins, phenolics, flavonoids, and steroids that have the potential to inhibit the growth and formation of *S. aureus* and *P. gingivalis* biofilms *in vitro*. Within the limitations of this preliminary study, we conclude that the addition of ethanolic extract of *M. indica* L. leaves to amoxicillin could potentially increase the antibacterial and

antibiofilm properties of amoxicillin against *S. aureus* and *P. gingivalis*.

## Financial support and sponsorship

Nil.

## Conflicts of interest

There are no conflicts of interest.

## References

1. Ministry of Health of the Republic of Indonesia. Report on Result of National Basic Health Research 2018. Jakarta: Ministry of Health; 2018. p. 197-207.
2. Japoni A, Vasin A, Noushadi S, Kiany F, Japoni S, Alborzi A. Antibacterial susceptibility patterns of *Porphyromonas gingivalis* isolated from chronic periodontitis patients. *Med Oral Patol Oral Cir Bucal* 2011;16:e1031-5.
3. Shweta N, Prakash SK. Dental abscess: A microbiological review. *Dent Res J (Isfahan)* 2013;10:585-91.
4. Mehrotra N, Singh S. Periodontitis. Treasure Island (FL): StatPearls Publishing; 2011.
5. Archer NK, Mazaitis MJ, Costerton JW, Leid JG, Powers ME, Shirtliff ME. *Staphylococcus aureus* biofilms: Properties, regulation, and roles in human disease. *Virulence* 2011;2:445-59.
6. Huang R, Li M, Gregory RL. Bacterial interactions in dental biofilm. *Virulence* 2011;2:435-44.
7. Periasamy S, Joo HS, Duong AC, Bach TH, Tan VY, Chatterjee SS, et al. How *Staphylococcus aureus* biofilms develop their characteristic structure. *Proc Natl Acad Sci U S A* 2012;109:1281-6.
8. Taylor TA, Unakal CG. *Staphylococcus aureus*. Treasure Island (FL): StatPearls Publishing; 2021.
9. How KY, Song KP, Chan KG. *Porphyromonas gingivalis*: An overview of periodontopathic pathogen below the gum line. *Front Microbiol* 2016;7:53.
10. Ciancio SG, Mariotti AJ. Systemic Anti-infective therapy for periodontal diseases. In: Carranza F, Newman M, Takei H, Klokkevold P, editors. Newman and Carranza's Clinical Periodontology. 13<sup>th</sup> ed. Philadelphia: Elsevier; 2019. p. 555-7.
11. Akhavan B, Khanna N, Vijhani P. Amoxicillin. Treasure Island (FL): StatPearls Publishing; 2020.
12. Foster TJ, Geoghegan JA. *Staphylococcus aureus*. In: Molecular Medical Microbiology. Massachusetts: Elsevier; 2015. p. 655-74.
13. Samaranayake L. Essential Microbiology for Dentistry. 5<sup>th</sup> ed. Poland: Elsevier Ltd; 2018. p. 75, 158, 293, 297.
14. Joshua M, Takudzwa M. Antibacterial properties of *Mangifera indica* on *Staphylococcus aureus*. *Afr J Clin Exp Microbiol* 2013;14:62-74.
15. Kalita P. An overview on *Mangifera indica*: Importance and its various pharmacological action. *Pharma Tutoring* 2014;2:72-6.
16. Sivakumar D, Jiang Y, Yahia EM. Maintaining mango (*Mangifera indica* L.) fruit quality during the export chain. *Food Res Int* 2011;44:1254-63.
17. Chiumarelli M, Ferrari CC, Sarantopoulos CI, Hubinger MD. Fresh cut *Tommy Atkins* mango pre-treated with citric acid and coated with cassava (*Manihot esculenta* Crantz) starch or sodium alginate. *Innov Food Sci Emerg Technol* 2011;12:381-7.
18. Gu C, Yang M, Zhou Z, Khan A, Cao J, Cheng G. Purification and characterization of four benzophenone derivatives from *Mangifera indica* L. leaves and their antioxidant,

- immunosuppressive and  $\alpha$ -glucosidase inhibitory activities. *J Funct Foods* 2019;52:709-14.
19. Hartanto R, Tran V, Khang G, Pham T, Trinh T, Denhara C. Effect of the addition of arumanis mango leaf extract (*Mangifera indica* L.) on clindamycin antibiotics in inhibiting the growth of *Staphylococcus aureus*. *J Prima Med Sains* 2020;2:14-7.
  20. Kurniasih R. Effect of Concentrations of Young Mango Arumanis Ethanolic Extract (*Mangifera indica* L.) leaves on growth inhibition of *Streptococcus mutans in vitro*. Surakarta: Fakultas Kedokteran Gigi Universitas Muhammadiyah (Faculty of Dentistry, Muhammadiyah University); 2016.
  21. Yamanaka T, Furukawa T, Matsumoto-Mashimo C, Yamane K, Sugimori C, Nambu T, et al. Gene expression profile and pathogenicity of biofilm-forming *Prevotella intermedia* strain 17. *BMC Microbiol* 2009;9:11.
  22. Sylvana D, Amir M, Purnamasari CB, Iskandar A, Asfirizal V. Antibacterial activity of ethanol extract of beluntas leaves on *Streptococcus mutans*, *Porphyromonas gingivalis*, and *Enterococcus faecalis*. *Padjadjaran J Dent* 2021;33:191-8.
  23. Sebastian J, Widyarman AS. Roselle flower petals extract inhibits periodontal pathogenic biofilms. *J Dentomaxillofacial Sci* 2021;6:102-5.
  24. Tirado-Kulieva V, Atoche-Dioses S, Hernandez-Martinez E. Phenolic compounds of mango (*Mangifera indica*) by-products: Antioxidant and antimicrobial potential, use in disease prevention and food industry, methods of extraction, and microencapsulation. *Sci Agropecu* 2021;12:283-93.
  25. Mazlan NA, Azman S, Ghazali NF, Zarith P, Yusri S. Synergistic antibacterial activity of mangiferin with antibiotics against *Staphylococcus aureus*. *Drug Invent Today* 2019;12:14-7.
  26. Ma Y, Ding S, Fei Y, Liu G, Jang H, Fang J. Antimicrobial activity of anthocyanins and catechins against foodborne pathogens *Escherichia coli* and *Salmonella*. *Food Control* 2019;106:78-80.
  27. Federika AS, Rukmo M, Setyabudi S. Antibiofilm activity of flavonoid mangosteen pericarp extract against *Porphyromonas gingivalis* bacteria. *Conserv Dent J* 2020;10:27-30.
  28. Madduluri S, Rao KB, Sitaram B. *In vitro* evaluation of antibacterial activity of five indigenous plants extracts against five bacteria pathogens of humans. *Int J Pharm Pharm Sci* 2013;5:679-84.
  29. Bjarnsholt T. The role of bacterial biofilms in chronic infections. *APMIS Suppl* 2013;121:1-51.
  30. Manzur AG, Sm Junior V, Morais-Costa F, Mariano EG, Careli RT, da Silva LM, et al. Extract of *Mangifera indica* L. leaves may reduce biofilms of *Staphylococcus* spp. in stainless steel and teatcup rubbers. *Food Sci Technol Int* 2020;26:11-20.

Author Queries???

AQ6: Kindly check the significant \* in the image.