

BUKTI KORESPONDENSI
ARTIKEL JURNAL NASIONAL BEREPUTASI

Judul Artikel : Antiproliferative Activity of *Borassus flabellifer* L. Seed Coat Extracts on Oral Carcinoma Cells

Jurnal : Journal Syifa Sciences and Clinical Research (JSSCR), Volume 7 Nomor 2, 2025

Journal Homepage: <http://ejurnal.ung.ac.id/index.php/jsscr>

E-ISSN: 2656-9612 P-ISSN: 2656-8187

DOI : <https://doi.org/10.37311/jsscr.v7i2.31570>

Penulis : Pretty Trisfilha*, Janti Sudiono, Komariah, Celine Chelsea Hartono,
Namita Prayorina Nugroho, Raynita Rosalinda Angeline

NO	PERIHAL	TANGGAL
1	Bukti registrasi sebagai penulis	9 Mei 2025
2	Bukti konfirmasi submit artikel dan artikel yang disubmit	9 Mei 2025
3	Bukti konfirmasi review	1 Juni 2025
4	Bukti konfirmasi submit review, respon kepada reviewer dan artikel yang disubmit kembali	7 Juni 2025
5	Bukti konfirmasi artikel accepted	14 Juni 2025
6	Bukti konfirmasi artikel published online	20 Juni 2025

1. Bukti Registrasi Sebagai Penulis

9 Mei 2025

[JSSCR] Validate Your Account

External

Inbox x



Muhammad Taupik, S.Farm., M.Sc. <jurnal@ung.ac.id>

to me ▾

Fri, May 9, 10:40 AM

Mrs Pretty Trisfilha

You have created an account with Journal Syifa Sciences and Clinical Research, but before you can start using it, you need to validate your email account. To do this, simply follow the link below:

<https://ejurnal.ung.ac.id/index.php/jsscr/user/activateUser/pretty01/394AdY6n>

Thank you,
Muhammad Taupik, S.Farm., M.Sc.

Jambura Journal Syifa Science and Clinical Research

<http://ejurnal.ung.ac.id/index.php/jsscr>

[JSSCR] Journal Registration

External

Inbox x



Muhammad Taupik, S.Farm., M.Sc. <jurnal@ung.ac.id>

to me ▾

Fri, May 9, 10:40 AM

Mrs Pretty Trisfilha

You have now been registered as a user with Journal Syifa Sciences and Clinical Research. We have included your username and password in this email, which are needed for all work with this journal through its website. At any point, you can ask to be removed from the journal's list of users by contacting me.

Username:

Password:

Thank you,
Muhammad Taupik, S.Farm., M.Sc.

Jambura Journal Syifa Science and Clinical Research

<http://ejurnal.ung.ac.id/index.php/jsscr>

2. Bukti konfirmasi submit artikel dan
artikel yang disubmit
9 Mei 2025

[JSSCR] Submission Acknowledgement

External

Inbox x

Muhammad Taupik, S.Farm.,M.Sc. <jurnal@ung.ac.id>

Fri, May 9, 8:27 PM

to me ▼

Mrs Pretty Trisfilha:

Thank you for submitting the manuscript, "Evaluation of the Antiproliferative Activity of Borassus flabellifer L. Seed Coat Extracts in Oral Carcinoma Cells" to Journal Syifa Sciences and Clinical Research. With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal web site:

Manuscript URL:

<https://ejurnal.ung.ac.id/index.php/jsscr/author/submission/31570>

Username: pretty01

If you have any questions, please contact me. Thank you for considering this journal as a venue for your work.

Muhammad Taupik, S.Farm.,M.Sc.

Journal Syifa Sciences and Clinical Research

Jambura Journal Syifa Science and Clinical Research

<http://ejurnal.ung.ac.id/index.php/jsscr>



Home About User Home Categories Search Current Archives Announcements Publication Ethics

Home > User > Author > **Active Submissions**

Active Submissions

ACTIVE ARCHIVE

ID	MM-DD SUBMIT	SEC	AUTHORS	TITLE	STATUS
31570	05-09	ART	Trisfilha	EVALUATION OF THE ANTIPROLIFERATIVE ACTIVITY OF BORASSUS...	Awaiting assignment

Evaluation of the Antiproliferative Activity of *Borassus flabellifer* L. Seed Coat Extracts in Oral Carcinoma Cells

Pretty Trisfilha^{1*}, Janti Sudiono¹, Celine Chelsea Hartono², Namita Prayorina Nugroho², Raynita Rosalinda Angeline²

¹ Department of Oral & Maxillofacial Pathology, Faculty of Dentistry, Universitas Trisakti, Jl. Kyai Tapa No. 260 Grogol, Jakarta Barat, Indonesia

² Undergraduated Student, Faculty of Dentistry, Universitas Trisakti, Jl. Kyai Tapa No 260 Grogol, Jakarta Barat, Indonesia

* Corresponding Author. Email: pretty@trisakti.ac.id (Phone/Whatshapp : 089688677796)

ABSTRAK

Latar Belakang: *Borassus flabellifer* L. merupakan tanaman dari famili Arecaceae yang tersebar luas di negara-negara tropis Asia. Berbagai bagian tanaman ini, seperti buah, biji, daun, dan akar, telah digunakan dalam pengobatan tradisional karena mengandung senyawa bioaktif yang beragam. **Tujuan:** Penelitian ini bertujuan untuk mengevaluasi aktivitas antiproliferatif *in vitro* dari ekstrak etanol kulit biji *Borassus flabellifer* L. (EBSC) terhadap sel karsinoma sel skuamosa oral (HSC-3). **Metode:** Ekstrak disiapkan menggunakan pelarut etanol, kemudian dilakukan analisis fitokimia untuk mengidentifikasi kandungan senyawa bioaktif. Aktivitas antiproliferatif EBSC dievaluasi menggunakan uji Cell Counting Kit-8 (CCK-8) pada sel HSC-3 dengan variasi waktu dan konsentrasi. **Hasil:** Analisis fitokimia menunjukkan bahwa EBSC mengandung senyawa fenolik, flavonoid, terpenoid, tanin, dan alkaloid. Hasil uji CCK-8 menunjukkan bahwa EBSC memiliki efek antiproliferatif yang bergantung pada dosis terhadap sel HSC-3. Hambatan proliferasi sel yang signifikan tercatat pada konsentrasi 750 µg/mL, dengan efektivitas sebanding dengan doxorubicin (3 µM, $p < 0,001$) setelah perlakuan selama 24, 72, dan 120 jam. **Kesimpulan:** EBSC menunjukkan potensi sebagai agen antikanker terhadap sel karsinoma sel skuamosa oral, sehingga diperlukan penelitian lanjutan untuk mengkaji mekanisme kerja serta efektivitasnya secara *in vivo*.

Kata Kunci:

Aktivitas Antiproliferative, *Borassus flabellifer* L., Karsinoma Sel Skuamosa Oral

Diterima:
04-01-20xx

Disetujui:
07-02-20xx

Online:
07-02-20xx

ABSTRACT

Introduction: *Borassus flabellifer* L. is a plant in Arecaceae family, widely distributed and cultivated in tropical Asian countries. Different parts of *Borassus flabellifer* L., including the fruit, seeds, leaves, and roots, are used in traditional medicine due to their rich bioactive compound content. **Objective:** The present study aim to investigate the *in vitro* anti-proliferative activity of EBSC on HSC-3 cells. **Methods:** The extract was prepared using an ethanolic solvent, and its phytochemical composition was analyzed to identify potential bioactive compounds. The antiproliferative effects of the extract were evaluated using the Cell Counting Kit-8 (CCK-8) assay. **Results:** The phytochemical analysis revealed the presence of phenolic, flavonoid, terpenoid, tannin, and alkaloid in EBSC. In CCK-8 assay, The EBSC exhibited dose-dependent antiproliferative effects on HSC-3 cells. Significant inhibition of cell proliferation was observed at 750 µg/mL, comparable to doxorubicin (3 µM, $p < 0.001$), following 24, 72, and 120 hours of treatment. **Conclusion:** These results suggest that EBSC holds potential as a therapeutic agent for the treatment of oral squamous cell carcinoma, warranting further investigation into its underlying mechanisms and *in vivo*

efficacy.

Copyright © 20xx Jsscr. All rights reserved.

Keywords:

Antiproliferative Activity, *Borassus flabellifer* L., Oral Squamous Cell Carcinoma

Received:
20xx -01-04

Accepted:
20xx -02-07

Online:
20xx -02-07

1. Introduction

Oral squamous cell carcinoma (OSCC) represents a significant global health issues (1), In 2020, lip and oral cavity cancer was estimated to be the 16th most common type of cancer globally in terms of both incidence and mortality, accounting for a substantial proportion of oral cancers (2). Despite advancements in treatment modalities, OSCC continues to have high morbidity and mortality rates(3), underscoring the need for novel therapeutic agents. Traditional medicine, with its rich repository of bioactive compounds, has emerged as a promising avenue for discovering new cancer therapies (4)(5).

Borassus flabellifer L.(*B.flabellifer*.L), a member of the Arecaceae family, is widely distributed and cultivated in tropical Asian countries (6). Known commonly as the palmyra palm, it has been used extensively in traditional medicine. Various parts of the plant, including the fruit, seeds, leaves, and roots, are valued for their medicinal properties, attributed to the presence of diverse bioactive compounds. The potential therapeutic effects of plant extracts, particularly their anti-proliferative properties, are often linked to the presence of phytochemicals such as phenolics, flavonoids, terpenoids, tannins, and alkaloids. These compounds are known to exhibit antioxidant, anti-inflammatory, and anticancer activities (7)(8)(9)(10).

Previous studies have highlighted the medicinal potential of *B.flabellifer* L. extracts in various contexts. Research has demonstrated that *B.flabellifer* L. male flower extracts has antimicrobial, antioxidant, and anti-inflammatory properties (6). Another study found that the aqueous extract from the endosperm of germinated *B. flabellifer* L. seeds exhibited strong antiproliferative activity against MOLT-4 blood cancer cell lines (10). However, studies focusing specifically on the anti-proliferative effects of *Borassus flabellifer* L. seed coat extract on oral cancer (HSC-3) cell lines are limited. Therefore in the present study, we investigated the effect of *B.flabellifer*. L seed coat ethanol extract (EBSC) on the proliferation of human oral squamous carcinoma-3 (HSC-3) cells. Understanding the mechanisms through which this extract affects cancer cell viability and proliferation could provide valuable insights into its potential therapeutic applications.

2. Metode

This study is an in vitro laboratory experimental research investigating the effects of EBSC on HSC-3 cell lines. The research includes the preparation of EBSC at various concentrations and the assessment of its effects on HSC-3 cell proliferation. This study has been approved by Universitas Trisakti of the Dentistry Ethics Commission under ethical clearance number 173/KEP-UY/EA.10/VII/2023.

Sample Collection

B.flabellifer L. were collected from our local area Bogor, Indonesia. Seed coat was taken out from the *B.flabellifer* L. fruit and was cut into small pieces. These small pieces of *B.flabellifer* L. seed coat were dried for 4-5 days. The dried seed coat samples were stored at room temperature in order to be used for this study.

Plant Extraction

The extraction process was conducted using the maceration technique, where 62.2 g of dried *B. flabellifer* L. seed coat was immersed in 1,500 mL of 70% ethanol for three days. After the maceration period, the mixture was filtered and then concentrated using a rotary vacuum evaporator to produce a thick extract. This extract was then subjected to serial dilution to achieve five different concentrations: 93.75, 187.5, 375, 750, and 1500 µg/mL.

Phytochemical Test

Phytochemical tests were carried out at the Integrated Research Laboratory of Yarsi University to identify the active compounds contained in EBSC. Phytochemical tests were carried out qualitatively.

Cell Culture

The human oral squamous cell carcinoma (OSCC) cell line (HSC-3) was acquired from Integrated laboratory of Yarsi University (Jakarta, Indonesia). Cells were maintained in complete DMEM medium, 10% fetal bovine serum (FBS), 1% Penicillin-Streptomycin, and 1% Amphotericin B.

Cell Proliferation Assay

A cell counting kit-8 (CCK-8; Sigma-Aldrich) was used to characterize the antiproliferative activity of EBSC on HSC-3 cells. In brief, HSC-3 cells (5×10^3 cells) were seeded in triplicate in a 96-well plate. Following incubating with various concentrations (93.75; 187.5; 375; 750; 1500 µg/mL) of EBSC for 24, 72, and 120 h. 10 mL of CCK-8 solution was added to each well and incubated for 60 min. Cell counts were quantified using a microplate reader at an absorbance of 450 nm. Doxorubicin 3 µM was used as the positive control, while DMEM medium served as the negative control.

Statistical Analysis

The Shapiro-Wilk test was used to assess normality. Differences between the experimental groups were analyzed using a one-way analysis of variance (ANOVA) followed by a Post-Hoc test. A p-value of less than 0.05 ($p < 0.05$) was considered statistically significant.

3. Result and Discussion

Phytochemical Test

The qualitative phytochemical analysis revealed that the ethanol extract of *B.flabellifer* L. seed coat contains phenolics, flavonoids, terpenoids, tannins, and alkaloids (Table 1).

Table 1. Phytochemical test results

Secondary Metabolites	Methods	Results
Phenolic	FeCl ₃ 5% Reagent	+
	a. Concentrated HCl + Mg Reagent	+
Flavonoid	b. H ₂ SO ₄ 2N Reagent	+
	c. NaOH 10% Reagent	+
Steroid	Lieberman-Burchard Reagent	-
Terpenoid	Lieberman-Burchard Reagent	+
Saponin	HCl + H ₂ O Reagent	-
Tannin	FeCl ₃ 1% Reagent	+
	a. Hager Reagent	+
Alkaloid	b. Wagner Reagent	+
	c. Dragendorff Reagent	+

Cell Proliferation Assay

The possible effects of different concentrations of *B.flabellifer* L. seed coat extract (93.75; 187.5; 375; 750; 1500 µg/mL) on the proliferation of OSCC cell line HSC-3 were investigated further using CCK-8 assay. *B.flabellifer* L. seed coat extract treatment significantly suppressed the proliferation of HSC-3 cells in concentration-dependent manner. The best results are at a concentration 750 µg/mL. (Fig.1)




Figure 1.

Representative image of the CCK-8 assay after 24 h of treatment. HSC-3 cells were treated with different concentration of EBSC. CCK-8 assays were performed at 24, 72 and 120 h. Data are presented as the mean $\pm$ standard error of the mean of 3 repeats.* $P < 0.05$ vs positive control. EBSC : a ethanol extract of *B.flabellifer* L. seed coat

At present, the pathogenesis of oral squamous cell carcinoma (OSCC) is considered a complex multifactorial process that includes genetic predisposition, environmental exposures, and lifestyle factors such as tobacco and alcohol consumption(11). Among various cell types involved in OSCC

progression, oral keratinocytes and carcinoma-associated fibroblasts play pivotal roles in promoting tumor growth, invasion, and metastasis through dysregulated proliferation, resistance to apoptosis, and increased inflammatory signaling(12). Despite advancements in surgical techniques, radiotherapy, and chemotherapy, the prognosis of OSCC remains unsatisfactory due to high rates of recurrence and metastasis(13). Thus, there is a pressing need for alternative and adjunctive therapeutic agents derived from natural sources. *Borassus flabellifer* L., commonly known as palmyra palm, has been reported to possess multiple pharmacological activities, including antioxidant, anti-inflammatory, and anticancer properties (10)(8). Based on these properties, we investigated the antiproliferative effects of *B. flabellifer* L. seed coat extract on human OSCC HSC-3 cells.

Previous studies showed that *B.flabellifer* L. not only exert significant cytotoxicity towards several cancer cell lines(8), but also showed significant reduction in the percent increase of body weight, tumor volume and tumor weight of Ehrlich ascites carcinoma and Dalton ascites lymphoma in Swiss albino (7). In addition, Abirami et al demonstrated that hydrogel containing *B.flabellifer* L. male flower extract can exert significant inhibition towards blood cancer (MOLT-4) cell lines proliferation(10).

In the present study, CCK-8 assays results showed that EBSC significantly decreased the cell proliferation of HSC-3 cell lines in a dose-dependent manner, comparable to those mediated by the clinic drug Doxorubicin. The optimal inhibitory effect on cell proliferation was observed at a concentration of 750 µg/mL of EBSC. At this concentration, the inhibitory effect on HSC-3 cell proliferation was not significantly different from that of the positive control. However, at a higher concentration of 1500 µg/mL, the inhibitory effect on cell proliferation was lower compared to the 750 µg/mL concentration (Figure 1). This may be due to the fact that the concentration of 750 µg/mL has already reached the maximum effective dose of the extract. Similarly, Duddukuri et al discovered that 50% inhibition of *B.flabellifer*. L against HeLa cancer cells was found at this concentration(8).

Species of the genus *Borassus* have been known for their richness in phenolic compounds. Our study also showed that EBSCs are rich in flavonoids compounds such as phenolics, flavonoids, terpenoids, tannins, and alkaloids (Table 1). Those phenolic compounds are non-harmful to human health and have many health benefits; therefore, there is an increasing use of plants with high phenolic content in cancer therapy. The antiproliferative activity of the plant was correlated with the content of polyphenolic compounds. Flavonoid compounds, saponins, quinones, steroids, terpenoids, and tannins are believed to have antiproliferative activity by direct and indirect mechanism. Flavonoids are the most important natural phenolic compounds and are thought to be responsible for the antioxidant and the antiproliferative activities of natural products (14). Flavonoid compounds can inhibit the cell cycle, either in G1/S or G2/M, by inhibiting cyclin-dependent kinases (CDKs), which are key regulators of cell cycle development (15). Steroids and terpenoids can inhibit the primary mechanisms in cell proliferation and trigger apoptosis and autophagy of cancer cells (16)(17). Kim et al, found that geraniol, a monoterpene with structural and functional similarities, can effectively induce apoptosis and autophagy in tumor cells. Their study also showed that inhibition of AMPK promotes cell proliferation and enhances malignant behavior. These findings suggest that activating AMPK may enhance the anticancer effects of AKT pathway inhibition. Furthermore, geraniol was shown to inhibit the AKT signaling pathway, activate the AMPK pathway, and suppress the mTOR signaling pathway (18). Tannin compounds inhibit cancer proliferation indirectly by inducing apoptosis in Caspase 3 and Caspase 9-dependent pathways, down-regulating TGF beta, and causing mitochondrial membrane potential in KB cells. Tannins' pro-apoptotic action in KB cells is most likely due to ROS production and cell cycle arrest mechanisms (19).

4. Conclusion

Natural compounds derived from medicinal plants have been pivotal in cancer treatment. This study demonstrated that the ethanol extract of *B.flabellifer* L. seed coat contains flavonoid capable of inhibiting the proliferation of HSC-3 cell lines. The optimal inhibition of HSC-3 cell proliferation was observed at a concentration of 750 µg/mL. These results suggest that *B. flabellifer* L. seed coat extract holds potential as a therapeutic agent for the treatment of oral squamous cell carcinoma. However, further functional studies are required to determine the underlying mechanism by which the

proliferation of oral cancer cells is inhibited.

Acknowledgment

The authors would like to express their sincere gratitude to Universitas Trisakti for providing financial support for this research through its research grant program. We also extend our appreciation to the Faculty of Dentistry, Universitas Trisakti, for facilitating this study. Additionally, we would like to thank all individuals who contributed to this research, either directly or indirectly, for their valuable assistance and support.

References

1. Bai G, Chen B, Xiao X, Li Y, Liu X, Zhao D, et al. Geniposide inhibits cell proliferation and migration in human oral squamous carcinoma cells via AMPK and JNK signaling pathways. *Exp Ther Med*. 2022 Oct 4;24(6):706.
2. Bouvard V, Nethan ST, Singh D, Warnakulasuriya S, Mehrotra R, Chaturvedi AK, et al. IARC Perspective on Oral Cancer Prevention. *N Engl J Med*. 2022 Nov 24;387(21):1999–2005.
3. Xu N, Tian H, Fung C, Lin Y, Chen Y, Zhu G, et al. Inhibition of human oral squamous cell carcinoma proliferation and migration by prodrug-activating suicide gene therapies. *Exp Ther Med*. 2023 Jan 9;25(2):92.
4. Dehghani Nazhvani A, Sarafriz N, Askari F, Heidari F, Razmkhah M. Anti-Cancer Effects of Traditional Medicinal Herbs on Oral Squamous Cell Carcinoma. *Asian Pac J Cancer Prev*. 2020 Feb 1;21(2):479–84.
5. Chang KW, Wu CY, Renn TY, Chang TM, Wei AIC, Kung YY, et al. Aqueous Extract of Sparganii Rhizoma and Curcumae Rhizoma Induces Apoptosis and Inhibits Migration in Human Oral Squamous Cell Carcinoma. Ray S, editor. *J Food Biochem*. 2023 May 25;2023:1–13.
6. Tunit P, Thammarat P, Okonogi S, Chittasupho C. Hydrogel Containing Borassus flabellifer L. Male Flower Extract for Antioxidant, Antimicrobial, and Anti-Inflammatory Activity. *Gels*. 2022 Feb 17;8(2):126.
7. Babu G, . J. Assessment of Anticancer Activity of Ethanolic Extract of Borassus Flabellifer Linn. Leaf Against Ehrlich Ascites Carcinoma (EAC) and Daltons Ascites Lymphoma (DAL) in Swiss Albino Mice. *Int Res J Pharm*. 2019 Jan 31;10(1):113–8.
8. Duddukuri GR, Sastry YN, Chaitanya KK. Preliminary Studies on In Vitro Anti-Tumor Activity of Tender Seed Coat Extract of Borassus flabellifer L. On HELA Cell Line. *Int J Curr Res*. 2011;3(7):075–7.
9. Banu SM, Vigasini N, Surenderan S. In Vitro Antibacterial, Anticancer and Antidiabetic Potential of Freeze-dried Aqueous Borassus flabellifer L. Seed Powder Extract. *Indian J Pharm Sci*. 2022;84(3):586–92.
10. Abirami S, Vinushia AJ, Samrot AV, Pachiyappan S, Yogeswaran. In Vitro Antimicrobial, Antiproliferative Activity of Aqueous Extract From Endosperm of Germinated Palmyra Palm Seed (Borassus flabellifer L). *J Pharm Negat Results*. 2022;13(7).
11. Warnakulasuriya S. Global epidemiology of oral and oropharyngeal cancer. *Oral Oncol*. 2009 Apr;45(4–5):309–16.
12. Pannone G, Santoro A, Papagerakis S, Lo Muzio L, De Rosa G, Bufo P. The role of human papillomavirus in the pathogenesis of head & neck squamous cell carcinoma: an overview. *Infect Agent Cancer*. 2011 Dec;6(1):4.
13. Jemal A, Bray F, Center MM, Ferlay J, Ward E, Forman D. Global cancer statistics. *CA Cancer J Clin*. 2011 Mar;61(2):69–90.
14. Amrati FEZ, Bourhia M, Slighoua M, Ibnemoussa S, Bari A, Ullah R, et al. Phytochemical Study on Antioxidant and Antiproliferative Activities of Moroccan Caralluma europaea Extract and Its Bioactive Compound Classes. Mahajna JA, editor. *Evid Based Complement Alternat Med*. 2020 Jan;2020(1):8409718.

15. Isnaini I, Achmadiyah RD, Awaeh G, Khatimah H, Yasmina A. Antioxidant and Antiproliferative Activities of Methanol Extract from *Melaleuca cajuputi* subsp. *Cumingiana* [Turcz.] Fruit. *J Berk Ilm Sains Dan Terap Kim*. 2023 Feb 14;17(1):21.
16. Man S, Gao W, Zhang Y, Huang L, Liu C. Chemical study and medical application of saponins as anti-cancer agents. *Fitoterapia*. 2010 Oct;81(7):703–14.
17. Zhao YZ, Zhang YY, Han H, Fan RP, Hu Y, Zhong L, et al. Advances in the antitumor activities and mechanisms of action of steroidal saponins. *Chin J Nat Med*. 2018 Oct;16(10):732–48.
18. Hwa Kim S, Jung Park E, Lee CR, Chun JN, Hyuk Cho N, Gyu Kim I, et al. Geraniol induces cooperative interaction of apoptosis and autophagy to elicit cell death in PC-3 prostate cancer cells. *Int J Oncol*. 2011 Dec 23;40:1683–90.
19. Ahmad F, Surapaneni KM, Kamaraj B, Kamaraj B. Anti-proliferative activity and apoptotic induction of tannins extracted from *Quercus infectoria* on oral cancer KB cell lines. *J Appl Pharm Sci* [Internet]. 2024 [cited 2025 May 7]; Available from: https://japsonline.com/abstract.php?article_id=4244&sts=2

3. Bukti konfirmasi review 1 Juni 2025

Home > User > Author > Archive

Archive

ACTIVE ARCHIVE

ID	MM-DD SUBMIT	SEC	AUTHORS	TITLE	STATUS
31570	05-09	ART	Trisfilha, Sudiono, Komariah, Chelsea...	ANTIPROLIFERATIVE ACTIVITY OF BORASSUS FLABELLIFER L....	Vol 7, No 2 (2025): Volume 7 Edisi 2 2025

1 - 1 of 1 Items

START A NEW SUBMISSION

[CLICK HERE](#) to go to step one of the five-step submission process.

REFBACKS

ALL NEW PUBLISHED IGNORED

	DATE ADDED	HITS	URL	ARTICLE	TITLE	STATUS	ACTION
☐	2025-06-21	1	https://www.google.com/	Antiproliferative Activity of Borassus flabellifer L. Seed Coat Extracts on Oral Carcinoma Cells	—	New	EDIT I DELETE
☐	2025-06-21	1	https://ejurnal.ung.ac.id/	Antiproliferative Activity of Borassus flabellifer L. Seed Coat Extracts on Oral Carcinoma Cells	—	New	EDIT I DELETE
☐	2025-07-01	1	https://scholar.google.com/scholar?as_ylo=2025&q...	Antiproliferative Activity of Borassus flabellifer L. Seed Coat Extracts on Oral Carcinoma Cells	—	New	EDIT I DELETE

1 - 3 of 3 Items

Focus and Scope

Editorial Board

Peer Reviewer

Copyright Notice

Author Guideline

Online Submission

Author Fees

This journal has been published by
Department of Pharmacy, State University of
Gorontalo collaboration with IAI (Ikatan
Apoteker Indonesia)



Template :





Evaluation of the Antiproliferative Activity of Borassus flabellifer L. Seed Coat Extracts in Oral Carcinoma Cells



ABSTRACT

Introduction: *Borassus flabellifer* L. is a plant in *Arecaceae* family, widely distributed and cultivated in tropical Asian countries. Different parts of *Borassus flabellifer* L., including the fruit, seeds, leaves, and roots, are used in traditional medicine due to their rich bioactive compound content. **Objective:** The present study [aim](#) to investigate the in vitro antiproliferative activity of EBSC on HSC-3 cells. **Methods:** The extract was prepared using an ethanolic solvent, and its phytochemical composition was analyzed to identify potential bioactive compounds. The antiproliferative effects of the extract were evaluated using the Cell Counting Kit-8 (CCK-8) assay. **Results:** The phytochemical analysis revealed the presence of phenolic, flavonoid, terpenoid, tannin, and alkaloid in EBSC. In CCK-8 assay, [The](#) EBSC exhibited dose-dependent antiproliferative effects on HSC-3 cells. Significant inhibition of cell proliferation was observed at 750 μ g/mL, comparable to doxorubicin (3 μ M, $p < 0.001$), following 24, 72, and 120 hours of treatment. **Conclusion:** These results suggest that EBSC holds potential as a therapeutic agent for the treatment of oral squamous cell carcinoma, warranting further investigation into its underlying mechanisms and in vivo efficacy.

Copyright © 20xx *JSSCR*. All rights reserved.

Keywords:

Antiproliferative Activity, *Borassus flabellifer* L., Oral Squamous Cell Carcinoma

Received:
20xx-01-04

Accepted:
20xx-02-07

Online:
20xx-02-07

1. Introduction

Oral squamous cell carcinoma (OSCC) represents a significant global health [issues](#) (1). In 2020, lip and oral cavity cancer was estimated to be the 16th most common type of cancer globally in terms of both incidence and mortality, accounting for a substantial proportion of oral cancers (2). Despite advancements in treatment modalities, OSCC continues to have high morbidity and mortality rates(3), underscoring the need for novel therapeutic agents. Traditional medicine, with its rich repository of bioactive compounds, has emerged as a promising avenue for discovering new cancer therapies (4)(5).

Borassus flabellifer L. (*B. flabellifer* L.), a member of the *Arecaceae* family, is widely distributed and cultivated in tropical Asian countries (6). Known commonly as the palmyra palm, it has been used extensively in traditional medicine. Various parts of the plant, including the fruit, seeds, leaves, and roots, are valued for their medicinal properties, attributed to the presence of diverse bioactive compounds. The potential therapeutic effects of plant extracts, particularly their anti-proliferative properties, are often linked to the presence of phytochemicals such as phenolics, flavonoids, terpenoids, tannins, and alkaloids. These compounds are known to exhibit antioxidant, anti-inflammatory, and anticancer activities (7)(8)(9)(10).



A

Author

...



The title is descriptive and identifies the research focus well.

Suggestion:

To make it more concise and publishable, consider:
Antiproliferative Activity of
Borassus flabellifer L. Seed
Coat Extract on Oral
Carcinoma Cells

Reply

A

Author

...



Scientifically accurate, but some grammar and syntax issues.

Inconsistent tense use (aim change to aims; was change to were).

Suggestions:

Revise ending sentence for impact:

EBSC shows promising potential as a natural therapeutic candidate for OSCC treatment and warrants further in vivo studies.

Reply

A

Author

...



aims

Reply

A

Author

...



no need for The

Reply

Previous studies have highlighted the medicinal potential of *B. flabellifer* L. extracts in various contexts. Research has demonstrated that *B. flabellifer* L. male flower extracts has antimicrobial, antioxidant, and anti-inflammatory properties (6). Another study found that the aqueous extract from the endosperm of germinated *B. flabellifer* L. seeds exhibited strong antiproliferative activity against MOLT-4 blood cancer cell lines (10). However, studies focusing specifically on the anti-proliferative effects of *Borassus flabellifer* L. seed coat extract on oral cancer (HSC-3) cell lines are limited. Therefore in the present study, we investigated the effect of *B. flabellifer* L. seed coat ethanol extract (EBSC) on the proliferation of human oral squamous carcinoma-3 (HSC-3) cells. Understanding the mechanisms through which this extract affects cancer cell viability and proliferation could provide valuable insights into its potential therapeutic applications.

2. Metode

This study is an in vitro laboratory experimental research investigating the effects of EBSC on HSC-3 cell lines. The research includes the preparation of EBSC at various concentrations and the assessment of its effects on HSC-3 cell proliferation. This study has been approved by Universitas Trisakti of the Dentistry Ethics Commission under ethical clearance number 173/KEP-UY/EA.10/VII/2023.

Sample Collection

B. flabellifer L. were collected from our local area Bogor, Indonesia. Seed coat was taken out from the *B. flabellifer* L. fruit and was cut into small pieces. These small pieces of *B. flabellifer* L. seed coat were dried for 4-5 days. The dried seed coat samples were stored at room temperature in order to be used for this study.

Plant Extraction

The extraction process was conducted using the maceration technique, where 62.2 g of dried *B. flabellifer* L. seed coat was immersed in 1,500 mL of 70% ethanol for three days. After the maceration period, the mixture was filtered and then concentrated using a rotary vacuum evaporator to produce a thick extract. This extract was then subjected to serial dilution to achieve five different concentrations: 93.75, 187.5, 375, 750, and 1500 µg/mL.

Phytochemical Test

Phytochemical tests were carried out at the Integrated Research Laboratory of Yarsi University to identify the active compounds contained in EBSC. Phytochemical tests were carried out qualitatively.

Cell Culture

The human oral squamous cell carcinoma (OSCC) cell line (HSC-3) was acquired from Integrated laboratory of Yarsi University (Jakarta, Indonesia). Cells were maintained in complete DMEM medium, 10% fetal bovine serum (FBS), 1% Penicillin-Streptomycin, and 1% Amphotericin B.

Cell Proliferation Assay

A cell counting kit-8 (CCK-8; Sigma-Aldrich) was used to characterize the antiproliferative activity of EBSC on HSC-3 cells. In brief, HSC-3 cells (5x10³ cells) were seeded in triplicate in a 96-well plate. Following incubating with various concentrations



A Author ...
consistent with British or American English

Reply



A Author ...
Clearly structured with subheadings.
Ethical approval mentioned – excellent.
Some sub-sections contain formatting inconsistencies (extra indentation, misaligned bullets).
Suggestions:
Rephrase for clarity:
HSC-3 cells (5x10³) were seeded in triplicates...
Define abbreviations the first time they appear (e.g., DMEM, EBSC).

Reply



A Author ...
Add references in the end paragraph

Reply



A Author ...
incubation

Reply

(93.75; 187.5; 375; 750; 1500 $\mu\text{g/mL}$) of EBSC for 24, 72, and 120 h. 10 mL of CCK-8 solution was added to each well and incubated for 60 min. Cell counts were quantified using a microplate reader at an absorbance of 450 nm. Doxorubicin 3 μM was used as the positive control, while DMEM medium served as the negative control.

Statistical Analysis

The Shapiro-Wilk test was used to assess normality. Differences between the experimental groups were analyzed using a one-way analysis of variance (ANOVA) followed by a Post-Hoc test. A p-value of less than 0.05 ($p < 0.05$) was considered statistically significant.

3. Result and Discussion

Phytochemical Test

The qualitative phytochemical analysis revealed that the ethanol extract of *B. flabellifer* L. seed coat contains phenolics, flavonoids, terpenoids, tannins, and alkaloids (Table 1).

Table 1. Phytochemical test results		
Secondary Metabolites	Methods	Results
Phenolic	FeCl ₃ 5% Reagent	+
	a. Concentrated HCl + Mg Reagent	+
	b. H ₂ SO ₄ 2N Reagent	+
Flavonoid	c. NaOH 10% Reagent	+
	Lieberman-Burchard Reagent	-
	Lieberman-Burchard Reagent	+
Steroid	HCl + H ₂ O Reagent	-
Terpenoid	FeCl ₃ 1% Reagent	+
Saponin	a. Hager Reagent	+
Tannin	b. Wagner Reagent	+
	c. Dragendorff Reagent	+

Cell Proliferation Assay

The possible effects of different concentrations of *B. flabellifer* L. seed coat extract (93.75; 187.5; 375; 750; 1500 $\mu\text{g/mL}$) on the proliferation of OSCC cell line HSC-3 were investigated further using CCK-8 assay. *B. flabellifer* L. seed coat extract treatment significantly suppressed the proliferation of HSC-3 cells in concentration-dependent manner. The best results are at a concentration 750 $\mu\text{g/mL}$ (Fig.1)

Author ...

Informative, but formatting is not reader-friendly. Column headers should be aligned; consider using consistent chemical reagent naming.

Reply

Author ...

at a concentration of 750 $\mu\text{g/mL}$ (Figure 1).

Reply

there is an increasing use of plants with high phenolic content in cancer therapy. The antiproliferative activity of the plant was correlated with the content of polyphenolic compounds. Flavonoid compounds, saponins, quinones, steroids, terpenoids, and tannins are believed to have antiproliferative activity by direct and indirect mechanism. Flavonoids are the most important natural phenolic compounds and are thought to be responsible for the antioxidant and the antiproliferative activities of natural products (14). Flavonoid compounds can inhibit the cell cycle, either in G1/S or G2/M, by inhibiting cyclin-dependent kinases (CDKs), which are key regulators of cell cycle development (15). Steroids and terpenoids can inhibit the primary mechanisms in cell proliferation and trigger apoptosis and autophagy of cancer cells (16)(17). Kim et al, found that geraniol, a monoterpene with structural and functional similarities, can effectively induce apoptosis and autophagy in tumor cells. Their study also showed that inhibition of AMPK promotes cell proliferation and enhances malignant behavior. These findings suggest that activating AMPK may enhance the anticancer effects of AKT pathway inhibition. Furthermore, geraniol was shown to inhibit the AKT signaling pathway, activate the AMPK pathway, and suppress the mTOR signaling pathway (18). Tannin compounds inhibit cancer proliferation indirectly by inducing apoptosis in Caspase 3 and Caspase 9-dependent pathways, down-regulating TGF beta, and causing mitochondrial membrane potential in KB cells. Tannins' pro-apoptotic action in KB cells is most likely due to ROS production and cell cycle arrest mechanisms (19). |

4. Conclusion

Natural compounds derived from medicinal plants have been pivotal in cancer treatment. This study demonstrated that the ethanol extract of *B. flabellifer* L. seed coat contains flavonoid capable of inhibiting the proliferation of HSC-3 cell lines. The optimal inhibition of HSC-3 cell proliferation was observed at a concentration of 750 µg/mL. These results suggest that *B. flabellifer* L. seed coat extract holds potential as a therapeutic agent for the treatment of oral squamous cell carcinoma. However, further functional studies are required to determine the underlying mechanism by which the proliferation of oral cancer cells is inhibited. |

Acknowledgment

The authors would like to express their sincere gratitude to Universitas Trisakti for providing financial support for this research through its research grant program. We also extend our appreciation to the Faculty of Dentistry, Universitas Trisakti, for facilitating this study. Additionally, we would like to thank all individuals who contributed to this research, either directly or indirectly, for their valuable assistance and support.

References

1. Bai G, Chen B, Xiao X, Li Y, Liu X, Zhao D, et al. Geniposide inhibits cell proliferation and migration in human oral squamous carcinoma cells via AMPK and JNK signaling pathways. *Exp Ther Med*. 2022 Oct 4;24(6):706.
2. Bouvard V, Nethan ST, Singh D, Warnakulasuriya S, Mehrotra R, Chaturvedi AK, et al. IARC Perspective on Oral Cancer Prevention. *N Engl J Med*. 2022 Nov 24;387(21):1999-2005.

Author ...

It is recommended to include a brief limitations section at the end of the discussion. This will help readers understand the scope of the study and acknowledge aspects that were not fully addressed, such as molecular mechanisms or the lack of in vivo validation.

Reply

Author ...

The conclusion summarizes the findings well, but it would be stronger with the addition of a recommendation. For example, the authors may suggest future in vivo studies, deeper exploration of the molecular pathways, or potential development of extract-based formulations

Reply

Author ...

a flavonoid capable

Reply

Author ...

Relevant and up to date. Properly cited in Vancouver style. Suggestions: Ensure consistent format: some DOIs missing periods or spacing. Cross-check in-text numbering matches reference list. Example : V. Zingales, M. Fernández-Franzón, and M.-J. Ruiz, "Sterigmatocystin: Occurrence, toxicity and molecular mechanisms of action—A review," *Food and Chemical Toxicology*, vol. 146, p. 111802, 2020. [Online]. Available: <https://doi.org/10.1016/j.fct.2020.111802>




Figure 1. Representative image of the CCK-8 assay after 24 h of treatment. HSC-3 cells were treated with different concentration of EBSC. CCK-8 assays were performed at 24, 72 and 120 h. Data are presented as the mean $\pm$ standard error of the mean of 3 repeats.* $P < 0.05$ vs positive control. **EBSC:** a ethanol extract of *B. flabellifer* L. seed coat

At present, the pathogenesis of oral squamous cell carcinoma (OSCC) is considered a complex multifactorial process that includes genetic predisposition, environmental exposures, and lifestyle factors such as tobacco and alcohol consumption(11). Among various cell types involved in OSCC progression, oral keratinocytes and carcinoma-associated fibroblasts play pivotal roles in promoting tumor growth, invasion, and metastasis through dysregulated proliferation, resistance to apoptosis, and increased inflammatory signaling(12). Despite advancements in surgical techniques, radiotherapy, and chemotherapy, the prognosis of OSCC remains unsatisfactory due to high rates of recurrence and metastasis(13). Thus, there is a pressing need for alternative and adjunctive therapeutic agents derived from natural sources. *Borassus flabellifer* L., commonly known as palmyra palm, has been reported to possess multiple pharmacological activities, including antioxidant, anti-inflammatory, and anticancer properties (10)(8). Based on these properties, we investigated the antiproliferative effects of *B. flabellifer* L. seed coat extract on human OSCC HSC-3 cells.

Previous studies showed that *B. flabellifer* L. not only exert significant cytotoxicity towards several cancer cell lines(8), but also showed significant reduction in the percent increase of body weight, tumor volume and tumor weight of Ehrlich ascites carcinoma and Dalton ascites lymphoma in Swiss albino (7). In addition, Abirami et al demonstrated that hydrogel containing *B. flabellifer* L. male flower extract can exert significant inhibition towards blood cancer (MOLT-4) cell lines proliferation(10).

In the present study, CCK-8 assays results showed that EBSC significantly decreased the cell proliferation of HSC-3 cell lines in a dose-dependent manner, comparable to those mediated by the clinic drug Doxorubicin. The optimal inhibitory effect on cell proliferation was observed at a concentration of 750 µg/mL of EBSC. At this concentration, the inhibitory effect on HSC-3 cell proliferation was not significantly different from that of the positive control. However, at a higher concentration of 1500 µg/mL, the inhibitory effect on cell proliferation was lower compared to the 750 µg/mL concentration (Figure 1). This may be due to the fact that the concentration of 750 µg/mL has already reached the maximum effective dose of the extract. Similarly, Duddukuri et al discovered that 50% inhibition of *B. flabellifer* L. against HeLa cancer cells was found at this concentration(8).

Species of the genus *Borassus* have been known for their richness in phenolic compounds. Our study also showed that EBSCs are rich in flavonoids compounds such as phenolics, flavonoids, terpenoids, tannins, and alkaloids (Table 1). Those phenolic compounds are non-harmful to human health and have many health benefits; therefore,

A Author ...
Figure 1 lacks clarity in the graph section. Consider separating microscopy images and CCK-8 data into two figures, improving the resolution, and revising the caption to correct grammar and increase readability. Scale bars should be clearly shown in all panels. Ensure that each microscopy panel includes a visible scale bar. Also, replace 'Doxo' with the full name 'Doxorubicin' for clarity.

Reply

A Author ...
Rich in references and mechanistic insights. Some claims need clearer links to the study's findings. Suggestions:
Use a figure or chart for proliferation trends over 24, 72, 120 h.
Clarify this part:
At 1500 µg/mL, the inhibitory effect was lower...
Add explanation like: This may suggest cytotoxicity threshold or off-target effects at higher doses.
Typo:
"These results are at a concentration 750 µg/mL." → should be "at a concentration of 750 µg/mL"

Reply

4. Bukti konfirmasi submit review, respon
kepada reviewer dan artikel yang disubmit
kembali

7 Juni 2025



Home > User > Author > Submissions > #31570 > Review

#31570 Review

SUMMARY REVIEW EDITING

SUBMISSION

Authors	Pretty Trisfilha, Janti Sudiono, Komariah Komariah, Celine Chelsea Hartono, Namita Namita Prayorina Nugroho, Raynita Rosalinda Angeline
Title	Antiproliferative Activity of Borassus flabellifer L. Seed Coat Extracts on Oral Carcinoma Cells
Section	Articles
Editor	Muhammad Taupik

PEER REVIEW

ROUND 1

Review Version	31570-89404-3-RV.docx 2025-06-01
Initiated	2025-06-01
Last modified	2025-06-02
Uploaded file	None

EDITOR DECISION

Decision	Accept Submission 2025-06-14
Notify Editor	Editor/Author Email Record 2025-06-18
Editor Version	31570-91423-1-ED.docx 2025-06-01 31570-91423-2-ED.docx 2025-06-01 31570-91423-3-ED.docx 2025-06-02 31570-91423-4-ED.docx 2025-06-18
Author Version	31570-91851-1-ED.docx 2025-06-07 DELETE
Upload Author Version	Choose File no file selected Upload

Antiproliferative Activity of *Borassus flabellifer* L. Seed Coat Extracts on Oral Carcinoma Cells

Pretty Trisfilha^{1*}, Janti Sudiono², Komariah Komariah³, Celine Chelsea Hartono⁴, Namita Prayorina Nugroho⁵, Raynita Rosalinda Angeline⁶

^{1,2} Department of Oral & Maxillofacial Pathology, Faculty of Dentistry, Universitas Trisakti, Jl. Kyai Tapa No. 260 Grogol, Jakarta Barat, Indonesia

³ Department of Oral Biology sub-division of Histology, Faculty of Dentistry, Universitas Trisakti, Jl. Kyai Tapa No. 260 Grogol, Jakarta Barat, Indonesia

^{4,5,6} Faculty of Dentistry, Universitas Trisakti, Jl. Kyai Tapa No 260 Grogol, Jakarta Barat, Indonesia

* Corresponding Author. Email: pretty@trisakti.ac.id

ABSTRACT

Borassus flabellifer L. is a plant in Arecaceae family, widely distributed and cultivated in tropical Asian countries. Different parts of *Borassus flabellifer* L., including the fruit, seeds, leaves, and roots, are used in traditional medicine due to their rich bioactive compound content. The present study aims to investigate the in vitro anti-proliferative activity of EBSC on HSC-3 cells. The extract were prepared using an ethanolic solvent, and its phytochemical composition were analyzed to identify potential bioactive compounds. The antiproliferative effects of the extract were evaluated using the Cell Counting Kit-8 (CCK-8) assay. The phytochemical analysis revealed the presence of phenolic, flavonoid, terpenoid, tannin, and alkaloid in EBSC. In CCK-8 assay, EBSC exhibited dose-dependent antiproliferative effects on HSC-3 cells. Significant inhibition of cell proliferation was observed at 750 µg/mL, comparable to doxorubicin (3 µM, $p < 0.001$), following 24, 72, and 120 hours of treatment. EBSC shows promising potential as a natural therapeutic candidate for OSCC treatment and warrants further in vivo studies.

Copyright © 2025 Jsscr. All rights reserved.

Keywords: *Borassus flabellifer* L.; Seed Coat Extract; Antiproliferative; OSCC; Natural Anticancer Agent; CCK-8 Assay

Received:

2025 -05-09

Accepted:

2025 -06-14

Online:

2025 -06-20

1. Introduction

Oral squamous cell carcinoma (OSCC) represents a significant global health issue [1], In 2020, lip and oral cavity cancer was estimated to be the 16th most common type of cancer globally in terms of both incidence and mortality, accounting for a substantial proportion of oral cancers [2]. Despite

advancements in treatment modalities, OSCC continues to have high morbidity and mortality rates [3], underscoring the need for novel therapeutic agents. Traditional medicine, with its rich repository of bioactive compounds, has emerged as a promising avenue for discovering new cancer therapies [4],[5].

Borassus flabellifer L. a member of the Arecaceae family, is widely distributed and cultivated in tropical Asian countries [6]. Known commonly as the palmyra palm, it has been used extensively in traditional medicine. Various parts of the plant, including the fruit, seeds, leaves, and roots, are valued for their medicinal properties, attributed to the presence of diverse bioactive compounds. The potential therapeutic effects of plant extracts, particularly their anti-proliferative properties, are often linked to the presence of phytochemicals such as phenolics, flavonoids, terpenoids, tannins, and alkaloids. These compounds are known to exhibit antioxidant, anti-inflammatory, and anticancer activities [7],[8],[9],[10].

Previous studies have highlighted the medicinal potential of *B.flabellifer* L. extracts in various contexts. Research has demonstrated that *B.flabellifer* L. male flower extracts possess antimicrobial, antioxidant, and anti-inflammatory properties [6]. Another study reported that the aqueous extract from the endosperm of germinated *B. flabellifer* L. seeds exhibited strong antiproliferative activity against MOLT-4 blood cancer cell lines [10]. However, studies focusing specifically on the antiproliferative effects of *Borassus flabellifer* L. seed coat extract on oral cancer cell lines remain limited. Therefore in the present study, we investigated the effect of *B.flabellifer*. L seed coat ethanol extract (EBSC) on the proliferation of human oral squamous carcinoma-3 (HSC-3) cells. Understanding the mechanisms by which this extract influences cancer cell viability and proliferation could provide valuable insights into its potential therapeutic applications.

2. Metode

This study is an in vitro laboratory experimental research investigating the effects of EBSC on HSC-3 cell lines. The research includes the preparation of EBSC at various concentrations and the assessment of its effects on HSC-3 cell proliferation. This study has been approved by Universitas Trisakti of the Dentistry Ethics Commission under ethical clearance number 173/KEP-UY/EA.10/VII/2023.

Sample Collection

B.flabellifer L. were collected from our local area Bogor, Indonesia. Seed coat was taken out from the *B.flabellifer* L. fruit and was cut into small pieces. These small pieces of *B.flabellifer* L. seed coat were dried for 4-5 days. The dried seed coat samples were stored at room temperature in order to be used for this study [8].

Plant Extraction

The extraction process was conducted using the maceration technique, where 62.2 g of dried *B. flabellifer* L. seed coat was immersed in 1,500 mL of 70% ethanol for three days. After the maceration period, the mixture was filtered and then concentrated using a rotary vacuum evaporator to produce a thick extract. This extract was then subjected to serial dilution to achieve five different concentrations: 93.75, 187.5, 375, 750, and 1500 µg/mL [7].

Phytochemical Test

Phytochemical tests were carried out at the Integrated Research Laboratory of Yarsi University to identify the active compounds contained in EBSC. Phytochemical tests were carried out qualitatively [6].

Cell Culture

The human oral squamous cell carcinoma (OSCC) cell line (HSC-3) was acquired from Integrated laboratory of Yarsi University (Jakarta, Indonesia). Cells were maintained in complete Dulbecco's Modified Eagle Medium (DMEM), 10% fetal bovine serum (FBS), 1% Penicillin-Streptomycin, and 1% Amphotericin B [5].

Cell Proliferation Assay

A cell counting kit-8 (CCK-8; Sigma-Aldrich) was used to characterize the antiproliferative activity of EBSC on HSC-3 cells. HSC-3 cells (5×10^3 cells) were seeded in triplicates in a 96-well plate. Following incubation with various concentrations (93.75; 187.5; 375; 750; 1500 µg/mL) of EBSC for 24, 72, and 120 h. 10 mL of CCK-8 solution was added to each well and incubated for 60 min. Cell counts were quantified using a microplate reader at an absorbance of 450 nm. Doxorubicin 3 µM was used as the positive control, while DMEM served as the negative control [5].

Statistical Analysis

The Shapiro-Wilk test was used to assess normality. Differences between the experimental groups were analyzed using a one-way analysis of variance (ANOVA) followed by a Post-Hoc test. A p-value of less than 0.05 ($p < 0.05$) was considered statistically significant [11].

3. Result and Discussion

Phytochemical Test

The qualitative phytochemical analysis revealed that the ethanol extract of *B.flabellifer* L. seed coat contains phenolics, flavonoids, terpenoids, tannins, and alkaloids (Table 1).

Table 1. Phytochemical Screening Results of EBSC

Secondary Metabolites	Detected
Phenolic	+
Flavonoid	+
Steroid	-

Terpenoid	+
Saponin	-
Tannin	+
Alkaloid	+

Note: (+) indicates presence; (-) indicates absence of the compound

Phytochemical screening of the ethanol extract of *B. flabellifer* L. seed coat (EBSC) revealed the presence of phenolics, flavonoids, terpenoids, tannins, and alkaloids, which are widely recognized for their anticancer and cytotoxic properties. Flavonoids and phenolics can act as antioxidants and modulators of cell signaling pathways, while alkaloids have been reported to induce apoptosis and interfere with DNA replication in cancer cells [7],[11],[13]. Terpenoids and tannins may also contribute to cytotoxicity by disrupting mitochondrial function and enhancing oxidative stress [8], [10]. The absence of steroids and saponins indicates that the observed antiproliferative effects are likely driven by the dominant phenolic and alkaloid constituents, in line with previous studies on plant-based anticancer agents [12],[14].

Cell Proliferation Assay

The possible effects of different concentrations of *B. flabellifer* L. seed coat extract (93.75; 187.5; 375; 750; 1500 $\mu\text{g/mL}$) on the proliferation of OSCC cell line HSC-3 were investigated further using CCK-8 assay. *B. flabellifer* L. seed coat extract treatment significantly suppressed the proliferation of HSC-3 cells in concentration-dependent manner. The best results are at a concentration of 750 $\mu\text{g/mL}$ (Figure 1 & 2).





Figure 1. Proliferation of HSC-3 cells measured using a CCK-8 assay. CCK-8 assays were performed at 24, 72 and 120 h. Data are presented as the mean $\pm$ standard error of the mean of 3 repeats.* $P < 0.05$ vs positive control.






Figure 2. Representative image of the CCK-8 assay after 24 h of treatment. HSC-3 cells were treated with different concentration of EBSC, with 750 µg/mL shows the most significant anti proliferative-effect. EBSC : a ethanol extract of *B.flabellifer* L. seed coat

At present, the pathogenesis of oral squamous cell carcinoma (OSCC) is considered a complex multifactorial process that includes genetic predisposition, environmental exposures, and lifestyle factors such as tobacco and alcohol consumption [12]. Among various cell types involved in OSCC progression, oral keratinocytes and carcinoma-associated fibroblasts play pivotal roles in promoting tumor growth, invasion, and metastasis through dysregulated proliferation, resistance to apoptosis, and increased inflammatory signaling [13]. Despite advancements in surgical techniques, radiotherapy, and chemotherapy, the prognosis of OSCC remains unsatisfactory due to high rates of recurrence and metastasis [14]. Thus, there is a pressing need for alternative and adjunctive therapeutic agents derived from natural sources. *Borassus flabellifer* L., commonly known as palmyra palm, has been reported to possess multiple pharmacological activities, including antioxidant, anti-inflammatory, and anticancer properties [10][8]. Based on these properties, we investigated the antiproliferative effects of *B. flabellifer* L. seed coat extract on human OSCC HSC-3 cells.

Previous studies showed that *B.flabellifer* L. not only exert significant cytotoxicity towards several cancer cell lines [8], but also showed significant reduction in the percent increase of body weight, tumor volume and tumor weight of Ehrlich ascites carcinoma and Dalton ascites lymphoma in Swiss albino [7]. In addition, Abirami et al demonstrated that hydrogel containing *B.flabellifer* L. male flower extract can exert significant inhibition towards blood cancer (MOLT-4) cell lines proliferation [10].

In the present study, CCK-8 assays results showed that EBSC significantly decreased the proliferation of HSC-3 cell lines in a dose-dependent manner, with effects comparable to those induced by the clinical drug Doxorubicin. The optimal inhibitory effect on cell proliferation was observed at a concentration of 750 µg/mL of EBSC. At this concentration, the inhibition of HSC-3 cell proliferation was not significantly different from that of the positive control. Interestingly, at a higher concentration of 1500 µg/mL, the inhibitory effect was reduced compared to the 750 µg/mL concentration (**Figure 1**). This may suggest that 750 µg/mL represents the maximal effective dose of the extract, and that higher concentrations may exceed a cytotoxicity threshold or trigger off-target effects that compromise efficacy. A similar trend was reported by Duddukuri et al., who discovered that 50% inhibition of *B.flabellifer* L against HeLa cancer cells occurred at this concentration [8].

Species of the genus *Borassus* have been known for their richness in phenolic compounds. Our study also showed that EBSCs are rich in flavonoids compounds such as phenolics, flavonoids, terpenoids, tannins, and alkaloids (**Table 1**). Those phenolic compounds are non-harmful to human health and have many health benefits; therefore, there is an increasing use of plants with high phenolic content in cancer therapy. The antiproliferative activity of the plant was correlated with the content of polyphenolic compounds. Flavonoid compounds, saponins, quinones, steroids, terpenoids, and tannins are believed to have antiproliferative activity by direct and indirect mechanism. Flavonoids are the most important natural phenolic compounds and are thought to be responsible for the antioxidant and the antiproliferative activities of natural products [15].

Flavonoid compounds can inhibit the cell cycle, either in G1/S or G2/M, by inhibiting cyclin-dependent kinases (CDKs), which are key regulators of cell cycle development [11]. Steroids and terpenoids can inhibit the primary mechanisms in cell proliferation and trigger apoptosis and autophagy of cancer cells [16][17]. Kim et al, found that geraniol, a monoterpene with structural and functional similarities, can effectively induce apoptosis and autophagy in tumor cells. Their study also

showed that inhibition of AMPK promotes cell proliferation and enhances malignant behavior. These findings suggest that activating AMPK may enhance the anticancer effects of AKT pathway inhibition. Furthermore, geraniol was shown to inhibit the AKT signaling pathway, activate the AMPK pathway, and suppress the mTOR signaling pathway [18]. Tannin compounds inhibit cancer proliferation indirectly by inducing apoptosis in Caspase 3 and Caspase 9-dependent pathways, down-regulating TGF beta, and causing mitochondrial membrane potential in KB cells. Tannins' pro-apoptotic action in KB cells is most likely due to ROS production and cell cycle arrest mechanisms [19].

Despite these promising findings, this study has several limitations. The molecular mechanisms underlying the antiproliferative effects of EBSC were not examined in detail, and the study was confined to in vitro testing using a single OSCC cell line without in vivo validation. Therefore, further studies involving mechanistic exploration, animal models, and isolation of specific active compounds are necessary to better elucidate the therapeutic potential of *B. flabellifer* L. seed coat extract as a candidate for OSCC treatment.

4. Conclusion

Natural compounds derived from medicinal plants have been pivotal in cancer treatment. This study demonstrated that the ethanol extract of *B. flabellifer* L. seed coat contains a flavonoid capable of inhibiting the proliferation of HSC-3 cell lines. The optimal inhibition of HSC-3 cell proliferation was observed at a concentration of 750 µg/mL. These results suggest that *B. flabellifer* L. seed coat extract holds potential as a therapeutic agent for the treatment of oral squamous cell carcinoma. The authors may suggest further functional and mechanistic studies to elucidate the molecular pathways involved in its anticancer activity, as well as in vivo studies to validate its efficacy and safety. It is therefore recommended that future research focuses on the isolation and characterization of the active constituents, alongside preclinical evaluations, to support the development of plant-derived adjunctive therapies for oral cancer.

Acknowledgment

The authors would like to express their sincere gratitude to Universitas Trisakti for providing financial support for this research through its research grant program. We also extend our appreciation to the Faculty of Dentistry, Universitas Trisakti, for facilitating this study. Additionally, we would like to thank all individuals who contributed to this research, either directly or indirectly, for their valuable assistance and support.

Conflicts of Interest:

The authors declare no conflicts of interest.

References

- [1] G. Bai et al., "Geniposide inhibits cell proliferation and migration in human oral squamous carcinoma cells via AMPK and JNK signaling pathways," *Exp. Ther. Med.*, vol. 24, no. 6, p. 706, Oct. 2022. [Online]. Available: <https://doi.org/10.3892/etm.2022.11642>
- [2] V. Bouvard et al., "IARC Perspective on Oral Cancer Prevention," *N. Engl. J. Med.*, vol. 387, no. 21, pp. 1999–2005, Nov. 2022. [Online]. Available: <https://doi.org/10.1056/NEJMSr2210097>
- [3] N. Xu et al., "Inhibition of human oral squamous cell carcinoma proliferation and migration by prodrug-activating suicide gene therapies," *Exp. Ther. Med.*, vol. 25, no. 2, p. 92, Jan. 2023. [Online]. Available: <https://doi.org/10.3892/etm.2023.11790>
- [4] A. D. Nazhvani, N. Sarafraz, F. Askari, F. Heidari, and M. Razmkhah, "Anti-Cancer Effects of Traditional Medicinal Herbs on Oral Squamous Cell Carcinoma," *Asian Pac. J. Cancer Prev.*, vol. 21, no. 2, pp. 479–484, Feb. 2020. [Online]. Available: <https://doi.org/10.31557/APJCP.2020.21.2.479>

- [5] K.-W. Chang et al., "Aqueous Extract of Sparganii Rhizoma and Curcumae Rhizoma Induces Apoptosis and Inhibits Migration in Human Oral Squamous Cell Carcinoma," *J. Food Biochem.*, vol. 2023, May 2023. [Online]. Available: <https://doi.org/10.1155/2023/6564124>
- [6] P. Tunit, P. Thammarat, S. Okonogi, and C. Chittasupho, "Hydrogel Containing Borassus flabellifer L. Male Flower Extract for Antioxidant, Antimicrobial, and Anti-Inflammatory Activity," *Gels*, vol. 8, no. 2, p. 126, Feb. 2022. [Online]. Available: <https://doi.org/10.3390/gels8020126>
- [7] G. Babu and J. J., "Assessment of Anticancer Activity of Ethanolic Extract of Borassus flabellifer Linn. Leaf Against Ehrlich Ascites Carcinoma and Daltons Ascites Lymphoma in Swiss Albino Mice," *Int. Res. J. Pharm.*, vol. 10, no. 1, pp. 113–118, Jan. 2019. [Online]. Available: <https://doi.org/10.7897/2230-8407.100120>
- [8] G. R. Duddukuri, Y. N. Sastry, and K. K. Chaitanya, "Preliminary Studies on In Vitro Anti-Tumor Activity of Tender Seed Coat Extract of Borassus flabellifer L. On HeLa Cell Line," *Int. J. Curr. Res.*, vol. 3, no. 7, pp. 075–077, 2011. [Online]. Available: <http://www.journalcra.com/?q=node/271>
- [9] S. M. Banu, N. Vigasini, and S. Surenderan, "In Vitro Antibacterial, Anticancer and Antidiabetic Potential of Freeze-dried Aqueous Borassus flabellifer L. Seed Powder Extract," *Indian J. Pharm. Sci.*, vol. 84, no. 3, pp. 586–592, 2022. [Online]. Available: <https://doi.org/10.36468/pharmaceutical-sciences.964>
- [10] S. Abirami et al., "In Vitro Antimicrobial, Antiproliferative Activity of Aqueous Extract From Endosperm of Germinated Palmyra Palm Seed (Borassus flabellifer L)," *J. Pharm. Negat. Results*, vol. 13, no. 7, 2022. [Online]. Available: <https://doi.org/10.47750/pnr.2022.13.S07.054>
- [11] I. Isnaini et al., "Antioxidant and Antiproliferative Activities of Methanol Extract from Melaleuca cajuputi subsp. cumingiana [Turcz.] Fruit," *J. Berk. Ilm. Sains Dan Terap. Kim.*, vol. 17, no. 1, p. 21, Feb. 2023. [Online]. Available: <https://doi.org/10.20527/jstk.v17i1.13055>
- [12] S. Warnakulasuriya, "Global epidemiology of oral and oropharyngeal cancer," *Oral Oncol.*, vol. 45, no. 4–5, pp. 309–316, Apr. 2009. [Online]. Available: <https://doi.org/10.1016/j.oraloncology.2008.06.002>
- [13] G. Pannone et al., "The role of human papillomavirus in the pathogenesis of head & neck squamous cell carcinoma: an overview," *Infect. Agent. Cancer*, vol. 6, no. 1, p. 4, Dec. 2011. [Online]. Available: <https://doi.org/10.1186/1750-9378-6-4>
- [14] A. Jemal et al., "Global cancer statistics," *CA. Cancer J. Clin.*, vol. 61, no. 2, pp. 69–90, Mar. 2011. [Online]. Available: <https://doi.org/10.3322/caac.20107>
- [15] F. E.-Z. Amrati et al., "Phytochemical Study on Antioxidant and Antiproliferative Activities of Moroccan Caralluma europaea Extract and Its Bioactive Compound Classes," *Evid. Based Complement. Alternat. Med.*, vol. 2020, p. 8409718, Jan. 2020. [Online]. Available: <https://doi.org/10.1155/2020/8409718>
- [16] S. Man, W. Gao, Y. Zhang, L. Huang, and C. Liu, "Chemical study and medical application of saponins as anti-cancer agents," *Fitoterapia*, vol. 81, no. 7, pp. 703–714, Oct. 2010. [Online]. Available: <https://doi.org/10.1016/j.fitote.2010.06.004>

- [17] Y.-Z. Zhao et al., "Advances in the antitumor activities and mechanisms of action of steroidal saponins," *Chin. J. Nat. Med.*, vol. 16, no. 10, pp. 732–748, Oct. 2018. [Online]. Available: [https://doi.org/10.1016/S1875-5364\(18\)30113-4](https://doi.org/10.1016/S1875-5364(18)30113-4)
- [18] S. H. Kim et al., "Geraniol induces cooperative interaction of apoptosis and autophagy to elicit cell death in PC-3 prostate cancer cells," *Int. J. Oncol.*, vol. 40, pp. 1683–1690, Dec. 2011. [Online]. Available: <https://doi.org/10.3892/ijo.2011.1318>
- [19] F. Ahmad, K. M. Surapaneni, B. Kamaraj, and B. Kamaraj, "Anti-proliferative activity and apoptotic induction of tannins extracted from *Quercus infectoria* on oral cancer KB cell lines," *J. Appl. Pharm. Sci.*, 2024. [Online]. Available: <https://doi.org/10.7324/JAPS.2024.174009>

5. Bukti konfirmasi artikel accepted
14 Juni 2025

Antiproliferative Activity of *Borassus flabellifer* L. Seed Coat Extracts on Oral Carcinoma Cells

Pretty Trisfilha^{1*}, Janti Sudiono², Komariah Komariah³, Celine Chelsea Hartono⁴,
Namita Prayorina Nugroho⁵, Raynita Rosalinda Angeline⁶

^{1,2} Department of Oral & Maxillofacial Pathology, Faculty of Dentistry, Universitas Trisakti,
Jl. Kyai Tapa No. 260 Grogol, Jakarta Barat, Indonesia

³ Department of Oral Biology sub-division of Histology, Faculty of Dentistry, Universitas Trisakti,
Jl. Kyai Tapa No. 260 Grogol, Jakarta Barat, Indonesia

^{4,5,6} Faculty of Dentistry, Universitas Trisakti, Jl. Kyai Tapa No 260 Grogol, Jakarta Barat, Indonesia

* Corresponding Author. Email: pretty@trisakti.ac.id

ABSTRACT

Borassus flabellifer L. is a plant in Arecaceae family, widely distributed and cultivated in tropical Asian countries. Different parts of *Borassus flabellifer* L., including the fruit, seeds, leaves, and roots, are used in traditional medicine due to their rich bioactive compound content. The present study aims to investigate the in vitro anti-proliferative activity of EBSC on HSC-3 cells. The extract were prepared using an ethanolic solvent, and its phytochemical composition were analyzed to identify potential bioactive compounds. The antiproliferative effects of the extract were evaluated using the Cell Counting Kit-8 (CCK-8) assay. The phytochemical analysis revealed the presence of phenolic, flavonoid, terpenoid, tannin, and alkaloid in EBSC. In CCK-8 assay, EBSC exhibited dose-dependent antiproliferative effects on HSC-3 cells. Significant inhibition of cell proliferation was observed at 750 µg/mL, comparable to doxorubicin (3 µM, $p < 0.001$), following 24, 72, and 120 hours of treatment. EBSC shows promising potential as a natural therapeutic candidate for OSCC treatment and warrants further in vivo studies.

Copyright © 2025 Jsscr. All rights reserved.

Keywords: *Borassus flabellifer* L.; Seed Coat Extract; Antiproliferative; OSCC; Natural Anticancer Agent; CCK-8 Assay

Received:
2025-05-09

Accepted:
2025-06-14

Online:
2025-06-20

6. Bukti konfirmasi artikel publish online 20 Juni 2025



Journal Syifa Sciences and Clinical Research (JSSCR)

Volume 7 Nomor 2, 2025

Journal Homepage: <http://ejurnalung.ac.id/index.php/jsscr>, E-ISSN: 2656-9612 P-ISSN: 2656-8187

DOI : <https://doi.org/10.37311/jsscr.v7i2.31570>



Antiproliferative Activity of *Borassus flabellifer* L. Seed Coat Extracts on Oral Carcinoma Cells

Pretty Trisfilha^{1*}, Janti Sudiono², Komariah Komariah³, Celine Chelsea Hartono⁴,
Namita Prayorina Nugroho⁵, Raynita Rosalinda Angeline⁶

^{1,2} Department of Oral & Maxillofacial Pathology, Faculty of Dentistry, Universitas Trisakti,
Jl. Kyai Tapa No. 260 Grogol, Jakarta Barat, Indonesia

³ Department of Oral Biology sub-division of Histology, Faculty of Dentistry, Universitas Trisakti,
Jl. Kyai Tapa No. 260 Grogol, Jakarta Barat, Indonesia

^{4,5,6} Faculty of Dentistry, Universitas Trisakti, Jl. Kyai Tapa No 260 Grogol, Jakarta Barat, Indonesia

* Corresponding Author. Email: pretty@trisakti.ac.id

ABSTRACT

Borassus flabellifer L. is a plant in Arecaceae family, widely distributed and cultivated in tropical Asian countries. Different parts of *Borassus flabellifer* L., including the fruit, seeds, leaves, and roots, are used in traditional medicine due to their rich bioactive compound content. The present study aims to investigate the in vitro anti-proliferative activity of EBSC on HSC-3 cells. The extract were prepared using an ethanolic solvent, and its phytochemical composition were analyzed to identify potential bioactive compounds. The antiproliferative effects of the extract were evaluated using the Cell Counting Kit-8 (CCK-8) assay. The phytochemical analysis revealed the presence of phenolic, flavonoid, terpenoid, tannin, and alkaloid in EBSC. In CCK-8 assay, EBSC exhibited dose-dependent antiproliferative effects on HSC-3 cells. Significant inhibition of cell proliferation was observed at 750 µg/mL, comparable to doxorubicin (3 µM, $p < 0.001$), following 24, 72, and 120 hours of treatment. EBSC shows promising potential as a natural therapeutic candidate for OSCC treatment and warrants further in vivo studies.

Copyright © 2025 Jsscr. All rights reserved.

Keywords: *Borassus flabellifer* L.; Seed Coat Extract; Antiproliferative; OSCC; Natural Anticancer Agent; CCK-8 Assay

Received:
2025-05-09

Accepted:
2025-06-14

Online:
2025-06-20