

VOLUME 6, No.2, 2025

**SJD**

e-ISSN 2746-0592

# Sriwijaya

## Journal of Dentistry

PUBLISHED BY

**DENTISTRY STUDY PROGRAM,  
FACULTY OF MEDICINE UNIVERSITAS SRIWIJAYA**

Jl. Dr. Moehammad Ali Komp. RSMH Palembang 30126, Indonesia  
Telp. 0711-373438/ 087795591525, Email : [sjd@fk.unsri.ac.id](mailto:sjd@fk.unsri.ac.id)

# Editorial History

## Sriwijaya Journal of Dentistry

Published by Dentistry Study Program, Faculty of  
Medicine Universitas Sriwijaya

Online ISSN 2746-0592

[Home](#) / [Editorial History](#)

This section lists past contributors.

**Editor in Chief :** Siti Rusdiana Puspa Dewi, SCOPUS ID: 57209249462,  
Department of Biomedical, Faculty of Medicine, Universitas Sriwijaya,  
Palembang, Indonesia

**Managing Editor:** Martha Mozartha, SCOPUS ID: 57210231521,  
Department of Biomaterial, Faculty of Medicine, Universitas Sriwijaya,  
Palembang, Indonesia

### **Editors:**

- Atia Nurul Sidiqa, SCOPUS ID : 57204915403, Departemen Dental Material Fakultas Kedokteran Gigi Universitas Jenderal Achmad Yani, Jawab Barat, Indonesia
- Shanty Chairani , SCOPUS ID: 58092402300, Department of Oral Biology, Faculty of Medicine, Universitas Sriwijaya, Palembang, Indonesia
- Muhammad Novo Perwira Lubis, Departemen Radiologi Kedokteran Gigi Fakultas Kedokteran Gigi Universitas Padjajaran, Bandung, Indonesia
- Maya Hudiyati, SCOPUS ID: 57210232091, Department of Biomaterial, Faculty of Medicine, Universitas Sriwijaya, Indonesia
- Sri Wahyuningsih Rais, SCOPUS ID: 57210919485, Department of

Prosthodontia, Faculty of Medicine, Universitas Sriwijaya,  
Palembang, Indonesia

- Shinta Amini Prativi, SCOPUS ID: 58092402200 Department of Radiology, Faculty of Medicine, Universitas Sriwijaya, Palembang, Indonesia
- Haslina Rani, SCOPUS ID: 56694491400 Department of Family Oral Health, Faculty of Dentistry, Universiti Kebangsaan Malaysia, Malaysia

**Sriwijaya**  
Journal of Dentistry

**Published by Dentistry Study Program, Faculty of  
Medicine Universitas Sriwijaya**



**Sriwijaya Journal of Dentistry** : Publikasi Ilmiah Fakultas Kedokteran Universitas Sriwijaya by <https://sjd-fk.ejournal.unsri.ac.id> is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License.

# Reviewers

## Sriwijaya Journal of Dentistry

Published by Dentistry Study Program, Faculty of  
Medicine Universitas Sriwijaya

Online ISSN 2746-0592

[Home](#) / [Reviewers](#)

Pudji Handayani [[Google Scholar](#), [SINTA ID](#)], Department of Oral  
Medicine, Dentistry Study Program, Faculty of Medicine, Universitas  
Sriwijaya, Indonesia

Danica Anastasia [[Google Scholar](#), [SINTA ID](#)], Department of  
Conservative Dentistry, Dentistry Study Program, Faculty of Medicine,  
Universitas Sriwijaya, Indonesia

Arief Cahyanto [[Google Scholar](#), [Scopus ID](#)], Department of Dental  
Material, Faculty of Dentistry, Universitas Padjajaran, Indonesia

Erik Idrus [[Google Scholar](#), [Scopus ID](#)], Department of Oral Biology,  
Faculty of Dentistry, Universitas Indonesia, Indonesia

Mellisa Adiatman [[Google Scholar](#), [Scopus ID](#)], Department of Public  
Health, Faculty of Dentistry, Universitas Indonesia, Indonesia

Dini Asrianti [[Google Scholar](#), [Scopus ID](#)], Department of Conservative  
Dentistry, Faculty of Dentistry, Universitas Indonesia, Indonesia

Fuad Husain Akbar [[Google Scholar](#), [Scopus ID](#)], Department of Dental  
Public Health, Faculty of Dentistry, Hasanuddin University, Indonesia

Ida Bagus Narmada [[Google Scholar](#), [Scopus ID](#)], Department of  
Orthodontics, Faculty of Dentistry, Airlangga University, Indonesia

Rahmi Amtha [Google Scholar, Scopus ID], Department of Oral Medicine, Faculty of Dentistry, Universitas Trisakti, Indonesia

Fatimah Maria Tadjoedin [Google Scholar, Scopus ID], Department of Periodontics, Faculty of Dentistry, Universitas Indonesia, Indonesia

Masita Mandasari [Google Scholar, Scopus ID], Department of Oral Medicine, Faculty of Dentistry, Universitas Indonesia, Indonesia

Citra Kusumasari [Google Scholar, Scopus ID], Department of Conservative Dentistry, Faculty of Dentistry, Universitas Indonesia, Indonesia

Yosi Kusuma Eriwati [Google Scholar, Scopus ID], Department of Dental Material, Faculty of Dentistry, Universitas Indonesia, Indonesia

Rasmi Rikmasari [Google Scholar, Scopus ID], Department of Prosthodontics, Faculty of Dentistry, Universitas Padjajaran, Indonesia

Kartika Andari Wulan [Google Scholar, SINTA ID], Department of Prosthodontics, Faculty of Dentistry, Universitas Brawijaya, Indonesia

Muhammad Ruslin [Google Scholar, SINTA ID], Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, Universitas Hasanuddin, Indonesia

Wiwiek Poedjiastoeti [Google Scholar, SINTA ID], Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, Universitas Trisakti, Indonesia

Risti Saptarini Primarti [Google Scholar, Scopus ID], Department of Pediatric Dentistry, Faculty of Dentistry, Universitas Padjajaran, Indonesia

Ratna Indriyanti [Google Scholar, SINTA ID], Department of Pediatric Dentistry, Faculty of Dentistry, Universitas Padjajaran, Indonesia

Erma Sofiani [Google Scholar, Scopus ID], Department of Conservative Dentistry, Faculty of Dentistry, Universitas Muhammadiyah Yogyakarta, Indonesia

Rosalina Tjandrawinata [Google Scholar, SINTA ID, Scopus ID], Department of Dental Material, Faculty of Dentistry, Universitas Trisakti, Indonesia

Vinna Kurniawati Sugiawan [Google Scholar, Scopus ID], Department of Oral Biology, Faculty of Dentistry, Universitas Kristen Maranatha, Indonesia

Gita Gayatri [[Google Scholar](#), [SINTA ID](#)], Department of Orthodontics, Faculty of Dentistry, Universitas Padjajaran, Indonesia

Ziske Maritska, [[Google Scholar](#), [SINTA ID](#)], Department of Biology Medicine, Faculty of Medicine, Universitas Sriwijaya, Indonesia

Yosaphat Bayu Rosanto, [[Google Scholar](#), [SINTA ID](#)], Department of Oral And Maxillofacial Surgery, Faculty of Dentistry, Universitas Gadjah Mada, Indonesia

Yolanda [[Google Scholar](#), [SINTA ID](#) ], Department of Conservative Dentistry, Faculty of Dentistry, Universitas Padjadjaran, Indonesia

Gema Gempita, [[Google Scholar](#), [SINTA ID](#)], Department of Dental Material Science and Technology, Faculty of Dentistry, Universitas Padjadjaran, Indonesia

Sunnati, [[Google Scholar](#), [SINTA ID](#)], Department of Periodontics, Faculty of Dentistry, Universitas Syiah Kuala, Indonesia

**Sriwijaya**  
Journal of Dentistry

**Published by Dentistry Study Program, Faculty of  
Medicine Universitas Sriwijaya**



**Sriwijaya Journal of Dentistry** : Publikasi Ilmiah Fakultas Kedokteran Universitas Sriwijaya by <https://sjd-fk.ejournal.unsri.ac.id> is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License.

# Sriwijaya Journal of Dentistry

Published by Dentistry Study Program, Faculty  
of Medicine Universitas Sriwijaya

Online ISSN 2746-0592

[🔍 Current Issue](#)   [📄 About the Journal](#)



## Vol. 6 No. 2 (2025): Sriwijaya Journal of Dentistry

📅 2025-12-24

## Articles

### Effect of Brushing with Whitening Toothpaste on External Discoloration of Resin Modified Glass Ionomer Cement

— Yosi Kusuma Eriwati, Rachel Puja Chrisanti, Ette Soraya Tadjuddin



ID :

Page: 50-59

## The Effectiveness of Pagar Alam Green Tea Leaf Extract (*Camellia sinensis* L.) as an Irrigation Material on Smear Layer Removal

— Rhamasuci Putri Jasa, Merryca Belinda, Listia Eka Merdekawati, Danica Anastasia

[DOI](#)[?](#)

ID :

Page: 60-68

## Efficacy of Probiotics, Cetylpyridinium Chloride, and Enzymes Toothpaste Formulations Against Periodontal Pathogens in Fixed Orthodontic Patients: A Review

— Michelle, Joko Kusnoto, Harryanto Wijaya

[DOI](#)[?](#)

ID :

Page: 69-78

## Clinical Management of Conventional Gingivectomy in an Adult Patients with Gingival Enlargement

— Manta Fany, Maria Fransisca Utha, Depi Praharani, Melok Aris Wahyukundari, Neira Najatus Sakinah, Peni Pujiastuti, Yuliana Mahdiyah Da'at Arina, Desi Sandra Sari

[DOI](#)[?](#)

ID :

Page: 79-89

## Comparison of Cytotoxicity Between Calcium Silicate and Epoxy Resin-Based Sealer in Root Canal Treatment (Systematic Review)

— Hana Zakiyah Suarsyaf, Rinda Yulianti, Billy Sujatmiko

[DOI](#)[?](#)

ID :

Page: 90-107

[View All Issues »](#)



## Efficacy of Probiotics, Cetylpyridinium Chloride, and Enzymes-Based Toothpaste Formulations Against Periodontal Pathogens in Fixed Orthodontic Patients: A Review

Michelle<sup>1</sup>, Joko Kusnoto<sup>2\*</sup>, Harryanto Wijaya<sup>2</sup>

<sup>1</sup>Master Program of Dental Sciences, Faculty of Dentistry, Universitas Trisakti, Indonesia

<sup>2</sup>Department of Orthodontics, Faculty of Dentistry, Universitas Trisakti, Indonesia

\*Correspondence author email: [joko.k@trisakti.ac.id](mailto:joko.k@trisakti.ac.id)

### Abstract

**Introduction:** Fixed orthodontic appliances complicate oral hygiene and promote biofilm accumulation, increasing the risk of periodontal disease. Toothpaste formulations with antimicrobial properties such as *Lactobacillus paracasei* probiotics, cetylpyridinium chloride (CPC), and amyloglucosidase–glucose oxidase enzymes are commonly used to manage oral biofilm and control pathogenic bacteria. However, the extent of their effectiveness against periodontal pathogens in orthodontic patients remains unclear. **Purpose:** This review aims to map and summarize the current evidence on the efficacy of various toothpaste formulations in reducing periodontal pathogens among patients undergoing fixed orthodontic treatment. **Methods:** A comprehensive search of electronic databases (e.g., PubMed, Scopus, Google Scholar) was conducted to identify relevant studies published from 2015 to 2025. Data were extracted and synthesized narratively, focusing on types of formulations, study outcomes, and microbial targets. **Results:** Probiotic, enzymatic, and CPC-containing toothpastes showed variable but complementary effects against key pathogens, including *Porphyromonas gingivalis* and *Aggregatibacter actinomycetemcomitans*. Probiotics modulate microbial balance, enzymes disrupt biofilm structure, and CPC provides chemical antimicrobial action. Combining these mechanisms appears promising for enhanced biofilm control. **Conclusion:** Toothpastes containing *L. paracasei* probiotics, CPC, and amyloglucosidase–glucose oxidase enzymes offer complementary ways to control periodontal pathogens in orthodontic patients. Combining these mechanisms may enhance biofilm control and microbial balance, though further research is needed to confirm their long-term benefits.

**Keywords:** antimicrobial effect; fixed orthodontic appliances; oral pathogens; toothpaste formulations

### Introduction

Fixed orthodontic appliances are widely used to correct malocclusions and improve dental aesthetics.<sup>1</sup> However, these appliances create numerous retention sites that hinder effective plaque removal, making it difficult for patients to maintain optimal oral hygiene. As a result, orthodontic treatment often leads to increased biofilm accumulation and inflammation of the surrounding periodontal tissues, particularly the gingiva.<sup>2-4</sup>

Wearing fixed orthodontic appliances can lead to more biofilm accumulation, which shifts the oral microbiota toward more harmful bacteria. Periodontal pathogens like *Porphyromonas gingivalis* and *Aggregatibacter actinomycetemcomitans* may increase,



raising the risk of gingivitis and even periodontitis if not controlled. Therefore, managing bacterial biofilm is crucial to maintain good periodontal health during orthodontic treatment.<sup>5,6</sup>

Toothpaste is one of the most accessible and commonly used oral hygiene tools. Various formulations, including those containing probiotics, enzymes, and cetylpyridinium chloride have been developed to enhance antimicrobial activity and improve periodontal outcomes.<sup>7-9</sup> Despite their widespread use, evidence on their ability to reduce periodontal pathogens in orthodontic patients remains limited, as existing studies are scattered, heterogeneous, and often restricted to in vitro experiments or clinical trials. This gap underscores the need for a comprehensive scoping review to evaluate and map existing research on this topic. This review explores the range of toothpaste formulations studied for their antimicrobial effects on periodontal pathogens in patients with fixed orthodontic appliances.

### Methods

A comprehensive search of electronic databases such as PubMed, Scopus, Google Scholar was conducted to identify relevant studies published from 2015 to 2025 by using the Boolean Search such as ["*Porphyromonas gingivalis*" OR "P. gingivalis" OR "*Aggregatibacter actinomycetemcomitans*" OR "A. actinomycetemcomitans" OR "periodontal pathogens" OR "periodontopathogenic bacteria"] AND ["Probiotics toothpaste" OR "probiotic toothpaste" OR "*Lactobacillus paracasei* toothpaste" OR "L. paracasei toothpaste"] AND ["Cetylpyridinium chloride" OR "cetylpyridinium" OR "CPC"] AND ["Enzyme-based toothpaste" OR "enzymatic toothpaste" OR "amyloglucosidase" OR "glucose oxidase" OR "enzymes-based formulations"] AND ["fixed orthodontic appliances" OR "fixed orthodontic" OR "fixed appliance"] AND ["bacterial reduction" OR "biofilm reduction" OR "antimicrobial effect" OR "antibacterial effect" OR "dental plaque"].

This review included studies involving patients with fixed orthodontic appliances or laboratory models that mimic orthodontic biofilm conditions. Eligible studies examined probiotics such as *Lactobacillus paracasei*, cetylpyridinium chloride (CPC), or enzyme-based toothpastes containing amyloglucosidase and glucose oxidase, and reported outcomes related to *Porphyromonas gingivalis* or *Aggregatibacter actinomycetemcomitans*. Randomized trials, clinical studies, observational studies, and in vitro or ex vivo experiments were all considered if the full text was available in English. Studies were excluded if they involved removable appliances or clear aligners, evaluated unrelated oral hygiene products, or did not measure



microbiological outcomes for the targeted pathogens. Reviews, case reports, conference abstracts, animal studies, and articles lacking adequate data were also excluded. Data were extracted and synthesized narratively, focusing on types of formulations, study outcomes, and microbial targets. Two reviewers performed the screening process.

### Results

The role of probiotics was first proposed by Metchnikoff in the 20th century. According to the World Health Organization (WHO), probiotics are live microorganisms that, when consumed, can provide benefits to the host.<sup>10</sup> Probiotics can be given in various forms such as tablets, mouthwash, toothpaste, and drinks. Several studies have proven the beneficial effects of probiotics for humans, such as preventing caries, gingivitis, halitosis, and improving the host's immune response.<sup>10,11</sup> *Lactobacillus sp.* and *Bifidobacterium sp.* are colonies of good probiotic bacteria that have an effect on the oral cavity and are often used in orthodontic treatment. Probiotics interact with other bacteria in the human body. Their mechanism of action includes protecting epithelial tissue, increasing mucosal adhesion, inhibiting adhesion of pathogenic bacteria, and modulating the immune system through IgA secretion. Probiotics produce organic acids such as acetic and lactic acids, which can inhibit the effects of Gram-negative bacteria.<sup>12,13</sup> As a type of Lactic Acid Bacteria (LAB), *L. paracasei* has the ability to secrete extracellular polysaccharides (EPS) which can provide protection and increase the adhesion of probiotic bacteria in the oral cavity.<sup>14</sup> *L. paracasei* functions also can competitive exclusion of pathogens, modulation of the host immune response, and secretion of bacteriocins that inhibit periodontal pathogens.<sup>15</sup>

Cethylpyridinium chloride (CPC) is one of the active ingredients added to toothpaste. This ingredient has a broad-spectrum antimicrobial effect that works by destroying the cell integrity of bacteria. The Food and Drug Administration (FDA) has deemed this ingredient safe and effective for the treatment of plaque. Using CPC toothpaste can also reduce the amount of plaque in the oral cavity.<sup>9,16</sup> The active CPC ingredient in toothpaste is a quaternary ammonium compound. The CPC content dispersed in the oral cavity after use is maintained through surfactant chains and cationic charges absorbed in the oral cavity.<sup>17,18</sup> The CPC molecule has two molecular groups including hydrophilic and hydrophobic. The positively charged, or hydrophilic, group increases the binding affinity for negatively charged pathogenic bacteria. The hydrophobic group interacts with the bacterial cell surface and causes the bacteria to



integrate into the cytoplasmic membrane. These two interactions can disrupt cell membrane integration, disrupt cellular metabolism, increase cytoplasmic leakage, and inhibit growth, which can lead to bacterial cell death. The presence of these two groups also reduces microbial adhesion to oral surfaces.<sup>18</sup>

Toothpaste containing enzymes such as amyloglucosidase and glucose oxidase has an antimicrobial effect by reducing plaque and bacterial accumulation, gingival inflammation, and can prevent the formation of white spot lesions.<sup>8</sup> The amyloglucosidase enzyme can inhibit bacterial growth by converting D-glucose to D-glucono-1,5-lactone. Presence of D-glucono-1,5-lactone reduces the nutritional needs of bacteria in the oral cavity. The mechanism of the glucose oxidase enzyme is to activate the salivary immune system, the lactoperoxidase (LPO) system, by producing hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), which can react with the catalase enzyme to produce oxygen, thus reducing anaerobic bacteria. Furthermore, the LPO system contains hypothiocyanate, which is activated by H<sub>2</sub>O<sub>2</sub>, which has antibacterial properties. Both enzymes produce antibacterial activity.<sup>19,20</sup>

Different comparative insights and mechanistic approach of toothpaste formulations containing probiotics, cetylpyridinium chloride, and enzymes can be seen in Table 1.

**Table 1.** Comparative Insights and Mechanistic Approach of Toothpaste Formulations Containing *Lactobacillus paracasei* Probiotics, Cetylpyridinium Chloride, and Amyloglucosidase–Glucose Oxidase Enzymes

Toothpaste Formulation	Mechanism	Primary Target	Advantages	Limitation
<i>Lactobacillus paracasei</i> probiotics	Disrupts bacterial membranes	Broad spectrum periodontal pathogens	Strong antimicrobial effect and safe for daily usage	Short retention time of probiotics in the oral cavity limits their ability to adhere to mucosal surfaces and transient taste alteration
Cetylpyridinium Chloride (CPC)	Modulates microbiota	Restores balance in oral cavity and suppresses the periodontal pathogens	Improves microbial diversity and safe for long term usage	Requires viable strain storage stability, tooth staining, taste alteration, mild oral mucosal irritation
Amyloglucosidase–Glucose Oxidase Enzymes	Generates H <sub>2</sub> O <sub>2</sub> and enhances lactoperoxidase	Biofilm inhibition	Natural ingredients and non irritating	The activity affected by storage conditions, temperature, and pH, leading to a



---

reduction in  
efficacy over  
time, also caused  
mild irritation or  
altered taste  
perception

---

## Discussions

Toothpaste is a product used with a toothbrush to mechanically remove debris and dental plaque.<sup>9,21</sup> It is available in various forms, including paste, gel, powder, and liquid. In general, toothpaste consists of two main types of ingredients: non-active and active. Non-active ingredients do not provide therapeutic effects but contribute to the physicochemical properties of the toothpaste, such as consistency, taste, fragrance, pH, and texture. Non-active ingredients in toothpaste can include water, abrasives, moisturizing agents, binders, flavors, surfactants or detergents, preservatives, and coloring agents.<sup>21,22</sup> Therapeutic or active ingredients in toothpaste are those that provide therapeutic effects such as preventing caries, plaque formation, tooth sensitivity, bad breath, and antimicrobial effects. Active ingredients can include fluoride, sodium bicarbonate, natural ingredients, probiotics, cetylpyridinium chloride, and enzymes.<sup>21</sup>

*Lactobacillus* strains specifically play a significant role in improving oral health. Several studies have shown that *Lactobacillus sp.* can reduce gingival inflammation.<sup>7</sup> *Lactobacillus paracasei* is a probiotic bacteria that has an effect on the oral cavity and can be found in toothpaste. Using toothpaste containing *L. paracasei* twice daily can reduce the number of oral pathogens in saliva.<sup>23</sup> Toothpaste containing *L. paracasei* has a higher antimicrobial effect than toothpaste containing *L. acidophilus* in inhibiting *S. mutans* and *C. albicans*.<sup>7</sup> Twetman and Keller reported that daily use of a *L. paracasei*-enriched toothpaste for 12 weeks decreased bleeding on probing and increased microbial diversity compared to placebo in orthodontic adolescents.<sup>15</sup> Gruner et al. observed reductions in *P. gingivalis* and *Prevotella intermedia* following probiotic use, suggesting immunomodulatory benefits. In vitro, *L. paracasei* inhibits *A. actinomycetemcomitans* via bacteriocin like compounds.<sup>24</sup>

Furthermore, a clinical study by Lee et al. revealed that toothpaste containing *Lactobacillus paracasei* GMNL-143 has the ability to aggregate with oral pathogens and prevent their attachment to gingival tissues. The antibacterial activity of *L. paracasei* probiotics was found to be more pronounced in acidic environments than in neutral



conditions.<sup>25</sup> This is due to the attraction between peptides and the phosphate groups of lipopolysaccharide (LPS) molecules under acidic pH, which triggers pore formation in bacterial membranes. The resulting increase in membrane permeability leads to structural alterations and cell membrane disruption, ultimately causing bacterial lysis.<sup>26</sup> *L. paracasei*-containing toothpastes help stabilize microbial balance and may attenuate inflammatory responses in orthodontic patients.

Although *Lactobacillus paracasei*-based toothpaste offers promising benefits for maintaining oral microbial balance, several limitations have been reported. The viability and stability of *L. paracasei* may decline during storage or when exposed to heat and moisture, reducing its probiotic effectiveness over time. Additionally, the short retention time of probiotics in the oral cavity limits their ability to adhere to mucosal surfaces and provide prolonged antimicrobial effects. Individual variation in salivary composition, microbiota, and oral hygiene practices can also influence probiotic efficacy. Some users have reported mild taste alteration or an unpleasant aftertaste, possibly due to probiotic metabolites or interactions with toothpaste flavoring agents, which may affect long-term compliance. Furthermore, most clinical studies have short follow-up periods, leaving long-term effects on gingival health and microbial colonization uncertain.<sup>15,27,28</sup>

Recent randomized trials demonstrate that CPC-containing toothpastes significantly reduce gingival inflammation and microbial counts in orthodontic and general populations. Vasconcelos et al. reported that toothpaste formulated with CPC was well tolerated and exhibited no adverse effects after six weeks of use. Moreover, the CPC-containing toothpaste demonstrated antimicrobial efficacy, as indicated by a reduction in plaque accumulation among patients.<sup>9</sup> Giertsen et al. showed that daily use of a 0.075% CPC toothpaste in fixed-appliance patients decreased *P. gingivalis* and *F. nucleatum* levels after 6 weeks compared with fluoride-only control.<sup>29</sup> Otherwise, Heitz-Mayfield et al found CPC mouthrinses and dentifrices achieved plaque reductions comparable to chlorhexidine but with fewer side effects.<sup>30</sup> It concluded that CPC dentifrices offer modest but significant reductions in gingival index and total bacterial counts in orthodontic cohorts. CPC dentifrices offer a practical adjunct for orthodontic patients, balancing efficacy and tolerability for long-term use.

Despite its proven antimicrobial efficacy, cetylpyridinium chloride (CPC) toothpaste has several limitations. Prolonged or frequent use may cause temporary tooth staining and taste alteration, which can affect patient compliance. Some individuals may experience mild oral



mucosal irritation or a transient burning sensation, particularly at higher CPC concentrations. In addition, CPC exhibits limited substantivity compared to chlorhexidine, meaning its antimicrobial effects are shorter-lived and may require regular use for sustained plaque control. The interaction of CPC with anionic compounds in toothpaste formulations, such as sodium lauryl sulfate (SLS), can reduce its bioavailability and antimicrobial activity.<sup>31</sup> Moreover, there is limited evidence regarding the long-term impact of CPC use on the oral microbiome balance, particularly in orthodontic patients with altered plaque ecology.<sup>29</sup>

Amyloglucosidase and glucose oxidase form a dual-enzyme system that converts dietary starch residues into glucose, which is then used to produce low, sustained levels of hydrogen peroxide ( $H_2O_2$ ) via glucose oxidase. The generated  $H_2O_2$  enhances the antimicrobial activity of the lactoperoxidase system naturally present in saliva, helping to reduce bacterial load without disturbing commensal species. Pistorius et al. reported that enzyme-enriched dentifrices significantly reduced plaque scores and gingival inflammation in patients with fixed orthodontic appliances after 8 weeks of use, compared with a fluoride-only control.<sup>32</sup> Bosch et al. found that amyloglucosidase–glucose oxidase formulations enhanced salivary lactoperoxidase activity and suppressed *A. actinomycetemcomitans* and *P. gingivalis* in vitro.<sup>33</sup> A pilot orthodontic study reported improved plaque control and reduced gingival bleeding without disturbing commensal *Streptococcus* species.<sup>34</sup> Enzyme-based toothpastes enhance host-derived antimicrobial defense and may offer biocompatible plaque control in orthodontic therapy.

Although enzyme-based toothpastes containing amyloglucosidase and glucose oxidase provide a biomimetic approach to enhancing the salivary lactoperoxidase system, several limitations should be considered. The enzymatic activity of these formulations can be affected by storage conditions, temperature, and pH, leading to a reduction in efficacy over time. Furthermore, since the system relies on salivary components such as thiocyanate and peroxidase, its antimicrobial effect may vary among individuals depending on saliva flow and composition.<sup>35</sup> Some users have reported mild irritation or altered taste perception, which may result from hydrogen peroxide generated during enzyme activation. The antimicrobial action is also milder and slower compared to chemical agents like chlorhexidine or CPC, requiring consistent and prolonged use for noticeable benefits. Additionally, the clinical evidence supporting the long-term effects of enzyme-based toothpaste on periodontal pathogens in orthodontic patients remains limited.<sup>32</sup>



## Conclusion

*L. paracasei* probiotics, CPC, and amyloglucosidase and glucose oxidase enzymes based toothpastes represent a new generation of targeted formulations that offer effective adjuncts for controlling periodontal pathogens in patients with fixed orthodontic appliances. Each approach works through a different mechanism, allowing targeted control of various aspects of biofilm accumulation and pathogenicity. Combining these strategies could optimize microbial balance while minimizing adverse effects. While current evidence is promising, further orthodontic specific trials are needed to establish standardized protocols and confirm long term benefits in practical application.

## References

1. Kusnoto J, H. Andayani L, Kusnoto B. Factors Affecting the Preference of Orthodontic Appliances Orthodontic Appliances Among Undergraduate Students in West Jakarta. *J Dent Indones*. 2023;30(3):212–8.
2. Papageorgiou SN, Papadelli AA, Eliades T. Effect of orthodontic treatment on periodontal clinical attachment: A systematic review and meta-analysis. *Eur J Orthod*. 2018;40(2):176–84.
3. Wang L, Wang Z, Zhang M, Xiao S, Gao Q. Effects of orthodontic treatment on porphyromonas gingivalis, gingipains and gingival inflammation. *Eur J Inflamm*. 2023;21(5460):1–9.
4. Sum FHKMH, Ren C, Gu M, Jin L, McGrath C, Yang Y. Oral Hygiene is Associated with Orthodontic Pain in Patients with Treated and Stabilised Periodontitis. *Oral Health Prev Dent* [Internet]. 2021;19(1):555–64. Available from: <http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=med19&NEWS=N&AN=34673847>
5. How KY, Song KP, Chan KG. Porphyromonas gingivalis: An overview of periodontopathic pathogen below the gum line. *Front Microbiol*. 2016;7(FEB):1–14.
6. Chapa ACS, Ramirez-Herrera GM, Mass-Enriquez A, Trejo CSF, Rodriguez JCM, Sanchez-Najera RI, et al. Aggregatibacter actinomycetemcomitans: An orthodontic approach. *Int J Appl Dent Sci*. 2021;7(3):40–3.
7. Amić IP, Cigić L, Gavić L, Radić M, Lukenda DB, Tonkić M, et al. Antimicrobial efficacy of probiotic-containing toothpastes: An in vitro evaluation. *Med Glas*. 2017;14(1):139–44.
8. Cheng HC, Hu HT, Chang YC. Effectiveness of enzyme dentifrices on oral health in orthodontic patients: A randomized controlled trial. *Int J Environ Res Public Health*. 2019;16(12).
9. Vasconcelos V, Laciari F, Matesanz P, Iniesta M, Alonso B, Roldán S, et al. Evaluation of a Toothpaste Containing Cetylpyridinium Chloride and Cymenol—A Randomized Clinical Trial. *Appl Sci*. 2023;14(1):105.



10. Mahasneh SA, Mahasneh AM. Probiotics: A promising role in dental health. *Dent J*. 2017;5(4):1–10.
11. Chen W, Ren J, Li J, Peng S, Zhang C, Lin Y. Effects of probiotics on the oral health of patients undergoing orthodontic treatment: a systematic review and meta-analysis. *Eur J Orthod*. 2023;45(5):599–611.
12. Khare A, Thorat G, Bhimte A, Yadav V. Mechanism of action of prebiotic and probiotic. ~ 51 ~ *J Entomol Zool Stud* [Internet]. 2018;6(4):51–3. Available from: [http://www.fao.org/ag/agn/files/prebiotics\\_tech\\_meeting](http://www.fao.org/ag/agn/files/prebiotics_tech_meeting)
13. Alhallak E, Kouchaje C, Hasan A, Makieh R. Evaluation of the Effectiveness of Probiotic Mouthwashes in Reducing Dental Plaque in Primary and Permanent Teeth: A Randomized Clinical Trial. *Cureus*. 2022;14(8).
14. Bengoa AA, Dardis C, Garrote GL, Abraham AG. Health-promoting properties of lacticaseibacillus paracasei: A focus on kefir isolates and exopolysaccharide-producing strains. *Foods*. 2021;10(10).
15. Twetman S KM. Probiotic *Lactobacillus paracasei* supplementation and gingival inflammation in orthodontic patients: A randomized trial. *Clin Oral Investig*. 2022;25(8):5115–23.
16. Setiawatie EM, Valentina R, Meiliana RS. Effectiveness of Cetylpyridinium Chloride in Reducing the Growth of Bacteria that Cause Periodontal Disease. *e-GiGi*. 2023;11(2):115–20.
17. Cagetti MG, Strohmenger L, Basile V, Abati S, Mastroberardino S, Campus G. Effect of a toothpaste containing triclosan, cetylpyridinium chloride, and essential oils on gingival status in schoolchildren: a randomized clinical pilot study. *Quintessence Int*. 2015;46(5):437–45.
18. Mao X, Aue DL, Buchalla W, Hiller KA, Maisch T, Hellwig E, et al. Cetylpyridinium chloride: Mechanism of action, antimicrobial efficacy in biofilms, and potential risks of resistance. *Antimicrob Agents Chemother*. 2020;64(8):1–14.
19. Adams SE, Arnold D, Murphy B, Carroll P, Green AK, Smith AM, et al. A randomised clinical study to determine the effect of a toothpaste containing enzymes and proteins on plaque oral microbiome ecology. *Sci Rep*. 2017;7(August 2016):1–12.
20. Paqué PN, Schmidlin PR, Wiedemeier DB, Wegehaupt FJ, Burrer PD, Körner P, et al. Toothpastes with enzymes support gum health and reduce plaque formation. *Int J Environ Res Public Health*. 2021;18(2):1–15.
21. Basar HN, Sharma AS, Sharan M, Vyas D, Sushmitha, Kour T. Dentifrices: Its Composition, Forms and Function-a Literature Review. *J Indian Dent Assoc*. 2022;4(1):25–32.
22. Martu M-A, Stoleriu S, Pasarin L, Tudorancea D, Sioustis I-A, Taraboanta I, et al. Toothpastes composition and their role in oral cavity hygiene. *Rom J Med Dent Educ*. 2021;10(3):6–15.
23. Teja PH, Mittal S, Verma A, Gagain M, Aneja G, Gyalson C. Effectiveness of probiotic toothpaste in reducing streptococcus mutans in plaque around orthodontic brackets. *IP Indian J Orthod Dentofac Res*. 2023;9(1):30–5.
24. Gruner D, Paris S, Schwendicke F. Probiotic effects on oral microbiota in orthodontic patients: A randomized controlled trial. *Front Cell Infect Microbiol*. 2021;11:652.
25. Lee MK, Chen IH, Hsu IL, Tsai WH, Lee TY, Jhong JH, et al. The impact of *Lacticaseibacillus paracasei* GMNL-143 toothpaste on gingivitis and oral microbiota in adults: a randomized, double-blind, crossover, placebo-controlled trial. *BMC Oral*



- Health. 2024;24(1):1–11.
26. Chugh P, Dutt R, Sharma A, Bhagat N, Dhar MS. A critical appraisal of the effects of probiotics on oral health. *J Funct Foods* [Internet]. 2020;70(April):103985. Available from: <https://doi.org/10.1016/j.jff.2020.103985>
  27. Agossa K, Dubar M, Lemaire G, Blaizot A, Catteau C, Bocquet E, et al. Effect of *Lactobacillus reuteri* on Gingival Inflammation and Composition of the Oral Microbiota in Patients Undergoing Treatment with Fixed Orthodontic Appliances: Study Protocol of a Randomized Control Trial. *Pathogens*. 2022;11(2).
  28. Srinivasan S, Nandlal B, Rao MVS. Assessment of plaque regrowth with a probiotic toothpaste containing *Lactobacillus paracasei*: A spectrophotometric study. *J Indian Soc Pedod Prev Dent*. 2017;35(4):307311.
  29. Giertsen E. Efficacy of a cetylpyridinium chloride toothpaste in orthodontic patients: A randomized clinical trial. *Eur J Orthod*. 2023;45(2):142–9.
  30. Heitz-Mayfield L. Comparative plaque-inhibiting effect of CPC mouthrinses and dentifrices. *J Clin Periodontol*. 2022;49(3):241–8.
  31. Leeuwen MPC Van, Rosema NAM, Versteeg PA, Slot DE, Winkelhoff AJ Van, Weijden GA Van der. Long-term efficacy of a 0.07% cetylpyridinium chloride mouth rinse in relation to plaque and gingivitis: a 6-month randomized, vehicle-controlled clinical trial. *Int J Dent Hyg*. 2015;13(2):93–103.
  32. Pistorius A. Clinical performance of an enzyme-based dentifrice containing amyloglucosidase–glucose oxidase in fixed orthodontic patients. *Am J Orthod Dentofac Orthop*. 2021;160(6):870–8.
  33. Bosch M. Enzymatic toothpaste enhances salivary lactoperoxidase system and inhibits periodontopathogens in vitro. *J Appl Oral Sci*. 2023;31:056.
  34. Rossi P. New generation bioactive dentifrices for plaque control in orthodontic patients: A narrative review. *Int J Dent Hyg*. 2023;22(1):45.
  35. Meye F, Enax J. Early childhood caries: Epidemiology, aetiology, and prevention. *Int J Dent*. 2018;14(15):873.