

# The Effect of 96% Ethanol Extract of Bidara Leaf (*Ziziphus Mauritiana*) on the Hemostatic Activity in Sprague-Dawley Rats

*by* Moehamad Orliando Roeslan

---

**Submission date:** 01-Jul-2025 08:59AM (UTC+0700)

**Submission ID:** 2667786210

**File name:** dara\_Leaf\_Ziziphus\_Mauritiana\_on\_the\_Hemostatic\_Activity\_in.pdf (263.46K)

**Word count:** 4268

**Character count:** 23044

## The Effect of 96% Ethanol Extract of Bidara Leaf (*Ziziphus Mauritiana*) on the Hemostatic Activity in Sprague-Dawley Rats

Irvan Septrian Syah Putra Rasad<sup>1</sup>, Wiwiek Poedjastoeti<sup>1\*</sup>, Moeahamad Oriando Roeslan<sup>2</sup>

1. Oral and Maxillofacial Surgery Department, Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia.  
2. Oral Biology Department, Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia.

### Abstract

Objective to investigate the efficacy of the 96% ethanol extract of *Ziziphus mauritiana* leaves on hemostasis activity in animal model study (Sprague-Dawley rats).

A laboratory experimental in vitro and in vivo study were conducted with Sprague-Dawley (SD) rats with randomized controlled design with a posttest-only control group design. Fifteen animals were used as research material, divided into 5 groups, negative control group (NCG), positive control group with golden standard hemostatic agent tranexamic acid 5% (TXA), 25% concentration of *Z. mauritiana* leaves extract (ZM25), 50% concentration (ZM50), and 100% concentration (ZM100). The bleeding time and the amount of bleeding were calculated by cutting the tip of the tails of the SD rats. PT, aPTT and TT were measured by taking blood samples from the SD rats. The data analysed using the Shapiro Wilk normality test, normally distributed data followed by a one-way ANOVA test with a significance level ( $p < 0.05$ ), continued with the Post-Hoc test.

The ZM100 group shows a significant difference with NCG, TXA and ZM25 groups in bleeding time. The NCG group shows a significant differences with all samples in the amount of bleeding. The ZM100 group shows a significant difference with all test groups in PT, aPTT and TT test.

The ethanol extract of *Z. mauritiana* leaves can accelerate bleeding time and the amount of bleeding, the extract can also stimulate the acceleration of PT, aPTT and TT factor in the bleeding cascade at high concentrations (100%).

Experimental article (J Int Dent Med Res 2023; 16(4): 1408-1413)

**Keywords:** Hemostasis, *Ziziphus mauritiana*, Bleeding Time, Prothrombin Time, Activated Partial Thromboplastin Time, Thrombin Time.

Received date: 22 July 2023

Accept date: 24 August 2023

### Introduction

Tooth extraction is one of the most frequently performed dental procedures, with bleeding as one of the most common complications.<sup>1</sup> Uncontrolled bleeding can threaten the patient's life and will interfere the dentists when performing tooth extraction procedure.<sup>2</sup> Incidence of bleeding after tooth extraction can be minimized by systemic or local interventions. Local post-extraction bleeding interventions can be carried out in various ways, some of which are by: pressure tampons, sutures, acrylic splints and the appliance of local hemostasis agents.<sup>3</sup>

In general, the wound healing process is

divided into four phases, hemostasis, inflammation, proliferation, and remodelling phase.<sup>4</sup> Hemostasis occurs at the beginning of the wound healing process, followed by an inflammatory phase characterized by swelling and pain, then soft tissue structures re-form in the proliferation phase and end with complete tissue structure restoration in the remodelling phase.<sup>4,5</sup> Hemostasis (the process of bleeding to stop) is a complex mechanism influenced by interactions between the plasma coagulation cascade, fibrinolytic proteins, the vascular system, and thrombin (platelets).<sup>6</sup> The process of hemostasis is characterized by the occurrence of vasoconstriction and blood clot formation through activation of platelets and the coagulation cascade.<sup>7</sup>

Indonesia is a country that is rich in a variety of herbal plants that can be used as medicine. One of the plants in Indonesia that has many benefits is the Bidara plant or Indian Jujube plant (*Ziziphus mauritiana*). These plants are

\*Corresponding author:

Wiwiek Poedjastoeti,  
Oral and Maxillofacial Surgery Department, Faculty of Dentistry,  
Universitas Trisakti, Jakarta, Indonesia.  
E-mail: wiwiek@trisakti.ac.id

found in deserts, tropical and subtropical climates country.<sup>8-10</sup> Dried *Z. mauritiana* leaves are known to contain complex secondary metabolites such as tannins.<sup>8</sup> In several different tree samples, it was found that the tannin content in *Z. mauritiana* plants had the highest content in the leaves.<sup>10</sup> *Ziziphus mauritiana* is known to have various benefits such as antimicrobial, analgesic, antipyretic, anti-inflammatory and anticancer agent. *Ziziphus mauritiana* works pharmacodynamically and pharmacokinetically by increasing cellular activities, such as migration, proliferation, adhesion, and differentiation to improve wound healing.<sup>11</sup> In the phytochemical test and quantification *Z. mauritiana* has secondary metabolites such as alkaloids, flavonoids, saponins, tannins and terpenoids. Bark, fruit flesh and leaves of *Z. mauritiana* are known to have astringent properties so that they can be used to accelerate cessation of bleeding and accelerate wound healing process.<sup>8,12</sup> The astringent substances contained in some plants have a contraction effect, causing tissues to occur shrinkage.<sup>13,14</sup>

Tannin is an astringent substance that can precipitate thrombin which will then convert fibrinogen into a collection of fibrin fibers thread in the area of bleeding, these fibrin fibers will become a place for blood clotting proteins to attach so that the coagulation process can be achieved.<sup>15</sup> Tannins are also known to have a vasoconstrictor effect so that bleeding time will be shortened and the wound healing process can be achieved.<sup>16,17</sup>

As far as the researchers know, there is not much literature discussing the hemostasis activity of *Z. mauritiana* leaves. Based on the descriptions, researchers are interested in researching on the activity of 96% ethanol extract of *Z. mauritiana* leaves against the hemostasis process in experimental animal model study Sprague-Dawley (SD) rats.

#### Materials and methods

The type of study is laboratory experimental research in vitro and in vivo with randomized controlled design with posttest-only control group. This study was cleared for ethics by the Ethical Commission of the Faculty of Dentistry at Universitas Trisakti (No. 636/S2/KEPK/FGK/2/2023). The study procedures are the preparation of extract ethanol

*Z. mauritiana* leaves, and phytochemical examination. The next procedures were in vivo and in vitro studies of the hemostasis activity of the 96% ethanol extract of *Z. mauritiana* leaves.

Preparation of *Z. mauritiana* Leaves Extract

*Ziziphus mauritiana* leaves weighing 4 kg were taken from the Indonesian Medicinal and Aromatic Crops Research Institute (IMACRI) also known as Balittro, the leaves had been dried and identified by the National Research and Innovation Agency (BRIN). The dried powder leaves are macerated in the BioCore Laboratory, Faculty of Dentistry, Universitas Trisakti with 96% ethanol (Supelco-Merck, German) for 3 days and stirred every 4 hours before decanted and filtered. Then, the filtrate was evaporated with a rotary vacuum evaporator (Buchi R-215, Flawil, Switzerland) until a concentrated extract (crude) was obtained. To remove the remaining ethanol, the crude was heated at 45°C, the crude then dissolved with aquadest, and divided into several concentrations, 25%, 50%, and 100% *Z. mauritiana* extract ethanol.

Phytochemical Assay of *Z. mauritiana* Leaves

A phytochemical test was conducted at BioCore Laboratory to identify the active compounds contained in the 96% ethanol extract of *Z. mauritiana* leaves. The phytochemical compounds investigated in this study are flavonoid, steroid, terpenoid, alkaloid, and tannin.

To perceive the flavonoid composition, 5 mL dilute ammonia is added to a portion of an aqueous filtrate of the extract and then 1 mL concentrated sulphuric acid was added, the yellow colouration disappears when left to stand indicates the presence of flavonoids. To perceive steroid composition, 3 drops of concentrated hydrochloric acid and 1 drop of concentrated sulfuric acid are added to 2 mL of the extract, the colour changes to bluish green indicates the presence of steroids. To perceive terpenoid composition, 2 mL of chloroform added to the 1 mL extract and then added 3 mL of concentrated sulfuric acid, the colour changes to reddish brown indicates the presence of terpenoid. To perceive alkaloid composition, dragendorff reagent is added to the extract, the depositions of orangish sediment indicates the presence of alkaloid. To perceive terpenoid. To perceive tannin, 0.1% ferric chloride added to 1 mL sampel, the colour changes to bluish indicates

the presence of tannin.

#### In Vivo Assessments of Hemostasis

A total of 15 male SD rats were adapted for about 1 week for the acclimatization process, the need to eat and drink ad libitum. Experimental animals were randomly divided into 5 groups (n=3), negative control group (NCG) using 0.9% sodium chloride (NaCl) solution (B. Braun, Melsungen, Germany), positive control group with the golden standard drug tranexamic acid (TXA) solution (Bernofarm, Sidoarjo, Indonesia), treatment group with 25% Z. mauritiana leaves extract (ZM25), treatment group with 50% Z. mauritiana leaves extract (ZM50) and the treatment group with 100% Z. mauritiana leaves extract (ZM100).

The SD rats on each groups were anesthetized with ketamine 40 mg/kg BW (Agrovot Market, Lima, Peru) and xylazine 5 mg/kg BW (Interchemie, Venray, Netherlands). Later, the surgical area was cleaned using a 2% chlorhexidine solution and 1 cm distal segment from the tip of the tail of the rats were amputated with a scalpel blade no 15. Then, 0.1 mL of the solution was applied using an insulin syringe (Onemed, Sidoarjo, Indonesia). A digital chronometer is used to measure how long it takes to achieve hemostasis. Timing begins at the time of incision and stops when the bleeding at the site stops completely. To determine the amount of bleeding, each filtration paper (Whatman - Cytiva, Marlborough, US) was weighed in gram before the procedure with a precision laboratory scale (Sartorius M-Prove, Göttingen, Germany). After the bleeding was achieved, the filtration paper was used to absorb the bleeding that occurred in the wound on the SD rat's tail. This process is carried out until the bleeding stops as indicated by no more blood spots adhering to the filter paper. The difference in the weight of the filtration paper before and after the experiment was considered as the amount of bleeding.

#### In Vitro Assessments of Hemostasis

In the in vitro study, Prothrombin Time (PT), Activated Partial Thromboplastin Time (aPTT), and Thrombin Time (TT) are the hemostasis factors assessed in this study. Blood samples (n=5) were taken from SD rats in the previous in vivo study, blood sampling was carried out through the aorta and directly added to plastic tubes containing 0.2 mL sodium citrate 3.2% as an anticoagulant in a 1:9 (v/v) ratio to

the blood. Blood samples were centrifuged at 3000 rpm for 10 minutes to separate blood plasma, then the supernatant plasma, which contains platelets, are divided in five sterile tubes for the coagulation test. The coagulation test such as PT, aPTT and TT were examined with reagents kit (Stago, Asnières, French) according to the manufacturer's instructions.

For the PT test, the plasma sample is divided into five tubes, each containing 0.1 mL of plasma which will then be treated by mixing the reagent with the research material. the first tube will only be given reagents as a negative control group, the second tube is given 0.1 mL 5% TXA as a positive control, the third tube is 0.1 mL ZM25, the fourth tube is 0.1 mL ZM50 and the fifth tube is 0.1 mL ZM100, all the tubes were incubated for two minutes at 37°C. Then, 0.2 mL of warmed PT reagent was added to the tube, and the clot formation time was calculated using digital chronometer from the start of the reagent drop until a coagulation formed at the bottom of the tube. The same protocols were delivered for aPTT and TT test depending on the reagents manufacturer's instructions, each protocol was repeated five times and assessed by blind investigators.

#### Statistical Analysis

The data were analysed using the Shapiro Wilk normality test, normally distributed data followed by a one-way ANOVA test with a significance level ( $p \leq 0.05$ ), continued with the Post-Hoc test.

#### Results

From the results of the phytochemical test of the ethanol extract of Z. mauritiana leaves it was found that it contains active substances such as terpenoids, flavonoids, alkaloids, and tannins, while steroids are absence in this phytochemical test. The results of the phytochemical test are shown in Table 1.

The results of the bleeding time test shows that NCG group is significantly different ( $p < 0.05$ ) with all groups and the ZM100 group shows a significant difference ( $p < 0.05$ ) with the negative control, TXA and ZM25 groups. The results of the bleeding amount test shows the NCG group is significantly different ( $p < 0.05$ ) with all groups (Table 2). In the PT test results, the NCG and TXA groups show a significant difference ( $p < 0.05$ ) with the ZM50 and ZM100

groups. In the aPTT test results, the NCG group and the ZM100 group are significantly difference ( $p < 0.05$ ) with all groups. In the TT test results, the ZM50 group is significantly different with the TXA and ZM100 groups (Table 3).

Metabolit Sekunder	Metode	Hasil Uji
Terpenoid	2mL CHCl <sub>3</sub> + 3mL H <sub>2</sub> SO <sub>4</sub>	+
Flavanoid	5mL NH <sub>3</sub> + 1mL H <sub>2</sub> SO <sub>4</sub>	+
Steroid	3 drops of CH <sub>3</sub> COOH + 1 mL H <sub>2</sub> SO <sub>4</sub>	-
Alkaloid	Dragendorff's reagent	+
Tanin	1 mL FeCl 0.1%	+

**Table 1.** The phytochemical analysis of 96% ethanol extract of *Ziziphus mauritiana* leaves.

Treatment Group (n=3)	Bleeding Time (s)	Amount Bleeding (g)
NCG	4.11 ± 0.84 <sup>a</sup>	1.74 ± 0.15 <sup>a</sup>
TXA	2.40 ± 0.15 <sup>b</sup>	1.33 ± 0.11 <sup>b</sup>
ZM25	2.08 ± 0.26 <sup>bc</sup>	1.27 ± 0.04 <sup>b</sup>
ZM50	1.50 ± 0.04 <sup>cd</sup>	1.36 ± 0.14 <sup>b</sup>
ZM100	0.94 ± 0.53 <sup>d</sup>	1.32 ± 0.17 <sup>b</sup>

**Table 2.** Hemostatic analysis of the bleeding time and amount of blood, for the different groups evaluated (in vivo).

Abbreviations: NCG, negative control (NaCl); TXA, positive control (5% Tranexamic Acid); ZM25, 25% concentration of *Z. mauritiana* leaves ethanol extract; ZM50, 50% concentration of *Z. mauritiana* leaves ethanol extract; ZM100, 100% concentration of *Z. mauritiana* leaves ethanol extract.

Note: The values represent the mean values and the standard deviation for the three animals evaluated in each group, different superscript letters (abcd) expressed statistically significant differences ( $p < 0.05$ ) in the lines.

Treatment Group (n=5)	PT (s)	aPTT (s)	TT (s)
NCG	37.58 ± 11.92 <sup>a</sup>	31.13 ± 0.89 <sup>a</sup>	14.64 ± 3.57 <sup>abc</sup>
TXA	36.43 ± 2.40 <sup>ab</sup>	20.14 ± 3.04 <sup>b</sup>	13.45 ± 2.57 <sup>b</sup>
ZM25	33.67 ± 4.99 <sup>abc</sup>	20.75 ± 1.39 <sup>b</sup>	18.78 ± 4.73 <sup>c</sup>
ZM50	26.21 ± 7.32 <sup>c</sup>	20.97 ± 2.69 <sup>b</sup>	21.53 ± 9.95 <sup>ac</sup>
ZM100	12.15 ± 1.90 <sup>d</sup>	17.10 ± 1.76 <sup>c</sup>	04.35 ± 2.43 <sup>d</sup>

**Table 3.** Hemostatic analysis of the PT, aPTT, and TT time, for the different groups evaluated (in vitro).

Abbreviations: NCG, negative control (Reagent); TXA, positive control (5% Tranexamic Acid); ZM25, 25% concentration of *Z. mauritiana* leaves ethanol extract; ZM50, 50% concentration of *Z. mauritiana* leaves ethanol extract; ZM100, 100% concentration of *Z. mauritiana* leaves ethanol extract.

Note: The values represent the mean values and the standard deviation for the three animals evaluated in each group, different superscript letters (abcd) expressed statistically significant differences ( $p < 0.05$ ) in the lines.

## Discussion

*Ziziphus mauritiana* plant is believed to have benefits against various diseases. The potential health benefits of *Z. mauritiana* are known as anti-carcinogenicity, antioxidant, anti-

inflammatory, immune stimulation, anti-obesity, gastrointestinal protective activity, hepatoprotective and inhibitor of foam cell formation in macrophages.<sup>18</sup> In this study, the leaves of *Z. mauritiana* is proven to contain several secondary metabolites such as terpenoids, flavonoids, alkaloids and tannins but not steroids. The findings of this study are in line with previous studies that reported *Z. mauritiana* leaves revealed the presence of alkaloids, terpenoids, phenolic substances, saponins, flavonoids, and tannins. The steroids in the studies are also absence.<sup>19-21</sup> In several different tree samples, it was found that the tannin content in *Z. mauritiana* plants had the highest content in the leaves.<sup>10,20</sup>

Tannins can accelerate the wound healing process through several cellular mechanisms, prevent free radicals and reactive oxygen species, promote wound vasoconstriction and increase the activity of forming capillaries and fibroblasts including the proliferation of keratinocytes.<sup>22,23</sup> Tannins have high hemostasis efficiency because they can conjugate with various proteins in the blood, causing instant coagulation.<sup>24</sup> Hemostasis is a process in which bleeding becomes reduced, and blood clots form to prevent further blood loss. Hemostasis occurs in two stages: the primary stage, in which the blood vessels contract causing a platelet plug to form, and the secondary stage, in which the plug is strengthened with fibrin strands to hold the clot in place through activation of the coagulation cascade.<sup>25</sup> Therefore, the astringent properties of tannins are expected to stop bleeding in damaged blood vessels, accelerate both the primary and secondary stages of hemostasis so the cessation of bleeding is completely achieved. Primary hemostasis can be assessed by bleeding time and amount of bleeding tests, while secondary hemostasis can be assessed by PT, aPTT and TT tests.<sup>26</sup>

In this study, the hemostasis activity of the ethanol extract of *Z. mauritiana* leaves was investigated using the rat tail tip amputation model for bleeding time and amount of bleeding test, which is a standard and reliable procedure for assessing the hemostasis properties of a new biomaterial.<sup>25,27</sup> The bleeding time test shows the negative control (NCG) had the longest bleeding time with 4.11 seconds and showed a significant difference ( $p < 0.05$ ) with all groups. While the ZM100 group was the group with the



fastest bleeding time with 0.94 seconds and showed a significant difference ( $p < 0.05$ ) with the NCG, TXA and ZM25 groups. The results of the bleeding amount test showed that the NCG group had the highest amount of bleeding with 1.74 grams, and showed a significant difference ( $p < 0.05$ ) with all groups.

These results are harmonious with studies investigating the therapeutic properties of tannins in *Z. mauritiana*. Tannins, expressively found in extracts, have demonstrated the ability to control bleeding through their astringent effects on contracting vessels and damaged tissues, accelerating blood proteins and supporting hemostasis.<sup>21,25,26</sup>

In the PT test shows, the negative control (reagent) and TXA groups show a significant difference ( $p < 0.05$ ) with the ZM50 and ZM100 groups. The ZM100 group is the group with the fastest PT with 12.15 seconds and shows a significant difference ( $p < 0.05$ ) with all groups. In the aPTT test shows, the reagent group was the group with the longest aPTT with 31.13 seconds, and shows a significant difference ( $p < 0.05$ ) with all groups. While the ZM100 group is the group with the fastest aPTT with 17.10 seconds and shows a significant difference ( $p < 0.05$ ) with all groups. In the TT test shows, the ZM50 group is the group with the longest TT with 21.53 seconds, and shows a significant difference with the TXA and ZM100 groups. While the ZM100 group was the group with the fastest TT with 4.35 seconds and shows a significant difference with all test groups.

This shows that the administration of ethanol extract of *Z. mauritiana* leaves can accelerate the process of secondary hemostasis at high concentrations (ZM100) compared to other control groups and lower concentrations. The higher the viscosity of the ethanol extract of *Z. mauritiana* leaves, the higher the composition of the active substance, so that wound healing can be improved due to its anti-inflammatory and antibacterial properties.<sup>19</sup>

Based on the results of the study, the ethanol extract of *Z. mauritiana* leaves showed therapeutic potential that could be used as a promising biomaterial for hemostasis agents in the clinical treatment of local bleeding. Further research is needed regarding the activity of the ethanol extract of *Z. mauritiana* leaves in a smaller concentration whether it can maintain the same efficacy and mechanism, and it is also

necessary to carry out clinical trials in humans to confirm the hemostasis potential shown in this study.

### Conclusions

The 96% ethanol extract of *Z. mauritiana* leaves can accelerate the cessation of bleeding time and the amount of bleeding, the extract can also stimulate the acceleration of extrinsic (PT) and intrinsic (aPTT) factors of the bleeding cascade, and accelerate the formation of fibrin clot (TT) at a high concentrations so that it can be developed as a topical hemostatic agent.

### Declaration of Interest

The authors report no conflict of interest.

### References

1. Kumbargere Nagraj S, Prashanti E, Aggarwal H, et al. Interventions for treating post-extraction bleeding. *Cochrane Database of Systematic Reviews*. 2018;2018(3):4-10. doi:10.1002/14651858.CD011930.pub3
2. Roeslan MO, Reskia SH, Firman FR, Dewa HS, Komariah K, Bustami DA. Effect of clinacanthus nutans leaf extract on oral mucosal burns and tongue wounds: An in-vivo study. *Scientific Dental Journal*. 2021;5(3):101-108. doi:10.4103/sdj.sdj\_92\_21
3. Gupta G, M.R. M. Kumar SP. Efficacy of Hemocoagulase as a Topical Hemostatic Agent After Dental Extractions: A Systematic Review. *Cureus*. Published online March 31, 2018. doi:10.7759/cureus.2398
4. Velnar T, Bailey T, Smrkolj V. The wound healing process: An overview of the cellular and molecular mechanisms. *Journal of International Medical Research*. 2009;37(5):1528-1542. doi:10.1177/147323000903700531
5. Alam G, Singh MP, Singh A. WOUND HEALING POTENTIAL OF SOME MEDICINAL PLANTS. *International Journal of Pharmaceutical Sciences Review and Research*. 2011;9(1):138-143.
6. Periyah MH, Halim AS, Saad AZM. Mechanism Action of Platelets and Crucial Blood Coagulation Pathways in Hemostasis. *International Journal of Hematology-Oncology and Stem Cell Research*. 2017;11(4):321-325.
7. Teller P, White TK. The Physiology of Wound Healing: Injury Through Maturation. *Surgical Clinics of North America*. 2009;89(3):599-610. doi:10.1016/j.suc.2009.03.006
8. Prakash O, Usmani S, Singh R, Singh N, Gupta A, Ved A. A panoramic view on phytochemical, nutritional, and therapeutic attributes of *Ziziphus mauritiana* Lam.: A comprehensive review. *Phytotherapy Research*. 2021;35(1):63-77. doi:10.1002/ptr.6769
9. Dhileepan K. Biological control of *Ziziphus mauritiana* (Rhamnaceae): feasibility, prospective agents and research gaps. *Annals of Applied Biology*. 2017;170(3):287-300. doi:10.1111/aab.12338
10. Yahia Y, Benabderrahim MA, Tili N, Bagues M, Nagaz K. Bioactive compounds, antioxidant and antimicrobial activities of extracts from different plant parts of two *Ziziphus* Mill. species. *PLoS One*. 2020;15(5):2-11. doi:10.1371/journal.pone.0232599.
11. Arma U, Widyawati W, Bakar A, Gani BA. Role of *Ziziphus mauritiana* Lam. on Oral Candidiasis and Its Relation to the Antibody Response, Blood Electrolyte, and Liver Profile. *J Int Dent Med Res (JIDMR)*. 2022;15(2):564-573.

12. Nairfana I, Nikmatullah A, Sarjan M, Tandeang A. Variability of secondary metabolites from leaves of *Ziziphus mauritiana* obtained from different locations in Sumbawa, Indonesia. *Biodiversitas Journal of Biological Diversity*. 2022;23(9):4948-4957. doi:10.13057/biodiv/d230965
13. Pizzorno JE, Murray MT, Joiner-Bey H. *Textbook Of Natural Medicine*. 3rd ed. Elsevier; 2015:1612-1615.
14. Wang M, Septier C, Brignot H, Martin C, Canon F, Feron G. Astringency Sensitivity to Tannic Acid: Effect of Ageing and Saliva. *Molecules*. 2022;27(5):232-233. doi:10.3390/molecules27051617
15. Sander Y, Ariesanti Y, Roeslan MO. Effect of moringa oleifera leaves on human blood coagulation process. 2021 IEEE International Conference on Health, Instrumentation & Measurement, and Natural Sciences (InHeNce). Published online 2021:1-5. doi:10.1109/Intence52833.2021.9537224.
16. Flanagan D. Tranexamic acid tamponade to control postoperative surgical hemorrhage. *Journal of Oral Implantology*. 2015;41(3):e82-e89. doi:10.1563/AID-JOI-D-13-00158
17. Kalalinia F, Amiri N, Mehrvarzian N, et al. Topical green tea formulation with anti-hemorrhagic and antibacterial effects. *Iran J Basic Med Sci*. 2020;23(8):1085-1090. doi:10.22038/IJBMS.2020.41397.9782
18. Paudel MR, Poudeyal MR, Devkota HP. *Ziziphus* spp. (*Ziziphus jujuba* mill., *Ziziphus Mauritiana* Lam.). Himalayan Fruits and Berries. 2023;1(45):491-497. doi:10.1016/b978-0-323-85591-4.00013-1
19. Maruf MT, Dewi PS, Nurlitasari DF. Efficacy of Bidara Leaf (*Ziziphus Mauritiana*) Viscous Extract to Gingival Wound Healing in Wistar Rats. *J Int Dent Med Res (JIDMR)*. 2021;14(4):1367-1371.
20. Abdallah EM, Elsharkawy ER, Ed-dra A. Biological activities of methanolic leaf extract of *Ziziphus mauritiana*. *Biosci Biotechnol Res Commun*. 2016;9(4):605-614. doi:10.21786/bbrc/9.4/6
21. Butt SZ, Hussain S, Munawar KS. Phytochemistry of *Ziziphus mauritiana*: An overview of its nutritional and pharmaceutical potential. *Sci Inquiry Rev*. 2021;5(2):01-15. doi:10.32350/sir.52.01
22. Sumanth M, Bhargavi YR. Evaluation of wound-healing effect of *Ziziphus mauritiana* L. leaf extract in rats. *International Journal of Green Pharmacy*. 2014;8(4):263-265. doi:http://dx.doi.org/10.4103/0973-8258.142687
23. Ariesanti Y, Rasad ISSP, Nimas M, Syabilla N. The effect of *Pearsea americana* Mill. seed extract on inflammatory cells and fibroblast formation in tooth extraction socket healing. *Dent J*. 2021;54(4):190-194. doi:10.20473/J.DJMKGV54.I4.P190-194
24. Chen C, Yang H, Yang X, Ma Q. Tannic acid: A crosslinker leading to versatile functional polymeric networks: A review. *RSC Adv*. 2022;12(13):7689-7711. doi:10.1039/d1ra07657d
25. Dantas MVO, Nogueira PL, De Oliveira Lima F, et al. In vivo Hemostatic Activity of *Jatropha mollissima*: A Triple-Blinded, Randomized, Controlled Trial in an Animal Model. *Eur J Dent*. 2021;15(4):741-745. doi:10.1055/s-0041-1729677
26. Gedar Totuk ÖM, Güzel ŞE, Ekici H, Kumandaş A, Emre Aydingöz S, Yılmaz EÇ, Kırdan T, Midi A. Effects of Algal Hemostatic Agent on bleeding time in a rat tail hemorrhage model. *Ulus Travma Acil Cerrahi Derg*. 2020;26(6):853-858. doi:10.14744/ijtes.2020.50384
27. Ebrahimi F, Mahmoudi J, Torbati M, Karimi P, Valizadeh H. Hemostatic activity of aqueous extract of *Myrtus communis* L. leaf in topical formulation: In vivo and in vitro evaluations. *J Ethnopharmacol*. 2020;249(1):1-13. doi:10.1016/j.jep.2019.112398

# The Effect of 96% Ethanol Extract of Bidara Leaf (Ziziphus Mauritiana) on the Hemostatic Activity in Sprague-Dawley Rats

## ORIGINALITY REPORT

17%

SIMILARITY INDEX

13%

INTERNET SOURCES

13%

PUBLICATIONS

7%

STUDENT PAPERS

## PRIMARY SOURCES

- |          |  |    |
|----------|--|----|
| <b>1</b> | <b>Submitted to iGroup</b><br>Student Paper  | 2% |
| <hr/>    |  |    |
| <b>2</b> | Fatemeh Ebrahimi, Javad Mahmoudi, Mohammadali Torbati, Poursan Karimi, Hadi Valizadeh. "Hemostatic activity of aqueous extract of Myrtus communis L. leaf in topical formulation: In vivo and in vitro evaluations", Journal of Ethnopharmacology, 2020<br>Publication | 2% |
| <hr/>    |  |    |
| <b>3</b> | web.mda.org.my<br>Internet Source  | 1% |
| <hr/>    |  |    |
| <b>4</b> | cdn.fortunepublish.com<br>Internet Source  | 1% |
| <hr/>    |  |    |
| <b>5</b> | Yessy Ariesanti, Irvan Septrian Syah Putra Rasad, Maylan Nimas, Nadira Syabilla. "The effect of Persea americana Mill. seed extract on inflammatory cells and fibroblast formation in tooth extraction socket healing", Dental Journal, 2021<br>Publication            | 1% |
| <hr/>    |  |    |
| <b>6</b> | T. Pullaiah, Maddi Ramaiah. "Handbook of Research on Herbal Liver Protection - Hepatoprotective Plants", CRC Press, 2021<br>Publication  | 1% |
| <hr/>    |  |    |
| <b>7</b> | Rondón María, Moncayo Shirley, Cornejo Xavier, Santos Jaime, Villalta David, Siguencia Rosa, Duche Jodie. "Preliminary phytochemical screening, total phenolic content and antibacterial activity of thirteen  | 1% |



native species from Guayas province  
Ecuador", Journal of King Saud University -  
Science, 2018

Publication

---

8	<a href="http://perpustakaan.poltekkes-malang.ac.id">perpustakaan.poltekkes-malang.ac.id</a> Internet Source	1 %
9	Submitted to Universiti Teknologi MARA Student Paper	1 %
10	<a href="http://www.escholar.manchester.ac.uk">www.escholar.manchester.ac.uk</a> Internet Source	1 %
11	<a href="http://www.topclassglobaljournals.org">www.topclassglobaljournals.org</a> Internet Source	1 %
12	Submitted to Herzing University Student Paper	1 %
13	Submitted to Fakultas Kedokteran Gigi Universitas Trisakti Student Paper	1 %
14	<a href="http://ijpsr.com">ijpsr.com</a> Internet Source	1 %
15	<a href="http://www.e-journal.trisakti.ac.id">www.e-journal.trisakti.ac.id</a> Internet Source	1 %
16	Abiodun, F, J Osakue, AS Uzoekwe, and S-X Qiu. "Phytochemical and Anticancer Studies on Ten Medicinal Plants Used in Nigeria", Bayero Journal of Pure and Applied Sciences, 2011. Publication	1 %
17	Atik Kurniawati, Yuli Dwi Kristanti, Naila Azifatur Rahmat, Yani Corvianindya Rahayu, Zainul Cholid, Agung Sosiawan. "The role of purple leaves extract (Graptophyllum Pictum (L.) Griff) on the number of fibroblasts and blood vessels in the socket after tooth extraction", Dental Journal, 2024 Publication	1 %

---

- 18 Submitted to Fakultas Teknologi Kebumian dan Energi Universitas Trisakti <1 %  
Student Paper
- 
- 19 Submitted to Far Eastern University <1 %  
Student Paper
- 
- 20 M.I. Builders, N.N. Wannang, G.A. Ajoku, P.F. Builders, A. Orisadipe, J.C. Aguiyi. "Evaluation of the Antimalarial Potential of Vernonia ambigua Kotschy and Peyr (Asteraceae)", International Journal of Pharmacology, 2011 <1 %  
Publication
- 
- 21 Priya Agrawal, Talever Singh, Devender Pathak, Himansu Chopra, Kuldeep Singh. "Evaluation of the anxiolytic activity of ethanolic extract of Ziziphus mauritiana Lam. in Swiss albino mice", Pharmacological Research - Modern Chinese Medicine, 2023 <1 %  
Publication
- 
- 22 Yessy Ariesanti, Wiwiek Poedjiastoeti, Gilma Ayu Sriyanto, Yenni Angraini. "Increase of Fibroblast Proliferation by Composite Membrane (Polyvinyl Alcohol - Collagen - Hydroxyapatite)", 2021 IEEE International Conference on Health, Instrumentation & Measurement, and Natural Sciences (InHeNce), 2021 <1 %  
Publication
- 
- 23 repository.unair.ac.id <1 %  
Internet Source
- 

Exclude quotes On

Exclude matches

< 15 words

Exclude bibliography On